ATP-driven Non-equilibrium Activation of Kinase Clients by the Molecular Chaperone Hsp90

Huafeng Xu[1,*]

[1]Silicon Therapeutics, Boston, MA 02210, USA.

*Correspondence: huafeng@gmail.com, +1 917 968 8176

Short Title: Hsp90-mediated client activation

Classification: BIOLOGICAL/PHYSICAL SCIENCES (Biophysics and Computational Biology)

Abstract

The molecular chaperone 90-kDa heat-shock protein (Hsp90) assists the late-stage folding and
activation of diverse types of protein substrates (called clients), including many kinases.
Previous studies have established that the Hsp90 homodimer undergoes an ATP-driven cycle
through open and closed conformations. Here I propose a model of client activation by Hsp90,
which predicts that this cycle enables Hsp90 to use ATP energy to drive a client out of
thermodynamic equilibrium toward its active conformation. My model assumes that an Hsp90-
bound client can transition between a deactivating conformation and an activating conformation.
It suggests that the cochaperone Cdc37 aids Hsp90 to activate kinase clients by differentiating
between these two intermediate conformations. My model makes experimentally testable
predictions, including how modulating the stepwise kinetics of the Hsp90 cycle—for example,
by various cochaperones—affects the activation of different clients. My model may inform
client-specific and cell-type-specific therapeutic intervention of Hsp90-mediated protein
activation.

Keywords: molecular chaperone, non-equilibrium protein folding, heat-shock protein 90, kinase
activation

Introduction

Many proteins depend on molecular chaperones for proper folding into their functional
structures. Hsp90 is responsible for the activation (sometimes referred to as maturation) and
stabilization of a wide range of client proteins—many involved in signal transduction—in the
cell (1, 2), including oncogenic kinases such as v-Src (3, 4). A client of Hsp90 is often first
processed by other chaperones such as Hsp70 into a partially folded state before Hsp90
completes its final maturation into the active state (5, 6). Pharmacological inhibition of Hsp90
ATPase activity results in *in vivo* degradation of its clients (7, 8), making therapeutic
intervention of Hsp90 function a promising approach to the treatment of a variety of diseases
caused by aberrant signal transduction, such as cancer (9), and by protein misfolding and

aggregation, such as Alzheimer's disease (10). Unraveling the mechanism by which Hsp90
activates its clients may help predict how the activities of different clients respond to different
Hsp90 modulations, which may enable the rational targeting of Hsp90 and its cochaperones in
order to achieve better potency and client or cell-type specificity, thus improving therapeutic
efficacy and safety (9).

Highly conserved from bacteria to eukaryotes, Hsp90 functions as a homodimer. Each protomer
consists of three domains: the C-terminal domain (CTD), which provides the site of
dimerization, the middle domain (MD), to which clients bind, and the N-terminal domain (NTD),
which harbors the ATPase capable of nucleotide-binding and ATP hydrolysis. Structural and
biochemical studies have revealed that the Hsp90 dimer undergoes an ATP-driven
conformational cycle (11). The nucleotide-free Hsp90 dimer preferentially adopts the open,
"V"-shaped conformation (Figure 1A). The client can bind to and unbind from the open Hsp90
dimer. ATP-binding induces a reconfiguration of the nucleotide lid in NTD (Figure 1B), which
exposes an additional dimerization interface and strongly biases Hsp90 toward a closed
conformation where the two NTDs also dimerize (Figure 1C). This has been clearly
demonstrated by electron microscopy (12) and single molecule experiments (13) in which Hsp90
incubated with an unhydrolyzable ATP analog 5'-adenylyl-β,γ-imidodiphosphate (AMPPNP) is
only seen in the closed conformation. In the closed conformation, a catalytic loop containing a
conserved arginine from MD is repositioned to complete the catalytic center for ATP hydrolysis.
ATP hydrolysis leads to a compact ADP-bound Hsp90 dimer (Figure 1D) and perhaps the loss of
the client-binding surface (12). Subsequent ADP release then returns the Hsp90 dimer to its
open conformation.

An unsettled question is how Hsp90 activates its clients through its ATP cycle (14). Compared
to other ATP-driven chaperones such as chaperonins and Hsp70, Hsp90 is a very slow ATPase,
with a turn-over rate of 0.05-1.2 ATP per minute and a large K_m in the range of 15 to
250 µM (11). ATP hydrolysis, however, appears indispensable to Hsp90 for its in vivo function
(15, 16) and client activation (17). What is the use of the spent ATP free energy? Some
cochaperones aid Hsp90 in the activation of a specific class of clients, such as Cdc37 in the case
of kinases (18, 19). Their roles in the client activation remain an open question. A number of
cochaperones, such as Aha1 and p23, modulate Hsp90 function by either accelerating or slowing
down the ATP cycle. It is puzzling why the altered cycle timing affects client activation (17).

Here I propose a model of Hsp90-mediated activation of client kinases, which predicts that
Hsp90 uses the free energy from ATP hydrolysis to elevate the active fraction of a client above
its Hsp90-free equilibrium value. A key assumption of my model is that the client can transition
between two intermediate conformations—one favoring activation and the other deactivation—
when it is clamped by Hsp90 in the closed conformation. I will discuss the structural basis of
this assumption and the likely types of conformational changes, using as an example two
hypothetical kinase intermediate conformations; I will also present a mathematical proof that
such conformational transition in an Hsp90-bound client is necessary for its non-equilibrium
activation. My model suggests a new thermodynamic criterion that distinguishes between client
and non-client proteins; it provides a mechanism by which Cdc37 expands Hsp90's kinase
clientele. It explains how cochaperones such as Aha1 affect client activation by changing the
kinetic rates of different steps in Hsp90's ATP cycle. My model implies that Hsp90 is not only
involved in the folding of nascent protein chains but also in the maintenance of their activity.
This and other predictions of my model subject it to experimental tests.

Results

Theoretical (20) and simulation studies (21) have suggested that proteins such as kinases go
through locally unfolded intermediate conformations in order to facilitate the inactive-to-active
transition. Hsp90 binds to the kinase clients in these intermediate conformations, as revealed by
the structures of the Hsp90-Cdc37-Cdk4 complex (22), including an atomic structure by cryoEM
at 3.9 Å resolution (23). In this structure, the closed Hsp90 dimer clamps its client kinase, which
adopts a bilobial structure with its N- and C-lobes separated to two sides of the clamp
(Figure 1K). The two individual lobes are largely folded, but their interface is completely
broken (Figure 1F), which is consistent with the observation that Hsp90 is mostly involved in the
final stage of protein folding and activation. Evidence has emerged that instead of adopting a
single structure, a client bound to Hsp90 is in a fluid conformational ensemble (24).

It is reasonable to presume that the intermediate conformations can be divided into two
categories based on whether they are more likely to transition into the active state or into the
inactive state. Here I assume in a simple model that the client can adopt two locally unfolded
intermediate states (Figure 1F, G; I will discuss later these hypothetical kinase structures in
detail): in the deactivating state (the d-state), the client is strongly disposed to fold into the
inactive state (the i-state); in the maturing state (the m-state), the client is poised to fold into the
active state (the a-state). The client can transition between the i- and the d-states, between the d-
and the m-states, and between the m- and the a-states (Figure 1E-H). Hsp90 can bind to the
client in the intermediate states but not in the inactive or active states.

Stable binding between the client and the open Hsp90 dimer will likely require extensive
contacts between the client and Hsp90 and thus hinder the $d \rightleftharpoons m$ conformational transition. In
the closed Hsp90 dimer, however, the client is clamped between the two Hsp90 protomers and it
cannot move away even if it break its contacts with Hsp90, thus allowing it to interconvert
between the d- and the m-states while still bound to Hsp90. The structures of the closed Hsp90
dimer—conserved across great evolutionary distances—indeed hint at such conformational
transitions. A large hole exists at the center of the dimer, but it is gated by two flexible src loops
(Figure 1C). These gating loops prevent folded domains from slipping through the hole and thus
keep the client trapped, but the remaining space appears large enough to permit unfolded amino
acid chains to thread through.

I thus propose a model (Figure 2 and Table 1) that couples Hsp90's ATP-driven conformational
cycle with the client's conformational changes. In my model, Hsp90 cycles through 4 states: 1)
the open and nucleotide-free state (Hsp90$_\text{open}$), which interconverts—by ATP binding—with 2)
the open and ATP-bound state (Hsp90$_\text{open}^\text{ATP}$), which overwhelmingly converts to the more stable
3) closed and ATP-bound state (Hsp90$_\text{closed}^\text{ATP}$), which, following ATP hydrolysis, converts to 4)
the compact and ADP-bound state (Hsp90$_\text{compact}^\text{ADP}$), which, upon ADP release, returns to the open
and nucleotide-free state. Although each state consists of multiple conformational sub-states
(25), and the two protomers hydrolyze ATP at a deterministic sequential order (26), for
simplicity these details are omitted from the current model. The client in the d- or the m-state,
using hydrophobic patches exposed by the local unfolding, can reversibly bind to Hsp90$_\text{open}$ and
Hsp90$_\text{open}^\text{ATP}$. When the client is in the solution or when it is clamped in Hsp90$_\text{closed}^\text{ATP}$, it can
interconvert between the d- and m-states.

My model implies the following pathway of client activation. First, an inactive client locally
unfolds to the d-state and binds to an open Hsp90 dimer, which closes upon ATP binding. The
client then transitions from the d- to the m-state. Hsp90 hydrolyzes ATP. After ADP-release,

the Hsp90 dimer reopens, allowing the maturing client to unbind and fold into its active

conformation. Denoting a complex between molecules x and y as $x \cdot y$, this d-to-m pathway can

be written as

$$i \rightleftharpoons d \quad \overset{+\text{Hsp90}_{\text{open}}}{\rightleftharpoons} \quad d \cdot \text{Hsp90}_{\text{open}} \quad \overset{+\text{ATP}}{\rightleftharpoons} \quad d \cdot \text{Hsp90}_{\text{open}}^{\text{ATP}} \rightleftharpoons d \cdot \text{Hsp90}_{\text{closed}}^{\text{ATP}}$$

$$\rightleftharpoons m \cdot \text{Hsp90}_{\text{closed}}^{\text{ATP}} \overset{-\text{Pi}}{\rightleftharpoons} m \cdot \text{Hsp90}_{\text{compact}}^{\text{ADP}} \overset{-\text{ADP}}{\rightleftharpoons} m \cdot \text{Hsp90}_{\text{open}} \overset{-\text{Hsp90}_{\text{open}}}{\rightleftharpoons} m \rightleftharpoons a$$

There is a corresponding m-to-d pathway—swapping i with a and d with m—that deactivates the

client:

$$a \rightleftharpoons m \quad \overset{+\text{Hsp90}_{\text{open}}}{\rightleftharpoons} \quad m \cdot \text{Hsp90}_{\text{open}} \quad \overset{+\text{ATP}}{\rightleftharpoons} \quad m \cdot \text{Hsp90}_{\text{open}}^{\text{ATP}} \rightleftharpoons m \cdot \text{Hsp90}_{\text{closed}}^{\text{ATP}}$$

$$\rightleftharpoons d \cdot \text{Hsp90}_{\text{closed}}^{\text{ATP}} \overset{-\text{Pi}}{\rightleftharpoons} d \cdot \text{Hsp90}_{\text{compact}}^{\text{ADP}} \overset{-\text{ADP}}{\rightleftharpoons} d \cdot \text{Hsp90}_{\text{open}} \overset{-\text{Hsp90}_{\text{open}}}{\rightleftharpoons} d \rightleftharpoons i$$

Note the directionality of the ATP-driven Hsp90 conformational cycle (one ATP molecule is

hydrolyzed in each pathway above); the second m-to-d pathway is not the reverse of the first d-

to-m pathway.

Hsp90 can drive the client toward the active state if the d-to-m pathway occurs more than the m-

to-d pathway. How can this happen at the steady state? One possible mechanism is that the

open Hsp90 dimer has a higher affinity for the client in the d-state than for the client in the m-

state. This promotes the d-to-m pathway over the m-to-d pathway in the steps of client binding

and release: a client molecule in the d-state is more likely than one in the m-state to bind to

Hsp90; a client molecule in the m-state is more likely than one in the d-state to unbind from

Hsp90. But it also means that the complex $d \cdot \text{Hsp90}_{\text{open}}$ is more stable than the complex $m \cdot$

$\text{Hsp90}_{\text{open}}$ and the external free energy from ATP is needed to drive the d-to-m pathway from $d \cdot$

$\text{Hsp90}_{\text{open}}$ to $m \cdot \text{Hsp90}_{\text{open}}$. Key to this mechanism is the overwhelming tendency of the ATP-

bound Hsp90 dimer to close: $d \cdot \text{Hsp90}_{\text{open}}^{\text{ATP}}$ is more stable than $m \cdot \text{Hsp90}_{\text{open}}^{\text{ATP}}$, which implies a

higher free energy penalty for the step of $d \cdot \text{Hsp90}_{\text{open}}^{\text{ATP}} \rightleftharpoons d \cdot \text{Hsp90}_{\text{close}}^{\text{ATP}}$ in the d-to-m pathway

than for $m \cdot \text{Hsp90}_{\text{open}}^{\text{ATP}} \rightleftharpoons m \cdot \text{Hsp90}_{\text{close}}^{\text{ATP}}$ in the m-to-d pathway, but this difference is negligible

in comparison to the large thermodynamic motive for the Hsp90 dimer to close in both

complexes.

To test whether this mechanism is consistent with experimentally observed client activation by

Hsp90, I used my model to analyze the activation of v-Src, a stringent Hsp90 client (4). The

normalized activity of v-Src—measured experimentally as the activity ratio between v-Src

incubated with and without Hsp90—is computed as the ratio of the active (*i.e. a*-state)

concentration at the steady state to that at the Hsp90-free equilibrium (Figure 3A). The

calculated normalized activity at the steady state is greater than 1 in the presence of Hsp90,

affirming that the proposed mechanism enables Hsp90 to convert ATP energy into non-

equilibrium stabilization of the active state of v-Src.

The more Hsp90 prefers the binding to a protein substrate in its d-state to the binding to the

protein in its m-state, the more Hsp90 elevates the substrate's steady state activity (Figure 4A).

Presence of such an affinity difference may be a thermodynamic criterion that distinguishes the

clients from the non-clients. But in the absence of a structural basis that systematically biases
Hsp90 to bind a protein in its d-state stronger than the protein in its m-state, it is unlikely that
Hsp90 solely depends on this difference in recognizing clients.

My model suggests a mechanism for Hsp90 to expand its clientele by cooperating with client-
class-specific cochaperones (14, 27). One such example is Cdc37, which plays a critical role in
activating Hsp90-dependent kinases (18). Experimental evidence (19) suggests that Cdc37 binds
to a locally unfolded conformation of the client kinase and stabilizes it for Hsp90-binding.
Cdc37 can form ternary complex with the locally unfolded client kinase and Hsp90 in the open
and closed states (18), but it may be expelled from the complex when Hsp90 changes to the more
compact conformation upon ADP hydrolysis. If Cdc37 selectively binds to the client in the d-
state but not to the client in the m-state (the structural basis for this selectivity is discussed
below), it will provide an additional pathway for the client to move from the d-state to the m-
state through the Hsp90 conformational cycle:

$$167 \quad d \;\underset{}{\overset{+\text{Cdc37}}{\rightleftharpoons}}\; d\cdot\text{Cdc37} \;\underset{}{\overset{+\text{Hsp90}_{\text{open}}^{\text{ATP}}}{\rightleftharpoons}}\; d\cdot\text{Cdc37}\cdot\text{Hsp90}_{\text{open}}^{\text{ATP}}$$

$$168 \quad \rightleftharpoons d\cdot\text{Cdc37}\cdot\text{Hsp90}_{\text{closed}}^{\text{ATP}} \;\underset{}{\overset{-\text{Cdc37}}{\rightleftharpoons}}\; d\cdot\text{Hsp90}_{\text{closed}}^{\text{ATP}}$$

$$169 \quad \rightleftharpoons m\cdot\text{Hsp90}_{\text{closed}}^{\text{ATP}} \;\underset{}{\overset{-\text{Pi}}{\rightleftharpoons}}\; m\cdot\text{Hsp90}_{\text{compact}}^{\text{ADP}} \;\underset{}{\overset{-\text{ADP}}{\rightleftharpoons}}\; m\cdot\text{Hsp90}_{\text{open}} \;\underset{}{\overset{-\text{Hsp90}_{\text{open}}}{\rightleftharpoons}}\; m$$

In this pathway, Cdc37 unbinds from the client in the d-state before the Hsp90 dimer enters the
compact conformation (the unbinding can happen either before or after the ATP hydrolysis),
allowing the client to transition from the d-state to the m-state in the closed Hsp90 dimer. Cdc37
thus selects the client molecules in the d-state to enter the Hsp90 cycle but allows some of them
to exit the cycle in the m-state, which boosts the d-to-m transition.

My model indeed captures Cdc37's augmentative effect on Hsp90's non-equilibrium activation
of kinases (Figures 3 and 4A). Cdc37 substantially increases the maximum normalized activity
at the steady state; it also decreases the Hsp90 concentration required to achieve the maximum
activation. The net reactive fluxes at the steady state (Figure 2) demonstrate that the client in the
d-state enters the Hsp90 cycle either alone or in complex with Cdc37. There is a net flux for the
client to convert from the d- to the m-state when clamped in the closed Hsp90 dimer, signifying
that the d-to-m pathway dominates the m-to-d pathway. Cdc37 enables Hsp90 to elevate the
steady state activity of kinases even when Hsp90 on its own cannot differentiate between the d-
and the m-states (Figure 4A).

Is there a structural basis for Cdc37 to differentiate between the d- and the m-states of a client
kinase? Kinase activation invariably entails the reconfiguration of its DFG activation loop from
a disordered ensemble of structures into a well-ordered structure bridging the N- and C-lobes of
the kinase (Figure 1E, H)(28, 29). The DFG loop of Cdk4 in the Hsp90-Cdc37-Cdk4 structure is
similar to that of an inactive kinase. In contrast, superimposing the C-lobe of the active Cdk2
kinase (a close homolog of Cdk4) onto Cdk4 in the same structure results in substantial steric
clashes between the DFG loop and Cdc37 (Figure 1J). These observations suggest a pair of
possible structural models for the d- and m-states of a client kinase: both have the N- and C-lobes
separated; the d-state structure has a disordered DFG loop, but the m-state structure has the DFG
loop in its active configuration, which prevents Cdc37 binding. In addition, the β4-αC loop may
move from the N-lobe side in the d-state to dock onto the C-lobe in the m-state, which would

also disrupt Cdc37 binding, because Cdc37 binds to the C-lobe using a 5-residue ($_{20}$HPNID$_{24}$)
loop mimicking the β4-αC loop; such a move seems permissible by the gated central hole in the
closed Hsp90 dimer. Cdc37 can thus discriminate the *d-* and the *m-*states by the positions of the
DFG-loop and perhaps of the β4-αC loop, a mechanism likely applicable across all kinase
clients.

Does a client indeed change conformations while bound to Hsp90? Much remains to be learned
about the structures of clients in complex with Hsp90 (30), but the molten globule state observed
for a Hsp90-bound client (24) and the gated hole in the closed Hsp90 dimer structure hint at the
proposed conformational change. To shore up the proposed mechanism, here I prove
mathematically that for Hsp90 to use ATP energy for non-equilibrium activation of the client,
the client must undergo a conformational transition while it is bound to Hsp90. This follows
from a general theorem that I prove in Methods: at the steady state, two molecular species
connected by a chemical reaction can only be out-of-equilibrium, *i.e.* a non-zero flux is sustained
in the reaction, if this reaction is part of a reaction cycle—defined as a set of reactions that
together return all molecular species to their original states (see Methods)—in which external
chemical energy is consumed. If the client does not undergo a transition between two
interconverting conformations, say x and y, during the time that it is bound to Hsp90, the reaction
$x \rightleftharpoons y$ cannot be part of an energy-consuming reaction cycle involving the client, Hsp90, and
their complexes, because it is the only reaction in which x and y appear on the different sides of
reactants and products. The steady state reactive flux between any two conformations of the apo
client is thus zero, their concentration ratio is the same as the equilibrium value, and non-
equilibrium activation is impossible. If non-equilibrium client activation is experimentally
demonstrated (see Discussion), this theorem implies that a conformational transition must take
place in the Hsp90-bound client.

I used my model to estimate ATP consumption by Hsp90 in non-equilibrium client activation.
Figure 3B shows the calculated time course of the activation of v-Src—starting from the
equilibrium concentrations—and the accompanying ATP consumption. In the initial phase of
activation (ending when the active concentration reaches midway between the starting and the
steady state values), roughly 14 ATP molecules are hydrolyzed per each activated v-Src
molecule. This is considerably smaller than the ATP consumption in Hsp70-mediated folding
(>100 ATP molecules per folded molecule of protein substrate)(31).

My model helps to unravel how cochaperones affect Hsp90 function by altering the kinetics of
different steps in the ATP cycle. To maintain non-equilibrium activation, the cycle time has to
be sufficiently fast to outpace the client's return to equilibrium. On the other hand, Hsp90 has to
dwell sufficiently long both in the open state to accommodate client binding and release and in
the closed, pre-hydrolysis state to allow client conformational transition. Calculations by my
model demonstrate that changes in the kinetic rates of different steps in the Hsp90 cycle affect
client activation differently (Figure 5A-C).

By considering a cochaperone's effects on the kinetics of the cycle, my model can predict its role
in the activation of different clients by Hsp90. For instance, the cochaperone Aha1 stimulates
the ATP turn-over by accelerating the change from the open to the closed Hsp90 dimer, but it
likely stabilizes a conformation that is not fully hydrolysis-competent (32) and it may have to
unbind from Hsp90 before ATP hydrolysis can proceed. It thus have two competing effects: 1)
accelerating the conformational closure of the Hsp90 dimer, which enhances client activation
(Figure 5A); 2) slowing down the step of ATP hydrolysis, which may compromise client
activation (Figure 5B). To analyze Aha1's overall effect on client activation, I included Aha1
into my model (Table 2), and calculated the normalized activities of two hypothetical clients at

different Aha1 concentrations (Figure 5D). Different clients either accelerate (33, 34) or slow
down (35) the cycle. For a client that on its own slows down the Hsp90 dimer closure, Aha1 can
augment its steady state activity by accelerating the closure. For a client that on its own
accelerates the closure, however, Aha1 may diminish its steady state activity by slowing down
the ATP hydrolysis. This analysis may explain the paradoxical observation that Aha1
downregulation promotes the Hsp90-dependent folding of a common disease variant of cystic
fibrosis transmembrane conductance regulator (36).

Discussion

My model predicts that Hsp90, like two other ATP-consuming chaperones Hsp70 (31) and
GroEL (37), is an agent of non-equilibrium protein-folding. It suggests that Hsp90 is not only
involved in the initial folding of nascent protein chains, but it is also responsible for maintaining
the activity of mature clients, which requires sustained ATP expenditure. This prediction is
consistent with Hsp90's role at buffering phenotypic variations against destabilizing mutations to
client proteins (38) and with the experimental observation that Hsp90 ATPase inhibitors such as
geldanamycin can suppress client protein activities and induce their degradation in cells (39).
Given that geldanamycin's effect *in vivo* may be more apparent on the nascent chains than on the
mature proteins (40, 41), it is worthwhile to directly test this prediction by an *in vitro*
experiment: first measure the activity of a client kinase maintained in the steady state with Hsp90
and an ATP regeneration system, then add geldanamycin to stop the Hsp90 cycle and measure
the kinase activity after a new steady state is established. A decrease in activity will demonstrate
ATP-driven non-equilibrium client activation by Hsp90.

The key assumption of my model is the existence of two interconverting intermediate client
conformations, one favoring activation and the other deactivation. Here I propose the following
experiment to test this assumption. A client kinase is first trapped in the closed Hsp90 dimer by
incubation with a nonhydrolysable ATP analog such as AMPPNP. An ATP competitive
inhibitor, perhaps geldanamycin, is then added in excess, opening the Hsp90 dimer and releasing
the client, and the activity of the client is measured. My model predicts that at increasing
concentrations of Cdc37 during the incubation, higher fractions of the trapped client will be in
the d-state (Figure 4B) and forming the ternary complex $d \cdot \text{Cdc37} \cdot \text{Hsp90}_{\text{closed}}^{\text{AMPPNP}}$, leaving
smaller fractions in the m-state for ready activation. Consequently, in the initial moments after
the client release, the measured activity will increase more slowly at higher Cdc37
concentrations.

My model predicts that client activation by Hsp90 may be promoted or suppressed by
modulating the stepwise kinetics of the ATP cycle. These predictions can be tested by designing
Hsp90 mutants that alter the stability of different conformational states of Hsp90 and testing their
effects on the timing of the ATP cycle and on client activation (17). My model implies that the
same cochaperone may affect different clients differently, depending on how the cochaperone
and how the client each alters the timings of the ATP cycle, which may be tested experimentally.
In addition, my model implies that the occupancies of Hsp90 conformational states will change
with the concentration levels of different clients and of various cochaperones, which may help
explain the observation that 17-allylaminogeldanamycin, a clinical-stage Hsp90 inhibitor, binds
to Hsp90 in tumor cells with about two-orders-of-magnitude higher affinity than to Hsp90 in

normal cells (42). These implications of my model raise the possibility of client-specific or cell-
type-specific targeting of Hsp90 for therapeutic applications.

Tables

Table 1. Reactions and kinetic parameters for Hsp90-activation of v-Src kinase. Hsp90 is
abbreviated as H90, and Cdc37 as C37. The boldfaced values are fitted to the experimental data
of Hsp90-activation of v-Src. The underlined values are uniquely determined from other
parameters so that the total free energy changes in any reaction cycle is zero. The kinetic
parameters of the Hsp90 cycle (reactions 1 to 4) are derived from kinetic analysis of nucleotide-
driven changes in fluorescence resonance energy transfer in labeled Hsp90 (25), with the k_f of
reaction 1 adjusted by a factor of 2 so that the ATPase has a K_m of 250 µM (11). The kinetic
parameters for v-Src conformational transitions—experimentally unavailable—are chosen
arbitrarily, except for k_f of reaction 7. The k_f of reactions 11d and 11m reflects a 20-fold
acceleration of the closing of the Hsp90 dimer induced by the client (33, 34). Other than this
acceleration, the client is assumed not to affect the kinetics of other steps of the ATP-driven
cycle (reactions 8 to 13). I also assume that the rates of client's conformational transition is the
same in the closed Hsp90 dimer (reaction 14) as in the solution (reaction 6). All protein-protein
association rate constants are taken to be 1 µM^{-1}s^{-1}, a reasonable value, except for reactions 19a
and 20, which should be slower because Cdc37 has to adopt a specific conformation in order to
simultaneously bind to both Hsp90 and the client. The dissociation rate constant of Cdc37 from
Hsp90 (reaction 15) is computed from its binding constant $K_D = 1.4$ µM (43, 44). The
dissociation rate constant of reaction 20 is arbitrary because the model predictions are insensitive
to its value.

No.	Reaction	k_f	k_r
1	$H90_{open} + ATP \rightleftharpoons H90^{ATP}_{open}$	6.7×10^{-4} µM^{-1}s^{-1}	1.45×10^{-1} s^{-1}
2	$H90^{ATP}_{open} \rightleftharpoons H90^{ATP}_{closed}$	1.7×10^{-2} s^{-1}	5×10^{-5} s^{-1}
3	$H90^{ATP}_{closed} \rightleftharpoons H90^{ADP}_{compact} + Pi$	3.3×10^{-1} s^{-1}	$\underline{9.3\times10^{-14}}$ µM^{-1}s^{-1}
4	$H90^{ADP}_{compact} \rightleftharpoons H90_{open} + ADP$	1 s^{-1}	1×10^{-1} µM^{-1}s^{-1}
5	$i \rightleftharpoons d$	1 s^{-1}	5×10^{2} s^{-1}
6	$d \rightleftharpoons m$	1×10^{-1} s^{-1}	1×10^{-1} s^{-1}
7	$m \rightleftharpoons a$	$\mathbf{6.5\times10^{1}}$ s^{-1}	1 s^{-1}
8d	$d + H90_{open} \rightleftharpoons d \cdot H90_{open}$	1 µM^{-1}s^{-1}	$\mathbf{5.6\times10^{-3}}$ s^{-1}
8m	$m + H90_{open} \rightleftharpoons m \cdot H90_{open}$	1 µM^{-1}s^{-1}	$\mathbf{6.2\times10^{-1}}$ s^{-1}
9d	$d + H90^{ATP}_{open} \rightleftharpoons d \cdot H90^{ATP}_{open}$	Same as 8d	Same as 8d
9m	$m + H90^{ATP}_{open} \rightleftharpoons m \cdot H90^{ATP}_{open}$	Same as 8m	Same as 8m
10d	$d \cdot H90_{open} + ATP \rightleftharpoons d \cdot H90^{ATP}_{open}$	Same as 1	Same as 1
10m	$m \cdot H90_{open} + ATP \rightleftharpoons m \cdot H90^{ATP}_{open}$	Same as 1	Same as 1
11d	$d \cdot H90^{ATP}_{open} \rightleftharpoons d \cdot H90^{ATP}_{closed}$	3.3×10^{-1} s^{-1}	$\mathbf{5.4\times10^{-3}}$ s^{-1}
11m	$m \cdot H90^{ATP}_{open} \rightleftharpoons m \cdot H90^{ATP}_{closed}$	Same as 11d	Same as 2

No.	Reaction	k_f	k_r
12d	$d \cdot H90_{closed}^{ATP} \rightleftharpoons d \cdot H90_{compact}^{ADP} + Pi$	Same as 3	Same as 3
12m	$m \cdot H90_{closed}^{ATP} \rightleftharpoons m \cdot H90_{compact}^{ADP} + Pi$	Same as 3	Same as 3
13d	$d \cdot H90_{compact}^{ADP} \rightleftharpoons d \cdot H90_{open} + ADP$	Same as 4	$\underline{1.8\times10^{-2}}$ $\mu M^{-1}s^{-1}$
13m	$m \cdot H90_{compact}^{ADP} \rightleftharpoons m \cdot H90_{open} + ADP$	Same as 4	$\underline{2.0}$ $\mu M^{-1}s^{-1}$
14	$d \cdot H90_{closed}^{ATP} \rightleftharpoons m \cdot H90_{closed}^{ATP}$	Same as 6	Same as 6
15	$C37 + H90_{open} \rightleftharpoons C37 \cdot H90_{open}$	1 $\mu M^{-1}s^{-1}$	1.4 s^{-1}
16	$C37 + d \rightleftharpoons d \cdot C37$	1 $\mu M^{-1}s^{-1}$	$\mathbf{5.1\times10^{-3}}$ s^{-1}
17	$d + C37 \cdot H90_{open} \rightleftharpoons d \cdot C37 \cdot H90_{open}$	Same as 16	$\underline{4.8\times10^{-4}}$ s^{-1}
18a	$d \cdot C37 + H90_{open} \rightleftharpoons d \cdot C37 \cdot H90_{open}$	1 $\mu M^{-1}s^{-1}$	$\underline{1.3\times10^{-1}}$ s^{-1}
18n	$d \cdot C37 + H90_{open}^{ATP} \rightleftharpoons d \cdot C37 \cdot H90_{open}^{ATP}$	Same as 18a	$\underline{2.8\times10^{-2}}$ s^{-1}
19a	$C37 + d \cdot H90_{open} \rightleftharpoons d \cdot C37{\cdot}H90_{open}$	2×10^{-1} $\mu M^{-1}s^{-1}$	$\mathbf{2.4\times10^{-2}}$ s^{-1}
19n	$C37 + d \cdot H90_{open}^{ATP} \rightleftharpoons d \cdot C37{\cdot}H90_{open}^{ATP}$	1 $\mu M^{-1}s^{-1}$	$\mathbf{2.5\times10^{-2}}$ s^{-1}
20	$C37 + d \cdot H90_{closed}^{ATP} \rightleftharpoons d \cdot C37{\cdot}H90_{closed}^{ATP}$	1×10^{-1} $\mu M^{-1}s^{-1}$	1×10^{-1} s^{-1}
21	$d \cdot C37 \cdot H90_{open} + ATP \rightleftharpoons d \cdot C37 \cdot H90_{open}^{ATP}$	Same as 1	$\underline{3\times10^{-2}}$ s^{-1}
22	$d \cdot C37 \cdot H90_{open}^{ATP} \rightleftharpoons d \cdot C37{\cdot}H90_{closed}^{ATP}$	Same as 11d	$\underline{2.2\times10^{-1}}$ s^{-1}

Table 2. Reactions and kinetic parameters for the cochaperone Aha1. The dissociation rate
constants are computed from the binding affinities measured by surface plasmon resonance (32).
I assume that Aha1 accelerates the closure of the Hsp90 dimer by 10-fold (45). The rate of
client-bound Hsp90 dimer closure in the presence of Aha1 (reactions 26d, 26m) is given by the
larger k_f of the reaction 11d and of the reaction 25; the reverse rate is computed from the reaction
cycle of the indicated reactions.

No.	Reaction	k_f	k_r
23a	$H90_{open} + Aha1 \rightleftharpoons Aha1 \cdot H90_{open}$	1 $\mu M^{-1}s^{-1}$	1.2 s^{-1}
23n	$H90_{open}^{ATP} + Aha1 \rightleftharpoons Aha1 \cdot H90_{open}^{ATP}$	Same as 1a	Same as 1a
24	$H90_{closed}^{ATP} + Aha1 \rightleftharpoons Aha1 \cdot H90_{closed}^{ATP}$	1 $\mu M^{-1}s^{-1}$	1.6×10^{-1} s^{-1}
25	$Aha1 \cdot H90_{open}^{ATP} \rightleftharpoons Aha1 \cdot H90_{closed}^{ATP}$	1.7×10^{-1} s^{-1}	$\underline{6.67\times10^{-5}}$ s^{-1}
26d	$d \cdot Aha1 \cdot H90_{open}^{ATP} \rightleftharpoons d \cdot Aha1 \cdot H90_{closed}^{ATP}$	Max(25 and 11d)	$\underline{\text{From 11d, 23n, 24}}$
26m	$m \cdot Aha1 \cdot H90_{open}^{ATP} \rightleftharpoons m \cdot Aha1 \cdot H90_{closed}^{ATP}$	Max(25 and 11m)	$\underline{\text{From 11m, 23n, 24}}$

312 ## Methods

313 **Numerical solution of the molecular concentrations at the steady state.** Consider a set of K
314 chemical reactions involving M molecular species:

$$\sum_{i=1}^{M} C_{ki}X_i = 0, \ k = 1,2,\cdots,K, \tag{1}$$

315   where X_i is the i'th molecular species and C_{ki} is the stoichiometry of X_i in the k'th reaction:
316     $C_{ki} > 0$ indicates that X_i is a reactant, $C_{ki} < 0$ indicates that X_i is a product, and $C_{ki} = 0$
317   indicates that X_i is not involved in the reaction. The reactive flux for the k'th reaction is

$$J_k = f_k \prod_{i|C_{ki}>0} x_i^{C_{ki}} - r_k \prod_{i|C_{ki}<0} x_i^{-C_{ki}}, \tag{2}$$

where $x_i = [X_i]$ is the concentration (or the activity in the case of non-ideal solutions) of X_i and
f_k and r_k are the forward and reverse rate constants.

Let C be the $K \times M$ matrix of elements C_{ki} and J be the K-vector of components J_k, the rate of
change for the molecular concentrations is given by $dx/dt = -C^t \cdot J$, with the superscript t
denoting matrix transpose.

Let $N = \text{null}(C)$ be the $M \times (M - \text{rank}(C))$ matrix representing the null space of C, then the
time derivative of $N^t \cdot x$ is

$$N^t \cdot dx/dt = -(C \cdot N)^t \cdot J = 0. \tag{3}$$

325   Thus

$$N^t \cdot x = N^t \cdot x(t = 0) \tag{4}$$

326   is constant, which reflects mass conservation; $N^t \cdot x(t = 0)$ gives the initial condition.

327   At the steady state,

$$C^t \cdot J = 0. \tag{5}$$

328   Equations 4 and 5 together constitute the steady state equations. To reduce them to linearly
329   independent equations, take the singular value decomposition $C = U \cdot \Sigma \cdot V^t$. Let Σ_s be the
330     $\text{rank}(C) \times \text{rank}(C)$ diagonal submatrix of Σ with the singular values being its diagonal elements
331   and let U_s be the $K \times \text{rank}(C)$ matrix consisting of the first $\text{rank}(C)$ column vectors of U (these
332   vectors span the range of C). Equation 5 becomes

$$0 = C^t \cdot J = V \begin{pmatrix} \Sigma_s \cdot U_s^t \cdot J \\ 0_{(M-\text{rank}(C)) \times K} \end{pmatrix}. \tag{6}$$

333   Because both V and Σ_s are full-rank, Equation 6 implies

$$U_s^t \cdot J = 0 \tag{7}$$

Equations 7 and 4 are the independent equations for the steady state concentrations.

Checking that there is no spurious free energy production in the modeled set of reactions.
Because my model is used to explore the conversion of ATP energy into non-equilibrium
activation of Hsp90 clients, it must ensure that the total free energy change for any reaction cycle
is zero. A reaction cycle is a subset of reactions that return all molecular species to their original
states. It can be represented by a set of coefficients Q_k such that

$$\sum_{k=1}^{K} Q_k C_{ki} = 0, \; i = 1,2,\cdots,M, \tag{8}$$

340   or in matrix form

$$C^t \cdot Q = 0. \tag{9}$$

341   The reaction cycles thus correspond to the null space of C^t. The total (unit-less) free energy
342   change for any reaction cycle Q,

$$\Delta\Delta G_Q = -\sum_{k=1}^{K} Q_k \ln(f_k/r_k), \tag{10}$$

must be zero. This has been verified for the reactions in Tables 1 and 2 and the reaction of ATP
hydrolysis in aqueous solution: $ATP + H_2O = ADP + Pi$, which together constitute a closed
chemical system with no external free energy input.

Proof that a reaction can have a non-vanishing steady state flux only if it is part of a
reaction cycle that consumes external free energy. Consider the set of reactions in Equation 1
at the steady state. The unit-less free energy change for the k'th reaction is

$$\Delta\Delta G_k = -\sum_i C_{ki}\ln x_i - \ln(f_k/r_k), \tag{11}$$

349   Equations 5 and 9 imply that for any non-vanishing steady state flux $J \neq 0$ we have a
350   corresponding reaction cycle $Q = J$. For any reaction cycle that does not consume external
351   chemical energy, the total change of free energy must be 0, that is

$$\sum_k J_k \Delta\Delta G_k = 0. \tag{12}$$

352   On the other hand, a reaction proceeds in the direction of free energy decrease, *i.e.*, $J_k\Delta\Delta G_k \leq 0$,
353   hence $\forall k, J_k = 0$. Q. E. D. To sustain $J \neq 0$ at the steady state, chemical energy must enter the
354   reaction cycle from molecules with an external source and sink.

Author Contributions

356   HX confirms being the sole contributor of this work and this paper's single author.

Additional Information

358   The author declares no competing interests.

References

1. M. Taipale, D. F. Jarosz, S. Lindquist, HSP90 at the hub of protein homeostasis: Emerging mechanistic insights. *Nat. Rev. Mol. Cell Biol.* **11**, 515–528 (2010).

2. F. H. Schopf, M. M. Biebl, J. Buchner, The HSP90 chaperone machinery. *Nat. Rev. Mol. Cell Biol.* **18**, 345–360 (2017).

3. Y. Xu, S. Lindquist, Heat-shock protein hsp90 governs the activity of pp60v-src kinase. *Proc. Natl. Acad. Sci.* **90**, 7074–7078 (1993).

4. E. E. Boczek, L. G. Reefschläger, M. Dehling, T. J. Struller, E. Häusler, A. Seidl, V. R. I. Kaila, J. Buchner, Conformational processing of oncogenic v-Src kinase by the molecular chaperone Hsp90. *Proc. Natl. Acad. Sci.* **112**, E3189–E3198 (2015).

5. U. Jakob, H. Lilie, I. Meyer, J. Buchner, Transient interaction of Hsp90 with early unfolding intermediates of citrate synthase IMPLICATIONS FOR HEAT SHOCK IN VIVO. *J. Biol. Chem.* **270**, 7288–7294 (1995).

6. E. Kirschke, D. Goswami, D. Southworth, P. R. Griffin, D. A. Agard, Glucocorticoid receptor function regulated by coordinated action of the Hsp90 and Hsp70 chaperone cycles. *Cell* **157**, 1685–1697 (2014).

7. L. Whitesell, E. G. Mimnaugh, B. De Costa, C. E. Myers, L. M. Neckers, Inhibition of heat shock protein HSP90-pp60v-src heteroprotein complex formation by benzoquinone ansamycins: essential role for stress proteins in oncogenic transformation. *Proc. Natl. Acad. Sci.* **91**, 8324–8328 (1994).

8. W. G. An, T. W. Schulte, L. M. Neckers, The heat shock protein 90 antagonist geldanamycin alters chaperone association with p210bcr-abl and v-src proteins before their degradation by the proteasome. *Cell Growth Differ.* **11**, 355–360 (2000).

9. J. Trepel, M. Mollapour, G. Giaccone, L. Neckers, Targeting the dynamic HSP90 complex in cancer. *Nat. Rev. Cancer* **10**, 537–549 (2010).

10. L. B. Shelton, J. D. Baker, D. Zheng, L. E. Sullivan, P. K. Solanki, J. M. Webster, Z. Sun, J. J. Sabbagh, B. A. Nordhues, J. Koren, S. Ghosh, B. S. J. Blagg, L. J. Blair, C. A. Dickey, Hsp90 activator Aha1 drives production of pathological tau aggregates. *Proc. Natl. Acad. Sci.* **114**, 9707–9712 (2017).

11. K. A. Krukenberg, T. O. Street, L. A. Lavery, D. A. Agard, Conformational dynamics of the molecular chaperone Hsp90. *Q. Rev. Biophys.* **44**, 229–255 (2011).

12. A. K. Shiau, S. F. Harris, D. R. Southworth, D. A. Agard, Structural analysis of E. coli hsp90 reveals dramatic nucleotide-dependent conformational rearrangements. *Cell* **127**, 329–340 (2006).

13. M. Mickler, M. Hessling, C. Ratzke, J. Buchner, T. Hugel, The large conformational changes of Hsp90 are only weakly coupled to ATP hydrolysis. *Nat. Struct. Mol. Biol.* **16**, 281–286 (2009).

14. L. H. Pearl, Review: The HSP90 molecular chaperone—an enigmatic ATPase. *Biopolymers* **105**, 594–607 (2016).

15. W. Obermann, H. Sondermann, A. A. Russo, N. P. Pavletich, U. F. Hartl, In Vivo Function of Hsp90 Is Dependent on ATP Binding and ATP Hydrolysis. *J. Cell Biol.* **143**,

400    901–910 (1998).

401 16. B. Panaretou, C. Prodromou, S. M. Roe, R. O'Brien, J. E. Ladbury, P. W. Piper, L. H.
402    Pearl, ATP binding and hydrolysis are essential to the function of the Hsp90 molecular
403    chaperone in vivo. *EMBO J.* **17**, 4829–4836 (1998).

404 17. B. K. Zierer, M. Rübbelke, F. Tippel, T. Madl, F. H. Schopf, D. A. Rutz, K. Richter, M.
405    Sattler, J. Buchner, Importance of cycle timing for the function of the molecular
406    chaperone Hsp90. *Nat. Struct. Mol. Biol.* **23**, 1020–1028 (2016).

407 18. K. A. Verba, D. A. Agard, How Hsp90 and Cdc37 lubricate kinase molecular switches.
408    *Trends Biochem. Sci.* **42**, 799–811 (2017).

409 19. D. Keramisanou, A. Aboalroub, Z. Zhang, W. Liu, D. Marshall, A. Diviney, R. W.
410    Larsen, R. Landgraf, I. Gelis, Molecular mechanism of protein kinase recognition and
411    sorting by the Hsp90 kinome-specific cochaperone Cdc37. *Mol. Cell* **62**, 260–271 (2016).

412 20. O. Miyashita, J. N. Onuchic, P. G. Wolynes, Nonlinear elasticity, proteinquakes, and the
413    energy landscapes of functional transitions in proteins. *Proc. Natl. Acad. Sci.* **100**, 12570–
414    12575 (2003).

415 21. Y. Shan, A. Arkhipov, E. T. Kim, A. C. Pan, D. E. Shaw, Transitions to catalytically
416    inactive conformations in EGFR kinase. *Proc. Natl. Acad. Sci.* **110**, 7270–7275 (2013).

417 22. C. K. Vaughan, U. Gohlke, F. Sobott, V. M. Good, M. M. U. Ali, C. Prodromou, C. V
418    Robinson, H. R. Saibil, L. H. Pearl, Structure of an Hsp90-Cdc37-Cdk4 complex. *Mol.*
419    *Cell* **23**, 697–707 (2006).

420 23. K. A. Verba, R. Y.-R. Wang, A. Arakawa, Y. Liu, M. Shirouzu, S. Yokoyama, D. A.
421    Agard, Atomic structure of Hsp90-Cdc37-Cdk4 reveals that Hsp90 traps and stabilizes an
422    unfolded kinase. *Science* **352**, 1542–1547 (2016).

423 24. S. J. Park, B. N. Borin, M. A. Martinez-Yamout, H. J. Dyson, The client protein p53
424    adopts a molten globule–like state in the presence of Hsp90. *Nat. Struct. Mol. Biol.* **18**,
425    537–541 (2011).

426 25. M. Hessling, K. Richter, J. Buchner, Dissection of the ATP-induced conformational cycle
427    of the molecular chaperone Hsp90. *Nat. Struct. Mol. Biol.* **16**, 287–293 (2009).

428 26. D. Elnatan, M. Betegon, Y. Liu, T. Ramelot, M. A. Kennedy, D. A. Agard, Symmetry
429    broken and rebroken during the ATP hydrolysis cycle of the mitochondrial Hsp90
430    TRAP1. *eLife* **6**, e25235 (2017).

431 27. M. B. Cox, J. L. Johnson, "Evidence for Hsp90 co-chaperones in regulating Hsp90
432    function and promoting client protein folding" in *Chaperones: Methods and Protocols*,
433    (Humana Press, 2017), pp. 397–422.

434 28. M. Huse, J. Kuriyan, The conformational plasticity of protein kinases. *Cell* **109**, 275–282
435    (2002).

436 29. V. Modi, R. L. Dunbrack, Defining a new nomenclature for the structures of active and
437    inactive kinases. *Proc. Natl. Acad. Sci.* **116**, 6818–6827 (2019).

438 30. M. Radli, S. G. D. Rüdiger, Dancing with the diva: Hsp90-client interactions. *J. Mol. Biol.*
439    **430**, 3029–3040 (2018).

440 31. H. Xu, Cochaperones enable Hsp70 to use ATP energy to stabilize native proteins out of

441   the folding equilibrium. *Sci. Rep.* **8**, 13213 (2018).

442   32.   J. Li, K. Richter, J. Reinstein, J. Buchner, Integration of the accelerator Aha1 in the Hsp90
443   co-chaperone cycle. *Nat. Struct. Mol. Biol.* **20**, 326–331 (2013).

444   33.   S. H. McLaughlin, H. W. Smith, S. E. Jackson, Stimulation of the weak ATPase activity
445   of human Hsp90 by a client protein. *J. Mol. Biol.* **315**, 787–798 (2002).

446   34.   T. O. Street, L. A. Lavery, D. A. Agard, Substrate binding drives large-scale
447   conformational changes in the Hsp90 molecular chaperone. *Mol. Cell* **42**, 96–105 (2011).

448   35.   O. R. Lorenz, L. Freiburger, D. A. Rutz, M. Krause, B. K. Zierer, S. Alvira, J. Cuéllar, J.
449   M. Valpuesta, T. Madl, M. Sattler, J. Buchner, Modulation of the Hsp90 chaperone cycle
450   by a stringent client protein. *Mol. Cell* **53**, 941–953 (2014).

451   36.   X. Wang, J. Venable, P. LaPointe, D. M. Hutt, A. V Koulov, J. Coppinger, C. Gurkan, W.
452   Kellner, J. Matteson, H. Plutner, J. R. Riordan, J. W. Kelly, J. R. III, W. E. Balch, Hsp90
453   cochaperone Aha1 downregulation rescues misfolding of CFTR in cystic fibrosis. *Cell*
454   **127**, 803–815 (2006).

455   37.   P. Goloubinoff, A. S. Sassi, B. Fauvet, A. Barducci, P. D. L. Rios, Chaperones convert the
456   energy from ATP into the nonequilibrium stabilization of native proteins. *Nat. Chem. Biol.*
457   **14**, 388–395 (2018).

458   38.   C. Queitsch, T. A. Sangster, S. Lindquist, Hsp90 as a capacitor of phenotypic variation.
459   *Nature* **417**, 618–624 (2002).

460   39.   W. Xu, E. Mimnaugh, M. F. N. Rosser, C. Nicchitta, M. Marcu, Y. Yarden, L. Neckers,
461   Sensitivity of mature ErbB2 to Geldanamycin is conferred by its kinase domain and is
462   mediated by the chaperone protein Hsp90. *J. Biol. Chem.* **276**, 3702–3708 (2001).

463   40.   M.-J. J. E. Bijlmakers, M. Marsh, Hsp90 is essential for the synthesis and subsequent
464   membrane association, but not the maintenance, of the Src-kinase p56 lck. *Mol. Biol. Cell*
465   **11**, 1585–1595 (2000).

466   41.   C. S. Gerbin, R. Landgraf, Geldanamycin selectively targets the nascent form of ERBB3
467   for degradation. *Cell Stress Chaperones* **15**, 529–544 (2010).

468   42.   A. Kamal, L. Thao, J. Sensintaffar, L. Zhang, M. F. Boehm, L. C. Fritz, F. J. Burrows, A
469   high-affinity conformation of Hsp90 confers tumour selectivity on Hsp90 inhibitors.
470   *Nature* **425**, 407–410 (2003).

471   43.   S. M. Roe, M. Ali, P. Meyer, C. K. Vaughan, B. Panaretou, P. W. Piper, C. Prodromou, L.
472   H. Pearl, The mechanism of Hsp90 regulation by the protein kinase-specific cochaperone
473   p50cdc37. *Cell* **116**, 87–98 (2004).

474   44.   A. M. Gaiser, A. Kretzschmar, K. Richter, Cdc37-Hsp90 complexes are responsive to
475   nucleotide-induced conformational changes and binding of further cofactors. *J. Biol.*
476   *Chem.* **285**, 40921–40932 (2010).

477   45.   B. Panaretou, G. Siligardi, P. Meyer, A. Maloney, J. K. Sullivan, S. Singh, S. H. Millson,
478   P. A. Clarke, S. Naaby-Hansen, R. Stein, R. Cramer, M. Mollapour, P. Workman, P. W.
479   Piper, L. H. Pearl, C. Prodromou, Activation of the ATPase activity of Hsp90 by the
480   stress-regulated cochaperone Aha1. *Mol. Cell* **10**, 1307–1318 (2002).

Figures

Figure 1. Structural basis for my model of non-equilibrium activation of client kinases by
Hsp90. The underlined texts below are PDB IDs and chain IDs. The Hsp90 homodimer goes
through four states in **A** to **D**. (**A**) $Hsp90_{open}$: nucleotide-free and open; 2IOQ. (**B**) $Hsp90_{open}^{ATP}$:
ATP-bound and open; 2IOQ but with the NTD of one chain replaced with the ATP-bound NTD
from the structure 2CG9. ATP-binding induces a structural change in the nucleotide lid in NTD.
(**C**) $Hsp90_{closed}^{ATP}$: ATP-bound and closed; 2CG9. The hole between the two MDs in the closed
Hsp90 dimer is gated by a flexible, often disordered src loop (the src loops from 2CG9, 5FWM,
2IOP, and 4IYN are superimposed). (**D**) $Hsp90_{compact}^{ADP}$: ADP-bound and compact; 2IOP. The
ADP-bound NTD may have a number of orientations with respect to MD: *e.g.*, the NTD of chain
A may move to the position occupied by the NTD of chain C in the crystal structure, resulting in
a compact conformation consistent with electron microscopy images (12) but clashing with the

bound kinase client. In B, C, and D, the nucleotides are shown in spheres. The kinase client
may transition through four states in **E** to **H**. (**E**) The inactive conformation: *i*; <u>1HCL:A</u>. (**F**)
The deactivating conformation: *d*; <u>5FWM:K</u>. (**G**) The hypothetical maturing conformation: *m*.
(**H**) The active conformation: *a*; <u>1FIN:A</u>. The deactivating structure is of the client kinase Cdk4
in the Hsp90-Cdc37-Cdk4 complex; the inactive and active structures are of Cdk2. (**I**) Complex
between Cdc37 and Cdk4 in the *d*-state (<u>5FWM:K+E</u>). (**J**) Steric clash prevents Cdc37 from
binding to Cdk4 in the *m*-state. Three residues (in spheres) in the DFG loop—in addition to
perhaps the β4-αC loop—would collide with Cdc37, whose clashing residues are highlighted in
red. (**K**) The ternary complex of Hsp90-Cdc37-Cdk4 (<u>5FWM</u>), where the client kinase is in the
d-state. (**L**) The hypothetical complex of the closed Hsp90 dimer with the client kinase in the *m*-
state.

Figure 2. The reactions in Hsp90-mediated non-equilibrium activation of v-Src and the reactive
fluxes at the steady state. Each circle represents a molecular species; the radius of the circle is
linear in the logarithm of its steady state concentration. The top four circles represent the free
client molecule in its four different conformations; the others represent complexes between the
client and Hsp90. The molecular species are divided by the vertical dashed lines according to
the client conformation, and into the colored horizontal bands according to the conformational
and nucleotide state of Hsp90. The ones that lie in the orange vertical band are ternary
complexes of Hsp90-client-Cdc37. The Hsp90 ATP cycles are highlighted in red. The
intermediate molecular species in the Cdc37-assisted activation pathway are labeled. The
directions of the steady state reactive fluxes are indicated by the green arrows. The reactions are
labeled with the reaction numbers in Table 1. The reaction conditions are [Hsp90] = [Cdc37] =
1.3 µM, [v-Src] = 0.32 µM, [ATP] = 20 µM, and [ADP] = [Pi] = 1 µM.

Figure 3. Activation of v-Src by Hsp90. (**A**) Normalized v-Src activity at the steady state at
different Hsp90 and Cdc37 concentrations. The experimental data (4) are shown as points, and
the results calculated from my model are shown as solid lines. (**B**) The time course of v-Src
activation by Hsp90 in the presence of Cdc37 and the concomitant ATP consumption.

Figure 4. Cdc37 expands Hsp90's clientele by recruiting deactivating kinases. (**A**) Kinase activation by Hsp90 at different ratios of its binding affinity for the *d*-state to that for the *m*-state of the kinase. Here, the K_D ratio is varied while keeping the conformational equilibrium constants of the apo and the Hsp90-bound client unchanged (reactions 6 and 14 in Table 1), which is satisfied by setting $K_{d,\text{closure}}/K_{m,\text{closure}} = K_{D,m\cdot\text{Hsp90}}/K_{D,d\cdot\text{Hsp90}}$, where $K_{s,\text{closure}}$ is the equilibrium constant of the reaction $s \cdot \text{Hsp90}_{\text{open}}^{\text{ATP}} \rightleftharpoons s \cdot \text{Hsp90}_{\text{closed}}^{\text{ATP}}$ for $s = d, m$. In the presence of Cdc37, kinases with low K_D ratios are activated by Hsp90. The reaction conditions differ from those in Figure 2 only in [Cdc37] and in that [ATP] = 250 µM. (**B**) The fraction of v-Src trapped in its *d*- and *m*-states by the closed Hsp90 dimer with the nonhydrolysable ATP analog AMPPNP at different Cdc37 concentrations. The reaction conditions are [Hsp90] = 15 µM, [v-Src] = 0.32 µM, and [AMPPNP] = 250 µM.

Figure 5. Changing the kinetic rates of the steps in the ATP cycle affects client activation by
Hsp90. Normalized activity (black line and left y-axis) and ATP cycle time (red dashed line and
right y axis) are calculated for different rates of (**A**) Hsp90 transition from the open to the closed
conformation, (**B**) ATP hydrolysis after Hsp90 adopts the closed conformation, and (**C**) ADP
release and the return of Hsp90 to the open conformation. The proportions of time spent in the
four Hsp90 states—open and nucleotide-free, open and ATP-bound, closed and ATP-bound, and
compact and ADP-bound—are shown as colored bands. Note the logarithmic scale of the times:
for the same apparent thickness, a higher band represents a much larger occupancy than a lower
band; in particular, the nucleotide-free open state has a higher occupancy than it seems. (**D**)
Modulation of client activation by the cochaperone Aha1. Aha1 can augment or diminish client
activation depending on whether the client itself accelerates (orange) or slows down (blue) the
rate that the Hsp90 dimer closes. Here [ATP] = 250 µM, [Cdc37] = 0 µM, [Hsp90] = 1.3 µM,
[client] = 0.32 µM, and [ADP] = [Pi] = 1 µM.