Phage infection mediates inhibition of bystander bacteria

Anushila Chatterjee[a]*, Julia L. E. Willett[b]*, Gary M. Dunny[b], Breck A. Duerkop[a,#]

[a]Department of Immunology and Microbiology, University of Colorado School of Medicine, Aurora,

CO, USA, 80045. [b]Department of Microbiology and Immunology, University of Minnesota Medical

School, Minneapolis, MN, USA, 55455.

[#]Correspondence: Breck A. Duerkop breck.duerkop@cuanschutz.edu

*A.C. and J.L.E.W. contributed equally to this work.

Running title: Phage induced T7SS promotes antibacterial antagonism

Key words: bacteriophages, *Enterococcus*, antibiotic resistance, phage–bacteria interactions,

bacterial secretion systems, type VII secretion, contact-dependent antagonism

Abstract

Bacteriophages (phages) are being considered as alternative therapeutics for the treatment of multidrug resistant bacterial infections. Considering phages have narrow host-ranges, it is generally accepted that therapeutic phages will have a marginal impact on non-target bacteria. We have discovered that lytic phage infection induces transcription of type VIIb secretion system (T7SS) genes in the pathobiont *Enterococcus faecalis*. Membrane damage during phage infection induces T7SS gene expression resulting in cell contact dependent antagonism of diverse Gram positive bystander bacteria. Deletion of *essB*, a T7SS structural component abrogates phage-mediated killing of bystanders. A predicted immunity gene confers protection against T7SS mediated inhibition and disruption of an upstream LXG toxin gene rescues growth of *E. faecalis* and *Staphylococcus aureus* bystanders. Phage induction of T7SS gene expression and bystander inhibition requires IreK, a serine/threonine kinase, and OG1RF_11099, a predicted transcriptional activator. Our findings highlight how phage infection of a target bacterium can affect non-target bystander bacteria and implies phage therapy could impose collateral damage to polymicrobial communities.

Introduction

Enterococci constitute a minor component of the healthy human microbiota (1). Enterococci, including *Enterococcus faecalis*, are also nosocomial pathogens that cause a variety of diseases, including sepsis, endocarditis, surgical-site, urinary tract and mixed bacterial infections (2, 3). Over recent decades, enterococci have acquired extensive antibiotic resistance traits, including resistance to "last-resort" antibiotics such as vancomycin, daptomycin, and linezolid (4-8). Following antibiotic therapy, multi-drug resistant (MDR) enterococci can outgrow to become a dominant member of the intestinal microbiota, resulting in intestinal barrier invasion and blood stream infection (7, 9). The ongoing evolution of MDR enterococci in healthcare settings (4-6, 10, 11) and their ability to transmit antibiotic resistance among diverse bacteria (9, 12-15), emphasize the immediate need for novel therapeutic approaches to control enterococcal infections.

Viruses that infect and kill bacteria (bacteriophages or phages) are receiving attention for their use as antibacterial agents (16). Recent studies have demonstrated the efficacy of anti-enterococcal phages in murine models of bacteremia (17-19) and the administration of phages to reduce *E. faecalis* burden in the intestine gives rise to phage resistant isolates that are resensitized to antibiotics (20). Considering phages are highly specific for their target bacterium, coupled with the self-limiting nature of their host-dependent replication, suggests that unlike antibiotics which have broad off-target antimicrobial activity, phages are likely to have nominal impact on bacteria outside of their intended target strain (21-23). However, our understanding of how phages interact with bacteria and the bacterial response to phage infection is limited.

While studying the transcriptional response of phage infected *E. faecalis* cells, we discovered that phage infection induced the expression of genes involved in the biosynthesis of a type VIIb secretion system (T7SS) (24). Firmicutes, including the enterococci, harbor diverse T7SS genes encoding transmembrane and cytoplasmic proteins involved in the secretion of protein substrates (25), and T7SSs promote antagonism of non-kin bacterial cells through contact-dependent secretion of antibacterial effectors and/or toxins (26, 27). The antibacterial activity of T7SSs from

Staphylococcus and *Streptococcus* are well characterized (25) but T7SS-mediated antibacterial

antagonism has not been described in *Enterococcus*. The environmental cues and regulatory

pathways that govern T7SS expression and activity are poorly understood, although recent

studies indicate that exposure to serum and membrane stresses triggered by pulmonary

surfactants, fatty acids and phage infection stimulate T7SS gene expression (24, 28-31). This

motivated us to determine if phage induced T7SS gene expression in *E. faecalis* results in the

inhibition of non-kin bacterial cells that are not phage targets (bystanders). We discovered that

phage infected *E. faecalis* produces potent T7SS antibacterial activity against bystander bacteria.

Expression of a T7SS antitoxin (immunity factor) gene in bystander cells and mutation of the LXG

domain containing gene located immediately upstream of the immunity factor confer protection

against phage mediated T7SS inhibition. Additionally, we discovered that membrane stress during

phage infection induces transcription of T7SS genes via a non-canonical IreK signaling pathway.

To our knowledge, the enterococcal T7SS is the only example of secretion system induction

during phage infection. These data shed light on how phage infection of a cognate bacterial host

can influence polymicrobial interactions and raises the possibility that phages may impose

unintended compositional shifts among bystander bacteria in the microbiota during phage

therapy.

Results

Phage mediated induction of *E. faecalis* T7SS leads to interspecies antagonism.

A hallmark feature of phage therapy is that because phages often have a narrow host range,

they will not influence the growth of non-susceptible bacteria occupying the same niche (22). We

discovered that infection of *E. faecalis* OG1RF by phage VPE25, induces the expression of T7SS

genes (24). The *E. faecalis* OG1RF T7SS locus is absent in the commonly studied vancomycin-

resistant strain V583, despite conservation of flanking genes (Fig. 1A) (32, 33). Homologs of the

E. faecalis T7SS gene *exsA* are found throughout three of the four *Enterococcus* species groups

(34), including *Enterococcus faecium*, suggesting a wide distribution of T7SS loci in enterococci

(Fig. S1). *In silico* analyses predict that the *E. faecalis* T7SS locus encodes multiple WXG100

family effectors and LXG family polymorphic toxins (27, 35). Hence, we hypothesized that

induction of T7SS genes during phage infection and consequently the heightened production of

T7SS substrates would indirectly influence the growth of non-kin phage-resistant bacterial cells.

To investigate if T7SS factors produced during phage infection of *E. faecalis* OG1RF interferes

with the growth of phage-resistant bystander bacteria, we generated a strain with an in-frame

deletion in the T7SS gene *essB*, encoding a transmembrane protein involved in the transport of

T7SS substrates (36). The *essB* mutant is equally susceptible to phage VPE25 infection

compared to wild type *E. faecalis* OG1RF (Fig. S2). We performed co-culture experiments where

phage susceptible wild type *E. faecalis* OG1RF or *ΔessB* were mixed with a phage resistant

bystander, a strain of *E. faecalis* V583 deficient in the production the VPE25 receptor (*Δpip$_{V583}$*)

(37), at a ratio of 1:1 in the absence and presence of phage VPE25 (multiplicity of infection [MOI]

= 0.01) (Fig. 1B). Since the *E. faecalis* V583 genome does not contain T7SS genes, this strain

should lack immunity to T7SS antagonism (33). The viability of *E. faecalis Δpip$_{V583}$*, was reduced

nearly 100-fold when co-cultured with *E. faecalis* OG1RF in the presence of phage VPE25 (Fig.

1C). However, growth inhibition of *E. faecalis Δpip$_{V583}$* was abrogated during co-culture with phage

infected *E. faecalis ΔessB*. Phage induced T7SS antagonism could be restored by

complementation (Fig. 1C), indicating that inhibition of phage resistant *E. faecalis Δpip$_{V583}$* is T7SS

dependent.

T7SS encoded antibacterial toxins secreted by Gram positive bacteria influence intra- and

interspecies antagonism (26, 27). While a nuclease toxin produced by *Staphylococcus aureus*

targets a closely related *S. aureus* strain (26), *Streptococcus intermedius* exhibits T7SS

dependent antagonism against a wide-array of Gram positive bacteria (27). To determine the

target range of *E. faecalis* OG1RF T7SS antibacterial activity, we measured the viability of a panel

of VPE25 resistant Gram positive and Gram negative bacteria in our co-culture assay (Fig. 1B).

Growth inhibition of the distantly related bacterial species *E. faecium* and Gram positive bacteria

of diverse genera, including *S. aureus* and *Listeria monocytogenes*, occurred following co-culture

with phage infected wild type *E. faecalis* OG1RF but not the Δ*essB* mutant (Fig. 1D). In contrast,

Gram positive streptococci were unaffected (Fig. 1D). Similarly, phage induced T7SS activity did

not inhibit any Gram negative bacteria tested (Fig. 1D). Collectively, these results show that phage

predation of *E. faecalis* promotes T7SS inhibition of diverse bystander bacteria.

Molecular basis of *E. faecalis* phage–triggered T7SS antagonism

Our data demonstrate that induction of *E. faecalis* OG1RF T7SS genes during phage infection

hinder the growth of select non-kin bacterial species. Antibacterial toxins deployed by Gram

negative bacteria via type VI secretion and Gram positive T7SS require physical contact between

cells to achieve antagonism (26, 27, 38, 39). Therefore, we investigated if growth inhibition of

bystander bacteria is contingent upon direct interaction with phage infected *E. faecalis* using a

trans-well assay (27). We added unfiltered supernatants from wild type *E. faecalis* OG1RF and

Δ*essB* mutant cultures grown for 24 hrs in the presence and absence of phage VPE25 (MOI =

0.01) to the top of a trans well and deposited phage resistant *E. faecalis* Δ*pip*$_{V583}$ in the bottom of

the trans well. The 0.4 μm membrane filter that separates the two wells is permeable to proteins

and solutes but prevents bacterial translocation. Supernatant from phage infected wild type *E.*

faecalis OG1RF did not inhibit *E. faecalis* Δ*pip*$_{V583}$ (Fig. 2A) indicating that T7SS mediated growth

interference relies on cell to cell contact. To exclude the possibility that T7SS substrates might

adhere to the 0.4 μm membrane filter in the trans-well assay, we administered unfiltered culture

supernatants directly to *E. faecalis* Δ*pip*$_{V583}$ cells (5x10^5 CFU/well) at a ratio of 1:10 and monitored

growth over a period of 10 hours. Growth kinetics of *E. faecalis* Δ*pip*$_{V583}$ remained similar

irrespective of the presence or absence of conditioned supernatant from wild type *E. faecalis*

OG1RF or Δ*essB* mutant cultures (Fig. 2B), further supporting the requirement of contact-

dependent engagement of phage mediated T7SS inhibition.

We discovered that *E. faecalis* OG1RF inhibits proliferation of non-kin bacterial cells through

increased expression of T7SS genes in response to phage infection, but the toxic effectors are

unknown. LXG domain containing toxins are widespread in bacteria with a diverse range of

predicted antibacterial activities (40, 41). The OG1RF T7SS locus encodes two LXG-domain

proteins, expressed from OG1RF_11109 and OG1RF_11121 (Fig. 3A). Polymorphic toxin

systems in Gram negative and Gram positive bacteria can encode "orphan" toxins in addition to

the primary secreted effectors (42). To identify putative orphan effectors, we aligned

OG1RF_11109 and OG1RF_11121 with downstream genes in the T7SS locus. OG1RF_11109

shared conserved sequence with OG1RF_11111 and OG1RF_11113, and OG1RF_11121 had

homology to OG1RF_11123 (Fig 3A). Homologs to these putative orphan toxins are found as full-

length LXG proteins in *E. faecalis* and *Listeria sp*. (Fig. S3), suggesting they are bonafide T7SS

effectors. Interestingly, we identified an additional LXG gene product, OG1RF_12414, in a distal

locus that is again notably absent from V583 (Fig. S4A and S4B). Although none of the five

putative toxins in the T7SS locus have recognizable C-terminal domains, OG1RF_12414 has

predicted structural homology to Tne2, a T6SS effector with NADase activity from *Pseudomonas*

protegens (Fig. S4C) (43). Additionally, we identified numerous C-terminal domains in LXG

proteins distributed throughout *Enterococcus* (Fig. S5). These include EndoU and Ntox44

nuclease domains (40, 44, 45), which have been characterized in effectors produced by other

polymorphic toxin systems.

Polymorphic toxins are genetically linked to cognate immunity proteins that neutralize

antagonistic activity and prevent self-intoxication (40, 45, 46). Each of the five putative toxins in

the OG1RF T7SS locus is encoded directly upstream of a small protein that could function in

immunity. Whitney *et al*. demonstrated that the cytoplasmic antagonistic activity of *S. intermedius*

LXG toxins TelA and TelB in *E. coli* can be rescued by co-expression of the corresponding

immunity factors (27). Therefore, we examined if OG1RF_11110, 11112, or 11122, confer

immunity to *E. faecalis* Δpip_{V583} during phage infection of *E. faecalis* OG1RF. Constitutive

expression of OG1RF_11122, and not OG1RF_11110 or 11112, partially neutralized phage

induced T7SS antagonism (Fig. 3B), confirming an essential role for the OG1RF_11122 gene

product in immunity, and suggesting that OG1RF_11121 is at least partly responsible for T7SS

mediated intra-species antagonism.

To determine the contribution of OG1RF_11121 on intra- and interbacterial antagonism during

phage infection, we measured the viability of phage resistant *E. faecalis* Δ*pip*$_{V583}$ and *S. aureus*

in co-culture with an *E. faecalis* OG1RF variant carrying a transposon insertion in OG1RF_11121

(OG1RF_11121-Tn). OG1RF_11121-Tn is equally susceptible to phage VPE25 infection

compared to wild type *E. faecalis* OG1RF (Fig. S2). Similar to the Δ*essB* mutant, phage infected

OG1RF_11121-Tn did not inhibit the growth of the bystander bacteria (Fig. 3C – D), implicating

that the OG1RF_11121 encoded LXG toxin drives *E. faecalis* T7SS mediated antagonism of

bystanders following phage infection.

Sub-lethal antibiotic stress promotes T7SS dependent antagonism

Considering two genetically distinct phages trigger the induction of T7SS genes in *E. faecalis*

(24), we reasoned that T7SS induction could be a result of phage mediated cellular damage and

not specifically directed by a phage encoded protein. Antibiotics elicit a range of damage induced

stress responses in bacteria (47-49), therefore, we investigated the effects of subinhibitory

concentrations of antibiotics on T7SS expression in *E. faecalis*.

To investigate the influence of sublethal antibiotic concentrations on *E. faecalis* OG1RF T7SS

transcription, we determined the minimum inhibitory concentrations (MIC) of ampicillin,

vancomycin, and daptomycin (Fig.S6A-S6C) and monitored T7SS gene expression in *E. faecalis*

OG1RF cells treated with a sub-lethal dose of antibiotic (50% of the MIC). We found that bacterial

T7SS genes were significantly upregulated in the presence of the cell membrane targeting

antibiotic, daptomycin, relative to the untreated control (Fig. 4A). In contrast, the cell wall

biosynthesis inhibitors ampicillin and vancomycin either did not induce or had a minor impact on

T7SS mRNA levels, respectively (Fig. 4A). Additionally, induction of T7SS transcription occurred

when bacteria are challenged with a sub-inhibitory concentration of the DNA targeting antibiotics

ciprofloxacin and mitomycin C (Fig. 4B, Fig. S6D–S6E). Collectively, these data show that T7SS

induction in *E. faecalis* occurs in response to cell envelope and DNA stress.

We next assessed the influence of daptomycin driven T7SS induction on inter-enterococal

antagonism. Since T7SS expression is less robust in the presence of daptomycin compared to

phage infection (24), a 10:1 ratio of daptomycin treated *E. faecalis* OG1RF was required for

growth inhibition of *E. faecalis* Δ*pip*$_{V583}$ during co-culture (Fig. 4C). Consistent with our previous

results, daptomycin induced T7SS inhibition of *E. faecalis* Δ*pip*$_{V583}$ was contact dependent (Fig.

4D). To facilitate T7SS mediated contact-dependent killing of the target strain during daptomycin

exposure, we performed the inhibition assay on nutrient agar plates. The sub-inhibitory

concentration of daptomycin (2.5 µg/ml) used in liquid culture was toxic to the cells on agar plates,

so we lowered the daptomycin concentration to 0.5 µg/ml to prevent drug toxicity in the agar-

based antagonism assay. Plating T7SS producing *E. faecalis* OG1RF cells and *E. faecalis*

Δ*pip*$_{V583}$ bystander cells at a ratio of 10:1 resulted in ~10–fold inhibition of bystander growth (Fig.

4E). These data show that in addition to phages, antibiotics can be sensed by *E. faecalis* thereby

inducing T7SS antagonism of non-kin bacterial cells. These data also suggest that the magnitude

of T7SS gene expression is directly related to the potency of T7SS inhibition.

IreK and OG1RF_11099 facilitate T7SS expression in phage infected *E. faecalis* OG1RF via

a non-canonical signaling pathway

Having established that phage and daptomycin mediated membrane damage stimulates

heightened *E. faecalis* OG1RF T7SS gene expression and antagonistic activity, we next sought

to identify the genetic determinants that sense this damage and promote T7SS transcription. Two-

component systems, LiaR/S and CroS/R, and the PASTA kinase family protein IreK are well-

characterized modulators of enterococcal cell envelope homeostasis and antimicrobial tolerance

(50-52). Aberrant cardiolipin microdomain remodeling in the bacterial cell membrane in the

absence of the LiaR response regulator results in daptomycin hypersensitivity and virulence

attenuation (53). CroS/R signaling and subsequent modulation of gene expression govern cell

wall integrity and promote resistance to cephalosporins, glycopeptides and beta—lactam

antibiotics (54-56). The *ireK* encoded transmembrane Ser/Thr kinase regulates cell wall

homeostasis, antimicrobial resistance, and contributes to bacterial fitness during long-term

colonization of the intestinal tract (51, 57, 58). Recently it has been shown that direct cross-talk

between IreK and the CroS/R system positively impacts enterococcal cephalosporin resistance

(59).

Wild type *E. faecalis* OG1RF, an *ireK* in-frame deletion mutant (51) and transposon (Tn)

insertion mutants of *liaR*, *liaS*, *croR*, and *croS* (60) all display similar growth kinetics in the

absence of phage VPE25 infection (Fig. S7A). Although *croR*-Tn and *croS*-Tn exhibit reductions

in the plaquing efficiency of VPE25 particles, none of these genetic elements of enterococcal cell

wall homeostasis and antibiotic resistance were required for VPE25 infection (Fig. S7B). We

queried the expression levels of T7SS genes in these isogenic mutants during phage VPE25

infection (MOI = 1). T7SS gene expression was not enhanced in the $\Delta ireK$ mutant during phage

infection (Fig. 5A), whereas *liaR*-Tn, *liaS*-Tn, *croR*-Tn, and *croS*-Tn produced heightened levels

of T7SS transcripts similar to the wild type *E. faecalis* OG1RF compared to the uninfected controls

(Fig. S8A-S8F). A sub-lethal concentration of the cephalosporin ceftriaxone did not induce T7SS

gene expression (Fig. S9), indicating that expression of T7SS genes following phage mediated

membrane damage signals through a pathway that is distinct from the IreK response to

cephalosporin stress. Additionally, the $\Delta ireK$ mutant phenocopies the $\Delta essB$ mutant strain in the

interbacterial antagonism co-culture assay, wherein the $\Delta ireK$ mutant is unable to mediate phage

induced T7SS dependent killing of the phage resistant *E. faecalis* Δpip_{V583} non-kin cells (Fig. 5B).

Collectively, these results indicate that IreK senses phage mediated membrane damage

promoting T7SS transcription independent of the CroS/R pathway.

OG1RF_11099, located immediately upstream of the T7SS cluster is predicted to encode a

GntR family transcriptional regulator, thus we sought to assess the contribution of *OG1RF_11099*

on T7SS transcription and functionality. OG1RF_11099-Tn is equally susceptible to phage VPE25

infection compared to wild type *E. faecalis* OG1RF (Fig. S2) In contrast to wild type *E. faecalis*

OG1RF, T7SS genes were not induced during phage predation in OG1RF_11099-Tn which

harbors a transposon insertion in this potential transcriptional regulatory gene (Fig. 5C).

Subsequently, we evaluated the influence of OG1RF_11099-dependent regulation on the activity

of enterococcal T7SS in intraspecies antagonism using our co-culture assays. Phage infected

OG1RF_11099-Tn behaved similar to the Δ*essB* mutant and was unable to inhibit the *E. faecalis*

Δ*pip*$_{V583}$ competitor strain during co-culture (Fig. 5D). Collectively, these results indicate that

OG1RF_11099 encodes a positive regulator of *E. faecalis* T7SS.

Discussion

Despite the fact that bacteria exist in complex microbial communities and engage in social

interactions (61, 62), phage predation studies have primarily been performed in monoculture (24,

63-65). Studies report phage-mediated effects on non-target bacteria linked to interbacterial

interactions and evolved phage tropism for non-cognate bacteria (66-68), whereas other studies

have identified minimal changes in microbiota diversity during phage therapy (66, 69).

Our results extend previous work that observed the induction of *E. faecalis* OG1RF T7SS

gene expression in response to phage infection (24). By using an *in vitro* antibacterial antagonism

assay, we discovered that phage predation of *E. faecalis* OG1RF has an inhibitory effect on non-

phage targeted bacterial species during co-culture. Our work shows that phage mediated

inhibition of Gram positive bystander bacteria relies on the expression and activity of T7SS genes.

This work emphasizes how phage infection of target bacteria can extend beyond intended phage

targets and inhibit other members of a microbial community. This discovery could have profound

ramifications on how microbial communities like the microbiota respond to phage therapy.

Our data suggest that membrane stress associated with phage infection or sub-lethal

daptomycin treatment stimulates T7SS mediated antibacterial antagonism of *E. faecalis* OG1RF.

Given that daptomycin is used to target vancomycin-resistant enterococcal infections, this finding

provides a model by which antibiotic-resistant enterococci may overgrow and dominate the

microbiota after antibiotic treatment. Although further investigation is required to understand how

T7SS induction might contribute to enterococcal fitness in polymicrobial environments, disruption

of T7SS loci in other bacteria compromises bacterial membrane integrity and attenuates virulence

(31, 70). It is possible that environmental conditions encountered in the intestinal tract, including

bile salts, antimicrobial proteins, and competition for nutrient resources could influence T7SS

activity in *E. faecalis* to facilitate niche establishment and/or persistence within a complex

microbial community. Indeed, *E. faecalis* T7SS mutants have a defect in their ability to colonize

the murine reproductive tract (manuscript submitted). Further, we discovered that transcriptional

activation of the T7SS during phage infection relies on IreK. Previously characterized IreK–

mediated stress response pathways, including cephalosporin stress or the CroS/R signaling, did

not contribute to T7SS expression. We hypothesize that IreK senses diverse environmental

stressors and coordinates distinct outputs in response to specific stimuli. Considering that IreK

signaling is important for *E. faecalis* intestinal colonization (58), it is possible that IreK–dependent

T7SS expression in response to intestinal cues modulate interbacterial interactions and

enterococcal persistence in the intestine. However, the molecular mechanism by which IreK

facilitates T7SS transcription remains unanswered.

Additionally, we discovered that *E. faecalis* OG1RF T7SS transcription is regulated by the

GntR domain containing transcriptional regulator encoded by *OG1RF_11099*, a gene found

immediately upstream of the T7SS cluster. Interestingly, *OG1RF_11099* is highly conserved

across enterococci, including *E. faecalis* V583 and other strains that lack T7SS. The presence of

a conserved transcriptional regulator in the absence of its target genetic region supports the idea

that certain strains of enterococci have undergone genome reduction as an evolutionary strategy

to adapt to unique host and non-host environments. It is possible that in *E. faecalis* V583, the

OG1RF_11099 homolog (EF1328) has been retained to regulate other genes within the regulon

that are less dispensable than T7SS.

Antibacterial properties of T7SS substrates have been demonstrated (26, 27). Here we

provide evidence that the expression of an immunity gene *OG1RF_11122* in T7SS targeted *E.*

faecalis cells confers protection from inhibition by the upstream LXG toxin encoded by

OG1RF_11121. Aside from its LXG domain, OG1RF_11121 does not harbor any other

recognizable protein domains, hence the mechanism underlying its toxicity is unclear. Whitney *et*

al. demonstrated that LXG toxin antagonism is contact–dependent, having minimal to no impact

on target cells in liquid media (27). Although we found that physical engagement is crucial for *E.*

faecalis T7SS mediated antagonism, we observed a significant reduction in target cell growth in

liquid media both during phage and daptomycin treatment of T7SS proficient *E. faecalis*.

Together, these data suggest that heightened T7SS transcription upon phage exposure

compared to daptomycin exposure may account for robust *E. faecalis* OG1RF inhibition of other

bacteria in liquid culture.

Enterococci occupy polymicrobial infections often interacting with other bacteria (71-74).

Although commensal *E. faecalis* antagonize virulent *S. aureus* through the production of

superoxide (75), the two species also exhibit growth synergy via exchange of critical nutrients

(76). Here, we show that phage treatment of *E. faecalis* OG1RF can indirectly impact the growth

of neighboring phage-resistant bacteria, including *S. aureus*, in a T7SS–dependent manner,

suggesting that phage therapy directed against enterococci driving T7SS activity could be useful

for the treatment of polymicrobial infections. However, the counter argument is that phage therapy

directed against enterococci could push a bacterial community toward dysbiosis, as phage

induced T7SS activity could directly inhibit beneficial bystander bacteria. This raises questions

about the consequences of phage mediated off-target effects on bacteria. Could phage induced

T7SS activity be used to reduce phage expansion into other closely related strains as a means to

dilute phages out of a population, or is it simply that phage induction of the T7SS serves as a

mechanism that benefits a select few within a population to aid in their reoccupation of a niche

upon overcoming phage infection? Future studies aimed at exploring enterococcal T7SS

antagonism in polymicrobial communities should help elucidate the impact of phages on microbial

community composition.

Materials and Methods

Bacteria and bacteriophage. Bacteria and phages used in this study are listed in Table S1.

Bacteria were grown with aeration in Todd-Hewitt broth (THB) or on THB agar supplemented with

10mM $MgSO_4$ at 37°C. The following antibiotic concentrations were added to media for the

selection of specific bacterial strains or species: *E. faecalis* OG1RF (25 µg/ml fusidic acid, 50

µg/ml rifampin), *E. faecalis* V583 Δpip_{V583} (100 µg/ml gentamycin), *S. aureus* AH2146 LAC

Φ11:LL29 (1 µg/ml tetracyclin), *L. monocytogenes* 10403S (100 µg/ml streptomycin), *V. cholerae*

C6706 int I4::TnFL63 and *S. enterica* serovar Typhimurium 140285 put::Kan (50 µg/ml

kanamycin). *S. agalactiae* COH1 was distinguished from *E. faecalis* on Chrome indicator Agar

(CHROMagar StrepB SB282). We were unable to differentially select *E. coli, S. pyogenes* and *S.*

mitis from *E. faecalis* based on antibiotic sensitivity. Therefore, colony counts of these bacteria in

co-culture experiments were acquired by subtracting the *E. faecalis* colony numbers on selective

media from total number of colonies on non-selective media. Strains harboring pLZ12A and its

derivatives were grown in the presence of 20 µg/ml chloramphenicol.

Bioinformatic analyses. Genome sequences of *E. faecalis* V583 (NC_004668.1) and OG1RF

(NC_017316.1) were obtained from NCBI. Alignments were generated and visualized using

EasyFig (77). OG1RF protein domains were identified using KEGG (78) and ExPASy PROSITE

(79). Structure modeling of OG1RF_12414 was done with Phyre2 (80). The EsxA phylogenetic

tree was constructed in MEGA version X (81) using non-redundant protein sequences obtained

from NCBI BLAST (82) with OG1RF_11100 as input and was edited using the Interactive Tree Of

Life browser (83). OG1RF_11109 was used as an input for the NCBI Conserved Domain

Architecture Retrieval Tool (84) to identify protein domains that co-occur with LXG domains in

Enterococcus.

Antibiotic sensitivity profiles. Antibiotic susceptibility profiles for ampicillin, vancomycin, and

daptomycin were determined using a broth microdilution assay. Overnight (O/N) *E. faecalis*

OG1RF cultures were diluted to 5×10^6 CFU/ml and 100 µl was added to each well of a 96-well

plate to give a final cell density of 5×10^5 CFU/ml. Antibiotic stocks were added to the first column

of each row, mixed thoroughly, and were serially diluted 2-fold across the rows. The last column

was used as a no drug control. Cultures containing daptomycin were supplemented with 50 µg/ml

$CaCl_2$. Bacterial growth was monitored by measuring absorbance (OD_{600}) using a Synergy H1

microplate reader set to 37°C with continuous shaking O/N. Growth curves are presented as the

average of three biological replicates. A concentration of antibiotic just below the drug amount

that inhibits bacterial growth was deemed sub-lethal and used to examine T7SS genes

expression.

Co-culture bacterial antagonism assays. For inter- and intraspecies antagonism assays in

liquid media, O/N cultures of different bacteria were diluted in THB containing 10mM $MgSO_4$ to

an OD_{600} of 0.2 and mixed together in a 1:1 or 10:1 ratio. The mixed cell suspensions were either

left untreated or treated with phage VPE25 (MOI 0.01) / daptomycin (2.5 µg/ml) and grown at

37°C with aeration. For antagonism experiments on agar plates, O/N cultures of different strains

were diluted to an OD_{600} of 0.2 and mixed together in a 1:1 or 10:1 ratio. A total of 10^7 cells from

mixed culture suspension was added to 5 ml THB + 0.35% agar at 55°C and were poured over

the surface of a THB agar plate in the absence or presence of daptomycin (0.5 µg/ml). The plates

were incubated at 37°C under static conditions for 24 hours. Cells were harvested by scraping off

the top agar, resuspending in 5 ml of PBS, and the cfus were obtained by plating serially diluted

cell suspension on appropriate selective or differentiating agar plates. Relative viability was

calculated from the ratio of target strain cfu in the treated versus the untreated co-culture. The

assays were performed in biological triplicates.

RNA extraction and quantitative PCR. RNA was extracted from phage treated or untreated *E.*

faecalis OG1RF cells by using an RNeasy Mini Kit (Qiagen) with the following published

modifications (24). cDNA was generated from 1 µg of RNA using qScript cDNA SuperMix

(QuantaBio) and transcript levels were analyzed by qPCR using PowerUp™ SYBR Green Master

Mix (Applied Biosystems). Transcript abundances were normalized to the 16S rRNA gene

transcripts and fold–change was calculated by comparing to untreated controls. All data are

represented as the average of three biological replicates.

Bacterial growth curves. 25 ml of 10mM $MgSO_4$ supplemented THB was inoculated with O/N

cultures of *E. faecalis* diluted to an OD_{600} of 0.025 and distributed to a 96-well plate in 0.1 ml

volumes. Cultures were incubated at 37° C with aeration. OD_{600} was measured periodically for 18

hours in a Synergy H1 microplate reader.

Efficiency of plating (EOP) assays. To investigate if phage VPE25 can infect and lyse *E.*

faecalis mutants and various other bacterial species, 10^7 PFU/ml of phage was serially diluted

and the phage was titered on each strain using a THB agar overlay plaque assay. EOP is

expressed as the percentage of phage titer from each strain relative to the wild type *E. faecalis*

OG1RF control. Data are presented as the average of three biological replicates.

Construction of *E. faecalis* mutants and complementation. Isolation of *E. faecalis* genomic

DNA was performed using a ZymoBIOMICS™ DNA Miniprep Kit (Zymo Research). All PCR

reactions used for cloning were performed with high fidelity KOD Hot Start DNA Polymerase (EMD

Millipore). *E. faecalis* Δ*essB* was generated by allelic replacement by cloning an in frame *essB*

deletion product into pLT06 using Gibson Assembly® Master Mix (New England Biolabs),

integrating this construct into the chromosome, and resolving the deletion mutant by homologous

recombination (85-87). For ectopic expression of putative immunity proteins, coding regions of

OG1RF_11110, OG1RF_11112, and OG1RF_11122 were cloned downstream of the *bacA*

promoter (P_{bacA}) by restriction digestion and ligation into the shuttle vector pLZ12A (20). Primer

sequences and restriction enzymes used for cloning are listed in Table S1. Plasmids were

introduced into electrocompetent *E. faecalis* cells as previously described (20).

Statistical analysis. Statistical tests were performed using GraphPad – Prism version 8.2.1. For

qPCR and bacterial competition assays, unpaired Student's t-tests were used. *P* values are

indicated in the figure legends.

References

1.   Lebreton F, Willems RJL, & Gilmore MS (2014) *Enterococcus* diversity, origins in nature,

and gut colonization. *Enterococci: from commensals to leading causes of drug resistant*

infection, eds Gilmore MS, Clewell DB, Ike Y, & Shankar NBoston).

2.   Onderdonk AB, Bartlett JG, Louie T, Sullivan-Seigler N, & Gorbach SL (1976) Microbial

synergy in experimental intra-abdominal abscess. *Infect Immun* 13(1):22-26.

3.   Weiner-Lastinger LM*, et al.* (2020) Antimicrobial-resistant pathogens associated with adult

healthcare-associated infections: Summary of data reported to the National Healthcare

Safety Network, 2015-2017. *Infect Control Hosp Epidemiol* 41(1):1-18.

4.   Arias CA*, et al.* (2011) Genetic basis for *in vivo* daptomycin resistance in enterococci. *N*
Engl J Med 365(10):892-900.

5.   Liu Y*, et al.* (2012) First report of the multidrug resistance gene *cfr* in *Enterococcus faecalis*
of animal origin. *Antimicrob Agents Chemother* 56(3):1650-1654.

6.   Patel SN*, et al.* (2013) Linezolid resistance in *Enterococcus faecium* isolated in Ontario,
Canada. *Diagn Microbiol Infect Dis* 77(4):350-353.

7.   Ubeda C*, et al.* (2010) Vancomycin-resistant *Enterococcus* domination of intestinal
microbiota is enabled by antibiotic treatment in mice and precedes bloodstream invasion
in humans. *J Clin Invest* 120(12):4332-4341.

8.   Zirakzadeh A & Patel R (2006) Vancomycin-resistant enterococci: colonization, infection,
detection, and treatment. *Mayo Clin Proc* 81(4):529-536.

9.   Taur Y*, et al.* (2012) Intestinal domination and the risk of bacteremia in patients
undergoing allogeneic hematopoietic stem cell transplantation. *Clin Infect Dis* 55(7):905-
914.

10.   Munoz-Price LS, Lolans K, & Quinn JP (2005) Emergence of resistance to daptomycin
during treatment of vancomycin-resistant *Enterococcus faecalis* infection. *Clin Infect Dis*
41(4):565-566.

11.   Palmer KL, Daniel A, Hardy C, Silverman J, & Gilmore MS (2011) Genetic basis for
daptomycin resistance in enterococci. *Antimicrob Agents Chemother* 55(7):3345-3356.

12.   Jasni AS, Mullany P, Hussain H, & Roberts AP (2010) Demonstration of conjugative
transposon (Tn5397)-mediated horizontal gene transfer between *Clostridium difficile* and
Enterococcus faecalis. *Antimicrob Agents Chemother* 54(11):4924-4926.

13.   Palmer KL, Kos VN, & Gilmore MS (2010) Horizontal gene transfer and the genomics of
463   enterococcal antibiotic resistance. *Curr Opin Microbiol* 13(5):632-639.

464   14.   Chang S*, et al.* (2003) Infection with vancomycin-resistant *Staphylococcus aureus*
465   containing the *vanA* resistance gene. *N Engl J Med* 348(14):1342-1347.

466   15.   Kurenbach B*, et al.* (2003) Intergeneric transfer of the *Enterococcus faecalis* plasmid
467   pIP501 to *Escherichia coli* and *Streptomyces lividans* and sequence analysis of its *tra*
468   region. *Plasmid* 50(1):86-93.

469   16.   Kortright KE, Chan BK, Koff JL, & Turner PE (2019) Phage therapy: a renewed approach
470   to combat antibiotic-resistant bacteria. *Cell Host Microbe* 25(2):219-232.

471   17.   Cheng M*, et al.* (2017) The vacteriophage EF-P29 efficiently protects against lethal
472   vancomycin-resistant *Enterococcus faecalis* and alleviates gut microbiota imbalance in a
473   murine bacteremia model. *Front Microbiol* 8:837.

474   18.   Gelman D*, et al.* (2018) Combined bacteriophages and antibiotics as an efficient therapy
475   against VRE *Enterococcus faecalis* in a mouse model. *Res Microbiol* 169(9):531-539.

476   19.   Uchiyama J, Rashel M, Takemura I, Wakiguchi H, & Matsuzaki S (2008) *In silico* and *in*
477   *vivo* evaluation of bacteriophage phiEF24C, a candidate for treatment of *Enterococcus*
478   *faecalis* infections. *Appl Environ Microbiol* 74(13):4149-4163.

479   20.   Chatterjee A*, et al.* (2019) Bacteriophage resistance alters antibiotic-mediated intestinal
480   expansion of enterococci. *Infect Immun* 87(6):e00085-00019.

481   21.   Lin DM, Koskella B, & Lin HC (2017) Phage therapy: an alternative to antibiotics in the
482   age of multi-drug resistance. *World J Gastrointest Pharmacol Ther* 8(3):162-173.

22.   Loc-Carrillo C & Abedon ST (2011) Pros and cons of phage therapy. *Bacteriophage*
1(2):111-114.

23.   Wu S, Zachary E, Wells K, & Loc-Carrillo C (2013) Phage Therapy: future inquiries.
Postdoc J 1(6):24-35.

24.   Chatterjee A*, et al.* (2020) Parallel genomics uncover novel enterococcal-bacteriophage
interactions. *mBio* 11(2).

25.   Unnikrishnan M, Constantinidou C, Palmer T, & Pallen MJ (2017) The enigmatic esx
proteins: looking beyond mycobacteria. *Trends Microbiol* 25(3):192-204.

26.   Cao Z, Casabona MG, Kneuper H, Chalmers JD, & Palmer T (2016) The type VII secretion
system of *Staphylococcus aureus* secretes a nuclease toxin that targets competitor
bacteria. *Nat Microbiol* 2:16183.

27.   Whitney JC*, et al.* (2017) A broadly distributed toxin family mediates contact-dependent
antagonism between Gram-positive bacteria. *Elife* 6.

28.   Burts ML, DeDent AC, & Missiakas DM (2008) EsaC substrate for the ESAT-6 secretion
pathway and its role in persistent infections of *Staphylococcus aureus*. *Mol Microbiol*
69(3):736-746.

29.   Ishii K*, et al.* (2014) Induction of virulence gene expression in *Staphylococcus aureus* by
pulmonary surfactant. *Infect Immun* 82(4):1500-1510.

30.   Lopez MS*, et al.* (2017) Host-derived fatty acids activate type VII secretion in
Staphylococcus aureus. *Proc Natl Acad Sci U S A* 114(42):11223-11228.

31.   Tchoupa AK*, et al.* (2020) The type VII secretion system protects *Staphylococcus aureus*

against antimicrobial host fatty acids. *bioRxiv*:572172.

32.   Bourgogne A*, et al.* (2008) Large scale variation in *Enterococcus faecalis* illustrated by the

genome analysis of strain OG1RF. *Genome Biol* 9(7):R110.

33.   Paulsen IT*, et al.* (2003) Role of mobile DNA in the evolution of vancomycin-resistant

Enterococcus faecalis. *Science* 299(5615):2071-2074.

34.   Lebreton F*, et al.* (2017) Tracing the enterococci from paleozoic origins to the hospital.

Cell 169(5):849-861 e813.

35.   Das C, Ghosh TS, & Mande SS (2016) *In silico* dissection of type VII secretion system

components across bacteria: new directions towards functional characterization. *J Biosci*

41(1):133-143.

36.   Burts ML, Williams WA, DeBord K, & Missiakas DM (2005) EsxA and EsxB are secreted

by an ESAT-6-like system that is required for the pathogenesis of *Staphylococcus aureus*

infections. *Proc Natl Acad Sci U S A* 102(4):1169-1174.

37.   Duerkop BA, Huo W, Bhardwaj P, Palmer KL, & Hooper LV (2016) Molecular basis for

lytic bacteriophage resistance in enterococci. *MBio* 7(4).

38.   Garcia EC (2018) Contact-dependent interbacterial toxins deliver a message. *Curr Opin*

Microbiol 42:40-46.

39.   Garcia-Bayona L & Comstock LE (2018) Bacterial antagonism in host-associated

microbial communities. *Science* 361(6408).

40. Zhang D, de Souza RF, Anantharaman V, Iyer LM, & Aravind L (2012) Polymorphic toxin

systems: comprehensive characterization of trafficking modes, processing, mechanisms

of action, immunity and ecology using comparative genomics. *Biol Direct* 7:18.

41. Klein TA, Pazos M, Surette MG, Vollmer W, & Whitney JC (2018) Molecular basis for

immunity protein recognition of a type VII secretion system exported antibacterial toxin. *J

Mol Biol* 430(21):4344-4358.

42. Jamet A & Nassif X (2015) New players in the toxin field: polymorphic toxin systems in

bacteria. *mBio* 6(3):e00285-00215.

43. Tang JY, Bullen NP, Ahmad S, & Whitney JC (2018) Diverse NADase effector families

mediate interbacterial antagonism via the type VI secretion system. *J Biol Chem*

293(5):1504-1514.

44. Michalska K*, et al.* (2018) Functional plasticity of antibacterial EndoU toxins. *Mol Microbiol*

109(4):509-527.

45. Zhang D, Iyer LM, & Aravind L (2011) A novel immunity system for bacterial nucleic acid

degrading toxins and its recruitment in various eukaryotic and DNA viral systems. *Nucleic

Acids Res* 39(11):4532-4552.

46. Klein TA, Ahmad S, & Whitney JC (2020) Contact-dependent interbacterial antagonism

mediated by protein secretion machines. *Trends Microbiol* 28(5):387-400.

47. Andersson DI & Hughes D (2014) Microbiological effects of sublethal levels of antibiotics.

Nat Rev Microbiol 12(7):465-478.

48. Bernier SP & Surette MG (2013) Concentration-dependent activity of antibiotics in natural

environments. *Front Microbiol* 4:20.

49. Yim G, Wang HH, & Davies J (2007) Antibiotics as signalling molecules. *Philos Trans R*
Soc Lond B Biol Sci 362(1483):1195-1200.

50. Kellogg SL & Kristich CJ (2016) Functional dissection of the CroRS two-component
system required for resistance to cell wall stressors in *Enterococcus faecalis*. *J Bacteriol*
198(8):1326-1336.

51. Kristich CJ, Wells CL, & Dunny GM (2007) A eukaryotic-type Ser/Thr kinase in
Enterococcus faecalis mediates antimicrobial resistance and intestinal persistence. *Proc*
Natl Acad Sci U S A 104(9):3508-3513.

52. Tran TT*, et al.* (2013) Daptomycin-resistant *Enterococcus faecalis* diverts the antibiotic
molecule from the division septum and remodels cell membrane phospholipids. *mBio* 4(4).

53. Reyes J*, et al.* (2015) A *liaR* deletion restores susceptibility to daptomycin and
antimicrobial peptides in multidrug-resistant *Enterococcus faecalis*. *J Infect Dis*
211(8):1317-1325.

54. Comenge Y*, et al.* (2003) The CroRS two-component regulatory system is required for
intrinsic beta-lactam resistance in *Enterococcus faecalis*. *J Bacteriol* 185(24):7184-7192.

55. Hancock L & Perego M (2002) Two-component signal transduction in *Enterococcus*
faecalis. *J Bacteriol* 184(21):5819-5825.

56. Hancock LE & Perego M (2004) Systematic inactivation and phenotypic characterization
of two-component signal transduction systems of *Enterococcus faecalis* V583. *J Bacteriol*
186(23):7951-7958.

57. Kristich CJ, Little JL, Hall CL, & Hoff JS (2011) Reciprocal regulation of cephalosporin
resistance in *Enterococcus faecalis*. *mBio* 2(6):e00199-00111.

58. Banla IL*, et al.* (2018) Modulators of *Enterococcus faecalis* cell envelope integrity and
antimicrobial resistance influence stable colonization of the mammalian gastrointestinal
tract. *Infect Immun* 86(1).

59. Kellogg SL & Kristich CJ (2018) Convergence of PASTA kinase and two-component
signaling in response to cell wall stress in *Enterococcus faecalis*. *J Bacteriol* 200(12).

60. Dale JL*, et al.* (2018) Comprehensive functional analysis of the *Enterococcus faecalis* core
genome using an ordered, sequence-defined collection of insertional mutations in strain
OG1RF. *mSystems* 3(5).

61. Madsen JS, Sorensen SJ, & Burmolle M (2018) Bacterial social interactions and the
emergence of community-intrinsic properties. *Curr Opin Microbiol* 42:104-109.

62. Xavier JB & Foster KR (2007) Cooperation and conflict in microbial biofilms. *Proc Natl*
Acad Sci U S A 104(3):876-881.

63. Leskinen K, Blasdel BG, Lavigne R, & Skurnik M (2016) RNA-sequencing reveals the
progression of phage-host interactions between phiR1-37 and *Yersinia enterocolitica*.
Viruses 8(4):111.

64. Mojardin L & Salas M (2016) Global transcriptional analysis of virus-host interactions
between phage Φphi29 and *Bacillus subtilis*. *J Virol* 90(20):9293-9304.

65. Sacher JC*, et al.* (2018) Transcriptomic analysis of the *Campylobacter jejuni* response to
T4-like phage NCTC 12673 infection. *Viruses* 10(6).

66. De Sordi L, Khanna V, & Debarbieux L (2017) The gut microbiota facilitates drifts in the
genetic diversity and infectivity of bacterial viruses. *Cell Host Microbe* 22(6):801-808 e803.

67.   Hsu BB*, et al.* (2019) Dynamic modulation of the gut microbiota and metabolome by

589   bacteriophages in a mouse model. *Cell Host Microbe* 25(6):803-814 e805.

590   68.   Reyes A, Wu M, McNulty NP, Rohwer FL, & Gordon JI (2013) Gnotobiotic mouse model

591   of phage-bacterial host dynamics in the human gut. *Proc Natl Acad Sci U S A*

592   110(50):20236-20241.

593   69.   Dissanayake U, Ukhanova M, Moye ZD, Sulakvelidze A, & Mai V (2019) Bacteriophages

594   reduce pathogenic *Escherichia coli* counts in mice without distorting gut microbiota. *Front*

595   *Microbiol* 10:1984.

596   70.   Bottai D*, et al.* (2012) Disruption of the ESX-5 system of *Mycobacterium tuberculosis*

597   causes loss of PPE protein secretion, reduction of cell wall integrity and strong attenuation.

598   *Mol Microbiol* 83(6):1195-1209.

599   71.   Flores-Mireles AL, Walker JN, Caparon M, & Hultgren SJ (2015) Urinary tract infections:

600   epidemiology, mechanisms of infection and treatment options. *Nat Rev Microbiol*

601   13(5):269-284.

602   72.   Giacometti A*, et al.* (2000) Epidemiology and microbiology of surgical wound infections. *J*

603   *Clin Microbiol* 38(2):918-922.

604   73.   Keogh D*, et al.* (2016) Enterococcal metabolite cues facilitate interspecies niche

605   modulation and polymicrobial infection. *Cell Host Microbe* 20(4):493-503.

606   74.   Goh HMS, Yong MHA, Chong KKL, & Kline KA (2017) Model systems for the study of

607   enterococcal colonization and infection. *Virulence* 8(8):1525-1562.

608   75.   King KC*, et al.* (2016) Rapid evolution of microbe-mediated protection against pathogens

609   in a worm host. *ISME J* 10(8):1915-1924.

76.   Hammer ND*, et al.* (2014) Inter- and intraspecies metabolite exchange promotes virulence

of antibiotic-resistant *Staphylococcus aureus*. *Cell Host Microbe* 16(4):531-537.

77.   Sullivan MJ, Petty NK, & Beatson SA (2011) Easyfig: a genome comparison visualizer.

Bioinformatics 27(7):1009-1010.

78.   Kanehisa M & Goto S (2000) KEGG: kyoto encyclopedia of genes and genomes. *Nucleic*

Acids Res 28(1):27-30.

79.   Sigrist CJ*, et al.* (2013) New and continuing developments at PROSITE. *Nucleic Acids*

Res 41(Database issue):D344-347.

80.   Kelley LA, Mezulis S, Yates CM, Wass MN, & Sternberg MJ (2015) The Phyre2 web portal

for protein modeling, prediction and analysis. *Nat Protoc* 10(6):845-858.

81.   Kumar S, Stecher G, Li M, Knyaz C, & Tamura K (2018) MEGA X: Molecular Evolutionary

Genetics Analysis across Computing Platforms. *Mol Biol Evol* 35(6):1547-1549.

82.   Coordinators NR (2018) Database resources of the National Center for Biotechnology

Information. *Nucleic Acids Res* 46(D1):D8-D13.

83.   Letunic I & Bork P (2019) Interactive Tree Of Life (iTOL) v4: recent updates and new

developments. *Nucleic Acids Res* 47(W1):W256-W259.

84.   Geer LY, Domrachev M, Lipman DJ, & Bryant SH (2002) CDART: protein homology by

domain architecture. *Genome Res* 12(10):1619-1623.

85.   Barnes WM (1994) PCR amplification of up to 35-kb DNA with high fidelity and high yield

from lambda bacteriophage templates. *Proc Natl Acad Sci U S A* 91(6):2216-2220.

86.   Gibson DG*, et al.* (2009) Enzymatic assembly of DNA molecules up to several hundred

631   kilobases. *Nat Methods* 6(5):343-345.

632   87.   Thurlow LR, Thomas VC, & Hancock LE (2009) Capsular polysaccharide production in

633   *Enterococcus faecalis* and contribution of CpsF to capsule serospecificity. *J Bacteriol*

634   191(20):6203-6210.

Acknowledgments

637   This work was supported by National Institutes of Health grants R01AI141479 (B.A.D.) and

638   R01AI122742 (G.M.D.). J.L.E.W. was supported by American Heart Association Grant

639   19POST34450124 / Julia Willett / 2018. We would like to thank Andrés Vázquez-Torres, Laurel

640   Lenz, Alex Horswill, Stefan Pukatzki, Kelly Doran, and their lab members for sharing bacterial

641   strains used in this study.

Author Contributions

644   A.C., J.L.E.W., G.M.D. and B.A.D. designed the study. A.C. and J.L.E.W. perfromed experiemtns

645   and bioinformatic analyses. A.C., J.L.E.W., G.M.D. and B.A.D. analyzed data. A.C., J.L.E.W. and

646   B.A.D. wrote the paper with input from G.M.D.

Corresponding author

649   Address coorespondence to Breck A. Duerkop breck.duerkop@cuanschutz.edu

Competing Interests

652   There are no competing interests to report for this work.

 Figure Legends

Figure 1. Phage mediated inhibition of bystander bacteria is dependent on enterococcal

T7SS. (A) Diagram showing the location of T7SS genes in *E. faecalis* OG1RF (NC_017316.1)

compared to *E. faecalis* V583 (NC_004668.1). Sequences were obtained from NCBI, and

homology comparisons were rendered in EasyFig. Nucleotide alignments generated by Clustal

Omega are enlarged for clarity (dashed lines). Stop codons of genes EF1328/OG1RF_11099 and

EF1337/OG1RF_11127 are boxed. **(B)** Schematic representation of the co-culture assay used to

assess the viability of bystander bacteria during phage induced T7SS activity of wild type *E.*

faecalis OG1RF and Δ*essB*. Relative viability of bystander strains is calculated by measuring the

ratio of bystander cfus in the phage infected culture compared to the bystander cfus from an

uninfected control culture. **(C)** The relative abundance of viable bystander bacterium *E. faecalis*

Δ*pip*$_{V583}$. Complementation of the *E. faecalis* Δ*essB* mutant restores T7SS dependent bystander

inhibition**. (D)** T7SS inhibition of other bacterial species in the presence and absence of phage

infected *E. faecalis* OG1RF or Δ*essB*. Data represent three biological replicates. *P < 0.00001 by

unpaired Student's t-test.

Figure 2. Phage induced T7SS inhibitory activity of *E. faecalis* is contact dependent. Intra-

species competition experiment performed in the presence of unfiltered supernatant from phage

treated and untreated *E. faecalis* wild type OG1RF or Δ*essB* added **(A)** to the top of a well

separated by a 0.4 μm membrane from the bottom well containing *E. faecalis* Δ*pip*$_{V583}$ culture or

(B) directly into *E. faecalis* Δ*pip*$_{V583}$ culture in microtiter plate wells.

Figure 3. Identification of *E. faecalis* T7SS toxin and immunity proteins that dictate

bystander growth inhibition bacteria. (A) LXG domains in OG1RF_11109 and OG1RF_11121

were identified using KEGG and ExPASy PROSITE. Other putative toxic effectors in OG1RF

were identified by homology to OG1RF_11109 or OG1RF_11121. Homology diagrams were

rendered in EasyFig. **(B)** *E. faecalis* OG1RF T7SS mediated growth inhibition of phage resistant

E. faecalis Δ*pip*$_{V583}$ during infection is alleviated by expressing OG1RF_11122 in *E. faecalis*

Δ*pip*$_{V583}$ but not in the presence of empty vector, or expressing OG1RF_11110, or OG1RF_11112.

E, represents empty vector. C, is expression vector. Disruption of OG1RF_11121 by a transposon

insertion rescues growth of phage resistant **(C)** *E. faecalis* Δ*pip*$_{V583}$ and **(D)** *S. aureus* strains

during co-culture. Data represent three biological replicates. *$P < 0.0001$ by unpaired Student's t-

test.

Figure 4. Sub-lethal antibiotic treatment enhances T7SS gene expression leading to

inhibition of bystander bacteria. Altered expression of T7SS genes upon exposure to sub-

inhibitory concentrations of **(A)** ampicillin (0.19 µg/ml), vancomycin (0.78 µg/ml) or daptomycin

(6.25 µg/ml) and **(B)** ciprofloxacin (2 µg/ml) or mitomycin C (4 µg/ml) for 40 minutes relative to

the untreated control. **(C-E)** Contact–dependent T7SS mediated inhibition of bystander bacteria

in the presence of daptomycin. Relative viability of *E. faecalis* Δ*pip*$_{V583}$ was measured during co-

culture with *E. faecalis* OG1RF or Δ*essB* antagonists in the presence and absence of daptomycin

treatment in **(C)** liquid culture (2.5 µg/ml), **(D)** trans-well plates to prevent physical engagement

between cells (2.5 µg/ml) and **(E)** on agar media (0.5 µg/ml). E – empty vector and C –

complementation vector. Data represent three biological replicates. *$P < 0.01$, **$P < 0.001$ to

0.0001 by unpaired Student's t-test.

Figure 5. IreK and OG1RF_11099 control transcription of enterococcal T7SS genes and

subsequent inhibition of bystander bacteria during phage infection. (A) Phage infection

leads to enhanced expression of T7SS genes in the wild type *E. faecalis* OG1RF but not in an

Δ*ireK* mutant strain. **(B)** Growth inhibition of *E. faecalis* Δ*pip*$_{V583}$ during phage infection of *E.*

faecalis OG1RF is abrogated in an Δ*ireK* mutant of OG1RF. **(C)** Disruption of OG1RF_11099

leads to reduced expression of T7SS genes during phage infection. The data are represented as

the fold change of normalized mRNA relative to uninfected samples at the same time points. **(D)**

T7SS dependent intraspecies antagonism during phage infection is alleviated in the presence of

OG1RF_11099-Tn mutant. Data represent three replicates *P < 0.00001 by unpaired Student's t-

test.

Figure S1. Phylogenetic tree of EsxA sequences in *Enterococcus*. Non-redundant

sequences (n=96) were identified using NCBI BLAST with OG1RF EsxA (OG1RF_11100) as the

input. The tree was constructed in MEGAX using the Maximum Likelihood method and JTT

matrix-based model and is drawn to scale, with branch lengths measured in the number of

substitutions per site. The tree with the highest log likelihood (-3544.39) is shown. *E. faecalis*

sequences are highlighted in purple, and the GenBank identifier for EsxA from OG1RF

(AEA93787.1) is shown in bold red font.

Figure S2. Phage VPE25 infects wild type *E. faecalis* OG1RF and ∆*essB* with similar

efficacy. The measurement of phage particles released from wild type *E. faeaclis* OG1RF, ∆*essB*,

OG1RF_11121-Tn, and OG1RF_11099-Tn mutant strains following phage VPE25 infection. Data

represent three biological replicates.

Figure S3. Putative orphan toxins in OG1RF are found as full-length LXG-domain proteins

in other bacteria.   OG1RF_11111, OG1RF_11113, and OG1RF_11123 sequences were used

as input for NCBI BLAST. Alignments and homology were rendered in EasyFig.

Figure S4. A distal OG1RF locus encodes an additional LXG-domain protein. **(A)** Schematic

showing homology between V583 (NC_004668.1, top) and OG1RF (NC_017316.1, bottom).

Sequences were obtained from NCBI, and homology comparisons were rendered in EasyFig. **(B)**

Cartoon depicting the LXG domain of OG1RF_12414 (identified using KEGG and ExPASy

PROSITE). **(C)** Predicted structural homology between OG1RF_12414 (lilac) and the

Pseudomonas protogens Pf-5 Tne2/Tni2 complex (PDB 6B12). Tne2 is shown in green, and Tni2

is shown in gray. Structural modeling was done using PHYRE2, and images were rendered in

Pymol.

Figure S5. Domain architecture of *Enterococcus* LXG proteins. Domain architectures were

identified using the NCBI Conserved Domain Architectural Retrieval Tool (DART) with

OG1RF_11109 as an input. Diagrams are drawn to scale.

Figure S6. Antibiotic susceptibility of *E. faecalis* OG1RF. Growth of wild type *E. faecalis*

OG1RF was monitored over 18 hours in the presence or absence of **(A)** ampicillin, **(B)**

vancomycin and **(C)** daptomycin in microtiter plates. The antibiotic concentrations highlighted with

a blue box were deemed sub-inhibitory and used to investigate T7SS expression levels. Early

log-phase cultures of *E. faecalis* OG1RF were grown in the presence or absence of **(D)** mitomycin

C (4 µg/ml) or **(E)** ciprofloxacin (2 µg/ml) to show that even though these routine concentrations

used to promote a stress response in other *E. faecalis*, they do not prevent bacterial growth.

Figure S7. *E. faecalis* mutants of cell wall homeostasis show no growth defects and

respond to phage VPE25 infection. (A) Optical density of wild type E. *faecalis* and isogenic

mutants were monitored for 18 hours. **(B)** While all the strains were susceptible to phage VPE25

infection, the proportion of released phage particles was diminished in the *croR* and *croS*

transposon mutant background. Data represent three biological replicates. *$P < 0.001$ by unpaired

Student's t-test.

Figure S8. Quantitative PCR demonstrates that LiaR/S and CroS/R two-component

systems do not influence T7SS gene expression during phage infection. (A- F) mRNA

transcript levels of T7SS genes are enhanced in the transposon mutants of *liaR*, *liaS*, *croR* and

croS strains similar to wild type *E. faecalis* OG1RF during phage infection (MOI = 1) compared to

untreated controls. Data represent three biological replicates. *$P < 0.01$, **$P < 0.0001$ by unpaired

Student's t-test.

Figure S9. Sub-lethal ceftriaxone exposure does not influence T7SS gene expression in *E.*

***faecalis* OG1RF.** Transcription of T7SS genes are not elevated 20 minutes post ceftriaxone

(128μg/ml) administration relative to the untreated controls.

Table S1. List of bacterial strains, phages, plasmids and primers used in this study.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. S1

Fig. S2

Fig. S3

Fig. S4

Fig. S5

Fig. S6

Fig. S7

Fig. S8

Fig. S9