Characterization of an α-glucosidase enzyme conserved in *Gardnerella* spp. isolated

from the human vaginal microbiome

Pashupati Bhandari[1], Jeffrey P. Tingley[2], D. Wade Abbott[2] and Janet E. Hill[1,*]

[1]Department of Veterinary Microbiology, Western College of Veterinary Medicine,

University of Saskatchewan, 52 Campus Drive, Saskatoon, Saskatchewan, S7N 5B4,

Canada

[2]Lethbridge Research and Development Centre, Agriculture and Agri-Food Canada,

Lethbridge, Alberta, T1J 4B1, Canada

*To whom correspondence should be addressed

Janet.Hill@usask.ca

Abstract

Gardnerella spp. in the vaginal microbiome are associated with bacterial vaginosis, a dysbiosis in which lactobacilli dominant microbial community is replaced with mixed aerobic and anaerobic bacteria including *Gardnerella* species. The co-occurrence of multiple *Gardnerella* species in the vaginal environment is common, but different species are dominant in different women. Competition for nutrients, particularly glycogen present in the vaginal environment, could play an important role in determining the microbial community structure. Digestion of glycogen into products that can be taken up by bacteria requires the combined activities of several enzymes collectively known as "amylases". In the present study, glycogen degrading abilities of *Gardnerella* spp. were assessed. We found that *Gardnerella* spp. isolates and filtered culture supernatants had amylase activity. Phylogenetic analyses predicted conserved Glycoside Hydrolase family 13 (GH13) members among *Gardnerella* spp. including a putative α-glucosidase. The gene for this enzyme was cloned and expressed, and recombinant protein was purified and functionally characterized. The enzyme was active on a variety of maltooligosaccharides over a broad pH range (4 - 8) with an optimum activity at pH 6-7. Glucose was released from maltose, maltotriose and maltopentose, however, no products were detected on thin layer chromatography (TLC) when the enzyme was incubated with glycogen. Our findings show that *Gardnerella* spp. produce a secreted α-glucosidase enzyme that can contribute to the complex and multistep process of glycogen breakdown by degrading smaller oligosaccharides into glucose, contributing to the pool of nutrients available to the vaginal microbiota.

Introduction

Gardnerella spp. in the vaginal microbiome are hallmarks of bacterial vaginosis, a condition characterized by replacement of the lactobacilli dominant microbial community with mixed aerobic and anaerobic bacteria including *Gardnerella*. This dysbiosis is associated with increased vaginal pH, malodorous discharge and the presence of biofilm (1). In addition to troubling symptoms, the presence of abnormal vaginal microbiota is associated with increased risk of HIV transmission and infection with other sexually transmitted pathogens such as *Neisseria gonorrhoeae* and *Trichomonas* spp. (2, 3). Historically, *Gardnerella* has been considered as single species genus. Jayaprakash *et al*. used cpn60 barcode sequences to divide *Gardnerella* spp. into four subgroups (A-D) (4), and this framework was supported by whole genome sequence comparison (5, 6). More recently, Vaneechoutte *et al*. emended the classification of *Gardnerella* based on whole genome sequence comparison, biochemical properties and matrix-assisted laser desorption ionization time-of-flight mass spectrometry and proposed the addition of three novel species: *Gardnerella leopoldii*, *Gardnerella piotii* and *Gardnerella swidsinskii* (7).

Colonization with multiple *Gardnerella* species is common, and different species are dominant in different women (8). Understanding factors that contribute to differential abundance is important since the species may differ in virulence (6) and they are variably associated with clinical signs (8, 9). Several factors, including inter-specific competition, biofilm formation and resistance to antimicrobials could contribute to the differential abundance of the different species. Khan *et al*. showed that resource-based scramble competition is frequent among *Gardnerella* subgroups (10). Glycogen is a significant

nutrient for vaginal microbiota, but previous reports on the growth of *Gardnerella* spp. on

glycogen containing medium are inconsistent (11). Species may differ in their ability to

digest glycogen and utilize the breakdown products, which may in turn contribute to

determining microbial community structure.

Glycogen is an energy storage molecule, which consists of linear chains of

approximately 13 glucose molecules covalently linked with α-1,4 glycosidic linkages, with

branches attached through α-1,6 glycosidic bonds (12, 13). A single glycogen molecule

consists of approximately 55,000 glucose residues with a molecular mass of ~ 10^7 kDa

(14). The size of glycogen particles can vary with source; from 10-44 nm in human skeletal

muscle to approximately 110-290 nm in human liver (15). Glycogen is deposited into the

vaginal lumen by epithelial cells under the influence of estrogen (16), and the concentration

of cell-free glycogen in vaginal fluid varies greatly, ranging from 0.1 - 32 μg/μl (17).

Digestion of this long polymeric molecule into smaller components is essential before it

can be taken up by bacterial cells (18, 19).

Glycogen digestion is accomplished by the coordinated action of enzymes that are

collectively described as "amylases" (20). α-glucosidase is an exo-acting enzyme that acts

on α-1,4 glycosidic bond from the non-reducing end to release glucose, while α-amylase is

an endo-acting amylase that cleaves α-1,4 glycosidic bonds to produce α-limit dextrins and

maltose. Debranching enzymes like pullulanase and isoamylase break α-1,6 glycosidic

bonds (15, 21). Glycogen degradation mechanisms active in the vaginal microbiome are

not completely described. Although it has been demonstrated that vaginal secretions

exhibit amylase activity that can degrade glycogen and the generation of these products

enable growth of vaginal lactobacilli (19), the role of bacterial enzymes in this process is

unknown and there is limited information about glycogen degradation by clinically

important bacteria such as *Gardnerella* spp..

The objective of the current study was to assess amylase activity of *Gardnerella*

spp. and to characterize the activity of a putative amylase protein conserved in all

Gardnerella spp..

Methods

***Gardnerella* isolates**

Isolates of *Gardnerella* used in this study were from a previously described culture

collection (6) and included representatives of *cpn*60 subgroup A (*G. leopoldii* (n = 4), *G.*

swidsinskii (n = 6), other (n = 1)), subgroup B (*G. piotii* (n = 5), Genome sp. 3 (n = 2)),

subgroup C (*G. vaginalis* (n = 6)), and subgroup D (Genome sp. 8 (n = 1), Genome sp. 9

(n = 1)). Whole genome sequences of these isolates had been determined previously

(BioProject Accessions PRJNA394757) and annotated by the NCBI Prokaryotic Genome

Annotation Pipeline (PGAP) (22).

Amylase activity assay

Amylase activity of *Gardnerella* isolates was assessed using a starch iodine test (23).

Isolates were revived from -80°C storage on Columbia blood agar plates with 5% sheep

blood, and incubated anaerobically using the GasPak system (BD, USA) at 37°C for 48

hours. Isolated colonies from blood agar plates were transferred in 0.85% saline using a

sterile swab and cell suspensions were adjusted to McFarland 1.0. The cell suspension (10

μL) was spotted on to BHI agar plates containing 1% (w/v) potato starch (Difco) or bovine

liver glycogen (Sigma-Aldrich), and plates were incubated anaerobically at 37°C for 24

hours. Lugol's iodine solution (1% w/v) was added to the plates to detect evidence of

amylase activity indicated by a clear halo around the bacterial colony.

To determine amylase activity of culture supernatant, overnight broth cultures were

centrifuged at $10,000 \times g$ for 10 mins and the cell-free supernatant was passed through a

0.45 μm filter. Filtered supernatant (10 μL) was spotted on to a 1% (w/v) starch agar plate

and incubated anaerobically at 37°C for 24 hours. Plates were flooded with 1% (w/v)

Lugol's iodine solution to detect evidence of amylase activity. α-amylase from *Bacillus*

licheniformis (0.05 mg/ml) (Sigma-Aldrich, Oakville, ON, CAT No. A455) was used as a

positive control.

Identification of putative secreted amylase sequences

The predicted proteomes of the 26 *Gardnerella* isolates used in this study were examined

with Phobius v1.01 (24) to identify cytoplasmic and non-cytoplasmic proteins based on the

detection of signal peptides. Within the predicted secreted proteins, putative amylase

enzymes were identified by annotation terms using PGAP annotation. Multiple sequence

alignments of predicted secreted amylase sequences were performed using CLUSTALw

with results viewed and edited in AliView (version 1.18) (25) prior to phylogenetic tree

building using PHYLIP (26).

Functional prediction of *Gardnerella* spp. amylases

Putative amylolytic enzymes in *Gardnerella* spp. genomes were identified using dbCAN,

and predicted sequences were run through the SACCHARIS pipeline (27). SACCHARIS

combines user sequences with sequences from the CAZy database and trims sequences to

the catalytic domain using dbCAN2 (28). Domain sequences were aligned with MUSCLE

(29), and a best-fit model was generated with ProtTest (30). Final trees were generated

with FastTree (31) and visualized with iTOL (32). Alignment of CG400_06090 from *G.*

leopoldii NR017 to orthologues from *Halomonas* sp. (BAL49684.1), *Xanthomonas*

campestris (BAC87873.1), and *Culex quinquefasciatus* (ASO96882.1) was done using

CLUSTALw and visualized in ESPript (33). A predicted structure model of CG400_06090

was aligned with *Halomonas* sp. HaG (BAL49684.1, PDB accession 3WY1) (34) using

PHYRE2 (35). BLASTp (36) alignment of CAZy derived sequences from reference

genomes to a database of the predicted proteomes of 26 study isolates was conducted to

identify orthologs.

Expression and purification of the CG400_06090 gene product

Genomic DNA from *G. leopoldi* NR017 was extracted using a modified salting out

procedure, and the CG400_06090 open reading frame (locus tag CG400_06090 in

GenBank accession NNRZ01000007, protein accession RFT33048) was PCR amplified

with primers JH0729-F (5'AT<u>G CAT GC</u>G CAT TAT ACG ATC ATG CTC-3') and

JH0730-R (5'AT<u>G GTA CC</u>T TAC ATT CCA AAC ACT GCA-3'). Underlined sequences

indicate *Sph*I and *Kpn*I restriction enzyme sites. PCR reaction contained 1 × PCR reaction

buffer (0.2 M Tris-HCl pH 8.4, 0.5 M KCl), 200 μM each dNTPs, 2.5 mM $MgCl_2$, 400 nM

each primer, 1 U/reaction of Platinum Taq DNA polymerase high-fidelity (5 units/μL in

50% glycerol, 20 mM Tris-HCl, 40 mM NaCl, 0.1 mM EDTA, and stabilizers) (Life-

technologies) and 2 µl of template DNA, in a final volume of 50 µl. PCR was performed

with following parameters: initial denaturation at 94°C for 3 minutes followed by

(denaturation at 94°C, 15 seconds; annealing at 60°C, 15 seconds and extension at 72°C, 2

minutes; 35 cycles), and final extension at 72°C for 5 minutes. Purified PCR products were

digested with *Kpn*I and *Sph*I and ligated into expression vector pQE-80 L (Qiagen,

Mississauga, ON) digested with same restriction endonucleases. The resulting recombinant

plasmid was used to transform One Shot TOP10 chemically competent *E. coli* cells

(Invitrogen, Carlsbad, California). Colony PCR was performed to identify transformants

containing vector with insert. Insertion of the putative amylase gene in-frame with the N-

terminal 6×Histidine tag was confirmed by sequencing of the purified plasmid.

E. coli containing the plasmid were grown overnight at 37°C in LB medium with

100 µg/ml ampicillin. Overnight culture was diluted 1:60 in fresh medium to a final volume

of 50 ml, and the culture was incubated at 37°C with shaking at 225 rpm until it reached

an OD_{600} of 0.6. At this point, expression was induced with 0.1 mM IPTG and a further 4

hours of incubation was done at 37°C at 225 rpm. Cells were harvested by centrifugation

at $10,000 \times g$ for 30 mins and the pellet was resuspended in lysis buffer (50 mM NaH_2PO_4,

300 mM NaCl, 10 mM imidazole, pH 8.0). Lysozyme was added to 1 mg/ml and cells were

lysed by sonication (8 mins total run time with 15 sec on and 30 sec off). The lysate was

clarified by centrifugation at $10,000 \times g$ for 30 mins at 4°C and the supernatant was applied

to an Ni-NTA affinity column according to the manufacturer's instructions (Qiagen,

Germany). Bound proteins were washed twice with wash buffer (50 mM NaH_2PO_4, 300

mM NaCl, 20 mM imidazole, pH 8.0) and eluted with elution buffer (50 mM NaH_2PO_4,

300 mM NaCl, 250 mM imidazole, pH 8.0). Eluted protein was buffer exchanged with the

same elution buffer but without imidazole using 30K MWCO protein concentrator

(ThermoFisher Scientific). Final protein concentration was measured by

spectrophotometry.

Measuring enzyme activity

Purified protein was tested for α-glucosidase activity by measuring the release of 4-

Nitrophenol from a chromogenic substrate 4-Nitrophenyl-α-D-glucopyranoside as

described elsewhere (37). Briefly, 25 µl of enzyme solution (50 µg/ml) was added to 25 µl

of 10 mM 4-Nitrophenylα-D-glucopyranoside substrate and the reaction mixture was

incubated at 37°C for 10 mins. Reactions were stopped by the addition of 100 µl of 1M

Na_2CO_3 solution and OD_{420} was measured. To determine activity at different pH values,

substrate was prepared in 50 mM citric acid-sodium citrate buffer (pH 3.0) or 50 mM

sodium phosphate buffer (pH 4.0-8.0).

Enzymatic activity on different substrates

The activity of the purified enzyme was examined with different substrates according to

Cockburn et *al.* (38). Briefly, 10 µl of enzyme solution (5.0 mg/ml) was added to 190 µl

of 0.1% (w/v) maltose, maltotriose, maltopentose, maltodextrins (MD 4-7, MD 13-17 and

MD 16.5-19.5) or glycogen in 50 mM sodium phosphate buffer, pH 6.0, and reaction

mixture was incubated at 37°C. Aliquots of the reaction mixture (10 µl) were taken out at

various time intervals of incubation and were frozen immediately until thin layer

chromatography (TLC) was performed. Standards and the reaction mixtures (0.5 µl) were

loaded on 0.25 mm silica gel plates (Sigma-Aldrich, Oakville, ON) and the chromatogram

was developed in a solvent system containing acetonitrile, ethyl acetate, 2-propanol and

water in (8.5:2:5:6 vol/vol) in a developing chamber until the mobile phase reached the top

of the plates. The plates were allowed to dry in a fume hood and products were resolved

by dipping the plates into the 0.3% solution of 1-napthyl ethylene diamine dihydrochloride

followed by drying and heating with the heat gun until spots became visible.

Results

Amylase activity of *Gardnerella* spp.

All 26 isolates tested showed amylase activity by degrading starch and glycogen as

indicated by the appearance of a clear zone around bacterial colonies after iodine staining.

A filter sterilized culture supernatant from *G. leopoldii* (NR017) was also able to digest the

starch and glycogen in the agar plate assay, indicating that amylase enzymes are secreted

(Figure 1).

Identification of sequences of secreted amylases

A total of 2,525 predicted proteins from the combined proteomes of the 26 study isolates

were determined to contain a signal peptide and predicted to be non-cytoplasmic. Of these

putative secreted proteins, 60 were annotated as amylases. After removal of severely

truncated sequences, a multiple sequence alignment of the remaining 42 sequences was

trimmed to the uniform length of 290 amino acids. Phylogenetic analysis of these predicted

extracellular amylase sequences revealed five clusters, but only one of these clusters

contained orthologous sequences from all 26 is1olates (Figure 2). Sequence identities

within this cluster were all 91-100%. This conserved sequence (corresponding to GenBank

accession WP_004104790) was annotated as an GH13 α-amylase (EC 3.2.1.1) in all

genomes, suggesting it was an endo-acting enzyme that releases maltooligosaccharides

from glycogen.

The GH13 family in the CAZy database is a large, polyspecific family targeting α-

glucose linkages. The functional range of members in this family make annotation of new

sequences challenging. In order to predict the activity of the conserved protein identified

among the study isolates, we employed SACCHARIS, which combines CAZyme family

trees generated from biochemically characterized proteins with related sequences of

unknown function. Currently, two *Gardnerella* reference genomes are available in CAZy:

G. vaginalis ATCC 14018 and *G. swidsinkii* GV37. GH13 family proteins encoded in these

Gardnerella genomes were identified and a family tree including all characterized GH13

sequences in the CAZy database was generated (Figure S1). SACCHARIS annotation of

the GH13 family in *Gardnerella* resulted in identification of protein domains belonging to

ten subfamilies. Alignment with characterized GH13 enzymes from the CAZy database

(39), suggested that the conserved amylase identified in the study isolates was an α-

glucosidase (EC 3.2.1.20), closely related to subfamilies 23 and 30 of GH13, and not an α-

amylase as annotated in GenBank.

Relationship to other α-glucosidases

A representative sequence of the predicted α-glucosidase was selected from *G.*

leopoldii (NR017). This sequence (CG400_06090) was combined with functionally

characterized members of the GH13 CAZy database using SACCHARIS (Figure 3A).

CG400_06090 partitioned with members of GH13 subfamilies 17, 23 and 30, of which the

prominent activity is α-glucosidase. This clade was expanded to better illustrate the

relationship of CG400_06090 with related characterized sequences as an apparently deep-

branching member of subfamily 23, although with low bootstrap support (Figure 3B).

When sequences of additional (not functionally characterized) members were included,

CG400_06090 clustered within subfamily 23 (Figure S2).

To identify conserved catalytic residues between CG400_06090 and characterized

members of GH13, CG400_06090 was aligned with structurally characterized members of

subfamily 23 (BAL49684.1 and BAC87873.1) and subfamily 17 (ASO96882.1) (Figure

3C). Of the GH13 catalytic triad, the aspartate nucleophile and the aspartate that stabilizes

the transition state are conserved among the four sequences (40). The glutamate general

acid/base, however, does not appear to be sequence conserved between CG400_06090 and

the GH13 subfamily 23 members. Upon modelling CG400_06090 in PHYRE2 and

aligning with BAL49684.1 (PDBID 3WY1), CG400_06090 Glu256 did appear to be

spatially conserved with the general acid/base (not shown).

Expression and purification of α-glucosidase protein

The *G. leopoldi* NR017 CG400_06090 open reading frame comprises 1701 base pairs

encoding a 567 amino acid protein, with a predicted mass of 62 kDa. The region of the

open reading frame encoding amino acids 129-418 was PCR amplified and ligated into

vector pQE-80L for expression in *E. coli* as an N-terminal 6xHis-tagged protein. A distinct

protein band in SDS-PAGE between 50 kDa and 75 kDa was obtained from IPTG induced

E. coli cells compared with non-induced cells, indicating the expression of the α-

glucosidase protein (Figure 4A). The recombinant protein was soluble, and was purified

using a Ni-NTA spin column (Figure 4B). From a 50 ml broth culture, approximately 700

µl of purified protein at 4.0 mg/ml was obtained.

Effect of pH on enzyme activity

α-glucosidase activity of the purified protein was demonstrated by release of 4-Nitrophenol

from chromogenic substrate 4-Nitrophenyl-α-D-glucopyranoside. This activity was

observed across a broad pH range (4.0 – 8.0), with maximum activity at pH 6.0 and 7.0

while no activity was seen at pH 3.0 (Figure 5).

Analysis of substrate hydrolysis

Production of glucose was detected when the purified α-glucosidase was incubated with

maltose, maltotriose and maltopentose at pH 6.0 (Figure 6). No appreciable activity,

however, was detected on maltodextrins (MD 4-7, MD 13-17 and MD 16.5-19.5) or

glycogen.

Discussion

Glycogen is a major available nutrient to vaginal microbiota and its utilization

likely plays an important role in the survival and success of *Gardnerella* spp. in the vaginal

microbiome. Glycogen is digested into smaller products, such as maltose and

maltodextrins, by a group of enzymes collectively known as "amylases". In this study, we

assessed the amylase activity of *Gardnerella* spp. and an α-glucosidase gene conserved in

Gardnerella spp. was expressed and its product was characterized.

Isolates of *G. leopoldii*, *G. swidsinskii*, *G. piotii* and *G. vaginalis* all showed

amylase activity by degrading starch and glycogen in an agar plate assay. Previously, Piot

et al. examined the starch fermentation ability of 175 *Gardnerella* isolates on Mueller

Hinton agar supplemented with horse serum and found that all *Gardnerella* strains were

capable of hydrolysing starch (41). They did not, however, assess glycogen degradation

and further, it is not clear if serum amylase contained in the media contributed to the

observed amylase activity. Robinson, in his review, reported that starch can be utilized by

Gardnerella (42) while Catlin in another review, reported that glycogen utilization is

inconsistent in *Gardnerella* (11). All *Gardnerella* spp. assessed in our study were amylase

positive, which suggests it is a conserved catalytic function. As these bacteria are found

almost exclusively in the glycogen-rich human vagina, it is not surprising that they produce

amylases to break down this substrate.

In the phylogenetic analysis of putative secreted amylase sequences, one cluster

contained sequences common to all 26 isolates. Although this protein sequence was

annotated as an α-amylase, suggesting it would produce maltooligosaccahrides from

glycogen, SACCHARIS predicted it to be closely related to α-glucosidases, which cleave

glucose from the non-reducing end of its substrate. SACCHARIS predicts enzyme activity

by comparing sequence domains to sequences deposited in the curated CAZy database for

which structural and / or biochemical information is available (27). Automated annotation

of sequences deposited in primary sequence databases, such as GenBank has resulted in

significant problems with functional mis-annotation (43). This problem is exacerbated by

the increasing rate at which whole genome sequences are being generated and deposited.

The genome sequences of the study isolates were annotated using the NCBI Prokaryotic

Genome Annotation Pipeline, which identifies genes and annotates largely based on

sequence similarity to previously annotated sequences, thus potentially propagating

incorrect functional annotations.

In the phylogenetic analysis of GH13 family enzymes, the conserved α-glucosidase

from *Gardnerella* spp. appears to be most closely related to GH13 subfamily 17, 23 and

30 characterized members, forming a monophyletic group. However, low bootstrap values

between CG400_06090 and other clade members make assignment of CG400_06090 to a

defined subfamily difficult based on comparison to sequences of functionally characterized

proteins (Figure 3B). Additional support for assignment to subfamily 23 was observed

when additional sequences of proteins lacking functional characterization from subfamilies

17, 23, and 30 were included (Figure S2). Regardless, the α-glucosidase activity observed

within this clade does extend to CG400_06090. Primary sequences of GH13 members of

amylases have seven highly conserved regions and three amino acids that form the catalytic

triad (21). Comparison of the primary sequence and three-dimensional model of

CG400_06090 with characterized α-glucosidases from GH13 subfamily 17 and 23 did

show sequence conservation of nucleophile and spatial conservation of the catalytic triad.

Our results demonstrate the value of using SACCHARIS to guide discovery of

CAZymes based on comparison at a catalytic domain level to functionally characterized

enzymes. All of the other proteins identified as putative secreted amylases were also

annotated as "alpha-amylase" in the GenBank records. Further investigation will be

required to demonstrate the actual functions of these proteins, and how they contribute to

carbohydrate degrading activities of *Gardnerella* spp..

Amylase enzymes that contribute to glycogen digestion in the vaginal environment

are not well studied. Spear *et al*. (19) suggested a role for a human amylase enzyme in the

breakdown of vaginal glycogen. They detected host-derived pancreatic α-amylase and

acidic α-glucosidase in genital fluid samples collected from women and showed that genital

fluid containing these enzymes can degrade the glycogen. The presence of these enzyme

activities alone in vaginal fluid, however, was not correlated with glycogen degradation,

suggesting contributions of activity from the resident microbiota. Many vaginal bacteria

likely produce amylases that contribute to this activity as various anaerobic taxa are known

to encode amylases and digest glycogen (44), however, the contributions of the vaginal

microbiota to this process and the importance of these processes to vaginal microbial

community dynamics are not yet well understood.

α-glucosidase enzymes are typically active over a broad pH range and exhibit

optimal activity at pH 4.5-6.0 (45). They have also been characterized in other bacteria

isolated from human body sites, particularly the intestinal tract. An extracellular α-

glucosidase form *Lactobacillus acidophilus* NCTC 1723 was reported to have activity at

pH 6.0-7.5 (46). Van den Broek *et al.* characterized an α-glucosidase (*algB*) in

Bifidobacterium adolescentis DSM20083 and reported its maximum activity at pH 6.8.

This protein had a molecular mass of 73 kDa and used maltose and maltotriose as substrates

(47). Purified α-glucosidase from *G. leopoldii* NR017 was active across a broad pH range

(4.0 – 8.0). By definition, the pH of the vaginal fluid in women with bacterial vaginosis is

> 4.5 (48), and has been reported as high as 7.0 - 7.5 (49), so it is not surprising for

Gardnerella to produce a secreted enzyme that functions optimally at pH 6-7. In fact, this

activity at high pH could help *Gardnerella* spp. liberate glucose at an increased rate in

dysbiotic microbiomes, thus increasing the nutrient pool available and contributing to

supporting the overgrowth of vaginosis-associated bacteria.

Complete glycogen digestion is a complex and multistep process that requires the

action of several different enzymes at the different stages. Protein sequences belonging to

eight different GH13 subfamilies, which could have roles in other steps of the glycogen

degradation, were identified in *G. vaginalis* ATCC 14018 and *G. swidsinskii* GV37

(Supplemental Figure 1), and multiple secreted amylase-like proteins were detected in the

proteomes of additional *Gardnerella* spp. (Figure 2). Functional characterization of these

proteins will be required to understand their roles in glycogen degradation.

Our findings show that *Gardnerella* spp. have a secreted α-glucosidase enzyme that

likely contributes to the complex and multistep process of glycogen breakdown by

releasing glucose from oligosaccharides, contributing to the pool of nutrients available to

the vaginal microbiota. Identification and biochemical characterization of additional

enzymes involved in glycogen digestion will provide insight into whether utilization of this

abundant carbon source is an important factor in population dynamics and competition

among *Gardnerella* spp..

Funding

This research is supported by an NSERC Discovery grant to JEH, and Agriculture and

Agri-Food Canada project number: J-001589. PB is supported by a Devolved Scholarship

from the University of Saskatchewan.

Conflicts of interest

The authors declare no competing interest.

Acknowledgements

The authors are grateful to Josseline Ramos-Figueroa, Douglas Fansher, Natasha Vetter and Dr. David Palmer (Department of Chemistry, University of Saskatchewan) for guidance with thin layer chromatography and Champika Fernando for technical support.

References

1.   Nugent RP, Krohn MA, Hillier SL. 1991. Reliability of diagnosing bacterial

vaginosis is improved by a standardized method of gram stain interpretation. J Clin

Microbiol 29:297–301.

2.   Martin HL, Richardson BA, Nyange PM, Lavreys L, Hillier SL, Chohan B,

Mandaliya K, Ndinya-Achola JO, Bwayo J, Kreiss J. 1999. Vaginal *Lactobacilli*,

microbial flora, and risk of Human Immunodeficiency virus Type 1 and sexually

transmitted disease acquisition. J Infect Dis 180:1863–1868.

3.   Taha TE, Hoover DR, Dallabetta GA, Kumwenda NI, Mtimavalye LAR, Yang L-P,

Liomba GN, Broadhead RL, Chiphangwi JD, Miotti PG. 1998. Bacterial vaginosis

and disturbances of vaginal flora: association with increased acquisition of HIV.

AIDS 12.

4.   Paramel Jayaprakash T, Schellenberg JJ, Hill JE. 2012. Resolution and

characterization of distinct cpn60-based subgroups of *Gardnerella vaginalis* in the

vaginal microbiota. PloS One 7:e43009–e43009.

5.   Ahmed A, Earl J, Retchless A, Hillier SL, Rabe LK, Cherpes TL, Powell E, Janto B,

Eutsey R, Hiller NL, Boissy R, Dahlgren ME, Hall BG, Costerton JW, Post JC, Hu

FZ, Ehrlich GD. 2012. Comparative genomic analyses of 17 clinical isolates of

Gardnerella vaginalis provide evidence of multiple genetically isolated clades

consistent with subspeciation into genovars. J Bacteriol 194:3922–3937.

409 6. Schellenberg JJ, Paramel Jayaprakash T, Withana Gamage N, Patterson MH,

410   Vaneechoutte M, Hill JE. 2016. *Gardnerella vaginalis* subgroups defined by cpn60

411   sequencing and Sialidase activity in isolates from Canada, Belgium and Kenya.

412   PloS One 11:e0146510.

413 7. Vaneechoutte M, Guschin A, Van Simaey L, Gansemans Y, Van Nieuwerburgh F,

414   Cools P. 2019. Emended description of *Gardnerella vaginalis* and description of

415   *Gardnerella leopoldii* sp. nov., *Gardnerella piotii* sp. nov. and *Gardnerella*

416   *swidsinskii* sp. nov., with delineation of 13 genomic species within the genus

417   *Gardnerella*. Int J Syst Evol Microbiol 69:679–687.

418 8. Hill JE, Albert AYK. 2019. Resolution and cooccurrence patterns of *Gardnerella*

419   *leopoldii*, *G. swidsinskii*, *G. piotii*, and *G. vaginalis* within the vaginal microbiome.

420   Infect Immun 87:e00532-19.

421 9. Balashov SV, Mordechai E, Adelson ME, Gygax SE. 2014. Identification,

422   quantification and subtyping of *Gardnerella vaginalis* in noncultured clinical

423   vaginal samples by quantitative PCR. J Med Microbiol 63:162–175.

424 10. Khan S, Voordouw MJ, Hill JE. 2019. Competition among *Gardnerella* subgroups

425   from the human vaginal microbiome. Front Cell Infect Microbiol 9:374.

426 11. Catlin BW. 1992. *Gardnerella vaginalis*: characteristics, clinical considerations, and

427   controversies. Clin Microbiol Rev 5:213–237.

12. Meléndez-Hevia E, Waddell TG, Shelton ED. 1993. Optimization of molecular

design in the evolution of metabolism: the glycogen molecule. Biochem J 295 (Pt

2):477–483.

13. Deng B, Sullivan MA, Chen C, Li J, Powell PO, Hu Z, Gilbert RG. 2016. Molecular

structure of human liver glycogen. PloS One 11:e0150540–e0150540.

14. Roach PJ, Depaoli-Roach AA, Hurley TD, Tagliabracci VS. 2012. Glycogen and its

metabolism: some new developments and old themes. Biochem J 441:763–787.

15. Adeva-Andany MM, González-Lucán M, Donapetry-García C, Fernández-Fernández

C, Ameneiros-Rodríguez E. 2016. Glycogen metabolism in humans. BBA Clin

5:85–100.

16. Cruickshank R, Sharman A. 1934. The biology of the vagina in the human subject.

BJOG Int J Obstet Gynaecol 41:208–226.

17. Mirmonsef P, Hotton AL, Gilbert D, Gioia CJ, Maric D, Hope TJ, Landay AL, Spear

GT. 2016. Glycogen levels in undiluted genital fluid and their relationship to

vaginal pH, estrogen, and progesterone. PloS One 11:e0153553.

18. Flint HJ, Scott KP, Duncan SH, Louis P, Forano E. 2012. Microbial degradation of

complex carbohydrates in the gut. Gut Microbes 3:289–306.

19. Spear GT, French AL, Gilbert D, Zariffard MR, Mirmonsef P, Sullivan TH, Spear

WW, Landay A, Micci S, Lee B-H, Hamaker BR. 2014. Human α-amylase present

in lower-genital-tract mucosal fluid processes glycogen to support vaginal

colonization by Lactobacillus. J Infect Dis 210:1019–1028.

20. Berg JM, Tymoczko JL, Stryer L. 2002. Glycogen breakdown requires the interplay

of several enzymes.Biochemistry, 5th ed. W H Freeman, New York.

21. Mehta D, Satyanarayana T. 2016. Bacterial and archaeal α-amylases: Diversity and

amelioration of the desirable characteristics for industrial applications. Front

Microbiol 7.

22. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L,

Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI prokaryotic genome

annotation pipeline. Nucleic Acids Res 44:6614–6624.

23. Ryan SM, Fitzgerald GF, van Sinderen D. 2006. Screening for and identification of

Starch-, Amylopectin-, and Pullulan-degrading activities in Bifidobacterial Strains.

Appl Environ Microbiol 72:5289.

24. Lukas Käll, Anders Krogh, Erik L.L. Sonnhammer. 2004. A combined

transmembrane topology and signal peptide prediction method. J Mol Biol

338:1027–1036.

25. Larsson A. 2014. AliView: a fast and lightweight alignment viewer and editor for

large datasets. Bioinforma Oxf Engl 30:3276–3278.

26. Felsenstein J. 1989. PHYLIP - phylogeny inference package (version 3.2). Cladistics

5:164–166.

27. Jones DR, Thomas D, Alger N, Ghavidel A, Inglis GD, Abbott DW. 2018.

SACCHARIS: an automated pipeline to streamline discovery of carbohydrate active

enzyme activities within polyspecific families and de novo sequence datasets.

Biotechnol Biofuels 11:27.

28. Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, Busk PK, Xu Y, Yin Y.

2018. dbCAN2: a meta server for automated carbohydrate-active enzyme

annotation. Nucleic Acids Res 46:W95–W101.

29. Edgar RC. 2004. MUSCLE: a multiple sequence alignment method with reduced

time and space complexity. BMC Bioinformatics 5:113.

30. Darriba D, Taboada GL, Doallo R, Posada D. 2011. ProtTest 3: fast selection of best-

fit models of protein evolution. Bioinformatics 27:1164–1165.

31. Price MN, Dehal PS, Arkin AP. 2010. FastTree 2 – Approximately Maximum-

Likelihood Trees for Large Alignments. PLOS ONE 5:e9490.

32. Letunic I, Bork P. 2019. Interactive Tree Of Life (iTOL) v4: recent updates and new

developments. Nucleic Acids Res 47:W256–W259.

33. Robert X, Gouet P. 2014. Deciphering key features in protein structures with the new

ENDscript server. Nucleic Acids Res2014/04/21. 42:W320–W324.

34. Shen X, Saburi W, Gai Z, Kato K, Ojima-Kato T, Yu J, Komoda K, Kido Y, Matsui

H, Mori H, Yao M. 2015. Structural analysis of the α-glucosidase HaG provides

new insights into substrate specificity and catalytic mechanism. Acta Crystallogr

Sect D 71:1382–1391.

35. Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. 2015. The Phyre2 web

portal for protein modeling, prediction and analysis. Nat Protoc 10:845–858.

36. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment

search tool. J Mol Biol 215:403–410.

37. Angelov A, Putyrski M, Liebl W. 2006. Molecular and biochemical characterization

of alpha-glucosidase and alpha-mannosidase and their clustered genes from the

thermoacidophilic archaeon *Picrophilus torridus*. J Bacteriol 188:7123–7131.

38. Cockburn D, Koropatkin N. 2015. Product analysis of starch active enzymes by TLC.

Bioprotocol 5:1–8.

39. Lombard V, Golaconda Ramulu H, Drula E, Coutinho PM, Henrissat B. 2014. The

carbohydrate-active enzymes database (CAZy) in 2013. Nucleic Acids Res

42:D490–D495.

40. Uitdehaag JCM, Mosi R, Kalk KH, van der Veen BA, Dijkhuizen L, Withers SG,

Dijkstra BW. 1999. X-ray structures along the reaction pathway of cyclodextrin

glycosyltransferase elucidate catalysis in the α-amylase family. Nat Struct Biol

6:432–436.

41. Piot P, Van Dyck E, Totten PA, Holmes KK. 1982. Identification of *Gardnerella*

(*Haemophilus*) *vaginalis*. J Clin Microbiol 15:19–24.

42. Taylor-Robinson D. 1984. The bacteriology of *Gardnerella vaginalis*. Scand J Urol

Nephrol Suppl 86:41.

43. Schnoes AM, Brown SD, Dodevski I, Babbitt PC. 2009. Annotation error in public

databases: misannotation of molecular function in enzyme superfamilies. PLoS

Comput Biol 5:e1000605–e1000605.

44. Nunn KL, Forney LJ. 2016. Unraveling the dynamics of the human vaginal

microbiome. Yale J Biol Med 89:331–337.

45. Del Moral S, Barradas-Dermitz DM, Aguilar-Uscanga MG. 2018. Production and

biochemical characterization of α-glucosidase from *Aspergillus niger* ITV-01

isolated from sugar cane bagasse. 3 Biotech 8:7–7.

46. Chan K, Li KB. 1981. Properties of alpha-glucosidase from *Lactobacillus*

acidophilus NCTC 1723. Microbios 32:47–61.

47. Van den Broek L, Struijs K, Verdoes J, Beldman G, Voragen A. 2003. Cloning and

characterization of two α-glucosidases from *Bifidobacterium adolescentis*

DSM20083. Appl Microbiol Biotechnol 61:55–60.

48. Amsel R, Totten PA, Spiegel CA, Chen KC, Eschenbach D, Holmes KK. 1983.

Nonspecific vaginitis. Diagnostic criteria and microbial and epidemiologic

associations. Am J Med 74:14–22.

49. O'Hanlon DE, Moench TR, Cone RA. 2013. Vaginal pH and microbicidal lactic acid

when lactobacilli dominate the microbiota. PloS One 8:e80074–e80074.

Figure 1. Amylase activity of filtered culture supernatant (CS) from *G. leopoldii* NR017 on starch (A) and glycogen (B) agar. PC: positive control (amylase enzyme from *B. licheniformis*, 0.05 mg/ml) and NC: negative control, BHI broth.

Figure 2. Phylogenetic analysis of predicted extracellular amylases protein sequences from 26 *Gardnerella* genomes. Trees are consensus trees of 100 bootstrap iterations. Bootstrap values are shown for major branch points. Species is indicated by label colour according to the legend.

Figure 3. (A) Phylogenetic tree of characterized GH13 members in the CAZy database and predicted *G. leopoldii* NR017 GH13 sequence CG400_06090. (B) Expanded tree of subfamilies

17, 23 and 30. Stars indicate structurally characterized sequences and all bootstrap values were included. (C) CLUSTAL alignment of the catalytic domain of CG400_06090 with catalytic domains of GH13 subfamily 23 proteins from *Halomonas* sp. (BAL49684.1) and *Xanthomonas campestris* (BAC87873.1), and subfamily 17 protein from *Culex quinquefasciatus* (ASO96882.1). Red arrows indicate the conserved members of the catalytic triad, and a red circle indicates the acid/base (Glu242) in BAL49684.1 and BAC87873.1. White circle indicates the predicted acid/base in CG400_06090. Invariant sequences are highlighted in black.

Figure 4. Production and purification of α-glucosidase protein. (A) A protein of the predicted mass of 62 kDa was observed in induced cultures. M: size marker, NP: *E. coli* with no vector, UN: uninduced culture, IN: induced with IPTG. Numbers on the left indicate the marker size in kDa. (B) Fractions from Ni-NTA affinity purification of His-tagged recombinant protein. M: size marker, L: lysate, P: pellet, F: flow-through, W1: first wash, W2: second wash, E1: first elution, E2: second elution.

Figure 5. pH profile of α-glucosidase enzyme. The enzyme was incubated with 10 mM paranitrophenyl α-D-glucopyranoside substrate at different pH and release of paranitrophenol was measured as described in the methods. Results from four independent experiments each with two technical replicates are shown.

Figure 6. TLC of products of maltose (A), maltotriose (B) and maltopentose (C) hydrolysis by α-glucosidase enzyme. B: Substrate with no enzyme. Reaction mixtures were assessed at 3 h, 6 h, 12 h, 24 h and 48 h.

Figure S1. (A) Phylogenetic tree of GH13 functionally characterized members with predicted GH13 domains from *G. swindsinskii* GV37 (red arrow) and *G. vaginalis* ATCC 14018 (blue arrow). Tree branches are coloured based on GH13 subfamily. Trees were generated using SACCHARIS and viewed in iTOL.

A.

Subfamily 17
Subfamily 23
Subfamily 30
▲ *G. leopoldii* NR017 sequence
☆ Characterized sequence(s)

Figure S2. (A) Phylogenetic tree of all GH13 subfamily 17, 23 and 30 members in the CAZy database and *G. leopoldii* CG400_06090. Branch colour is based on GH13 subfamily and functionally characterized proteins are indicated with stars. The clade highlighted in grey including *G. leopoldii* CG400_06090 and its closest relatives is expanded in (B).