# Genetic determinants of receptor-binding preference and zoonotic potential of H9N2 avian influenza viruses

Thomas P. Peacock[a,b#], Joshua E. Sealy[b,c#], William T. Harvey[d#], Donald J. Benton[e], Richard Reeve[d], and Munir Iqbal[b*]

[a]Department of Infectious Diseases, Imperial College London, UK, W2 1NY
[b]Avian Influenza Group, The Pirbright Institute, Woking, UK, GU24 0NF
[c]Royal Veterinary College, University of London, UK, NW1 0TU
[d]Boyd Orr Centre for Population and Ecosystem Health, Institute of Biodiversity, Animal Health and Comparative Medicine, College of Medical, Veterinary and Life Sciences, University of Glasgow, UK, G12 8QQ
[e]The Francis Crick Institute, London, UK, NW1 1ST
#TPP, JES and WTH contributed equally to this work
*Corresponding author: Professor Munir Iqbal, The Pirbright Institute, Pirbright, Woking, GU24 0NF, United Kingdom (E-mail. munir.iqbal@pirbright.ac.uk, tel: +44 (0)1635-577273)

## Abstract

Receptor recognition and binding is the first step of viral infection and a key determinant of host specificity. The inability of avian influenza viruses to effectively bind human-like sialylated receptors is a major impediment to their efficient transmission in humans and pandemic capacity. Influenza H9N2 viruses are endemic in poultry across Asia and parts of Africa where they occasionally infect humans and are therefore considered viruses with zoonotic potential. We previously described H9N2 viruses, including several isolated from human zoonotic cases, showing a preference for human-like receptors. Here we take a mutagenesis approach, making viruses with single or multiple substitutions in H9 haemagglutinin to determine the genetic basis of preferences for alternative avian receptors and for human-like receptors. We describe amino acid motifs at positions 190, 226 and 227 that play a major role in determining receptor specificity, and several other residues such as 159, 188, 193, 196, 198 and 225 play a smaller role. Furthermore, we show changes at residues 135, 137, 147, 157, 158, 184, 188, and 192 can also modulate virus receptor avidity and that substitutions that increased or decreased the net positive charge around the haemagglutinin receptor-binding site show increases and decreases in avidity, respectively. The motifs we identify as increasing preference for the human-receptor will help guide future H9N2 surveillance efforts and facilitate our understanding of the emergence of influenza viruses with high zoonotic potential.

## Author Summary

As of 2020, over 60 infections of humans by H9N2 influenza viruses have been recorded in countries were the virus is endemic. Avian-like cellular receptors are the primary target for these viruses. However, given that human infections have been detected on an almost monthly basis since 2015, there may be a capacity for H9N2 viruses to evolve and gain the ability to target human-like cellular receptors. Here we identify molecular signatures that can cause viruses to bind human-like receptors, and we identify the molecular basis for the distinctive preference for sulphated receptors displayed by the majority of recent H9N2 viruses. This work will help guide future surveillance by providing markers that signify the emergence of viruses with enhanced zoonotic potential as well as improving understanding the basis of influenza virus receptor-binding.

## Introduction

44
In the 1990s avian influenza virus subtypes H5N1 and H9N2 underwent a host-switch from
wild birds to domestic poultry where they have circulated ever since. H9N2 has since become one
of the most widespread strains in poultry, infecting domestic fowl throughout Asia and North Africa,
where it circulates hyper-endemically [1-4]. Zoonotic H9N2 cases are also occasionally detected,
with human infections reported in Hong Kong, mainland China, Bangladesh, Egypt, Pakistan, Oman,
Indian and Senegal; over half of human infections have been reported in the last 4 years alone, all
of which indicates a growing pandemic threat from these viruses [4-12]. Although no human-to-
human transmission has been recorded, some H9N2 virus strains have shown a high propensity for
airborne transmission between ferrets [13, 14], the most commonly used model for human
influenza transmission.

The influenza glycoprotein, haemagglutinin (HA), mediates attachment of virus to host cells
through binding of glycans with terminal sialic acids moieties. The human upper respiratory tract
(URT) is rich in glycans with terminal α2,6-linked sialic acid (SA) and is the primary site of replication
for human influenza viruses. An important determinant for adaptation to the human URT is the
ability of HA to bind α2,6-linked sialylated glycans [15]. However, most avian influenza viruses
preferentially bind to glycans with terminal α2,3-linked SAs, which are common in the avian
gastrointestinal and respiratory tracts [16]. Therefore, for avian influenza viruses to be able to
efficiently infect and transmit between humans they must gain the ability to bind to α2,6-linked SA.

Several molecular determinants have been shown to influence the receptor-binding profile
of H9N2 viruses, including position 226 (H3 HA numbering used throughout; position 216 in mature
polypeptide H9 numbering) – in one strain, Q226L alone could facilitate greater replication of an
H9N2 isolate in human epithelial airway cells [17]. Furthermore, substitutions at positions 155, 190
and 227 (145, 180 and 217 in H9 numbering) have also been shown to play a role in receptor-binding
preference in some H9N2 viruses [13, 18-20]. However, understanding of H9N2 receptor-binding
preference remains piecemeal, with no studies having systematically looked at the roles of multiple
residues, or residue combinations, in the variation in H9N2 receptor preference.

In a previous study, we tested the receptor-binding preference of several H9N2 viruses and
described notable receptor preference variability amongst circulating H9N2 viruses which we
hypothesised was due to amino acid variability at residues 190, 226 and 227 [21]. In this study we
take three H9N2 viruses that are representative of different receptor-binding profiles, including a
virus isolated from a human with a natural preference for α2,6-linked SA, and generate recombinant
virus libraries with HA amino acid substitutions that represent reciprocal changes between the
progenitor H9N2 viruses. We test the receptor-binding of these libraries and show residues 190,
226, 227, and to a lesser extent 159, 188, 193, 196, 198, and 225, explain this receptor preference
variability. We further show several antibody escape mutants have changed receptor preference or
avidity, and we describe a correlation between the electrostatic charge of the HA head and receptor
avidity. Finally, we use the insights from these experiments to predict that certain H9N2 lineages
may have a naturally high propensity to bind human receptors and therefore possess a higher
zoonotic potential within the general viral population.

## Results

84
Three previously characterised viruses were chosen to act as mutagenesis backgrounds due
to their distinct receptor-binding phenotypes. Receptor-binding profiles were measured using three
receptor analogues: sulphated and non-sulphated 3'sialyllactosamine (3SLN(6su) and 3SLN,
respectively) and 6'sialyllactosamine (6SLN). Both 3SLN(6su) and 3SLN are analogues for avian-like
receptors while 6SLN is an analogue for human-like receptors. The virus A/chicken/Pakistan/UDL-

01/2008 (UDL1/08) displays high binding avidity to 3SLN(6su), but not 3SLN (avian-like), with
residual binding to the human-like receptor 6SLN (Figure 1A), similar to the majority of
contemporary H9N2 viruses [20-22]. The virus A/chicken/Emirates/R66/2002 (Em/R66) binds to
both 3SLN(6su) and 3SLN but has no detectable binding to 6SLN (Figure 1B), similar to conventional
avian-adapted H5N1 and H7 viruses [21, 23]. Finally, A/Hong Kong/33982/2009 (HK/33982) binds to
all three receptor analogues, but with an appreciable preference for 6SLN, similar to early human
pandemic H3N2 viruses and zoonotic H7N9 viruses (Figure 1C) [21, 24]. To test the molecular basis
of these different receptor preferences, libraries of individual or multiple reciprocal mutants were
generated between these viruses with a particular focus on positions 190, 226 and 227, as well as
several other nearby RBS residues.

100
***Figure 1 – Varying receptor binding profiles of naturally occurring viruses.*** *Biolayer*
*interferometry was used to determine the receptor binding profiles of H9N2 viruses UDL1/08*
*(Panel A), Em/R66 (B), and HK/33982 (C). Binding was measured to three receptor analogues*
*representative of: 1) an avian-like receptor (α2,3-sialyllactosamine or 3SLN, red lines), 2) a*
*sulphated avian-like receptor (Neu5Ac α2,3 Gal β1-4(6-HSO3)GlcNAc or 3SLN(6su), green lines),*
*and 3) a human-like receptor (α2,6-sialyllactosamine or 6SLN, blue lines). Where no binding was*
*observed, the lines are omitted.*

*Molecular basis of preference for sulphated and non-sulphated avian receptors*
We previously showed that contemporary H9N2 viruses generally have a strong preference
for sulphated avian-like receptors, a property not shared with non-H9N2 avian influenza viruses. To
explore the molecular basis of this preference, we investigated amino acid differences between
UDL1/08 and Em/R66, which do and do not show this preference respectively. Generally, these
viruses had few differences near the RBS, though they did differ at positions 190, 226, and 227.
Mutants were tested with individual reciprocal substitutions at positions 190, 226, and 227 and
combinations thereof.

Substituting all three variable RBS residues (190/226/227) led to an approximate switch of
the receptor-binding phenotypes between the two viruses. Substituting these three residues
increased UDL1/08 3SLN binding and eliminated its 6SLN binding and abolished Em/R66's 3SLN
binding while increasing its 6SLN binding (Figure 2, purple lines). In general, 190/226 reciprocal
mutants expressed similar binding phenotypes to the triple mutants (Figure 2, brown lines),
suggesting the L227I and I227L substitutions had only a modest influence on receptor-binding. The
exception to this was that Em/R66 E190A/Q226L/L227I did not show the increased binding to 6SLN
observed with Em/R66 E190A/Q226L (Figure 1E). However, in both cases the exact binding avidities
to the three receptor analogues were not completely recapitulated; both triple mutants (UDL1/08
A190E/L226Q/I227L and Em/R66 E190A/Q226L/L227I) had reduced avidity compared to their wild-
type parental viruses indicating that further substitutions are required to fully recover the receptor-
binding profile of the donor viruses. Nonetheless, positions 190, 226, and to a lesser extent, 227

appeared to be the primary determinants of variation in receptor-binding preference phenotypes
of UDL1/08 and Em/R66s (Table 1).

130
*Figure 2 - Receptor binding profiles of reciprocal UDL1/08-Em/R66 mutants. Biolayer*
*interferometry was used to determine the receptor binding profiles mutant UDL1/08 (Panels A-C,*
*UDL1/08 A190E could not be rescued) and Em/R66 viruses (panels D-F). Binding was measured to 3*
*receptor analogues: Neu5Ac α2,3 Gal β1-4(6-HSO3)GlcNAc (3SLN(6su), Panels A,D), α2,6-*
*sialyllactosamine (6SLN, Panels B,E) and α2,3-sialyllactosamine (3SLN, Panels C,F). Dashed black*
*and red lines show wild-type UDL1/08 and Em/R66 binding, respectively.*

Further investigating the contribution of individual RBS residues, position 190 exerted a
strong influence on the preference for sulphated or non-sulphated avian-like receptors; viruses with
A190 showed increased binding to 3SLN(6su) and decreased binding to non-sulphated 3SLN
analogues while those with E190 showed the opposite (Figure 2A,C,D,F). This is potentially due to a
charge repulsion between the negatively charged side-chain of glutamate and the sulphate group
of sulphated-3SLN. Additionally, viruses containing A190 generally retained or had enhanced
human-like 6SLN binding as seen in UDL1/08 wild-type and the mutants Em/R66 E190A and Em/R66
E190A/Q226L (Figure 2B,E). This was also exemplified in the differences in binding between
UDL1/08 mutants: A190E/L226Q led to a loss of 6SLN binding while L226Q alone did not (Figure 1B
– mint and brown lines).

| Residue (H3 no) | Residue (H9 no) | WT Virus | Substitution | Receptor | |
|---|---|---|---|---|---|
| | | | | 3SLN(6su) | 3SLN |
| 190 | 180 | UDL1/08 | A->E | DNR | DNR |
| | | Em/R66 | E->A | + + | - - |
| 190/226 | 180/216 | UDL1/08 | A->E, L->Q | - - | + |
| | | Em/R66 | E->A, Q->L | = | - - - |
| 190/226/227 | 180/216/217 | UDL1/08 | A->E, L->Q, I->L | - - | + |
| | | Em/R66 | E->A, Q->L, L->I | - | - - - |

**Table 1. Haemagglutinin amino acid differences modulating preference for sulphated and non-sulphated avian**
**receptor.** [a] '=' indicates <2 fold difference, '+/- ' indicates 2-10 fold increase/decrease, '++/- -' indicates 10-100-fold
increase/decrease, '+++/- - -' indicates >100-fold increase/decrease in binding relative to the wild type virus. 'null'
indicates no difference was able to be seen because no binding to this analogue was detected with or without the
substitution. 'DNR' indicates the virus was unable to be rescued. '/' indicates a mutant was not made in that virus
background.

At position 226, we determined that substitutions were exerting an effect on receptor-
binding avidity, consistent with the results of our previous work with red blood cell based avidity
assays [25]. Viruses with Q226 bound with higher avidity, compared with L226, to each of the three
analogues tested (Figure 2A-F). Additionally, Q226 favoured 3SLN binding as can be seen by the
mutant UDL1/08 L226Q and the difference in avidity between Em/R66 E190A/Q226L and E190A
alone (Figure 2C,F – blue and brown lines). In the background of Em/R66, Q226L showed a large
increase in 6SLN binding (Figure 2E – mint line), however a matching reduction in 6SLN binding by
UDL1/08 L226Q was not seen (Figure 2B), indicating this is probably dependent on the context of
the other amino acids in the H9 HA RBS as we had previously predicted [21].

Finally, substitutions at position 227 were identified as playing a minor role in modulation of
avidity, though did not change receptor preference or regulate complete gain or loss of binding to
any analogue. Relative to the associated parental virus, UDL1/08 I227L showed higher avidity to all
receptors while Em/R66 L227I showed lower avidity (Figure 2, orange lines), consistent with
previous inferences made from indirect measurements of avidity [25].

*Molecular basis of preference for the human receptor*
To investigate the molecular basis of human-like receptor-binding as seen in some H9N2
viruses, a further reciprocal library was generated between UDL1/08 and HK/33982, a virus isolated
from a human case of H9N2. UDL1/08 naturally binds to sulphated 3SLN with slight binding to 6SLN,
while HK/33982 binds strongly to the human-like receptor 6SLN with moderate binding to both
sulphated and non-sulphated 3SLN (Figure 1A, C). The amino acids at the three residues shown
above to largely determine preference for sulphated vs non-sulphated (190, 226, and 227) also vary
between these two viruses. In addition to these three positions, reciprocal mutations were
introduced at positions 188 and 193 (178 and 183 in H9 numbering) on the basis that these residues
are located next to the binding site and vary between UDL1/08 and HK/33982.

Introducing reciprocal mutations at the three residues (190/226/227) shown above to
modulate preference for the sulphated vs non-sulphated avian receptor also influenced receptor
preference in this case. Introducing the substitutions A190D, L226Q and I227Q from HK/33982 into
UDL1/08 led to a loss of most of its 3SLN(6su) binding and a slight gain in 6SLN binding (Figure 2A,
B, purple lines), however this binding phenotype did not resemble that of HK/33982. This suggests
additional substitutions are required for a full gain of the human-adapted receptor-binding profile.
The reciprocal triple mutant was unable to be recovered in the HK/33982 background, indicating
additional residues must play a role in stabilising residues 190, 226 and 227 between these two
viruses (Table S1).

Introduction of reciprocal substitutions at residues 190 and 227 together indicated that
these positions likely play an important role in modulation of preference between the sulphated
avian and human-like receptors. UDL1/08 with A190D/I227Q showed slightly decreased binding to
3SLN(6su) and appreciably increased 6SLN binding (Figure 3A,B – grey lines). HK/33982
D190A/Q227I had reduced receptor-binding to 6SLN and increased binding to sulphated 3SLN,
creating an approximation of the receptor-binding preference of wild-type UDL1/08 (Figure 3D,E –
grey lines). From this mutant library it is clear that switches at position 226 create an incompatibility
in both virus backgrounds, with several viruses losing nearly all receptor-binding (UDL1/08
A190D/L226Q and UDL1/08 A190D/L226Q/I227Q) or infectious virus unable to be rescued entirely
(HK/33982 D190A/Q226L and HK/33982 D190A/Q226L/Q227I) (Table S1).

***Figure 3. Receptor binding profiles of reciprocal UDL1/08-HK/33982 mutants*** *Biolayer interferometry was used to determine the receptor binding profiles mutant UDL1/08 (Panels A-C) and HK/33982 viruses (panels D-F). Binding was measured to 3 receptor analogues: Neu5Ac α2,3 Gal β1-4(6-HSO3)GlcNAc (3SLN(6su), Panels A,D), α2,6-sialyllactosamine (6SLN, Panels B,E) and α2,3-sialyllactosamine (3SLN, Panels C,F). Dashed black and green lines show wild-type UDL1/08 and HK/33982 binding, respectively.*

The effects of single amino acid substitutions made between UDL1/08 and HK/33982 were also evaluated. Amino acid substitutions at residue 190 in the single reciprocal mutants of UDL1/08 and HK/33982 showed potential incompatibilities. UDL1/08 A190D showed a mixed population also including D190G while HK/33982 D190A gained the additional substitution G225D, as previously described [25]; HK/33982 with the G225D substitution alone was also unable to produce infectious virus (Table S1). It is, therefore, hard to draw conclusions from these viruses upon the influence of these mutations in isolation. HK/33982 D190A/G225D did appear to increase 6SLN binding and 3SLN(6su) binding while decreasing 3SLN binding (similar to Em/R66 E190A) (Figure 3D-F – blue lines). UDL1/08 A190D/G showed no difference in binding compared to wild-type UDL1/08 (Figure 2A-C – blue lines), however it is difficult to interpret these data, given the mixed population.

# Genetic basis of H9N2 receptor-binding variability

| Residue (H3 no) | Residue (H9 no) | WT Virus | Substitution | Avian receptor | | Human-like receptor |
|---|---|---|---|---|---|---|
| | | | | 3SLN(6su) | 3SLN | 6SLN |
| 193 | 183 | UDL1/08 | N->T | = | null | + |
| | | HK/33982 | T->N | + | = | = |
| 196 | 186 | UDL1/08 | T->K | - - | null | + + |
| 198 | 188 | UDL1/08 | T->A | - - | null | + |
| 226 | 216 | Em/R66 | Q->L | - | - - - | + + + |
| 227 | 217 | UDL1/08 | I->M | = | null | + + |
| | | UDL1/08 | I->Q | = | null | + + |
| | | HK/33982 | Q->I | - - | - - - | - |
| 190/227 | 180/217 | UDL1/08 | A->D, I->Q | - - - | null | + + |
| | | HK/33982 | D->A, Q->I | + + | - - | - - |
| 190/226/227 | 180/216/217 | UDL1/08 | A->D, L->Q, I->Q | - - - | null | = |
| | | HK/33982 | D->A, Q->L, Q->I | DNR | DNR | DNR |

**Table 2. Haemagglutinin amino acid differences modulating preference for avian and human-like receptors.** [a] '=' indicates <2 fold difference, '+/-' indicates 2-10 fold increase/decrease, '++/- -' indicates 10-100-fold increase/decrease, '+++/- - -' indicates >100-fold increase/decrease in binding relative to the wild type virus. 'null' indicates no difference was able to be seen because no binding to this analogue was detected with or without the substitution. 'DNR' indicates the virus was unable to be rescued. '/' indicates a mutant was not made in that virus background.

Similar to the reciprocal mutants of UDL1/08 and Em/R66, amino acid substitutions at residue 226 between UDL1/08 and HK/33982 displayed a clear avidity effect, with viruses possessing 226L appearing to show higher avidity compared with 226Q. HK/33982 with Q226L gained the additional compensatory substitution H184E (Table S1); HK/33982 with H184E alone showed a large reduction in avidity to all receptor analogues, whereas HK/33982 with H184E/Q226L increased avidity to both 6SLN and 3SLN(6su) relative to H184E alone, suggesting Q226L increased binding avidity (Figure 3D-F – purple and mint lines). The mutant HK/33982 H184E/Q226L also showed a strong preference for 6SLN.

At position 227, the introduction of I227Q in the UDL1/08 background resulted in a large increase in 6SLN binding, a drop in binding to UDL1/08's preferred receptor 3SLN(6su), and did not alter binding to 3SLN (Figure 3A-C, orange lines). The reciprocal substitution, HK/33982 Q227I, showed a general avidity effect with lower binding to all analogues (Figure 3D-F – orange line), however when introduced alongside D190A, the impact of Q227I in the HK/33982 D190A/Q227I double mutant facilitated reduced 6SLN binding but slightly increased 3SLN(6su), relative to the parental wild-type and the D190A(+G225D) mutant (Figure 3D-F – grey and orange lines).

Reciprocal substitutions at position 188 were found to influence receptor-binding while those at position 193 were not. These residues have not previously been described as affecting receptor-binding in H9 HA, though they are at the edge of the RBS and 193 has been described as playing a vital role in the modulation of binding of sulphated 3SLN by H5 and H7 HA [26, 27]. Viruses with Y188 showed higher avidity relative to viruses with D188, regardless of the virus background (Figure 3 – red lines). Additionally, HK/33982 Y188D appeared to have minor effects on specific receptor analogue preference; this substitution reduced binding for the 3SLN analogues to a greater extent than the 6SLN analogue. Amino acid swaps at position 193 showed a very minor, non-reciprocal receptor preference effect (Figure 3), consistent with our previous estimates that swaps of residue 193 between UDL1/08 and HK/33982 would only have a minimal receptor avidity effect [25].

*Molecular basis of variation in receptor-binding avidity*

In addition to showing preference for different receptors, influenza viruses vary in receptor-
binding avidity. In several previous studies we have inferred that escape mutants with the largest
impact of polyclonal antisera binding may be driven by avidity effects [20, 25, 28]. To further
investigate the genetic basis of variation in receptor-binding avidity we constructed a library of
mutants in the UDL1/08 background and assessed their receptor-binding phenotypes. To
complement this, we analysed a large dataset of haemagglutination inhibition (HI) titres and HA
sequences from natural H9N2 viruses to identify amino acid variants correlating with avidity effects
apparent in the measured HI titres.

Testing of UDL1/08 mutants showed several to exhibit a general decrease in avidity to the
tested analogues including K137I, F147L, K157T, N158D, N193D, and G225D (Figure 4A,B). Only a
single substitution, T189N, showed a negligible effect on receptor-binding (Figure 4C,D - mint lines).
Two substitutions, T135K and T192R, appeared to facilitate a general increase in avidity (i.e.
increased binding to all receptor analogues tested) in a similar manner to UDL1/08 I227L and D188Y
from the reciprocal mutant libraries (Figure 4C,D, 2A-C and 3A-C). R82G showed a reduction in
3SLN(6su) binding while not having an effect on 6SLN binding (Figure 4E,F – blue lines). Finally, one
group of mutants showed changes in receptor-binding preference with increases in 'human-like'
6SLN binding relative to 'avian-like' 3SLN(6su): I227M showed a large increase in 6SLN binding
without changing sulphated 3SLN binding (orange lines); G159K showed a modest increase in avidity
to sulphated 3SLN but a much larger relative increase in 6SLN binding (purple lines); T196I, T196K,
and T198A all decreased sulphated 3SLN binding while increasing 6SLN binding to varying degrees
(Figure 4E,F – light red, dark red and light brown lines). This last group of mutations represent single
amino acid changes that could act as markers for viruses with greater zoonotic potential, along with
the previously described Q226L (in the background of Em/R66 and HK/33982), and I227Q in a
UDL1/08-like background (Figure 2E, 3B,E).

**Figure 4. Receptor binding modulation by non-reciprocal mutations in the background of UDL1/08.** *Biolayer interferometry was used to determine the receptor binding profiles of each mutant virus. Binding was measured to 3 receptor analogues, α2,6-sialyllactosamine (6SLN), α2,3-sialyllactosamine (3SLN) and Neu5Ac α2,3 Gal β1-4(6-HSO3)GlcNAc (3SLN(6su)). No viruses had any detectable 3SLN binding. Dashed black lines show wild-type UDL1/08 binding. Panels indicate binding by mutants that show an avidity reduction (A,B), an increase in avidity (C,D) and changes in receptor binding preference (E,F). No mutants had any detectable binding to 3SLN.*

In general, amino acid replacements that increased positive charge in the HA head domain tended to cause an increase in receptor-binding avidity while the opposite was true for substitutions that increased negative charge. In Figure 5A, the impact of substitutions introduced to the UDL1/08 backbone on avidity for its preferred receptor, 3SLN(6su), is plotted by the change in net charge of the HA head domain. The proximity of the residues at which substitutions were introduced to the RBS is shown in Figure 5B.

***Figure 5 - Mapping of residues tested for receptor binding changes and correlation between charge and 3SLN(6su) binding.*** *A) The fold-increase in binding to 3SLN(6su) is plotted along with the introduced change in net charge difference. Wild type UDL1/08 binding to 3SLN(6su) is set to 1 with relative binding calculated as described previously [21]. Annotated P values calculated using a non-parametric Mann-Whitney U test – lines indicate mean fold increase. B) Residues which effect receptor binding preference (i.e. effect binding to different analogues in different ways) shown in red, residues that affect receptor binding avidity across analogues shown in orange. Residue that when mutated have no effect on receptor binding shown in green. LSTa, shown in yellow, is an α2,3-linked (avian-like) receptor analogue. H3 HA numbering used throughout. Figure made using structure PDBID:1JSH [41], made using PyMol [42].*

To investigate variation in avidity among natural H9N2 viruses, we modelled variation in HI titres for a dataset of large number of viruses covering all major H9 lineages. In addition to measuring antigenic similarity of influenza viruses, HI titres are influenced by varying avidity. Viruses that bind receptors on the surface of red blood cells used in the assay with higher avidity require more antibodies to inhibit haemagglutination manifesting as a tendency towards lower HI titres regardless of antigenic relationships to the antisera used, and vice-versa. To identify amino acid variants correlating with such variation in titres, we adapted a model we previously developed to explore antigenic variation in both human and avian influenza viruses, identifying several substitutions predicted to cause antigenic variation and validating these using mutagenesis [25, 29].

Variation in HI titres resulting both from antigenic differences and from variation in virus avidity was mapped to branches of the H9 HA phylogenetic tree as previously described [25]. To explore the genetic basis of variation caused by differences in avidity, phylogenetic terms representing branches leading to clades of viruses with systematically higher or lower titres were replaced with terms representing amino acid identity in the test virus at each variable HA position in turn. Under a forward selection procedure, terms representing positions 190, 196 and 198 were selected (180, 186 and 188 in H9 numbering, respectively). Each of these positions are in the 190-helix, proximal to the RBS. Position 190 has already been shown to play a role in receptor-binding in this and other studies [18, 20], and we see the effect of 196 and 198 on receptor-binding preference in this study (Figure 4E,F), further validating this modelling approach (Table 3).

## Genetic basis of H9N2 receptor-binding variability

| Residue (H3 no) | Residue (H9 no) | Distinct amino acid residues[b] | Number of HI titres | Effect on 3SLN(6su) binding |
|---|---|---|---|---|
| 190 | 180 | A | 1,145 | |
| | | T | 309 | 50x increase |
| | | V | 494 | 1400x increase |
| 196 | 186 | T | 1,912 | |
| | | I | 331 | 41x decrease |
| | | K | 296 | 7x decrease |
| 198 | 188 | T | 1,913 | |
| | | A | 351 | 41x decrease |

**Table 3. Amino acid residues predicted to explain variation in HI titres as a result differences in receptor-**
**binding avidity.** Effect on 3SLN(6su) binding is shown relative to most common amino acid in meta-analysis
dataset: 190A, 196T, and 198T.

*Sequence-based prediction of receptor-binding preference of H9N2 viruses*
Finally, we extrapolated the results of this study to predict the receptor-binding preferences
of circulating H9N2 viruses on the basis of amino acid identity at positions 190, 226 and 227. Viruses
were predicted as possessing one of three receptor-binding phenotypes (Figure 6): 1) a strong
preference for sulphated avian-like receptors, the canonical chicken-adapted H9N2 virus phenotype
(shown in green and represented by UDL1/08), 2) a receptor-binding phenotype more similar to
chicken-adapted H5Nx or H7N1 viruses and binds both sulphated and non-sulphated avian-like
receptors (shown in red and represented by Em/R66), or 3) an preference for the human-like
receptor with concurrent binding to avian receptors which we hypothesise may be an adaptation to
minor poultry (shown in blue and represented by HK/33982). Viruses were predicted to exhibit
preference for the sulphated avian-like receptor if at positions 190-226-227 they possessed either
A, T, or V at 190, L at 226 and either I, L, M, or Q at 227, or if they possessed the motifs A-Q-I, A-Q-
T, or I-Q-F. Viruses possessing either E-Q-L or E-Q-Q were predicted to bind both sulphated and non-
sulphated avian-like receptors. Viruses possessing A, D, T, or V at 190 and Q-Q at 226-227 were
predicted to show preference for the human-like receptor, as were viruses possessing the motif E-
L-Q. Predictions for motifs without direct evidence but made based on combinations of other motifs
results are shown in lighter shades to indicate reduced confidence (details in Materials and
Methods), while predicted receptor preference was considered unknown for viruses with
incomplete sequence information at positions 190, 226 and 227 or alternative motifs. The
substitutions at positions 196 and 198 that increased relative 6SLN binding were not included at this
stage as it is not known with which other substitutions must occur to elevate 6SLN binding above
binding for the avian receptor.

***Figure 6. Sequence variation and predicted receptor preference across H9N2 phylogeny****. HA phylogeny for 2,440 H9N2 viruses generated using and rooted with a sequence from H9-like bat influenza virus (starred). Amino acid identity at positions 190, 226 and 217 is shown by colour, grouped by side-chain property, according to the legend. Predicted receptor preference for α2,6-sialyllactosamine (6SLN), α2,3-sialyllactosamine (3SLN) and Neu5Ac α2,3 Gal β1-4(6-HSO3)GlcNAc (3SLN(6su)) is shown in blue, red and green respectively, with lighter shades indicating reduced confidence. Predicted receptor preference is based on amino acid identity at positions 190, 226 and 227 based on extrapolation of the amino acid sequence of viruses tested using BLI and the results of mutagenesis experiments.*

Amino acid sequence at positions 190, 226, and 227 and the resulting prediction of receptor preference is shown across a phylogeny constructed using all available H9N2 HA sequences in Figure 6. Almost all viruses in Y439-like lineage, prevalent in wild birds and poultry in Korea, as well as a few viruses in the G1-Eastern sub-lineage are predicted to show an Em/R66-like preference for any avian-like receptor. The vast majority of viruses in the chicken-adapted BJ94 and G1 lineages are predicted to maintain a canonical H9N2 sulphated avian receptor preference. A significant number of viruses belonging to the G1-Eastern sub-lineage, prevalent in minor poultry in China, are predicted to show a preference for human-like receptors, as are a number of viruses interspersed within a clade of viruses belonging to the BJ94 lineage.

## Discussion

Receptor-binding is an important determinant of host specificity and modification of receptor binding properties is often a critical step in cross-species virus transmission. In this study we have comprehensively investigated residues in and around the receptor binding site of HA from H9N2 viruses for their ability to influence receptor preference and avidity. We have shown that different combinations of the residues 190, 226 and 227 account for much, but not all, of the

variability in H9N2 receptor binding between viruses representative of the major binding
phenotypes. Furthermore, we described several other residues that have strong influences on H9N2
receptor binding preference including positions 159, 188, 193, 196 and 198. We propose a model
whereby residues in the influenza HA1 that don't directly coordinate receptor binding can play a
delocalised avidity role through modulating the charge of the head domain with increases in positive
charge giving a non-specific increase in avidity and vice versa. We hypothesise this effect may be
exaggerated for H9N2 viruses where sulphated, siaylated glycans appear to be the preferred
receptor owing to the greater amount of negative charge in these receptors (compared to non-
sulphated, siaylated glycans). Finally, we have applied the results of this study to try and predict the
receptor binding phenotypes of different circulating H9N2 viruses as a way of predicting strains that
may have a heightened zoonotic potential.

Several of the residues identified in this study have been previously described as directly or
indirectly affecting H9 receptor binding, including position 155, 190, 225, 226 and 227 [13, 17-20,
30, 31]. In previous studies, generally only one or two of these residues are measured in isolation
for their receptor binding effect or are tested in a non- or only semi-quantitative manner. Here we
perform a comprehensive, quantitative analysis often testing changes at residues in multiple
different virus backbones and in multiple combinations to investigate strain-specific and
compensatory effects. For example, residue 226 has been shown several times in an older G1-
Eastern lineage virus to increase binding to human-like receptors [17, 32], with a similar effect when
we investigate a contemporary virus of the same lineage (i.e. HK/33982), however in a G1-Western
backbone (i.e. UDL1/08) we find a completely different effect whereby there is a general avidity
increase and an overall increase in 6SLN-binding when a L226Q substitution is incorporated. This
highlights the importance of the overall context of receptor binding residues when looking at
influenza receptor binding mutants.

In our previous study we predicted that one of the groups of substitutions that had the
largest effect on immune escape in H9N2 viruses were substitutions that affect avidity, as has been
predicted for human influenza viruses [25, 33, 34]. Here we test a wide variety of escape mutants
that were previously shown to robustly modulate polyclonal antisera binding and show a large
number of them also modulate receptor binding avidity. This suggests that many escape mutations
that have a large influence on antigenicity may be exerting this effect through a receptor binding
avidity effect as has been shown for human influenza viruses [33, 34]. We also performed an analysis
of matched genetic and antigenic data for 330 H9N2 viruses, covering each of the major H9 lineages
[25]. Using a modified version of a model used to identify antigenic determinants, we predicted that
different residues at positions 190, 196 and 198 would explain variation in HI titres as a result of
differing avidity, as they tended to be associated with lower or higher HI titres irrespective of
antigenic relationships between the viruses and antisera being compared. Each of these three
positions are confirmed in this study to play an important role in regulation of receptor-binding
avidity. In addition to the important role for 190 in receptor preference, the substitutions T196I,
T196K and T198A all showed relative increases in human-like receptor binding. These results further
demonstrate there is a strong case for using integrated modelling approaches to reanalyse large
data sets and predict residues that affect receptor binding and antigenic phenotypes.

We present evidence suggesting that residues in the influenza HA1 that don't directly
coordinate receptor binding play a delocalised avidity role through modulating the charge of the
head domain. For mutations introduced in the UDL1/08 background and measured in binding to its
preferred receptor, 3SLN(6su), we found a significant trend in avidity change between substitutions
that introduced a negative charge and those that introduced a positive charge. In general,
substitutions observed to increase avidity tended to increase the net positive charge around the

RBS (e.g. T135K, G159K, T192R, D188Y, Em/R66 E190A, Em/R66 Q226L/E190A) whilst avidity
decreasing mutants usually decreased the net positive charge around the RBS (e.g. R82G, K137I,
K157T, N158D, N193D, G225D, HK/33982 Y188D, HK/33982 H184E). This effect is likely due to non-
specific charge interactions with the negatively charge sialic acid and we hypothesise may be more
pronounced for H9N2 viruses as the negative charge of sulphated, siaylated glycans, which appear
to be their preferred receptor, is greater due to the negatively charged sulphate group.

An important contribution of this study is the identification of several substitutions that, in
one or more of our virus backbones, resulted in viruses with increased or *de novo* human-like 6SLN
binding. These residues will be useful for future surveillance efforts to identify newly sequenced
viruses with elevated zoonotic potential. These particular substitutions include the previously well
characterised Q226L substitution in HK/33982 or Em/R66-like backbones (but not the contemporary
G1-Western UDL1/08-like backbone) as well as the newly characterised R82G, T135K, G159K, T196I,
T196K, T198A, I227M and I227Q substitutions in the UDL1/08-like backbone. Several of these
mutations are already commonly found in the field, further suggesting that H9N2 virus variants
naturally circulate with a likely heightened zoonotic potential.

In conclusion, we have quantified the impact of single and multiple amino acid substitutions
on receptor binding phenotypes in the context of several H9N2 viruses with varying receptor binding
preferences, identifying seven novel mutations that increase binding to the human-like receptor.
We further highlight the importance of mutations that impact receptor binding avidity. Avidity
modulation has a dramatic impact on antigenicity and an equally important role in receptor binding
phenotype, thus viruses that gain avidity enhancing mutations may present multiple challenges in
that both vaccine efficacy may be compromised, and zoonotic potential may be increased
concurrently. As well as helping better understand the molecular basis of avian influenza receptor
binding, the results generated here will help future surveillance efforts to identify viruses which may
potentially have an augmented zoonotic potential and/or greater vaccine escape potential.

## Materials and Methods

### *Ethics statement*

446 Use of embryonated eggs in this study was carried out in strict accordance with European
447 and United Kingdom Home Office regulations and the Animals (Scientific Procedures) Act 1986
448 Amendment Regulations, 2012. These studies were carried out under the United Kingdom Home
449 Office approved project license numbers 30/2683 and 30/2952.

### *Cell lines and eggs*

451 HEK 293Ts and MDCKs were maintained in Dulbecco's modified Eagle medium (DMEM),
452 supplemented with 10% foetal calf serum, 37°C, 5% $CO_2$. Viruses were propagated in 10-day-old
453 embryonated eggs; allantoic fluid was harvest 48 hours post-inoculation.

### *Viruses*

Throughout this study recombinant viruses, generated by standard 8 plasmid influenza
reverse genetics were used [35]. All viruses contained the named HA gene (whether wild type or
mutant), the NA of A/chicken/Pakistan/UDL-01/2008 (UDL1/08) and the remaining genes from
A/Puerto Rico/8/1934 (PR8), allowing for high viral titres from eggs. Mutant HA plasmids were
generated by site directed mutagenesis. Viruses were attempted to be rescued a minimum of three
independent times and left for 7 days post-co-culture before being determined to be un-rescuable.

*Virus sequencing*

Viruses were sequenced to confirm no reversions or additional substitutions had occurred
upon production and propagation. The HA1 region of HA was sequenced for each virus as previously
described [36].

*Virus purification*

Low speed centrifugation was initially used to remove large debris from virus containing egg
allantoic fluid. Virus particles were next pelleted by ultracentrifugation at 27,000rpm for 2 hours.
Virus pellets were subsequently homogenised by glass homogeniser, resuspended and purified with
a 30-60% sucrose gradient. The visible band containing virus was then isolated, diluted into PBS and
centrifuged for another 2 hours at 27,000rpm. The final virus pellet was then resuspended in PBS,
0.01% azide. Concentration of purified viruses was determined using a nucleoprotein ELISA as
described previously [37].

*Biolayer interferometry*

Purified virus binding to different sialylated receptor analogues was tested using an Octet
RED bio-layer interferometer (Pall ForteBio) as described previously [21]. Receptor analogues
contained 30kDa polyacrylamide backbones conjugated to 20 mol % trisaccharides and 5 mol %
biotin (Lectinity Holdings). The three analogues used in this study were α2,6-sialyllactosamine
(6SLN), α-2,3-sialyllactosamine (3SLN) and Neu5Ac α-2,3Galβ1-4(6-HSO3)GlcNAc(3SLN(6su)).
Sialoglycopolyemers were bound onto streptavidin coated biosensors (Pall ForteBio) at ranges of
concentrations from 0.01-0.55µg/ml in 10mM HEPES, pH 7.4, 150mM NaCl, 3mM EDTA, 0.005%
Tween-20 (HBS-EP). Virus was diluted to a concentration of 100pM in HBS-EP, 10µM oseltamivir
carboxylate (Roche), 10µM zanamivir (GSK). Virus association to the bound sialoglcopolymers was
measured at 20°C for 30 minutes. Virus binding curves were normalised to fractional saturation and
plotted as a function of sugar loading. Relative dissociation constants were calculated as described
previously [21, 38].

*Modelling of potential receptor binding residues*

To identify amino acid positions where substitutions correlated with differences in receptor-
binding avidity apparent in HI titres, a modelling approach previously used to identify substitutions
causing antigenic differences among influenza viruses was adapted [25, 29, 39, 40]. Following the
methodology described in the aforementioned studies, branches of the HA phylogenetic tree
correlated with variation in HI titres when the branch 1) separated test virus and antisera, 2)
descended the test virus, or 3) descended the virus used to generate antisera. These phylogenetic
terms are interpreted as being associated with changes in 1) antigenicity, 2) receptor-binding
avidity, and 3) immunogenicity, respectively. Internal branches of the phylogeny leading to clades
of two or more viruses associated with systematically higher or lower titres (numbered 2 in previous
sentences) and containing at least one virus also used as an antisera strain were removed,
effectively dropping any terms from the model that explained variation in HI associated with
differences in virus avidity. In their place, terms representing amino acid identity in the assayed virus
at each variable HA position were tested. At each position, these terms allowed for titres to vary
according to which amino acid residue the virus possessed to account for potential differences in
contributions to avidity. These position-specific terms were added to the model under a forward
selection procedure until the addition of further terms ceased to improve the model, as assessed
by likelihood ratio test (p < 0.05) with a Holm-Bonferroni correction for multiple testing. At selected
positions, effect sizes were estimated for each alternative amino acid relative to the amino acid
found most commonly in the dataset.

*Prediction of receptor binding profiles from sequence*
All available HA sequences from H9 viruses were downloaded from GIASID. A phylogenetic
tree was generated from aligned nucleotide sequences using MEGA. Receptor-binding profiles were
predicted across the phylogeny according to amino acid identify at positions 190, 226, and 227 on
the basis of the BLI results derived during this study. Viruses were predicted as possessing either: 1)
a strong preference for sulphated avian-like receptors, 2) a receptor binding phenotype more similar
to chicken-adapted H5Nx or H7N1 viruses that bind both sulphated and non-sulphated avian-like
receptors or 3) a preference for the human-like receptor with concurrent binding to avian receptors.
Viruses were predicted to exhibit preference for the sulphated avian-like receptor if at positions
190-226-227 they possessed the motifs A-L-I, A-L-L, A-L-M, A-L-Q, A-Q-I, A-Q-T, I-Q-F, T-L-I, V-L-I (or
with reduced confidence T-L-L, T-L-M, T-L-Q, V-L-L, V-L-M, or V-L-Q), for any avian-like receptor if
they possessed the motifs E-Q-L or E-Q-Q, and for the human-like receptor if they possessed A-Q-Q,
D-Q-Q, E-L-Q (or with reduced confidence T-Q-Q or V-Q-Q). Predictions made with reduced
confidence indicate that we have not tested the exact combination of amino acids but that the
prediction is consistent with other combinations barring unforeseen interactions between sites. For
viruses with incomplete sequence information at positions 190, 226 and 227 or alternative motifs,
no prediction was made.

## Funding statement

This research was supported by the Biotechnology and Biological Sciences Research Council
under grants BBS/E/00001759 (TPP), BBS/E/I/00001708 (JES), BB/L018853/1 (MI, JES),
BBS/E/I/00007031 (MI), BBS/E/I/00007035 (MI), BBS/E/I/00007038 (MI), BBS/E/I/00007039 (MI),
BB/L004828/1 (RR), BB/P004202/1 (RR) and BB/R012679/1 (MI, RR) and the Medical Research
Council under grant numbers MR/J50032X/1 (1097258) and MR/R024758/1 (WTH). The funders had
no role in study design, data collection and analysis, decision to publish or preparation of the
manuscript. The Francis Crick Institute receives its core funding from Cancer Research UK, the UK
Medical Research Council, and the Wellcome Trust.

## Competing Interests

533      The authors state they have no conflict of interest.

## Acknowledgements

535      The authors would like to thank Dr Stephen Martin of The Francis Crick institute for use of
536 his software for analysing the bio-layer interferometry results and Dr Jürgen Stech of the Friedrick
537 Loeffler Institute for the haemagglutinin reverse genetics plasmid of A/chicken/Emirates/R66/2002
538 (H9N2).

## Author contributions

540      Conceptualisation - TP, WH. Data curation, Formal analysis, funding acquisition – RR & MI
541 Investigation - TP, JES & WH. Methodology, Resources, Software- WH & RR Supervision – WB, RR &
542 MI. Writing original draft – TP, JES & WH. Writing – review and editing – All

## References

543
1.      Thuy DM, Peacock TP, Bich VT, Fabrizio T, Hoang DN, Tho ND, et al. Prevalence and diversity
of H9N2 avian influenza in chickens of Northern Vietnam, 2014. Infect Genet Evol. 2016;44:530-40.
doi: 10.1016/j.meegid.2016.06.038. PubMed PMID: 27340015.

2.      Negovetich NJ, Feeroz MM, Jones-Engel L, Walker D, Alam SM, Hasan K, et al. Live bird
markets of Bangladesh: H9N2 viruses and the near absence of highly pathogenic H5N1 influenza.
PLoS One. 2011;6(4):e19311. doi: 10.1371/journal.pone.0019311. PubMed PMID: 21541296;
PubMed Central PMCID: PMC3082571.
3.      Chen LJ, Lin XD, Guo WP, Tian JH, Wang W, Ying XH, et al. Diversity and evolution of avian
influenza viruses in live poultry markets, free-range poultry and wild wetland birds in China. J Gen
Virol. 2016;97(4):844-54. Epub 2016/01/14. doi: 10.1099/jgv.0.000399. PubMed PMID: 26758561.
4.      Peacock THP, James J, Sealy JE, Iqbal M. A Global Perspective on H9N2 Avian Influenza Virus.
Viruses. 2019;11(7). Epub 2019/07/10. doi: 10.3390/v11070620. PubMed PMID: 31284485;
PubMed Central PMCID: PMCPMC6669617.
5.      Peiris M, Yuen KY, Leung CW, Chan KH, Ip PL, Lai RW, et al. Human infection with influenza
H9N2. Lancet. 1999;354(9182):916-7. PubMed PMID: 10489954.
6.      Guo Y, Li J, Cheng X. [Discovery of men infected by avian influenza A (H9N2) virus]. Zhonghua
shi yan he lin chuang bing du xue za zhi = Zhonghua shiyan he linchuang bingduxue zazhi = Chinese
journal of experimental and clinical virology. 1999;13(2):105-8. PubMed PMID: 12569771.
7.      International Centre for Diarrhoeal Disease Research (ICDDRB). Outbreak of mild respiratory
disease caused by H5N1 and H9N2 infections among young children in Dhaka, Bangladesh. Health
and Science Bulletin: 2011.
8.      World Health Organisation. Influenza at the human-animal interface, Summary and
assessment, 3 March 2015. www.who.int: 2015 3 March 2015. Report No.
9.      Ali M, Yaqub T, Mukhtar N, Imran M, Ghafoor A, Shahid MF, et al. Avian Influenza A(H9N2)
Virus in Poultry Worker, Pakistan, 2015. Emerg Infect Dis. 2019;25(1):136-9. Epub 2018/12/19. doi:
10.3201/eid2501.180618. PubMed PMID: 30561309.
10.     Influenza at the human-animal interface, Summary and assessment, 10 April to 10 May 2019.
Report. Geneva, Switzerland: World Health Organisation, 2019.
11.     World Health Organisation. Influenza at the human-animal interface, Summary and
assessment, from 21 January to 28 February 2020. www.who.int: 2020 28 February 2020. Report
No.
12.     World Health Organisation. Influenza at the human-animal interface, Summary and
assessment, from 28 September to 25 November 2019. www.who.int: 2019 25 November 2019.
Report No.
13.     Li X, Shi J, Guo J, Deng G, Zhang Q, Wang J, et al. Genetics, receptor binding property, and
transmissibility in mammals of naturally isolated H9N2 Avian Influenza viruses. PLoS Pathog.
2014;10(11):e1004508. doi: 10.1371/journal.ppat.1004508. PubMed PMID: 25411973; PubMed
Central PMCID: PMC4239090.
14.     Uyeki TM, Chong YH, Katz JM, Lim W, Ho YY, Wang SS, et al. Lack of evidence for human-to-
human transmission of avian influenza A (H9N2) viruses in Hong Kong, China 1999. Emerg Infect Dis.
2002;8(2):154-9. doi: 10.3201/eid0802.010148. PubMed PMID: 11897066; PubMed Central PMCID:
PMC2732440.
15.     Rogers GN, Paulson JC. Receptor determinants of human and animal influenza virus isolates:
differences in receptor specificity of the H3 hemagglutinin based on species of origin. Virology.
1983;127(2):361-73. Epub 1983/06/01. doi: 10.1016/0042-6822(83)90150-2. PubMed PMID:
6868370.
16.     Pillai SP, Lee CW. Species and age related differences in the type and distribution of influenza
virus receptors in different tissues of chickens, ducks and turkeys. Virol J. 2010;7:5. Epub
2010/01/14. doi: 10.1186/1743-422X-7-5. PubMed PMID: 20067630; PubMed Central PMCID:
PMCPMC2821308.

17.    Wan H, Perez DR. Amino acid 226 in the hemagglutinin of H9N2 influenza viruses determines
cell tropism and replication in human airway epithelial cells. J Virol. 2007;81(10):5181-91. doi:
10.1128/JVI.02827-06. PubMed PMID: 17344280; PubMed Central PMCID: PMC1900221.
18.    Teng Q, Xu D, Shen W, Liu Q, Rong G, Li X, et al. A Single Mutation at Position 190 in
Hemagglutinin Enhances Binding Affinity for Human Type Sialic Acid Receptor and Replication of
H9N2 Avian Influenza Virus in Mice. J Virol. 2016. doi: 10.1128/JVI.01141-16. PubMed PMID:
27558420.
19.    Sang X, Wang A, Ding J, Kong H, Gao X, Li L, et al. Adaptation of H9N2 AIV in guinea pigs
enables efficient transmission by direct contact and inefficient transmission by respiratory droplets.
Sci Rep. 2015;5:15928. doi: 10.1038/srep15928. PubMed PMID: 26552719.
20.    Sealy JE, Yaqub T, Peacock TP, Chang P, Ermetal B, Clements A, et al. Association of Increased
Receptor-Binding Avidity of Influenza A(H9N2) Viruses with Escape from Antibody-Based Immunity
and Enhanced Zoonotic Potential. Emerg Infect Dis. 2018;25(1):63-72. Epub 2018/12/19. doi:
10.3201/eid2501.180616. PubMed PMID: 30561311.
21.    Peacock TP, Benton DJ, Sadeyen JR, Chang P, Sealy JE, Bryant JE, et al. Variability in H9N2
haemagglutinin receptor-binding preference and the pH of fusion. Emerg Microbes Infect.
2017;6(3):e11. doi: 10.1038/emi.2016.139. PubMed PMID: 28325922.
22.    Gambaryan AS, Tuzikov AB, Pazynina GV, Desheva JA, Bovin NV, Matrosovich MN, et al. 6-
sulfo sialyl Lewis X is the common receptor determinant recognized by H5, H6, H7 and H9 influenza
viruses of terrestrial poultry. Virol J. 2008;5:85. doi: 10.1186/1743-422X-5-85. PubMed PMID:
18652681; PubMed Central PMCID: PMC2515299.
23.    Xiong X, Tuzikov A, Coombs PJ, Martin SR, Walker PA, Gamblin SJ, et al. Recognition of
sulphated and fucosylated receptor sialosides by A/Vietnam/1194/2004 (H5N1) influenza virus.
Virus research. 2013;178(1):12-4. doi: 10.1016/j.virusres.2013.08.007. PubMed PMID: 24036174.
24.    Xiong X, Martin SR, Haire LF, Wharton SA, Daniels RS, Bennett MS, et al. Receptor binding by
an H7N9 influenza virus from humans. Nature. 2013;499(7459):496-9. doi: 10.1038/nature12372.
PubMed PMID: 23787694.
25.    Peacock TP, Harvey WT, Sadeyen JR, Reeve R, Iqbal M. The molecular basis of antigenic
variation among A(H9N2) avian influenza viruses. Emerg Microbes Infect. 2018;7(1):176. Epub
2018/11/08. doi: 10.1038/s41426-018-0178-y. PubMed PMID: 30401826; PubMed Central PMCID:
PMCPMC6220119.
26.    Gambaryan AS, Matrosovich TY, Philipp J, Munster VJ, Fouchier RA, Cattoli G, et al. Receptor-
binding profiles of H7 subtype influenza viruses in different host species. J Virol. 2012;86(8):4370-9.
doi: 10.1128/JVI.06959-11. PubMed PMID: 22345462; PubMed Central PMCID: PMC3318648.
27.    Gambaryan A, Tuzikov A, Pazynina G, Bovin N, Balish A, Klimov A. Evolution of the receptor
binding phenotype of influenza A (H5) viruses. Virology. 2006;344(2):432-8. Epub 2005/10/18. doi:
10.1016/j.virol.2005.08.035. PubMed PMID: 16226289.
28.    Peacock TP, Benton DJ, James J, Sadeyen JR, Chang P, Sealy JE, et al. Immune escape variants
of H9N2 influenza viruses containing deletions at the haemagglutinin receptor binding site retain
fitness in vivo and display enhanced zoonotic characteristics. J Virol. 2017. doi: 10.1128/JVI.00218-
17. PubMed PMID: 28468875.
29.    Harvey WT, Benton DJ, Gregory V, Hall JP, Daniels RS, Bedford T, et al. Identification of Low-
and High-Impact Hemagglutinin Amino Acid Substitutions That Drive Antigenic Drift of Influenza
A(H1N1) Viruses. PLoS Pathog. 2016;12(4):e1005526. doi: 10.1371/journal.ppat.1005526. PubMed
PMID: 27057693; PubMed Central PMCID: PMCPMC4825936.
30.    Mancera Gracia JC, Van den Hoecke S, Saelens X, Van Reeth K. Effect of serial pig passages
on the adaptation of an avian H9N2 influenza virus to swine. PLoS One. 2017;12(4):e0175267. Epub
2017/04/07. doi: 10.1371/journal.pone.0175267. PubMed PMID: 28384328; PubMed Central
PMCID: PMCPMC5383288.

31.    Kaverin NV, Rudneva IA, Ilyushina NA, Lipatov AS, Krauss S, Webster RG. Structural
differences among hemagglutinins of influenza A virus subtypes are reflected in their antigenic
architecture: analysis of H9 escape mutants. J Virol. 2004;78(1):240-9. PubMed PMID: 14671105;
PubMed Central PMCID: PMC303415.

32.    Obadan AO, Santos J, Ferreri L, Thompson AJ, Carnaccini S, Geiger G, et al. Flexibility in vitro
of amino acid 226 in the receptor-binding site of an H9 subtype influenza A virus and its effect in
vivo on virus replication, tropism, and transmission. J Virol. 2018. Epub 2018/12/21. doi:
10.1128/JVI.02011-18. PubMed PMID: 30567980.

33.    Hensley SE, Das SR, Bailey AL, Schmidt LM, Hickman HD, Jayaraman A, et al. Hemagglutinin
receptor binding avidity drives influenza A virus antigenic drift. Science. 2009;326(5953):734-6. doi:
10.1126/science.1178258. PubMed PMID: 19900932; PubMed Central PMCID: PMC2784927.

34.    Li Y, Bostick DL, Sullivan CB, Myers JL, Griesemer SB, St George K, et al. Single Hemagglutinin
Mutations that Alter Both Antigenicity and Receptor Binding Avidity Influence Influenza Virus
Antigenic Clustering. J Virol. 2013;87(17):9904-10. doi: 10.1128/JVI.01023-13. PubMed PMID:
23824816.

35.    Hoffmann E, Neumann G, Kawaoka Y, Hobom G, Webster RG. A DNA transfection system for
generation of influenza A virus from eight plasmids. Proc Natl Acad Sci U S A. 2000;97(11):6108-13.
doi: 10.1073/pnas.100133697. PubMed PMID: 10801978; PubMed Central PMCID: PMC18566.

36.    Peacock T, Reddy K, James J, Adamiak B, Barclay W, Shelton H, et al. Antigenic mapping of
an H9N2 avian influenza virus reveals two discrete antigenic sites and a novel mechanism of immune
escape. Sci Rep. 2016;6:18745. doi: 10.1038/srep18745. PubMed PMID: 26738561; PubMed Central
PMCID: PMC4704030.

37.    Lin YP, Xiong X, Wharton SA, Martin SR, Coombs PJ, Vachieri SG, et al. Evolution of the
receptor binding properties of the influenza A(H3N2) hemagglutinin. Proc Natl Acad Sci U S A.
2012;109(52):21474-9. doi: 10.1073/pnas.1218841110. PubMed PMID: 23236176; PubMed Central
PMCID: PMC3535595.

38.    Xiong X, Coombs PJ, Martin SR, Liu J, Xiao H, McCauley JW, et al. Receptor binding by a ferret-
transmissible H5 avian influenza virus. Nature. 2013;497(7449):392-6. doi: 10.1038/nature12144.
PubMed PMID: 23615615.

39.    Reeve R, Borley DW, Maree FF, Upadhyaya S, Lukhwareni A, Esterhuysen JJ, et al. Tracking
the Antigenic Evolution of Foot-and-Mouth Disease Virus. PLoS One. 2016;11(7):e0159360. doi:
10.1371/journal.pone.0159360.    PubMed    PMID:    27448206;    PubMed    Central    PMCID:
PMCPMC4957747.

40.    Reeve R, Blignaut B, Esterhuysen JJ, Opperman P, Matthews L, Fry EE, et al. Sequence-based
prediction for vaccine strain selection and identification of antigenic variability in foot-and-mouth
disease virus. PLoS Comput Biol. 2010;6(12):e1001027. doi: 10.1371/journal.pcbi.1001027. PubMed
PMID: 21151576; PubMed Central PMCID: PMCPMC3000348.

41.    Ha Y, Stevens DJ, Skehel JJ, Wiley DC. H5 avian and H9 swine influenza virus haemagglutinin
structures: possible origin of influenza subtypes. The EMBO journal. 2002;21(5):865-75. doi:
10.1093/emboj/21.5.865. PubMed PMID: 11867515; PubMed Central PMCID: PMC125880.

42.    Schrodinger, LLC. The PyMOL Molecular Graphics System, Version 1.3r1. 2010.