# Arthropod ILGF, Relaxin and Gonadulin, putative orthologs of *Drosophila* insulin-like peptides 6, 7 and 8, likely originated from an ancient gene triplication

Jan A. Veenstra[1],

[1] INCIA UMR 5287 CNRS, University of Bordeaux, Bordeaux, Pessac, France

Corresponding Author:
Jan A. Veenstra[1]
INCIA UMR 5287 CNRS, Université de Bordeaux, allée Geoffroy St Hillaire, CS 50023, 33 615
Pessac Cedex, France
Email address: jan-adrianus.veenstra@u-bordeaux.fr

## Abstract

**Background.** Insects have several genes coding for insulin-like peptides and they have been

particularly well studied in *Drosophila.* Some of these hormones function as growth hormones

and are produced by the fat body and the brain. These act through a typical insulin receptor

tyrosine kinase. Two other *Drosophila* insulin-like hormones are either known or suspected to act

through a G-protein coupled receptor. Although insulin-related peptides are known from other

insect species, *Drosophila* insulin-like peptide 8, one that uses a G-protein coupled receptor, has

so far only been identified from *Drosophila* and other flies. However, its receptor is widespread

within arthropods and hence it should have orthologs. Such putative orthologs were recently

identified in decapods and have been called gonadulins.

**Methodology**. In an effort to identify gonadulins in other arthropods public genome assemblies

and short-read archives from insects and other arthropods were explored for the presence of

genes and transcripts coding insulin-like peptides and their putative receptors.

**Results.** Gonadulins were detected in a number of arthropods. In those species for which

transcriptome data from the gonads is available insect gonadulin genes are expressed in the

ovaries and at least in some species also in the testes. In some insects differences in gonadulin

expression in the ovary between actively reproducing and non-reproducing females differs more

than 100-fold. Putative orthologs of *Drosophila* ilp 6 were also identified. In several non-

Dipteran insects these peptides have C-terminally extensions that are alternatively spliced. The

predicted peptides have been called arthropod insulin-like growth factors. In cockroaches,

termites and stick insects genes coding for the arthropod insulin-like growth factors, gonadulin

and relaxin, a third insulin-like peptide, are encoded by genes that are next to one another

suggesting that they are the result of a local gene triplication. Such a close chromosomal
association was also found for the arthropod insulin-like growth factor and gonadulin genes in
spiders. Phylogenetic tree analysis of the typical insulin receptor tyrosine kinases from insects,
decapods and chelicerates shows that the insulin signaling pathway evolved differently in these
three groups. The G-protein coupled receptors that are related to the *Drosophila* ilp 8 receptor
similarly show significant differences between those groups.

**Conclusion.** A local gene triplication in an early ancestor likely yielded three genes coding
gonadulin, arthropod insulin-like growth factor and relaxin. Orthologs of these genes are now
commonly present in arthropods and almost certainly include the *Drosophila* insulin-like peptides
6, 7 and 8.

## Introduction

50
Insulin may well be the best studied and perhaps the best known hormone, due to its essential
role in the regulation of glucose homeostasis and its effective and widespread use to treat
diabetes. Insulin is related to a number of other hormones with different functions, such as
insulin-like growth factor, relaxin and Ins3. One of the interesting aspects of these hormones is
their use of two structurally very different receptors, receptor tyrosine kinases (RTKs) and
leucine-rich repeat G-protein coupled receptors (LGRs). Thus, whereas insulin and insulin-like
growth factors (ILGFs) act through an RTK, relaxin and Ins3 use an LGR for signal transduction.
An intriguing question remains as to how this switch was made from one type of receptor to
another, or alternatively whether the ancestral insulin used perhaps both types of receptors and
during evolution its descendants became specific ligands for only one of the two receptors.
Like other hormones and neuropeptides, insulin was already present in the ancestral bilaterian
that gave rise to both protostomes and deuterostomes. The first indication that an insulin-like
peptide (ilp) might exist in protostomes was the observation that insulin enhances cell
differentiation in cultured *Drosophila* cells (Seecof & Dewhurst, 1974). The identification of one
ilp in the silkworm *Bombyx mori* that can break diapause (Nagasawa et al., 1984, 1986) and
another one in neuroendocrine cells known to produce a growth hormone in the pond snail
*Lymnaea stagnalis* (Smit et al., 1988) reinforced the hypothesis that insulin may functions as
growth hormones in protostomes. Since then a large variety of invertebrates has yielded a still
larger variety of ilps (*e.g.* Murphy & Hu, 2013; Mizoguchi & Okamoto, 2013; Nässel & Vanden
Broeck, 2016; Yu, Han & Liu, 2020).

In insects the ilps of *Drosophila* and the silkworm *Bombyx mori* have been extensively studied

and these hormones are best known in fruit fly due to the genetic power that can be employed in

this species. There are eight ilp genes in *Drosophila melanogaster*, which are referred to as

*Drosophila* ilps 1-8. *Drosophila* ilps 1, 2, 3 and 5 are co-expressed in a single cell type of

neuroendocrine cells of the brain (Brogiolo et al., 2001; Grönke et al., 2010). Of these ilp 2 seems

to be the most important and it also seems to be expressed exclusively or predominantly in these

brain neuroendocrine cells. *Drosophila* ilps 3 and 5 are also expressed in other tissues, *e.g.* ilp 3

is expressed in midgut muscle of both larvae and adults where its expression stimulates midgut

growth in response to feeding (O'Brien et al., 2011). *Drosophila* ilp 1 has been shown to be

expressed in the brain neuroendocrine cells, but its expression is largely limited to stages when

the animal does not feed, *i.e.* metamorphosis and diapause (Liu et al., 2016). The expression of

dilp 4 seems limited to the embryonic stage, while ilp 6 is expressed predominantly if not

exclusively by the fat body (Slaidina et al., 2009; Okamoto et al., 2009b). All these ilps are

believed to activate the single *Drosophila* insulin RTK, while *Drosophila* ilps 7 and 8 are either

known (ilp 8) or suspected (ilp 7) to activate *Drosophila* LGRs 3 and 4 respectively (Vallejo et

al., 2015; Gontijo & Garelli, 2018; Veenstra et al., 2012). *Drosophila* ilp 7 is expressed by

neurons in the abdominal neuromeres in a sex specific manner (Miguel-Aliaga, Thor & Gould,

2008; Yang et al., 2008; Castellanos, Tang & Allan, 2013), while ilp 8 is expressed by the

imaginal disks as well as the ovary and testes as shown by flyatlas (Chintapalli, Wang & Dow,

2007; Gontijo & Garelli, 2018).

The primary amino acid sequences of the *Drosophila* ilps vary considerably and this is also the

case in other arthropod species that have multiple genes coding insulin-related peptides. There is

not only large sequence variability within a species, but also between species. Only when species

are relatively closely related is it possible to reliably identify orthologous genes in different

species. However, while in most insects the A- and B-chains have thus quite variable amino acid

sequences, this not the case for orthologs of *Drosophila* ilp 7. The strong conversation of the

primary amino acid sequence of these peptides allows for easy identification of its orthologs, not

only in other insect species, but also in other protostomes like various mollusks and even in some

deuterostomes (Veenstra et al., 2012). The strongly conserved primary amino acid sequence of

these peptides suggests that it interacts with another receptor than the other ilps, perhaps in

addition to the RTK. As some ilps act through a G-protein coupled receptor (GPCR), it seemed a

distinct possibility that *Drosophila* ilp 7 and their orthologs might also stimulate a GPCR.

Interestingly, genes coding LGR4 and its orthologs are present in the same genomes as those that

have genes coding orthologs of *Drosophila* ilp 7. This holds not only for insects, but also other

arthropods, mollusks and even some basal deuterostomes. Every genome that has a *Drosophila*

ilp 7 ortholog also has a LGR4 ortholog and *vice versa* (Veenstra et al., 2012; Veenstra, 2014,

2019). Furthermore, LGR3 and LGR4 are holomologs of vertebrate LGRs that use ilps as ligands.

This means that the ligands for the LGR4s must be the *Drosophila* ilp 7 orthologs. Since these

peptides are so different from the typical insect neuroendocrine insulins, it made sense to give it a

different name. Earlier work on *Drosophila* suggested that it might have a role similar to relaxin

in vertebrates (Yang et al., 2008) and since LGR4 is an ortholog of the relaxin receptor (Veenstra,

2014), has also been called relaxin, but it might be better to call them arthropod or invertebrate

relaxin.

*Drosophila* ilp 8 (for review see Gontijo & Garelli, 2018) is another ilp (for review see Gontijo &

Garelli, 2018) that acts through a leucine-rich repeat GPCR, LGR3 (Garelli et al., 2015; Vallejo

et al., 2015; Colombani et al., 2015). However, whereas *Drosophila* ilp 7 orthologs have well

conserved primary amino acid sequences, this is not the case for *Drosophila* ilp 8. Indeed, if it

were not for the common presence of LGR3 orthologs in insect and other arthropod genomes one

might believe that this peptide hormone evolved within the higher flies and is absent from other

insects. The imaginal disks in *Drosophila* produce and release ilp 8 as long as they develop and

also when they get damaged. When it is no longer released this is used by the brain as a signal to

initiate metamorphosis (Garelli et al., 2012; Colombani, Andersen & Léopold, 2012; Jaszczak et

al., 2016). *Drosophila* ilp 8 is furthermore produced by the testes and ovaries (Chintapalli, Wang

& Dow, 2007) and since imaginal disks are only present in holometabolous insects, it is tempting

to speculate that the gonads are the original site of expression of orthologs of this peptide. I had

previously suggested that the crustacean androgenic insulin-like peptide that stimulates premature

sexual maturation in male crustaceans and can induce sex reversal in females, might be an

ortholog of *Drosophila* ilp 8 (Veenstra, 2016b). However, more recently a fourth type of ilp was

identified in two decapod species, that seem to be structurally more similar to *Drosophila* ilp 8

than the androgenic insulin-like peptides (Chandler et al., 2017). It has now been shown that

these peptides, which have been called gonadulins, are generally present in decapods and

commonly expressed by the gonads (Veenstra, 2020). Since gonadulins might be orthologs of

*Drosophila* ilp 8 (Veenstra, 2020), it seemed worthwhile to look for this hormone in other

arthropods. Analysis of arthropod genome and transcriptome sequences revealed that such

peptides are not limited to decapods but are also present in insects as well as chelicerates.

During this analysis interesting new details of the putative orthologs of *Drosophila* ilp 6 were

also encountered as well as evidence suggesting that the putative orthologs of *Drosophila* ilps 6,

7 and 8 arose from an ancestral gene triplication.

## Materials & Methods

The sratoolkit (https://trace.ncbi.nlm.nih.gov/Traces/sra/sra.cgi?view=software) in combination

with Trinity (Grabherr et al., 2011) was used in the search for transcripts encoding peptides that

might be somewhat similar to insulin in insect gonad transcriptome short read archives (SRAs).

Once such transcripts had been found, it was often possible to find orthologs from related species.

Alternatively genes coding such transcripts were identified in genome assemblies using Artemis

(Rutherford et al., 2000). For example, once the honeybee gonadulin was found, it was much

easier to find it in other Hymenoptera. Details of the methods used have been previously

described (Veenstra and Khammasi, 2017; Veenstra, 2019) and the only detail that is

significantly different is the large -evalue option of 100 in the tblastn_vdb command from the

sratoolkit. This implied that subsequently many sequences needed to be checked for possible

gonadulin sequence similarity. The same method was used to identify relaxin and C-terminally

extended ilps, which have much better conserved primary amino acid sequences and

consequently are more easily identified, as well as their putative receptors. Whenever possible all

sequences were confirmed in both genome assemblies and in transcriptome SRAs. In many cases

transcripts for the various ilps and receptors were already present in genbank, although they were

not always correctly identified. All these sequences are listed in Suppl. Spreadsheet 1.

Expression was estimated by counting how many RNAseq reads in each SRA contained coding

sequence for each of the genes. In order to avoid untranslated sequences of the complete

transcripts, that sometimes share homologous stretches with transcripts from other genes and can

cause false positives, only the coding sequences were used as query in the blastn_vdb command

from the sratoolkit. This yielded the blue numbers in Suppl. Spreadsheet 2. In order to more

easily compare the different SRAs these numbers were then expressed as per million spots in

each particular SRA. These are the bold black numbers in Suppl. Spreadsheet 2.

For the expression of alternative aILGF splice forms reads for each splice variant were first

separately identified. Unique identifiers in these two sets were determined to obtain the total

number of reads for aILGF. Those identifiers that were present in the initial counts for both splice

forms were counted separately and subtracted from the initial counts of the two splice variants to
obtain the number of reads specific for each isoform.
The various SRAs that were used are listed in the supplementary pdf file and were downloaded
from https://www.ncbi.nlm.nih.gov/sra/. The following genome assemblies were also analyzed:
*Aedes aegypti* (Matthews et al., 2018), *Blattella germanica* (Harrison et al., 2018), *Bombyx mori*
(Kawamoto et al., 2019), *Galleria melonella* (Lange et al., 2018), *Glossina morsitans* (Attardo et
al., 2019), *Hermetia illucens* (Zhan et al., 2020), *Latrodectus hesperus* (unpublished),
*Mesobuthus martensii* (Cao et al., 2013), *Oncopeltus fasciatus* (Panfilio et al., 2019),
*Parasteatoda tepidariorum* (Schwager et al., 2017), *Pardosa pseudoannulata* (Yu et al., 2019),
*Periplaneta americana* (Li et al., 2018), *Stegodyphus dumicola* (Liu et al., 2019), *Tetranychus*
*urticae* (Grbić et al., 2011), *Timema cristinae* (Riesch et al., 2017), *Tribolium castaneum*
(Herndon et al., 2020) and *Zootermopsis nevadensis* (Terrapon et al., 2014). All genomes were
downloaded from https://www.ncbi.nlm.nih.gov/genome/.
181

**Phylogenetic and sequence similarity trees**

For the phylogenetic tree of the insulin RTKs sequences were aligned with clustal omega
(Sievers et al., 2011). Using Seaview (Gouy, Guindon & Gascuel, 2010) only well aligned
sequences were retained and saved as a fasta file. Fasttree2 (Price et al., 2010) was then used to
produce a phylogenetic tree using the ./FastTreeDbl command with the following options: -spr 4 -
mlacc 2 -slownni. The phylogenetic GPCR tree was constructed in the same way using only the
transmembrane regions.
The amino acid sequences of arthropod ilps are not well conserved and hence one can not make
phylogenetic trees, as it is impossible to know which amino acid residues can be reliably aligned
apart from the cysteines. Even the latter can cause problems when the spacing between is not the
same in different peptides. To compare the different ilps an unbiased method is needed. For this
clustal omega was used to align the complete precursor sequences and even though visual
inspection with Seaview reveals very poor alignments, the alignment was not changed but saved
as a fasta file and Fasttree was used with the same parameters as above to construct trees.
Although such trees are not phylogenetic trees they do allow for an unbiased comparison of the
various sequences. The trees so produced have been called sequence similarity trees. Note that

the branch probabilities of such trees give useful information as to how reliable the grouping of

the various ilp precursors is.

**Prediction of prepropeptides processing**

Signal peptides were predicted using signal P-5.0 (Almagro Armenteros et al., 2019). No

attempts were made to predict convertase cleavage sites in the gonadulin and aILGF precursors.

Both of these putative hormones are likely not made by neuroendocrine cells. This implies that

these hormones may not be exposed to neuroendocrine convertases and hence rules that describe

how these convertases might cleave would not be applicable.

# Results

**Gonadulin-like peptides are present in many arthropods**

Peptides that share the typical location of six cysteine residues with insulin but are insufficiently

similar to known insect ilps to be easily recognized as such, were identified in a large number of

arthropods. Their sequences differ not only from other ilps, but are also very variable between

them. As a consequence they are difficult to find in genome assemblies, unless a sequence from a

not too distantly related species is available. This explains why searches were most successful

when done in ovary and/or testes transcriptomes. Putative gonadulin orthologs were identified not

only in insects, but also in several chelicerates, notably spiders, a spider mite and scorpions. A

list of gonadulin propeptides in representative species is given in Figure 1 and additional

sequences are provided in Suppl. Spreadsheet 1. It is evident that although these peptides have

been given the same name, their sequences diverge even more than the neuroendocrine arthropod

insulins. When constructing a sequence similarity tree from the insect ilps the gonadulins are well

separated from the relaxins and the other insect ilps. Interestingly even the insulins and aILGFs

are reasonably well separated, except for the precursors from highly evolved flies (*Drosophila*

and *Glossina*) and the head louse (Fig. 2). When this is repeated on sequences from a set of

arthropod ilps, the gonadulins are once again well separated from the other insulin related

peptides (Fig. 3, Fig. S1).

Publicly available transcriptomes were used to explore in which tissues they are expressed. By

nature such data is imperfect, as these transcriptomes were not made to answer the question

where gonadulin, other insulin-related peptides or their putative receptors are expressed and

hence such data is limited. Some of the more salient examples are illustrated in Table 1. In

honeybees the ovaries of virgin queens do not express significant amounts of gonadulin, but those

of egg-laying queens produce it in large quantities. In a single queen bumblebee larva

transcriptome the gonadulin reads are at least 15 times more numerous than those in the three

transcriptomes each for male and worker larvae (Suppl. Spreadsheet 2). In the tsetse fly *Glossina*

*morsitans* the gene is strongly expressed in ovaries of non-pregnant females, *i.e.* those that

mature an egg, and hardly at all in pregnant/lactating females when egg maturation is arrested. In

the bugs *Rhodnius prolixus* and *Oncopeltus fasciatus* the ovaries also express this gene, as do

ovaries and testes of the stick insect *Timema cristinae,* while short read archives of unfecundated

eggs from *Blattella germanica* similarly contain large amounts of gonadulin reads. In the termite

*Zootermopsis nevadensis,* gonadulin reads are abundant in reproducing females but rare or absent

in alate females or reproducing males. The gene is also expressed in the testis of the American

cockroach and possibly in the ovary as well, since it can be detected in whole body

transcriptomes from females (Suppl. Spreadsheet 2). However, as in decapods (Veenstra, 2020),

in insects gonadulin expression is not limited to the gonads (Table 1; Suppl. Spreadsheet 2). In

the spider *Parasteatoda tepidariorum* ovary expression of gonadulin varied significantly between

different samples (Table 1), and even larger variability in gonadulin expression has previously

been reported for the crab *Portunus trituberculatus* (Veenstra, 2020). This shows that data from a

single SRA are not necessarily informative as to the level of gonadulin expression in this organ.

Interestingly, in some spider transcriptomes gonadulin expression is also observed in silk glands

(Table1, Suppl. Spreadsheet 2).

**251    Arthropod insulin-like growth factors**

Most insect ilps contain only a few amino acid residues after the sixth cysteine residue in the

precursor and sometimes there are none, however some ilps have a long C-terminal extension.

Such ilps are commonly present in hemimetabolous insects as well as several holometabolous

species (Fig. 4, Fig. S2 , Suppl. Spreadsheet 1). In some species the C-terminal extension of these

peptides are easily missed since if one ignores an intron donor site in the genome sequence the

conceptual translation of such sequences predicts much smaller ilps that look similar to the well

known *Drosophila* peptides. Nevertheless, analysis of RNAseq SRAs from several species shows

that such intron donor sites are functional. These C-terminal extensions are coded by two

additional exons that are not present in the typical insect neuroendocrine insulin genes. These

extensions are not only commonly present, but also look similar to one another, providing further

evidence that they are genuine parts of these ilps (Fig. 4). Furthermore, Trinity analysis of SRAs

containing reads for such transcripts reveals that they are alternatively spliced which leads to the

production of precursors with different C-terminal extensions. The difference often consists of

the inclusion or exclusion of a sequence rich in dibasic amino acid residues, mostly arginines,

that in many species has two characteristic cysteine residues. The alternative splice site is in the

middle in the first additional exon, the last coding exon of these genes is less well conserved but

contains a sequence that conforms more or less to the $GTVX_1PX_2(F/Y)$ consensus sequence. Such

genes are present in species as diverse as cockroaches, termites, stick insects, beetles, bees, ants

and moths (Fig. 5). Interestingly, in the stick insect *Timema cristinae* this gene underwent a local

gene duplication, with one gene coding a peptide with the arginine-rich peptide sequence and the

second one lacking it.

In hemiptera ilp genes exist that similarly code for C-terminally extended ilps that are

alternatively spliced, but in those species the extended C-terminals of the predicted peptides are

not as well conserved (Fig. S3). Other C-terminally extended ilps were found in spiders and

scorpions, but in those species no evidence was found for alternative splicing. Such C-terminally

extended ilps appear absent from decapods (Veenstra, 2020).

The term insulin-like growth factors was initially used as a description of substances in plasma

that had insulin-like biological activity, but it is now mostly used as a name for the vertebrate

hormones that are predominantly made in the liver. The use of the same term for both a group of

molecules that have similar characteristics as well two specific hormones is confusing. This is

particularly the case for insects, since hormones that have been called insulin-like growth factors

are not necessarily orthologs of one another nor of these vertebrate hormones. One of the two

types insect insulin-like growth factors, *Bombyx* IGF-like peptide (BIGFLP; Okamoto et al.,

2009a), has only been found in *Bombyx mori*, although it can be expected to be present in other

Lepidoptera as well. The insulin-like growth factor described above seems to be commonly

present in arthropods, including *Bombyx mori* and I propose to call it arthropod insulin-like

growth factor (aILGF).

The data from an extended set of *Bombyx* transcriptome SRAs shows that in this species both

insulin-like growth factors are expressed by the fat body, but the temporal patterns of expression

of the two differ. Thus, aILGF is also expressed in larvae, when there is very little expression of

BIGFLP while during the pupal stage their peaks of expression do not coincide (Fig. 6).

Alternative splicing of aILGF mRNA in the silkworm and the honey bee is variable between the

different samples (Fig. 7, Fig S4). In *Bombyx* SRAs from adults show a relatively higher

expression of the longer isoform that has the additional sequence sequence rich in dibasic amino

acid residues (Fig. S5), while in the honeybee those are more common in SRAs from the bodies

of larvae destined to become queens as well as in samples from the Nasonov glands from nurse

bees and in the only available sample of queen heads (Fig. S4). In most species there are

insufficient data or a single isoform may be predominantly expressed such as in the bumblebee

(Suppl. Spreadsheet 2).

**Is Drosophila ilp 6 an aILGF ortholog ?**

Given the absence of prominent sequence similarity between *Drosophila* ilp 6 and the aILGFs

from more basal insect species, the obvious question is whether or not *Drosophila* ilp 6 is an

ortholog of aILGF or whether it arose independently. The archetype aILGF gene contains four

coding exons, the first two of which code for the insulin structure. The third consists of two parts,

the first half which is present in both isoforms and the last part of this exon which is alternatively

sliced. The fourth and last coding exon contains the $GTVX_1PX_2(F/Y)$ consensus sequence.

During evolution, this basic pattern has been modified on several occasions. In *Oncopeltus* and

other hemiptera the third coding exon was modified, while in both the mosquito *Aedes aegypti*

and the honeybee, the intron between the first two coding exons was eliminated. In *Aedes* other

changes occurred as well, but the gene can still be recognized as an aILGF gene by the presence

of an exon that shows sequence similarity to the fourth coding exon of the aILGF genes. Soldier

flies and robber flies have aILGF genes that are complete except for the third coding exons. It

thus seems that the third coding exon was lost when Diptera evolved (Fig. 8). However, whereas

the aILGF genes in those flies can still be recognized as such by the presence of what once was

the fourth coding exon of a classical aILGF gene, this exon is not only absent from *Drosophila*,

but is missing from all Erenomeura (Fig. 8). There are two possible explanations for this absence,

either this exon was also lost, or the entire aILGF gene was lost and a novel insulin-like growth

factor-like gene evolved in those flies, perhaps not unlike the origin of BIGFLP in *Bombyx*. The

amino acid sequences of the aILGFs from the most highly developed non-Eremoneura flies are

very similar to those from the least evolved Erenomeura flies (Fig. S6). So the aILGF gene

persisted but it lost the last coding exon reminiscent of a classical aILGF gene and the only

physico-chemical characteristic that separates aILGF from the neuroendocrine insulins is the very

short sequence connecting the putative A- and B-chains.

### Arthropod relaxins

The arthropod relaxins are known from a large number of arthropods, although they are lacking

in many insect species. As noted previously they have by far the best conserved amino acid

sequences in both the A- and B-chains of all the protostomian ilps and paralogs of this particular

peptide are also found in other protostomians and even some basal deuterostomes. Neuropeptides

containing cysteine residues typically have them in pairs, since as soon a cysteine containing

peptide enters the endoplasmatic reticulum these residues get oxidized and form disulfide

bridges. For this reason it is very surprising to see a decapod relaxin having seven cysteine

residues (Fig. 9). The presence of this particular cysteine is unlikely to be an artifact as it is found

in relaxins from a number of decapod species from different orders (Fig. S7). Although most

arthropods appear to have a single relaxin gene, two scorpion genomes have two such genes; in

both cases the predicted amino acid sequence of one of the relaxins seems less conserved

(Veenstra, 2016a).

### Gonadulin, relaxin and aILGF likely evolved from a local gene triplication

The very large primary amino acid sequence variability of arthropod ilps makes it difficult to

establish their phylogenetic affinities. Sequence similarity trees show that the gonadulins

resemble one another and hence may share a common ancestor, but such trees do not provide

details. Synteny offers another means to establish evolutionary relationships and, as chance has it,

the genomes of some species suggests that aILGF, gonadulin and arthropod relaxin originated

from an ancient gene triplication. Thus the three genes coding gonadulin, aILGF and relaxin are

located next to one another in the genomes of the German cockroach and the termite

*Zootermopsis nevadensis*. In the stick insect *Timema cristinae* the aILGF underwent a local

duplication (see above) and the four genes coding the two aILGFs, gonadulin and relaxin, are

also next to one another. Furthermore, in the genomes from the spiders *Parasteatoda*

*tepidariorum* and *Pardosa pseudoannulata* the gonadulin genes are next to an aILGF gene, which

in these spiders has undergone one or more local duplications (Fig. 10). Such a close genomic

association of these genes strongly suggests that they arose from a local gene triplication.

## Receptors

Two different types of receptors are used by ilps, LGRs and RTKs. Insect have two different types of insulin RTKs, that have been labeled A and B. Most species have one gene coding an type A and one gene coding for an B type B RTK. However, cockroaches, termites and at least some stick insects have two B type RTKs and the American cockroach has also two type A RTKs. The genome assembly of the American cockroach shows that the two type A RTKs originated from a local gene duplication, as did the two type B RTKs. In Lepidoptera and Diptera all insulin RTKs are type A and there is no type B RTK. However, Lepidoptera do have a second gene coding for a protein with clear structural similarity to insulin RTKs, but these proteins lack the tyrosine kinase domain. I have called those proteins Insulin RTK-like (IRTKL). This gene is more similar to the type B RTKs than type A and it may have evolved from those RTKs. The presence of an IRTKL coding gene in at least two genomes from Trichoptera, suggests that it may have evolved before the two groups diverged. Interestingly, the genome of the beetle *Tribolium castaneum* has genes for both a type A and a type B RTK, as well as for an IRTKL. No IRTKL orthologs was detected in any of other sequenced beetle genomes, but such an ortholog was found in the transcriptome data of *Tenebrio molitor,* a species that is closely related to *Tribolium*. A phylogenetic tree of the various RTKs and IRTKL proteins suggests that the origin of the IRTKL proteins in Lepidoptera is ancient, but that of *Tribolium* is quite recent, indicating convergent evolution (Fig. 11). In chelicerates there are also two types of RTK, while there are four in decapods (Veenstra, 2020), however the various gene duplications of the arthropod RTKs occurred after these groups diverged (Fig. 11).

*Drosophila* LGR3 is activated by *Drosophila* ilp 8 and orthologs of this receptor are present in many but not all arthropods, always in a single copy. As described LGR4 is likely the arthropod relaxin receptor and is similarly present in many arthropods, usually in a single copy, but chelicerates have two copies and some spider have even three. LGR3 and LGR4 are closely related GPCRs together with another receptor that was first described from the pond snail *Lymnaea stagnalis* as GRL101 (Tensen et al., 1994) to stress its similarity with LGR3 and LGR4 it is called here LGR5. The strong sequence similarity of LGR5 with LGR3 and LGR4 suggests that it, like LGR3 and LGR4 might have an insulin-like ligand and for this reason it is included here. LGR5 is commonly present in arthropods. Unlike LGR3 and LGR4, that are missing in many insect species, LGR5 is consistently present in hemimetabolous insects, but it is completely absent from all holometabolous insects (the GRL101 from *Rhagoletis zephyria* [XP_017487580]

386    is appears to be a mite GPCR). In both chelicerates and decapods there are usually several

387    paralogs (Fig. 12). All three of these receptors are widely expressed and it is difficult to discern

388    clear expression patterns (Suppl. Spreadsheet 2).

## Discussion

I describe a number of novel arthropod ilps, the gonadulins and aILGFs, that are putative

orthologs of *Drosophila* ilp 8 and ilp 6 respectively and I present evidence that the genes coding

these peptides and relaxin are commonly present in arthropods and most likely originated from an

ancient gene triplication.

There are four arguments that together suggests that the various gonadulins and *Drosophila* ilp 8

are indeed orthologs. First, the gonadulins cluster together with *Drosophila* ilp 8 on on a

sequence similarity tree that bundles peptides with similar structures. It is clear that even though

the amino acid sequences of the gonadulins are poorly conserved, they do resemble one another

better than each one of them resembles any of the better known arthropod ilps. Secondly, for

those members of this group where this could be determined, they are all made by the gonads,

although this is not the only tissue expressing these peptides and expression by the gonads is

variable. Thirdly, all species for which a gonadulin could be identified also have an ortholog of

*Drosophila* LGR3, even though not all arthropod species have such a receptor. Unfortunately,

due to the large structural variability of the gonadulins, it was not always possible to demonstrate

the existence of a gonadulin gene in each species that has an ortholog of LGR3. Nevertheless, no

gonadulins were found in species lacking such a receptor. Finally, peptides that were identified as

putative gonadulins, but that are present in species as distantly related as spiders on one hand and

stick insects and cockroaches on the other, are produced by genes for which orthology is

independently confirmed by synteny with aILGF genes.

The physiological significance of the presence of both aILGF and BIGFLP in *Bombyx* is an

intriguing question, perhaps even more so as there seems to be only a single insulin receptor in

this species that might induce an intracellular response. As shown by the data of an impressive set

of transcriptome SRAs from this species, expression of the two differs during development (Fig.

6). Although the physiological meaning of this remains to be investigated, it is tempting to

speculate that it is related to oogenesis and metamorphosis taking place simultaneously. In most

insect species juvenile hormone and ecdysone play preponderant roles in the regulation of both

processes, but using the same hormones for two different processes at the same time might be

counterproductive and may have led to the evolution of BIGFLP.

It has been reported previously that, unlike *Drosophila,* most insect species have more than one

insulin RTK (Kremer, Korb & Bornberg-Bauer, 2016). This is confirmed here and in the

American cockroach there are actually four insulin RTKs. The presence of three insulin RTKs in

termites has led to the suggestion that they might be involved in caste determination (Kremer,

Korb & Bornberg-Bauer, 2016). Although this may be so, it is somewhat surprising in this

context that the American cockroach, a close relative, has even four such receptors, yet has no

caste system. In Lepidoptera and at least two tenebrionid beetles one of the putative insulin

receptors lacks the tyrosine kinase domain, like the vertebrate ILGF2 receptor and the

*Drosophila* decoy insulin receptor (Okamoto et al., 2013). The *Drosophila* receptor also lacks the

transmembrane domaine and is released into the hemolymph. These receptors may well have

similars role in regulating the hemolymph ilp concentrations.

In some insects the expression of insulin RTKs have been studied sometimes together with the

effects of inactivating one or both insulin RTKs by RNAi (Wheeler, Buck & Evans, 2006; Sang

et al., 2016; Okada et al., 2019). Interestingly, in honeybee larvae that are changed from worker

food to royal yelly, that is rich in proteins, it is the RTK A that is upregulated (Wheeler, Buck &

Evans, 2006), whereas in two beetle species the effects of RTK inactivation by RNAi is much

stronger for type A than for type B (Sang et al., 2016; Okada et al., 2019). In the honeybee and

the beetle *Gnatocerus* these receptors have been indirectly linked to an aILGF peptide. In Diptera

and in the head louse there is only a single RTK, which in both case are also type A and this is

also the type that is most abundantly expressed in insects. This suggests that the A type insulin

RTK is the more important than the B type RTK.

The structural difference between neuroendocrine ilps and aILGFs, as illustrated by the large C-

terminal extensions absent in the former but present in the latter, suggest the use of different

receptors. Indeed, the similarities between insulin and vertebrate ILGF on one hand and insect

neuroendocrine ilps and aILGF on the other, are striking. Thus whereas the liver is the major

tissue expressing ILGF, in insects aILGF is expressed by the fat body, the tissue that performs the

functions of both the vertebrate liver and adipose tissue. Furthermore, both ILGF and aILGF are

C-terminally extended insulin-like molecules and in both cases the primary transcripts produced

are alternatively spliced (Roberts et al., 1987). Finally, insulin and ILGF act on two RTKs and

many insects also seem to have two insulin RTKs, although structural similarity is insufficient

proof that these are insulin receptors, as illustrated by the insulin receptor–related receptor that

functions as an alkali receptor (Deyev et al., 2011). Nevertheless, this suggests that the C-

terminal extension of aILGF may allow for the differential activation of two different types of

insulin RTKs and the presence of only one insulin receptor in Diptera may well explain why

*Drosophila* ilp 6 has lost this C-terminal extension. Furthermore, it is perhaps no coincidence that

the ilps that did not separately clearly into either an aILGF or an insulin branch on the sequence

similarity tree (Fig. 2) are from *Drosophila*, *Glossina* and *Pediculus.* These are species that have

only one insulin RTK and thus there would be no physiological need to maintain different

molecular structures for peptides in order to preferentially activate one or the other of the two

insulin RTKs.

The close genomic association of the gonadulin, relaxin and ILGF genes in some arthropod

genomes suggest that they originated from an ancient gene triplication. Although the primary

amino acid sequences of the different gonadulins is limited, as discussed above, there is reason to

believe they are also orthologs and the same holds for the various aILGFs. The *Drosophila*

gonadulin (dilp 8) receptor has been shown to be *Drosophila* LGR3, while LGR4 must be an

arthropod relaxin receptor. As demonstrated by the sequence similarity of their transmembrane

regions, these LGRs and LGR5 are evolutionary cousins. Insect aILGFs on the other hand are

known to stimulate RTKs. When neuropeptide genes undergo a local duplication the ligands they

encode either keep using the same receptor or use paralogs that have their origin in a gene

duplication of the receptor. The gene coding aILGF that uses an RTK is flanked on both sides by

genes that code LGR ligands. The only reasonable explanation is that the original gene coding for

an insulin-like peptide (the one that got triplicated) used both types of receptors, *i.e.* it had two

receptors, both an LGR and an RTK.

It is clear that during evolution at least holometabolous species no longer have such a GPCR, as

these species have at most only two such receptors, LGR3 and LGR4 (for gonadulin and relaxin

respectively), while some species have none. Nevertheless, it is interesting to note that the

hemimetabolous insects have LGR5, a receptor that is evolutionarily closely related to the

receptors for gonadulin and relaxin. Thus LGR5 could be a second receptor for aILGF. If this

were so, then the remarkable absence of LGR5 from holometabolous species might be related to

the switch to holometaboly itself. Maggots and caterpillars undergo essentially linear growth,

while in non-holometabolous species, growth is accompanied by development at the same time.

Perhaps, LGR5 stimulated by aILGF is responsible for this.

Arthropod relaxin has an primary amino acid sequence that is much better conserved than that of

the typical arthropod insulins or ILGFs. As arthropod relaxin shares what seem to be the

structural characteristics common to both aILGF and neuroendocrine insulins, it seems plausible

that relaxin can also stimulate the insulin RTK. It is of interest in this respect that *Drosophila*

relaxin, ilp7, binds to the decoy insulin receptor as well other *Drosophila* ilps (Okamoto et al.,

2013) and previous work suggested that it acts through the *Drosophila* RTK (Linneweber et al.,

2014). When there is only a single receptor for a ligand, both can co-evolve and over time

structures of both the ligand and its receptor may change. However, when the ligand activates

two different receptors, all three components have to coevolve and one would expect that this

would restrain the structures of all three elements much more than when there is only a single

receptor and such restraints would be the strongest on the ligand that activates both receptors.

This might explain why arthropod relaxin is so well conserved.

It is tempting to speculate that orthologous genes have similar expression patterns and functions.

In related species this is a reasonable hypothesis and the observed expression of aILGFs in

cockroaches (Castro-Arnau et al., 2019) and beetles (Okada et al., 2019) is predominantly in the

fat body and this seems to be the case for *Bombyx* aILGF too. What is likely the *Rhodnius* aILGF

is also expressed by the fat body (Defferrari, Orchard & Lange, 2015) as is *Drosophila* ilp 6

(Okamoto et al., 2009b). This suggest that within insects these peptides have the same function. It

is notable however, that at least in *Pardosa* spiders the fat body does not express aILGF but a

specific insulin (Yu et al., 2020), while aILGFs expression is limited to the cephalothorax. This

suggests that in spiders aILGFs are expressed in the brain, is indeed observed in the spider

*Stegodyphus dumicola* (Suppl. Spreadsheet 2). In decapods, which are more closely related to

insects than chelicerates, no aILGFs were found (Veenstra, 2020). The phylogenetic tree of the

various arthropod insulin RTKs also shows that the various paralogs of this receptor are not direct

orthologs of one another, but must have evolved independently in each subphylum or even class.

This within arthropods the functions of the various insulin-like peptides may be significantly

different. It suggests that the apparent resemblance between insect neuroendocrine insulins and

aILGF on one hand and insulin and ILGF on the other could reflect a case of convergent

evolution rather than one of orthology.

In the beetle *Gnatocerus cornutus* it has been shown that aILGF specifically stimulates the

growth of a sexual ornament (Okada et al., 2019), while higher levels of aILGF are observed in

honeybee larvae that are destined to become queens and thus develop functional ovaries

(Wheeler, Buck & Evans, 2006). In *Gnatocerus* aILGF release depends on nutrition status and in

honeybees protein rich royal yelly is associated with in increase of aILGF. Although we don't

know as much detail for *Blattella* aILGF, its expression is strongly inhibited during starvation

(Castro-Arnau et al., 2019). This suggests that in insects aILGF is released by the fat body in

response to nutritious food.

The physiological function of gonadulin is less clear. Insulin and related peptides typically

stimulate growth and reproduction, so its presence in the ovaries and testes suggests a function in

reproduction. Its presence in unfecundated eggs of *Blattella* suggests that within the ovary it are

the oocytes themselves that express gonadulin, likely the follicle cells that in *Drosophila* have

been shown to express *Drosophila* ilp 8 (Liao & Nässel, 2020). In the crab *Portunus*

*trituberculatus* gonadulin expression is on occasion very high in the gonads (Veenstra, 2020), but

the very variable degree of expression makes it difficult to see this hormone as merely

stimulating reproduction. The expression of gonadulin in hematopoetic tissue and the anterior

proliferation center of the brain in *Procambarus clarkii* (Veenstra, 2020), neither suggest a role

limited to reproduction but hints at a more general role in promoting growth. Spider silks are

proteins and thus its production requires plenty of amino acids, not unlike vitellogenesis, or the

development and reparation of imaginal disks. Gonadulin secreted by these organs might

therefore suggest that, not unlike insulin, it stimulates growth, but more intensely and/or more

localized. Such an intensified stimulation of growth might be achieved by increasing not only the

uptake of glucose as an energy substrate but also that of amino acids. Under this hypothesis, it

might act as both an autocrine to stimulate uptake of metabolites and an endocrine to make these

available and by doing so it might be able to stimulate growth of specific organs, such as

imaginal disks and/or gonads that secrete it.

**Conclusions**

A local gene triplication in an early ancestor likely yielded three genes coding gonadulin,

arthropod insulin-like growth factor and relaxin. Orthologs of these genes are now commonly

present in arthropods and almost certainly include the *Drosophila* insulin-like peptides 6, 7 and 8.

550  **Acknowledgements**

551 I gracefully acknowledge the contributions made by all those who not only produced but also
552 made available the numerous SRAs that I analyzed here. This work would neither have been
553 possible without the various programs employed.
554
555

556
557
## References
Almagro Armenteros JJ, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak S, von
Heijne G, Nielsen H. 2019. SignalP 5.0 improves signal peptide predictions using deep neural
networks. Nat Biotechnol. 37:420-423. doi: 10.1038/s41587-019-0036-z.
Attardo GM, Abd-Alla AMM, Acosta-Serrano A, Allen JE, Bateta R, Benoit JB, Bourtzis K,
Caers J, Caljon G, Christensen MB, Farrow DW, Friedrich M, Hua-Van A, Jennings EC, Larkin
DM, Lawson D, Lehane MJ, Lenis VP, Lowy-Gallego E, Macharia RW, Malacrida AR, Marco
HG, Masiga D, Maslen GL, Matetovici I, Meisel RP, Meki I, Michalkova V, Miller WJ, Minx P,
Mireji PO, Ometto L, Parker AG, Rio R, Rose C, Rosendale AJ, Rota-Stabelli O, Savini G,
Schoofs L, Scolari F, Swain MT, Takáč P, Tomlinson C, Tsiamis G, Van Den Abbeele J,
Vigneron A, Wang J, Warren WC, Waterhouse RM, Weirauch MT, Weiss BL, Wilson RK, Zhao
X, Aksoy S. 2019. Comparative genomic analysis of six *Glossina* genomes, vectors of African
trypanosomes. Genome Biol. 20:187. doi: 10.1186/s13059-019-1768-2.
Brogiolo W, Stocker H, Ikeya T, Rintelen F, Fernandez R, Hafen E. 2001. An evolutionarily
conserved function of the *Drosophila* insulin receptor and insulin-like peptides in growth control.
Curr Biol. 11:213-221. doi: 10.1016/s0960-9822(01)00068-9.
Cao Z, Yu Y, Wu Y, Hao P, Di Z, He Y, Chen Z, Yang W, Shen Z, He X, Sheng J, Xu X, Pan B,
Feng J, Yang X, Hong W, Zhao W, Li Z, Huang K, Li T, Kong Y, Liu H, Jiang D, Zhang B, Hu
J, Hu Y, Wang B, Dai J, Yuan B, Feng Y, Huang W, Xing X, Zhao G, Li X, Li Y, Li W. 2013.
The genome of *Mesobuthus martensii* reveals a unique adaptation model of arthropods. Nat
Commun. 4:2602. doi: 10.1038/ncomms3602.
Castellanos MC, Tang JC, Allan DW. 2013. Female-biased dimorphism underlies a female-
specific role for post-embryonic Ilp7 neurons in *Drosophila* fertility. Development. 140:3915-
3926. doi: 10.1242/dev.094714.
Castro-Arnau J, Marín A, Castells M, Ferrer I, Maestro JL. 2019. The expression of cockroach
insulin-like peptides is differentially regulated by physiological conditions and affected by
compensatory regulation. J Insect Physiol. 114:57-67. doi: 10.1016/j.jinsphys.2019.02.010.
Chandler JC, Gandhi NS, Mancera RL, Smith G, Elizur A, Ventura T. 2017. Understanding
Insulin Endocrinology in Decapod Crustacea: Molecular Modelling Characterization of an
Insulin-Binding Protein and Insulin-Like Peptides in the Eastern Spiny Lobster, *Sagmariasus
verreauxi.* Int J Mol Sci. 18. pii: E1832. doi: 10.3390/ijms18091832.
Chintapalli VR, Wang J, Dow JA. 2007. Using FlyAtlas to identify better *Drosophila
melanogaster* models of human disease. Nat Genet. 39:715-20. doi: 10.1038/ng2049.
Colombani J, Andersen DS, Léopold P. 2012. Secreted peptide Dilp8 coordinates *Drosophila
tissue growth with developmental timing. Science. 336:582-5. doi: 10.1126/science.1216689.

Colombani J, Andersen DS, Boulan L, Boone E, Romero N, Virolle V, Texada M, Léopold P.

2015. *Drosophila* Lgr3 Couples Organ Growth with Maturation and Ensures Developmental

Stability. Curr Biol. 25:2723-2729. doi: 10.1016/j.cub.2015.09.020.

Defferrari MS, Orchard I, Lange AB. 2015. An insulin-like growth factor in *Rhodnius prolixus* is

involved in post-feeding nutrient balance and growth. Front Neurosci. 10:566. doi:

10.3389/fnins.2016.00566.

Deyev IE, Sohet F, Vassilenko KP, Serova OV, Popova NV, Zozulya SA, Burova EB, Houillier

P, Rzhevsky DI, Berchatova AA, Murashev AN, Chugunov AO, Efremov RG, Nikol'sky NN,

Bertelli E, Eladari D, Petrenko AG. 2011. Insulin receptor-related receptor as an extracellular

alkali sensor. Cell Metab. 13:679-689. doi: 10.1016/j.cmet.2011.03.022.

Garelli A, Gontijo AM, Miguela V, Caparros E, Dominguez M. 2012. Imaginal discs secrete

insulin-like peptide 8 to mediate plasticity of growth and maturation. Science. 336:579-582. doi:

10.1126/science.1216735.

Garelli A, Heredia F, Casimiro AP, Macedo A, Nunes C, Garcez M, Dias ARM, Volonte YA,

Uhlmann T, Caparros E, Koyama T, Gontijo AM. 2015. Dilp8 requires the neuronal relaxin

receptor Lgr3 to couple growth to developmental timing. Nat Commun. 6:8732. doi:

10.1038/ncomms9732.

Gontijo AM, Garelli A. 2018. The biology and evolution of the Dilp8-Lgr3 pathway: A relaxin-

like pathway coupling tissue growth and developmental timing control. Mech Dev. 154:44-50.

doi: 10.1016/j.mod.2018.04.005.

Gouy M, Guindon S, Gascuel O. 2010. SeaView version 4: A multiplatform graphical user

interface for sequence alignment and phylogenetic tree building. Mol Biol Evol. 2010

Feb;27(2):221-4. doi: 10.1093/molbev/msp259.

Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L,

Raychowdhury R, Zeng Q, Chen Z, Mauceli E, Hacohen N, Gnirke A, Rhind N, di Palma F,

Birren BW, Nusbaum C, Lindblad-Toh K, Friedman N, Regev A. 2011. Full-length

transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol.

15;29:644-652. doi: 10.1038/nbt.1883.

Grbić M, Van Leeuwen T, Clark RM, Rombauts S, Rouzé P, Grbić V, Osborne EJ, Dermauw W,

Ngoc PC, Ortego F, Hernández-Crespo P, Diaz I, Martinez M, Navajas M, Sucena É, Magalhães

S, Nagy L, Pace RM, Djuranović S, Smagghe G, Iga M, Christiaens O, Veenstra JA, Ewer J,

Villalobos RM, Hutter JL, Hudson SD, Velez M, Yi SV, Zeng J, Pires-daSilva A, Roch F,

Cazaux M, Navarro M, Zhurov V, Acevedo G, Bjelica A, Fawcett JA, Bonnet E, Martens C,

Baele G, Wissler L, Sanchez-Rodriguez A, Tirry L, Blais C, Demeestere K, Henz SR, Gregory

TR, Mathieu J, Verdon L, Farinelli L, Schmutz J, Lindquist E, Feyereisen R, Van de Peer Y.

2011. The genome of *Tetranychus urticae* reveals herbivorous pest adaptations. Nature. 479:487-

492. doi: 10.1038/nature10640.

Grönke S, Clarke DF, Broughton S, Andrews TD, Partridge L. 2010. Molecular evolution and

functional characterization of *Drosophila* insulin-like peptides. PLoS Genet. 6:e1000857. doi:

10.1371/journal.pgen.1000857.

Harrison MC, Jongepier E, Robertson HM, Arning N, Bitard-Feildel T, Chao H, Childers CP,

Dinh H, Doddapaneni H, Dugan S, Gowin J, Greiner C, Han Y, Hu H, Hughes DST, Huylmans

AK, Kemena C, Kremer LPM, Lee SL, Lopez-Ezquerra A, Mallet L, Monroy-Kuhn JM, Moser

A, Murali SC, Muzny DM, Otani S, Piulachs MD, Poelchau M, Qu J, Schaub F, Wada-

Katsumata A, Worley KC, Xie Q, Ylla G, Poulsen M, Gibbs RA, Schal C, Richards S, Belles X,

Korb J, Bornberg-Bauer E. 2018. Hemimetabolous genomes reveal molecular basis of termite

eusociality. Nat Ecol Evol. 2:557-566. doi: 10.1038/s41559-017-0459-1.

Herndon N, Shelton J, Gerischer L, Ioannidis P, Ninova M, Dönitz J, Waterhouse RM, Liang C,

Damm C, Siemanowski J, Kitzmann P, Ulrich J, Dippel S, Oberhofer G, Hu Y, Schwirz J,

Schacht M, Lehmann S, Montino A, Posnien N, Gurska D, Horn T, Seibert J, Vargas Jentzsch

IM, Panfilio KA, Li J, Wimmer EA, Stappert D, Roth S, Schröder R, Park Y, Schoppmeier M,

Chung HR, Klingler M, Kittelmann S, Friedrich M, Chen R, Altincicek B, Vilcinskas A,

Zdobnov E, Griffiths-Jones S, Ronshaugen M, Stanke M, Brown SJ, Bucher G. 2020. Enhanced

genome assembly and a new official gene set for *Tribolium castaneum*. BMC Genomics. 21:47.

doi: 10.1186/s12864-019-6394-6.

Jaszczak JS, Wolpe JB, Bhandari R, Jaszczak RG, Halme A. 2016. Growth coordination during

*Drosophila melanogaster* imaginal disc regeneration is mediated by signaling through the relaxin

receptor LGR3 in the prothoracic gland. 204:703-709. doi: 10.1534/genetics.116.193706.

Kawamoto M, Jouraku A, Toyoda A, Yokoi K, Minakuchi Y, Katsuma S, Fujiyama A, Kiuchi T,

Yamamoto K, Shimada T. 2019. High-quality genome assembly of the silkworm, *Bombyx mori*.

Insect Biochem Mol Biol. 107:53-62. doi: 10.1016/j.ibmb.2019.02.002.

Kremer LPM, Korb J, Bornberg-Bauer E. 2016. Reconstructed evolution of insulin receptors in

insects reveals duplications in early insects and cockroaches. J Exp Zool B Mol Dev Evol.

330:305-311. doi: 10.1002/jez.b.22809.

Lange A, Beier S, Huson DH, Parusel R, Iglauer F, Frick JS. 2018. Genome Sequence of

*Galleria mellonella* (Greater Wax Moth). Genome Announc. 6. pii: e01220-17. doi:

10.1128/genomeA.01220-17.

Li S, Zhu S, Jia Q, Yuan D, Ren C, Li K, Liu S, Cui Y, Zhao H, Cao Y, Fang G, Li D, Zhao X,

Zhang J, Yue Q, Fan Y, Yu X, Feng Q, Zhan S. 2018. The genomic and functional landscapes of

developmental plasticity in the American cockroach. Nat Commun. 9:1008. doi: 10.1038/s41467-

018-03281-1.

Liao S, Nässel DR. 2020. Drosophila insulin-like peptide 8 (DILP8) in ovarian follicle cells

regulates ovulation and metabolism. bioRxiv. doi: https://doi.org/10.1101/2020.05.02.073585.

Linneweber GA, Jacobson J, Busch KE, Hudry B, Christov CP, Dormann D, Yuan M, Otani T,

Knust E, de Bono M, Miguel-Aliaga I. 2014. Neuronal control of metabolism through nutrient-

dependent modulation of tracheal branching. Cell. 156:69-83. doi: 10.1016/j.cell.2013.12.008.

Liu Y, Liao S, Veenstra JA, Nässel DR. 2016. *Drosophila* insulin-like peptide 1 (DILP1) is

transiently expressed during non-feeding stages and reproductive dormancy. Sci Rep. 6:26620.

doi: 10.1038/srep26620.

Liu S, Aageaard A, Bechsgaard J, Bilde T. 2019. DNA Methylation Patterns in the Social Spider,

*Stegodyphus dumicola*. 2019. Genes (Basel). 10. pii: E137. doi: 10.3390/genes10020137.

Matthews BJ, Dudchenko O, Kingan SB, Koren S, Antoshechkin I, Crawford JE, Glassford WJ,

Herre M, Redmond SN, Rose NH, Weedall GD, Wu Y, Batra SS, Brito-Sierra CA, Buckingham

SD, Campbell CL, Chan S, Cox E, Evans BR, Fansiri T, Filipović I, Fontaine A, Gloria-Soria A,

Hall R, Joardar VS, Jones AK, Kay RGG, Kodali VK, Lee J, Lycett GJ, Mitchell SN, Muehling
J, Murphy MR, Omer AD, Partridge FA, Peluso P, Aiden AP, Ramasamy V, Rašić G, Roy S,
Saavedra-Rodriguez K, Sharan S, Sharma A, Smith ML, Turner J, Weakley AM, Zhao Z, Akbari
OS, Black WC 4th, Cao H, Darby AC, Hill CA, Johnston JS, Murphy TD, Raikhel AS, Sattelle
DB, Sharakhov IV, White BJ, Zhao L, Aiden EL, Mann RS, Lambrechts L, Powell JR,
Sharakhova MV, Tu Z, Robertson HM, McBride CS, Hastie AR, Korlach J, Neafsey DE,
Phillippy AM, Vosshall LB. 2018. Improved reference genome of *Aedes aegypti* informs
arbovirus vector control. Nature. 563:501-507. doi: 10.1038/s41586-018-0692-z.
Miguel-Aliaga I, Thor S, Gould AP. 2008. Postmitotic specification of *Drosophila* insulinergic
neurons from pioneer neurons. PLoS Biol 6: e58. doi:10.1371/journal.pbio.0060058.
Mizoguchi A, Okamoto N. 2013. Insulin-like and IGF-like peptides in the silkmoth Bombyx
mori: discovery, structure, secretion, and function. Front Physiol. 2013 4:217. doi:
10.3389/fphys.2013.00217.
Murphy CT, Hu PJ. 2013. Insulin/insulin-like growth factor signaling in *C. elegans*. WormBook,
ed. The C. elegans Research Community, WormBook. 2013. doi: 10.1895/wormbook.1.164.1.
Nagasawa H, Kataoka H, Isogai A, Tamura S, Suzuki A, Ishizaki H, Mizoguchi A, Fujiwara Y,
Suzuki A. 1984. Amino-terminal amino Acid sequence of the silkworm prothoracicotropic
hormone: homology with insulin. Science.226:1344-1345. doi: 10.1126/science.226.4680.1344.
Nagasawa H, Kataoka H, Isogai A, Tamura S, Suzuki A, Mizoguchi A, Fujiwara Y, Suzuki A,
Takahashi SY, Ishizaki H. 1986. Amino acid sequence of a prothoracicotropic hormone of the
silkworm *Bombyx mori.* Proc Natl Acad Sci U S A. 83:5840-5843. doi: 10.1073/pnas.83.16.5840.
Nässel DR, Vanden Broeck J. 2016. Insulin/IGF signaling in *Drosophila* and other insects:
factors that regulate production, release and post-release action of the insulin-like peptides. Cell
Mol Life Sci. 2016 Jan;73(2):271-90. doi: 10.1007/s00018-015-2063-3.
O'Brien LE, Soliman SS, Li X, Bilder D. 2011. Altered modes of stemcell division drive
adaptive intestinal growth. Cell 147:603–614. doi: 10.1016/j.cell.2011.08.048.
Okada Y, Katsuki M, Okamoto N, Fujioka H, Okada K. 2019a. A specific type of insulin-like
peptide regulates the conditional growth of a beetle weapon. PLoS Biol. 17:e3000541. doi:
10.1371/journal.pbio.3000541.
Okamoto N, Yamanaka N, Satake H, Saegusa H, Kataoka H, Mizoguchi A. 2009a. An
ecdysteroid-inducible insulin-like growth factor-like peptide regulates adult development of the
silkmoth *Bombyx mori.* FEBS J. 276:1221-1332. doi: 10.1111/j.1742-4658.2008.06859.x.
Okamoto N, Yamanaka N, Yagi Y, Nishida Y, Kataoka H, O'Connor MB, Mizoguchi A. 2009b.
A fat body-derived IGF-like peptide regulates postfeeding growth in *Drosophila.* Dev Cell.
17:885-891. doi: 10.1016/j.devcel.2009.10.008.
Okamoto N, Nakamori R, Murai T, Yamauchi Y, Masuda A, Nishimura T. 2013. A secreted
decoy of InR antagonizes insulin/IGF signaling to restrict body growth in *Drosophila.* Genes
Dev. 27:87-97. doi: 10.1101/gad.204479.112.
Panfilio KA, Vargas Jentzsch IM, Benoit JB, Erezyilmaz D, Suzuki Y, Colella S, Robertson HM,
Poelchau MF, Waterhouse RM, Ioannidis P, Weirauch MT, Hughes DST, Murali SC, Werren JH,
Jacobs CGC, Duncan EJ, Armisén D, Vreede BMI, Baa-Puyoulet P, Berger CS, Chang CC, Chao
H, Chen MM, Chen YT, Childers CP, Chipman AD, Cridge AG, Crumière AJJ, Dearden PK,

Didion EM, Dinh H, Doddapaneni HV, Dolan A, Dugan S, Extavour CG, Febvay G, Friedrich

M, Ginzburg N, Han Y, Heger P, Holmes CJ, Horn T, Hsiao YM, Jennings EC, Johnston JS,

Jones TE, Jones JW, Khila A, Koelzer S, Kovacova V, Leask M, Lee SL, Lee CY, Lovegrove

MR, Lu HL, Lu Y, Moore PJ, Munoz-Torres MC, Muzny DM, Palli SR, Parisot N, Pick L, Porter

ML, Qu J, Refki PN, Richter R, Rivera-Pomar R, Rosendale AJ, Roth S, Sachs L, Santos ME,

Seibert J, Sghaier E, Shukla JN, Stancliffe RJ, Tidswell O, Traverso L, van der Zee M, Viala S,

Worley KC, Zdobnov EM, Gibbs RA, Richards S. 2019. Molecular evolutionary trends and

feeding ecology diversification in the Hemiptera, anchored by the milkweed bug genome.

Genome Biol. 20:64. doi: 10.1186/s13059-019-1660-0.

Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-likelihood trees for

large alignments. PLoS One. 5:e9490. doi: 10.1371/journal.pone.0009490.

Riesch R, Muschick M, Lindtke D, Villoutreix R, Comeault AA, Farkas TE, Lucek K, Hellen E,

Soria-Carrasco V, Dennis SR, de Carvalho CF, Safran RJ, Sandoval CP, Feder J, Gries R, Crespi

BJ, Gries G, Gompert Z, Nosil P. 2017. Transitions between phases of genomic differentiation

during stick-insect speciation. Nat Ecol Evol. 1:82. doi: 10.1038/s41559-017-0082.

Roberts CT Jr, Lasky SR, Lowe WL Jr, Seaman WT, LeRoith D. 1987. Molecular cloning of rat

insulin-like growth factor I complementary deoxyribonucleic acids: differential messenger

ribonucleic acid processing and regulation by growth hormone in extrahepatic tissues. Mol

Endocrinol. 1:243-248. doi: 10.1210/mend-2-6-528.

Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream MA, Barrell B. 2000. Artemis:

sequence visualization and annotation. Bioinformatics. 16:944-945. doi: 10.1093/bioinformatics/

16.10.944.

Sang M, Li C, Wu W, Li B. 2016. Identification and evolution of two insulin receptor genes

involved in Tribolium castaneum development and reproduction. Gene. 585:196-204. doi:

10.1016/j.gene.2016.02.034.

Schwager EE, Sharma PP, Clarke T, Leite DJ, Wierschin T, Pechmann M, Akiyama-Oda Y,

Esposito L, Bechsgaard J, Bilde T, Buffry AD, Chao H, Dinh H, Doddapaneni H, Dugan S,

Eibner C, Extavour CG, Funch P, Garb J, Gonzalez LB, Gonzalez VL, Griffiths-Jones S, Han Y,

Hayashi C, Hilbrant M, Hughes DST, Janssen R, Lee SL, Maeso I, Murali SC, Muzny DM,

Nunes da Fonseca R, Paese CLB, Qu J, Ronshaugen M, Schomburg C, Schönauer A, Stollewerk

A, Torres-Oliva M, Turetzek N, Vanthournout B, Werren JH, Wolff C, Worley KC, Bucher G,

Gibbs RA, Coddington J, Oda H, Stanke M, Ayoub NA, Prpic NM, Flot JF, Posnien N, Richards

S, McGregor AP. 2017. The house spider genome reveals an ancient whole-genome duplication

during arachnid evolution. BMC Biol. 15:62. doi: 10.1186/s12915-017-0399-x.

Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert

M, Söding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of high-quality protein

multiple sequence alignments using Clustal Omega. Mol Syst Biol. 7:539. doi:

10.1038/msb.2011.75.

Slaidina M, Delanoue R, Gronke S, Partridge L, Léopold P. 2009. A *Drosophila* insulin-like

peptide promotes growth during nonfeeding states. Dev Cell. 17:874-884. doi:

10.1016/j.devcel.2009.10.009.

Seecof RL, Dewhurst S. 1974. Insulin is a *Drosophila* hormone and acts to enhance the
differentiation of embryonic *Drosophila* cells. Cell Differ 3: 63-70. doi: 10.1016/0045-
6039(74)90041-4.
Smit AB, Vreugdenhil E, Ebberink RH, Geraerts WP, Klootwijk J, Joosse J.1988. Growth-
controlling molluscan neurons produce the precursor of an insulin-related peptide. Nature.
31:535-538. doi: 10.1038/331535a0.
Tensen CP, Van Kesteren ER, Planta RJ, Cox KJ, Burke JF, van Heerikhuizen H, Vreugdenhil E.
1994. A G protein-coupled receptor with low density lipoprotein-binding motifs suggests a role
for lipoproteins in G-linked signal transduction. Proc Natl Acad Sci U S A. 91:4816-4820. doi:
10.1073/pnas.91.11.4816.
Terrapon N, Li C, Robertson HM, Ji L, Meng X, Booth W, Chen Z, Childers CP, Glastad KM,
Gokhale K, Gowin J, Gronenberg W, Hermansen RA, Hu H, Hunt BG, Huylmans AK, Khalil
SM, Mitchell RD, Munoz-Torres MC, Mustard JA, Pan H, Reese JT, Scharf ME, Sun F, Vogel
H, Xiao J, Yang W, Yang Z, Yang Z, Zhou J, Zhu J, Brent CS, Elsik CG, Goodisman MA,
Liberles DA, Roe RM, Vargo EL, Vilcinskas A, Wang J, Bornberg-Bauer E, Korb J, Zhang G,
Liebig J. 2014. Molecular traces of alternative social organization in a termite genome. Nat
Commun. 5:3636. doi: 10.1038/ncomms4636.
Vallejo DM, Juarez-Carreño S, Bolivar J, Morante J, Dominguez M. 2015. A brain circuit that
synchronizes growth and maturation revealed through Dilp8 binding to Lgr3. Science.
50:aac6767. doi: 10.1126/science.aac6767.
Veenstra JA. 2014. The contribution of the genomes of a termite and a locust to our
understanding of insect neuropeptides and neurohormones. Front Physiol. 5:454. doi:
10.3389/fphys.2014.00454.
Veenstra JA. 2016a. Neuropeptide evolution: Chelicerate neurohormone and neuropeptide genes
may reflect one or more whole genome duplications. Gen Comp Endocrinol. 229:41-55. doi:
10.1016/j.ygcen.2015.11.019.
Veenstra JA. 2016b. Similarities between decapod and insect neuropeptidomes. PeerJ :e2043.
doi: 10.7717/peerj.2043.
Veenstra JA. 2019. Coleoptera genome and transcriptome sequences reveal numerous differences
in neuropeptide signaling between species. PeerJ. 7:e7144. doi: 10.7717/peerj.7144. eCollection
2019.
Veenstra JA. 2020. Gonadulins, the fourth type of Insulin-related peptides in Decapods. bioRxiv.
doi: https://doi.org/10.1101/2020.03.19.998484.
Veenstra JA, Khammassi H. 2017. Rudimentary expression of RYamide in *Drosophila*
*melanogaster* relative to other *Drosophila* species points to a functional decline of this
neuropeptide gene. Insect Biochem Mol Biol. 83:68-79. doi: 10.1016/j.ibmb.2017.03.001.
Veenstra JA, Rombauts S, Grbić M. 2012. In silico cloning of genes encoding neuropeptides,
neurohormones and their putative G-protein coupled receptors in a spider mite. Insect Biochem
Mol Biol. 42:277-295. doi: 10.1016/j.ibmb.2011.12.009.
Wheeler DE, Buck N, Evans JD. 2006. Expression of insulin pathway genes during the period of
caste determination in the honey bee, *Apis mellifera.* Insect Mol Biol. 15:597-602. doi:
10.1111/j.1365-2583.2006.00681.x.

Yang CH, Belawat P, Hafen E, Jan LY, Jan YN. 2008. *Drosophila* egg-laying site selection as a

system to study simple decision-making processes. Science 319: 1679–1683. doi:

10.1126/science.1151842.

Yu N, Li J, Liu M, Huang L, Bao H, Yang Z, Zhang Y, Gao H, Wang Z, Yang Y, Van Leeuwen

T, Millar NS, Liu Z. 2019. Genome sequencing and neurotoxin diversity of a wandering spider

*Pardosa pseudoannulata* (pond wolf spider). BioRxiv. doi: https://doi.org/10.1101/747147.

Yu N, Han C, Liu Z. 2020. *In silico* identification of the neuropeptidome of the pond wolf spider

*Pardosa pseudoannulata.* Gen Comp Endocrinol. 285:113271. doi:

10.1016/j.ygcen.2019.113271.

Zhan S, Fang G, Cai M, Kou Z, Xu J, Cao Y, Bai L, Zhang Y, Jiang Y, Luo X, Xu J, Xu X,

Zheng L, Yu Z, Yang H, Zhang Z, Wang S, Tomberlin JK, Zhang J, Huang Y. 2020. Genomic

landscape and genetic manipulation of the black soldier fly *Hermetia illucens*, a natural waste

recycler. Cell Res. 2020 30:50-60. doi: 10.1038/s41422-019-0252-6.

# Figure 1

Gonadulin sequence alignment.

Note the very large sequence variability of the gonadulin propeptides. Cysteine residues are indicated in red and sequences are from Suppl. Spreadsheet 1.

```
Drosophila       -----------------------SFCSLERMKKFAMEACEHLFQADEGARRDRRSIEFAH--
Apis             ----------------ESTRQSLQCKN-IARDVIINSCKGPRMKRSPEKSDTSN-TDQIVK
Athalia          -----------------IDIYNCKM-KVRDMALKFCSDPAGTRKKRDIS--------IG
Pediculus        -------------------CDTKWIRLVYMMSCGKKKRDSSTSSFNE---------
Acyrthosiphon    ------------------CGDEFIKNMIVNVCGMGGIKRSFPD---NSRPDS---
Cimex            ------------------FMKPTCSKDYLRFMVATACSYAKREVSNDDVMKVVEEEFDKH
Gryllus          APAAAAALRSGGSDWEEQRVLHPTCDHDQFRNFMRLSCQYHRRREVRGAA----------
Timema           -----------------------CDKARIMNYIMYSCSKKKRSPADH----------LIY
Blattela         ----------------RPEYEDCNR-KIRRQILESCSSEKGKRSAE------------
Daphnia          -----------SIVAPGAPTKTICDPRAFRQAIVSICTFQRRDIHPISASPDR-------
Paralithodes-1   ------RPQQQHQHYHQEKRGVRLCSARDVKLIATYVCNLHRRSVRSVALDENSEGDTYIM
Paralithodes-2   ----------WPALRGRRSTSDLCTPREIRRIANDVCNIARRSI-----------------
Latrodectus      ------------FSRREKLFELVHVCTGHDLLNVIKRACAIHKRNE--------------
Pardosa-1        ---------------QQQRLDEMANVCSGQDLVHVIARACSIFKRSP--------------
Tetranychus      ------------------ENLIRVCTIKELQSARSRICSLYKRSY---------------

Drosophila       ---------------------------HHLNRLGSGKT------HNKHHYISRS----
Apis             ATISDV-IEENPVSIEIIEPEGSYHVQARMPLQHGFYGGYGLMPSRFGGIGQNYQS-TNFH
Athalia          ---AES------ESEVQ------RS-RDLEIGDPRF---------------------
Pediculus        ------------------------------TLLRNMPSDFYGA---------------
Acyrthosiphon    ------SLGLHFR---------------------------GMVT----DVE
Cimex            FDDVGNYVTADNEIFHAPLNYRNHKHWREVPFKLFKNKRTTDKFGNFQSDWQNFMKISDNP
Gryllus          ---------------------------------------------------------
Timema           PS-----------------------------ENWHL---------------------
Blattela         ---------------------------------------------------------
Daphnia          ------------------------------------YFYNVN-----------
Paralithodes-1   PGVGGSSVILPWRSLPPCDGPDGDCGGLNSADSPVQQTQRASPPQAPFNMAKRWYALHRFG
Paralithodes-2   -----RT---------------QL----TPSHHETDWPLLRRTPNG--IPAGHEGTNNYRFS
Latrodectus      ---------------------------------------------------------
Pardosa-1        ---------------------------------------------------------
Tetranychus      ---------------------------------------------------------

Drosophila       --------SYPMGGYLKV------TREHFNRLSELDIFPRY----------KPIKPHHEK
Apis             QSELITPGLEYMDTQYNGLSGGMYGSSLPMRPKYLLRSVKSDDLLGFNL--SPEELEELHD
Athalia          ----------------------GFASLNSGADLGLTL-TQNEE---------EDLFMTLSQ
Pediculus        VKYLKEN------DPQNPMFK-----DDQFLYR-ILEDGIHGFPD---QWEVDEINDD---
Acyrthosiphon    REFKHLVGDIEEVDLLLK-TSNNKNKLKEPNVTPLWR--PINLPGFRDD---RSRHPRIR
Cimex            GFGRFRRLLG-LEGASMPSGDIAGIPVPWMNREE------------------------
Gryllus          ---------------------------------------------------------
Timema           ----------DLPDAPSHNGGKLSEMLVVA--------MNPSHPVLYEWMMEGEEPTIFD
Blattela         --YDTPSLPMHDE--PLQNAPSSALLGRILGVPSQWT------ADDV-------------
Daphnia          ----------------ESESGTESGSQGERQTYD----------------KVVFTQLGK
Paralithodes-1   TAKGMQAEKENDITEDYRADSNSIDYAPSLSGALPLSELFPENVSKFIRVNYPKYVTSLLP
Paralithodes-2   LPSPLTLSSDVRPGVHSLIRSNTKLSQVLVPPLSNTIKARLASQAMVWR--LMLAGMVRRK
Latrodectus      ---------------------------------------------------------
Pardosa-1        ---------------------------------------------------------
Tetranychus      ---------------------------------------------------------

Drosophila       K------HRFKRDHSSRSY-NNIPYCCLNQ----CEEEFF---C---------------
Apis             EIGD-RMPRNSKNLNKKIFQQIAMKCCPN--ARLCYDNPRIIPCMGY-------------
Athalia          R---LSRAAKNKNQTMRMINAFAAKCCGENRHDECNDAANVIACP--------------
Pediculus        ---KIN----ENHRRSSKIDKLIEECCKVPPVRTCSEKTFKGAC-------------
Acyrthosiphon    G------DSGKATIIKRELMNDFRECCNKN----CSLKDLKRICGKK-----------
Cimex            ----LHSKGINGNLLYRDIRDTVMKCCTRN----CTVDEFKHLCG------------
Gryllus          ---AADDGGARVARSPSPEADLFAMCCERP----CNVEDFVGVCP------------
Timema           ---DLSKKTRSIFVYPEKAQRVIEQCCDK--EQYCDVNTFLGACK------------
Blattela         ---AVNNANRQVKRSPETIRQLMIDCCLAN----CSPDRFLGMC------------
Daphnia          ETLFFPVEVYTAKVKRHREFALHRKCCLSG----CYPMDFASVCMH----------
Paralithodes-1   HKGKRDKEMSLPVVVPRSLSAIRQDCCVK----CNAEDFFGAC------------
Paralithodes-2   ---RLNRRQTQHQSTTMTLEELRETCCNAQ----CSEQDFLAACS-----------
Latrodectus      ----HDPT-QPTHPLSIPLQDVSKVCCETP----CRISLFVRLCY-----------
Pardosa-1        ----SEPE-DILYPRSIPLKDIALVCCKMP----CRMSLFFRLCS-----------
Tetranychus      ----NPNSLMNWIRERRAISSIASSCCSPG----CPESLLTVGC-------------
```

# Figure 2

Insect ilp sequence similarity tree.

Note that the gonadulins and relaxins are well differentiated from the insulins and aILGFs, but that the latter are only partially separated from one another on the tree. Sequences are from Suppl. Spreadsheet 1.

# Figure 3

Radial arthropod ilp sequence similarity tree.

Note that the gonadulins cluster and are well separated from the other arthropod ilps. A more detailed sequence similarity tree with species names is present in the supplementary data as Fig. S1; sequences are from Suppl. Spreadsheet 1.

# Figure 4

aILGF sequence alignment.

Sequence alignment of some insect aILGF propeptides; where applicable only the isoforms with the arginine-rich sequences are shown. Note that these proteins have three different regions with sequence similarity: the insulin subsequence, the arginine-rich sequence containing an additional disulfide bridge and the C-terminal parts (underlined in blue), that is also somewhat conserved in *Gryllus rubens*, even though that species lacks an arginine-rich subsequence. Sequences are from Suppl. Spreadsheet 1; cysteine residues are indicated in red and the predicted disulfide bridges by lines.

# Figure 5

Intron-exon structures of aILGF genes.

(A) *Bombyx mori,* (B) *Tribolium castaneum,* (C) *Apis mellifera,* (D) *Blattella germanica*. In the top part for each species the gene structure with the exon represented as small colored boxes and the introns as a black line. The orange brown color in the first exon corresponds to the coding sequence of the signal peptide and the blue corresponds to the insulin sequence, while the red color corresponds to the coding sequence of the arginine-rich region that is alternatively spliced in the two isoforms produced from these genes. The yellow exons contains coding sequence for the $GTVX_1PX_2(F/Y)$ consensus sequence. The amino acid sequence coded by this alternatively spliced DNA sequence is indicated. The numbers 1 and 2 show the coding sequences of the two mRNA species produced from these gene using the same colors as for the gene structures. Note that the structures of these genes are very similar, with the major differences being the size of the introns, some of which are very large, as indicated by interruption sign in the gene structures, and the loss of an intron in *Apis*. The only other notable difference is that in the honey bee the second transcript is produced in a different fashion and only consists of one coding exon. The alternative splice site in this species have been indicated together with how this results in either splicing or the inclusion of a stop codon in the mRNA.

# Figure 6

Expression of aILGF and BIGFLP during development in *Bombyx mori*.

Number of reads per million present in SRAs of a single series of experiments where there are three experiments for each developmental stage. The three values corresponds to the maximum, minimun and average of those three. Scale to the left is for aILGF, scale to the right for BIGFLP; note that the number of BIGFLP reads is much larger than those for aILGF.

# Figure 7

Differential expression of *Bombyx* aILGF alternative splice forms.

Counts of reads specific for the long isoform of aILGF were plotted agains the total number of aILGF reads in all *Bombyx* SRAs (n=253). Samples with high total aILGF counts show different levels of expression of the long isoform. Identification of some of the salient samples in this figure is provided in Fig. S5.

# Figure 8

Evolution of the insect aILGF gene.

Schematic organization of insect aILGF genes in relation to their place on a simplified phylogenetic tree. The archetype of the insect aILGF gene consists of four coding exons. The first codes for the signal peptide (orange brown color) and the B-chain and part of the C-peptide, the second codes for the remainder of the C-peptide and the A-chain. The third coding exon codes for the first part of the C-terminal extension and has in the middle an alternative splice site allowing alternative splicing of the arginine rich sequene (in red). The last coding exon (yellow) codes for final part of the C-terminal extension that contains the $GTVX_1PX_2(F/Y)$ consensus sequence. Diptera have lost the third coding exon and the Eremoneura have also lost the fourth coding exon. Sequence comparison of the aILGF precursors from *Dasypogon, Hermetia, Condylostylus* and *Platypeza* are very similar, except for the loss of the last part (Fig. S6). The structures of the *Blattella, Apis, Tribolium* and *Bombyx* genes are from Fig. 5 and those of the *Oncopeltus* and *Aedes* genes from Fig. S3. The *Timema, Hermetia, Dasypogon, Ceratitis, Drosophila* and *Glossina* genes were produced in the same fashion. The *Condylostylus* gene could only partially constructed from genome and transcriptome SRAs. The *Platypeza* gene has been made transclucent, as is only inferred; it is on a transcript (GCGU01007956.1) and assumes that its structure is identical to those of its closest relatives.

# Figure 9

Arthropod relaxin sequence alignment.

Note how the relaxin propeptides are much better conserved than the either the gonadulins or the aILGFs. Also not that the exception is the single decapod relaxin that has also an additional cysteine residue. This is not a sequencing or other technical error, as the same residue is also found in several others decapods from different decapod orders (Fig. S7). Cysteine residues are indicated in red; sequences are provided in Suppl. Spreadsheet 1 and from Veenstra (2020).

# Figure 10

Synteny of arthropod ilp genes.

Schematic representation of relaxin, gonadulin and aILGF genes in three insect and two spider species. Genbank accession numbers are indicated below the species name. Arrowheads indicate the direction of transcription. The numbers in Kbp indicate the distances between the coding regions of neighboring genes.

# Figure 11

Phylogenetic tree of arthropod insulin RTKs.

Phylogenetic tree of various arthropod RTKs. Note that the decpods, chelicerates and insect RTKs evolved independently. Only branch probabilities of more than 0.900 have been indicated. Sequences are from (Veenstra, 2020) and others provided in Supp. Spreadsheet 1.

# Figure 12

Phylogenetic tree of putative arthropod insulin GPCRs.

Phylogenetic tree based exclusively on the transmembrane regions of various decapod LGRs that might function as receptors for insulin-related peptides. Only branch probabilities of more than 0.900 have been indicated. Sequences are from (Veenstra, 2020) and others provided in Supp. Spreadsheet 1.

## Table 1(on next page)

Gonadulin expression.

Numbers are the number of gonadulin half reads per million in one or more transcriptome SRAs. This is a selection of the data from Suppl. Spreadsheet 2.

# Table1. Gonadulin expression.

| Species | Gonadulin | Tissue/organ |
|---|---|---|
| *Blattella germanica* | 0.00 | Male heads |
| | 7.30 | Female gonads and fat body |
| | 4.10 | Male gonads and fat body |
| | 493.70 | Non-fecundated eggs |
| *Timema cristinae* | | |
| | 0.00 | Head |
| | 23.30 | Testis |
| | 46.90 | Ovary |
| *Rhodnius prolixus* | | |
| | 78.40 | Ovary |
| | 1.40 | Testis |
| | 0.20 | CNS |
| | 3.00 | Antenna |
| *Apis mellifera* | | |
| | 0.20 | Ovary of virgin queen |
| | 61.60 | Ovary of normal egg-laying queen |
| | 96.70 | Ovary of normal egg-laying inhibted queen |
| | 59.00 | Ovary of normal egg-laying recovered queen |
| | 7.10 | Antenna |
| | 4.10 | Second thoracic ganglion |
| *Steatoda grossa* | | |
| | 0.00 | Cephalothorax |
| | 7.10 | Ovary with eggs |
| | 5.90 | Minor ampullate silk glands |
| | 3.30 | Tubuliform silk glands |
| *Parasteatoda tepidariorum* | | |
| | 2.00 | Ovary from SRR1824489 |
| | 0.00 | Ovary from SRR8755633 |
| | 0.00 | Ovary from SRR8755634 |

Numbers are the number of gonadulin half reads per million in one or more transcriptome SRAs. This is a selection of the data from Suppl. Spreadsheet 2.