1   **TITLE:**

2   Onion thrips *Thrips tabaci* [Thysanoptera: Thripidae] reduces yields and THC in indoor grown

3   cannabis

4   **IDENTIFICATION OF AUTHORS:**

5   Frédéric McCune[1], Chad Murphy[2], James Eaves[3], and Valérie Fournier[1]

---

[1] Centre de recherche et d'innovation sur les végétaux, Université Laval, Québec, QC, Canada G1V 0A6; corresponding author: frederic.mccune.1@ulaval.ca.

[2] GreenSeal Cannabis Co., Stratford, ON, Canada N4Z 1H3

[3] Département de management/ Department of Management, FSA ULaval – Faculté des sciences de l'administration, Université Laval, Québec, QC, Canada G1V 0A6

6   **ABSTRACT**

Cannabis (*Cannabis sativa* L. [Rosales: Cannabaceae]) is a newly legalized crop and requires

deeper insights on its pest communities. In this preliminary study, we identified a thrips species

affecting indoor grown cannabis in Canada and tested its impact on plant yield. We used three

levels of initial infestation (zero, one, and five thrips) on individual plants grown in two growing

mediums: normal substrate or substrate containing the biostimulant *Bacillus pumilus*, Meyer and

Gottheil [Bacillales: Bacillaceae]. We found that the onion thrips, *Thrips tabaci* (Lindeman)

[Thysanoptera: Thripidae] is proliferating in indoor grown cannabis. Furthermore, our results

showed that fresh yields were higher for the plants that initially received zero thrips compared to

those that initially received five thrips. Moreover, the biostimulant did not help reduce the impact

of thrips. We highlight the importance for growers to carefully monitor thrips infestations in indoor

grown cannabis. Finally, we emphasize the need for more research related to the impact of pests

on cannabis yields and safe means of pest control for this strictly regulated crop.

19   **RÉSUMÉ**

Le cannabis (*Cannabis sativa* L. [Rosales: Cannabaceae]) est une culture nouvellement légalisée

et qui requiert des connaissances approfondies sur ses ravageurs. Dans cette étude préliminaire,

nous avons identifié une espèce de thrips affectant le cannabis cultivé à l'intérieur au Canada et

testé son impact sur le rendement des plants. Nous avons testé trois niveaux initiaux d'infestation

(zéro, un, et cinq thrips) sur des plants individuels cultivés dans deux terreaux : un substrat normal

ou un substrat contenant le biostimulant *Bacillus pumilus*, Meyer and Gottheil [Bacillales:

Bacillaceae]. Nous avons observé que le thrips de l'oignon, *Thrips tabaci* (Lindeman)

[Thysanoptera: Thripidae] prolifère dans le cannabis cultivé à l'intérieur. Nos résultats montrent

que le rendement des plants de cannabis est plus élevé pour les plants n'ayant pas reçu de thrips

comparativement aux plants sur lesquels cinq thrips ont initialement été inoculés. De plus, le

biostimulant n'a pas permis de réduire l'impact des thrips. Nous mettons en lumière l'importance

pour les producteurs de cannabis cultivé à l'intérieur de faire un suivi rigoureux de leurs

populations de thrips. Finalement, nous soulignons les besoins importants en recherches concernant

les ravageurs du cannabis, leurs impacts et le développement de méthodes de lutte dans cette culture

hautement règlementée.

INTRODUCTION

Cannabis (*Cannabis sativa* L. [Rosales: Cannabaceae]) was legalized for recreational purposes in

October 2018 in Canada and is still under strict prohibition in most of the world. Thus, there is a

severe lack of information regarding its growing practices (Eaves, Eaves, Morphy, & Murray,

2020; Wilson et al., 2019). This includes research related to the impact of pest species and the

means of controlling them (Cranshaw et al., 2019). Under the Cannabis Regulations and the Pest

Control Products Act, Health Canada only allows cannabis growers to use a limited number of

pesticide products. Consequently, companies rely mostly on biological control, but these

techniques are very costly, increase the risk of contaminating the final product with dead insect

parts, and yield uneven results.

More than 300 arthropod species have been identified on hemp and cannabis (Cranshaw et al.,

2019; McPartland, 1996a). On cannabis, the most predominant ones are sap-sucking arthropods

such as aphids, whiteflies, leafhoppers, mealybugs and various mites (Lago & Stanford, 1989;

McPartland, 1996a; Wilson et al., 2019). Recent reports of potential pests in cannabis include the

marmorated stink bug (*Halyomorpha halys*) (Britt, Pagani, & Kuhar, 2019) and two aphid species

(*Phorodon cannabis* and *Rhopalosiphum rufiabdominale*) (Lagos-Kutz, Potter, DiFonzo, Russell,

& Hartman, 2018). Despite this, it is believed that very few insects can actually cause significant

losses in commercial cannabis production (Dewey, 1913; McPartland, 1996a). In a recent survey,

growers from California reported from zero to over 25% crop damage caused by arthropods

(Wilson et al., 2019). Nonetheless, a large proportion of cannabis production occurs indoor or in

greenhouses, which provide environments that are particularly favourable for pests. If fact, in

Canada, Health Canada only started licensing outdoor area in October of 2019. Since most

published studies have focused on outdoor production, our current estimates of pest-risk posed to

cannabis producers may greatly underestimate the actual risk.

Thrips have been shown to be a major pest for many crops, most notably in greenhouses (Stuart,

Gao, & Lei, 2011) and can inflict both direct and indirect damage (Hao, Shipp, Wang,

Papadopoulos, & Binns, 2002; Pereira et al., 2017). Damage resulting from sucking or ovipositing

in the marketable plant parts, like fruits, correspond to direct damage (Shipp, Hao, Papadopoulos,

& Binns, 1998), while damage caused on non-marketable plant parts, like leaves, are considered

indirect damage (Diaz-Montano, Fuchs, Nault, Fail, & Shelton, 2011; German, Ullman, & Moyer,

1992). Thrips are found in indoor cannabis facilities (McPartland, 1996a) but we are not aware of

any studies investigating their impact on cannabis yields. Nevertheless, Cranshaw et al. (2019)

reports that thrips are common pests in hemp farms. For instance, Onion thrips, *Thrips tabaci*

(Lindeman) [Thysanoptera: Thripidae] have been frequently found on hemp in Colorado and can

cause important foliage damage on indoor grown plants (Cranshaw et al., 2019). Western flower

thrips, *Frankliniella occidentalis* (Pergande), tobacco thrips, *Frankliniella fusca* (Hinds) and

greenhouse thrips, *Heliothrips haemorrhoidalis* (Bouché) have also been found in hemp farms

(Cranshaw et al., 2019; Lago & Stanford, 1989; McPartland, 1996a).

Biostimulants are biological products that improve the productivity of plants. These products

are often a mixture of compounds derived from various organisms, such as bacteria, fungi, algae,

higher plants or animals, and frequently possess unexplained modes of action (Calvo, Nelson, &

Kloepper, 2014; Conant, Walsh, Walsh, Bell, & Wallenstein, 2017; Yakhin, Lubyanov, Yakhin, &

Brown, 2017). Specifically, the bacterium *Bacillus pumilus*, Meyer and Gottheil, [Bacillales:

Bacillaceae] is known for its growth promoting (de-Bashan, Hernandez, Bashan, & Maier, 2010;

Gutiérrez-Mañero et al., 2001; Probanza, Lucas, Acero, & Gutierrez Mañero, 1996) and antifungal

(Pérez-García, Romero, & de Vicente, 2011) properties. Furthermore, *B. pumilu*s successfully

suppressed larvae of *Scirpophaga incertulas* and *Bruchus dentipes* in laboratory conditions

(Rishad, Rebello, Shabanamol, & Jisha, 2017; Tozlu, Dadasoglu, Kotan, & Tozlu, 2011). These

results are likely explained by its high production of chitinase, an enzyme that can degrade the

chitin containing cell walls of insects and thus induce death (Rishad et al., 2017). Chitinase has

shown insecticidal properties against weevils (Laribi-Habchi, 2014) and aphids (Kim & Je, 2010).

Growing mediums enhanced with entomopathogenic bacteria represents a promising avenue

toward pest control and reduced use of pesticides. When added to a growing medium, *B. pumilus*

reduces the infestation level of fungus gnats (Diptera) in greenhouses, but shows inconclusive

results for the western flower thrips (Gravel & Naasz, 2019).

Hence, the objectives of this preliminary study were to identify the thrips species affecting

indoor cannabis production in Ontario, to determine the potential yield and quality losses associated

with their infestations, and, finally, to evaluate the impact of adding the biostimulant *B. pumilus* to

the growing substrate on the impacts of thrips.

MATERIALS AND METHODS

Experiment

The experiment was conducted in the autumn of 2019 in the commercial cannabis production

facility of GreenSeal Cannabis Company located in Stratford, Ontario, Canada. We used 60 clones

(approximately two weeks old) of cannabis (*C. sativa* var. Green Crack) to test the impact of three

initial levels of thrips infestation (zero, one or five thrips) and two growing substrates (normal or

biostimulant). Ten cannabis clones, acting as ten replicas, were randomly assigned to each

combination of infestation level and growing substrate. All clones were planted in seven inches

square pots (4L) using one of the two types of substrate. The first ("normal substrate") was a

fibrous, peat-moss substrate with perlite (Pro-Mix HP Mycorrhizae, Premier Tech). The second

("biostimulant substrate") was the normal substrate with the addition of the biostimulant *Bacillus*

pumilus (strain GHA180) (Pro-Mix HP Biostimulant + Mycorrhizae, Premier Tech). Plants were

propagated in a quarantine room and we visually inspected them for predatory mite or thrips. As

an additional precaution, we carefully used a spray-bottle and cloth to wipe each individual leaf to

ensure no arthropods were on them.

As thrips can reproduce asexually (Morison, 1957; Stuart et al., 2011) inoculating a single

immature thrips can lead to an significant population overtime. It is thus not possible to control for

their final infestation levels. Even though we do not think that controlling the initial infestation

levels will result in consistent levels of infestation at the end of the experiment, we consider that it

provides a valuable insight about the impact of a pest (Torres-Vila, Rodríguez-Molina, & Lacasa-

Plasencia, 2003). In this way, we inoculated the plants with zero, one and five thrips to represent

respectively control, low and high levels of infestation (Hao et al., 2002). Thrips used for the

experiment were collected directly from the production area of the facility with entomological

mouth aspirators. We targeted what we believed to be late instar larvae. Thrips were carefully

inoculated on each plant with fine brushes. All plants, including the controls with no inoculated

thrips, were then covered with Nitex (150 µm mesh) bags that were supported by stainless-steel

frames and tightly secured around the pots by elastics bands. The fine meshes of the Nitex bags

help prevent thrips from escaping and non-experimental pests or predatory mites from entering.

Nonetheless, since this experiment was conducted in a production facility, rather than inside a

university lab (It is still very difficult to receive a cannabis research license in Canada.), we did

expect some cross contamination and thus recorded leaf damage for all plants. Two drippers were

threaded under the elastic bands for irrigation purpose. We considered monitoring the thrips levels

over the course of the experiment but decided the risk of allow arthropodes to enter or escape was

too high.

The plants were grown in the facility's quarantine room, where all plants were placed on two-

levels shelves. Five plants of each treatment were placed on each level of the shelves following a

completely randomized design, so that 30 plants were located on the top level and 30 plants on the

bottom level. Plants were placed in two rows of 15 plants on each level. Shelves were equipped

with broad-spectrum LED lighting (Voltserver High-Intensity Lighting Platform). Light intensity

was gradually increased each week from 25% intensity (average of around 500 PPFD) to a

maximum of 50% intensity (average of around 1,080 PPFD). Plants were then kept under

commercial cultivation conditions (day/night temperature of 25°C/21°C +/- 2, 12 hours of daylight,

50% +/- 5 RH, and CO2 at ambient levels). Using a short vegetative period combined with these

environmental conditions are typical for growers who follow a "sea of green" growing strategy, as

GreenSeal Cannabis does. The exception is CO2 concentration levels, which were maintained at

ambient levels for the experiment. Plants were watered by drip irrigation about every two days

with approximately 1 L of water/plant.

All 60 plants were grown in the cages under aforementioned conditions for eight weeks. At the

end of the eight-week period, the plants had reached the end of the flower stage. The fresh

inflorescences were then harvested following normal commercial methods and weighed for each

plant (Pennsylvania 7600 Series Bench Scale 4536 g x 0.5 g). In order to measure total THC levels,

we took flower samples from three plants from each treatment with zero or five thrips. The samples

for each treatment were then blended together and analyzed using HPLC analysis conducted by

A&L Canada Laboratories Inc. located in London, Ontario. We therefore obtained a single THC

measure for each control and high infestation treatment.

Multiple rows of plants in the two production rooms of the cannabis facility were checked to

collect and identify all thrips species occurring in the facility. All collected specimens were

observed under a stereomicroscope and appeared to be similar. Multiple thrips specimens at both

adult and larval stages were collected and sent to the expert insects taxonomist from the Laboratoire

d'expertise et de diagnostic en phytoprotection (LEDP) of the Ministère de l'Agriculture, des

Pêcheries et de l'Alimentation du Québec for identification (Palmer, Mound, Du Heaume, & Betts,

1989), who has stored the vouchers. No other pests than thrips were observed in the facility.

Statistical analyses
Fresh yield data were analyzed with R (R Core Team, 2019). We used generalized least squares

fitted linear models and linear mixed-effects models (package "nlme" (Pinheiro, Bates, DebRoy,

Sarkar, & R Core Team, 2019)). Fresh yield was used as the response variable, while both the

number of thrips initially inoculated and the substrate type were explanatory variables. All models

included all interactions between our explanatory variables. We first computed generalized least

squares fitted linear models and then compared these to linear mixed-effects models in which the

shelves' level (upper or lower) was treated as a random effect. This allows us to take into account

a potential difference in temperature or growing conditions between levels. We thereafter

compared both types of models based on their Akaike information criterion (AIC). The more

complex linear mixed-effects models including the shelves' level had a higher AIC than the more

simple models, indicating less accuracy of the model. We therefore only present results from the

generalized least squares fitted linear models in the result section. We used an ANOVA to test the

effect of both the number of inoculated thrips and the type of substrate on the fresh yield. We

changed the reference level and fitted models once more to evaluate the effect of the multiple levels

of our explanatory variables that were identified as significant in the ANOVA.

172   **RESULTS**

All thrips specimens collected in the GreenSeal Cannabis Company's facility were identified as

being onion thrips, *Thrips tabaci* (Lindeman). The final fresh yield of our individual plants varied

between 36.81 g and 195.84 g. Three plants died at the start of the experiment from transplant

shock. We did not have replacement plants. They were respectively under treatments zero thrips-

biostimulant, one thrips-normal, and five thrips-normal. Five others grew far more slowly than the

average plant, which may indicate they were somehow stunted. Those were respectively under

treatments zero thrips-biostimulant, zero thrips-normal (two plants), one thrips-biostimulant, and

one thrips-normal. The plants that died or had a reduced growth were thus represented in almost

all treatments, but slightly more in controls and low infestation treatments. Nonetheless, a boxplot

revealed that only three of these observations were actual outliers. We removed the dead plants

from our dataset and performed all analyses with and without the stunted plants. We obtained very

similar results both times and thus decided to include the stunted plants in all analyses. We observed

thrips on most plants at the end of the experiment, even on many control plants. However, we are

not very concerned about this contamination since the relative amount of damage is representative

of our desired levels of infestation. For example, control plants had very little damage compared

to what was observed on the treatment plants. For the plants inoculated with five thrips, the total

THC level was of 17.6% for the normal substrate and 17.77% for the biostimulant substrate. For

the zero thrips treatment, the total THC level was 19.34% for the normal substrate and 19.62% for

the biostimulant substrate. We limited our THC measurements to those four samples and thus

cannot provide statistics here. Those THC levels nevertheless indicate a possible reduction in THC

when plants are under high infestation of thrips.

The number of thrips had a significant effect on the final yield of the cannabis plants (ANOVA

$F(2, 51) = 7.1062$, $P = 0{,}0019$; Fig. 1). Specifically, the yields were lower for plants that were

initially inoculated with five thrips compare to the plants that had no thrips ($t(51) = -2.569502$,

$P = 0.0132$). Average yield was 30.68% higher for plants that were not inoculated with thrips

(130.15 g per plant compared to 90.22 g). This result highlights the relevance of keeping stunted

plants in the analysis. Indeed, as the prevalence of stunted plants was slightly higher in the control

and low infestation treatment, the effect of including these low yielding plants in the models was

to reduce the potential negative impact of thrips. As we found an opposite trend, it therefore

reinforces the hypothesis that thrips negatively impact yields. The effect of the substrate type was

not significant (ANOVA $F(1, 51 = 3.5769$, $P = 0.0643$; Fig. 2) while the interaction between the

infestation level and the substrate type (ANOVA $F(2, 51 = 0.0506$, $P = 0.9507$) had no effect on

the final yields.

206   **DISCUSSION**

In this preliminary article, we report the first quantification of yield loss from damage caused by

onion thrips for indoor grown cannabis. We estimated yield losses that are higher than those

reported for outdoor cannabis growers in California in a survey on all pests (Wilson et al., 2019).

Indoor growing environments are particularly favourable for thrips and, thus, likely increase the

risk associated with thrips' outbreaks. A similar study using three initial infestation levels of thrips

found that the western flower thrips (*F. occidentalis* (Pergande)) can significantly reduce yields for

greenhouse grown cucumbers, as well as the plant's growth and photosynthesis rates (Hao et al.,

2002). A multitude of factors such as the crop nutritional status, its growing condition and the

prevalence of pests and diseases influence yields in a crop and thus make describing yields as a

function of a precise pest infestation very difficult (Pereira et al., 2017). This is even more true for

pests inflicting indirect damage such as thrips (Hao et al., 2002; Pereira et al., 2017). However, as

growing conditions are highly controlled in indoor cannabis production, as they were standardized

in between our plants, as climatic variations are minimal, and as our plants were all clones equally

treated, we believe the differences observed in this study are most likely due to differences in

infestations rates.

The onion thrips have a very diversified range of hosts (Diaz-Montano et al., 2011; Nault, Kain,

& Wang, 2014; Stuart et al., 2011) and has long been recognized as a greenhouse pest (Morison,

1957). Considering it has been found on hemp (Cranshaw et al., 2019), it is not surprising that

indoor grown cannabis can be added to this long list of hosts. Nonetheless, this is a new piece of

valuable information for producers. In onions, it can notably impair bulb weight (Ghosheh & Al-

Shannag, 2000) and reduce yields, sometime by more than 50% (Diaz-Montano et al., 2011;

Fournier, Boivin, & Stewart, 1995), especially since as little as 10 thrips per plant is sufficient to

decrease yields by 7% in greenhouses (Kendall & Capinera, 1987). Onion thrips are particularly

known to feed on leaves, causing photosynthesis reduction, distorted plant parts, and reduced bulbs

size as well as transmitting viruses, such as the Iris yellow spot virus (family Bunyaviridae, genus

Tospovirus, IYSV) (Diaz-Montano et al., 2011; Gent et al., 2006; Wu et al., 2013). Damage

consisting of yellow dot-shaped scars were observed both on the leaves of our experimental plants

and on production plants under outbreak pressure (Fig. 3). These injuries can be considered indirect

damage and were almost certainly inflicted by the onion thrips. We believe our observations

correspond to the "serious foliage damage" reported by Cranshaw et al. (2019) on hemp. Even

though we did not investigate damage extent or photosynthetic rate in our cannabis plants, it can

be expected that our reduced yields originate from those indirect feeding damage. Similar injuries

and scars are known to reduce the photosynthetic ability of leaves in onions (Diaz-Montano et al.,

2011). Little information is available about the transmission of viruses to cannabis plants by thrips

but McPartland (1996b) mention that viruses can greatly reduce yields in cannabis and that the

onion thrips is one of the worst vector of viruses in this crop. Besides yield losses, we also highlight

potential decreases in product value through reduced levels of total THC from highly infested

plants.

The use of a growing medium with *B. pumilus* did not improved the plants' strength. It is

consistent with previous experiments with the same enhanced growing medium on the control of

the western flower thrips (*F. occidentalis*) (Gravel & Naasz, 2019). However, our results were

nearly significant, and we still consider that growing medium enhanced with microbial

biostimulants represents a promising avenue for integrated pest management in greenhouse or

indoor productions. Their use should be investigated more in indoor cannabis production and for

various pests.

In conclusion, we showed that the onion thrips is present in indoor grown cannabis in Canada

and that it represents an economic threat. We observed damage caused by thrips feeding on leaves

and experimentally found a link between infestation levels and final fresh yields, in addition to a

possible reduction of total THC levels under high infestation. This study was preliminary and

should motivate further experiments. As chemical means of control are very limited for indoor

grown cannabis, we recommend strict monitoring programs for indoor cannabis producers to avoid

economic losses. We show that thrips potentially represent a major threat to product quality and

yields. In this way, we suggest more research on cannabis pests to identify all pest species in various

growing setting, including the range of damage they can cause and their economic thresholds. Only

this can subsequently lead to the development of management programs and the development of

safe and affordable control methods.

263   **ACKNOWLEDGMENTS**

264   We thank the team from GreenSeal Cannabis Co., especially Kurt Fisher for providing plant care,

265   and Chris Murray and Peter Reeves for helping build the cages. We are also grateful to Premier

266   Tech for providing the Pro-Mix HP Biostimulant + Mycorrhizae. This research was supported by

267   funding from the NSERC's Engage Grants for Universities program, no 536612-2018.

268   **REFERENCES**

Britt, K. E., Pagani, M. K., & Kuhar, T. P. (2019). First report of brown marmorated stink bug (Hemiptera:
Pentatomidae) associated with *Cannabis sativa* (Rosales: Cannabaceae) in the United States. *Journal of*
Integrated Pest Management, 10(1), 3. doi:10.1093/jipm/pmz014

Calvo, P., Nelson, L., & Kloepper, J. (2014). Agricultural uses of plant biostimulants. *Plant and Soil, 383*(1), 3-41.
doi:10.1007/s11104-014-2131-8

Conant, R., Walsh, R., Walsh, M., Bell, C., & Wallenstein, M. J. J. H. F. (2017). Effects of a microbial biostimulant,
Mammoth PTM, on *Cannabis sativa* bud yield. *Journal of Horticulture, 4*, 2376-0354.1000191.

Cranshaw, W., Schreiner, M., Britt, K., Kuhar, T. P., McPartland, J., & Grant, J. (2019). Developing insect pest
management systems for hemp in the United States: A work in progress. *Journal of Integrated Pest*
Management, 10(1), 11. doi:10.1093/jipm/pmz023

de-Bashan, L. E., Hernandez, J.-P., Bashan, Y., & Maier, R. M. (2010). *Bacillus pumilus* ES4: Candidate plant growth-
promoting bacterium to enhance establishment of plants in mine tailings. *Environmental and Experimental*
Botany, 69(3), 343-352. doi:10.1016/j.envexpbot.2010.04.014

Dewey, L. H. (1913). Hemp. In *Yearbook of the United States Department of Agriculture* (pp. 283-346). Washington,
D. C.

Diaz-Montano, J., Fuchs, M., Nault, B. A., Fail, J., & Shelton, A. M. (2011). Onion thrips (Thysanoptera: Thripidae):
a global pest of increasing concern in onion. *Journal of economic entomology, 104*(1), 1.
doi:10.1603/EC10269

Eaves, J., Eaves, S., Morphy, C., & Murray, C. (2020). The relationship between light intensity, cannabis yields, and
profitability. *Agronomy Journal, 112*(2), 1466-1470. doi:10.1002/agj2.20008

Fournier, F., Boivin, G., & Stewart, R. K. (1995). Effect of *Thrips tabaci* (Thysanoptera, Thripidae) on yellow onion
yields and economic thresholds for its management. *Journal of economic entomology, 88*(5), 1401-1407.
doi:10.1093/jee/88.5.1401

Gent, D., Du Toit, L., Fichtner, S., Mohan, S., Pappu, H., & Schwartz, H. (2006). Iris yellow spot virus: An emerging
threat to onion bulb and seed production. *PLANT DISEASE, 90*(12), 1468-1480. doi:10.1094/PD-90-1468

German, T. L., Ullman, D. E., & Moyer, J. W. (1992). Tospoviruses: diagnosis, molecular biology, phylogeny, and
vector relationships. *Annual review of phytopathology, 30*, 315.

Ghosheh, H. Z., & Al-Shannag, H. K. (2000). Influence of weeds and onion thrips, *Thrips tabaci* (Thysanoptera :
Thripidae), on onion bulb yield in Jordan. *Crop Protection, 19*(3), 175-179. doi:10.1016/s0261-
2194(99)00084-8
Gravel, A., & Naasz, R. (2019). Development of new insect suppression solutions for greenhouse production. *Acta
Horticulturae, 1266*(1266), 121-128. doi:10.17660/ActaHortic.2019.1266.17
Gutiérrez-Mañero, F. J., Ramos-Solano, B., Probanza, A. N., Mehouachi, J., R. Tadeo, F., & Talon, M. (2001). The
plant-growth-promoting rhizobacteria *Bacillus pumilus* and *Bacillus licheniformis* produce high amounts of
physiologically active gibberellins. *Physiologia Plantarum, 111*(2), 206-211. doi:10.1034/j.1399-
3054.2001.1110211.x
Hao, X., Shipp, J. L., Wang, K., Papadopoulos, A. P., & Binns, M. R. (2002). Impact of western flower thrips on
growth, photosynthesis and productivity of greenhouse cucumber. *Scientia Horticulturae, 92*(3-4), 187-203.
doi:10.1016/s0304-4238(01)00301-6
Kendall, D. M., & Capinera, J. L. (1987). Susceptibility of onion growth stages to onion thrips (Thysanoptera:
Thripidae) damage and mechanical defoliation. *Environmental Entomology, 16*(4), 859-863.
doi:10.1093/ee/16.4.859
Kim, J., & Je, Y. (2010). A novel biopesticide production: attagel-mediated precipitation of chitinase from *Beauveria
bassiana* SFB-205 supernatant for thermotolerance. *Applied Microbiology and Biotechnology, 87*(5), 1639-
1648. doi:10.1007/s00253-010-2543-1
Lago, P. K., & Stanford, D. F. (1989). Phytophagous insects associated with cultivated marijuana, *Cannabis sativa*, in
Northern Mississippi. *Journal of Entomological Science, 24*(4), 437-445. doi:10.18474/0749-8004-24.4.437
Lagos-Kutz, D., Potter, B., DiFonzo, C., Russell, H., & Hartman, G. L. (2018). Two aphid species, *Phorodon cannabis
and *Rhopalosiphum rufiabdominale*, identified as potential pests on industrial hemp, *Cannabis sativa* L., in
the US Midwest. *Crop Forage & Turfgrass Management, 4*(1), 3. doi:10.2134/cftm2018.04.0032
Laribi-Habchi, H. (2014). Effectiveness of a biological insecticide derived from scorpion fish offal against the chickpea
weevil. *Journal of Bioprocessing & Biotechniques, 04*(02). doi:10.4172/2155-9821.1000148
McPartland, J. M. (1996a). Cannabis pests. *Journal of the International Hemp Association, 3*(2), 49, 52-55.
McPartland, J. M. (1996b). A review of Cannabis diseases. *Journal of the International Hemp Association, 3*(1), 19-
23.
Morison, G. D. (1957). A review of British glasshouse Thysanoptera. *Transactions of the Royal Entomological Society
of London, 109*(16), 467-520. doi:10.1111/j.1365-2311.1957.tb00334.x
Nault, B. A., Kain, W. C., & Wang, P. (2014). Seasonal changes in *Thrips tabaci* population structure in two cultivated
hosts. *PLoS ONE, 9*(7). doi:10.1371/journal.pone.0101791
Palmer, J. M., Mound, L. A., Du Heaume, G. J., & Betts, C. (1989). *CIE guides to insects of importance to man. 2.
Thysanoptera.*: CAB, International Institute of Entomology.
Pereira, P. S., Sarmento, R. A., Galdino, T. V. S., Lima, C. H. O., dos Santos, F. A., Silva, J., . . . Picanco, M. C.
(2017). Economic injury levels and sequential sampling plans for Frankliniella schultzei in watermelon crops.
Pest Management Science, 73(7), 1438-1445. doi:10.1002/ps.4475
Pérez-García, A., Romero, D., & de Vicente, A. (2011). Plant protection and growth stimulation by microorganisms:
biotechnological applications of Bacilli in agriculture. *Current Opinion in Biotechnology, 22*(2), 187-193.
doi:10.1016/j.copbio.2010.12.003
Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., & R Core Team. (2019). _nlme: Linear and nonlinear mixed effects
models_. R package version 3.1-140. Retrieved from https://CRAN.R-project.org/package=nlme
Probanza, A., Lucas, J., Acero, N., & Gutierrez Mañero, F. (1996). The influence of native rhizobacteria on european
alder (*Alnus glutinosa* (L.) Gaertn.) growth. *Plant and Soil, 182*(1), 59-66. doi:10.1007/BF00010995
R Core Team. (2019). R: A language and environment for statistical computing. Vienna, Austria. Retrieved from
https://www.R-project.org/
Rishad, K. S., Rebello, S., Shabanamol, P. S., & Jisha, M. S. (2017). Biocontrol potential of halotolerant bacterial
chitinase from high yielding novel *Bacillus Pumilus* MCB-7 autochthonous to mangrove ecosystem. *Pesticide
Biochemistry and Physiology, 137*, 36-41. doi:10.1016/j.pestbp.2016.09.005
Shipp, J. L., Hao, X., Papadopoulos, A. P., & Binns, M. R. (1998). Impact of western flower thrips (Thysanoptera:
Thripidae) on growth, photosynthesis and productivity of greenhouse sweet pepper. *Scientia Horticulturae,
72*(2), 87-102. doi:10.1016/S0304-4238(97)00130-1
Stuart, R. R., Gao, Y.-L., & Lei, Z.-R. (2011). Thrips: Pests of concern to China and the United States. *Agricultural
Sciences in China, 10*(6), 867-892. doi:10.1016/S1671-2927(11)60073-4

Torres-Vila, L. M., Rodríguez-Molina, M. C., & Lacasa-Plasencia, A. (2003). Impact of Helicoverpa armigera larval
density and crop phenology on yield and quality losses in processing tomato: developing fruit count-based
damage thresholds for IPM decision-making. *Crop Protection, 22*(3), 521-532. doi:10.1016/S0261-
2194(02)00205-3
Tozlu, E., Dadasoglu, F., Kotan, R., & Tozlu, G. (2011). Insecticidal effect of some bacteria on *Bruchus dentipes* Baudi
(Coleoptera: Bruchidae). *FRESENIUS ENVIRONMENTAL BULLETIN, 20*(4), 918-923.
Wilson, H., Bodwitch, H., Carah, J., Daane, K., Getz, C., Grantham, T. E., & Van, B. (2019). First known survey of
cannabis production practices in California. *California Agriculture, 73*(3-4), 119-127.
doi:10.3733/ca.2019a0015
Wu, S., Gao, Y., Xu, X., Zhang, Y., Wang, J., Lei, Z., & Smagghe, G. (2013). Laboratory and greenhouse evaluation
of a new entomopathogenic strain of *Beauveria bassiana* for control of the onion thrips *Thrips tabaci*.
Biocontrol Science and Technology, 23(7), 794-802. doi:10.1080/09583157.2013.794896
Yakhin, O. I., Lubyanov, A. A., Yakhin, I., & Brown, P. (2017). Biostimulants in plant science: A global perspective.
Frontiers in Plant Science, 7. doi:10.3389/fpls.2016.02049

LEGENDS OF FIGURES

Fig. 1 Mean final fresh yields of the cannabis plants after eight weeks according to the

number of inoculated onion thrips (*Thrips tabaci*). Error bars represent 95% confidence intervals.

Fig. 2 Mean final fresh yields of the cannabis plants after eight weeks according to the

substrate type used (normal or with *Bacillus pumilus* biostimulant). Error bars represent 95%

confidence intervals.

Fig. 3 Example of leaf damage observed on the experimental plants and on production

plants in the facility.

374   **RUNNING TITLE:**

375   Onion thrips in cannabis