An Optimal Lysis Time Maximizes Bacteriophage Fitness in Quasi-continuous Culture

Sherin Kannoly[1], Abhyudai Singh[2,*], John J. Dennehy[1,3,*]
[1]Biology Department, Queens College of the City University of New York, NYC, NY, USA
[2]Department of Electrical and Computer Engineering, University of Delaware, Newark, DE,
USA
[3]The Graduate Center of the City University of New York, NYC, NY, USA

17   *For correspondence: absingh@udel.edu, john.dennehy@qc.cuny.edu

18   **Key words**: adaptation, bacteriophage lambda, event timing, holin, life history traits, optimality

19   models, optimization, timing of reproduction

20   **Running Head:** Phage λ Optimal Lysis Time

ABSTRACT

Optimality models have a checkered history in evolutionary biology. While optimality models have been successful in providing valuable insight into the evolution of a wide variety of biological traits, a common objection is that optimality models are overly simplistic and ignore organismal genetics. We revisit evolutionary optimization in the context of a major bacteriophage life history trait, lysis time. Lysis time refers to the period spanning phage infection of a host cell and its lysis, whereupon phage progeny are released. Lysis time, therefore, directly determines phage fecundity assuming progeny assembly rate is maximized. Noting that previous tests of lysis time optimality rely on batch culture, we implemented a quasi-continuous culture system to observe productivity of a panel of isogenic phage λ mutants differing in lysis time. We report that λ phage productivity in our experiments is maximized around an optimal lysis time of 63 min, which is the lysis time of the λ "wildtype" strain. We discuss this finding in light of recent results that lysis time variation is also minimized in the λ "wildtype" strain.

INTRODUCTION

Evolutionary biologists have long been interested in the power of natural selection to refine biological adaptations (Smith 1978; Endler 1986; Lewontin 1989; Crow 1993). Generally, views on the power of natural selection fall into two camps: one emphasizing the ability of natural selection to optimize biological traits (Endler 1986; Parker and Smith 1990) and the other stressing the limitations of natural selection in face of evolutionary tradeoffs and genetic details (Lewontin 1989; Crow 1993) affecting phenotypic traits. Several studies have employed bacteriophages applied optimization principles to theoretically predict how these traits will evolve in response to changing ecological conditions (Abedon 1989; Wang et al. 1996; Abedon et al. 2001; Bull et al. 2004; Bull 2006; Bonachela and Levin 2014).

However, despite clear predictions, experimental tests have generally failed to confirm theoretical predictions. T7, ST-1, and ΦX174 phages with theoretically predicted suboptimal lysis times generally failed to evolve predicted optimal lysis times (Heineman et al. 2007; Chantranupong and Heineman 2012). While the evolution of T7 phage with a deleted lysin gene qualitatively supported optimality predictions, it was noted that abolition of the lysis function did not extend the latent period, thus had no effect on the expected tradeoff between latent period and burst size (Heineman et al. 2007). ST-1 qualitatively, but not quantitatively, adapted as predicted by modeling, and ΦX174 failed to adapt at all (Chantranupong and Heineman 2012). In another study, experimental evolution of phage RB69 did result in selection for shorter lysis time mutants in high host density cultures, but it is unclear whether the lysis time of these mutants were optimal for the experimental conditions (Abedon et al. 2003). A study employing isogenic lambda phages with different lysis times found the model predictions for both the optimum and fitness to be different from the experimental estimations (Wang 2006).

However, we note that these studies used batch cultures to examine host-phage

interactions, and phage fitnesses were estimated using absolute growth rates. These conditions

may not effectively mimic the kinds of conditions experienced by phages in their natural habitats.

To revisit the question of phage optimal lysis time, we employed a different approach, which was

inspired by Bonachela *et. al* (2014), who modeled host-virus interactions in natural environments

in terms of steady state conditions, such as those seen in continuous culture systems. A typical

example of a continuous culture system is a two-stage chemostat. In the first stage, a chemostat

receives a flow of fresh media to maintain a continuous culture of bacteria. Fresh host bacteria

from the first chemostat are fed into a second chemostat containing a population of phages. Both

host cells and phages are washed out from the second chemostat at a specified rate. This system

can mimic steady states that exist in natural environments, such as the gastrointestinal tract where

peristaltic movements can maintain a continuous flow.

In this study, we maintained phage λ and its host, *Escherichia coli*, under quasi-steady state

conditions. To initiate experiments, λ holin mutants differing in their lysis times (Dennehy and

Wang 2011; Singh and Dennehy 2014; Kannoly et al. 2020) were added to exponentially

replicating batch cultures of *E. coli*. After a short period of growth, λ phages were filtered from

host cells, and a small fraction was transferred into a fresh exponential culture. At no time did

these phages experience stationary phase host cells during the course of the experiment. In

principle, differences in lysis times should result in differences in phage productivity (Levin et al.

1977; Wang 2006). We predicted that there exists an optimum for lysis time that maximizes phage

production under these quasi-steady state conditions. By contrast, strains with suboptimal lysis

times should show declining phage populations in this experimental context. Our results show that

phage production in these quasi-steady state conditions is maximized at lysis times approaching

63 min, which, perhaps not coincidentally, is the lysis time of the phage λ wildtype. These results

suggest an optimum lysis time that maximizes phage progeny production.

RESULTS AND DISCUSSION

Lysis is the last step in the infectious cycle of lytic bacteriophage, resulting in the release

of virions. Lysis timing in many phage is controlled by a single protein, holin, which accumulates

in the cell membrane up to a genetically determined time, whence it nucleates generating

membrane-permeabilizing holes (Wang et al. 2000, 2003; Dennehy and Wang 2011; Singh and

Dennehy 2014; Kannoly et al. 2020).

To study the effect of lysis time on phage productivity (i.e., total phage reproduction during

a fixed growth period), we employed a panel of λ phages harboring mutations in the holin gene *S*,

which delay or hasten the lysis event following induction of the lytic cycle. These λ mutants were

used to initiate batch cultures containing a fixed concentration of *E. coli* cells at a multiplicity of

infection (MOI) of 0.001, then purified, diluted 1000-fold, and transferred to fresh cultures at 3 h

intervals, generating quasi-steady state conditions. A 3 h infection allows at least three generations

for wild type λ such that the MOI ≈ 1 at the end of phage growth. A low MOI also minimizes

multiple infections per cell and ensures multiple generations without depleting host cells to such

an extent that the cell number becomes a limiting factor for phage growth. Furthermore, the short

duration of growth prevents *E. coli* densities from being sufficiently high enough to enter

stationary phase.

Ideally, this study would be performed using actual chemostats, but we choose not to

pursue this option because it would limit the number of holin mutants and degree of replication

that we could easily manage. Moreover, it is well known that *E. coli* growth is consistent (i.e., in

a steady state) as long as exponential growth conditions are maintained (Fishov et al. 1995) so, in

physiological terms, *E. coli* growth is not inherently different between chemostats and our quasi-

steady state transfers. Such a steady state achieved by introducing fresh batch cultures ensures a

balanced exponential phase throughout the transfers. The cells in exponential phase are in the same

physiological state characterized by relatively similar cell volumes and macromolecular

composition. This aspect ensures that the growth parameters such as cell density (number of cells

per milliliter) and the concentration of key macromolecules such as proteins, RNA, DNA, etc.,

increase exponentially with precisely the same doubling time (Sezonov et al. 2007).

Pilot studies were performed by initiating wild type λ infections of host cells in exponential

phase. Cells exceeded phage by 1000-fold minimizing the time required for phage to encounter

host cells for attachment and ensuring that the MOI was close to one at the end of the first 3-h

transfer. Experiments where the growth period was less than 3 h per transfer and the dilution

between transfers was than 1000-fold resulted in lower phage numbers and increased likelihood

of phage population extinction after a few transfer cycles, especially in fast lysing strains (data not

shown). On the other hand, experiments where the growth period was longer than 3 h resulted in

higher phage yields, and the host populations were depleted before the end of the growth period,

especially in strains with long lysis times.

For the subsequent serial transfers after the initial growth period, phages were diluted 1000-

fold and this dilution rate was kept constant throughout the transfers. The transfer conditions were

adjusted using the wild type strain to obtain increasing productivity after each transfer until it peaks

after the fifth transfer. Subsequent transfers did not result in any significant increase in productivity

primarily due to coinfections and faster host cell depletion (data not shown). Moreover, five

transfers would result in approximately fifteen phage generations, beyond which phage adaptation

would begin to impact our results. It is important to note that unlike previous studies, our study does not aim to explore evolution of an optimum lysis time as a result of phenotypic adaptation. Instead, we aim to test the productivity (i.e., total phage reproduction during a fixed growth period) of genotypes varying in their lysis times under dynamic environments, which are subject to constant dilution or wash out. To this end, the experimental protocol was designed to mimic conditions were both phages and cells would be washed out at regular intervals.

Five such serial transfers were completed, and the phage titers were obtained after every transfer (Fig. 1). Fig. 2 shows phage titers after every transfer for all the strains and Fig. 3 compares phage productivity only for the fifth transfer for all strains. Our data show that phage progeny production following five transfers was maximized across a broad range spanning 60–100 min (Fig. 2; Fig. 3). Interestingly the phage λ wildtype strain falls within this range (Fig. 2; Fig. 3). Outside this range, phage progeny production drops precipitously up to six orders of magnitude (Fig. 2; Fig. 3). This outcome demonstrates the strong influence of lysis timing on the reproductive success of the phage.

Conceptually, phage production per generation depends on the burst size, which is positively correlated to the lysis time (Wang 2006; Baker et al. 2016). The exact shape of this correlation is a matter of debate (Baker et al. 2016). A phage genotype with a short lysis time and hence a low burst size would result in an overall reduction in phage productivity. Thus, fewer phages are carried over to the next transfer. Conversely, one might intuitively assume that for delayed-lysing strains, the phage productivity will be significantly higher. However, the delay in lysing traps a significant number of phages within cells, which fail to make it to the next transfer. Although delaying lysis increases the number of progenies, it also delays the generation time. This trade-off makes the optimum sensitive to environmental conditions such as dilution due to a

constant flow rate. Our experimental set up simulates such a dilution, and the strains with lysis

times that deviate from the optimum show reduced phage productivity over subsequent transfers

(Figs. 2 and 3).   In a previous study using phage λ holin mutants, we demonstrated that holin

accumulation thresholds generate precision in lysis timing. We showed that noise in lysis timing

is reduced in mutants with lysis times closer to that of the wild type (Ghusinga et al. 2017; Kannoly

et al. 2020). Current results support the idea of an optimal lysis time presumably derived from

minimizing the variation in lysis timing. This is in stark contrast to previous studies, which have

suggested that mutations that increase lysis time variance would be favored by selection (Bull et

al. 2011; Baker et al. 2016). It was hypothesized that since phage burst size increase linearly

(Hutchison and Sinsheimer 1966; Wang 2006), early lysis events would contribute more to

population growth than late lysis events detract (Bull et al. 2011; Baker et al. 2016). Interestingly,

Storms *et. al* (2014) have reported that T4 phage productivity in a cell about to undergo cell

division was almost three times greater than the productivity in a young, newly formed cell. We

speculate that if the contribution of early lysis events to population growth is balanced by larger

bursts in the later stages of infection, mutations that increase lysis time variance may not be favored

after all (Baker et al. 2016).

Taken together, minimizing lysis time variance may factor in the evolution of holin. In this

study, we used holin mutants to further explore the effect of lysis time on phage productivity under

quasi-steady state conditions. Again, mutants with lysis times closer to that of the wildtype showed

a consistent increase in phage productivity compared to those with suboptimal lysis times.

Optimality models can reveal the underlying ecological contexts influencing evolutionary

processes. Our results suggest that the wildtype λ might already be selected for conditions where

a constant dilution results in a continuous wash out. Further experiments to test this hypothesis

might involve direct competition of wildtype against mutants under similar conditions. Also, long-

term experimental evolution under the same conditions should reveal if phage strains with

suboptimal lysis times could evolve towards optimality.

Our results add additional support for the idea that life history traits evolve to optimal values.

Much work has been conducted experimentally assessing the validity of optimality models with

varying degrees of success (Kozlowski 1999; Dekel and Alon 2005; Miller et al. 2006; Camps et al.

2007; Poelwijk et al. 2011*b*, 2011*a*; Nakabayashi 2012; Graves and Weinreich 2017). At this point, it is

clear that an unnuanced account of optimality evolution may fail because of genetic details or

constraints (Bull et al. 2004). Nonetheless, optimality models have provided valuable insights

into biological traits ranging from the genetic code to animal behavior (Charnov 1976; Parker

and Smith 1990; Jensen et al. 2006; Roff et al. 2006; Noblin et al. 2008; Baranov et al. 2009;

Henshaw 2018; Ruess et al. 2019).

MATERIALS AND METHODS

Strains

All bacteria and lysogen strains used in this study are listed in Table 1.

Plaque assays

The plaque assays were designed for minimizing variation in plaque size and enabling

plaque counting. The *E. coli* strain MC4100 was grown overnight in TB broth (5 g NaCl and 10 g

tryptone in 1 L water) plus 0.2% maltose at 37°C. The overnight culture was diluted with equal

volume of TB + maltose and grown for another 1.5 h. 100 ul of these cells were mixed with

appropriately diluted phage lysates and incubated at room temperature for 20 min to allow pre-

adsorption. This mixture was then added to 2.5 ml of molten H-top agar (Miller 1992), gently

vortexed, and overlaid onto freshly prepared plates containing 35 ml LB agar. The plates were then

incubated at 37°C and plaques were counted after 18–22 h. Each assay was triplicated to estimate

the average plaque forming units per milliliters (PFU/ml).

Thermal induction of lysogens to obtain phage lysates

Lysogens were grown overnight in LB media at the permissive temperature of 30°C.

Overnight cultures were diluted 100-fold and grown in a 30°C shaking incubator till an OD_{600} of

0.3–0.4 was reached. For heat induction, the cultures were transferred to a 42°C shaking water

bath for 20 min. Following induction, the cultures were transferred to a 37°C shaking incubator

until lysis. The phage lysates were filtered and titered using plaque assay.

Single-cell lysis time determination

We induced the lytic cycle in lysogens and recorded the lysis times of single cells to estimate the

mean lysis time. The protocol for determining single-cell lysis times has been described previously

(Dennehy and Wang 2011). Briefly, a 200-µl aliquot of exponentially growing culture (OD_{600} =

0.3–0.4) of lysogens was chemically fixed to a 22 mm square glass coverslip, which was pretreated

with 0.01% poly-L-lysine (mol. wt. 150 K–300 K; Millipore Sigma, St. Louis, MO) at room

temperature for 10 min. Using this coverslip, a perfusion chamber (RC-21B, Warner Instruments,

New Haven, CT) was assembled and immediately placed on a heated platform (PM2; Warner

Instruments, New Haven, CT). The heated platform was mounted on an inverted microscope stage

(TS100, Nikon, Melville, NY), and the perfusion chamber was infused with heated LB at 30°C

(Inline heater: SH-27B, dual channel heating controller: TC-344B; Warner Instruments, New

Haven, CT). The chamber temperature was elevated to 42°C for 20 min and thereafter maintained

at 37°C until ~95% cells were lysed. Videos of cell lysis were recorded using an eyepiece camera

(10X MiniVID™; LW Scientific, Norcross, GA). The lysis times of individual cells were visually

determined using VLC™ media player. Lysis time was defined as the time required for a cell to

disappear after the temperature was increased to 42°C. For each lysogen, an average of

approximately 100 cells was used to calculate the lysis time.

Serial Transfers

To initiate the transfers, an *E. coli* ($\sim 10^7$ CFU/ml) culture growing exponentially at 37°C

in a shaking incubator (200 rpm) was infected with a phage lysate ($\sim 10^4$ PFU/ml). The infection

could proceed for 3 h, after which phages were separated by filtration using a 0.2 μ syringe filter

(Pall Corp.). The filtered lysate was diluted 1000-fold to start the next transfer using freshly

growing cells ($\sim 10^7$ CFU/ml). These steps were repeated four more times for a total of five serial

transfers (Fig. 1). Plaque assays were performed at the end of each transfer to determine the titers

of phage lysates. The transfers for all strains were duplicated.

Acknowledgements

This work was funded by National Institutes of Health NIGMS grant 1R01GM124446-01 to AS

and JJD. We thank Ry Young for stimulating conversations about phage λ, and for the phage and

bacterial strains described herein. We also appreciate thought-provoking discussions about phage

life history with Stephen Abedon, Jim Bull, Khem Ghusinga, Cesar Vargas-Garcia, Ing-Nang

Wang, and Daniel Weinreich.

References

Abedon, S. T. 1989. Selection for bacteriophage latent period length by bacterial density: A theoretical examination. Microbial ecology 18:79–88.

Abedon, S. T., ed. 2008. Bacteriophage Ecology. Cambridge University Press, Cambridge.

Abedon, S. T., T. D. Herschler, and D. Stopar. 2001. Bacteriophage latent-period evolution as a response to resource availability. Applied and Environmental Microbiology 67:4233–4241.

Abedon, S. T., P. Hyman, and C. Thomas. 2003. Experimental examination of bacteriophage latent-period evolution as a response to bacterial availability. Applied and Environmental Microbiology 69:7499–7506.

Baker, C. W., C. R. Miller, T. Thaweethai, J. Yuan, M. H. Baker, P. Joyce, and D. M. Weinreich. 2016. Genetically determined variation in lysis time variance in the bacteriophage φX174. G3: Genes, Genomes, Genetics 6:939–955.

Baranov, P. v., M. Venin, and G. Provan. 2009. Codon size reduction as the origin of the triplet genetic code. PLoS ONE 4:e5708.

Bonachela, J. A., and S. A. Levin. 2014. Evolutionary comparison between viral lysis rate and latent period. Journal of Theoretical Biology 345:32–42.

Bull, J. J. 2006. Optimality models of phage life history and parallels in disease evolution. Journal of theoretical biology 241:928–38.

Bull, J. J., R. H. Heineman, and C. O. Wilke. 2011. The phenotype-fitness map in experimental evolution of phages. PLoS ONE 6:e27796.

Bull, J. J., D. W. Pfennig, and I. N. Wang. 2004. Genetic details, optimization and phage life histories. Trends in Ecology and Evolution 19:76–82.

Camps, M., A. Herman, E. Loh, and L. A. Loeb. 2007. Genetic constraints on protein evolution. Critical Reviews in Biochemistry and Molecular Biology. NIH Public Access.

Casadaban, M. J. 1976. Transposition and fusion of the lac genes to selected promoters in Escherichia coli using bacteriophage lambda and Mu. Journal of Molecular Biology 104:541–555.

Chantranupong, L., and R. H. Heineman. 2012. A common, non-optimal phenotypic endpoint in experimental adaptations of bacteriophage lysis time. BMC Evolutionary Biology 12:37.

Charnov, E. L. 1976. Optimal foraging: attack strategy of a mantid. The American Naturalist. The University of Chicago PressThe American Society of Naturalists.

Crow, J. F. 1993. Natural selection: Domain, levels, and challenges. Trends in Ecology & Evolution 8:151–152.

Dekel, E., and U. Alon. 2005. Optimality and evolutionary tuning of the expression level of a protein. Nature 436:588–592.

Dennehy, J. J., and I.-N. Wang. 2011. Factors influencing lysis time stochasticity in bacteriophage λ. BMC microbiology 11:174.

Endler, J. A. 1986. Natural selection in the wild. Princeton University Press.

Fishov, I., A. Zaritsky, and N. B. Grover. 1995. On microbial states of growth. Molecular Microbiology
15:789–794.

Ghusinga, K. R., J. J. Dennehy, and A. Singh. 2017. First-passage time approach to controlling noise in the
timing of intracellular events. Proceedings of the National Academy of Sciences 114:693–698.

Graves, C. J., and D. M. Weinreich. 2017. Variability in Fitness Effects Can Preclude Selection of the
Fittest. Annual Review of Ecology, Evolution, and Systematics 48:399–417.

Heineman, R. H., J. J. Bull, and T. Hansen. 2007. Testing optimality with experimental evolution: lysis
time in a bacteriophage. Evolution; international journal of organic evolution 61:1695–709.

Henshaw, J. M. 2018. Finding the one: optimal choosiness under sequential mate choice. Journal of
Evolutionary Biology 31:1193–1203.

Hutchison, C. A., and R. L. Sinsheimer. 1966. The process of infection with bacteriophage ΦX174: X.
Mutations in a ΦX lysis gene. Journal of Molecular Biology 18:429–447.

Jensen, K. H., T. Little, A. Skorping, and D. Ebert. 2006. Empirical support for optimal virulence in a
castrating parasite. PLoS Biology 4:1265–1269.

Kannoly, S., T. Gao, S. Dey, I.-N. Wang, A. Singh, and J. J. Dennehy. 2020. Optimum threshold minimizes
noise in timing of intracellular events. bioRxiv 2020.02.14.949891.

Kozlowski, J. 1999. Adaptation: A Life History Perspective. Oikos 86:185.

Levin, B. R., F. M. Stewart, and L. Chao. 1977. Resource-Limited Growth, Competition, and Predation: A
Model and Experimental Studies with Bacteria and Bacteriophage. The American Naturalist. The
University of Chicago PressThe American Society of Naturalists.

Lewontin, R. C. 1989. A natural selection. Nature 339:107–107.

Miller, J. H. 1992. A short course in bacterial genetics: a laboratory manual and handbook for *Escherichia*
coli and related bacteria. Volume 1. CSHL Press.

Miller, S. P., M. Lunzer, and A. M. Dean. 2006. Direct demonstration of an adaptive constraint. Science
(New York, N.Y.) 314:458–61.

Nakabayashi, J. 2012. Optimal ratio of scaffold complex to free Fus3 to maximise the accumulation of
phosphorylated Fus3 in yeast pheromone signalling pathway. IET Systems Biology 6:9–21.

Noblin, X., L. Mahadevan, I. A. Coomaraswamy, D. A. Weitz, N. M. Holbrook, and M. A. Zwieniecki. 2008.
Optimal vein density in artificial and real leaves. Proceedings of the National Academy of Sciences of the
United States of America 105:9140–9144.

Parker, G. A., and J. M. Smith. 1990. Optimality theory in evolutionary biology. Nature. Nature Publishing
Group.

Poelwijk, F. J., M. G. J. de Vos, and S. J. Tans. 2011*a*. Tradeoffs and optimality in the evolution of gene
regulation. Cell 146:462–470.

Poelwijk, F. J., P. D. Heyning, M. G. J. de Vos, D. J. Kiviet, and S. J. Tans. 2011*b*. Optimality and evolution
of transcriptionally regulated gene expression. BMC Systems Biology 5:128.

Roff, D. A., E. Heibo, and L. A. VØllestad. 2006. The importance of growth and mortality costs in the
evolution of the optimal life history. Journal of Evolutionary Biology 19:1920–1930.

Ruess, J., M. Pleška, C. C. Guet, and G. Tkačik. 2019. Molecular noise of innate immunity shapes bacteria-
phage ecologies. (O. A. Igoshin, ed.)PLOS Computational Biology 15:e1007168.

Sezonov, G., D. Joseleau-Petit, and R. D'Ari. 2007. Escherichia coli physiology in Luria-Bertani broth.
Journal of Bacteriology 189:8746–8749.

Singh, A., and J. J. Dennehy. 2014. Stochastic holin expression can account for lysis time variation in the
bacteriophage λ. Journal of the Royal Society, Interface 11:20140140.

Smith, J. M. 1978. Optimization Theory in Evolution. Annual Review of Ecology and Systematics 9:31–56.

Wang, I. N., D. E. Dykhuizen, and L. B. Slobodkin. 1996. The evolution of phage lysis timing. Evolutionary
Ecology 10:545–558.

Wang, I.-N. 2006. Lysis timing and bacteriophage fitness. Genetics 172:17–26.

Wang, I.-N., J. Deaton, and R. Young. 2003. Sizing the holin lesion with an endolysin-beta-galactosidase
fusion. Journal of bacteriology 185:779–87.

Wang, I.-N., D. L. Smith, and R. Young. 2000. Holins: the protein clocks of bacteriophage infections.
Annual Review of Microbiology 54:799–825.

Table 1. All bacteria and lysogen strains used in this study are listed along with the genotype of

the prophage's holin gene and the source of the strain. [a]Coli Genetic Stock Center.

Strain	Genotype	Lysis Time (min)	Source
CGSC#: 6152[a]	*E. coli* MC4100 (λ-)	NA	Casadaban 1976
JJD3	MC4100 (λ cI857 S)	63	Wang 2006
JJD246	MC4100 (λ cI857 S105$_{H7D}$)	69	Kannoly et al., 2020
JJD248	MC4100 (λ cI857 S105$_{F94C}$)	56	Kannoly et al., 2020
JJD391	MC4100 (λ cI857 S105$_{A16G/K92Q}$)	140	Kannoly et al., 2020
JJD404	MC4100 (λ cI857 S105$_{I21V}$)	30	Kannoly et al., 2020
JJD405	MC4100 (λ cI857 S105$_{V45G}$)	32	Kannoly et al., 2020
JJD414	MC4100 (λ cI857 S105$_{L90I}$)	45	Kannoly et al., 2020
JJD423	MC4100 (λ cI857 S105$_{F27Y}$)	122	Kannoly et al., 2020
JJD432	MC4100 (λ cI857 S105$_{S89W}$)	107	Kannoly et al., 2020
JJD436	MC4100 (λ cI857 S105$_{K92N}$)	100	Kannoly et al., 2020

FIG. 1. Serial transfers. For serial transfers, λ phage strains were cocultured with exponentially growing host cells, filtered to separate phages, and diluted 1000-fold to start the next transfer with freshly growing cultures. After every transfer, the filtered phage lysates were titered using plaque assays.

346
347 **FIG. 2. Titers of λ strains with different lysis times (LT) after each transfer.** Lysis times are
348 from single-cell estimates as reported in Kannoly et al., 2020. WT lysis time = 63 min. Error
349 bars, mean ± SEM.

354
Fɪɢ. 3. Titers of λ strains with different lysis times after the fifth transfer (Log scale). Titers
increase as the lysis time approaches optima. Inset shows the same data on a linear scale. Error
bars, mean ± SEM.