The IQGAP1-hnRNPM interaction links tumour-promoting alternative splicing

to heat-induced signals

Andrada Birladeanu[1,‡], Malgorzata Rogalska[2,‡], Myrto Potiri[1,‡], Vassiliki

Papadaki[1], Margarita Andreadou[1], Dimitris Kontoyiannis[1,3], Zoi Erpapazoglou[1],

Joe D. Lewis[4], Panagiota Kafasla[*,1]

[1]Institute for Fundamental Biomedical Research, B.S.R.C. "Alexander Fleming", 34

Fleming st., 16672 Vari, Athens, Greece

[2]Centre de Regulació Genòmica, The Barcelona Institute of Science and Technology

and Universitat Pompeu Fabra, Dr. Aiguader 88, 08003, Barcelona, Spain

[3]Department of Biology, Aristotle University of Thessaloniki, Greece

[4]European Molecular Biology Laboratory, 69117 Heidelberg, Germany

[‡]These authors contributed equally to this work

[*]Correspondence: kafasla@fleming.gr

Abstract (150 words):

Alternative Splicing (AS) is extensively regulated during the cell cycle and is also

involved in the progression of distinct cell cycle phases. Stressing agents, such as heat

shock, halts AS affecting mainly post-transcriptionally spliced genes. Stress-dependent

regulation of AS relies possibly on the subnuclear location of its determinants, such us

Serine-Arginine rich (SR) and heterogeneous nuclear ribonucleoproteins, hnRNPs.

Spliceosomal components like SRSF1, SRSF9, SRp38, hnRNPM have been connected

to such responses of AS, but how they are directed to respond to stress remains largely

unknown. Here, we report that upon heat stress, the cytosolic scaffold protein IQGAP1

translocates into the nucleus of gastric cancer cells, where it acts as a tethering module

for hnRNPM and other spliceosome components, mediating their subnuclear

positioning and their response to stress. Moreover, together, they regulate the AS of the

anaphase promoting complex (ANAPC) subunit 10 to promote gastric cancer cell

growth.

Introduction

In humans, more than 95% of multi-exonic genes are potentially alternatively spliced[1,2].

Precise modulation of Alternative Splicing (AS) is essential for shaping the proteome

of any given cell and altered physiological conditions can change cellular function via

AS reprogramming[3]. The importance of accurate AS in health and disease, including

cancer[4–7], has been well documented. However, evidence connecting AS regulation to

signalling comes mostly from few cases where localization, expression, or post-

translational modifications of specific splicing factors such as Serine-Arginine-rich

(SR) proteins or heterogeneous nuclear ribonucleoproteins (hnRNPs)[3,8] are altered.

An abundant, mainly nuclear protein of the hnRNP family is hnRNPM, with four

isoforms that arise from AS and/or post-translational modifications[9]. The only nuclear

role so far reported for hnRNPM is in spliceosome assembly and splicing itself[10,11],

including regulation of AS of other as well as its own pre-mRNAs[12–14]. The association

of hnRNPM with the spliceosome is abolished under heat-induced stress[11,15], known to

affect largely post-transcriptional splicing events[16], suggesting hnRNPM's response in

stress-related signalling pathways.

HnRNPM-regulated AS events have been linked to disease development and progression. Specifically, hnRNPM-mediated AS is deterministic for the metastatic potential of breast cancer cells, due to cell-type specific interactions with other splicing factors[17–19]. Furthermore, in Ewing sarcoma cells, hnRNPM abundance and subnuclear localization change in response to a chemotherapeutic inhibitor of the PI3K/mTOR pathway, resulting in measurable AS changes[20]. Very recently, hnRNPM was shown to respond to pattern recognition receptor signalling, by modulating the AS outcome of an RNA-regulon of innate immune transcripts in macrophages[21].

Despite such growing evidence on the cross-talk between hnRNPM-dependent AS and cellular signalling, how distinct signals are transduced to hnRNPM and the splicing machinery remains unclear. Here, we present conclusive evidence of how this could be achieved. We describe a novel interaction between hnRNPM and the scaffold protein IQGAP1 (IQ Motif Containing GTPase Activating Protein 1) in the nucleus of gastric cancer cells. Cytoplasmic IQGAP1 acts as a signal integrator in a number of signalling pathways[22], but there is no defined role for the nuclear pool of IQGAP1. With *IQGAP1* mRNA being overexpressed in many malignant cell types, the protein seems to regulate cancer growth and metastatic potential[23–25]. Moreover, aged mice lacking IQGAP1 develop gastric hyperplasia suggesting an important *in vivo* role for IQGAP1 in maintaining the gastric epithelium[26]. In the present study we show that this novel, nuclear interaction between hnRNPM and IQGAP1 links heat-induced signals to hnRNPM-regulated AS in gastric cancer, a cancer type that has been associated with a significantly high incidence of AS changes[6,7]. Additionally, we show that depletion of both interacting proteins results in alternative splicing changes that disfavour tumour growth, which makes them and their interaction interesting cancer drug targets.

Results

hnRNPM and IQGAP1 expression levels are significantly altered in gastric cancer.

Analysis of the hnRNPM and IQGAP1 protein levels by immunofluorescence on commercial tissue microarrays revealed increased hnRNPM and IQGAP1 staining in tumour as compared to normal tissue, especially in adenocarcinoma samples (**Fig. 1a and Supplementary Fig. 1a-c).** This agrees with TCGA data analysis indicating significantly increased expression of both *IQGAP1* and *hnRNPM* mRNAs in stomach adenocarcinoma (STAD) vs normal tissue samples (**Fig. 1b**). Furthermore, a strong correlation between the expression of the two mRNAs (*HNRNPM* and *IQGAP1*) in normal tissues (GTex) was revealed, which was reduced in normal adjacent tissues (Normal) and eliminated in STAD tumours (**Fig. 1c**).

Detection of hnRNPM and IQGAP1 protein levels in a number of gastric cancer cell lines by immunoblotting (**Fig. 1d**) identified cell lines with similar levels of hnRNPM and low (MKN45, AGS) or high (NUGC4, KATOIII) levels of IQGAP1, corroborating the TCGA data. Two of those STAD cell lines were used for further studies on the interaction between hnRNPM and IQGAP1 (NUGC4 and MKN45) since they have similar hnRNPM, but different IQGAP1 levels.

IQGAP1 and hnRNPM regulate gastric cancer cell growth *in vitro* and *in vivo*

To test the significance of IQGAP1 and hnRNPM in tumour development, we used a CRISPR-Cas9 approach to generate *hnRNPM*-KnockOut (KO) cell lines, *IQGAP1*^{KO} and double KO. We knocked-out successfully *IQGAP1* in both MKN45 and NUGC4 cells without significant change in hnRNPM protein levels (**Supplementary Fig. 2a**). However, numerous attempts to disrupt the ORF of hnRNPM resulted in ~75%

reduction, as we could not isolate single hnRNPM^{KO} clones. Thus, for the subsequent

experiments we either worked with mixed cell populations (e.g. MKN45-*hnRNPM*^{KO}-

IQGAP1^{KO}) with 75% reduced hnRNPM expression levels (**Supplementary Fig. 2a**),

or, where stated, we used siRNAs for down-regulation of the hnRNPM levels.

To evaluate the role of the two proteins in gastric cancer progression, we first assayed

the STAD cell lines and the derivative KO/Knock-Down cell lines for their metabolic

activity (**Fig. 2a**). Downregulation of hnRNPM alone or in combination with IQGAP1

impairs cellular metabolic activity, compared to the parental lines, as indicated by MTT

assays performed over a period of 5 days. Interestingly, we detected an increased

metabolic rate in IQGAP1^{KO} cells, which can be attributed to hnRNPM-independent

changes in metabolic activity in the absence of IQGAP1 (data not shown). In a 2D

colony formation assay, cells with reduced levels of both IQGAP1 and hnRNPM

proteins generated a significantly reduced number of colonies compared to parental

cells (**Fig. 2b**). Furthermore, cell cycle analyses using propidium iodide combined with

flow cytometry showed that unsynchronized IQGAP1^{KO} cells depicted a small but

significant increase of cell population at the S and G2/M phases with subsequent

reduction of G1 cells (**Fig. 2c**). hnRNPM^{KO} cells showed a similar phenotype, whereas

depletion of both interacting proteins (hnRNPM^{KO}-IQGAP1^{KO}) enhanced the effect

(**Fig. 2c**). These differences were stronger after cell synchronization (**Supplementary**

Fig. 2b). Wound healing assays did not reveal significant differences in the migratory

ability of these cell lines, only an increase in wound healing rate for hnRNPM^{KO} cells

compared to the parental line. Importantly, this expedited wound healing in hnRNPM^{KO}

cells   was   completely   abolished   upon   concomitant   absence   of   IQGAP1

(**Supplementary Fig. 2c**).

To examine the *in vivo* effect of abrogating IQGAP1 and hnRNPM on tumour

development and progression, we injected MKN45, MKN45-*IQGAP1*^KO, MKN45-

hnRNPM^KO and MKN45-*hnRNPMKO-IQGAP1*^KO cells subcutaneously into the flanks

of NOD/SCID mice. Measurements of tumour dimensions throughout the experiment

demonstrated that cells lacking both IQGAP1 and hnRNPM result in significantly

reduced tumour growth compared to the parental and the single KO cells (**Fig. 2d**).

Immunohistochemical analysis of the tumours confirmed greatly reduced levels of

hnRNPM and/or IQGAP1 in the mutant cell line-derived xenografts. Furthermore, Ki-

67 staining was significantly reduced in the single and double KO tumours compared

to the parental cell line-derived ones, showing the involvement of the two proteins in

the *in vivo* proliferation of gastric cancer cells (**Fig. 2e**). Collectively, these results

demonstrate that co-operation of IQGAP1 and hnRNPM is required for gastric cancer

cell growth and progression both *in vitro* and *in vivo*.

IQGAP1 and hnRNPM interact in the nucleus of gastric cancer cell lines

An interaction between hnRNPM and IQGAP1 was detected in nuclear extracts from

both STAD cell lines using immunoprecipitation with anti-IQGAP1 antibodies (**Fig.**

3a). Treatment with RNaseA showed that the two proteins interact independently of the

presence of RNA. This interaction also appeared to be DNA-independent

(**Supplementary Fig. 3a**). Immunoprecipitation using anti-IQGAP1 antibodies from

cytoplasmic extracts did not reveal interaction with the minor amounts of cytoplasmic

hnRNPM (**Supplementary Fig. 3b and data not shown**), indicating that if the proteins

do interact in the cytoplasm, these complexes are less abundant compared to the nuclear

ones.

In agreement with previous reports of a small fraction of IQGAP1 entering the nucleus

in a cell cycle-dependent manner[27], we confirmed the presence of IQGAP1 in the

nucleus of both NUGC4 and MKN45 cells using confocal imaging (**Fig. 3b**) The

interaction of nuclear IQGAP1 with hnRNPM *in situ* was confirmed using proximity

ligation assays (PLA) (**Fig. 3c).** Quantification of the PLA signal per cell demonstrated

that the interaction between the two proteins was clearly detectable in the nucleus of

both cell lines (**Fig. 3d**). Some cytoplasmic interaction sites were also detected, but they

were minor compared to the nuclear ones (**Fig. 3c, d**) in agreement with the

immunoprecipitation results from cytoplasmic extracts (**Supplementary Fig. 3b**).

Together these results demonstrate the RNA-independent, nuclear interaction between

hnRNPM and IQGAP1 in gastric cancer cells.

Nuclear IQGAP1-containing RNPs are involved in splicing regulation

To identify other possible nuclear IQGAP1 interacting partners, we performed

immunoprecipitation with anti-IQGAP1 Abs from nuclear extracts of NUGC4 cells and

analysed the resulting proteins by LC-MS/MS (**Supplementary Table 1**). GO-term

enrichment analysis of the co-precipitated proteins showed a significant enrichment in

biological processes relevant to splicing regulation (**Supplementary Fig. 4a**).

Construction of an IQGAP1 interaction network revealed that IQGAP1 can interact not

only with the majority of the hnRNPs, but also with a large number of spliceosome

components (mainly of U2, U5snRNPs) and RNA-modifying enzymes (**Fig. 4a**). The

interactions between IQGAP1 and selected hnRNPs (A1, A2B1, C1C2, L, M) and with

selected spliceosome components and RNA processing factors (SRSF1, CPSF6,

DDX17, DHX9, ILF3/NF90)[28] were independently validated in both NUGC4 and

MKN45 cells (**Fig. 4b, Supplementary Fig. 4b**). The interactions of IQGAP1 with

hnRNPs A1, A2B1 are RNA-dependent, whereas only a subset of hnRNPs (L, C1/C2) interact with IQGAP1 in the absence of RNA (**Fig. 4b**). These observations pinpointed to the RNA-independent interaction with hnRNPM as a distinct one, worthy of further investigation.

In the nuclear extracts used in our immunoprecipitations hnRNPM is enriched together with the majority of hnRNPs[29,30] (e.g. A2B1, K), other nuclear speckle components like SRSF1, and nuclear matrix associated proteins like SAFB and MATRIN3 (**Fig. 4c and Supplementary Fig. 4c**), but not histones such as H3, which are present mainly in the insoluble nuclear material (**Supplementary Fig. 4c**). To further explore the interactions of nuclear IQGAP1 with the splicing machinery, we queried whether itself and its interacting partners are present in described splicing-related subnuclear fractions[31]. This was justified by the recent identification of hnRNPM as a significant component of the Large Assembly of Spliceosome Regulators (LASR). This complex is assembled via protein-protein interactions, lacks DNA/RNA components, and appears to function in co-transcriptional AS[31]. In MKN45 cells, IQGAP1 and hnRNPM co-exist mainly in the soluble nuclear fraction together with hnRNPs K, C1/C2 and other spliceosome components[31]. Significantly smaller IQGAP1 and hnRNPM amounts were detected in the proteins released from the HMW material upon DNase treatment (D), together with hnRNPC1/C2 and other spliceosome components, including SF3B3 (**Supplementary Fig. 4d).** Thus, the interacting pools of IQGAP1 and hnRNPM are not major LASR components and possibly participate in post-transcriptional splicing events.

To assess the possible functional involvement of IQGAP1 in splicing, we used the hnRNPM responsive DUP51M1, DUP50M1 and control DUP51-ΔM and DUP50-ΔM mini-gene reporters[31]. DUP51M1 is a three exon minigene, where exon 2 contains a unique UGGUGGUG hnRNPM consensus binding motif that results in skipping of this

exon upon hnRNPM binding. The DUP51-ΔM minigene carries a mutation in the

hnRNPM binding site, resulting in increased inclusion of exon 2[31]. The DUP50M1

reporter was derived from DUP51M1 by slightly altering the hnRNPM binding site on

exon 2 (to UGUUGUGUUG). The DUP50-ΔM reporter has also a mutation in the

hnRNPM binding site, that does not allow hnRNPM binding[31]. Transfection of the two

gastric cancer cell lines with the pDUP51M1 plasmid and subsequent RT-PCR analysis

with primers that allow detection of the two possible mRNA products, revealed

different splicing patterns of the reporter: significantly more inclusion of exon 2 was

detected in NUGC4, which express IQGAP1 at higher levels, than MKN45

(**Supplementary Fig. 4e**). Moreover, analysis of the splicing pattern of the DUP51-ΔM

reporter showed that the dependence of exon 2 inclusion on hnRNPM was more

pronounced in MKN45 compared to NUGC4 (**Supplementary Fig. 4e**). The same

results were obtained when we used the DUP50M1 and DUP50ΔM sets of reporters.

These results prompted us to focus on MKN45 cells to further probe the involvement

of the hnRNPM-IQGAP1 interaction in AS. In the IQGAP1^{KO} cells, the DUP51M1

reporter presented a significant ~2-fold increase in exon 2 inclusion compared to the

parental cells (**Fig. 4d**). As expected from the interaction of IQGAP1 with other splicing

factors, splicing of the DUP51-ΔM reporter was also affected upon IQGAP1 loss.

However, this change was smaller (~1.3-fold increase) compared to the impact on

hnRNPM-dependent splicing (**Fig. 4d**). Attempts to restore splicing efficiency by

expressing GFP-IQGAP1 were inconclusive, as the recombinant protein localized very

efficiently in the nucleus[27], thus inhibiting rather than rescuing exon 2 skipping

(**Supplementary Fig. 4f**), likely due to sequestration of hnRNPM from the splicing

machinery. These results denote that IQGAP1 interacts with a large number of splicing

factors, of which hnRNPM stands out as a unique, RNA-independent interaction, important for hnRNPM's activity in alternative splicing.

A broad role of IQGAP1 in hnRNPM-dependent alternative splicing

To determine whether IQGAP1^{KO} results in broad splicing deregulation, we profiled AS changes between MKN45 and MKN45-*IQGAP1^{KO}* cells by RNA-seq (**Fig. 5a-c**). A number of significantly altered AS events were detected (**Fig. 5a** and **Supplementary Table 2a**) more than 50% of which were alternative exons (**Fig. 5b**). Statistical analysis showed that down-regulated exons are more likely to have the hnRNPM binding motif than up-regulated exons (chi-square $(1, 86) = 36.4848$, $P < 0.00001$; **Fig. 5c, d**). The presence of hnRNPM consensus binding motifs[13] was significantly enriched downstream of 81% of the down-regulated alternative exons. GO term enrichment analysis of the affected genes yielded significant enrichment of the biological processes of cell cycle (GO:0007049, P: 3.75E-04) and cell division (GO:0051301, P: 3.33E-04) (**Supplementary Table 2b**), connecting these results to the detected cell cycle defects (**Fig. 2c**).

Splicing events selected for validation were required to adhere to most of the following criteria: 1) high difference in Psi (ΔPsi) between the two cell lines (where [Psi] is the Percent Spliced In, i.e. the ratio between reads including or excluding alternative exons), 2) involvement of the respective proteins in the cell cycle, 3) previous characterization of pre-mRNAs as hnRNPM targets[32,33] and/or 4) presence of the hnRNPM-consensus motif up- or downstream of the alternative exon. 12 out of 19 AS events (63%) selected based on the above criteria were validated by RT-PCR **(Fig. 5e, Supplementary Fig. 5a-c, Supplementary Table 3).** Therefore, the above indicate that IQGAP1 and hnRNPM co-regulate the AS of alternative exons in a cell-cycle-

245 related RNA regulon. Within this regulon, ANAPC10 pre-mRNA was singled out for

246 further analysis as it had the highest change in |ΔPsi|/Psi combination (**Fig. 5a, e**) and

247 it is an hnRNPM-eCLIP target[33] with the major hnRNPM binding sites downstream of

248 the regulated exon, where the predicted hnRNPM consensus binding motif is also

249 located (**Supplementary Fig. 5d**).

IQGAP1 and hnRNPM co-operatively regulate the function of APC/C through AS

of the *ANAPC10* pre-mRNA

Importantly for cancer cells, ANAPC10 plays a crucial role in cell cycle and cell

division[34–37] as a substrate recognition component of the anaphase promoting

complex/cyclosome (APC/C), a cell cycle-regulated E3-ubiquitin ligase that controls

progression through mitosis and the G1 phase of the cell cycle. ANAPC10 interacts

with the co-factors CDC20 and/or CDH1 to recognize targets to be ubiquitinated and

subsequently degraded by the proteasome. In *IQGAP1^{KO}* cells, increased levels of

ANAPC10 exon 4 skipping were detected and simultaneous knock-down of hnRNPM

using siRNAs led to further increase in *ANAPC10* exon 4 skipping compared to scr-

siRNA transfected cells (**Fig. 5e** and **6a**). Skipping of exon4 is predicted to result in the

preferential production of an isoform lacking amino acid residues important for

interaction with the D-box of the APC/C targets[36,37]. To verify that this is the case, using

LC-MS/MS analyses of the proteomes of the parental and the *IQGAP1*^{KO} cell lines, we

compared the levels of known targets of the APC/C complex. We detected increased

abundance of anaphase-specific targets of the APC/C-CDH1 [35], namely RRM2, TPX2,

ANLN, and TK1, but not of other APC/C known targets (**Supplementary Fig. 6a).**

Immunoblotting showed that TPX2, RRM2 and TK1 levels were indeed increased in

IQGAP1^{KO} cells and even more after concomitant siRNA mediated *hnRNPM* knock-

down (**Fig. 6b**). The same was true for CDH1/FZR, an APC/C co-factor, which is also

a target of the complex, as is ANLN (**Supplementary Fig. 6b**). Interestingly,

ANAPC10 levels were reduced in the *IQGAP1^{KO}* cells after knock-down of *hnRNPM*.

Given the role of APC/C and its targets, TK1, RRM2 and TPX2 in the progression of

mitosis and cell division[35,38–40], we determined the impact of the downregulation of both

IQGAP1 and *hnRNPM* on cell division. Using DAPI staining and anti-β-tubulin

cytoskeleton immunostaining we detected a significant number of double *IQGAP1-*

hnRNPM^{KO} cells being multinucleated (2 or more nuclei; **Fig. 6d-e**). A similar

phenotype was detected when we used siRNAs to downregulate hnRNPM levels

(**Supplementary Fig. 6c**). These results denote that IQGAP1 interacts with hnRNPM

in the nucleus of gastric cancer cells and co-operatively they generate at least an

alternatively spliced isoform of ANAPC10. This, in turn, tags cell cycle-promoting

proteins for degradation, thus contributing to the accelerated proliferation phenotype of

tumour cells.

IQGAP1 alters hnRNPM's ability to respond to heat-shock in cancer cells

To detect the mechanism behind the IQGAP1-hnRNPM co-regulation of AS events, we

investigated whether binding of hnRNPM on its pre-mRNA target is altered in the

absence of IQGAP1. For this, we used the previously mentioned minigene reporter and

tested the association of hnRNPM with the DUP51M1 transcript using crosslinking,

immunoprecipitation with anti-hnRNPM antibodies, and RT-PCR of the associated pre-

mRNA. No significant differences were detected between the parental and the

IQGAP1^{KO} cells in the amount of RNA that was crosslinked to hnRNPM

(**Supplementary Fig. 7a**), indicating that IQGAP1 does not regulate the binding of

hnRNPM to its RNA targets.

The activity of hnRNPM in splicing is altered upon heat-shock, when hnRNPM is

shifted from the nucleoplasm towards the insoluble nuclear matrix[15]. To detect an

involvement of IQGAP1 in a similar regulation of hnRNPM's splicing activity through

changes of its subnuclear distribution, we compared the localization of hnRNPM

between parental and *IQGAP1*[KO] cells, both untreated and after heat-shock (**Fig. 7a-b,**

8c and Supplementary Fig. 7b) using immunofluorescence and confocal microscopy.

A clear difference in the subnuclear distribution of hnRNPM was detected between

paternal and IQGAP1[KO] cells, with the perinuclear enriched localization in parental

cells changing to a more diffuse distribution, not only at the periphery of the nuclei, but

also towards the inside of the nuclei (**Fig. 7a** and **Supplementary Fig. 7b**). As

expected[15], hnRNPM's localization also changed from its perinuclear pattern to a

granular one in the nucleus of parental cells upon heat-shock. Surprisingly, hnRNPM's

localization and staining pattern did not further change upon heat-shock in cells lacking

IQGAP1 (**Fig. 7a** and **Supplementary Fig. 7b**). The effect of heat shock on the

alternative splicing of the DUP50 minigene reporter was also obvious in the parental

cells, however such an effect was not apparent in cells lacking IQGAP1

(**Supplementary Fig. 7c**). To assay the localization of hnRNPM in relation to

spliceosomal components, we compared it to that of SR proteins in untreated and heat-

shocked parental and IQGAP1[KO] cells (**Fig. 7b, c**). Upon heat-shock, the colocalization

between hnRNPM and SR proteins was reduced in the parental cells. hnRNPM and SR

showed also decreased colocalization in untreated cells lacking IQGAP1, and no further

change was induced upon heat-shock. Furthermore, the localization of SR proteins

changes upon heat shock, and these changes seem to depend as well on the presence of

IQGAP1, suggesting a more general role of IQGAP1 for heat-shock response of the spliceosome machinery (**Fig. 7a-c**).

Since in heat-shocked cells hnRNPM moves away from spliceosomal components towards the nuclear matrix[15], we compared nuclear matrix preparations from parental and *IQGAP1*[KO] cells before and after heat-shock. hnRNPM was detected elevated in the nuclear matrix of the parental cells after heat-shock compared to untreated cells, whereas this change was not detected in the *IQGAP1*[KO] cells (**Fig. 8a**). Interestingly, IQGAP1 levels were also increased in nuclear matrix fractions prepared from heat-shocked cells (**Fig. 8a**). In agreement to this, increased nuclear IQGAP1 staining was detected in heat-shocked cells, compared to the untreated controls (**Supplementary Fig. 8a**). Using confocal microscopy and immunofluorescence staining, we compared the localization of hnRNPM to its interacting partner SFPQ (PSF) which is a known component of the nuclear matrix, interacting with the splicing machinery in soluble nucleoplasm[41]. The colocalization of hnRNPM and PSF was partial in untreated parental cells, and was significantly increased upon heat shock. Though, in untreated cells lacking IQGAP1, there was a higher percentage of colocalization between hnRNPM and SFPQ compared to parental cells, this was not affected upon heat shock (**Fig. 8b, c**).

Therefore, IQGAP1 is important for the proper subnuclear distribution of hnRNPM close to spliceosomal components (**Fig. 9)**. Upon heat-shock, hnRNPM translocates closer to nuclear matrix components and away from the spliceosomal components, only if IQGAP1 is present. Collectively, these results indicate the involvement of IQGAP1 in the response of alternative splicing regulating, nuclear ribonucleoproteins (RNPs) to stress, with a more significant role in regulating the splicing activity of hnRNPM-containing RNPs.

Discussion

Splicing regulatory networks are subject to signals modulating alternative exon choice. One of the best characterized AS changes in response to stress-signals is the shutdown of post-transcriptional pre-mRNA splicing observed in heat-shocked cells[42,16]. However, how this occurs mechanistically is still unknown. Here, we report the first insights for the roles of hnRNPM and IQGAP1 in mediating the response of alternative splicing factors to heat-induced stress. The fact that IQGAP1 is a scaffold protein with well-known roles in the cytoplasm as an integrator of many signalling cascades suggests that this may be a more general phenomenon. We show that in gastric cancer cells, the interaction of the two proteins, happens in fractions that are involved in post-transcriptional splicing and is necessary for the response of hnRNPM to heat-induced stress signals. Furthermore, we show here that nuclear IQGAP1 interacts with a large number of splicing regulators mostly in an RNA-dependent manner, in addition to its RNA-independent interaction with hnRNPM. Thus, based on our data, we propose that upon heat, and possibly other stressors, IQGAP1 and hnRNPM are removed from the spliceosome and move towards the less-well-defined nuclear matrix. IQGAP1 is not required for binding of hnRNPM to a reporter pre-mRNA *in vivo*, however, it is necessary for the function of hnRNPM, and possibly other factors in splicing, regulating the interaction of hnRNPM with spliceosomal components at distinct subnuclear compartments.

The nuclear translocation and localization of IQGAP1 appears to be cell-cycle dependent, since it is significantly increased in response to replication stress and subsequent G1/S arrest[27] and as presented herein, upon heat-shock. This finding complements prior reports suggesting IQGAP1 localization at the nuclear envelope during late mitotic stages[43]. Cyclebase data[44] suggest that hnRNPM is required for

progression of the cell cycle G1 phase. On the basis of our data, we propose that in the absence of IQGAP1, a number of pre-mRNAs involved in cell cycle regulation undergoes altered alternative splicing.

Specifically, IQGAP1 and hnRNPM seem to co-operatively regulate the AS of a cell-cycle RNA regulon. 5 out of the 10 cell division-related AS events that are deregulated in the absence of IQGAP1 are exon skipping events, characterized by the presence of the hnRNPM binding motif downstream of the alternative exon. Of these 5 events, ANAPC10 has the highest change in AS pattern upon IQGAP1 abrogation and a significant role in cell cycle regulation and cell division[34,35]. In cells with reduced amounts of both IQGAP1 and hnRNPM, ANAPC10 levels are reduced and at least a group of APC/C-CDH1 targets are specifically stabilized (TPX2, RRM2, TK1, CDH1 itself). Given the role of the controlled degradation of these proteins for cell cycle progression[35,45], we posit that these observations explain the aberrant cell cycle effect in the double KO cell lines, the multinucleated cells phenotype and the importance of the two proteins for gastric cancer development and progression as detected by the xenograft experiments.

Currently, the literature on alternative splicing, cell cycle control, multinucleated cancer cells and tumour growth is rather unclear and often conflicting. There is evidence connecting these cell cycle effects manifested as a balance between up-regulation or down-regulation of tumour growth[45–48]. Mitotic errors, such as mitotic exit defects can have distinct impacts at different points during tumour development[45]. Thus, targeting of the APC/C[Cdh1] activity has been suggested as a possible tumour suppressive approach[34,45]. On the other hand, alternative splicing is subject to extensive periodic regulation during the cell cycle in a manner that is highly integrated with distinct layers of cell cycle control. Our data put these into context, leading to an experimentally

testable model in which IQGAP1 and hnRNPM co-operatively drive the balance

towards tumour growth, and when they cannot interact, the balance is shifted towards

tumour suppression. They accomplish this by adding a new level of control in the

activity of the APC/C complex: alternative splicing regulation. If so, it is of critical

importance to identify additional protein partners involved in this balancing

mechanism, test its connection to specific cell-cycle stages and investigate whether this

mechanism is involved both in additional cancer types and also in normal cells.

Methods

Cell culture. The human gastric cancer cell lines AGS, KATOIII, MKN45 and NUGC4

were a kind gift from P. Hatzis (B.S.R.C. "Al. Fleming", Greece). Cells were grown

under standard tissue culture conditions (37^0C, 5% CO2) in RPMI medium (GIBCO),

supplemented with 10% FBS, 1% sodium pyruvate and 1% penicillin–streptomycin.

NUGC4 originated from a proximal metastasis in paragastric lymph nodes, and

MKN45 was derived from liver metastasis. According to the GEMiCCL database,

which incorporates data on cell lines from the Cancer Cell Line Encyclopedia, the

Catalogue of Somatic Mutations in Cancer and NCI60[49], none of the gastric cancer cell

lines tested have altered copy number of hnRNPM or IQGAP1. Only NUGC4 has a

silent mutation c.2103G to A in *HNRNPM*, which is not included in the Single

Nucleotide Variations (SNVs) or mutations referred by cBioportal in any cancer

type[50,51].

Transfection of MKN45 and NUGC4 cells. Gastric cancer cell lines were transfected

with plasmids pDUP51M1 or pDUP51-ΔM[31] (a gift from D. L. Black, UCLA, USA)

and pCMS-EGFP (Takara Bio USA, Inc) or pEGFP-IQGAP1[52] [a gift from David

Sacks   (Addgene   plasmid   #   30112;   http://n2t.net/addgene:30112;

RRID:Addgene_30112)], using the TurboFect transfection reagent (Thermo Fisher

Scientific, Inc., MA). For RNA-mediated interference, cells were transfected with

control siRNA (sc-37007, Santa Cruz Biotechnology, Inc., CA) or hnRNPM siRNA

(sc-38286) at 30 nM final concentration, using the Lipofectamine RNAiMAX

transfection reagent (Thermo Fisher Scientific), according to manufacturer's

instructions.

Subcellular fractionation. The protocol for sub-cellular fractionation was as described

before[30]. Briefly, for each experiment, approximately 1.0×10^7-1.0×10^8 cells were

harvested. The cell pellet was re-suspended in 3 to 5 volumes of hypotonic Buffer A

(10 mM Tris-HCl, pH 7.4, 100 mM NaCl, 2.5 mM $MgCl_2$) supplemented with 0.5 %

Triton X-100, protease and phosphatase inhibitors (1 mM NaF, 1 mM Na_3VO_4) and

incubated on ice for 10 min. Cell membranes were sheared by passing the suspension

4-6 times through a 26-gauge syringe. Nuclei were isolated by centrifugation at 3000 x

g for 10 min at 4°C, and the supernatant was kept as cytoplasmic extract. The nuclear

pellet was washed once and the nuclei were resuspended in 2 volumes of Buffer A and

sonicated twice for 5s (0.2A). Then, samples were centrifuged at 4000 x g for 10 min

at 4°C. The upper phase, which is the nuclear extract, was collected, while the nuclear

pellet was re-suspended in 2 volumes of 8 M Urea and stored at -20°C. Protein

concentration of the isolated fractions was assessed using the Bradford assay[53].

For the subnuclear fractionation protocol that allows for analysis of the LASR

complex[31] cells were harvested, incubated on ice in Buffer B (10 mM HEPES-KOH pH

7.5, 15 mM KCl, 1.5 mM EDTA, 0.15 mM spermine) for 30 min and lysed with the

addition of 0.3 % Triton X-100. Nuclei were collected by centrifugation and further

purified by re-suspending the pellet in S1 buffer (0.25M Sucrose, 10 mM $MgCl_2$) and

laid over an equal volume of S2 buffer (0.35 M Sucrose, 0.5 mM $MgCl_2$). Purified

nuclei were lysed in ten volumes of ice-cold lysis buffer (20 mM HEPES-KOH pH 7.5,

150 mM NaCl, 1.5 mM $MgCl_2$, 0.5 mM DTT and 0.6 % Triton X-100) and nucleosol

was separated via centrifugation from the high molecular weight fraction (pellet). The

high molecular weight (HMW) fraction was subsequently resuspended in Buffer B and

treated with either DNase I (0.1 mg/ml) or RNase A (0.1 mg/ml). The supernatant was

collected by centrifugation at 20,000 x g for 5 min, as the HMW treated sample.

The nuclear matrix fractionation was as previously described[54]. Briefly, cells were

harvested and washed with PBS. The cell pellet obtained was re-suspended in a five

packed cell pellet volumes of buffer A (10 mM Tris-HCl pH 7.5, 2.5 mM $MgCl_2$, 100

mM NaCl, 0.5% Triton X-100, 0.5 mM DTT and protease inhibitors) and incubated on

ice for 15 min. The cells were then collected by centrifugation at 2000rpm for 10 min

and re-suspended in 2 volumes of buffer A. To break the plasma membrane a Dounce

homogenizer (10 strokes) was used and the cells were checked under the microscope.

After centrifugation at 2000 rpm for 5 min, supernatant was gently removed and kept

as cytoplasmic fraction, while the pellet containing the nuclei was re-suspended in 10

packed nuclear pellet volumes of S1 solution (0.25 M sucrose, 10 mM $MgCl_2$), on top

of which an equal volume of S2 solution (0.35 M sucrose, 0.5 mM $MgCl_2$) was layered.

After centrifugation at 2800 x g for 5 min, the nuclear pellet was re-suspended in 10

volumes of buffer NM (20 mM HEPES pH 7.4, 150 mM NaCl, 2.5 mM $MgCl_2$, 0.6 %

Triton X-100 and Protease inhibitors) and lysed on ice for 10 minutes, followed by

centrifugation as above. The supernatant was removed and kept as nuclear extract while

the pellet was re-suspended in buffer A containing DNase I (0.5mg/mL) or RNase A

(0.1 mg/mL) and Protease inhibitors and stirred gently at room temperature for 30

minutes. The upper phase defining the nuclear matrix fraction was quantified and stored

at -20˚C.

Immunoprecipitation. Co-immunoprecipitation of proteins was performed using

Protein A/G agarose beads (Protein A/G Plus Agarose Beads, Santa-Cruz

Biotechnologies, sc-2003) as follows: 20 µl of bead slurry per immunoprecipitation

reaction was washed with NET-2 buffer (10 mM Tris pH 7.5, 150 mM NaCl, 0.05 %

NP-40). 4-8 µg of antibody were added to a final volume of 500-600 µL in NET-2

buffer per sample. Antibody binding was performed by overnight incubation at 4ºC on

a rotating wheel. Following the binding of the antibody, beads were washed at least 3

times by resuspension in NET-2. For each IP sample, 500-1000 µg of protein were

added to the beads, in a final volume of 800 µL with NET-2 buffer and incubated for 2

hrs at 4ºC on a rotating wheel. After sample binding, beads were washed 3 times with

NET-2 buffer, and twice with NET-2 buffer supplemented with 0.1% Triton X-100 and

a final concentration of 0.1% NP-40. For the UV-crosslinking experiments, beads were

washed five times with wash buffer containing 1M NaCl and twice with standard wash

buffer. Co-immunoprecipitated proteins were eluted from the beads by adding 15-20

µL of 2x Laemmli sample buffer (0.1 M Tris, 0.012% bromophenol blue, 4 % SDS,

0.95 M β-mercapthoethanol, 12 % glycerol) and boiled at 95ºC for 5 min. Following

centrifugation at 10.000 x g for 2 min, the supernatant was retained and stored at -20ºC

or immediately used.

Western Blot analysis. Cell lysate (7-10 µg for nuclear lysates and 15-20 µg for the

cytoplasmic fraction) was resolved on an 8%, 10% or 12 % SDS-polyacrylamide gel

and transferred to a polyvinylidinedifluoride membrane (PVDF, Millipore). Primary

antibodies were added and the membranes were incubated overnight at 4˚C. Antibodies

were used against: hnRNPM (1:500, clone 1D8, sc-20002), IQGAP1 (1:1000, clone

H109, sc-10792; 1:1000, clone C-9, sc-379021; 1:1000 clone D-3, sc-374307, all from

Santa Cruz), beta-actin (1:1000, clone 7D2C10, 60008-1-Ig, ProteinTech), Lamin B1

(1:1000, clone A-11, sc-377000, Santa Cruz), GAPDH (1:2000, 60004-1-Ig,

ProteinTech), hnRNP A2/B1 (1:1000, clone DP3B3, sc-32316, Santa Cruz), SRSF1

(SF2/ASF; 1:1000, SC-33652, Santa Cruz), hnRNP A1 (1:1000, clone 4B10, sc-32301,

Santa Cruz), hnRNPL (1:1000, clone 4D11, sc-32317, Santa Cruz), hnRNP C1/C2

(1:1000, clone 4F4, sc-32308, Santa Cruz), DHX9/RHA (1:1000, ab26271, Abcam),

hnRNP K/J (1:1000, clone 3C2, sc-32307, Santa Cruz), SF3B3 (1:1000, clone B-4, sc-

398670, Santa-Cruz), beta tubulin (1:1000, clone 2-28-33, Sigma-Aldrich), TPX2

(1:200, clone E-2, sc-271570, Santa Cruz), RRM2 (1:200, clone A-5, sc-398294, Santa

Cruz), SAFB (1:1000, F-3, sc-393403, Santa Cruz), Matrin3 (1:1000, 2539C3a, Santa

Cruz), Histone-H3 (1:2000, 17168-AP, ProteinTech), DDX17 (clone H-7, sc-398168,

Santa Cruz), CPSF6 (1:1000, sc-292170, Santa Cruz), NF90 (1:1500, clone A-3, sc-

377406, Santa Cruz), anillin (1:1000, CL0303, ab211872, Abcam) , FZR (1:200, clone

DCS-266, sc-56312, Santa Cruz), TK1 (1:5000, clone EPR3193, ab76495, Abcam),

ANAPC10 (1:100, clone B-1, sc-166790, Santa Cruz). HRP-conjugated goat anti-

mouse IgG (1:5000, 1030-05, SouthernBiotech) or HRP-conjugated goat anti-rabbit

IgG (1:5000, 4050-05, SouthernBiotech) were used as secondary antibodies. Detection

was carried out using Immobilon Crescendo Western HRP substrate (WBLUR00500,

Millipore).

Generation of knockouts. The CRISPR/Cas9 strategy was used to generate IQGAP1

knockout cells[55]. Exon1 of the *IQGAP1* transcript was targeted using the following pair

of synthetic guide RNA (sgRNA) sequences: Assembly 1: 5'-

CACTATGGCTGTGAGTGCG-3' and Assembly 2: 5'-

CAGCCCGTCAACCTCGTCTG-3'. The sequences were identified using the CRISPR

Design tool (http://crispr.mit.edu/). These sequences and their reverse complements

were annealed and ligated into the BbSI and BsaI sites of the All-In-One vector [AIO-

Puro, a gift from Steve Jackson (Addgene plasmid #74630;

http://n2t.net/addgene:74630; RRID:Addgene_74630)][56]. The two pairs of

complementary DNA-oligos (Assemblies 1 and 2 including a 4-mer overhang + 20-mer

of sgRNA sequence) were purchased from Integrated DNA technologies (IDT). The

insertion of sgRNAs was verified via sequencing. MKN45 and NUGC4 cells were

transfected using Lipofectamine 2000, and clones were selected 48 h later using

puromycin. Individual clones were plated to single cell dilution in 24 well-plates, and

IQGAP1 deletion was confirmed by PCR of genomic DNA using the following

primers: Forward: 5'-GCCGTCCGCGCCTCCAAG-3';   Reverse: 5'-

GTCCGAGCTGCCGGCAGC-3' and sequencing using the Forward primer. Loss of

IQGAP1 protein expression was confirmed by Western Blotting. MKN45 and NUGC4

cells transfected with AIO-Puro empty vector were selected with puromycin and used

as a control during the clone screening process.

For the generation of the *hnRNPM* [KO] cells we used a different approach. We ordered a

synthetic guide RNA (sgRNA) (5'- CGGCGTGCCGAGCGGCAACG-3'), targeting

exon 1 of the *hnRNPM* transcript, in the form of crRNA from IDT, together with

tracrRNA. We assembled the tracrRNA:crRNA duplex by combining 24pmol of

tracRNA and 24pmol of crRNA in a volume of 5μl, and incubating at 95°C for 5 min,

followed by incubation at room temperature. 12pmol of recombinant Cas9 (Protein

Expression and Purification Facility, EMBL, Heidelberg) were mixed with 12pmol of

the tracrRNA:crRNA duplex in OPTIMEM I (GIBCO) for 5min at room temperature

and this RNP was used to transfect MKN45 cells in the presence of Lipofectamine

RNAiMax. Cells were harvested 48 h later and individual clones were isolated and

assayed for hnRNPM downregulation as described above for IQGAP1. The primers

used were: Forward: 5'- CACGTGGGCGCGCAGG -3'; Reverse: 5'-

GCAAAGGACCGTGGGATACTCAC -3.

Splicing assay. Splicing assays with the DUP51M1 and DUP50M1 mini-gene reporters

were performed as previously described[31]. Briefly, cells were co-transfected with

DUP51M1 (or DUP50M1) or DUP51-ΔM (or DUP50-ΔM) site plasmids and pCMS-

EGFP at 1:3 ratio for 40 h. Total RNA was extracted using TRIzol Reagent® (Thermo

Fisher Scientific) and cDNA was synthesized in the presence of a DUP51-specific

primer (DUP51-RT, 5'-AACAGCATCAGGAGTGGACAGATCCC-3'). Analysis of

alternative spliced transcripts was carried out with PCR (15-25 cycles) using primers

DUP51S_F (5'-GACACCATCCAAGGTGCAC-3') and DUP51S_R (5'-

CTCAAAGAACCTCTGGGTCCAAG-3'), followed by electrophoresis on 8%

acrylamide-urea gel. Quantification of percentage of exon 2 inclusion was performed

with ImageJ or with ImageLab software (version 5.2, Bio-Rad Laboratories) when ^{32}P-

labelled DUP51S_F primer was used for the PCR. For the detection of the RNA

transcript bound on hnRNPM after UV crosslinking, PCR was performed using primers

DUP51UNS_F (5'-TTGGGTTTCTGATAGGCACTG-3') and DUP51S_R (see

above).

For the validation of the AS events identified by RNA-seq, cDNA was synthesized from

total RNA of appropriate cells in the presence of random hexamer primers and used as

a template in PCR with the primers listed in **Supplementary Table 4**. % inclusion for

each event in 3 or more biological replicates was analysed in 8% acrylamide-urea gel

and quantified by ImageJ.

UV-crosslinking experiments were performed as described[31]. Briefly, monolayer

MKN45 cell cultures after transfection with the minigene reporters, as described above,

were irradiated with UV (254 nm) at 75 mJ/cm^2 on ice in a UV irradiation system BLX

254 (Vilber Lourmat). UV-irradiated cells were lysed for 5 min on ice with ten packed

cell volumes of buffer [20 mM HEPES-KOH pH 7.5, 150 mM NaCl, 0.5 mM DTT, 1

mM EDTA, 0.6% Triton X-100, 0.1% SDS, and 50mg/ml yeast tRNA] and centrifuged

at 20,000 x g for 5 min at 4^{0}C. The supernatants were 5 x diluted with buffer [20 mM

HEPES-KOH pH 7.5, 150 mM NaCl, 0.5 mM DTT, 1 mM EDTA, 1.25x Complete

protease inhibitors (Roche), and 50 µg/ml yeast tRNA]. Lysates were centrifuged for

10 min at 20,000 x g, 4^{0}C prior to IP.

MTT cell proliferation assay. In 96-well plates, cells were seeded at a density of 2×10^3

cells/well in complete RPMI. After 24 hrs, the medium was replaced with serum-free

RPMI supplemented with 0.5mg/ml MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl

tetrazolium bromide; Sigma-Aldrich) colorimetric staining solution for 2h. After the

removal of MTT, the cells were mixed with 200µl DMSO and incubated for 5-10min

at RT on a shaking platform. The absorbance was read at 570 nm using the SUNRISE^{TM}

Absorbance Reader (Tecan Trading AG).

Colony-Formation assay. In 6-well plates, 200 cells/well were placed and allowed to

grow for 7 days at 37^{o}C with 5% CO_2. The formed colonies were fixed with 0.5mL of

100% methanol for 20min at RT. Methanol was then removed and cells were carefully

rinsed with H_2O. 0.5ml crystal violet staining solution (0.5% crystal violet in 10%

ethanol) was added to each well and cells were left for 5min at RT. The plates were

then washed with H_2O until excess dye was removed and were left to dry. The images

were captured by Molecular Imager® ChemiDoc^{TM} XRS+ Gel Imaging System (Bio-

Rad) and colonies were quantified using ImageJ software.

Wound healing assay. Cells were cultured in 24-well plates at 37^{o}C with 5% CO_2 in a

monolayer, until nearly 90% confluent. Scratches were then made with a sterile 200µl

pipette tip and fresh medium without FBS was gently added. The migration of cells in

the same wound area was visualized at 0, 8, 24, 32 and 48 hrs using Axio Observer A1

(Zeiss) microscope with automated stage.

Cell Cycle Analysis. The cells were seeded in 6-well plates at a density of 3×10^5

cells/well. When cells reached 60-80% confluence, they were harvested by

trypsinization into phosphate-buffered saline (PBS). The pellets were fixed in 70%

ethanol and stored at -20°C till all time-points were collected. On the day of the FACS

analysis, cell pellets were washed in phosphate-citrate buffer and centrifuged for 20min.

250μl of RNase/propidium iodide (PI) solution were then added to each sample (at

concentrations of 100μg/ml for RNase and 50μg/ml for PI) and cells were incubated at

37°C for 30min. Finally, the cells were analysed through flow cytometric analysis using

FACSCanto[TM] II (BD-Biosciences).

Mouse Xenograft study. Experiments were performed in the animal facilities of

Biomedical Sciences Research Center (BSRC) "Alexander Fleming" and were

approved by the Institutional Committee of Protocol Evaluation in conjunction with the

Veterinary Service Management of the Hellenic Republic Prefecture of Attika

according to all current European and national legislation and performed in accordance

with the guidance of the Institutional Animal Care and Use Committee of BSRC

"Alexander Fleming". 1 x10^6 cells of MKN45, MKN45-IQGAP1[KO], MKN45-hnRNP-

M[KO] or double KO cells were injected into the flank of 8-10-week-old NOD-SCID[57]

both male and female mice randomly distributed among groups. Groups of 11 mice

were used per cell type, based on power analysis performed using the following

calculator: https://www.stat.ubc.ca/~rollin/stats/ssize/n2.html. Tumour growth was

monitored up to 4 weeks and recorded by measuring two perpendicular diameters using

the formula 1/2(Length × Width^2) bi-weekly[58]. During the experiment, the investigator

contacting the measurements was unaware of the sample group allocation (blinded

experiment). At end-point, mice were euthanized and tumours were collected and

enclosed in paraffin for further analyses.

Immunostaining. For immunofluorescence, cells were seeded on glass coverslips and

were left to adhere for 24 hrs. Cells were next fixed for 10 minutes with 4%

paraformaldehyde PFA (Alfa Aesar), followed by permeabilization with 0.25% (w/v)

Triton X-100. Cells were then incubated for 30 min in 5% BSA/PBS (phosphate buffer

saline). The primary antibodies used for immunostaining were: anti-hnRNPM (1:300,

clone 1D8, NB200-314SS, Novus or 1:100, NBP1-84555, Novus), anti-IQGAP1

(1:500, ab86064, Abcam or 1:250 sc-374307, Santa-Cruz), anti-SR (1:100, clone 1H4,

sc-13509, Santa-Cruz), recognizing SRp75, SRp55, SRp40, SRp30a/b and SRp20, anti-

PSF (1:100, clone H-80, sc-28730, Santa-Cruz). For β-TUBULIN staining, cells were

fixed in -20°C with ice-cold methanol for 3 minutes, blocked in 1% BSA/PBS solution

and incubated overnight with the primary antibody (1:250, clone 2-28-33, Sigma-

Aldrich). After washing with PBS, cells were incubated with secondary antibodies

(anti-rabbit-Alexa Fluor 555 or anti-mouse Alexa Fluor 488, both used at 1:500;

Molecular Probes) at room temperature for 1h followed by the staining of nuclei with

DAPI for 5 min at RT. For mounting Mowiol mounting medium (Sigma-Aldrich) was

used and the images were acquired with Leica DM2000 fluorescence microscope or a

LEICA SP8 White Light Laser confocal system and were analysed using the Image J

software.

Tissue Microarrays (TMA) slides were purchased from US Biomax, Inc (cat. no.

T012a). The slides were deparaffinized in xylene and hydrated in different alcohol

concentrations. Heat-induced antigen retrieval in citrate buffer pH 6.0 was used.

Blocking, incubation with first and secondary antibodies as well as the nuclei staining

and mounting, were performed as mentioned above.

Microscopy and image analysis. Fluorescent images were acquired with a Leica TCS

SP8 X confocal system equipped with an argon and a supercontinuum white light laser

source, using the LAS AF software (Leica). The same acquisition settings were applied

for all samples. Pixel-based colocalization analysis was performed with the Image J

software, using the "Colocalization Threshold" plugin[59] (Costes et al, 2002) to calculate

the Pearson correlation coefficient. Image background was subtracted using the

"Substract background" function of Image J (50px ball radius). For each image, the

middle slices representing the cell nuclei (selected as regions of interest (ROI) based

on the DAPI signal) were chosen for analysis and at least 30 cells or more were analysed

for each cell line. Intensity plot profiles (k-plots) were generated using the "Plot profile"

function of Image J. After background substraction (as mentioned above), a line was

drawn across each cell and the pixel gray values for hnRNPM, SR & PSF signals were

acquired. Adobe Photoshop CS6 was used for merging the final images, where

brightness and contrast were globally adjusted.

Immunohistochemistry and H&E staining. At the end of the xenograft experiment

tumours were dissected from the mice, fixed in formalin and embedded in paraffin.

Sections were cut at 5 μm thickness, were de-paraffinized and stained for haematoxylin

and eosin. For IHC, after de-paraffinization serial sections were hydrated, incubated in

3% H_2O_2 solution for 10 minutes, washed and boiled at 95°C for 15 minutes in sodium

citrate buffer pH 6.0 for antigen retrieval. Blocking was performed with 5% BSA for 1

hr and sections were then incubated with the following primary antibodies overnight at

4°C diluted in BSA: anti Ki-67 (1:200, clone 14-5698-82, ThermoFisher Scientific),

hnRNP-M (1:100, clone 1D8, sc-20002, Santa Cruz), IQGAP1 (1:100, ab86064,

Abcam). Sections were subsequently washed and incubated with the appropriate

secondary antibody conjugated to HRP, HRP-conjugated goat anti-mouse IgG (1:5000,

1030-05, SouthernBiotech) or HRP-conjugated goat anti-rabbit IgG (1:5000, 4050-05,

SouthernBiotech) and the DAB Substrate Kit (SK-4100, Vector Laboratories) was used

to visualise the signal. The sections were counterstained with hematoxylin and imaged

with a NIKON Eclipse E600 microscope, equipped with a Qcapture camera.

Proximity ligation assay. Cells were grown on coverslips (13 mM diameter, VWR)

and fixed for 10 min with 4% PFA (Alfa Aesar), followed by 10 min permeabilization

with 0.25% Triton X-100 in PBS and blocking with 5% BSA in PBS for 30 min.

Primary antibodies: anti-hnRNPM (1:300, clone 1D8, NB200-314SS, Novus), anti-

IQGAP1 (1:500, ab86064, Abcam or 1:500, 22167-1-AP, Proteintech) diluted in

blocking buffer were added and incubated overnight at 4˚C. Proximity ligation assays

were performed using the Duolink kit (Sigma-Aldrich DUO92102), according to

manufacturer's protocol. Images were collected using a Leica SP8 confocal

microscope.

RNA isolation and reverse transcription. Total RNA was extracted with the TRIzol®

reagent (Thermo Fisher Scientific). DNA was removed with RQ1 RNase-free DNase

(Promega, WI) or DNase I (RNase-free, New England Biolabs, Inc, MA), followed by

phenol extraction. Reverse transcription was carried with 0.4-1 µg total RNA in the

presence of gene-specific or random hexamer primers, RNaseOUT™ Recombinant

Ribonuclease Inhibitor (Thermo Fisher Scientific) and SuperScript® III (Thermo

Fisher Scientific) or Protoscript II (New England Biolabs) reverse transcriptase,

according to manufacturer's instructions.

Mass spectrometry and Proteomics analysis. Anti-IQGAP1 immunoprecipitation

samples were processed in collaboration with the Core Proteomics Facility at EMBL

Heidelberg. Proteomics analysis[60] was performed as follows: samples were dissolved

in 2x Laemmli sample buffer, and underwent filter-assisted sample preparation (FASP)

to produce peptides with proteolytic digestion[61]. These were then tagged using 4

different multiplex TMT isobaric tags (ThermoFisher Scientific, TMTsixplex™

Isobaric Label Reagent Set): one isotopically unique tag for each IP condition, namely

IQGAP1 IP cancer (NUGC4) and the respective IgG control. TMT-tagged samples

were appropriately pooled and analysed using HPLC-MS/MS. Three biological

replicates for each IP condition were processed.

Samples were processed using the ISOBARQuant[62], an R-package platform for the

analysis of isobarically labelled quantitative proteomics data. Only proteins that were

quantified with two unique peptide matches were filtered. After batch-cleaning and

normalization of raw signal intensities, fold-change was calculated. Statistical analysis

of results was performed using the LIMMA[63] R-package, making comparisons between

each IQGAP1 IP sample and their respective IgG controls. A protein was considered

significant if it had a *P*value < 5% (Benjamini-Hochberg FDR adjustment), and a fold-

change of at least 50% between compared conditions. Identified proteins were

classified into 3 categories: Hits (FDR threshold= 0.05, fold change=2), candidates

(FDR threshold = 0.25, fold change = 1.5), and no hits (see **Supplementary Table 1**).

For the differential proteome analysis of MKN45 and MKN45-*IQGAP1^{KO}* cells, whole

cell lysates were prepared in RIPA buffer [25 mM Tris-HCl (pH 7.5), 150 mM NaCl,

1% NP-40, 0.5% sodium deoxycholate, 0.1% SDS). Samples underwent filter-assisted

sample preparation (FASP) to produce peptides with proteolytic digestion[61] and

analysed using HPLC-MS/MS. The full dataset is being prepared to be published

elsewhere.

RNA-seq analysis. Total TRIzol-extracted RNA was treated with RQ1-RNase free

DNase (Promega). cDNA libraries were prepared in collaboration with Genecore, at

EMBL, Heidelberg. Alternative splicing was analyzed by using VAST-TOOLS

v2.2.2[64] and expressed as changes in percent-spliced-in values (ΔPSI). A minimum read

coverage of 10 junction reads per sample was required, as described[64]. Psi values for

single replicates were quantified for all types of alternative events. Events showing

splicing change (|ΔPSI|> 15 with minimum range of 5% between control and

IQGAP1[KO] samples were considered IQGAP1-regulated events.

ORF impact prediction. Potential ORF impact of alternative exons was predicted as

described[64]. Exons were mapped on the coding sequence (CDS) or 5'/3' untranslated

regions (UTR) of genes. Events mapping on the CDS were divided in CDS-preserving

or CDS-disrupting.

RNA maps. We compared sequence of introns surrounding exons showing more

inclusion or skipping in IQGAP1[KO] samples with a set of 1,050 not changing alternative

exons. To generate the RNA maps, we used the rna_maps function[65], using sliding

windows of 15 nucleotides. Searches were restricted to the affected exons, the first and

last 500 nucleotides of the upstream and downstream intron and 50 nucleotides into the

upstream and downstream exons. Regular expression was used to search for the binding

motif   of   hnRNPM   (GTGGTGG|GGTTGGTT|GTGTTGT|TGTTGGAG   or

GTGGTGG|GGTTGGTT|TGGTGG|GGTGG)[13].

Gene Ontology. Enrichment for GO terms was analyzed using ShinyGO v0.61[66] with

P value cut-off (FDR) set at 0.05.

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange

Consortium via the PRIDE68 partner repository[67] with the dataset identifier

PXD017842.

RNA-seq data have been deposited in GEO: GSE146283.

Furthermore, the data and/or reagents that support the findings of this study are

available from the corresponding author, P.K., upon reasonable request.

Source data for Figs. 1-7 and Supplementary Data Figs. 1-6 will be provided online.

References

1.   Wang, Z. & Burge, C. B. Splicing regulation: From a parts list of regulatory

elements to an integrated splicing code. RNA 14, 802–813 (2008).

2.   Pan, Q., Shai, O., Lee, L. J., Frey, B. J. & Blencowe, B. J. Deep surveying of

alternative splicing complexity in the human transcriptome by high-throughput

sequencing. Nat. Genet. 40, 1413–1415 (2008).

3.   Heyd, F. & Lynch, K. W. Degrade, move, regroup: signaling control of

splicing proteins. Trends Biochem. Sci. 36, 397–404 (2011).

4.   El Marabti, E. & Younis, I. The Cancer Spliceome: Reprograming of

Alternative Splicing in Cancer. Front. Mol. Biosci. 5, (2018).

5.   Oltean, S. & Bates, D. O. Hallmarks of alternative splicing in cancer.

Oncogene (2013) doi:10.1038/onc.2013.533.

6.   Sveen, A., Kilpinen, S., Ruusulehto, A., Lothe, R. A. & Skotheim, R. I.

Aberrant RNA splicing in cancer; expression changes and driver mutations of splicing

factor genes. Oncogene (2015) doi:10.1038/onc.2015.318.

7.   Kahles, A. et al. Comprehensive Analysis of Alternative Splicing Across

Tumours from 8,705 Patients. Cancer Cell 34, 211-224.e6 (2018).

8.   Shkreta, L. & Chabot, B. The RNA Splicing Response to DNA Damage.

Biomolecules 5, 2935–2977 (2015).

9.   Datar, K. V., Dreyfuss, G. & Swanson, M. S. The human hnRNP M proteins:

identification of a methionine/arginine-rich repeat motif in ribonucleoproteins.

Nucleic Acids Res. 21, 439–446 (1993).

10.   Kafasla, P., Patrinou-Georgoula, M., Lewis, J. D. & Guialis, A. Association of

the 72/74-kDa proteins, members of the heterogeneous nuclear ribonucleoprotein M

group, with the pre-mRNA at early stages of spliceosome assembly. Biochem. J. 363,

793–799 (2002).

11.   Llères, D., Denegri, M., Biggiogera, M., Ajuh, P. & Lamond, A. I. Direct

interaction between hnRNP-M and CDC5L/PLRG1 proteins affects alternative splice

site choice. EMBO Rep. 11, 445–451 (2010).

12.   Hovhannisyan, R. H. & Carstens, R. P. Heterogeneous ribonucleoprotein m is

a splicing regulatory protein that can enhance or silence splicing of alternatively

spliced exons. J. Biol. Chem. 282, 36265–36274 (2007).

13.   Huelga, S. C. et al. Integrative Genome-wide Analysis Reveals Cooperative

Regulation of Alternative Splicing by hnRNP Proteins. Cell Rep. 1, 167–178 (2012).

14.   Pervouchine, D. et al. Integrative transcriptomic analysis suggests new

autoregulatory splicing events coupled with nonsense-mediated mRNA decay.

Nucleic Acids Res. 47, 5293–5306 (2019).

15.   Gattoni, R. et al. The human hnRNP-M proteins: structure and relation with

early heat shock-induced splicing arrest and chromosome mapping. Nucleic Acids

Res. 24, 2535–2542 (1996).

16.   Shalgi, R., Hurt, J. A., Lindquist, S. & Burge, C. B. Widespread Inhibition of

Posttranscriptional Splicing Shapes the Cellular Transcriptome following Heat Shock.

Cell Reports **7**, 1362–1370 (2014).

17.   Xu, Y. et al. Cell type-restricted activity of hnRNPM promotes breast cancer

metastasis via regulating alternative splicing. Genes Dev. 28, 1191–1203 (2014).

18.   Harvey, S. E. et al. Coregulation of alternative splicing by hnRNPM and

ESRP1 during EMT. RNA 24, 1326–1338 (2018).

19.   Hu, X. et al. The RNA-binding protein AKAP8 suppresses tumour metastasis

by antagonizing EMT-associated alternative splicing. Nat. Commun. 11, 486 (2020).

20.   Passacantilli, I., Frisone, P., De Paola, E., Fidaleo, M. & Paronetto, M. P.

hnRNPM guides an alternative splicing program in response to inhibition of the

PI3K/AKT/mTOR pathway in Ewing sarcoma cells. Nucleic Acids Res. 45, 12270–

12284 (2017).

21.   West, K. O. et al. The Splicing Factor hnRNP M Is a Critical Regulator of

Innate Immune Gene Expression in Macrophages. Cell Rep. 29, 1594-1609.e5 (2019).

22.   Smith, J. M., Hedman, A. C. & Sacks, D. B. IQGAPs choreograph cellular

signaling from the membrane to the nucleus. Trends Cell Biol. 25, 171–184 (2015).

23.   Osman, M. A., Sarkar, F. H. & Rodriguez-Boulan, E. A molecular rheostat at

the interface of cancer and diabetes. Biochim. Biophys. Acta 1836, 166–176 (2013).

24.   White, C. D., Brown, M. D. & Sacks, D. B. IQGAPs in cancer: A family of

scaffold proteins underlying tumourigenesis. FEBS Lett. 583, 1817–1824 (2009).

25.   Hu, W. et al. IQGAP1 promotes pancreatic cancer progression and epithelial-

mesenchymal transition (EMT) through Wnt/β-catenin signaling. Sci. Rep. 9, 7539

(2019).

26.   Li, S., Wang, Q., Chakladar, A., Bronson, R. T. & Bernards, A. Gastric

Hyperplasia in Mice Lacking the Putative Cdc42 Effector IQGAP1. Mol. Cell. Biol.

20, 697–701 (2000).

27.   Johnson, M., Sharma, M., Brocardo, M. G. & Henderson, B. R. IQGAP1

translocates to the nucleus in early S-phase and contributes to cell cycle progression

after DNA replication arrest. Int. J. Biochem. Cell Biol. 43, 65–73 (2011).

28.   Cvitkovic, I. & Jurica, M. S. Spliceosome database: a tool for tracking

components of the spliceosome. Nucleic Acids Res. 41, D132-141 (2013).

29.   Choi, Y. D. & Dreyfuss, G. Isolation of the heterogeneous nuclear RNA-

ribonucleoprotein complex (hnRNP): a unique supramolecular assembly. Proc. Natl.

Acad. Sci. U. S. A. 81, 7471–7475 (1984).

30.   Kafasla, P., Patrinou-Georgoula, M. & Guialis, A. The 72/74-kDa

polypeptides of the 70-110 S large heterogeneous nuclear ribonucleoprotein complex

(LH-nRNP) represent a discrete subset of the hnRNP M protein family. Biochem. J.

350 Pt 2, 495–503 (2000).

31.   Damianov, A. et al. Rbfox Proteins Regulate Splicing as Part of a Large

Multiprotein Complex LASR. Cell 165, 606–619 (2016).

32.   Zhu, Y. et al. POSTAR2: deciphering the post-transcriptional regulatory

logics. Nucleic Acids Res. 47, D203–D211 (2019).

33.   Van Nostrand, E. L. et al. Robust transcriptome-wide discovery of RNA-

binding protein binding sites with enhanced CLIP (eCLIP). Nat. Methods 13, 508–

514 (2016).

34.   Yamano, H. APC/C: current understanding and future perspectives.

F1000Research 8, (2019).

35.   Zhou, Z., He, M., Shah, A. A. & Wan, Y. Insights into APC/C: from cellular

function to diseases and therapeutics. Cell Div. 11, 9 (2016).

36.   da Fonseca, P. C. A. et al. Structures of APC/C(Cdh1) with substrates identify

Cdh1 and Apc10 as the D-box co-receptor. Nature 470, 274–278 (2011).

37.   Alfieri, C., Zhang, S. & Barford, D. Visualizing the complex functions and

mechanisms of the anaphase promoting complex/cyclosome (APC/C). Open Biol. 7,

(2017).

38.   Engström, Y. et al. Cell cycle-dependent expression of mammalian

ribonucleotide reductase. Differential regulation of the two subunits. J. Biol. Chem.

260, 9114–9116 (1985).

39.   Neumayer, G., Belzil, C., Gruss, O. J. & Nguyen, M. D. TPX2: of spindle

assembly, DNA damage response, and cancer. Cell. Mol. Life Sci. CMLS 71, 3027–

3047 (2014).

40. Sherley, J. L. & Kelly, T. J. Regulation of human thymidine kinase during the

cell cycle. J. Biol. Chem. 263, 8350–8358 (1988).

41. Meissner, M. *et al.* Differential nuclear localization and nuclear matrix

association of the splicing factors PSF and PTB. *J. Cell. Biochem.* **76**, 559–566

(2000).

42. Biamonti, G. & Caceres, J. F. Cellular stress and RNA splicing. Trends

Biochem. Sci. 34, 146–153 (2009).

43. Lian, A. T., Hains, P. G., Sarcevic, B., Robinson, P. J. & Chircop, M. IQGAP1

is associated with nuclear envelope reformation and completion of abscission. Cell

Cycle 14, 2058–2074 (2015).

44. Cyclebase 3.0: a multi-organism database on cell-cycle regulation and

phenotypes | Nucleic Acids Research | Oxford Academic.

https://academic.oup.com/nar/article/43/D1/D1140/2437426.

45. Penas, C., Ramachandran, V. & Ayad, N. G. The APC/C Ubiquitin Ligase:

From Cell Biology to Tumourigenesis. Front. Oncol. 1, 60 (2011).

46. Levine, M. S. & Holland, A. J. The impact of mitotic errors on cell

proliferation and tumourigenesis. Genes Dev. 32, 620–638 (2018).

47. Sansregret, L. et al. APC/C Dysfunction Limits Excessive Cancer

Chromosomal Instability. Cancer Discov. 7, 218–233 (2017).

48. Gijn, S. E. van et al. TPX2/Aurora kinase A signaling as a potential

therapeutic target in genomically unstable cancer cells. Oncogene 38, 852–867

(2019).

49.   Jeong, I. et al. GEMiCCL: mining genotype and expression data of cancer cell

lines with elaborate visualization. Database 2018, (2018).

50.   Gao, J. et al. Integrative analysis of complex cancer genomics and clinical

profiles using the cBioPortal. Sci. Signal. 6, pl1 (2013).

51.   Cerami, E. et al. The cBio cancer genomics portal: an open platform for

exploring multidimensional cancer genomics data. Cancer Discov. 2, 401–404 (2012).

52.   Ren, J.-G., Li, Z., Crimmins, D. L. & Sacks, D. B. Self-association of

IQGAP1: characterization and functional sequelae. J. Biol. Chem. 280, 34548–34557

(2005).

53.   Bradford, M. M. A rapid and sensitive method for the quantification of

microgram quantities of protein utilizing the principle of protein-dye binding. Anal.

Biochem. 72, 248–254 (1976).

54.   Mähl, P., Lutz, Y., Puvion, E. & Fuchs, J. P. Rapid effect of heat shock on two

heterogeneous nuclear ribonucleoprotein-associated antigens in HeLa cells. J. Cell

Biol. 109, 1921–1935 (1989).

55.   Ran, F. A. et al. Genome engineering using the CRISPR-Cas9 system. Nat.

Protoc. 8, 2281–2308 (2013).

56.   Chiang, T.-W. W., le Sage, C., Larrieu, D., Demir, M. & Jackson, S. P.

CRISPR-Cas9D10A nickase-based genotypic and phenotypic screening to enhance

genome editing. Sci. Rep. 6, 24356 (2016).

57.   Shultz, L. D. et al. Multiple defects in innate and adaptive immunologic

function in NOD/LtSz-scid mice. J. Immunol. Baltim. Md 1950 154, 180–191 (1995).

58.   Euhus, D. M., Hudd, C., LaRegina, M. C. & Johnson, F. E. Tumour

measurement in the nude mouse. J. Surg. Oncol. 31, 229–234 (1986).

59.   Costes, S. V. *et al.* Automatic and Quantitative Measurement of Protein-

Protein Colocalization in Live Cells. *Biophysical Journal* **86**, 3993–4003 (2004).

60.   Aebersold, R. & Mann, M. Mass spectrometry-based proteomics. Nature 422,

198–207 (2003).

61.   Wiśniewski, J. R., Zougman, A., Nagaraj, N. & Mann, M. Universal sample

preparation method for proteome analysis. Nat. Methods 6, 359–362 (2009).

62.   Breitwieser, F. P. et al. General Statistical Modeling of Data from Protein

Relative Expression Isobaric Tags. J. Proteome Res. 10, 2758–2766 (2011).

63.   Smyth, G. K. Linear models and empirical bayes methods for assessing

differential expression in microarray experiments. Stat. Appl. Genet. Mol. Biol. 3, 1–

25 (2004).

64.   Irimia, M. et al. A highly conserved program of neuronal microexons is

misregulated in autistic brains. Cell 159, 1511–1523 (2014).

65.   Gohr, A. & Irimia, M. Matt: Unix tools for alternative splicing analysis.

Bioinforma. Oxf. Engl. 35, 130–132 (2019).

66.   Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool

for animals and plants. Bioinformatics doi:10.1093/bioinformatics/btz931.

67.   Perez-Riverol, Y., et al. The PRIDE database and related tools and resources

in 2019: improving support for quantification data. Nucleic Acids Res 47(D1):D442-

D450 (2019).

68.   McQuin, C. et al. CellProfiler 3.0: Next-generation image processing for

biology. PLOS Biol. 16, e2005970 (2018).

69.   Rando, O. J., Zhao, K. & Crabtree, G. R. Searching for a function for nuclear

actin. Trends Cell Biol. 10, 92–97 (2000).

Acknowledgements:

We thank N. Boni-Kazantzidou (Centre for Proteome Research, Liverpool, UK) and

G.-R. Manikas (European Commission, Brussels, Belgium) for the generation of crucial

preliminary data; P. Hantzis, M. Fousteri, V. Koliaraki (Institute for Fundamental

Biomedical Research, B.S.R.C. "Al. Fleming") and N. Balatsos (Dept. of Biochemistry

and Biotechnology, University of Thessaly, Greece) for cell lines and reagents; D.

Black and A. Damianov (Department of Microbiology, Immunology, and Molecular

Genetics, UCLA, USA) for plasmids and technical advice on minigene reporter splicing

assays; A. Guialis (N.H.R.F., Athens, Greece) for antibodies and reagents; Per

Haberkant and the EMBL Proteomics Core Facility for LC-MS/MS analyses and

advice; Sofia Grammenoudi and the Flow cytometry facility of B.S.R.C. "Al. Fleming"

for help with cell cycle analyses and disussions; Vladimir Benes, Jonathan Landry and

the EMBL Genecore for RNA-seq analyses and discussions; Martina Samiotaki,

George Stamatakis at the Proteomics Facility of B.S.R.C. "Al. Fleming" for LC-

MS/MS analyses and discussions. We also thank George Panayotou and Efthimios

Skoulakis (B.S.R.C. "Al. Fleming") for critical reading of the manuscript; Juan

Valcarcel for help with the analysis of the RNA-seq data; Skarlatos G. Dedos

(Department of Biology, National and Kapodistrian University of Athens, Greece) for

reagents, plasmids, discussions and critical reading of the manuscript.

A.B. was supported by InfrafrontierGR/Phenotypos Infrastructure, co-funded by Greece and the European Union (European Regional Development Fund), under NSRF 2014-2020, MIS 5002135 and by the start-up fund of P.K. M.P. was supported by the start-up fund of P.K. V.P. and Z.E. were supported by the Hellenic Foundation for Research & Innovation (HFRI) and the General Secretariat for Research and Technology (GSRT) to Z.E., under grant agreement 846. J.L. was supported by EMBL Core Facilities. M.R. was supported by the European Research Council (ERC AdvG 670146). M.A. and D.K were supported by InfrafrontierGR/Phenotypos (MIS 5002135). This work was funded by a European Commission Grant FP7-PEOPLE-2010-IEF (274837) to P.K and was supported by a start-up fund to P.K. from a Stavros Niarchos Foundation donation to BSRC "Al. Fleming" and by the Hellenic Foundation for Research & Innovation (HFRI) and the General Secretariat for Research and Technology (GSRT), under grant agreement 846.

Author contributions

These authors contributed equally: Andrada Birladeanu, Malgorzata Rogalska, Myrto Potiri.

A.B. performed most of the immunofluorescence, subcellular fractionation, immunostaining and *in vitro* splicing experiments. M.R. performed all the statistical and bioinformatics analyses. M.P. performed the cellular assays (cell cycle, wound healing, colony formation, MTT assays). Z.E. performed RT-PCR analyses and supervised A.B in *in vitro* splicing assays and M.P in cell cycle analyses and edited the manuscript. V.P. performed immunofluorescence analyses, western blotting analyses, advised A.B. on image acquisition and analyses, supervised M.P. on cellular assays and

edited the manuscript. M.A. and P.K. performed the xenograft experiment. D.K.

advised M.A. and helped with the cellular assays. J.D.L. advised and edited the

manuscript. P.K conceived, designed and supervised the study, performed the

immunoprecipitation experiments and wrote the manuscript. All authors commented on

the manuscript.

Competing interests

The authors declare no competing interests.

Materials & Correspondence

Correspondence and material requests should be addressed to P.K.

Fig. 1

Fig. 1: hnRNPM and IQGAP1 expression levels are altered in gastric cancer. a
Representative epifluorescence images of normal and adenocarcinoma gastric tissues
on a commercial tissue microarray. Tissues were immunostained with mouse anti-
hnRNPM and rabbit anti-IQGAP1 antibodies. DAPI was used for nuclei staining. The
same settings for hnRNPM and IQGAP1 signal acquisition were applied in all
samples. **b** Expression matrix plot showing the *IQGAP1* and *HNRNPM* mRNA levels
in tumour samples (T) from Stomach Adenocarcinoma (STAD) patients in
comparison to TCGA normal and GTex data. The density of colour in each block
represents the median expression value of a gene in a given tissue, normalized by the
maximum median expression value across all blocks. The expression levels are
indicated in $\log_2(TPM + 1)$ values. The plot was generated using the gepia2 website
(http://gepia2.cancer-pku.cn/#indexn). Sample numbers: 408 tumours, 36 normal and
175 stomach GTex. **c** Correlation plots of the expression of *hnRNPM* and *IQGAP1*
mRNAs in normal gastric tissues (GTEx database), normal-adjacent tissues (TCGA
STAD Normal database) and tumours from Stomach Adenocarcinoma (STAD)
patients (TCGA STAD Tumour database). The plots were generated using the gepia2
website. R: Spearman correlation coefficient. Sample numbers were as in (**b**). **d**
Immunoblotting of crude protein extracts from different gastric cancer cell lines
against hnRNPM and IQGAP1. β-actin was used to normalize hnRNPM and IQGAP1
levels. MKN45 cells were used as a reference for the comparison of the expression
levels between cell lines (ratios shown at the bottom). Quantification was performed
using ImageLab software version 5.2 (Bio-Rad Laboratories). AGS: gastric
adenocarcinoma; MKN45: poorly differentiated gastric adenocarcinoma, liver
metastasis; KATOIII: gastric carcinoma, pleural effusion and supraclavicular and

axillary lymph nodes and Douglas cul-de-sac pleural; NUGC4: poorly differentiated

gastric adenocarcinoma, gastric lymph node.

Fig. 2

Fig. 2 hnRNPM and IQGAP1 act co-operatively to regulate tumour growth *in vitro* and *in vivo*. a MTT cell proliferation assays to measure the metabolic activity of gastric cancer cell lines (MKN45, MKN45-*IQGAP1^{KO}*, MKN45-*hnRNPM^{KO}* and double MKN45-*IQGAP1^{KO}*-*hnRNPM^{KO}*, left panel; NUGC4 and NUGC4-*IQGAP1^{KO}* transfected with siRNAs for scrambled control (scr) or hnRNPM (siM); right panel). Growth curves show the mean absorbance values/day from 3 independent experiments. *P* values were calculated using two-tailed, unpaired t-tests, where **$p < 0.01$. **b** 2D colony formation assays to measure the ability of the MKN45-derived cell lines used in (**a**) to form colonies. Bar graph shows the average fold difference ± SD between the numbers of colonies formed across the different groups from 3 independent experiments. *P*-values were calculated using two-tailed, unpaired t-tests, where *$p < 0.05$, ***$p < 0.001$. Representative images for each cell line are shown at the bottom. **c** Cell cycle analysis of asynchronous MKN45-derived cell lines using propidium iodide staining followed by FACS analysis. Quantification of the percentage of cells in each cell cycle phase was performed with FlowJo software. Data represent mean values ± SD of two independent experiments. ***$p < 0.001$, ****$p < 0.0001$. **d** MKN45-derived cell lines were subcutaneously injected into the flanks of NOD/SCID mice and tumours were left to develop over a period of 28 days. The tumour growth graph shows the increase of tumour volume (mm^3) over time. Tumour size was measured in anesthetised mice with a digital caliper twice per week, and at the end-point of the experiment when tumours were excised. Data presented are average values ± SD, from 11 mice per group. *P* values were calculated using one-way ANOVA, where *$p < 0.05$, **$p < 0.01$, ****$p < 0.0001$. **e** Excised tumours from all four cell lines were embedded in paraffin, sectioned and stained for H&E and also

probed with antibodies against hnRNPM, IQGAP1 and Ki67. Representative pictures from one tumour from every cell group are shown.

Fig. 3

Fig. 3: IQGAP1 and hnRNPM interact in the nucleus of gastric cancer cell lines.

a Immunoprecipitation with an anti-IQGAP1 antibody or control IgG using nuclear extracts from MKN45 and NUGC4 cells, in the presence or absence of RNase A (0.1 mg/ml for 30 min). Immunoblot detection of IQGAP1 and hnRNPM is shown in comparison to $1/70^{th}$ of the input used. **b** Representative confocal images of MKN45 and NUGC4 cells stained with an anti-IQGAP1 antibody and DAPI to visualise the nuclei. Single confocal slices are shown for each fluorescence signal and for the merged image. Cross sections of the xz and yz axes show the presence of IQGAP1 within the cell nuclei. **c, d** Proximity ligation assay (PLA) in MKN45 and NUGC4 cells showing the nuclear interaction between hnRNPM and IQGAP1. Representative images for the 2 cell lines displaying a central plane from confocal z-stacks are shown in **(c)**. Negative control (secondary antibody only, MKN45 C and NUGC4 C) samples show minimal background signal. Quantification (**d**) of the nuclear (n.MKN45 and n.NUGC4) and cytoplasmic signal (c.MKN45 and c.NUGC4) was performed per cell, using the DuoLink kit-associated software. Each plot represents at least 15 cells analysed. *P* values were calculated using ANOVA multiple comparisons tests; ****p < 0.0001.

a

other RBPs: SFPQ, ELAVL1

U2 snRNP: SF3B3, SF3A, SF3B1, SF3B2

A complex: DHX15, DDX5, THRAP3, RBM25

U1 snRNP: SNRPA

SM proteins: SNRPD2, SNRPA1, SNRPB

SYNCRIP, HNRNPD, HNRNPUL1, HNRNPM, PTBP1, HNRNPC, RBMX, PCBP1, HNRNPR, HNRNPL, HNRNPAB, HNRNPA3, RALY, HNRNPA1, HNRNPA2B1, ILF3, MATR3, HNRNPDL, HNRNPU, HNRNPUL2, HNRNPH1, HNRNPK

IQGAP1

EJC: EIF4A3, ALYREF

SR proteins: SRSF1, SRSF3, TRA2A, TRA2B, EEF1A1

PRP19 complex: PRPF19, HSPA8, EFTUD2, PRPF8, PRPF6, SNRNP200

U5 snRNP

helicase: DDX17, DHX9

b

IP: input | anti-IQGAP1 | IgG
RNaseA: − + −

IQGAP1, hnRNPA1, hnRNPL, hnRNPC1/C2, DHX9/RHA, hnRNPA2B1

NUGC4

MKN45

-170, -31, -72, -42, -125, -31

c

	MKN45	NUGC4	MKN45	NUGC4	MKN45	NUGC4	
170-							IQGAP1
72-							hnRNPM
42-							β-ACTIN
72-							LAMIN b1
31-							hnRNPA2B1
31-							SRSF1

cytoplasmic | soluble nuclear | insoluble nuclear : Fraction

d

MKN45 | MKN45-*IQGAP1^KO*

51M1 51ΔM | 51M1 51ΔM

% inclusion of exon 2

p=0,0009 ***
p=0,0054 **

51M1 51ΔM | 51M1 51ΔM
MKN45 | MKN45-*IQGAP1^KO*

Fig. 4

Fig. 4: Nuclear IQGAP1-containing RNPs are involved in splicing regulation. a Network of protein interactions generated from the proteins that were pulled down by anti-IQGAP1 from nuclear extracts of NUGC4 cells classified as spliceosomal components. The network was generated using the igraph R package. Colours represent classes of spliceosomal components according to SpliceosomeDB[28]. Vertices are scaled according to *P* values and ordered according to known spliceosomal complexes. **b** Validation of representative IQGAP1-interacting partners presented in (**a**). Anti-IQGAP1 or control IgG pull down from nuclear extracts of NUGC4 and MKN45 cells were immunoprobed against IQGAP1, hnRNPA1, hnRNPA2/B1, hnRNPC1/C2, hnRNPL and DHX9/RHA. The immunoprecipitated proteins were compared to 1/70th of the input used in the pull down. Where indicated, RNase A was added in the pull down for 30 min. Numbers indicate MW in kDa. **c** Subcellular distribution of hnRNPM and IQGAP1. MKN45 and NUGC4 cells were subjected to subcellular fractionation as described in Methods. Equal amounts of the resulting cytosolic and soluble nuclear fractions as well as the insoluble nuclear material were analysed by Western blot for the presence of IQGAP1, hnRNPM, hnRNPA2/B1 and SRSF1. LAMIN b1 and β-ACTIN were used as fractionation and loading controls, respectively. Numbers indicate MW in kDa. **d** MKN45 and MKN45-*IQGAP1^KO* cells were transfected with the DUP51M1 or DUP51ΔM1 minigene splicing reporters[31] for 40 hrs. Exon 2 (grey box) splicing was assessed by RT-PCR using primers located at the flanking exons. Quantification of exon 2 inclusion was performed using ImageJ. Data shown represent the average exon 2 inclusion values ± SD from at least 3 independent experiments. *P* values were calculated using unpaired, two-tailed, unequal variance Student´s t-test.

Fig 5

Fig. 5: A broad role of IQGAP1 in hnRNPM-dependent alternative splicing. a Scatterplot with the distribution of the Psi values for the AS events detected by VAST-TOOLS in RNA-seq in IQGAP1^{KO} and control cells. In yellow are the significantly changed AS events between MKN45 and MKN45-IQGAP1^{KO} cells (|ΔPsi|>15, range 5), in ochre and orange are events with detected iClip binding for hnRNPM or predicted RNA-binding motif, respectively. The gene names of the events that were selected for validation are indicated. The ANAPC10 event that was selected for further analysis is shown in bold. BG: background. **b** Pie chart presenting the frequency of the different types of AS events (exon skipping, intron retention, alternative splice donor and alternative splice acceptor) regulated by IQGAP1 in MKN45 cells. **c** Classification of the up-regulated (Up) and down-regulated (Down) exons in MKN45-IQGAP1KO cells compared to MKN45, as i. having the hnRNPM consensus binding motif downstream of the regulated exon, ii. having been identified as iCLIP targets, iii. belonging to both (i.) and (ii.) and iv. not belonging in any of the above categories. **d** Schematic showing the position of the hnRNPM binding motif enrichment relative to the alternatively spliced exon (grey box). The reported hnRNPM motifs[13] were identified only down-stream of the down-regulated exons. **e** IQGAP1^{KO} affects AS of ANAPC10 in MKN45 cells, by promoting exon 4 skipping. RT-PCR (see **Supplementary Table 4**) followed by electrophoresis were used to monitor the rate of *ANAPC10* exon 4 inclusion in MKN45 and MKN45-IQGAP1^{KO} cells. Exon 4 inclusion was quantified with ImageJ in at least 3 biological replicates. *P* value was calculated with unpaired t-test.

Fig 6

Fig. 6: IQGAP1 and hnRNPM co-operatively regulate the function of APC/C through AS of the ANAPC10 pre-mRNA. a *ANAPC10* exon 4 inclusion was analysed in MKN45 and MKN45-IQGAP1^{KO} cells transfected with siRNAs for hnRNPM or scrambled (scr) control siRNAs. RT-PCR (see **Supplementary Table 4**) followed by electrophoresis were used to monitor the rate of ANAPC10 exon 4 inclusion. Exon 4 inclusion was quantified with ImageJ in at least 3 biological replicates. P value was calculated with unpaired t-test. **b** Immunoblot analysis to monitor the steady-state levels of TPX2 (2 isoforms), RRM2 and TK1 in crude protein extracts from MKN45 and MKN45-*IQGAP1*^{KO} cells transfected with siRNAs for hnRNPM or scrambled control (scr) as indicated. *IQGAP1*^{KO} and the efficiency of hnRNPM down-regulation were also confirmed by immunoblotting. β-ACTIN was used as loading control for the TK1 immunoblot and GAPDH was the loading control for all other immunoblots shown. Numbers indicate MW in kDa. **c-d** Non-synchronized cells from all four cell groups were stained for β-tubulin and DAPI, to visualize the cell cytoplasm and nucleus, respectively. A representative confocal image of the double MKN45-*IQGAP1*^{KO}-*hnRNPM*^{KO} stained cells in (**c**) shows the pronounced multi-nucleation observed in these cells. Quantification of the percentage of cells having 1x, 2x or >2x nuclei (**d**) was performed in 20 images from each cell line, reaching a minimum number of 250 cells analysed per group.

Fig. 7

Fig. 7: IQGAP1 alters hnRNPM's ability to respond to heat-shock by changing its localization relative to SR proteins. a Single confocal planes of MKN45 and MKN45-*IQGAP1^{KO}* cells stained for hnRNPM, IQGAP1 and DAPI after heat-shock (HS) stress induction for 1 hour at 42°C. hnRNPM signal alone is shown in grey for better visualisation and merged images with all three coloured signals are shown on the side. **b** Representative stacks from confocal images of MKN45 and MKN45-*IQGAP1^{KO}* cells stained for hnRNPM and SR proteins. For each condition the single and merged signals of the 2 proteins are shown on top. A single cell stained for hnRNPM and SR is shown on the bottom together with the plot profile line drawn in Image J, while the accompanying pixel grey value graphs are visible on the right of the image. **c** Histogram showing the Pearson's coefficient values for hnRNPM and SR co-localisation, for MKN45 and MKN45-*IQGAP1^{KO}* cells before and after heat-shock stress induction for 1h at 42°C. Pixel-based co-localisation was performed in 36 cells for each of the MKN45 and MKN45-*IQGAP1^{KO}* cells, untreated of after HS, and data represent mean values ± SD. *P* values were calculated using ANOVA multiple comparisons tests; ****$p < 0.0001$.

Fig 8

Fig. 8: IQGAP1 alters hnRNPM's ability to respond to heat-shock by changing its localization relative to SFPQ. a Immunoblot of nuclear matrix extracts from MKN45 and MKN45-*IQGAP1^{KO}* cells before and after heat-shock stress induction for 1hr at 42^0C. β-ACTIN is used as a loading control[69]. Quantification of the relevant amounts, in arbitrary units, was performed using ImageLab software version 5.2 (Bio-Rad Laboratories). **b** Pearson's coefficient values for hnRNMP and PSF co-localisation, for MKN45 and MKN45-*IQGAP1^{KO}* cells before and after heat-shock stress induction for 1h at 42°C (exemplar images of cells analysed are shown in c). Pixel-based co-localisation was performed in 30 cells for each of the MKN45 and MKN45-*IQGAP1^{KO}* cells, untreated of after HS, and data represent mean values ± SD. *P* values were calculated using ANOVA multiple comparisons tests; ****p < 0.0001. **c** Representative confocal planes of MKN45 and MKN45-*IQGAP1^{KO}* cells stained for hnRNPM and PSF. For each cell type and condition both the single and merged signals of the 2 proteins are shown on top. A slice from a single cell stained for hnRNPM and PSF is visible on the bottom together with the plot profile line drawn in Image J, while the accompanying pixel grey value graphs are shown on the right of the image.

Figure 9

Fig. 9: **IQGAP1 regulates hnRNPM-mediated alternative splicing and its response to heat-induced stress.** In gastric cancer cells, IQGAP1 (IQ) interacts with hnRNPM (M) and other alternative splicing regulators, such as SR proteins (AS-SR) to promote exon 4 skipping of the *ANAPC10* pre-mRNA. This leads to the formation of a cell cycle-promoting isoform of ANAPC10 that efficiently targets RRM2, TK1, TPX2 and ANLN for degradation. Upon inactivation of IQGAP1, hnRNPM maintains its ability to bind to its pre-mRNA target, but translocates to the nuclear matrix (NM), thus dissociating from the spliceosomal machinery. This favours the formation of a cancer impeding ANAPC10 isoform that recognizes less efficiently RRM2, TK1, TPX2 and ANLN, thus casing their accumulation, cell cycle delay and cell division defects (multinucleation). IQGAP1 responds to heat-induced stress (HS) by translocating to the nucleus, and is required for redistribution of hnRNPM to the nuclear matrix and its dissociation from the spliceosomal machinery.