1 A Darwinian Selection Based Explanation of Peacock's Tail

2 Dinkar Wadhwa*

3 *The author is currently unaffiliated.

Abstract

There is no satisfactory explanation for why peacock possesses a tail, presence and especially courtship display of which makes the organism vulnerable to predation. Here, I present a model according to which in a polygynous mating system a mechanism which increases vulnerability to predation, a Zahavian handicap, evolves when other two mechanisms to identify high-quality males are either absent or are not sufficiently strong. The two mechanisms are: 1) male resource acquisition ability, and 2) male-male competition for females. The three mechanisms are not necessarily mutually exclusive. It is shown through stochastic simulation that sexual selection, mediated by the tail (a Zahavian handicap), leads to higher rate of increase in the quality of the population of tailed peacocks and tailed-choosy peahens (which exclusively mate with tailed peacocks) as compared to the population of tailless peacocks and tailless-choosy peahens (which exclusively mate with tailless peacocks), through a positive feedback, as daughters of tailed-choosy peahens are of higher average quality and, by virtue of not carrying the tail's handicap, also fitness than daughters of tailless-choosy peahens. Also, the fold-change in the population of tailed peacocks and tailless-choosy peahens are higher than the fold-change in the population of tailless peacocks and tailless-choosy peahens, for all combinations of the initial conditions. The same results were got, though in milder form, when tailless-choosy peahens were replaced by undiscriminating peahens (which mate with tailless and tailed peacocks in proportion to their frequencies in the population). Although sons of tailed-choosy peahens have lower average fitness than sons of undiscriminating peahens, this difference is inconsequential, because in a polygynous mating system a single male can potentially mate with every female. In peacock, increased vulnerability to predation because of tail exists throughout its life, but the model only demands the presence of heightened vulnerability to predation during mate selection process. The work presented here reconciles Zahavi's handicap principle with Darwin's theories of natural and sexual selection. Further, it is hypothesized that a genotype responsible for producing in males a reliable indicator of high quality (a Zahavian handicap) or paternal care ability generates mating desirability in females toward males possessing the indicator. It

32 is demonstrated through simulation that this cross-gender pleiotropy expedites the evolution of

33 tailed peacocks and tailed-choosy peahens and leads to higher rate of increase in their quality.

34 **Keywords**: Peacock's tail, Zahavi's handicap principle, sexual selection, cross-gender pleiotropy

1.   Introduction

In many species, males possess characteristics such as antlers, ornate tail, and colouration,

and behaviours such as courtship display and song which reduce their fitness and thus cannot

be explained by the theory of natural selection. In order to explain the presence of such

characteristics, Darwin [1] proposed the theory of sexual selection, according to which the

lowered fitness of males possessing such characteristics gets compensated by their being more

attractive to females and thus producing more and/or superiour quality progeny than those

males which lack them. But the sexual selection theory did not explain what advantage do

progeny of females which choose males possessing such characteristics have over progeny of

those females which choose other males, so that the evolution of the female choosiness itself

could be accounted for.

According to Fisher [2], initially the male characteristic and male quality were correlated;

thus, those females which chose males possessing the characteristic left more descendants

than those females which chose males without the characteristic. Thus, in the subsequent

generations, males had evolved heightened form of the characteristic and females had evolved

higher attractiveness for it. Then a point arrived when further increase in the characteristic's

size had no correlation with male fitness, but still the increase in the characteristic's size and

female attractiveness for it continued because of the selective pressure created by the latter.

In other words, males possessing the characteristic had more and/or superiour quality progeny

simply because of the female attractiveness for the characteristic. A few questions can be raised

on Fisher's argument: a) in what way the characteristic was correlated with male quality?, b)

how that correlation came to an end?, and c) what factor compensates for the cost of the extra

size of the characteristic?

For a characteristic to undergo positive selection in a polygynous mating system, it should

increase fitness of at least female offspring in terms of them getting good genes and/or better

parental care. The characteristic may decrease fitness of male offspring to the extent that

only few males survive. The lower survivability of males is inconsequential because in a

polygynous mating system a single male can potentially mate with every female. Natural

selection can be looked at as a feedback mechanism. Genes that increase fitness ("good genes")

enable the organism to leave more descendants, hence higher presence of the good genes in

the subsequent generation. Genes that decrease fitness ("bad genes") cause the organism to

leave fewer descendants, hence lower presence of the bad genes in the subsequent generation.

Sexual selection for a characteristic can only escape this loop if the characteristic has no effect

on fitness. Thus, Fisher's argument is tantamount to assuming that the characteristics have no

effect on fitness, which is clearly not true.

In 1975, Amotz Zahavi [3] proposed the handicap hypothesis to explain the evolution of the

male characteristics and behaviours which, although reduce their fitness, increase their mating

success. According to Zahavi, the presence of such fitness-reducing traits are tests, as only

high quality males can survive despite carrying their burden. Thus, the presence of such traits

reliably signals male quality, for males not possessing such traits may or may not be of high

quality. It has been demonstrated that offspring of peacocks with more elaborate train grow and

survive better under experimental conditions approximately the same as natural conditions [4].

In 1976, Maynard Smith [5] analyzed the handicap hypothesis and concluded that the proposed

mechanism does not work, even when the handicap is absent in females. In 1978, Bell [6]

carried out a more extensive study of the handicap principle, considering haploid population in

monogamous and polygynous mating system. The study found out that the principle can work

in a certain condition related to the initial frequency of choosy females. The results of Bell's

paper are discussed below. A study by Davis and O'Donald [7] concluded that Zahavi's model

cannot work; but because the analysis in the paper is beyond the scope of the present work,

results of that work will not be discussed here.

2.   Results and Discussion

2.1.   Fundamental biological functions

In a naive consideration, life can be thought of as an organism's struggle to leave as many

descendants as possible. This, then, demands the organism to succeed in performing at least the

following functions: 1) acquire and protect resources, 2) acquire mate, and 3) protect oneself

and progeny. For the male gender, the function of mate protection may also be added. In

many cases, the female gender has the additional function of gestation to perform. Because

resource protection is needed only when resources have been acquired, the function of resource

protection may be subsumed into the function of resource acquisition. Further, the functions

of protecting progeny and mate (in case mate protection is present) may be subsumed into

the need to protect oneself. Arguably, the function of self-protection should also include

protecting oneself from parasites and pathogens, because abstractly, parasites and pathogens

can be considered equivalent to predators. Also, gestation may be subsumed into resource

acquisition. Thus, we are left with three fundamental biological functions, namely resource

acquisition, mate acquisition, and self-protection.

In any mating system, female cannot possess characteristics as indicators of quality that reduce

their fitness, because reproduction primarily depends on them. In a given species, females

which display their quality by a fitness-reducing feature will be outnumbered by females which

don't, because the latter are more likely to survive until the reproductive age and could invest

advantage due to higher fitness in reproduction (higher quality eggs or better gestation) and

parental care. In monogamous mating system, because each male can mate with only one

female, male should not indicate their quality by characteristics which reduce fitness. In

polygynous mating system, however, because a high-quality male can mate with several females, characteristics that reduce fitness can serve as indicator of quality. This is because a few high-quality males may still survive, and consequently all females can get to mate. Because the present work seeks to explain the evolution of peacock's tail, the focus will be on how females determine male quality. Thus, a mechanism, or test, needs to exist by means of which females can determine male quality. Naturally, fundamental biological functions of males can serve as the tests. However, because male's resource acquisition ability may also contribute to his mate acquisition ability, an alternate formulation may be better conceptually. Thus, because the process of mate acquisition consists of a male displaying his ability to acquire resources and/or fighting off rival males, it may be better to replace "resource and mate acquisition" with "resource acquisition and male-male competition".

a) Resource acquisition: Males may demonstrate their resource acquisition ability in three ways. First, by stockpiling resources. Second, by possessing high-investment characteristics, such as antlers, which are shed and regrown annually. Only those males which are good at resource acquisition will be able to grow and maintain antlers. Third, by building an architecture, e.g. nest or bower. Although vast majority of bird species are monogamous, a bird species may be imagined in which males build many nests in order to attract multiple mates. b) Male-male competition: The male which defeats rival males in a physical confrontation demonstrates his high quality. c) Self-protection: This test is implemented by way of males possessing a characteristic or displaying a behaviour which increases their vulnerability to predation. The examples are the peacock's tail plumes, bright colouration of many bird species, and risky courtship displays and songs. Males which are able to survive in spite of being more vulnerable to predation are of higher quality than those which are not. It should be noted that the presence of increased vulnerability to predation post-mating will hamper paternal care in case it is present. A hypothetical phenomenon can be considered in which males, as part of courtship ritual, put them in a situation which makes them highly vulnerable to parasitic/pathogenic infection. Only high-quality males (i.e., having strong immune system) will be able to survive

such ritual. The three functions can also be present in combination. For example, a) antlers are
used as weapons in male-male combat for mate acquisition, and b) male-male fighting may also
be involved in resource acquisition directly or indirectly, that is, in competition for territorial
possession. In the peacock, because the first two mechanisms for determining male quality are
absent or weakly present, males are tested on their ability to protect themselves. Therefore,
a characteristic, the tail, is evolved in males, which makes them more vulnerable to predation
than females.

Figure 1: Fundamental biological functions for males in a polygynous mating system.

148 **2.2.   The evolution of peacock's tail and peahen's choosiness**

A haploid population is considered which has i) an autosomal locus for quality, ii) male-specific
locus for tail, and iii) female-specific locus which determines the peahen type. The three loci
have two alleles. The first locus has Q (whose values are 1.1, 1.2, 1.5, 2, 5, 10) and q (whose
value is 1). The second locus has T (tailed peacock) and t (tailless peacock). For the third
locus two cases are considered. First in which the two alleles are tailed-choosy, peahens which
mate only with tailed peacocks, and tailless-choosy, peahens which mate only with tailless
peacocks. Second in which the two alleles are tailed-choosy, peahens which mate only with
tailed peacocks, and undiscriminating, peahens which mate with both types of peacocks in
proportion to their frequencies in the population. In the present model, an additive model of
fitness is employed where fitness is equal to quality minus test (the tail's burden is set equal to
q). Because peahens do not have tail, their fitness is equal to their quality, Q. Table 1 shows the
equivalence of this model with the additive fitness model of Bell's paper [6]. The simulation

was started with initial population of 10^5 male and female offspring each. The initial frequency

of the allele Q was taken to be 0.01, 0.05, and 0.1. For simplicity, the initial frequency of

tailed peacocks and tailed-choosy peahens were considered to be equal, and were taken to be

0.01, 0.05, and 0.1. A tailed peacock having the allele Q reached the reproductive age with the

probability $(Q-q)/(Q+q)$, whereas a tailed peacock with the allele q reached the reproductive

age with the probability $(q-q)/(Q+q)$; this means that tailed peacocks with the allele q died

before being able to reproduce. A tailless peacock or a peahen having the allele Q reached the

reproductive age with the probability $Q/(Q+q)$, whereas a tailless peacock or s peahen with the

allele q reached the reproductive age with the probability $q/(Q+q)$.

For a given value of the fitness differential, the initial frequency of the allele Q, and the initial

frequency of tailed peacocks, simulation was carried out for the number of generations at which

the value of the below-stated quantities stabilized, because of the fixation of the allele Q by

natural selection. The quantities are: 1) the ratio of the fold-change in the population of

tailed peacocks to the fold-change in the population of tailless peacocks, 2) the ratio of the

fold-change in the population of tailed-choosy peahens to the fold-change in the population of

tailless-choosy or undiscriminating peahens, 3) the ratio of the fold-change in the quality of

the population of tailed peacocks to the fold-change in the quality of the population of tailless

peacocks, and 4) the ratio of the fold-change in the quality of the population of tailed-choosy

peahens to the fold-change in the quality of the population of tailless-choosy or undiscriminating

peahens. The quality of a population of peafowl is equal to the number of peafowl having the

allele Q multiplied by the fitness value of the allele Q plus the number of peafowl having the

allele q multiplied by the fitness value of the allele q. In other words, the quality of a peafowl

population = (Number of Q peafowls)·Q + (Number of q peafowls)·q.

As argued above, increased vulnerability to predation in peacocks due to the tail enables

tailed-choosy peahens to mate with high-quality peacocks, thereby expediting the increase

in the quality of the population of tailed peacocks and tailed-choosy peahens. To test this

hypothesis, two endogamous populations were considered. The first consisted of tailed peacocks

and tailed-choosy peahens, which mate only with tailed peacocks. The second consisted of

tailless peacocks and tailless-choosy peahens, which mate only with tailless peacocks. Thus,

the second population serves as the control, for natural selection is the only evolutionary force

acting upon it. The ratio of the fold-change in the quality of the population of tailed peacocks

to the population of tailless peacocks was interpreted as the rate of increase in the quality of

peacocks due to sexual selection mediated by the tail. The ratio of the fold-change in the

quality of the population of tailed-choosy peahens to the population of tailless-choosy peahens

was interpreted as the rate of increase in the quality of peahens due to sexual selection mediated

by the tail. The ratio of the fold-change in the population of tailed peacocks to the population

of tailless peacocks was interpreted as the rate of the evolution of the tail. The ratio of the

fold-change in the population of tailed-choosy peahens to the population of tailless-choosy

peahens was interpreted as the rate of the evolution of the female choosiness.

As figure 2 shows, for every combination of the initial conditions all the four fold-change ratios

are more than one, demonstrating that the Zahavian handicap leads to higher rate of increase

in the quality of peacocks and peahens and expedites the evolution of tailed peacocks and

tailed-choosy peahens. The explanation of these results are as follows. The average quality

of tailed peacocks which survive to reproduce despite heightened vulnerability to predation is

higher than the average quality of tailless peacocks. Therefore, the average fitness of daughters

of tailed-choosy peahens is higher than the average fitness of daughters of tailless-choosy

peahens. Consequently, more daughters of tailed-choosy peahens survive until the reproductive

age as compared to daughters of tailless-choosy, thereby amounting to higher percentage of

tailed-choosy peahens in the subsequent generation, and so on. In honour of Amotz Zahavi,

I name this positive feedback loop the Zahavian runaway. Alternately, from the perspective

of higher survivability of daughters of tailed-choosy peahens, the mechanism proposed here

may also be called the fitter-daughter principle. Because these daughters will produce tailed

peacocks, the average quality of tailed peacocks also will be higher than the average quality of

tailless peacocks, and the percentage of tailed peacocks will also increase with each generation.

It should be noted, however, that the peacock percentage is not as crucial a factor as the peahen

percentage because the mating system is polygynous, where a single male can potentially mate

with every female. The fixation of the allele Q and consequent stabilization of the fold-change

ratios does not create a conceptual problem, because the struggle between prey and predator

is perpetual. In other words, the prey needs to evolve continuously in order to keep up with

the predator's evolution, and vice-versa. Richard Dawkins reasoned that according to Zahavi's

argument, selection should favour 'the evolution of males with only one leg and only one eye'

[8]. In the light of the work presented here, the reply is that selection could have favoured the

evolution of males with only one leg and only one eye if the genetic mechanism had been such

that the defect was phenotypically restricted to males.

Notably, all the fold-change ratios decrease as the fitness differential increases. This happens

because with higher fitness differential, effect of natural selection becomes stronger (higher

percentage of tailless peacocks with the allele Q survives), thereby diminishing the role of

the handicap principle, which is mediated through sexual selection. The four fold-change

ratios also decrease as the initial frequency of the allele Q increases. The explanation for

this trend is as follows. The average quality of tailed peacocks is higher than that of tailless

peacocks because tailed peacocks having the allele q do not survive to reproduce, whereas

tailless peacocks with the same allele do. With higher initial frequency of the allele Q, fewer

tailless peacocks with the allele q are present, thereby reducing the advantage due to the tail.

The same analysis was carried out by replacing tailless-choosy peahens with undiscriminating

peahens, which mate with tailed and tailless peacocks in proportion to their frequencies in

the population. As previously, for every combination of the initial conditions all the four

fold-change ratios are more than one, though there are a few differences. For all combinations

of the initial conditions, the ratio of the fold-change in the population of tailed-choosy peahens

to the population of undiscriminating peahens is lower than the ratio of the fold-change in

the population of tailed-choosy peahens to the population of tailless-choosy peahens. The

explanation for this is as follows. Because the average quality of tailed peacocks is higher than

the average quality of tailless peacocks, the average quality (and hence also average fitness)

of daughters of undiscriminating peahens is higher than the average quality of daughters of

tailless-choosy peahens. Consequently, more daughters of undiscriminating peahens survive

to reproduce, which lowers the fold-change ratio. Further, as the initial fraction of tailed

peacocks increases, the ratio of the fold-change in the population of tailed-choosy peahens

to the population of tailless-choosy peahens remains constant, but the ratio of the fold-change in

the population of tailed-choosy peahens to the population of undiscriminating peahens decreases.

This is because the likelihood of mating between undiscriminating peahens and tailed peacocks

goes up as the initial fraction of tailed peacocks increases.

For lower values of the fitness differential, the ratio of the fold-change in the population of tailed

peacocks to the population of tailless peacocks is lower for the case where undiscriminating

peahens are present as compared to the case where tailless-choosy peahens are present. For

higher values of the fitness differential, the reverse is observed. The reason for this result

could be the following. As the initial fraction of tailed peacocks increases more mating events

occur between tailed peacocks and undiscriminating peahens. This has two opposing effects

with respect to the population growth of tailless peacocks: 1) it reduces the chances of mating

between tailless peacocks and undiscriminating peahens, which produces tailless male offspring,

and 2) it increases the average fitness of tailless peacocks, as they receive the higher quality

allele, Q, from tailed peacocks via undiscriminating peacocks. When the fitness differential is

high, the former effect dominates because the quality of tailless peacocks increases much more

due to natural selection, whose intensity is higher due to large fitness differential, than due

to the coupling between tailless peacocks and tailed peacocks via undiscriminating peahens,

and vice-versa. The reasoning may also explain the result that although the fold-change in

the population of undiscriminating peahens increases for all fitness differentials as the initial

fraction of tailed peacocks goes up, the fold-change in the population of tailless peacocks, as the

initial fraction of tailed peacocks goes up, decreases only when the fitness differential is equal

to 2. However, for all fitness differentials, the fold-change in the population of tailed peacocks

does not increase as the initial fraction of tailed peacock goes up. This was unexpected because

as the initial fraction of tailed peacock increases, more mating events occur between tailed

peacocks and undiscriminating peahens. It seems that this discrepancy might have something to

do with very high variance of the fold-change in the population of tailed peacocks. The variance

decreased as the initial frequency of tailed peacocks increased. For the initial frequency of

tailed peacocks equal to 0.01, the order of magnitude of the variance was the same as that of the

mean, whereas for the initial frequency equal to 0.1, the variance was one order of magnitude

less than that of the mean. The same trend was observed for the fold-change in the population

of tailed peacocks for the undiscriminating peahens case and also for tailed-choosy peahens in

both cases. In any case, because the ratio of the fold-change in the population of tailed-choosy

peahens to the population of undiscriminating peahens is more than one, the evolution in this

mating scenario also occurs through Zahavian runaway.

The study by Bell [6] showed that the handicap principle can work within a certain range of

the initial frequency of choosy females. Specifically, for a given value of the fitness differential

(which is equal to Q-q in the present work), there will be a value of the initial frequency

of tailed-choosy females, below which the handicap principle cannot not work. However,

according to the mechanism proposed here, whatever may the initial frequency of tailed-choosy

females be, the handicap principle will always work. That's because the average fitness of

daughters of tailed-choosy females will always be greater than the average fitness of daughters

of undiscriminating females. This leads to higher percentage of tailed-choosy peahens, and

therefore lower percentage of undiscriminating peahens, in the subsequent generation.

A study [9] showed that peacocks which exhibited frequent behavioural display and had a large

number of tail eyespots had better immune system. Peacocks which displayed frequently put

themselves at predation risk more often than those which displayed less frequently. Arguably,

Figure 2: (In the first column, the initial fraction of tailed peacocks and tailed-choosy peahens (f) is equal to 0.01, in the second column f=0.05, and in the third f=0.1. **A**) The ratio of the fold-change in the population of tailed peacocks to the population of tailless peacocks. (**B**) The ratio of the fold-change in the population of tailed-choosy peahens to the population of tailless-choosy peahens. (**C**) The ratio of the fold-change in the quality of the population of tailed peacocks to the quality of the population of tailless peacocks. (**D**) The ratio of the fold-change in the quality of the population of tailed-choosy peahens to the quality of the population of tailless-choosy peahens.

Table 1: Equivalence of Bell's additive fitness model [6] and the model in the present work

Allele	Tailed Peacock	Tailless Peacock	Peahen
B	$\alpha + \beta$	$\alpha + \beta + \epsilon$	$\alpha + \beta + \epsilon$
b	α	$\alpha + \epsilon$	$\alpha \epsilon$
Q	Q - q	Q	Q
q	q - q	q	q

The fitness differential β=Q-q; the tail's burden ϵ=q; α=0

Figure 3: (In the first column, the initial fraction of tailed peacocks and tailed-choosy peahens (f) is equal to 0.01, in the second column f=0.05, and in the third f=0.1. **A**) The ratio of the fold-change in the population of tailed peacocks to the population of tailless peacocks. (**B**) The ratio of the fold-change in the population of tailed-choosy peahens to the population of undiscriminating peahens. (**C**) The ratio of the fold-change in the quality of the population of tailed peacocks to the quality of the population of tailless peacocks. (**D**) The ratio of the fold-change in the quality of the population of tailed-choosy peahens to the quality of the population of undiscriminating peahens.

this behaviour should be heritable. A tail eyespot most probably gives a semblance of a face,

and consequently invites more attention from predators. Thus, frequent behavioural display

and more tail eyespots translate to stronger test. Only high quality peacocks are able to pass

this test, and daughters of those peahens which mate with such peacocks have higher fitness.

Overtime, this will lead to a correlation between high quality (such as better immune system)

and frequent behavioural display and more tail eyespots. This result may seem to support the

Hamilton-Zuk hypothesis, but an alternative, more general explanation is provided below.

309 2.3. The Hamilton-Zuk hypothesis

According to the Hamilton-Zuk hypothesis, male secondary sexual characteristics signal resistance

to parasites and pathogens [10]. Recently, a comparative genome analysis of Indian peacock

[11] reported that the immune system genes of the bird displayed signs of adaptive evolution.

The authors observed this result of being supportive of the Hamilton-Zuk hypothesis. However,

a more general reasoning can account for the evolution of the immune system genes in peacock

[11] and the correlation reported between better immune system and higher frequency of

behavioural display and large number of tail eyespots [9], if infection by parasites or pathogens

are considered to be a test. A peacock with weak immune system and carrying an infection

will sluggishly respond to a predatory attack and consequently will be less likely to survive.

This will lead to selection of higher quality immune system. Further, all faculties involved

in evading predation, such as the sensory systems and muscles and bones involved in taking

wing and flying, will undergo selection for improvement. This may explain the findings of

the comparative genome analysis [11] that genes involved in bone morphogenesis and skeletal

muscle development showed signs of adaptive evolution.

2.4.   Cross-gender pleiotropy

The analysis above is based on the assumption that the locus for the tail is male-specific and the locus which determines peahen type (tailed-choosy, tailless-choosy, and undiscriminating) is female-specific. However, because an ovariectomized peahen develops peacock-type plumage [12], the locus for the tail has to be autosomal. This leads us to the question: what could be a purpose of the tail's locus being autosomal when normal peahens do not have the tail? I hypothesize that genotype for the tail in peahens contributes, partially or fully, to their desirability to mate with a tailed peacock. Generalizing, a genotype responsible for producing a Zahavian handicap in males generates mating desirability in females toward males possessing the Zahavian handicap. Because, besides good genes, parental care can also increase fitness of the progeny, the hypothesis can be extended. A genotype responsible for producing in males a reliable indicator of high quality (a Zahavian handicap) or paternal care ability generates mating desirability in females toward males possessing the indicator.

In the earlier simulation, all daughters of a mating between a tailed peacock and an undiscriminating peahens were undiscriminating peahens. By contrast, when genotypes for the tail and female choosiness are assumed to be the same, half of the daughters from mating between a tailed peacock and an undiscriminating peahen are tailed-choosy. Although, half of the sons also are tailless, but that is immaterial in a polygynous mating system. Thus, when genotype for the tail and female choosiness are identical, the evolution of tailed peacocks and tailed-choosy peahens should be faster. Also, the rate of increase in the quality of tailed peacock and tailed-choosy peahens should be higher. If these predictions are correct, then the four fold-change ratios of the case where genotype for the tail and female choosiness are identical should be higher than the respective four fold-change ratios of the case where one fraction of peahens are tailed-choosy while another fraction is undiscriminating, for all combinations of the initial conditions. Therefore, the ratio of the two ratios should be greater than one. The simulation could not be carried out for the number of generations at which the four fold-change ratio

stabilize, because of the lack of computing power. But even for lower number of generations, the prediction turns out to be true, though the result are not presentable.

References

[1] Darwin C The Descent of Man and Selection in Relation to Sex. 2nd edn. London: John Murray.

[2] Fisher RA The Genetical Theory of Natural Selection 1930; Oxford: Clarendon Press

[3] Zahavi A Mate Selection-A Selection for a Handicap J. theor. Biol. 1975; 53 205–214

[4] Petrie M Improved growth and survival of offspring of peacocks with more elaborate trains Nature 1994; 371 598–599

[5] Maynard Smith J Sexual Selection and the Handicap Principle J. theor. Biol. 1976; 57 239–242

[6] Bell G The Handicap Principle in Sexual Selection Evolution 1978; 32(4) 872–885

[7] Davis JWF and O'Donald P Sexual Selection for a Handicap : A Critical Analysis of Zahavi's Model

[8] Dawkins R The Selfish Gene 1976 (p. 172) Oxford University Press, Oxford.

[9] Loyau et al. Multiple sexual advertisements honestly reflect health status in peacocks (Pavo cristatus) 2005 Behavioral Ecology and Sociobiology 58(6) 552–557

[10] Hamilton WD and Zuk M Heritable true fitness and bright birds: a role for parasites? 1982 Science 218 384–387

[11] Jaiswal SK et al Genome Sequence of Peacock Reveals the Peculiar Case of a Glittering Bird 2018 Frontiers in Genetics 9, 392

374 [12] Owens IP and Short RV Hormonal basis of sexual dimorphism in birds: implications for

375   new theories of sexual selection 1995 Trends Ecol. Evol. 10 44–47