GROEL/ES BUFFERS ENTROPIC TRAPS IN FOLDING PATHWAY DURING EVOLUTION OF A MODEL SUBSTRATE

Anwar Sadat[1,3,#], Satyam Tiwari[1,3,#], Kanika Verma[1,3], Arjun Ray[2], Mudassar Ali[5], Vaibhav Upadhyay[4], Anupam Singh[3], Aseem Chaphalkar[1,3], Asmita Ghosh[1,3], Rahul Chakraborty[1,3], Kausik Chakraborty[1,3,$], Koyeli Mapa[1,5,$,≠]

[1] Academy of Scientific and Innovative Research, CSIR-HRDG, Ghaziabad, Uttar Pradesh 201002, India.

[2] Indraprastha Institute of Information Technology-Delhi, Okhla Industrial Estate, Phase III, New Delhi 110020, India.

[3] CSIR-Institute of Genomics and Integrative Biology, Mathura Road, New Delhi 110025, India.

[4] National Institute of Immunology, Aruna Asaf Ali Marg, New Delhi 110067, India.

[5] Department of Life Sciences, School of Natural Sciences, Shiv Nadar University, NH91, Greater Noida, Gautam Buddha Nagar, Uttar Pradesh 201314, India.

[#] Equally contributing first authors

[$] Equally contributing senior authors

[≠] Correspondence: koyeli.mapa@snu.edu.in

ABSTRACT

The folding landscape of proteins can change during evolution with the accumulation of

mutations that may introduce entropic or enthalpic barriers in the protein folding pathway,

making it a possible substrate of molecular chaperones *in vivo*. Can the nature of such

physical barriers of folding dictate the feasibility of chaperone-assistance? To address

this, we have simulated the evolutionary step to chaperone-dependence keeping

GroEL/ES as the target chaperone and GFP as a model protein in an unbiased screen.

We find that the mutation conferring GroEL/ES dependence *in vivo* and *in vitro* encode

an entropic trap in the folding pathway rescued by the chaperonin. Additionally, GroEL/ES

can edit the formation of non-native contacts similar to DnaK/J/E machinery. However,

this capability is not utilized by the substrates *in vivo*. As a consequence, GroEL/ES caters

to buffer mutations that predominantly cause entropic traps, despite possessing the

capacity to edit both enthalpic and entropic traps in the folding pathway of the substrate

protein.

INTRODUCTION

Mutations in a protein sequence may subtly change either thermodynamics of the folding polypeptide or protein-solvent interactions. *In vivo*, mutations that arise spontaneously may lead to problems in the folding pathway or stability of proteins. This, in turn, may make the proteins either non-functional or dependent on molecular chaperones for folding (Chiti and Dobson, 2017; Hartl, 2017)

In *E. coli*, the most abundant cytosolic chaperone systems consist of the Hsp70 system (DnaK/DnaJ/GrpE), the Hsp60 system (GroEL/GroES), Hsp90 (HtpG), Trigger Factor (Tig) and SecB along with other less abundant chaperones and small heat shock proteins. Properties of the substrates assisted by these chaperone systems have been explored by multiple groups (Calloni et al., 2012; Dunn et al., 2001; Houry et al., 1999; Kerner et al., 2005; Knoblauch et al., 1999; Rüdiger et al., 1997). While the DnaK system binds to many proteins and has the potential to stabilize thermosensitive proteome (Zhao et al., 2019), GroEL/ES binds and helps in the folding of a much smaller subset of cellular proteins (Kerner et al., 2005; Niwa et al., 2016). The mechanism of substrate targeting to the right chaperone is an active field of research underlining the significance of the conformations adopted by proteins in their non-folded states (Mapa et al., 2012; Nagpal et al., 2015).

While canonical chaperone-targeting is important for chaperone-dependent wild type proteins, the accumulation of mutations during evolution may create additional substrates requiring chaperone assistance. Can we predict the type of mutations on a GroEL/ES independent protein that would make it GroEL/ES dependent? Mechanistically, some mutations may increase the propensity of formation of non-native contacts (intramolecular or intermolecular) that need to melt before the protein can fold to its native state (enthalpic trap) while other mutations may increase the flexibility of the non-native states (entropic trap), both of which are folding problems at opposite ends of the spectrum. Do different chaperone systems differ in their capacity to edit the two types of mutations?

GroEL/ES system is capable of removing the entropic trap in its substrates (Chakraborty et al., 2010; Georgescauld et al., 2014). It has also been shown to unfold proteins (Sharma et al., 2008) and remove enthalpic traps in the folding pathway *in vitro*. Does the chaperonin possess both these activities *in vivo*? When proteins accumulate mutations, which type of mutations would be preferentially accommodated by GroEL/ES? This is difficult to answer with the previous model substrates as neither the authentic substrates (which precludes understanding the mutational steps that led to its chaperone dependence) nor the slow-folding model substrates (that were not evolved for GroEL/ES dependence *in vivo* in an unbiased manner) gave us the handle to

quantitate folding assistance by GroEL/ES *in vivo*. Although, Horovitz group made advances in

this direction by identifying that mutations in frustrated sequence regions lead to GroEL/ES

dependence *in vivo*, but the biophysical consequence of these mutations on the folding landscape

was not investigated (Bandyopadhyay et al., 2017).

To address this, we sought an unbiased screen using mutagenesis to obtain an *in vivo* GroEL/ES

substrate starting from a spontaneously folding GroEL/ES-independent substrate. We show that

the identified substrate is exclusively dependent upon the GroEL/ES system *in vivo* and *in vitro*.

We find that the unique mutation present in the pool of GroEL/ES dependent protein resulted in

an entropic trap in folding that is corrected by GroEL/ES system. We also show that

DnaK/DnaJ/GrpE (KJE) system or the GroEL chaperone can take care of enthalpic traps

effectively *in vitro* in an ATP dependent manner. This function of GroEL is essential for folding the

substrate protein *in vivo*. Thus, we posit that the proteins that acquire entropic traps during

evolution would be assisted by GroEL/ES system *in vivo*. While GroEL/ES also possesses the

capability to edit folding landscape by preventing the formation of non-native contacts, this

chaperoning capacity is not exclusive to GroEL/ES but is also shared by more abundant KJE

system.

RESULTS

Isolation of a synthetic GroEL-dependent substrate

A single mutational step may make a spontaneously folding protein chaperone-dependent during

evolution. To learn the physico-chemical basis of the mutational step that confers chaperone-

dependence, we wanted to mimic this evolutionary step and develop a synthetic substrate

dependent on the canonical chaperone GroEL/ES *in vivo* from a GroEL/ES-independent protein.

Comparison of the mutant and Wt protein would help in understanding the type of folding problems

that GroEL/ES tends to edit, and the mechanism of its chaperoning action. We chose yeGFP

(yeast enhanced GFP- referred to as Wt GFP hereafter), a fast-folding form of GFP, as the starting

protein as, 1) folding of the Wt protein was independent of GroEL/ES *in vivo* and *in vitro* (Figure

1A), and 2) it was easy to quantify the amount of soluble well-folded protein in single-cells using

flow-cytometry (Verma et al., 2020). Importantly, to normalize for expression levels, plasmid copy

number and induction, we used Wt mCherry in an operonic construct containing Wt GFP followed

by ribosome-binding-site (RBS) and Wt mCherry, under the control of an arabinose-inducible

system (Figure 1B). The expression levels of mCherry were used as a reference to obtain the

relative expression levels of GFP (Verma et al., 2020).

To make a GroEL/ES dependent substrate, we used a random mutant library of GFP, constructed

using the backbone of this GFP-mCherry operonic construct (Verma et al., 2020). The mutant

library showed populations of cells containing varying degrees of GFP fluorescence starting from

very low fluorescence to near-Wt GFP fluorescence (Figure 1C, upper panel). Using

Fluorescence Assisted Cell Sorting (FACS) we first purified a population of clones (Population-

LF) that exhibit extremely low GFP fluorescence compared to Wt GFP (Figure 1C). These clones

would either have problems in folding or have quenched fluorescence due to the alteration of the

fluorophore environment. We co-transformed Population-LF in a K-12 strain of *E. coli* (BW35113,

referred to as WT hereafter) overexpressing GroEL/ES from a plasmid under the control of an

IPTG-inducible *tac* promoter. We isolated mutant clones that exhibited higher GFP fluorescence

upon GroEL/ES overexpression (Figure 1D and S1A). GFP-mCherry plasmids were isolated from

individual clones after sorting and re-transformed to confirm the dependence of GFP fluorescence

on the overexpression levels of GroEL/ES *in vivo*. Ten of the isolated clones were sequenced and

all of them were found to have the mutation K45E; the mutation did not map to any residues

around the GFP fluorophore (Figure 1E). The purified mutant protein had similar *in vitro*

fluorescence as that of Wt GFP (Figure S1B). *In vivo* fluorescence of K45E mutant (hereafter

referred to as slow-folding GFP or sGFP) increased upon co-expression of GroEL/ES (Figure 1F

and S1C). sGFP showed compromised folding and the nascent chains were rapidly cleared *in*

vivo in the absence of GroEL/ES overexpression in WT *E.coli* cells (Figure S1D). The dependence

of sGFP on GroEL/ES for folding was further confirmed with by the drastically enhanced solubility

of the protein along with GroEL/ES overexpression (Figure S1E) in BL21(DE3) cell lacking major

protease systems. This corroborated well with the increased fluorescence and indicated that

sGFP was a folding mutant of Wt GFP that depends upon the GroEL/ES system for folding, *in*

vivo.

Furthermore, to check the folding dependence of sGFP on other chaperone systems, we

measured the *in vivo* fluorescence of sGFP in the presence of DnaK/J/E overexpression (Figure

1G and S1F). The fluorescence of sGFP did not increase rather decreased in the presence of

DnaK/J/E overexpression probably due to the binding of sGFP to DnaK/J/E routing it for

degradation. Additionally, deletion of canonical molecular chaperones *Tig*, *dnaK*, and *dnaJ* did

not significantly decrease sGFP fluorescence suggesting that increase in fluorescence of sGFP

was not routed through *Tig*, *dnaK*, and *dnaJ* (Figure 1H and S1G). In fact, GFP fluorescence

mildly increased in Δ*dnaK,* which is known to overexpress GroEL/ES system (McCarty and

Walker, 1994; Verma et al., 2020). This suggested that sGFP folding was primarily dependent on

the cellular GroEL/ES system. Taken together we found that a single mutation on Wt GFP
conferred sGFP with a stringent GroEL/ES dependence *in vivo.*
To check if K45E conferred GroEL/ES dependence specifically or any marginally active mutant
of GFP would be GroEL/ES substrate, we picked 10 clones randomly from the LF-GFP mutant
pool. Out of these 10 clones 6 clones were finally taken to check their dependence on GroEL/ES
for their folding. The basal level fluorescence of these mutants was extremely low similar to that
of sGFP. Upon overexpression of GroEL/ES with these mutants only three of the six mutants M4,
M5, M6 showed an increase in GFP fluorescence like that of sGFP (Figure 1I and S1H). All the
others showed lower fluorescence indicating that mutations that confer GroEL/ES dependence
are special and are present in ~50% of the mutant pool. That GroEL/ES expression affects a sub-
set of the mutant GFP pool, indicating that GroEL/ES most likely rescues only specific problems
in folding pathways. K45E mutation captured a rare step that would confer GroEL/ES dependence
to a GroEL/ES independent folder such as Wt GFP used in the study.

Refolding of sGFP is limited by a flexibility dependent kinetic trap

To understand the folding barrier introduced by K45E, we characterized the spontaneous
refolding pathway of sGFP. Wt GFP and sGFP were unfolded in 6M GuHCl for one hour, causing
complete unfolding of the protein (Figure S2A) and refolded by a hundred-fold dilution of the
unfolded protein in refolding buffer. We found that sGFP refolded with a much slower rate than
Wt GFP (Figure 2A) and was fit well to single exponential kinetics (Figure S2B) with an apparent
rate of ~$0.2 \times 10^{-3} s^{-1}$. Additionally, we found that the refolding rate and amplitude of sGFP at 25ºC
were independent of the concentration of the protein between 12.5 nM to 400 nM (Figure 2B,
S2C, and S2D). Similarly, the rates and yield were independent of protein concentration even at
37ºC (Figure 2C) suggesting that the protein refolding was not limited by off-pathway aggregation
in the temperatures used for this study. This suggested that the mutation K45E perturbed the
unimolecular rate of folding in sGFP.

To understand the type of folding-barrier incorporated by the K45E mutation in sGFP, we
analyzed the Arrhenius plots for the folding of sGFP and Wt GFP (Figure 2D, Table S1). We fitted
temperature-dependent refolding rates of the proteins using a full model of the Arrhenius equation
containing the $\Delta Cp^{\#}$ term (to account for the difference in heat capacity of the folding intermediate
and the transition state) (Dandage et al., 2015). sGFP did not have a temperature-dependent
slope suggesting the absence of any enthalpic folding barrier ($\Delta H^{\#}$) while Wt GFP had a high $\Delta H^{\#}$
indicating an enthalpic barrier to folding (Figure 2E, Table S1). Since sGFP refolded slower than
Wt GFP, this suggested that the rate-limiting barrier to folding in sGFP is non-enthalpic (Figure
S3A). Complementary to $\Delta H^{\#}$ is the value of ρ that is a composite function of the frequency across

the transition barrier (prefactor for Arrhenius function) and the entropic assistance towards
reaching the transition state (TS) (Figure S3A); lower ρ is associated with higher entropic barrier
(Dandage et al., 2015). ρ was higher for Wt GFP compared to sGFP (Figure 2E, Table S1)
demonstrating that sGFP faced an entropic barrier in folding trajectory. This was consistent with
earlier findings with other GroEL substrates (Chakraborty et al., 2010; Georgescauld et al., 2014).
sGFP was likely trapped in a folding intermediate I_1 that was stabilized by flexible regions that
prevented the formation of native contacts. Analysis of GFP structure revealed that the residue
mutated in sGFP, K45, interacted with multiple negatively charged residues in Wt GFP (E32,
D210, and E213) (Figure 2F). This lysine residue also formed a core for nucleating long-range
interactions in a loop region. The substitution of lysine (K45) to a negatively charged glutamate in
sGFP could have led to the loss of these interactions and introduced repulsive interactions that
would destabilize this region. Spontaneous refolding rate of sGFP increased with an increase in
salt concentration (salt shields charges and prevents repulsive and attractive interactions) (Figure
2G) and charge repulsion in sGFP likely prevented folding by increasing flexibility around this
region. Similar salt-dependence was absent in Wt GFP (Figure S2E) demonstrating that
substitution of lysine with glutamate introduced electrostatic repulsions that limit sGFP refolding.
Taken together, the folding landscape of sGFP had a rate-limiting entropic trap driven by flexibility
of I_1 that accounts for the slower folding rate of sGFP compared to Wt GFP.
166

GroEL/ES alters the folding pathway of sGFP

To check if the folding of sGFP was altered by GroEL/ES system, we tested for its chaperone-
dependence *in vitro* by reconstituting chaperone-assisted refolding reactions. The refolding of
sGFP was strongly accelerated (3~4 fold) by GroEL/ES (Figure 3A) system while that of Wt GFP
remained unchanged (Figure 1A, lower panel) at 25ºC. Of importance, the *in vivo* chaperonin-
dependence of sGFP was found to be intact even at 25ºC, the temperature used for *in vitro*
chaperone-assisted refolding experiments (Figure S3B). Chaperonin-dependent acceleration of
sGFP refolding was reliant on the presence of all the components; GroEL, GroES, and ATP
(Figure 3B). Thus, like authentic class-III (or IV) substrates (Kerner et al., 2005; Niwa et al., 2016)
this lab-evolved *in vivo* substrate depended on the full-cycle of the GroEL/ES system. Moreover,
the DnaK/J/E system did not alter the refolding rate of sGFP *in vitro* (Figure S3C) in line with the
in vivo data described earlier (Figure 1G). This demonstrated that the isolated mutant sGFP, was
a specific substrate of GroEL/ES *in vivo* and *in vitro,* and GroEL/ES accelerates the refolding rate
of sGFP. A single mutation K45E on Wt GFP conferred GroEL/ES dependence for its *in vivo*
folding as well as accelerated refolding rate *in vitro.*

To find the mode of acceleration by GroEL/ES, we obtained the $\Delta H^{\#}$ and ρ for GroEL/ES assisted

folding from modified Arrhenius analysis (Dandage et al., 2015). Notably, the GroEL/ES system

did not decrease the enthalpic barrier ($\Delta H^{\#}$) thereby working through the route of entropic

destabilization (Figure 3C, Table S1). A higher value of ρ for GroEL/ES assisted refolding than

for spontaneous refolding indicated a lower entropic barrier to refolding in the presence of

GroEL/ES (Figure 3C, Table S1). This was consistent with other substrates of the GroEL/ES

system (Chakraborty et al., 2010; Georgescauld et al., 2014) suggesting that entropic traps in

folding landscapes may characterize a GroEL/ES substrate in general, and GroEL/ES could have

a general role of affecting entropic destabilization to assist protein folding. The difference in

GroEL/ES-assisted and the spontaneous refolding pathway further corroborated with the

difference in m-value (dependence of refolding rate on GuHCl concentration) of folding (Figure

3D and 3E).

To check if the allosteric cycle of GroEL/ES or iterative annealing played a role in accelerating

sGFP refolding, we used the single-ring version of GroEL (SR-EL) (Horwich et al., 1998) that

lacks the negative allostery between rings, encapsulates the substrate and completes folding

inside the cavity. SR-EL/ES could accelerate the refolding of sGFP like GroEL/ES *in vitro* (Figure

S3D) and the Arrhenius parameters (Figure 3C) and m-value (Figure 3D and 3E) from SR-EL/ES

assisted refolding were similar to that assisted by GroEL/ES. The similarity of Arrhenius

parameters between SR-EL/ES and GroEL/ES dependent folding of sGFP suggested that the

temperature dependence of the refolding rates were independent of the temperature effects on

GroEL/ES allostery and was primarily determined by the folding landscape of sGFP. The similarity

of the Arrhenius parameters and the m-value suggested that the folding mechanism of sGFP had

maximal contribution from the encapsulated folding environment in the cage that had the potential

to change the folding landscape. Thus, a decrease in the entropic barrier of the refolding-substrate

was primarily affected by the GroEL/ES cavity and could rescue the entropically trapped state of

the mutant GFP.

sGFP refolding is also limited by a non-native contact formation

Interestingly the yield of spontaneous refolding of sGFP drastically reduced in the absence of any

reducing agent in the refolding buffer (Figure 4A) and reduced further in the presence $CuCl_2$, a

disulfide-assisting catalyst (Figure S4A). Since each molecule of sGFP has two cysteines (C48

and C70) that are distal in the native structure (Figure S4B), there were two possibilities, 1) the

proteins formed intermolecular-disulfides and formed covalently bonded aggregates or 2) the

proteins formed an intramolecular-disulfide bond that trapped the molecules in the non-native
state. Formation of an intramolecular-disulfide would result in the co-migration of two peptides
that were disulfide-bonded if the protein was digested with trypsin (Figure 4B). Mass spectrometry
confirmed the presence of the expected product during sGFP refolding in non-reducing conditions
and proved the formation of an intramolecular-disulfide (Figure 4C and S4C); this peptide was not
detectable in the presence of a reducing agent. Consistently, we observed the non-disulfide
bonded peptide fragments in the presence of DTT (Figure 4D). This peptide was ~100 fold
enriched in the presence of DTT than in its absence (Table S2). This proved that the intermediate
was trapped with an intramolecular-disulfide while refolding spontaneously in non-reducing
conditions. Non-reducing gel electrophoresis confirmed that intermolecular-disulfide was not a
major species when refolding was initiated in the absence of DTT (Figure 4E). Indicating that an
intramolecular-disulfide trapped the folding intermediate in a non-native state and not an
intermolecular-disulfide. Thus, the most likely model for spontaneous refolding of sGFP was
through a refolding intermediate I_1 that was free to fold to its native state with an apparent rate of
~2e-3 s^{-1}. In oxidizing conditions, I_1 converted to a quasi-stable species with non-native contacts
(I_1^*) which rapidly converted to the terminally misfolded intramolecular-disulfide bonded species
(I_2) (Figure 4F, and S4D).
Since sGFP folds poorly in the absence of DTT, we asked if delayed addition of DTT could rescue
the disulfide-trapped refolding intermediate I_2. When the refolding was initiated in the presence of
2 μM $CuCl_2$, the delayed addition of DTT could not restore the sGFP folding (Figure 5A),
suggesting that sGFP in the disulfide-bonded state enters an irreversibly misfolded state I_2 that
was refractile to reducing agents. Since more than 95% of the molecules reached I_2 within 5
minutes of the start of the refolding reaction, it revealed that there was a rapid equilibrium between
I_1 and I_1^* (Figure 4F, dotted arrows) that was much faster than the time scale of folding.
Suggesting that the formation of I_1^* did not affect the apparent rate of folding from I_1 to N.
Additionally, under conditions that favor disulfide formation, sGFP folding was limited by an
enthalpic trap that was driven by non-native contact formation in the regions surrounding the two
cysteines. The traps for this substrate could be switched on/off: the enthalpic trap could be
switched off by DTT (reducing agent), and the entropic trap could be attenuated in the presence
of salts.
246
GroEL/ES accelerates refolding rate by decreasing the entropic barrier
Could the enthalpic traps generated by the formation of non-native contacts be rescued by
GroEL/ES? Notably, the GroEL/ES system was able to accelerate the folding of sGFP and

increase the yield of refolding even in the absence of DTT (Figure 5B). Since the cysteines were
not proximal in the native structure, disulfide formation was dependent on non-native contact
formation and hence misfolding. This suggested that GroEL/ES prevents the formation of a non-
native contact during the conversion from I_1 to I_2. To further understand this, we performed the
refolding with delayed addition of the GroEL/ES system after initiating the refolding of sGFP in
the presence of 2 µM $CuCl_2$ (Figure 5C). Delayed addition of GroEL/ES led to a near-complete
loss of refolding, indicating that the GroEL/ES system was unable to break the preformed
disulfide, as expected, and could only prevent the formation of the non-native disulfide by altering
the folding path (Figure 5D). Taken together, the refolding intermediate (I_1) formed non-native
contacts to form a quasi-stable species (I_1*) that rapidly formed non-native disulfide (under
oxidizing conditions) to form the intermediate I_2. This intermediate was terminally misfolded and
refractile to folding assistance by DTT or GroEL/ES system. Implying that GroEL/ES system was
able to prevent the formation of non-native contacts in I_1 (Figure 5D, pink box) and hence the
formation of I_2, channeling folding of sGFP through a productive route.
GroEL-binding is known to cause partial unfolding of proteins, and can potentially remove non-
native interactions (Mapa et al., 2012; Sharma et al., 2008). To test this, we initiated the refolding
of sGFP in the presence of GroEL in a non-reducing buffer, but in the absence of GroES or ATP.
GroEL alone did not assist refolding (Figure S5A). Delayed addition of GroES/ATP restarted
refolding with the same rate as observed when GroEL/ES/ATP is added without any delay (Figure
S5B). In contrast to delayed addition of the full GroEL/ES/ATP system, the presence of GroEL
prevented misfolding and the amplitude of refolding did not drop even after a 20 minute of delay
in addition of GroES/ATP to GroEL-bound unfolded sGFP. (Figure 5E, and S5C). This proved
that GroEL-binding to the non-native state of sGFP prevented the protein from forming non-native
interactions, thereby maintaining it in a folding-competent state for a long time. Taken together,
the GroEL/ES system or GroEL alone can prevent cysteines from coming in close contact; thereby
averting the formation of non-native contacts that would result in enthalpic traps during folding.
While GroEL/ES was able to prevent I_1 to I_2 conversion, the refolding rate of GroEL/ES assisted
folding was same in the presence or absence of DTT in the refolding reaction (Figure 5F).
Essentially, GroEL/ES assisted refolding rate was faster than the spontaneous folding rate of
sGFP even in the presence of DTT (Figure 5G). The refolding rate of sGFP did not increase as a
function of DTT concentrations in the range used for the refolding here (Figure S5D) negating the
argument that partial reduction of disulfide was the reason for slower spontaneous folding than
with GroEL/ES system. This clearly demonstrated that the GroEL/ES system could accelerate the
refolding rate of sGFP even under (reducing) conditions where I_1 to I_2 conversion was completely

prevented. Thus, in addition to preventing I_1 to I_2 conversion, GroEL/ES assistance increased the
folding rate from I_1 to N irrespective of the redox condition of the refolding reaction (Figure 5D).
However, the enthalpic trap was not specific to the K45E mutation of GFP, rather it was the
property of the GFP backbone that showed a redox-dependent change in refolding amplitude and
rate (Figure S5E). Thus, the entropic component of the folding barrier possibly rendered the K45E
mutation amenable to GroEL/ES dependent folding *in vivo* and *in vitro*.
Collectively, GroEL/ES worked in a bipartite manner to assist sGFP refolding by (Figure S3A,
upper panel), 1) preventing non-native contact formation and hence the formation of non-
productive off-pathway intermediates that had the potential to form a terminally misfolded
conformation and 2) decreasing the entropic barrier in the folding pathway to increase the folding
rate (Figure S3A, lower panel).
295
Implications
While proteins evolve, they accumulate mutations. A subset of these mutations may enhance the
existing activity or impart new functions, they are also more likely to destabilize proteins than
mutations on other regions of the protein surface (Tokuriki et al., 2008). Chaperones have been
proposed to aid these transitional sequences allowing them to cross fitness barriers.
Aiming to link the molecular mechanism of chaperone-dependent buffering to the specific
perturbations in protein-folding landscapes we used GroEL/ES as the model chaperone. We
found that the GroEL/ES system buffers the entropic traps that can arise due to mutations that
stabilize the folding intermediates entropically.
Interestingly, GroEL/ES could also prevent the formation of incorrect contacts in the folding
polypeptide (rescue of enthalpic traps). However, this was not an exclusive activity of GroEL/ES
as we show that even the DnaK/DnaJ/GrpE chaperone machinery shares this capability. Given
that binding of non-native proteins with DnaK (Banerjee et al., 2016; Mattoo et al., 2014; Rüdiger
et al., 1997) or DnaJ (Tiwari et al.) or GroEL (Lin et al., 2008; Sharma et al., 2008) can unfold
protein chains and melt preformed contacts, it is conceivable that the holdase and the unfoldase
action is required for removing enthalpic traps from folding landscapes. DnaK machinery in *E. coli*
cytosol is more abundant than the GroEL/ES machinery and hence the mutations that introduce
enthalpic traps in the folding pathway are more likely to be channeled through the former for
efficient rescue.
While the current knowledge can be used in protein redesigning, our proposition, that GroEL/ES
may specifically buffer mutations that traps flexible folding intermediates in entropic traps, will
have interesting implications if it is found to be generally true in natural evolution.

METHODS

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Strains, Plasmids, and Proteins

E. coli strain DH5α was used for cloning, WT *E. coli* K-12 (BW25113) strain was used for expression of arabinose inducible pBAD GFP and BL21 (DE3) was used for protein expression and purification. Protein concentrations were determined spectrophotometrically at 562 nm using BCA kit (Pierce-ThermoFisher Scientific). Deletion strains were obtained from CGSC as part of Keio collection (Baba et al., 2006).

METHOD DETAILS

Construction of mutant GFP library

Mutant GFP library was made in arabinose inducible pBAD vector using a random mutagenesis approach by error-prone polymerase Mutazyme II (Agilent) that incorporated 7 to 11 mutations per kb of the template. The said library has a total complexity of around 10,000 mutants. The reporter is constructed such that GFP and mCherry are under the same arabinose inducible pBAD promoter in an operon to give a readout of GFP according to the mutation created on it but the mCherry readout will remain similar thus serving as an internal control for transcription, translation, and inducibility.

Screening of mutant GFP library for GroEL/ES dependent GFP mutant

Wild type *E. coli* cells (WT) were transformed with the mutant GFP library maintaining 10-fold converge for preserving complexity. 0.4 OD600 cells were induced with 0.1% arabinose and fluorescence was observed three hours post-induction at 37°C after diluting cells in 1X PBS and incubating at 37°C for 1 hour. Fluorescence of the mutant library was studied in a pooled manner against wild type GFP. The entire library was sorted into populations of very low fluorescent, low fluorescent, and mildly less fluorescent according to the GFP fluorescence. Each of these populations was purified and plasmids prepared. WT cells were co-transformed with GroEL/ES over-expressing plasmid (Castanie et al., 1997) and pool of GFP mutant plasmids in a sequential manner, maintaining the minimum 10X coverage. The pool of transformants were grown and

GroEL/ES induced by adding 0.5 mM IPTG 30 minutes before induction of GFP by 0.1%
arabinose. After that the induced cultures were grown for another 3 and half hours. FACS was
performed to sort *E.coli* cells showing high fluorescence upon GroEL/ES overexpression. To
confirm the dependence of sorted clones on GroEL/ES, plasmid pool was prepared from the
sorted cells (harboring pBAD GFP as well as pOFX GroEL/ES) and digested using SacII
(linearizes pOFX GroEL/ES only) to obtain GFP clones post-transformation in wild type *E.coli*
cells. Single clones of GFP were picked from here and checked for their fluorescence in the
presence and absence of GroEL/ES overexpression. The isolated mutant having higher
fluorescence in GroEL/ES overexpression was identified by Sanger sequencing. Isolated
GroEL/ES dependent mutant of GFP (K45E) was cloned in pET SUMO between BamHI and
HindIII restriction sites and purified using *E. coli* BL21 (DE3) for further characterization.

***E.coli* GroEL/ES, DnaK, DnaJ, GrpE Expression, and Purification**

GroEL/ES, GroEL chimeras, DnaK, DnaJ, GrpE were purified using *E. coli* BL21 (DE3) as
described (Kerner et al., 2005; Mapa et al., 2012; Tiwari et al., 2013). GroEL/ES was expressed
from pOFX plasmid for co-expression studies (Castanie et al., 1997).

Solubility of sGFP *in vivo*

WT *E.coli* K-12 (BW25113) cells containing pBAD sGFP were transformed with pOFX GroEL/ES.
0.1% inoculation was done in 10 ml LB medium added with chloramphenicol (35 μg/ml) and
ampicillin (100 μg/ml) from overnight grown cultures and grown till OD600-0.5 at 37°C, 200 rpm.
0.5 mM IPTG was added to cells with pOFX GroEL/ES 30 minutes before the induction of sGFP
with 0.1% arabinose for 3 hours. The cell type in which we transformed only pBAD sGFP were
directly induced with 0.1% arabinose for 3 hours after reaching OD600- 0.5. Cells were harvested
at 4000 rpm for 10 minutes and resuspended in 1 ml PBS pH 7.4, 2 mM DTT. Lysis was done
using sonication followed by high-speed centrifugation to separate soluble and pellet fraction
which was separately loaded onto 12% SDS-PAGE. Gel visualized by Coomassie staining.

Spontaneous and chaperonin assisted *in vitro* refolding of sGFP

Wt GFP and sGFP (20 μM each) were denatured in buffer containing 6 M GuHCl in buffer A (20
mM Tris, 20 mM KCl, 5 mM $MgCl_2$, 2 mM DTT, pH 7.4) for 1 hour at 25°C and refolded upon 100
fold dilution into buffer A. Either of three refolding buffers were used for refolding, buffer A to mimic
reducing conditions, buffer B (20 mM Tris, 20 mM KCl, 5 mM $MgCl_2$, pH 7.4) to mimic non-

reducing conditions and buffer C (20 mM Tris, 20 mM KCl, 5 mM MgCl$_2$, 2 μM CuCl$_2$, pH 7.4) to

mimic oxidizing conditions.

GroEL/ES assisted refolding was done in the presence of (400 nM) GroEL (tetradecamer), (800

nM) GroES (heptamer), (substrate:GroEL:ES :: 1:2:4) and the refolding was started by addition

of 2 mM ATP. SR-EL/ES assisted refolding was done in the presence of (800 nM) of SR-EL

(heptamer) (800 nM) GroES (heptamer) and the refolding was started by the addition of 2 mM

ATP. GFP fluorescence at 480 nm excitation (slit width 2 nm) and 515 nm emission (10 nm slit

width) was monitored as a readout of refolding using Fluorolog 3 Spectrofluorometer (Horiba).

Buffer conditions are described in the figures. All the unfolding and refolding experiments were

carried out at 25°C unless specified.

GuHCl concentration-dependent spontaneous and chaperonin assisted *in vitro* refolding

of sGFP

sGFP (80 μM) were denatured in buffer containing 6 M GuHCl in buffer A for 1 hour at 25°C and

was refolded upon 400 times dilution in buffer A alone or buffer A containing (400 nM) GroEL

(tetradecamer), (800 nM) GroES (heptamer) and the refolding was started by addition of 2 mM

ATP. GuHCl present in unfolded sGFP was diluted to 15 mM upon 400 times dilution in buffer A.

GuHCl concentration was increased to 30 mM, 45 mM, 60 mM by adding GuHCl from outside in

refolding mixture.

Analysis of temperature-dependent refolding using Arrhenius equation

To obtain thermodynamic parameters that define the barrier between the refolding intermediate

I$_1$ and the transition state (TS) of folding we used the following equation essentially as defined in

(Dandage et al., 2015).

400
$$k_f = \rho \, T e^{-\frac{\Delta H^{\#} - \Delta C_p^{\#}\left(T - T_0 - T ln\left(\frac{T}{T_0}\right)\right)}{RT}}$$

401   Where

402
$$\rho = \left(\frac{k_B}{h}\right)\left(\kappa. e^{\frac{\Delta S^{\#}}{R}}\right)$$

and k$_f$ is refolding rate, R is the universal gas constant, ΔH$^{\#}$, ΔS$^{\#}$ and ΔC$_p^{\#}$ are the differences in

enthalpy, entropy and heat capacity at constant pressure and the reference temperature T$_0$

between TS and I$_1$, respectively, T is the temperature of refolding reaction, T$_0$ is the reference

temperature at which the parameters are determined (here itis 298.15 K). κ is the transmission

factor, that reports the proportion of activations that lead to the formation of the native state (N),

and k_B and h are Boltzmann and Planck constant respectively. Since $\Delta S^{\#}$ and A linearly combine,

it is not possible to obtain independent estimates of these two parameters by non-linear

regression. Hence, we combine it to obtain ρ. This term indicates the ease of barrier crossing,

either because the diffusion is faster or because the barrier is less broad (due to lower $\Delta S^{\#}$). A

high $\Delta H^{\#}$ indicates a higher enthalpic barrier, and a low ρ indicates an entropic (diffusion-limited)

barrier. The equations were fitted using standard non-linear regression fitting using R or Octave.

Fitting was performed by varying the starting parameters by 4-fold within the rage of expected

values reported for globular proteins (Dandage et al., 2015), and the fitting was deemed

satisfactory only when the r-squared values were above 0.9 and the dependencies were above

0.9 for the different parameters that were floated during fitting. The floating parameters were, ρ,

ΔC_p, ΔH.

Mass spectrometry analysis for disulfide bond detection:

5 µg of control or DTT treated sGFP was digested by using sequencing grade trypsin (1:10 ratio,

Trypsin:protein) for 16-18 hours at 37°C. Before the digestion DTT treated sample was alkylated

by using 55 mM iodoacetamide. Tryptic digested peptides were reconstituted in 5µl of LC-MS

grade water containing 0.1% formic acid and run on a quadrupole-TOF hybrid mass spectrometer

(TripleTOF 6600, Sciex, USA) coupled to a nano-LC system (Eksigent NanoLC-400). Two

microliters of sample was injected and loaded onto a reverse-phase peptide Chromo LC trap (200

mm 0.5 mm) column and peptides were separated using a C18 column (75 mm 15 cm, Eksigent).

The samples were run using a gradient method using buffer X (99.9% LC-MS water + 0.1% formic

acid) and buffer Y (99.9% acetonitrile + 0.1% formic acid). The gradient consists of 95% of buffer

X for 2 minutes, and then shifted to 90% of buffer X for 8 minutes, and then decreased to 20% of

buffer X in 42 minutes and finally shifted to 95% of buffer X again for 16 minutes at a consistent

flow rate of 250 nl min-1. Data was acquired with a NanoSpray source installed in the TripleTOF

6600 System using a nebulizing gas of 20 psi, a curtain gas of 25 psi, an ion spray voltage of

2000 V, and a heater interface temperature of 75 0C. Information-dependent acquisition (IDA)

mode was set up with a TOF/MS survey scan (350–1600 m/z) with an accumulation time of 250

ms. For fragmentation, a maximum of ten precursor ions per cycle was selected with each MS/MS

spectrum (200–1800 m/z) accumulated for 70 ms with a total cycle time of approximately 2.05

seconds. Parent ions with a charge state from +2 to +5 and an abundance of more than 150 cps

were selected for MS/MS fragmentation. Once an ion had been fragmented by MS/MS, its mass

and isotopes were excluded for 3 seconds. The wiff files generated from Triple TOF 6600 (which

contain both MS and MS/MS spectra) were analyzed using the protein pilot v5.0 for the

identification of our desire protein. For disulfide bond detection and further spectra analysis was

done using Biopharma view v2.0 and peak view 2.2 software (Sciex).

QUANTIFICATION AND STATISTICAL ANALYSIS

Student t-test and R package for non-linear regression was used for statistical analysis. Flow-

cytometry data was analyzed using octave.

DATA AND SOFTWARE AVAILABILITY

All data are provided in the manuscript. We did not develop any new software.

KEY RESOURCE TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit anti-GFP	Abcam	AB290
Bacterial and Virus Strains		
E.coli BL21 (DE3)		
E.coli DH5α		
E.coli WT (K-12, BW25113) F-, DE(araD-araB)567, lacZ4787(del)::rrnB-3, LAM-, rph-1, DE(rhaD-rhaB)568, hsdR514	(Baba et al., 2006)	CGSC#: 7636

Δ*Tig* (JW0426-1) *F-, Δ(araD-araB)567, ΔlacZ4787(::rrnB-3), Δtig-722::kan, λ-, rph-1, Δ(rhaD-rhaB)568, hsdR514*	(Baba et al., 2006)	CGSC#: 8589
Δ*dnaK* (JW0013-4) F-, Δ(araD-araB)567, ΔlacZ4787(::rrnB-3), λ-, rph-1, ΔdnaK734::kan, Δ(rhaD-rhaB)568, hsdR514	(Baba et al., 2006)	CGSC#: 8342
Δ*dnaJ* (JW0014-1) F-, Δ(araD-araB)567, ΔlacZ4787(::rrnB-3), λ-,rph-1, ΔdnaJ735::kan,Δ(rhaD-rhaB)568, hsdR514	(Baba et al., 2006)	CGSC#: 8343
Chemicals, Peptides, and Recombinant Proteins		
Proteinase K	MP Biomedicals	Cat#193981
ATP	Sigma	Cat#A2383
Oligonucleotides		
yeGFP Forward primer: CCCGGATCCATGTCTAAAGGTGAAGAATTATTCACTGGTGTTGTCCC	This Paper	
yeGFP Reverse primer: CCCAAGCTTTTATTTGTACAATTCATCCATACCATGGGTAATA	This Paper	
Recombinant DNA		
pOFX ELES (pOFX tac-SL2)	(Castanie et al., 1997)	
pOFX DnaK/J/GrpE (pOFX tac-KJE1)	(Castanie et al., 1997)	
pET duet1 dnaK	(Tiwari et al., 2013)	

pET duet1 dnaJ	(Tiwari et al., 2013)	
pET duet1 grpE	(Tiwari et al., 2013)	
pET SUMO sGFP	This Paper	
Software and Algorithms		
Chimera		
ABPS		

ACKNOWLEDGEMENT

We thank Prof. Hideki Taguchi for the kind gift of pET21 SR1 plasmid. We also thank Kanika Saxena for sharing the starting plasmid with GFP and mCherry. KM acknowledges the funding from Department of Biotechnology (DBT), Government of India, grant number (BT/PR28386/BRB/10/1671/2018) and Science and Engineering Research Board (SERB), Government of India, for Core Research Grant (SERB/CRG/2019/006281) and SNU core funding. The work in KC Lab was supported by Swarnajayanthi Fellowship Grant from DST, and partly from BSC0124 from CSIR. Instrument support was also obtained from Wellcome Trust-DBT India Alliance and CSIR. KM and MA acknowledge SNU for infrastructural support and KC acknowledges CSIR and CSIR-IGIB for infrastructural support. A. Sadat and ST acknowledges DBT; AR, KV, AC, A. Singh acknowledge CSIR; AG and RC acknowledge UGC for fellowship support. MA acknowledges SNU doctoral fellowship.

AUTHOR CONTRIBUTION

Conceptualization: Koyeli Mapa

Supervision: Koyeli Mapa, Kausik Chakraborty

Reagent Generation: Satyam Tiwari, Kanika Verma, Anwar Sadat, Aseem Chaphalkar,

Experiments: Anwar Sadat, Satyam Tiwari, Kanika Verma, Mudassar Ali, Vaibhav Upadhyay,

Anupam Singh, Aseem Chaphalkar, Asmita Ghosh, Rahul Chakraborty, Kausik Chakraborty

Analysis: Arjun Ray (computational), Anwar Sadat (biophysics), Satyam Tiwari (biophysics),

Kanika Verma (FACS), Rahul Chakraborty (MS), Kausik Chakraborty, Koyeli Mapa

Manuscript writing: Koyeli Mapa and Kausik Chakraborty with inputs from Anwar Sadat, Kanika

Verma, and Aseem Chaphalkar. All the authors read and edited the manuscript.

DECLARATIONS OF INTEREST

The authors declare no competing interests.

References

FIGURE LEGENDS:

Figure 1: Laboratory evolution of an authentic substrate of GroEL/ES

(A) Upper panel: Histogram for *in vivo* fluorescence of Wt GFP (represented as the ratio of
GFP/mCherry) in the presence and absence of plasmid-based overexpression of GroEL/ES.

Lower panel: Refolding kinetics of Wt GFP in the presence and absence of GroEL/ES/ATP.
Spontaneous refolding of Wt GFP was initiated by a 100-fold dilution of the unfolded proteins in
buffer A (20 mM Tris, 20 mM KCl, 5 mM MgCl$_2$, 2 mM DTT, pH 7.4) at 25°C. For GroEL/ES
assisted refolding unfolded protein was diluted 100-fold in buffer A containing 400 nM of GroEL
(all concentrations of GroEL are in terms of tetradecamer), 800 nM GroES (all concentration of
GroES are in terms of heptamer) so that the final concentration of unfolded protein is 200 nM.
Subsequently, refolding was initiated by adding 2 mM ATP. Recovery of GFP fluorescence over
time was followed to monitor refolding.

(B) Schematic of the bicistronic construct of GFP and mCherry drove by an Arabinose inducible
system. RBS indicates the position of the additional ribosome binding site (RBS) to initiate
translation of mCherry.

(C) Upper panel: Scatter plot of *E. coli* WT cells expressing random mutant library of GFP. The

red box highlights the population of cells with low GFP fluorescence (LF-GFP) compared to Wt

GFP, with the same mCherry fluorescence as that of Wt GFP.

Lower panel: Histogram for flow cytometry of cells harboring low-fluorescent (LF) mutant pool of

GFP and the Wt GFP. GFP/mCherry ratio is monitored at a single-cell level.

(D) Upper panel: Schematic of FACS to isolate GroEL/ES-dependent mutant GFP clones.

Lower panel: The cells harboring LF-GFP library were induced to express GFP mutants either

alone or along with the GroEL/ES system. Negative gating (gate I) was defined by the single-cell

GFP/mCherry ratio of the LF-GFP when expressed alone (red dots). Separately, LF-GFP

expression was induced in the presence of overexpressed GroEL/ES (blue dots), Positive clones

were sorted by gating for cells that had higher GFP/mCherry ratio than Gate I (Positive gate).

Gate I included cells only with high GFP fluorescence along with a high GFP/mCherry ratio.

(E) K45 residue depicted on the crystal structure of GFP (1GFL (Xu et al., 1997; Yang et al.,

1996)) showing its surface exposure away from the buried fluorophore. Some of the residues in

beta-sheets facing the viewer are not shown to make the fluorophore underneath clearly visible.

Only one chain of GFP is shown for clarity. Depiction made using Chimera (Pettersen et al., 2004).

(F) Histogram of single-cell GFP/mCherry ratio of sGFP (GFP (K45E)) in the presence and

absence of GroEL/ES over-expression. For individual histograms of GFP and mCherry please

see Figure S1C.

(G) Histogram of single-cell GFP/mCherry ratio of sGFP in the presence and absence of

overexpressed DnaK/J/E. For individual histogram of GFP and mCherry please see Figure S1F.

(H) Histogram for in vivo GFP/mCherry fluorescence of sGFP in wild type E.coli K12 (WT) and

knockout strains for canonical molecular chaperones dnaK (∆*dnaK*), dnaJ (∆*dnaJ*) and trigger

factor (∆*tig*). For individual histogram of GFP and mCherry please see Figure S1G.

(I) Bar plot for in vivo GFP/mCherry fluorescence of sGFP and 6 independent GFP mutants

isolated from LF-GFP library in wild type *E.coli* K12 (WT) overexpressing GroEL/ES.

See Figure S1.

Figure 2: A comprehensive folding landscape for sGFP

(A) Comparison of the refolding kinetics of WtGFP and sGFP. WtGFP and sGFP were unfolded in 6 M GuHCl in buffer A for 1 hour at 25°C. Refolding was initiated by a 100-fold dilution of the unfolded proteins in buffer A (20 mM Tris, 20 mM KCl, 5 mM MgCl$_2$, 2 mM DTT, pH 7.4) at 25°C so that the final concentration of the proteins was 200 nM. Refolding was monitored by following the fluorescence of GFP with time.

(B) Refolding rate and amplitude as a function of sGFP concentration. sGFP was unfolded as described earlier at 25°C but at different concentrations. Refolding was initiated from each of these by a 100-fold dilution into buffer A at 25°C such that the final protein concentrations were 12.5, 100, 200, or 400 nM. Refolding was monitored by measuring the recovery of GFP fluorescence over time. The refolding traces were fitted to obtain the apparent folding rate (red circles) and the percentage of expected amplitude (black circles). These were plotted as a function of protein concentration. The expected amplitude was calculated based on the fluorescence recovery upon refolding protein at 400 nM concentration.

(C) Refolding rate as a function of sGFP concentration. sGFP was unfolded as described earlier at 37°C but at different concentrations. Refolding was initiated from each of these by a 100-fold dilution into buffer A at 37°C such that the final protein concentrations were 100, 200, or 400 nM. Refolding was monitored by measuring the recovery of GFP fluorescence over time.

(D) Arrhenius plot for sGFP and Wt GFP refolding. Refolding of sGFP or Wt GFP was initiated as described earlier. The refolding reactions were initiated in buffer A that were preincubated at different temperatures (15°C, 20°C, 25°C, 30°C and 35°C) and monitored by measuring the recovery of GFP fluorescence. Rates obtained by fitting the refolding traces obtained at different temperatures are plotted against (1/T) for sGFP (grey circles), and Wt GFP (black circles). The rates were obtained from independent replicates of 3 refolding reactions at each temperature. Line plots are the fits obtained by fitting the temperature-dependent refolding rates to the Arrhenius equation as described in the supplemental text.

(E) The calculated enthalpy of activation ($\Delta H\#$), and ρ obtained from Arrhenius fitting of sGFP and Wt GFP refolding shown in (D).

(F) Close up of K45 (orange:side-chain carbons of K45, navy blue:εN) region on GFP (using 1GFL) (Yang et al., 1996), and its interacting amino acids. The model was prepared using Chimera.

557 (G) Comparison of the refolding rate of sGFP in the presence of different salts of equal ionic
558 strength. Spontaneous refolding of sGFP was initiated by a 100-fold dilution of the unfolded
559 proteins in buffer A at 25°C. For salt assisted refolding unfolded protein was diluted 100-fold in
560 buffer A containing different salts (100 mM $MgCl_2$, or 300 mM NaCl, or 300 mM KCl) of equal ionic
561 strength so that the final concentration of unfolded protein is 200 nM. Recovery of GFP
562 fluorescence over time was followed to monitor refolding.

563 See Figure S2 and S3.

564 **Figure 3: GroEL/ES alters the folding pathway of sGFP**

565 (A) Left panel: Comparison of the refolding kinetics of sGFP in the presence and absence of
566 GroEL/ES/ATP. Spontaneous refolding of sGFP was initiated as mentioned earlier either in buffer
567 A. For GroEL/ES assisted refolding unfolded protein was diluted 100-fold in buffer A containing
568 400 nM of GroEL (all concentrations of GroEL are in terms of tetradecamer), 800 nM GroES (all
569 the concentrations of GroES are in terms of heptamer) so that the final concentration of unfolded
570 protein is 200 nM. Subsequently, refolding was started by adding 2 mM ATP. Recovery of GFP
571 fluorescence over time was followed to monitor refolding.

572 Right panel: Refolding traces were fitted to single exponential kinetics to obtain the refolding rates
573 shown as bar plots. Three independent refolding runs were fitted separately to obtain the standard
574 deviation in rates (shown as errors on the bar graph).

575 (B) Monitoring the dependence of sGFP refolding on the different components of EL/ES. Unfolded
576 sGFP was diluted 100-fold either in the presence of GroEL (400 nM), or GroEL (400 nM) + GroES
577 (800 nM) or GroEL (400 nM) + ATP (2 mM) or GroEL (400 nM) + GroES (800 nM) + ATP (2 mM)
578 and refolding was monitored by the recovery of GFP fluorescence.

579 (C) The calculated enthalpy of activation ($\Delta H^{\#}$), and ρ obtained from Arrhenius fitting of
580 GroEL/ES/ATP and SR-EL/ES/ATP assisted refolding of sGFP.

581 (D) Effect of GuHCl on refolding rate upon spontaneous, GroEL/ES/ATP and SREL/ES/ATP
582 assisted refolding of sGFP. The rates [$\ln(k_f)$] of spontaneous, GroEL/ES/ATP, and SREL/ES/ATP
583 assisted refolding of sGFP at different GuHCl concentration are plotted against different GuHCl
584 concentrations.

585 (E) The m-value (dependence of refolding rate on GuHCl concentration) of folding upon
586 spontaneous, GroEL/ES/ATP, and SR-EL/ES/ATP assisted refolding of sGFP.

587 **Figure 4: GroEL/ES prevents non-native contact formation**

(A) Refolding of sGFP was initiated in buffer A (shown as +DTT) or buffer B (20 mM Tris, 20 mM KCl, 5 mM MgCl$_2$, pH 7.4) (shown as -DTT). Refolding was monitored by the recovery of GFP fluorescence over time.

(B) Schematic representation of a hypothetical fragment pattern when an intramolecular disulfide is formed (in the absence of DTT) and in its absence (in the presence of DTT).

(C) The expected peptide sequence and mass upon the formation of the intramolecular disulfide is shown along with the experimentally observed isotopic mass distribution obtained for the peptide when refolding of sGFP is initiated in the absence of DTT.

(D) The peptide fragment expected when disulfide bonding is prevented during sGFP refolding in DTT, and the experimentally observed isotopic mass distribution for the peptide is shown.

(E) Non-reducing PAGE to monitor the formation of intermolecular disulfide bonds. Folding of sGFP was initiated as described earlier in buffer B or buffer A at a final concentration of 500 nM sGFP. The proteins were then boiled and loaded onto a non-reducing polyacrylamide gel to resolve the proteins. Immunoblotting was performed to detect sGFP using anti-GFP antibody.

(F) A simplistic model of the spontaneously refolding sGFP. The unfolded state reaches the folding intermediate I$_1$ which then converts to the native state N. I$_1$ is in rapid equilibrium with I$_1$* where the cysteines (shown as orange circles) come close. I$_1$*, in turn, can form the disulfide in oxidizing conditions and form a terminally misfolded state I$_2$ that is refractile to refolding even in the presence of DTT.

See Figure S4.

Figure 5: GroEL/ES accelerates refolding rate by decreasing the entropic barrier

(A) Effect of delayed addition of DTT to spontaneously refolding sGFP. sGFP refolding was initiated in buffer C (20 mM Tris, 20 mM KCl, 5 mM MgCl$_2$, 2 μM CuCl$_2$ pH 7.4) at a final concentration of 200 nM of sGFP. DTT was either added before unfolded sGFP was diluted in the buffer to start refolding (no delay) or after 5 min or 10 min. Refolding traces for 5 min and 10 minutes delayed addition of DTT are shown from the time DTT was added.

(B) Spontaneous or GroEL/ES dependent refolding of sGFP was initiated as described earlier, except that the refolding was performed in buffer B for both the refolding reactions.

(C) Effect of delayed addition of GroEL/ES/ATP to spontaneously refolding sGFP. Refolding of sGFP was initiated as earlier in buffer C, and GroEL/ES/ATP (400 nM/800 nM/2 mM respectively)

was added after a time-delay of 5 min or 10 min. For zero delay, GroEL/ES/ATP was added to buffer C before adding unfolded sGFP.

(D) Schematic of the refolding pathway of sGFP in the presence and absence of GroEL/ES. GroEL/ES/ATP efficiently prevents the formation of I_1^* (shown by the red arrow) and also allows more efficient conversion of I_1 to N (shown by the green arrow).

(E) GroEL-binding is sufficient to prevent sGFP misfolding. sGFP refolding was initiated as earlier, in buffer C containing 400 nM GroEL. GroES/ATP (800 nM/2 mM, respectively) was added after time delays of 5 min, 10 min, or 20 min. Refolding was monitored although the time course, traces shown are after the addition of GroES/ATP. For zero delay addition, GroES/ATP was present along with GroEL in buffer C before unfolded sGFP was added to the buffer.

(F) Effect of DTT on GroEL/ES assisted refolding rate of sGFP. Refolding of sGFP was initiated as described in buffer C (black line), buffer C containing GroEL/ES/ATP (400 nM, 800 nM, 2 mM, respectively) (red line), buffer A (blue line), and buffer A containing GroEL/ES/ATP (400 nM, 800 nM, 2 mM, respectively) (maroon circles).

(G) Rates for refolding in the latter three (from Figure 5F) were obtained by fitting to the exponential rate equation and is shown as bar graphs. Errors shown are standard deviations over three independent replicates.

See Figure S5.

Supplementary Figures:

Figure S1: related to Figure 1

(A) Bar graph for *in vivo* fluorescence of GroEL/ES dependent 10 single GFP clones isolated through FACS with (red) and without (black) GroEL/ES overexpression.

(B) Bar graph for native fluorescence of sGFP (K45E) compared to Wt GFP.

(C) Independent fluorescence of GFP and mCherry channel for sGFP with (red) and without (black) GroEL/ES overexpression. GroEL/ES overexpression leads to an increase in GFP fluorescence (left panel) with similar mCherry fluorescence (right panel).

(D) Western blot showing the degradation of sGFP after translation halt in the absence of GroEL/ES overexpression. Over time slow folding sGFP is degraded after translation is halted in

absence of GroEL/ES overexpression. The western blot is representative of three independent
experiments.

(E) *in vivo* solubility of sGFP in the absence (left panel) and presence (right panel) of GroEL/ES
over-expression. Partitioning of sGFP is shown in the total lysate (T), pellet fraction (P), and
soluble fraction (S).

(F) Independent fluorescence of GFP and mCherry channel for sGFP with (red) and without
(black) plasmid-based DnaK/J/GrpE overexpression. Overexpression of DnaK/J/GrpE decreases
GFP fluorescence (upper panel) with no effect on mCherry fluorescence (lower panel).

(G) Independent fluorescence of GFP (upper panel) and mCherry (lower panel) channel for sGFP
in WT cells and in cells with the single-gene knockout of the most abundant chaperones (*Δtig*,
ΔdnaK, and *ΔdnaJ*)

(H) Sequences of 6 GFP mutants obtained after Sanger sequencing were aligned with the Wt
GFP sequence. All the different mutations that are present in the GFP mutants are marked with
a red-colored box.

Figure S2: related to Figure 2

(A) sGFP was unfolded in 6 M GuHCl in buffer A for 1 hour at 25°C. Unfolding was monitored by
following the decrease in fluorescence of GFP with time.

(B) Refolding kinetics of spontaneously refolding sGFP in buffer A as followed by GFP
fluorescence (upper panel open circles). Single exponential fit to the refolding is shown as a red
line in the upper panel. The scatter from residuals are shown in the bottom panel.

(C-D) sGFP refolding kinetics at different concentrations of sGFP (C). sGFP was unfolded as
described earlier but at different concentrations. Refolding was initiated from each of these by a
100-fold dilution into buffer A such that the final protein concentrations were 12.5, 100, 200, or
400 nM. Refolding was monitored by measuring the recovery of GFP fluorescence over time. The
plot of normalized GFP fluorescence/time (D).

(E) Comparison of the refolding rate of sGFP (left panel) and Wt GFP (right panel) in the presence
of different concentrations of $MgCl_2$. Spontaneous refolding of both the proteins was initiated by
a 100-fold dilution of the unfolded proteins in buffer A at 25°C containing different concentrations

of MgCl$_2$ as shown in the figure. Recovery of GFP fluorescence over time was followed to monitor

refolding.

Figure S3: Schematic showing barriers in the folding landscape, related to Figure 2

(A) A section of the folding funnel is represented in all cases. Unfolded state (U) forms the folding

intermediate (I) that are separated by a transition state (TS) from the native state (N) of the protein.

Inset shows the axis - vertical axis represents enthalpy, whereas the cylindrical axis (distance

from the center of rotation) represents entropy of the protein chain.

The upper panel schematically shows an enthalpic barrier in folding, essentially a hill in the

enthalpy axis, that leads to slower folding. A decrease in the height of the hill (decreasing $\Delta H^{\#}$)

leads to faster folding (right panel).

The lower panel schematically depicts an entropic barrier. The intermediate (I) is a flat area in the

landscape that delays folding as the conformational search is undirected and it takes time to reach

TS. The flatness is inversely proportional to ρ and hence a lower ρ indicates slow folding (left

panel) while an increase in ρ (decrease in the flat basin) leads to accelerated folding (right panel).

(B) Bar graph showing GFP/mCherry ratio obtained from *in vivo* fluorescence of sGFP in the

presence (green) and (red) absence of GroEL/ES at 25^{0}C in the flow cytometer.

(C) Refolding kinetics of sGFP in the presence and absence of DnaK/J/GrpE/ATP. Unfolded

protein was diluted 100-fold in buffer A containing 800 nM of DnaK, 400 nM DnaJ, 800 nM GrpE

(2:1:2), so that the final concentration of unfolded protein is 200 nM. Subsequently, refolding was

initiated by adding 5 mM ATP. Recovery of GFP fluorescence over time was followed to monitor

refolding.

(D) Comparison of the refolding rate of sGFP in the presence of GroEL/ES and SR-EL/ES in

buffer A at 25°C.

Figure S4: related to Figure 4

(A) Refolding kinetics of sGFP in the presence and absence of CuCl$_2$. Spontaneous refolding of

sGFP was initiated by a 100 fold dilution of the unfolded proteins in buffer B at 25^{0}C. For CuCl$_2$

assisted refolding unfolded protein was diluted 100-fold in buffer B containing 2 μM of CuCl$_2$ so

that the final concentration of unfolded protein is 200 nM. Recovery of GFP fluorescence over

time was followed to monitor refolding.

(B) Figure showing the distance between C48 and C70 position (Cysteine 48 and 70 (orange),

connecting line between two cysteines (red)) on native GFP (using 1GFL). The figure made using

Chimera.

(C) Schematic showing the preparation of the sample for LC-MS. sGFP was unfolded as

described earlier and refolded by 100-fold dilution in buffer A and B for 30 minutes. Buffer

exchange of sGFP was done in 500 mM TEAB buffer. Then sGFP was subjected to digestion by

trypsin for 18 hours at 37°C. The digested product was vacuum dried and run in LC-MS.

(D) Qualitative one-dimensional energy landscape for spontaneously refolding sGFP. Black

denotes the energy landscape of sGFP while refolding in the presence of DTT. The Red dashed

line shows the change in the landscape in the presence of $CuCl_2$. The rest of the landscape

remains unchanged.

Figure S5: related to Figure 5

(A) Comparison of the refolding kinetics of sGFP in the presence of different components of

GroEL/ES/ATP. Spontaneous refolding of sGFP was initiated as mentioned earlier in buffer B at

25°C. For the different components of GroEL/ES/ATP assisted refolding unfolded protein was

diluted 100-fold in buffer C containing either GroEL alone or GroEL/ATP or GroEL/ES/ATP so

that the final concentration of unfolded protein is 200 nM. Recovery of GFP fluorescence over

time was followed to monitor refolding.

(B) Comparison of the refolding rate of sGFP upon delayed addition of GroES/ATP to GroEL

bound sGFP. Refolding of sGFP and was initiated as earlier in buffer C in the presence of GroEL,

and GroES/ATP was added after a time-delay of 5, 10, or 20 minutes. For no delay,

GroEL/ES/ATP was added to buffer C before adding unfolded sGFP.

(C) Refolding kinetics of sGFP upon delayed addition of GroES/ATP to GroEL bound sGFP

(similar to figure S5B).

(D) Refolding rate as a function of DTT concentration. sGFP was unfolded as described earlier

and refolding was initiated by a 100-fold dilution into buffer B containing different concentrations

of DTT (2,3 and 4 mM) such that the final protein concentrations were 200 nM. Refolding was

monitored by measuring the recovery of GFP fluorescence over time. Refolding traces were fitted

to single exponential kinetics to obtain the refolding rates shown as bar plots with standard

deviation in rates (shown as errors on the bar graph).

(E) Effect of non-reducing (in buffer B, shown as - DTT) and reducing environment (in buffer A,

shown as + DTT) on GFP backbone, in terms of refolding rate. Error bars indicate the standard

deviation from three independent replicates.

References:

Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., Datsenko, K.A., Tomita, M., Wanner, B.L.,
and Mori, H. (2006). Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the
Keio collection. Mol Syst Biol *2*, 2006 0008.
Bandyopadhyay, B., Goldenzweig, A., Unger, T., Adato, O., Fleishman, S.J., Unger, R., and Horovitz, A.
(2017). Local energetic frustration affects the dependence of green fluorescent protein folding on the
chaperonin GroEL. The Journal of biological chemistry *292*, 20583-20591.
Banerjee, R., Jayaraj, G.G., Peter, J.J., Kumar, V., and Mapa, K. (2016). Monitoring conformational
heterogeneity of the lid of DnaK substrate-binding domain during its chaperone cycle. FEBS J *283*, 2853-
2868.
Calloni, G., Chen, T., Schermann, S.M., Chang, H.C., Genevaux, P., Agostini, F., Tartaglia, G.G., Hayer-Hartl,
M., and Hartl, F.U. (2012). DnaK functions as a central hub in the E. coli chaperone network. Cell reports
1, 251-264.
Castanie, M.P., Berges, H., Oreglia, J., Prere, M.F., and Fayet, O. (1997). A set of pBR322-compatible
plasmids allowing the testing of chaperone-assisted folding of proteins overexpressed in Escherichia coli.
Anal Biochem *254*, 150-152.
Chakraborty, K., Chatila, M., Sinha, J., Shi, Q., Poschner, B.C., Sikor, M., Jiang, G., Lamb, D.C., Hartl, F.U.,
and Hayer-Hartl, M. (2010). Chaperonin-catalyzed rescue of kinetically trapped states in protein folding.
Cell *142*, 112-122.
Chiti, F., and Dobson, C.M. (2017). Protein Misfolding, Amyloid Formation, and Human Disease: A
Summary of Progress Over the Last Decade. Annual review of biochemistry *86*, 27-68.
Dandage, R., Bandyopadhyay, A., Jayaraj, G.G., Saxena, K., Dalal, V., Das, A., and Chakraborty, K. (2015).
Classification of chemical chaperones based on their effect on protein folding landscapes. ACS Chem Biol
10, 813-820.
Dunn, A.Y., Melville, M.W., and Frydman, J. (2001). Review: cellular substrates of the eukaryotic
chaperonin TRiC/CCT. Journal of structural biology *135*, 176-184.
Georgescauld, F., Popova, K., Gupta, A.J., Bracher, A., Engen, J.R., Hayer-Hartl, M., and Hartl, F.U. (2014).
GroEL/ES chaperonin modulates the mechanism and accelerates the rate of TIM-barrel domain folding.
Cell *157*, 922-934.
Hartl, F.U. (2017). Protein Misfolding Diseases. Annual review of biochemistry *86*, 21-26.

Horwich, A.L., Burston, S.G., Rye, H.S., Weissman, J.S., and Fenton, W.A. (1998). [11] Construction of
single-ring and two-ring hybrid versions of bacterial chaperonin GroEL. In Methods in enzymology
(Elsevier), pp. 141-146.

Houry, W.A., Frishman, D., Eckerskorn, C., Lottspeich, F., and Hartl, F.U. (1999). Identification of in vivo
substrates of the chaperonin GroEL. Nature *402*, 147-154.

Kerner, M.J., Naylor, D.J., Ishihama, Y., Maier, T., Chang, H.C., Stines, A.P., Georgopoulos, C., Frishman, D.,
Hayer-Hartl, M., Mann, M*., et al.* (2005). Proteome-wide analysis of chaperonin-dependent protein
folding in Escherichia coli. Cell *122*, 209-220.

Knoblauch, N.T., Rüdiger, S., Schönfeld, H.J., Driessen, A.J., Schneider-Mergener, J., and Bukau, B. (1999).
Substrate specificity of the SecB chaperone. The Journal of biological chemistry *274*, 34219-34225.

Lin, Z., Madan, D., and Rye, H.S. (2008). GroEL stimulates protein folding through forced unfolding. Nat
Struct Mol Biol *15*, 303-311.

Mapa, K., Tiwari, S., Kumar, V., Jayaraj, G.G., and Maiti, S. (2012). Information encoded in non-native states
drives substrate-chaperone pairing. Structure *20*, 1562-1573.

Mattoo, R.U., Farina Henriquez Cuendet, A., Subanna, S., Finka, A., Priya, S., Sharma, S.K., and Goloubinoff,
P. (2014). Synergism between a foldase and an unfoldase: reciprocal dependence between the
thioredoxin-like activity of DnaJ and the polypeptide-unfolding activity of DnaK. Frontiers in molecular
biosciences *1*, 7.

McCarty, J.S., and Walker, G.C. (1994). DnaK mutants defective in ATPase activity are defective in negative
regulation of the heat shock response: expression of mutant DnaK proteins results in filamentation. J
Bacteriol *176*, 764-780.

Nagpal, S., Tiwari, S., Mapa, K., and Thukral, L. (2015). Decoding Structural Properties of a Partially
Unfolded Protein Substrate: En Route to Chaperone Binding. PLoS computational biology *11*, e1004496.

Niwa, T., Fujiwara, K., and Taguchi, H. (2016). Identification of novel in vivo obligate GroEL/ES substrates
based on data from a cell-free proteomics approach. FEBS Lett *590*, 251-257.

Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C., and Ferrin, T.E.
(2004). UCSF Chimera--a visualization system for exploratory research and analysis. J Comput Chem *25*,
1605-1612.

Rüdiger, S., Germeroth, L., Schneider-Mergener, J., and Bukau, B. (1997). Substrate specificity of the DnaK
chaperone determined by screening cellulose-bound peptide libraries. The EMBO journal *16*, 1501-1507.

Sharma, S., Chakraborty, K., Muller, B.K., Astola, N., Tang, Y.C., Lamb, D.C., Hayer-Hartl, M., and Hartl, F.U.
(2008). Monitoring protein conformation along the pathway of chaperonin-assisted folding. Cell *133*, 142-
153.

Tiwari, S., Kumar, V., Jayaraj, G.G., Maiti, S., and Mapa, K. (2013). Unique structural modulation of a non-
native substrate by cochaperone DnaJ. Biochemistry *52*, 1011-1018.

Tokuriki, N., Stricher, F., Serrano, L., and Tawfik, D.S. (2008). How protein stability and new functions trade
off. PLoS computational biology *4*, e1000002.

Verma, K., Saxena, K., Donaka, R., Chaphalkar, A., Rai, M.K., Shukla, A., Zaidi, Z., Dandage, R., Shanmugam,
D., and Chakraborty, K. (2020). Distinct metabolic states of a cell guide alternate fates of mutational
buffering through altered proteostasis. Nature Communications *11*, 2926.

Xu, Z., Horwich, A.L., and Sigler, P.B. (1997). The crystal structure of the asymmetric GroEL-GroES-(ADP)7
chaperonin complex. Nature *388*, 741-750.

Yang, F., Moss, L.G., and Phillips, G.N., Jr. (1996). The molecular structure of green fluorescent protein.
Nat Biotechnol *14*, 1246-1251.

Zhao, L., Vecchi, G., Vendruscolo, M., Körner, R., Hayer-Hartl, M., and Hartl, F.U. (2019). The Hsp70
Chaperone System Stabilizes a Thermo-sensitive Subproteome in E. coli. Cell reports *28*, 1335-
1345.e1336.

FIGURE 1

FIGURE 2

ΔH# (Kcal/mole)
wt GFP   +9.2±0.4
sGFP   -0.4 ±1.4

Figure 3

FIGURE 4

A

B

Trypsin Digestion

Without intramolecular disulfide

Trypsin Digestion

With intramolecular disulfide

Expected S-S peptide mass 3555.8298 amu

C

Trypsin (-DTT)

Disulfide

LTLEFICTTGKLPVPWPTLVTTFGYGVQCFAR

m/z = 4
Observed mass =889.9647(4)
Expected mass= 889.96(4)

D

+DTT+Iodoacetamide+Trypsin

LTLEFICTTGK   LPVPWPTLVTTFGYGVQCFAR

m/z = 2
Observed mass = 641.8467(2)
Expected mass= 641.8392 (2)

E

Spontaneous
+DTT -DTT

70KD
50KD

30KD
25KD

anti-GFP

F

Unfolded (U) → Intermediate1 (I_1) → $2\times10^{-3}(s^{-1})$ → Folded (F)

Intermediate1* (I_1*) → Oxidising → Terminally Misfolded Intermediate (I_2)

FIGURE 5

FIGURE S1

A. GFP/mCherry of different GroEL/ES dependent clones isolated by FACS

B. Native baseline

C. Fluorescence of GFP and mCherry channel upon expressing sGFP in the presence and absence of GroEL/ES overexpression

G. Fluorescence of GFP and mCherry channel upon expressing sGFP in the presence and absence of chaperones

D.

Ponceau (Loading control)

E.

sGFP   sGFP+GroEL/ES OE

GroEL 57 kDa

GFP 26 kDa

F. Fluorescence of GFP and mCherry channel upon expressing sGFP in the presence and absence DnaK/J/E overexpression

Sequence alignment of 6 GFP mutants with Wt GFP sequence

H.

```
Wt MSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTLKFICTT
M1 MSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTLKFICTT
M2 MSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTLKFICTT
M3 MSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTLKFICTT
M4 MSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTLKFICTT
M5 MSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYDKLTLKFICTT
M6 MSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTLKFICTT

Wt GKLPVPWPTLVTTFGYGVQCFARYPDHMKQHDFFKSAMPEGYVQERTIFF
M1 GKLPVPWPTLVTTFGYGVQCFARYPDHMKQHDFFKSAMPEGYVQERTIFF
M2 GKLPVPWPTLVTTFGYGVQCFARYPDHMKQYDFFRSAMPEGYVQERTIFF
M3 GNLPVPWPTLVTTFGYGVQCYARYPDHMKQHDFFKSAMPEGYVQERTIFF
M4 GKLPVPWPTLVTTFGYGVQCFARYPDHMKQHDFFKSAMPEGYVQERTIFF
M5 GKLPVPWPTLVTTFGYGVQCFARYPDHMKQHDFFKSAMPEGYVQERTIFF
M6 GKLPVPWPTLVTTFGYGVQCFARYPDHMKQHDFFKSAVPEGYVQRTIFF

Wt KDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYNYNSHNV
M1 KDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYNYNSHNV
M2 EDDGDYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYNYNSHNV
M3 KDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYNYNSHNV
M4 KDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYNYNSHNV
M5 KDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYNYNSHNV
M6 KDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYNYNSHNV

Wt YIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHY
M1 YIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHY
M2 YIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHY
M3 YIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHY
M4 YIMADKQKNGIKVNFKIRQNIEDGSVQLADHYQQNTPIGDGPVLLPDNHY
M5 YIMADNQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHY
M6 YIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHY

Wt LSTQSALSKDPNEKRDHMVLLEFVTAAGITHGMDELYK
M1 LSTQSALSKDPNEKRYHMVLLEFVTAAGITHGMDELYK
M2 LSTQSALSKDPNEKRDHMVLLEFVTAAGITHGMDELYK
M3 LSTQSTLSKDPNEKRDHMVLLEFVTAAGITHGMDELYK
M4 LSTQSALSKDPNEKRDHMVLLEFVTAADITHGMDELYK
M5 LSTQSALSKDPNEKRDHMVLLEFVTAAGITHGMDEL
M6 LSTQSALSKDPNEKRDHMVLLEFVTAAGITHGMDELYK
```

FIGURE S2

A. Unfolding kinetics of sGFP in 6M GuHCl

B. Refolding kinetics of sGFP with single exponential fit and residuals

C. Raw spontaneous refolding traces of sGFP at different concentrations.

D. Normalized spontaneous refolding traces of sGFP at different concentrations.

——— 400nM   ——— 200nM   ——— 100nM   ——— 12.5nM

E. Refolding rate of sGFP and Wt GFP in increasing salt concentrations

FIGURE S3

A Schematic of an enthalpic barrier in folding funnel

Schematic of an entropic or diffusion-limited barrier in folding funnel (flatness factor)

B. GFP/mCherry ratio of sGFP with and without GroEL/ES over expression

C. Refolding kinetics of sGFP in the presence and absence of DnaK/J/E

D. Comparison of refolding rate of GroEL/ES and SR-EL/ES

FIGURE S4

A. Effect of CuCl$_2$ on spontaneous folding of sGFP in non-reducing conditions

B. Distance between the cysteine residues in the native state

C48

C70

24.2 Å

C. Strategy for detecting disulfide formed during sGFP refolding

D. Qualitative simplified energy land-scape comparing spontaneous refolding in oxidizing and reducing conditions

FIGURE S5

A. sGFP refolding in oxidizing conditions in the presence of different components of the GroEL/ES/ATP system

B. Refolding rate as a function of delay in addition of GroES/ATP to GroEL-bound sGFP

C. Refolding yield of GroEL-bound unfolded sGFP upon addition of GroES/ATP after different delay times

D. Spontaneous refolding rates of sGFP in different concentrations of DTT

E. Comparison of refolding rate of Wt GFP with and without DTT

Table S1. Fitted parameters from Arrhenius analysis of temperature dependent refolding reactions in different conditions as given in first column. The standard errors reported are errors obtained after fitting the triplicates of refolding rates obtained at different temperatures for each of the conditions.

Refolding Condition	$\Delta H^{\#}$ (kCal/mole)	Std error ($\Delta H^{\#}$)	$\Delta Cp^{\#}$ (kCal/K)	Std error ($\Delta Cp^{\#}$)	ρ (kCal/(mole.K))	Std error (ρ)
Spontaneous (+DTT)	-0.41	1.42	1.70	0.4505	5.88e-6	1.40e-5
GroEL/ES/ATP	6.20	0.31	-0.18	0.1	1.26	0.65
SREL/ES	6.06	0.30	-0.42	0.09	0.65	0.32
wtGFP (spontaneous)	9.26	0.41	-0.24	0.12	497.30	339.40

Table S2. Relative quantitation and enrichment of the peptide fragment LTLEFICTTGK that is obtained from the non-disulfide bonded sGFP in refolding reactions with and without DTT. Normalization of the area of the peptide Ion count (IC) is normalized with the area under Total Ion Count (TIC) of both the LS-MS runs.

Sample	Total TIC area	area of LTLEFICTTGK	TIC area normalized	Enrichment (+DTT/-DTT)
sGFP (spontaneous)	3.28E+10	4.74E+05	1.44E-05	---
sGFP+DTT (spontaneous)	1.36E+10	2.15E+07	1.58E-03	109.3