Rewiring of peatland plant-microbe networks outpaces species turnover

Bjorn J.M. Robroek[1,†], Magalí Martí[2,†], Bo H. Svensson[2], Marc G. Dumont[3], Annelies J. Veraart[1], Vincent E.J. Jassey[4]

[1]Aquatic Ecology & Environmental Biology, Institute for Water and Wetland Research, Faculty of Science, Radboud University Nijmegen, AJ 6525 Nijmegen, The Netherlands.

[2]Thematic Studies - Environmental Change, Linköping University, SE-581 83 Linköping, Sweden.

[3]School of Biological Sciences, Faculty of Environmental and Life Sciences, University of Southampton, SO17 1BJ Southampton, United Kingdom.

[4]Laboratoire d'Ecologie Fonctionnelle et Environnement, Université de Toulouse, CNRS, 31062 Toulouse Cedex, France.

[†]These authors contributed equally to this work

Author for correspondence: B.J.M. Robroek: b.j.m.robroek@ru.nl

Abstract   Enviro-climatological changes are thought to be causing alterations in ecosystem processes through shifts in plant and microbial communities; however, how links between plant and microbial communities change with enviro-climatological change is likely to be less straightforward but may be fundamental for many ecological processes. To address this, we assessed the composition of the plant community and the prokaryotic community –using amplicon-based sequencing– of three European peatlands that were distinct in enviro-climatological conditions. Bipartite networks were used to construct site-specific plant-prokaryote co-occurrence networks. Our data show that between sites, plant and prokaryotic communities differ and that turnover in interactions between the communities was complex. Essentially, turnover in plant-microbial interactions is much faster than turnover in the respective communities. Our findings suggest that network rewiring does largely result from novel associations between species that are common and shared across the networks. Turnover in network composition is largely driven by novel interactions between a core community of plants and microorganisms. Taken together our results indicate that plant-microbe associations are context dependent, and that changes in enviro-climatological conditions will likely lead to network rewiring. Integrating turnover in plant-microbe interactions into studies that assess the impact of enviro-climatological change on peatland ecosystems is essential to understand ecosystem dynamics and must be combined with studies on the impact of these changes on ecosystem processes.

Keywords   microbial and plant diversity, plant-microbe interactions, bipartite networks, 16S rRNA, 16S amplicon sequencing, peatlands.

Introduction

Throughout the present interglacial period, northern peatlands have acted as carbon (C) sinks, resulting in over 500 Gt of C (Yu et al. 2010) to be locked up in these ecosystems. Recently it has even been suggested that northern peatlands may contain twice as much C (Nichols & Peteet 2019), i.e. > 1000 Gt, yet these numbers are debated (Ratcliffe et al. 2020). Nevertheless, maintaining a positive carbon balance (C uptake > C release) in peatlands is of great relevance to mitigate carbon-climate feedbacks. The carbon sink function of peatlands largely depends on the imbalance between gross primary production and decomposition. In undisturbed peatlands, the balance of these two processes is largely in favour of primary production. Indeed, decomposition rates in natural peatlands are low. Peatland plants, and *Sphagnum* mosses in particular, produce recalcitrant litter (Clymo 1965, Dorrepaal et al. 2005) and release anti-microbial compounds (Fudyma et al. 2019, Hamard et al. 2019), which together with acidic and anoxic environmental conditions impede microbial breakdown of organic matter and thus facilitate the accumulation of plant remains. Hence, the C sink function of peatlands is controlled by an interplay between abiotic and biotic factors.

Direct effects of environmental change on peatland ecosystem function have been studied intensively. Examples of these direct effects include changes in nitrogen deposition (Aldous 2002, Bragazza et al. 2006, Olid et al 2014), drought (Fenner & Freeman 2011; Estop-Aragonés et al. 2016), and warming (Dorrepaal et al. 2009, Olid et al 2014, Wilson et al. 2016), amongst others, which have been linked to carbon loss. Environmental change can however also indirectly affect ecosystem function, through alterations in the biotic composition of ecosystems (Walther et al. 2002, Pimm et al. 2014). Such changes in community composition more often than not have consequences for ecosystem processes (Bardgett et al. 2013, Rillig et al. 2019). Indeed, the effects of warming, drought and nutrient deposition on carbon cycling in peatlands have previously shown to manifest via changes in the composition of the vegetation (Limpens et al. 2003, Dieleman et al. 2015, Rupp et al. 2019). However, the effect of environmental change on carbon cycling can also be mitigated by changes in the vegetation through plant-controlled metabolomic and biogeochemical mechanisms that protects C loss during drought (Wang et al. 2015; Fenner & Freeman, 2020). Following the train of thought that changes in environmental conditions cause shifts in the composition of peatland plant communities

(Vitt & Slack 1975, Robroek et al. 2017, Norby et al. 2019), convergent shifts in the microbial

66   community can be expected. Indeed, peatland microbial community structure and activity are strongly

        connected to plant community assemblage (Bragina et al. 2014, Chronakova et al. 2019, Martí et al.

68   2019b, Ivanova et al. 2020). Hence, interactions between plant and microorganisms are the

        cornerstone that shape carbon-related processes in peatlands (Lindo et al. 2013) and it is critical that

70   we understand how enviro-climatological change affects these interactions to forecast the

        consequences for ecosystem processes (Kostka et al. 2016).

72   Most species do not exist in isolation but are embedded in trophic, non-trophic, mutualistic or

        antagonist interaction networks (Tylianakis et al. 2008, Bascompte 2009, Kéfi et al. 2016). In these

74   networks, the number of species and interactions vary depending on species identity, functional traits

        present and enviro-climatologic niche (Schleuning et al. 2016). Network assemblage, in turn, can

76   affect the resilience and stability of the network, which ultimately underlies the robustness of

        ecosystem functions to enviro-climatic change (Allesina et al. 2009, Morriën et al. 2017, de Vries et

78   al. 2018). The effects of changes in plant community composition on soil microbial communities, are

        however thought to be dependent on the environmental context (De Vries et al. 2012) – but see Fanin,

80   et al. (2019) – , leading to complex responses of plant-microbial interactions to changes in

        environmental conditions (Hagedorn et al. 2019). Adding to this complexity, recent work in peatlands

82   has shown that plant and soil communities exhibit non-linear responses to changing environmental

        conditions (Jassey et al. 2018, Lamentowicz et al. 2019), yet no studies have tested the co-response of

84   peatland plant and microbial communities to environmental change. To overcome this knowledge

        gap, it is essential to gain understanding on how environmental conditions regulate plant-microbe

86   associations (i.e. networks) and how this relates to within-community responses (i.e. plant and

        microbial communities). Such knowledge would ultimately lead to improved understanding on the

88   effects of environmental change on peatland processes.

            To assess how enviro-climatological conditions affect plant-microbe interactions, we

90   characterized plant and microbial communities in three European *Sphagnum*-dominated peatlands.

        We assessed how enviro-climatological conditions relate to compositional differences in plant and

92   16S rRNA-derived microbial communities (i.e. occurrence of taxonomic units with active function)

and how this affects plant-microbial networks, including network topology. We postulate that enviro-

94   climatic change will trigger both communities to turnover, leading to changes in the network

structure. The influence of enviro-climatologic conditions on plant-microbe associations can manifest

96   in various ways (Fig. 1): (i) no difference in biotic communities, nor in network associations, (ii) no

difference in biotic communities but a rewiring in network association with altered enviro-

98   climatological conditions, (iii) differences in the composition in one or both communities with

congruent differences in network associations, and (iv) differences in biotic community compositions

100   with a disproportionate rewiring of interaction networks. Congruent with the effects of environmental

conditions on peatland plant communities (Robroek et al. 2017), we expect that differences in plant-

102   microbe network structure are largely driven by turnover in one or both communities.

104   **Figure 1.** Examples of potential, i.e. hypothesised, effects of enviro-climatological change on associations
between species in the plant community and 'species' in the microbial community: plant-microbe networks.
106   Effects of enviro-climatological change (environmental filter) on network structure (pre-change network) are
postulated to manifest in four potential ways: (i) no alterations in post-change biotic communities, nor in
108   network associations; (ii) no alterations in biotic communities but a rewiring in network associations [depicted
as red lines]; (iii) alterations in the composition in one or both communities [depicted as changes in composition
110   of coloured symbols] with congruent alterations in network associations, and (iv) alterations in biotic
community compositions with a disproportionate rewiring of interaction networks.

Material and Methods

114   *Site description and enviro-climatic data.*    Three geographically distinct European *Sphagnum-*

dominated peatlands were selected for this study. Degerö Stormyr (64°88′N, 19°56′E, 277 m above

116   sea level (asl)), Sweden, is a minerogenic peatland with ombrotrophic elements. Cena Mire (56°25'N,

23°81'E, 12 m asl), Latvia, is an ombrotrophic raised bog. Dosenmoor, (54°95'N, 10°02'E, 28 m asl),

118   Germany, is a restored ombrotrophic raised bog. These sites vary in degree of conservation status, but

all sampling locations were located in an ombrotrophic part of each peatland with vegetation

120   characteristic for natural peatlands. Full vegetation characteristics of the sites can be found on

https://doi.org/10.5061/dryad.g1pk3.

122    We collected variables that describe climatic and environmental conditions in the selected

peatlands previously being shown to be the most important drivers for the turnover in peatland plant

124   community composition (Robroek et al. 2017). Four bioclimatic variables (mean annual temperature,

temperature seasonality, mean annual precipitation, precipitation seasonality) were extracted from the

126   *WorldClim* database (Hijmans et al. 2005), and averaged over a 5-year (2005-2009) period that

preceded our sampling campaign. *WorldClim* (http://www.worldclim.org/bioclim) calculates

128   temperature seasonality as the standard deviation of the mean monthly temperature. Larger standard

deviations refer to a larger variability in monthly temperature. Precipitation seasonality is a measure

130   of the variation in monthly precipitation totals and calculated as the standard deviation of the monthly

precipitation estimates expressed as a percentage of the mean of those estimates (i.e. the annual

132   mean). Atmospheric deposition data were produced using the EMEP (European Monitoring and

Evaluation Programme)-based IDEM (Integrated Deposition Model) model (Pieterse et al. 2007) and

134   consisted of grid cell (50 × 50 km) averages of total nitrogen and sulphur deposition. pH was

measured (Mettler Toledo MP220 pH meter) in five wells in each peatland site. These wells were

136   situated on the lawn-hummock interface.

138   *Field sampling of plant and microbial communities.*    Abundance data for all vascular plant and

bryophyte species from randomly selected lawn (n =3) and hummock (n = 3) microhabitats (0.25 m^2

140   quadrats; six in total) were collected in the summer of 2010. Microhabitats were sampled in pairs,

with each lawn-hummock pair separated < 2 m. This sampling strategy was chosen as to represent the

142 nature of the ombrotrophic part of the peatland as much as possible. Vascular plants and *Sphagnum* mosses were identified to the species level. Non-*Sphagnum* bryophytes were identified to the family

144 level. Rarefaction analysis indicated that our sampling adequately captured species richness in our peatlands.

146    In all plots (3 sites, 2 microhabitats, 3 replicates; 18 in total), we collected a 5 cm^3 peat sample from the acrotelm – defined as the peat layer between the peatland surface and the lowest

148 water table level, which largely contains living plants – and the catotelm – the peat layer that underlies the acrotelm, is largely anoxic, and contains mostly dead plant remains (Clymo 1984).

150 Hence, we collected a total of 36 samples from 18 plots. Sampling depth for the acrotelm samples was 5 cm below the surface, while sampling depth of the catotelm samples was 12.5 cm and 37.5 cm

152 below the surface for the lawns and hummocks, respectively (Marti et al. 2015; Suppl. Fig. 2). Sampling was conducted using a 10-cm wide Holmen auger, which was carefully cleaned with 70%

154 alcohol between sampling plots. Samples were refrigerated at 4°C, transported to the laboratory, and stored at -20°C prior to RNA extraction.

RNA isolation, amplicon sequencing and data handling.   Total RNA was extracted from 0.75

g of wet peat (7.8% ± 2.5 dry peat) using the FastRNA®Pro Soil-Direct Kit and FastPrep® Instrument (MP Biomedicals, Santa Ana, CA) as per the manufacturer's instructions. Quality and

concentration were determined on a Qubit fluorometer (Invitrogen, Eugene, OR, USA). Extracted RNA was reverse transcribed to complementary DNA (cDNA) using the lllustraTM Ready-to-Go RT-

PCR Bead (GE-Healthcare, Uppsala, Sweden). Briefly, 2 μl of the extracted RNA and 2.5 μl of the random primer pd(N)6 (1.0 μg/μl) (GE-Healthcare, Uppsala, Sweden) were mixed, denatured at 97°C

for 5 min, and then chilled on ice. One IllustraTM Ready-to-Go RT-PCR Bead and RNAse-free water were added to a final volume of 50 μl. Next, reverse transcription, consisting of a chain of 30 mins at

42°C followed by 5 mins at 95°C was used to create cDNA. RNA and cDNA were stored at -80°C and -20°C, respectively.

A 485 base pair (bp) fragment flanking the V3-V4 region of prokaryotic 16S rRNA was amplified using the primers 341F (5'-CCTAYGGGRBGCASCAG-3') and 806R (5'-

GGACTACNNGGGTATCTAAT-3') modified from (Yu et al. 2005). Each 25-µl PCR mixture contained one Illustra PuReTaq Ready-To-Go PCR Bead (GE Healthcare, Uppsala, Sweden), 0.5 µM

of each primer, 5 µl of cDNA and RNAse-free water. The 35 PCR cycles, performed on a MyCycler™ Thermal Cycler (Bio-Rad Laboratories), consisted of 90 s at 95°C, 30 s at 95°C, 30 s at

56°C, 30 s at 72°C and a final 5 mins at 72°C. PCR products were verified on 1% agarose gel and purified with a GeneJET PCR Purification Kit (Fermentas, Vilnius, Lithuania). Purified PCR product

concentrations were determined using the Quant-iT dsDNA HS Assay Kit and the Qubit fluorometer (Invitrogen) and adjusted to 5 ng µl^{-1}. To link sequence reads to samples, a second PCR on the initial

product was performed using the primer pair 341F/806R with adapters and a unique barcode. The number of cycles was reduced to 10. Bands of ca. 530 bp were cut out and purified by High Pure PCR

Cleanup Micro Kit (Roche Diagnostics GmbH, Mannheim, Germany) as per the manufacturer's instructions. These fragments were quantified using a Qubit™ dsDNA HS Assay Kit and the Qubit™

fluorometer (Invitrogen) and qPCR (Mx-3000, Stratagene). The sample amplicons were mixed to equal concentrations (4×10^{5} copies µl^{-1}) and subjected to two-region 454-pyrosequencing on a $70 \times$

75 GS PicoTiterPlate (PTP) using a GS FLX pyrosequencing system.

      Sequence data were sorted, trimmed, filtered and quality checked ($QS \geq 25$ for trimming, and

186   lengths ≥ 150 bp) using the Qiime 1.6.0 data analysis pipeline (Caporaso et al. 2010). Chimeras were detected and removed using UCHIME in reference mode with the MicrobesOnline gold set (Edgar et

188   al. 2011). Briefly, operational taxonomical units (OTUs) were selected at 97% identity with a confidence threshold of 50%. Representative sequences were selected randomly and classified against

190   the SILVA database version 132 (https://www.arb-silva.de/). Unifrac distances for beta diversity were calculated from trees generated with Fasttree. The input multiple sequence alignments were generated

192   using PyNAST against gg2011 template alignments. From the 36 libraries, we obtained a total of 772,742 sequences with an average of 21,465 sequences per library, except for one sample from Cena

194   Mire –lawn, acrotelm– which had 135 reads and was therefore removed from the analysis. OTUs that were present in fewer than 20% of the samples were removed from the analyses as well as OTUs

identified as chloroplast. A total of 464,327 sequences were left, corresponding to 4707 OTUs. From those, 74% (3690) of the OTUs corresponded to bacterial taxa and 20% (141) to archaeal taxa.6% of the OTUs were unclassified.

Plant-prokaryote bipartite networks.   Bipartite networks (i.e. a network with two distinct sets of nodes, plant nodes and prokaryotic nodes, with associations only defined between nodes from different sets) were used to detect co-occurrence between plants and prokaryotes as a proxy for site-specific plant-prokaryote associations (Faust & Raes 2012). Co-occurrence networks were computed at site level. Hence data for the hummock and lawn microhabitats were pooled; however, separate networks were constructed to link the plant community with the acrotelm and with the catotelm prokaryote communities. Hellinger-transformed matrices, comprising data of bacterial and archaeal OTUs, vascular plants and *Sphagnum* species, were used to calculate Spearman's correlation coefficients for plant-prokaryote associations (Berry & Widder 2014). Significance for each permutation was assessed using a permutation test with 10,000 repetitions. Only correlations > 0.6 (positive and negative) were adopted in the co-occurrence matrix (Sander et al. 2017). Each co-occurrence matrix was transformed into an adjacency (binary) matrix based on the presence or absence of links. Networks were then produced using the *igraph* and *bipartite* R packages (references in Supplementary), where each network comprised positive and negative plant-prokaryote associations. Each network was further analysed in terms of topology, and network dissimilarity (Berry & Widder 2014, Williams et al. 2014).

First, we evaluated the topological attributes of the networks using a set of indices focussed on species interactions, which were calculated for interaction between the plant and prokaryotic communities. *Effective partners* was defined as the number average of links for individuals in a specific network. *Partner diversity* is the diversity (PD; Shannon-based) of interactions between species from the two levels of the network. *Partner specificity* (PS) was calculated as the coefficient in variation of interactions, normalised to values between 0 and 1 (Poisot et al. 2012b), where PS = 0 indicates low specificity and PS = 1 indicates high specificity. These three indices were calculated from a top-down and bottom up perspective. Top-down accounts for the number of interactions plant

224   species have with prokaryotic 'species'. Bottom-up accounts for the number of interactions

prokaryotes have with species in the plant community. Essentially, in a network top-down and bottom

226   up networks do not necessarily have to match (Suppl. Fig. 3). Next, we identified network attributes

using a set of plant-prokaryote interaction indices. *Linkage density* (LD) is the marginal total-

228   weighted diversity of interactions between species. *Generality* of the plant community (G_p) is

calculated as the mean effective number of prokaryotic 'species' per plant species weighted by the

230   total number of interactions. It describes the degree of specialization in the plant community. Low

values indicate a high degree of specialization. *Generality* for the prokaryotic community (G_m) is

232   calculated as the mean effective number of plant species per prokaryotic 'species' weighted by the

total number of interactions, and represents the degree of specialization in the prokaryotic community.

234   *Robustness* (R) is a measure of robustness of the network to species loss, calculated as the area under

the extinction curve of the network. The curve, essentially, describes the dependency –for survival– of

236   species from one level on species of another level. R = 1 corresponds to a situation where most

prokaryotes survive while most plant species are eliminated, or vice versa. R = 0 corresponds to a

238   situation where species loss form one level results in the loss of most to all network connections.

Partner diversity in the networks (PDn) is the diversity (Shannon-based) of the number of interactions

240   for the species in each level of the bipartite network. *Species richness* (SR) is the number of species in

the bipartite network community. *Niche specialization* (N) represents the site scores from a NMDS

242   analysis, and represent the niche characteristics of each community (Devictor et al. 2010). The indices

G, R, PD SR and N were calculated separately for the two levels –plants and prokaryotes– in the

244   bipartite networks.

246   *Network turnover.*   To test if enviro-climatic conditions affected species turnover and turnover in

bipartite networks, we calculated pairwise β-diversity in species composition and plant-prokaryote

248   networks in each site using the methods proposed by Legendre & De Cáceres (2013) and Poisot et al.

(2012a), respectively. For pairwise β-diversity in species composition, we used a site $\times$ species

250   abundance matrix. Total β-diversity (BD) was partitioned into species contributions (BD_S; degree of

variation of individual species in each site) and local contribution (BD_L; comparative indicators of the

252   ecological uniqueness of the sites) to β-diversity (Legendre & De Cáceres 2013). For β-diversity in

co-occurrence networks, we use a site × species pairs matrix (adjacency binary matrix). As

254   differences in networks can arise either through changes in species composition and/or realized

interactions between species, we specifically focused on the dissimilarities in interactions between

256   networks (BDN) that originates from differences in species interactions due to species turnover

(BDN_{NS}) or from novel interactions between common network species (BDN_{CS}) (Koleff et al. 2003,

258   Canard et al. 2012, Poisot et al. 2012a).

260   *Data handling and statistical analyses*. All statistical analyses were performed in *R* version 3.1.2

using the *vegan, diverse, phyloseq, nlme, dplyr and tidyr* packages (references in Supplementary).

262   Analysis of Variance tests were used to test for differences in pore water nutrients, plant species and

prokaryote richness, diversity (Shannon H'), evenness (Pielou's J), and plant-prokaryote network

264   indices between sites and sampling depth (for prokaryote data only). Prior to testing for turnover in

prokaryotic community composition across the different habitats (β-diversity), relative abundance of

266   each OTU was calculated within each sample.

We used redundancy analysis (RDA) to relate enviro-climatologic factors to plant and

268   prokaryotic communities (Hellinger-transformed matrices). We used *ClustOfVar* package in R to

reduce the enviro-climatological covariates of interest (i.e. mean annual temperature, temperature

270   seasonality, mean annual precipitation, precipitation seasonality, total nitrogen and sulphur

depositions), and select the most representative and least collinear variables. We further tested the

272   variance inflation factor among selected variables to test the independence of the variables in the

ordination space. According to *ClustOfVar*, the variables that best described environmental

274   differences between sites were seasonality in temperature, sulphur deposition and pH (Suppl. Fig. 3).

The robustness and significance of the RDAs was then tested using permutation tests.

276   We tested the relationships between network indices and the enviro-climatological conditions

that were returned by the RDA models as most significant – temperature seasonality and sulphur

278   deposition – using regression analyses. We grouped network indices into three categories: network

composition (linkage density, species richness, partner diversity), network structure (robustness, niche

280 specialization, modularity) and network generality (generality indices). Network indices were

previously standardized to allow direct comparison among the different categories.

Results

Mean annual temperature (MAT) was lowest in Degerö Stormyr and highest in Dosenmoor, while seasonality in temperature (TS) followed an opposite pattern (Table 1). Mean annual precipitation (MAP) varied from 710 mm in the Cena Mire to 760 mm in Dosenmoor. Seasonality in precipitation (PS) did not vary much between sites (Table 1). Total nitrogen and sulphur deposition were lowest in Degerö Stormyr. Total nitrogen deposition values in Dosenmoor were almost double as compared to those in Cena Mire. For sulphur, Cena Mire and Dosenmoor had similar deposition values (Table 1). pH in all sites was below 4.5 but differed slightly between sites (F = 4.5, P = 0.034) and was highest in Degerö Stormyr and lowest in the Dosenmoor (Table 1).

Table 1. Bioclimatic data for Degerö Stormyr, Cena Mire, and Dosenmoor. MAT = Mean annual temperature (°C), TS = Seasonality in temperature (%), MAP = Mean annual precipitation (mm), PS = Seasonality in precipitation (%); N_{tot} = Total nitrogen deposition (mg m^{-2} yr^{-1}), S_{tot} = Total sulphur deposition (mg m^{-2} yr^{-1}), and pH of the porewater (n=5). Different letters represent significant difference (P ≤ 0.05)

	MAT	TS	MAP	PS	N_{tot}	S_{tot}	pH
Degerö Stormyr	1.2	9.3	730	26	130	96	4.01 ± 0.15^a
Cena Mire	6.4	7.9	710	31	760	344	3.71 ± 0.17^{ab}
Dosenmoor	8.0	5.9	760	19	1260	6144	3.63 ± 0.29^b

Plant and prokaryote community composition.   Vascular plant and bryophyte species richness were highest in Cena Mire, while evenness in this site was lowest (Table 2). Detected archaeal richness did not differ among the three sites. Richness of the acrotelm archaeal communities was at least three times lower as compared to richness of the catotelm archaeal communities (Table 2). Bacterial richness, opposite to the patterns observed in the plant community, was lowest in Cena Mire, for both acrotelm and catotelm (Table 2).

Redundancy analysis (RDA) showed that all communities (plants, acrotelm prokaryotes, and catotelm prokaryotes) well separated in the ordination space (Suppl. Fig. 4). RDA analysis explained 28% (vascular plants), 7.2% (prokaryote, acrotelm) and 19% (prokaryote, catotelm) of the variance (adjusted R^2) of each community. Sulphur deposition and temperature seasonality both explained much of the differences in community structure for all three communities (P ≤ 0.05), while pH was

not significant. Notably, the three communities showed similar patterns in their species-environment

310   relationships. Communities in Degerö Stormyr were related to high seasonality in temperature, and

communities in Dosenmoor were related to high sulphur deposition (Suppl. Fig. 4).

Table 2. Plant species and prokaryote richness (OTUs) and evenness (Pielou's J). Values are means ± standard
314   errors. Prokaryotic richness and evenness have been described for the acrotelm and catotelm communities
separately. Different letters represent significant difference (P ≤ 0.05) in richness or evenness data for the sites
316   as tested by analysis of variance within four sub-communities (vascular plants, bryophytes, archaea and
bacteria).

Plant community		Richness	Evenness		
Vascular plants	Degerö Stormyr	6.5 ± 1.23^{b}	0.8 ± 0.05^{a}		
	Cena Mire	12.0 ± 1.94^{a}	0.2 ± 0.05^{b}		
	Dosenmoor	7.2 ± 0.40^{b}	0.8 ± 0.06^{a}		
Bryophytes	Degerö Stormyr	1.8 ± 0.17^{b}	0.5 ± 0.15		
	Cena Mire	3.3 ± 0.56^{a}	0.5 ± 0.12		
	Dosenmoor	1.7 ± 0.21^{b}	0.7 ± 0.12		
Prokaryotic community		Acrotelm (oxic)		Catotelm (anoxic)	
		Richness	Evenness	Richness	Evenness
Archaea	Degerö Stormyr	14 ± 2.0	0.8 ± 0.05	51 ± 4.5	0.4 ± 0.06
	Cena Mire	30 ± 13.3	0.6 ± 0.08	74 ± 7.0	0.4 ± 0.05
	Dosenmoor	38 ± 14.2	0.7 ± 0.05	84 ± 11.9	0.5 ± 0.06
Bacteria	Degerö Stormyr	1367 ± 95.0^{a}	0.8 ± 0.00	1511 ± 99.4^{a}	0.8 ± 0.01^{a}
	Cena Mire	1231 ± 109.0^{b}	0.8 ± 0.02	838 ± 105.9^{b}	0.7 ± 0.04^{b}
	Dosenmoor	1615 ± 88.3^{a}	0.8 ± 0.01	1246 ± 139.2^{a}	0.7 ± 0.01^{a}

320   Across sites, and pooled for the acro- and catotelm communities, bacterial to archaeal ratios

exceeded 3:1 (Fig. 2 a). Acidobacteria (35%), Proteobacteria (29%), Verrucomicrobia (13%), and

322   Actinobacteria (9%) were the most abundant bacterial phyla across sites (Fig. 2 b). Relative

abundances in bacterial phyla were similar across sites. Composition of the archaeal community, on

324   the other hand, seemed site-specific (Fig. 2 c). In Degerö Stormyr, methanogens of the order

Methanomicrobiales (50%) and Methanobacteriales (37%) were the dominant archaeal orders.

326   Archaeal communities in Cena Mire were dominated by one single order, Methanomicrobiales (83%).
In Dosenmoor, Thermoplasmata (39%) and an uncultured archaeon (28%) likely associated to the
328   phyla Thaumarchaeota and Crenarchaeota were most dominant. Of the classified orders in this site,
Nitrosotaleales (30%), Methanomicrobiales (22%) and Methanosarcinales (19%) had highest
prevalence.

Figure 2. Comprehensive view of the sequence content of peat soil prokaryotic libraries per site: DES = Degerö Stormyr, CEN = Cena Mire, DOS = Dosenmoor. Segments that compose each bar represent the total number of
sequences normalised to the total number of sequences in the libraries per site. a) View of the relative abundance of bacteria (blue) and archaea (red) in the site-specific communities. b) View of the relative
abundance of bacterial 'species' in the respective site-specific communities. Each segment represents identified bacterial phyla. c) View of the relative abundance of archaeal 'species' in the respective site-specific
communities. Each bar segment represents identified archaeal orders.

Network structure.   The number of plant-prokaryote pairs in bipartite networks differed between

the three sites (site effect, $F_{1,2} = 65$, $P = 0.01$; site x depth, $F_{1,2} = 2.2$, $P = 0.27$), with > 70% fewer

342   interactions in Cena Mire and Dosenmoor as compared to Degerö Stormyr (Fig. 3). A lower number

of bipartite interactions results from lower numbers of both positive and negative associations (Fig.

3).

Figure 3. Bipartite network visualisation for the three European peatlands. Indicated are interactions between the plant community and acrotelm prokaryotic community (left) and between the plant community and catotelm
prokaryotic community below (right). Plants are visualised at the top of the networks and coloured dark and light green for vascular plants and bryophytes, respectively. Prokaryotes are illustrated on the bottom of the
networks and follow the colour scheme as in Fig.1. A complementary list of prokaryotes and their classification can be found in Suppl. Table 1. Size of the bars scales with relative abundance in the respective communities.
For all sites, networks are filtered to only visualise interactions with 'keystone' prokaryotes, defined as those that had > 5 interactions in Degerö Stormyr. Interaction lines represent Spearman's rank correlations > 0.6. Dark
grey interactions indicate vascular plant-prokaryote interactions, light grey indicates bryophyte-prokaryote interactions. Inset numbers indicate the full number of bi-partite (bold) interactions, further broken down in the
number of positive / negative interactions.

In the vascular plant community, only *Rhynchospora alba* could be identified to play a major role in

all plant-prokaryote networks, except in the Dosenmoor catotelm network (Fig. 3). In the moss

360 community, no such key species seems present (Fig. 3). OTUs classified as belonging to the families Acidobacteriaceae (Bac 4), WD2101 soil group (Bac 11), Acidothermaceae (Bac 12),

362 Pedosphaeraceae (Bac 13), and Methylacidiphilaceae (Bac 14) were present in all three networks of the acrotelm prokaryotic communities (Fig. 3, Suppl. Table 1). In the catotelm prokaryotic

364 communities, bacteria in the families Koribacteraceae (Bac 22), Acetobacteraceae (Bac 33), and Pedosphaeraceae (Bac 34) were important network species (Fig. 3, Suppl. Table 1).

For all sites, β-diversity for bipartite (plant-prokaryote) networks (BDN) exceeded β-diversity for

community composition (BD), indicating that network turnover is higher than species turnover between sites ($P \leq 0.001$; Fig. 4). Indeed, network compositions (i.e. pairwise comparison between

networks) between sites were more dissimilar (acrotelm, BDN = 0.91; catotelm, BDN = 0.89) than species compositions (i.e. pairwise comparison between communities) between sites (acrotelm, BD =

0.32; catotelm, BD= 0.31). Turnover in species composition between sites, β-diversity (BD), was equally driven by species and local contributions (BD_S and BD_L). Network β-diversity (BDN) was

mostly driven by turnover in interactions (both direction and number) between species that are common in the three sites (BDN_{CS}), rather than by novel interactions driven by changes in the

community composition (BDN_{NS}; Fig. 4).

Figure 4. Turnover in species composition (species β-diversity) and association (network β-diversity) and their respective components across three peatlands (see methods). Red circles indicate plant and microbial

communities in the acrotelm. Blue circles indicate plant and microbial communities in the catotelm.

Differences in the number of bi-partite interactions between sites seem to relate to differences in the

numbers of effective partners between the respective communities, which were always highest in

384   Degerö Stormyr (acrotelm: $P < 0.001$; catotelm: $P < 0.001$; Fig. 5 a, b). This is further supported by

differences in diversity in plant-prokaryotic partners, which were higher in Degerö Stormyr as

386   compared to Cena Mire and Dosenmoor (acrotelm: $P < 0.001$; catotelm: $P < 0.001$, Fig 4 c, d).

Moreover, the number, but also diversity, of prokaryotic OTUs that are associated (i.e. effective

388   partners) to specific plant species (top-down links) were higher than the number of plant species that

associate with a specific OTU in the prokaryote community (bottom up links). Essentially this means

390   that plant species link to a higher number of prokaryotic species, than prokaryotes to plants. Despite

the relative low number of links of prokaryotes with plants, we found a larger specificity of these

392   links, as compared to the specificity of plants in their association with prokaryotes (acrotelm: $P <$

0.001; catotelm: $P < 0.001$). Partner specificity was lowest in Degerö Stormyr for both levels of

organisation in the bi-partite networks (acrotelm: $P < 0.001$; catotelm: $P < 0.05$, Fig. 5 e, f).

Figure 5. Boxplots of the species associations and network properties for the acrotelm (a, c, e) and catotelm (b, d, f) plant-prokaryote interactions. The number of specific partners (a, b), partner diversity (c, d) and partner

specificity (e, f) have been calculated for plants (white boxplots: top down links; Suppl. Fig. 2) and prokaryotes (grey boxplots: bottom up links; Suppl. Fig. 2). Horizontal solid lines indicate median values. Different letters

indicate differenced at the $P \leq 0.05$ level, based on analysis of variance using mean values.

At the network level, the overall linkage density (LD), generality (G), partner diversity (PD) and species richness (SR) in the bipartite networks was higher in Degerö Stormyr as compared to

Cena Mire and Dosenmoor; a pattern found in both acrotelm and catotelm bipartite networks (Suppl. Fig. 5). The bipartite networks, acro- and catotelm, in Degerö Stormyr were strongly sensitive to the

loss of prokaryotes (low robustness) but not to the loss of plants. Oppositely, bipartite networks in Cena Mire and Dosenmoor were more sensitive to the loss of plants as compared to prokaryote loss

(Suppl. Fig. 5). As compared to Degerö Stormyr, networks in Cena Mire and Dosenmoor show a high degree in niche specialization, especially in the plant communities (Suppl. Fig. 5). These differences

in network indices between the sites were driven mostly by temperature seasonality and sulphur deposition in a lesser extent (Fig. 6). While compositional network indices (linkage density, partner

diversity, number of effective partners) and generality indices increased with increasing temperature seasonality, structural network indices (robustness, niche extinctions, modularity) decreased. Opposite

patterns were found with increasing

sulphur deposition (Fig. 6). These results

416 indicate that plant-microbial interactions

change in complexity, specificity and

418 evenness in the distribution of species

interactions with changes in

420 environmental conditions, and that these

patterns are likely driven by the turnover

of species between peatland sites.

Figure 6. Linear regressions between the key enviro-climatological variables –total nitrogen

deposition and temperature seasonality– and network indices. Network indices were

grouped into three categories, network composition (linkage density, species

richness, partner diversity), network structure (robustness, niche specialization, modularity)

and network generality (generality indices). The grey area represents the 95% confidence

interval of the mean. Data on specific network

indices can be found in Suppl. Fig. 5.

Discussion

Using data on plant and prokaryotic community composition in three *Sphagnum*-dominated peatlands, we evidence that plant-prokaryotic networks are site-specific. As these peatlands are situated in three distinct environmental zones (Metzger et al. 2005) our data suggest that these networks are vulnerable to the effects of enviro-climatological change. Notably, our work indicates that turnover in plant-microbe links outpaces turnover in biotic composition of the plant and microbial communities that comprise the plant-microbe networks. This suggests a disproportionate rewiring of plant-microbe networks along an enviro-climatological gradient, as hypothesized in scenario iv (Fig. 1). Hence, a turnover in the plant community as shown in earlier research (Gunnarsson et al. 2002, Robroek et al. 2017, Pinceloup et al. 2020) will lead to complex changes in plant-microbe interactions that could affect ecosystem processes in the longer term (Morriën et al. 2017).

Turnover in community composition.   In line with earlier work where it was suggested that changes in enviro-climatological conditions in peatland ecosystems leads to a significant turnover in the composition of the plant community (Robroek et al. 2017), plant community composition was highly distinctive between sites. Moreover, our study confirms previous suggestions that enviro-climatological conditions are important drivers that underlie the composition of the vegetation, and that changes in these drivers result in alterations of the plant community (Gunnarsson et al. 2002, Pinceloup et al. 2020) and associated biotic interactions (Wiedermann et al. 2007). Indeed, we note that site-specificity of plant communities extends to the prokaryotic community, a pattern that is likely explained by the strong links between plant and microbial communities (Chronakova et al. 2019, Ivanova et al. 2020). Despite the variation in prokaryotic community composition between sites, the archaea to bacteria ratio remains strikingly stable, suggesting that both microbial groups are equally affected by alterations in enviro-climatological conditions.

Bacterial community composition was remarkably similar between sites. Archaeal communities, on the other hand, were highly dissimilar between the three peatland sites. Bacteria in the genera *Occallatibacter*, *Acidothermus*, *Verrucomicrobium,* bacteria in the Phycisphaera-related

464   group WD2101 (yet uncultured Planctomycetes of the order Tedisphaerales) and bacteria in the

family Methylacidiphilaceae (phylum Verrucomicrobia) occurred in all three acrotelm networks.

466   These genera have been previously found in soils, and have been characterised as strictly aerobic,

moderately acidophilic, and chemoorganotrophic mesophiles. Methylacidophilaceae include methane-

468   oxidizing genera, but there is no evidence that the members of this family present in peatlands are

involved in methane cycling (Ivanova et al. 2020). This underlies the fact that it can be difficult to

470   confidently predict the function of uncultivated bacterial species. The bacterium "*Candidatus*

Koribacter", a bacterium in the family Acidobacteriaceae, and a bacterium in the family

472   Pedosphaeraceae were important network species in catotelm prokaryotic communities. These results

mirror earlier findings, where bacterial community composition was reported to be rather stable under

474   experimental warming (Weedon et al. 2017). The archaeal communities in Degerö Stormyr and Cena

mire were dominated by methanogens. In these sites, biogeochemical drivers likely promoted a

476   classical anaerobic degradation of organic matter to methane in the absence of other terminal electron

acceptors. The dominating methanogens, Methanobacteriales and Methanomicrobiales, characteristic

478   for acidic peatlands, are hydrogenotrophic and form methane from hydrogen/CO_2 and/or formate

(Kotsyurbenko et al. 2007, Martí et al. 2015). Interestingly, the Methanomicrobiales family

480   *Methanoregulaceae* identified in the present study dominated the methanogen population in a

comparable set of peatlands along a nitrogen deposition gradient (Martí et al. 2015). Reductions in the

482   relative abundance of methanogens in Dosenmoor may be the result of higher deposition levels of

nitrogen and sulphur. The presence of oxidized forms of these elements, may favour nitrogen oxide

484   and sulphate reduction, resulting in increased competition for methanogens. This is supported by the

higher relative abundance of the phyla Nitrosotaleales and Thaumarchaeota (Suppl. Table 1), which

486   are known ammonia-oxidizing archaea in acidic environments (Lehtovirta-Morley et al. 2011, Lin et

al. 2015), in the Dosenmoor. Furthermore, Thermoplasmata – highly abundant in the Dosenmoor –

488   are suggested to be involved in the transformation of sulphite and possibly organosulphonates (Lin et

al. 2015). Given the high sulphur deposition in the Dosenmoor, these compounds prospectively are

490   abundant. Here, it should be stressed that relative abundance does not necessarily correspond to the

actual abundance and/or the physiological activity by the microorganisms. To exemplify, increased

492   methanogen activity has been shown in peatlands with high nitrogen deposition and low relative

abundance in methanogens (Martí et al. 2019a). Hence, the main part of the organic matter

494   degradation likely follows the methanogenic route, also at the Dosenmoor. In this context it should be

noted that the relative abundance might discriminate against prokaryotes at very low abundance,

496   which may have very high activity and thus process biogeochemical reactions at high speed as

exemplified for sulphur in peat soil (cf. Pester et al. 2010, 2012)

Turnover in plant-microbe network composition. Beyond differences in community composition, our

500   work shows that plant-microbe network structure differs between sites. Notably, and in line with

scenario iv in our hypotheses, turnover of plant-microbe interactions is disproportionate when

502   compared to the turnover in plant and prokaryote communities. These results would suggest that

changes in enviro-climatological conditions are key in driving the observed rewiring of plant-

504   prokaryote interactions. Interestingly, the highest numbers of effective partners and partner diversity

were found in the plant community. While prokaryotes have significantly lower numbers of plant

506   partners, their associations with plant species tend to be more specific. The observation that diversity

and numbers in top-down interactions (i.e. number of interactions plant species have with

508   prokaryotes) exceeds diversity and numbers in bottom-up interactions (interactions prokaryotes have

with species in the plant community) can simply be attributed to the fact that species richness in the

510   prokaryotic community exceeds plant species richness by a couple of orders of magnitude, which by

definition leads to higher diversity of plant-associated prokaryotes. Our analysis, however also

512   suggests that species in the soil prokaryotic community in peatlands form specific links to species in

the plant community (cf. Bragina et al. 2012, Hamard et al. 2019). Three non-exclusive scenarios can

514   explain the patterns we observe. First, a particular plant species provides resources or niche space for

a wide array of microbes. Second, a particular plant species under particular environmental conditions

516   produce and release specific metabolites that can attract, deter or kill targeted microbes. Third, these

microbes are highly specific to a particular plant (Suppl. Fig. 2). Taken together, our findings indicate

518   that plants are important drivers for plant-prokaryote interactions and is in line with the general idea

that the plant community has a direct effect on the prokaryotic species occurrences and community

520 composition and activity (Fisk et al. 2003, Wardle et al. 2004, Robroek et al. 2015, Yavitt & Williams

2015), but the opposite is not necessarily true.

522   Network size and complexity differ between sites and decrease from the most northern

(Degerö Stormyr) to the most southern (Dosenmoor) site. These findings are in line with earlier

524 findings where, as a response to warming, networks were observed to become less complex (Galiana

et al. 2014). Wei et al. (2013) have connected nutrient enrichment to a decoupling of grassland plant-

526 soil-microbe interactions and attribute this to differences in responses in the composition of plant and

microbial communities to changes in nutrient load. Hence, the observed decrease in network

528 complexity along the enviro-climatologic gradients in our study may indicate that plant-microbe

associations, and perhaps even feedbacks, go down with increased nutrient deposition and decreased

530 temperature seasonality.

A faster loss of network interactions as compared to species loss has earlier been suggested to

532 have major implications on ecosystem processes and services (Valiente-Banuet et al. 2014). Indeed,

increased network complexity has been connected to enhanced efficiency in soil carbon uptake, that is

534 increased C uptake by the soil microbial communities without substantial changes in microbial

biomass, even without major changes in the composition of plant communities (Morriën et al. 2017).

536 Our data suggest that seasonality in temperature and nutrient deposition modulate network indices

that are related to composition, structure and generality of plant-microbe networks. In other words,

538 our findings indicate that decreased seasonality and increased nutrient deposition, which reflect a

north-south gradient in our study, leads to decreases in network indices related to composition and

540 generality, but simultaneously increases structure related indices. A number of plant-prokaryotic

associations that were identified in bi-partite networks in Degerö Stormyr were persistent in the other

542 two sites despite differences in enviro-climatological different. These results would suggest a certain

resistance and robustness of plant-prokaryotic associations when facing radical environmental change

544 (Mandakovic et al 2018). Furthermore, while networks become less diverse and complex while

moving from the northern to the southern end of our enviro-climatologic gradient, network structure

546 becomes less prone to species loss. Essentially this means that we observe an increased level of

specialization in ecological networks, potentially caused by local adaptations that favour network

548   specialization (Dalsgaard et al. 2011, 2013) under harsh conditions (e.g.. increased nutrient load) for

peatland ecosystems. Such increase in specialization indicates an increased level of nestedness and

550   would in the long term weaken network stability to changing enviro-climatological conditions (de

Vries et al. 2018).

Network rewiring – new connections between common species. As correlation networks along

554   environmental gradients between vegetation and microbial communities may not necessarily be

causal, care should be taken in drawing firm conclusions from our data. Additionally, space-for-time

556   substitutions, as used in this study, have been reported to show a greater magnitude of response in

community turnover than climate manipulations or long-term monitoring studies (Elmendorf et al.

558   2015). Space-for-time approaches nevertheless are extremely useful for gauging the direction of

change of communities to decadal scales in enviro-climatological change (Elmendorf et al. 2015).

560   Despite these caveats, our findings show that while differences in environmental properties, such as

temperature and nutrient deposition, between sites markedly relate to differences in the composition

562   of plant and prokaryote, acro- and catotelm, communities, turnover in plant-prokaryote network

composition exceeds species turnover in either one of both communities. Specifically, we show that a

564   reshuffling of plant and microbial communities, most likely driven by enviro-climatological changes,

has resulted in a significant rewiring of plant-microbial interactions (scenario 4; Fig 1). This rewiring

566   was not just the result of changes in spatial patterns of diversity across plant and microbe trophic

levels (cf. Galiana et al. 2019). The rewiring between the peatland plant-microbe networks that we

568   have assessed was to a large degree caused by the creation of novel associations between species that

are common and shared across the sites. Hence, turnover in network composition is largely driven by

570   novel interactions between plants and microbes that are common across networks and to a lower

extent by species new to a metacommunity due to species turnover. These results would suggest that

572   changes in environmental conditions can lead to network rewiring, without clear and strong effects in

plant or microbial community per se. We propose that the next step is to link changes in plant-

574   microbe network structure to peatland carbon cycling and to identify key plant and microbial groups

that underpin the carbon uptake function of northern peatlands.

Data and materials availability   All data that support this manuscript are available through DANS, the Netherlands institute for permanent access to digital research resources (https://dans.knaw.nl/en). Nucleotide sequence accession numbers have been deposited in the European Nucleotide Archive (ENA) at EMBL-EBI under accession number PRJEB36623.

582 **Declarations**

Acknowledgments – We thank the Danish National High-Throughput DNA-Sequencing Centre (Copenhagen, Denmark) for using their sequencing facilities. We also thank the members of the PEATBOG project for valuable input in the planning phase of this project, and the landowners for field access. This paper is dedicated to late Dr Richard Payne, who tragically lost his life while attempting to climb Peak 6477 in the Indian Himalayas.

Funding – The BiodivERsA-PEATBOG project was funded as an ERA-net project within the European Union's 6th Framework Programme for Research funded though through the Dutch Research Council (NWO: 832.09.003), FORMAS (215-2008-1879) and the Swedish Research Council (3550141100). This specific project was supported by the British Ecological Society research grant (SR17\1427; BJMR), the Dutch Foundation for the Conservation of Irish bogs (BJMR) and the French National Research Agency (Mixopeat; grant number ANR☐17☐CE01☐0007; VEJJ). AJV is supported by a Mohrmann fellowship.

Author contributions – BJMR and BHS initiated the data collection. BJMR and MM led the field sampling. VEJ, MM and BJMR analysed all data, with help from MGD, AJV and BHS, and led the writing of the manuscript to which all co-authors contributed. The first and second author contributed equally to this paper.

Conflict of interests – The authors declare no conflicts of interest.

References

602   Aldous, A. R. 2002. Nitrogen translocation in *Sphagnum* mosses: effects of atmospheric nitrogen deposition. – New Phytol. 156: 241–253.

604   Allesina, S. et al. 2009. Functional links and robustness in food webs. – Philos. Trans. R. Soc. Lond. B. Biol. Sci. 364: 1701–1709.

606   Bardgett, R. D. et al. 2013. Hierarchical responses of plant-soil interactions to climate change: consequences for the global carbon cycle. – J. Ecol. 101: 334–343.

608   Bascompte, J. 2009. Disentangling the Web of Life. – Science 325: 416–419.

Berry, D. and Widder, S. 2014. Deciphering microbial interactions and detecting keystone species with co-
610   occurrence networks. – Front. Microbiol. 5: 219

Bragazza, L. et al. 2006. Atmospheric nitrogen deposition promotes carbon loss from peat bogs. – Proc. Natl.
612   Acad. Sci. U. S. A. 103: 19386–19389.

Bragina, A. et al. 2012. *Sphagnum* mosses harbour highly specific bacterial diversity during their whole
614   lifecycle. – ISME J. 6: 802–813.

Bragina, A. et al. 2014. The *Sphagnum* microbiome supports bog ecosystem functioning under extreme
616   conditions. – Mol. Ecol. 23: 4498–4510.

Canard, E. et al. 2012. Emergence of structural patterns in neutral trophic networks. – PLoS ONE 7: e38295

618   Caporaso, J. G. et al. 2010. QIIME allows analysis of high-throughput community sequencing data. – Nat. Methods 7: 335–336.

620   Chronakova, A. et al. 2019. Spatial heterogeneity of belowground microbial communities linked to peatland microhabitats with different plant dominants. – FEMS Microbiol. Ecol. 95: fiz130.

622   Clymo, R. S. 1965. Experiments on breakdown of *Sphagnum* in two bogs. – J. Ecol. 53: 747–758.

Clymo, R. S. 1984. The limits to peat bog growth. – Philos. Trans. R. Soc. Lond. B Biol. Sci. 303:605–654.

624   Dalsgaard, B. et al. 2011. Specialization in plant-hummingbird networks is associated with species richness, contemporary precipitation and quaternary climate-change velocity. – PLoS ONE 6: e25891.

626   Dalsgaard, B. et al. 2013. Historical climate-change influences modularity and nestedness of pollination networks. – Ecography 36: 1331–1340.

628   Devictor, V. et al. 2010. Defining and measuring ecological specialization. – J. Appl. Ecol. 47: 15–25.

Dieleman, C. M. et al. 2015. Climate change drives a shift in peatland ecosystem plant community: implications
630   for ecosystem function and stability. – Glob. Change Biol. 21: 388–395.

Dorrepaal, E. et al. 2005. Are growth forms consistent predictors of leaf litter quality and decomposability
632   across peatlands along a latitudinal gradient? – J. Ecol. 93: 817–828.

Dorrepaal, E. et al. 2009. Carbon respiration from subsurface peat accelerated by climate warming in the subarctic. – Nature 460: 616–619.

Edgar, R. C. 2011. UCHIME improves sensitivity and speed of chimera detection. – Bioinformatics 27: 2194–2200.

Estop-Aragonés, C. et al. 2016. Effects of extreme experimental drought and rewetting on CO_2 and CH_4 exchange in mesocosms of 14 European peatlands with different nitrogen and sulfur deposition. – Glob. Change Biol. 22: 2285–2300.

Elmendorf, S.C. et al. 2015. Experiment, monitoring, and gradient methods used to infer climate change effects on plant communities yield consistent patterns – Proc. Natl. Acad. Sci. USA 112: 448–452.

Fanin, N. et al. 2019. Effects of plant functional group removal on structure and function of soil communities across contrasting ecosystems. – Ecol. Lett. 22: 1095–1103.

Faust, K. and Raes, J. 2012. Microbial interactions: from networks to models. – Nat. Rev. Microbiol. 10: 538–550.

Fenner, N. and Freeman, C. 2011. Drought-induced carbon loss in peatlands. – Nat. Geosci. 4: 895–900.

Fenner, N. and Freeman, C. 2020. Woody litter protects peat carbon stocks during drought. – Nat. Clim. Change 10:363–369.

Fisk, M. C. et al. 2003. Microbial activity and functional composition among northern peatland ecosystems. – Soil Biol. Biochem. 35: 591–602.

Fudyma, J. D. et al. 2019. Untargeted metabolomic profiling of *Sphagnum fallax* reveals novel antimicrobial metabolites. – Plant Direct 3: 1-17.

Galiana, N. et al. 2014. Invasions cause biodiversity loss and community simplification in vertebrate food webs. – Oikos 123: 721–728.

Galiana, N. et al. 2019. The geographical variation of network structure is scale dependent: understanding the biotic specialization of host–parasitoid networks. – Ecography 42: 1175–1187.

Gunnarsson, U. et al. 2002. Dynamics or constancy in *Sphagnum* dominated mire ecosystems? A 40-year study. – Ecography 25: 685–704.

Hagedorn, F. et al. 2019. Above- and belowground linkages shape responses of mountain vegetation to climate change. – Science 365: 1119-1123

Hamard, S. et al. 2019. Effects of *Sphagnum* leachate on competitive *Sphagnum* microbiome depend on species and time. – Front. Microbiol. 10: 2042.

Hijmans, R. J. et al. (2005). Very high resolution interpolated climate surfaces for global land areas. – Int. J.
664   Climatol. 25: 1965–1978.

Ivanova, A. A. et al. 2020. Closely located but totally distinct: highly contrasting prokaryotic diversity patterns
666   in raised bogs and eutrophic fens. – Microorganisms 8: E484.

Jassey, V. E. J. et al. 2018. Tipping point in plant–fungal interactions under severe drought causes abrupt rise in
668   peatland ecosystem respiration. – Glob. Change Biol. 24: 972–986.

Kéfi, S. et al. 2016. How structured is the entangled bank? The surprisingly simple organization of multiplex
670   ecological networks leads to increased persistence and resilience. – PLoS Biol. 14: e1002527.

Koleff, P. et al. 2003. Measuring beta diversity for presence-absence data. – J. Anim. Ecol. 72: 367–382.

672 Kostka, J. E. et al. 2016. The *Sphagnum* microbiome: new insights from an ancient plant lineage. – New Phytol.
      211: 57-64

674 Kotsyurbenko, O. R. et al. 2007. Shift from acetoclastic to H_2-sependent methanogenesis in a West Siberian
      peat bog at low pH values and isolation of an acidophilic Methanobacterium strain. – Appl. Environ.
676   Microbiol. 73: 2344–2348.

Lamentowicz, M. et al. 2019. Unveiling tipping points in long-term ecological records from *Sphagnum*-
678   dominated peatlands. – Biol. Lett. 15: 20190043.

Legendre, P. and De Cáceres, M. 2013. Beta diversity as the variance of community data: dissimilarity
680   coefficients and partitioning. – Ecol. Lett. 16: 951–963.

Lehtovirta-Morley, L. E. et al. 2011. Cultivation of an obligate acidophilic ammonia oxidizer from a nitrifying
682   acid soil. – Proc. Natl. Acad. Sci. USA 108: 15892–15897.

Limpens, J. et al. 2003. N deposition affects N availability in interstitial water, growth of *Sphagnum* and
684   invasion of vascular plants in bog vegetation. – New Phytol. 157: 339–347.

Lin, X. et al. 2015. Metabolic potential of fatty acid oxidation and anaerobic respiration by abundant members
686   of Thaumarchaeota and Thermoplasmata in deep anoxic peat. –ISME J. 9: 2740–2744.

Lindo, Z. et al. 2013. Bryophyte-cyanobacteria associations as regulators of the northern latitude carbon balance
688   in response to global change. – Glob. Change Biol. 19: 2022–2035.

Mandakovic, D. et al. Structure and co-occurrence patterns in microbial communities under acute environmental
690   stress reveal ecological factors fostering resilience. – Sci. Rep. 8: 5875

Martí, M. et al. 2015. Nitrogen and methanogen community composition within and among three *Sphagnum*
692   dominated peatlands in Scandinavia. – Soil Biol. Biochem. 81: 204–211.

Martí, M. et al. 2019a. Effects of long-term warming and enhanced nitrogen and sulfur deposition on microbial communities in a boreal peatland. – bioRxiv 704411.

Martí, M. et al. 2019b. Strong long-term interactive effects of warming and enhanced nitrogen and sulphur deposition on the abundance of active methanogens in a boreal oligotrophic mire. – Mires Peat 24: 1–14.

Metzger, M. J. et al. 2005. A climatic stratification of the environment of Europe: a climatic stratification of the European environment. – Glob. Ecol. Biogeogr. 14: 549–563.

Morriën, E. et al. 2017. Soil networks become more connected and take up more carbon as nature restoration progresses. – Nat. Commun. 8: 14349,

Nichols, J. E. and Peteet, D. M. 2019. Rapid expansion of northern peatlands and doubled estimate of carbon storage. – Nat. Geosci. 12: 917–921.

Norby, R. J et al. 2019. Rapid loss of an ecosystem engineer: *Sphagnum* decline in an experimentally warmed bog. Ecol Evol. 9: 12571–12585

Olid, C. et al. 2014. The effects of temperature and nitrogen and sulfur additions on carbon accumulation in a nutrient-poor boreal mire: Decadal effects assessed using ^{210}Pb peat chronologies. – J Geophys. Res.: Biogeosciences 119: 392–403.

Pester, M. et al. 2010. A 'rare biosphere' microorganism contributes to sulfate reduction in a peatland. – ISME J. 4: 1591-1602.

Pester, M. et al. 2012. Sulfate-reducing microorganisms in wetlands - fameless actors in carbon cycling and climate change. – Front. Microbiol 3: 72.

Pieterse, G. et al. 2007. High resolution modelling of atmosphere-canopy exchange of acidifying and eutrophying components and carbon dioxide for European forests. – Tellus B 59: 412–424.

Pimm, S. L. et al. 2014. The biodiversity of species and their rates of extinction, distribution, and protection. – Science 344: 1246752–1246752.

Pinceloup, N. et al. 2020. Vegetation changes in temperate ombrotrophic peatlands over a 35 year period. – PLoS ONE 15: e0229146.

Poisot, T. et al. 2012a. The dissimilarity of species interaction networks. – Ecol. Lett. 15:1353–1361.

Poisot, T. et al. 2012b. A comparative study of ecological specialization estimators. – Methods Ecol. Evol. 3:537–544.

Ratcliffe, J. L. et al. 2020. Northern peatland depth and lateral expansion is inconsistent with a 1055 GtC estimate of carbon storage. – EarthArXiv. DOI:10.31223/osf.io/kafcu.

Rillig, M. C. et al. 2019. The role of multiple global change factors in driving soil functions and microbial biodiversity. – Science 366: 886–890.

Robroek, B.J.M. et al. 2015. Peatland vascular plant functional types affect methane dynamics by altering microbial community structure. – J. Ecol. 103:925–934.

Robroek, B. J. M. et al. 2017. Taxonomic and functional turnover are decoupled in European peat bogs. – Nat. Commun. 8:1181.

Rupp, D. et al. 2019. Plant functional group effects on peat carbon cycling in a boreal rich fen. – Biogeochemistry 144: 305–327.

Sander, E. L. et al. 2017. Ecological network inference from long-term presence-absence data. – Sci. Rep. 7: 7154.

Schleuning, M. et al. 2016. Ecological networks are more sensitive to plant than to animal extinction under climate change. – Nat. Commun 7: 13965.

Tylianakis, J. M. et al. 2008. Global change and species interactions in terrestrial ecosystems. – Ecol. Lett. 11: 1351–1363.

Valiente-Banuet, A. et al. 2015. Beyond species loss: the extinction of ecological interactions in a changing world. – Functional Ecology 29: 299–307.

Vitt, D. H. and Slack N. G. 1975. An analysis of the vegetation of *Sphagnum* -dominated kettle-hole bogs in relation to environmental gradients. – Can. J. Bot. 53: 332–359.

de Vries, F .T. et al. 2012. Abiotic drivers and plant traits explain landscape-scale patterns in soil microbial communities. – Ecol. Lett. 15: 1230–1239

de Vries, F. T. et al. 2018. Soil bacterial networks are less stable under drought than fungal networks. – Nat. Commun. 9: 3033.

Walther, G.-R. et al. 2002. Ecological responses to recent climate change. – Nature 416: 389–395.

Wang, H. et al. 2015. Dual controls on carbon loss during drought in peatlands. – Nat. Clim. Change 5: 584–587.

Wardle, D. A. et al. 2004. Ecological linkages between aboveground and belowground biota. – Science 304:1629–1633.

Weedon, J. T. et al. 2017. Compositional stability of the bacterial community in a climate-sensitive sub-Arctic peatland. – Front. Microbiol. 8: 317.

752   Wei, C. et al. 2013. Nitrogen deposition weakens plant–microbe interactions in grassland ecosystems. – Glob. Change Biol, 19: 3688–3697.

754   Wiedermann, M. M. et al. 2007. Global change shifts vegetation and plant-parasite interactions in a boreal mire. – Ecology 88:454–464.

756   Williams, R. J. et al. 2014. Demonstrating microbial co-occurrence pattern analyses within and between ecosystems. – Front. Microbiol. 5: 358.

758   Wilson, R. M. et al. 2016. Stability of peatland carbon to rising temperatures. – Nat. Commun. 7: 13723

Yavitt, J. B. and Williams, C. J. 2015. Linking tree species identity to anaerobic microbial activity in a forested

760   wetland soil via leaf litter decomposition and leaf carbon fractions. – Plant Soil 390: 293–305.

Yu, Y. et al. 2005. Group-specific primer and probe sets to detect methanogenic communities using quantitative

762   real-time polymerase chain reaction. – Biotechnol. Bioeng. 89: 670–679.

Yu, Z. et al. 2010. Global peatland dynamics since the Last Glacial Maximum. – Geophys. Res. Lett. 37:

764   L13402

Peatland plant-microbe network rewiring outpaces species turnover

Bjorn JM Robroek[1,†], Magalí Martí[2,†], Bo H Svensson[2], Marc G Dumont[3], Annelies J. Veraart[1], Vincent EJ Jassey[4]

[1]Aquatic Ecology & Environmental Biology, Institute for Water and Wetland Research, Faculty of Science, Radboud University Nijmegen, AJ 6525 Nijmegen, The Netherlands; [2]Thematic Studies - Environmental Change, Linköping University, SE-581 83 Linköping, Sweden; [3]School of Biological Sciences, Faculty of Environmental and Life Sciences, University of Southampton, SO17 1BJ Southampton, United Kingdom; [4]Laboratoire d'Ecologie Fonctionnelle et Environnement, Université de Toulouse, CNRS, 31062 Toulouse Cedex, France; [†]These authors contributed equally to this work

Supplementary Tables and Figures

Supplementary Table 1 Key prokaryotes in the networks (OTUs with > 8 interactions). * indicate uncultured prokaryotes.

Abbr	Kingdom	Phylum	Class	Order	Family
-- acrotelm layer (above water table) ----------------------------					
Bac 1	Bacteria	Verrucomicrobia	Verrucomicrobiae	Chthoniobacterales	Chthoniobacteraceae
Bac 2	Bacteria	Acidobacteria	Acidobacteriia	Acidobacteriales	Acidobacteriaceae
Bac 3	Bacteria	Acidobacteria	Acidobacteriia	Acidobacteriales	Acidobacteriaceae
Bac 4	Bacteria	Acidobacteria	Acidobacteriia	Acidobacteriales	Acidobacteriaceae
Bac 5	Bacteria	Acidobacteria	Acidobacteriia	Acidobacteriales	Acidobacteriaceae
Bac 6	Bacteria	Actinobacteria	Acidimicrobiia	IMCC26256	*
Bac 7	Bacteria	Acidobacteria	Holophagae	Subgroup 7	proteobacterium*
Bac 8	Bacteria	Acidobacteria	Acidobacteriia	Acidobacteriales	Acidobacteriaceae
Bac 9	Bacteria	Proteobacteria	Gammaproteobacteria	WD260	*
Bac 10	Bacteria	Acidobacteria	Acidobacteriia	Acidobacteriales	*
Bac 11	Bacteria	Planctomycetes	Phycisphaerae	Tepidisphaerales	WD2101 soil group
Bac 12	Bacteria	Actinobacteria	Actinobacteria	Frankiales	Acidothermaceae
Bac 13	Bacteria	Verrucomicrobia	Verrucomicrobiae	Pedosphaerales	Pedosphaeraceae
Bac 14	Bacteria	Verrucomicrobia	Verrucomicrobiae	Methylacidiphilales	Methylacidiphilaceae
-- catotelm layer (below water table) ----------------------------					
Arc 1	Archaea	Euryarchaeota	Methanomicrobia	Methanomicrobiales	Methanoregulaceae
Arc 2	Archaea	Thaumarchaeota	Group 1.1c	uncultured archaeon	*
Bac 15	Bacteria	Verrucomicrobia	Verrucomicrobiae	Opitutales	Opitutaceae
Bac 16	Bacteria	Acidobacteria	Acidobacteriia	Solibacterales	Solibacteraceae
Bac 17	Bacteria	WPS-2	Forest soil bacterium	*	
Bac 18	Bacteria	Proteobacteria	Alphaproteobacteria	Rhizobiales	Xanthobacteraceae
Bac 19	Bacteria	Actinobacteria	Thermoleophilia	Solirubrobacterales	Solirubrobacteraceae
Bac 20	Bacteria	Verrucomicrobia	Verrucomicrobiae	Methylacidiphilales	Methylacidiphilacea
Bac 21	Bacteria	Proteobacteria	Gammaproteobacteria	JG36-TzT-191	*
Bac 22	Bacteria	Acidobacteria	Acidobacteriia	Acidobacteriales	Koribacteraceae
Bac 23	Bacteria	Acidobacteria	Acidobacteriia	Acidobacteriales	*
Bac 24	Bacteria	Acidobacteria	Acidobacteriia	Acidobacteriales	Acidobacteriaceae
Bac 25	Bacteria	Acidobacteria	Acidobacteriia	Acidobacteriales	Koribacteraceae
Bac 26	Bacteria	Acidobacteria	Acidobacteriia	Solibacterales	Solibacteraceae
Bac 27	Bacteria	Actinobacteria	Actinobacteria	Frankiales	Acidothermaceae
Bac 28	Bacteria	Acidobacteria	Acidobacteriia	Acidobacteriales	*
Bac 29	Bacteria	Acidobacteria	Acidobacteriia	Solibacterales	Solibacteraceae
Bac 30	Bacteria	Actinobacteria	Acidimicrobiia	*	
Bac 31	Bacteria	Planctomycetes	Phycisphaerae	Tepidisphaerales	WD2101 soil group
Bac 32	Bacteria	Proteobacteria	Alphaproteobacteria	Rhizobiales	Beijerinckiaceae
Bac 33	Bacteria	Proteobacteria	Alphaproteobacteria	Acetobacterales	Acetobacteraceae
Bac 34	Bacteria	Verrucomicrobia	Verrucomicrobiae	Pedosphaerales	Pedosphaeraceae
Bac 35	Bacteria	Acidobacteria	Acidobacteriia	Acidobacteriales	*

Bac 36	Bacteria	Planctomycetes	Phycisphaerae	Tepidisphaerales	WD2101 soil group
Bac 11	Bacteria	Planctomycetes	Phycisphaerae	Tepidisphaerales	WD2101 soil group
Bac 37	Bacteria	Actinobacteria	Acidimicrobiia	IMCC26256	*
Bac 38	Bacteria	WPS-2			
Bac 39	Bacteria	Verrucomicrobia	Verrucomicrobiae	Pedosphaerales	Pedosphaeraceae
Bac 40	Bacteria	Proteobacteria	Alphaproteobacteria	Rhizobiales	Beijerinckiaceae
Bac 41	Bacteria	Acidobacteria	Acidobacteriia	Solibacterales	Solibacteraceae

Supplementary Figure 1 Schematic cross-cut of a typical *Sphagnum*-dominated peatland with a distinct lawn and hummock (i.e. peat mound) microhabitat pattern. In the both microhabitats, samples were extracted from the acrotelm and the catotelm peat. Acrotelm peat samples were takes 5 cm below the peat surface in both microhabitats. Catotelm peat samples were taken 12.5 cm and 37.5 cm below the peatland surface in the lawn and hummock microhabitats, respectively.

Top-down plant-microbe interactions   Bottom-up plant-microbe interactions

852 **Supplementary Figure 2** Plant-microbe bipartite network interactions. Each plant (I) in the plant community (I-IV) interacts with a number of prokaryotic 'species' (a,b,c,f,g & h). Hence, top-down interactions (left) can be

854 quantified for all plant-microbe interactions in terms of network attributes (see main text). In the same network (right), a prokaryotic 'species' (c) can interact with species in the plant community (I & II). Again, these

856 bottom-up interactions can be quantified for all microbe-plant interactions in terms of network attributes. It is important to note that top-down and bottom-up network attributes can differ significantly (see Fig. 2, and main

858 text) as the number interactions of plant species with microbial 'species' does not necessarily parallel the number of interactions of microbial species with plant species.

Supplementary Figure 3 Enviro-climatological variable reduction using the *ClustOfVar* function (Chavent et al. 2012). Following *ClustOfVar*, three variables were selected (bold) from the different clusters (in red): SO_x, pH and TS.

Supplementary Figure 4 Redundancy analyses (RDA; axes 1 and 2) biplot of site scores for the plant species (left), prokaryotic OTUs sampled from the acrotelm peat (centre), and prokaryotic OTUs sampled from the catotelm peat (right). RDA scores are constrained by data on pH, seasonality of temperature (TS) and sulphur deposition (N). * indicates enviro-climatological variable that were returned as significant ($P \leq 0.05$) by the RDA models.

Supplementary Figure 5 Network topology for the (a) plant–acrotelm prokaryote and (b) plant–catotelm

882 prokaryote bi-partite networks, calculated at network level. LD = Linkage density (LD), Gp = Generality of

plant links with prokaryotes. Gm = Generality of prokaryote links with plants, R = Robustness, PDn = Partner

884 diversity in the networks, SR = Species richness (SR), N= Niche specialization (N). The indices G, R, PDn SR

and N were calculated separately for the two levels –plants (p) and prokaryotes (m)– in the bipartite networks.

Additional references

888 *R packages*

bipartite: Dormann, C.F. et al. 2009. Indices, graphs and null models: analyzing bipartite ecological networks. –

890   The Open Ecology Journal 2: 7-24.

ClustOfVar: Chavent, M. et al. 2012. *ClustOfVar*: An R package for the clustering of variables. *J Stat Softw*. 50:

892   1–16.

diverse: Guevara, M.R. et al. 2016. diverse: an R package to measure diversity in complex systems. –The R

894   Journal 8: 60–78.

dplyr: Wickham, H. et al. 2019. dplyr: A Grammar of Data Manipulation. – R package version 0.8.3.

896 *igraph*: Csardi, G. and Nepusz, T. 2006. The igraph software package for complex network research. –

    InterJournal, Complex Systems 1695.

898 *nmle*: Pinheiro, J. et al. 2020. nlme: Linear and Nonlinear Mixed Effects Models. – R package version 3.1-147.

phyloseq: McMurdie, P.J and Holmes, S. 2013. phyloseq: An R Package for Reproducible Interactive Analysis

900   and Graphics of Microbiome Census Data. – PLoS ONE 8: e61217.

tidyr: Wickham, H. and Henry, L. 2020. Tidyr: Tidy Messy Data.

902 *vegan*: Oksanen, J. et al. 2019. vegan: Community Ecology Package.– R package version 2.5-6