Mediocremonas mediterraneus, a New Member within the Developea

Bradley A. Weiler[a], Elisabeth L. Sa[b] Michael E. Sieracki[c], Ramon Massana[b], Javier del Campo[a*]

a Department of Marine Biology and Ecology, Rosenstiel School of Marine and Atmospheric Science, University of Miami, Miami, 33149, Florida, USA

b Department of Marine Biology and Oceanography, Institut de Ciències del Mar (CSIC), Barcelona, 08003, Catalonia, Spain

c National Science Foundation, Alexandria, 22314, Virginia, USA

* **Correspondence**
J. del Campo, Department of Marine Biology and Ecology, Rosenstiel School of Marine and Atmospheric Science, University of Miami, Miami, 33149, Florida, USA
Telephone number; e-mail jdelcampo@rsmas.miami.edu

ABSTRACT

The Stramenopiles are a large and diverse group of eukaryotes that possess various lifestyles required to thrive in a broad array of environments. The stramenopiles branch with the alveolates, rhizarians, and telonemids, forming the supergroup TSAR. Here, we present a new genus and species of aquatic nanoflagellated stramenopile: *Mediocremonas mediterraneus*, a free-swimming heterotrophic predator. *M. mediterraneus* cell bodies measure between 2.0-4.0 μm in length and 1.2-3.7 μm in width, possessing two flagella and an oval body morphology. The growth and grazing rate of *M. mediterraneus* in batch cultures ranges from 0.68 to 1.83 d^{-1} and 1.99 to 5.38 bacteria h^{-1}, respectively. *M. mediterraneus* was found to be 93.9% phylogenetically similar with *Developayella elegans* and 94.7% with *Develorapax marinus*, two members within the class Developea. The phylogenetic position of the Developea and the ability of *M. mediterraneus* to remain in culture makes it a good candidate for further genomic studies that could help us to better understand phagotropy in marine systems as well as the transition from heterotrophy to phototrophy within the stramenopiles.

KEYWORDS

TSAR; stramenopiles; developea; heterotrophic nanoflagellates; phagotrophy; bacterivory; culturing; phylogeny, taxonomy; microscopy

INTRODUCTION

Stramenopiles, also known as Heterokonts (Cavalier-Smith 1986), are a diverse group of eukaryotes found in marine, limnetic and terrestrial systems (Andersen 2004; Massana et al. 2004, 2014; Simon et al. 2015). The stramenopiles branch with Alveolata, Rhizaria, and Telonemia forming a clade supergroup termed TSAR (Adl et al. 2019; Strassert et al. 2019). Within the stramenopiles there are two large monophyletic clades Bigyra and Gyrista, that each contain two monophyletic groups Opalozoa/Sagenista and Ochrophyta/Oomycota, respectively (Derelle et al. 2016). However, these taxonomic classifications are prone to change due to molecular environmental surveys that uncover the expansive diversity previously impossible to characterize by morphological and culturing approaches. These molecular surveys using environmental DNA have uncovered an array of uncultured eukaryotes such as the marine stramenopiles (MAST) clades from diverse aquatic habitats, including also limnetic ecosystems despite their name (Massana et al. 2014).

Stramenopiles have various lifestyles, such as photoautotrophic (most of the ochrophytes, eg. diatoms), heterotrophic predators (e.g. bicosoecids), osmotrophic (e.g. oomycetes) and parasitic (e.g. *Blastocystis*) (Arndt et al. 2000; Andersen 2004; Tan 2008). In some cases, certain stramenopiles have a combination of both photoautotrophic and heterotrophic feeding behaviours, termed mixotrophic that both possess chloroplasts and may also phagocytize prey (Jürgens and Massana 2008). Heterotrophic flagellates capture their prey by raptorial feeding, and prey items are subsequently consumed by phagocytosis (Hansen et al. 1994; Jürgens and Massana 2008). The functional and numerical responses, as discussed in detail by Weisse et al. (2016), are models used to predict the functional ecology of prey consumption by mixo- and heterotrophic aquatic protists. These responses depict how the flagellate growth and grazing rates vary with prey abundance, and can generally be adjusted to Michaelis-Menten kinetics (Weisse et al. 2016).

A novel class within the superphylum Stramenopiles was suggested, termed Developea (Aleoshin et al. 2016), which is composed of biflagellated non-amoeboid heterotrophic nanoflagellates (Cavalier-Smith 2018). Developea was classified under the subphylum Bigyromonada (that includes Pirsonea), within the phylum Gyrista. Within Developea, *Developayella elegans* is a free-swimming aquatic heterotrophic nanoflagellate predator which was previously purported to be grouped within the Oomycetes (AKA Pseudofungi) (Leipe et al. 1996; Cavalier-Smith and Chao 2006). *D. elegans* was recently observed to group with a novel species, *Develorapax marinus*, both forming the Developea class (Aleoshin et al. 2016). Despite not being very abundant in the environment based on molecular surveys, Developea is a very interesting group because of its phylogenetic position. Recent phylogenomic studies have placed *D. elegans* at the base of the Ochrophyta (Leonard et al. 2018), so Developea could represent a key group to understand the transition from heterotrophy to phototrophy within the Stramenopiles. In a previous culturing effort (del Campo et al. 2013), we isolated several heterotrophic nanoflagellates, and one of them remained poorly characterized. Here, we focus on that culture and present a new genus and species of aquatic heterotrophic nanoflagellate: *Mediocremonas mediterraneus*, that clustered closely with both *D. elegans* and *D. marinus* within the Developea class in our 18S rRNA gene phylogenetic analyses. In addition, we present the functional and numerical response of *M. mediterraneus*.

93
94 **MATERIAL AND METHODS**
95
Isolation by single cell sorting
The HNF described here was isolated in a previous culturing effort (del Campo et al. 2013), and
the isolation procedure is explained below. Seawater from the Blanes Bay Microbial Observatory
sampled on September 30[th], 2008 was filtered through 3 µm and sent to Bigelow Laboratory for
Ocean Sciences (Boothbay Harbor, ME, USA) for cell sorting in a MoFlo™ Flow Cytometer
(Dako-Cytomation, Denmark). Digestive vacuoles of heterotrophic protists were stained using
the vital stain LysoTracker® (Life Technologies, NY, USA) and cells were sorted based on their
green fluorescence (LysoTracker® fluorescence) and the absence of chlorophyll fluorescence.
Details of the staining protocol and flow cytometer setup are described elsewhere (Rose et al.
2004; Heywood et al. 2011). Side scatter was used to select only the smallest protists,
approximately <10 µm in diameter. Individual target cells were deposited into 24-well plates, in
which some wells were dedicated for positive controls (10 cells per well) and negative controls
(0 cells per well). All wells on the microplates contained 1ml aged seawater together with natural
bacteria at a final concentration of 5×10^6 bacteria ml^{-1}. Multi-well plates were hand carried
back to the Institut de Ciències del Mar (Barcelona, Catalonia, Spain) by plane at room
temperature (12 hours).
112
Culture maintenance
Cultures were maintained in 50 mL flasks and transferred every two to four weeks to fresh media
(aged sea water with added bacteria) at 1/10 dilution. The full culturing protocol is outlined in
del Campo *et al*. (2013), however in short batch cultures were prepared by adding small
aliquotes of the *M. mediterraneus* culture to sterile seawater (about 2 liters) containing *Dokdonia*
donghaensis MED134 at varying cell abundances. Bacterial and flagellate counting was
conducted once or twice a day using epifluorescence microscopy and DAPI stain by fixing 3 mL
aliquots of the batch culture with glutaraldehyde directly after sampling. Growth rate was
calculated from the exponential increase of flagellate cells. Grazing rate was calculated based on
the calculations by Frost (1972) using the resultant growth rates, the exponential decrease of
bacterial cells and the geometric average concentration of flagellates and bacteria during the
incubation (Frost 1972). Growth efficiency was calculated from the ratio of protist biomass
produced as compared with bacterial biomass consumed.
126
Sequencing and phylogenetic analysis
For molecular analysis, the whole culture was filtered on 0.6 µm polycarbonate filters, DNA was
extracted by standard procedures, and 18S rDNA genes were PCR amplified with eukaryote-
specific primers (Díez et al. 2001). Complete sequences of 18S rDNA were obtained with five
internal primers by the MACROGEN Genomics Sequencing Services (accession numbers
JX272636 & XXXXXXXX). The longest sequence of *M. mediterraneus (*JX272636), a subset of
previously published Stramenopiles and two outgroups were aligned using MAFFT v7.453
(setting: -auto) (Katoh and Standley 2013). Regions of poor alignment were removed using
trimAl v1.4 (Capella-Gutierrez et al. 2009) with a gap and similarity threshold of 0.3 and 0.001,
respectively (settings: -gt 0.3 -st 0.001). Finally, to explore *M. mediterraneus* phylogeny,
maximum likelihood (ML) inference was conducted with RAxML v8.2.12 (Stamatakis 2014)
assuming the GTRCAT model using rapid bootstrap analysis for 1,000 iterations.

139
Scanning Electron Microscopy
Scanning Electron Microscopy was performed as in Garcés et al. (2006). Samples were fixed for
2 h in 2% OsO_4 diluted in seawater, or with 2% glutaraldehyde. Cells were subsequently washed
with distilled water (2 h) and filtered onto a 0.6 µm pore size Nuclepore™ filter (Whatman,
Maidstone, UK). Samples were dehydrated in an ethanol series (30, 50, 70, 96, and 100%) for 15
min each, followed by an acetone series (25, 50, 75, and 100%) for 15 min each. Samples were
critical point-dried in liquid CO_2 using a BAL-TEC CPD 030 critical point drying apparatus
(Balzers Union, Balzers, Germany). Filters were subsequently glued to SEM-stubs with colloidal
silver, sputter coated with gold palladium, and examined with a Hitachi S-3500N (Nissei Sangyo
Co. Ltd., Tokyo, Japan) SEM operating at 5 kV.
150
151 **RESULTS AND DISCUSSION**
152
In terms of gross morphology *M. mediterraneus* is relatively similar to the other members of the
Developea clade (Aleoshin et al. 2016; Tong 1995). Ovoid cell bodies measure 2.0 – 4.0 μm in
length and 1.2 – 3.7 μm in width (Fig. 1A) and possesses a depression on the anterior ventral
surface (Fig. 1B, 1D). Protruding from the depression are two flagella of equal length, unlike
those of both *D. elegans* and *D. marinus*, measuring roughly twice the cell body length. The
anterior flagellum possesses mastigonemes and projects from the cell in an arc (Fig. 1). The
posterior flagellum projects posteriorly out from the cell depression relatively straight, not
possessing mastigonemes. The function of the Developea flagella have been previously
described for both locomotion (posterior) and current generation (anterior) to aid in prey capture
(Aleoshin et al. 2016; Tong 1995).
163
Sequences were analyzed using BLAST for phylogenetic similarity at the 18S rRNA
gene; it was 93.9% similar with *Developayella elegans* (Leipe et al. 1996) and 94.7% with
Develorapax marinus (Aleoshin et al. 2016). When analyzing the 18S rRNA gene sequence of
M. mediterraneus using maximum likelihood (ML), there is supported clustering with previously
described members of the Developea class (Leipe et al. 1996; Aleoshin et al. 2016), as well as
supported clustering with two other uncultured eukaryotes within the class Developea, both
retrieved from deep-sea sediments (Fig. 2). The next closest cluster observed in the ML tree is
the "Abyssal" group (as in Aleoshin et al. 2016) including eukaryotes found in the Pacific
abyssal plains (Takishita et al. 2010). Additionally, sister clade to the Developea+Abyssal
grouping are the Pseudofungi from the phylum Gyrista. As mentioned in the introduction the
most recent phylogenomic analysis of this part of the eukaryotic tree of life shows *D. elegans* at
the base of the Ochrophyta (Leonard et al. 2018), however this position is not strongly supported
in our 18S rDNA tree and in other phylogenetic reconstructions (Aleoshin et al. 2016). Obtaining
in the future the genome of *M. mediterraneus* can help to resolve this area of the Stramenopiles
tree within Ochrophyta and Pseudofungi.
179
We prepared several batch cultures where we added an aliquote of *M. mediterraneus* on
flasks containing varying cell abundances of bacterial strain and followed the dynamics over
time of both components (Fig. S1). By following the exponential predator increase and prey
decrease, we calculated the growth rate and the grazing rate in each of these batch cultures. The
grazing rate of *M. mediterraneus* ranged from 1.99 to 5.38 bacteria h^{-1}, and the growth rate

ranged from 0.68 to 1.83 d^{-1} (Table 1). The yield of bacteria to flagellate (Yield $_{b/f}$) ranged
between 54.8 and 84.1 (mean of 74.8, SD = 11.3). The doubling time in days observed for *M.*
mediterraneus , ranging from 0.38 to 1.02, was relatively longer than that of other heterotrophic
nanoflagellates (HNF), as *Paraphysomonas imperforata, Minorisa minuta*, and *Cafeteria*
roenbergensis have doubling times of 0.28, 0.44, and 0.45, respectively (Goldman and Caron
1985; del Campo et al. 2013; De Corte et al. 2019). The functional and numerical responses
showed a relatively clear trend (Fig. 3), wherein the growth and grazing rates increase as
bacterial prey abundance increases, up to a theoretical asymptotic maximum in terms of grazing
rate (I_{max}) and growth rate (R_{max}), inferred to be 6.23 (h^{-1}) and 2.42 (d^{-1}), respectively (Fig. 3).
194
Bacterivory in aquatic ecosystems is a crucial functional role played predominantly by
pico- and nanoflagellates (up to 5 µm) to maintain bacterial populations (Sherr and Sherr 2002).
Grazing on bacteria is necessary for nutrient recycling by releasing waste products back into the
environment in the form of inorganic compounds (such as iron, ammonium and phosphate), and
particulate and dissolved organic compounds (Sherr and Sherr 2002; Massana et al. 2009). The
grazing rate of *M. mediterraneus* ~4.1 bacteria h^{-1} is greater than some HNFs such as the closely
related MAST-1C group (Fig. 2), which has a grazing rate of 3.6 bacteria h^{-1} and the more
distantly related MAST-4 group, 1.5 bacteria h^{-1} (Table 2). In contrast, the grazing rate of *M.*
mediterraneus is much lower than several other cultured HNFs, including various gluttonous
members of the genus *Bodo spp.* (34.3 – 160 bacteria h^{-1}), and members of the genera *Spumella*
sp. and *Ochromonas sp.* (37 and 63 bacteria h^{-1}, respectively) (Table 2). In general, grazing rates
of cultured HNFs (Eccleston-Parry and Leadbeater 1994), which are suggested to be poor models
of natural and dominant taxa (Massana et al. 2009), are much higher than community grazing
rates typically between 2 and 20 bacteria h^{-1} (Jürgens and Massana 2008).
209
Blanes Bay is an oligotrophic coastal area of the NW Mediterranean and resides within
proximity to a continental submarine canyon. The microbial food web dynamics of Blanes Bay,
similar to other regions, are largely influenced by environmental factors (i.e. salinity, Chl a,
temperature, nutrients, etc…) and may change given a shift in just one of those factors (e.g.
increased sea surface temperature, Vázquez-Domínguez et al. (2012)). Given the oligotrophic
nature of Blanes Bay, and considering the low abundance of *M. mediterraneus* (Giner et al.
2019), it may be present perhaps as opportunistic consumers in microbial "hotspots" that arise in
the oligotrophic ecosystem (Stocker et al. 2008; Dang and Lovell 2016).
218
219   The ability to have *M. mediterraneus* in culture and its interesting phylogenetic position
220 make this organism a good candidate for further genomic studies that could help us to better
221 understand phagotropy in marine systems as well as the transition from heterotrophy to
222 phototrophy within the stramenopiles.
223
224 **TAXONOMIC SUMMARY**
225 **Phylum Gyrista, Cavalier-Smith, 1997**
226   **Class Developea, Aleoshin, 2016**
227   **Order Developayellales, Cavalier-Smith, 1997**
228   **Family Developayellaceae, Cavalier-Smith, 1997**
229   **Genus *Mediocremonas, del Campo & Weiler, 2020***
230   **Species *Mediocremonas mediterraneus, del Campo & Weiler, 2020***

231
Mediocremonas **n. gen. del Campo & Weiler**

Zoobank ID: XXXXX

Diagnosis: Heterotrophic oval-shaped protist possessing two flagella with a free-swimming predatorial feeding behaviour. Similar to other genera in the Developea (Aleoshin et al. 2016; Tong 1995), *Mediocremonas* flagella are ventrally located, with the anterior flagellum held in an arc and the posterior flagellum projecting posterior from a conspicuous depression. The anterior flagellum is observed to possess mastigonemes while generating current by performing a slow sweeping motion and the posterior beats rapidly for locomotion. The genus type species, *M. mediterraneus,* was isolated from Blanes Bay (Catalonia, Spain) and the 18S sequence was observed to be 93.9% similar to *Developayella elegans* from surface estuarine water of Southampton Water, UK (Leipe et al. 1996) and 94.7% similar to *Develorapax marinus* from coastal waters of the Red Sea (Aleoshin et al. 2016).

Type species: *M. mediterraneus* n. sp. del Campo & Weiler

Etymology: Mediocre – [average (l)] + monas – [unicellular organism (l)]

M. mediterraneus **n. sp. del Campo & Weiler**

Zoobank ID: XXXXX

Diagnosis: Cell bodies measure between 2.0-4.0 μm in length and 1.2-3.7 μm in width, possessing an oval body morphology. The general body size is comparatively smaller than other members of Developea, wherein *Developayella elegans* is roughly twice as large (3.5-8.5 μm in length and 2.0-6.0 μm in width) and the body size of *Develorapax marinus* is relatively largest (7.0-10.0 μm in length and 4.0-6.0 μm in width). Each cell contains two flagella originating from a ventral anterior depression in the cell body. Within the cell depression, the two flagella lay either anterior or posterior relative to one another, both with specific form and function. The anterior flagellum of *Developea* has been previously described to be held in an arc containing mastigonemes that sweeps slowly ventral posterior to the cell, while the posterior beats rapidly for locomotion. The anterior and posterior flagella of *M. mediterraneus* measure between 5.5-6.0 μm in length and 0.2-0.3 μm in width dissimilar to the unequal flagella measurements in both *D. elegans* and *D. marinus*.

Type material: Fig. 1 from cultures 9 and 11, isolated from Blanes Bay Microbial Observatory (NW Mediterranean Sea) and maintained at the Institut de Ciències del Mar (Barcelona, Catalonia, Spain). Scanning microscopy samples are also preserved there.

Type sequence: The SSU rRNA gene sequence is Genbank JX272636

Type locality: Blanes Bay (Catalonia, Spain)

Etymology: mediterraneus [from the Mediterranean (l)]

ACKNOWLEDGEMENTS

This work was supported by two grants from the Spanish government, FLAME (CGL2010-16304, MICINN) and ALLFLAGS (CTM2016-75083-R, MINECO), and by the National Science Foundation award DEB-1031049 to MES. BW was supported by the Natural Sciences and Engineering Research Council of Canada Postgraduate Scholarships-Doctoral program (NSERC PGS-D). BW and JdC were supported by startup funds from University of Miami, Rosenstiel School of Marine and Atmospheric Sciences.

277
278 **REFERENCES**
279
Adl S. M., Bass D., Lane C. E., Lukeš J., Schoch C. L., Smirnov A., Agatha S., Berney C.,
Brown M. W., Burki F., Cárdenas P., Čepička I., Chistyakova L. L. L., del Campo J.,
Dunthorn M., Edvardsen B., Eglit Y., Guillou L., Hampl V., Heiss A. A., Hoppenrath M.,
James T. Y., Karnkowska A., Karpov S. A., Kim E., Kolisko M., Kudryavtsev A., Lahr D.
J. G., Lara E., Le Gall L., Lynn D. H., Mann D. G., Massana R., Mitchell E. A. D. D.,
Morrow C., Park J. S., Pawlowski J. W., Powell M. J., Richter D. J., Rueckert S., Shadwick
L., Shimano S., Spiegel F. W., Torruella G., Youssef N., Zlatogursky V. & Zhang Q. 2019.
Revisions to the Classification, Nomenclature, and Diversity of Eukaryotes. *J. Eukaryot.*
Microbiol., 66:jeu.12691.
Aleoshin V. V., Mylnikov A. P., Mirzaeva G. S., Mikhailov K. V. & Karpov S. A. 2016.
Heterokont Predator Develorapax marinus gen. et sp. nov. – A Model of the Ochrophyte
Ancestor. *Front. Microbiol.*, 7:1–14.
Andersen R. A. 2004. Biology and systematics of heterokont and haptophyte algae. *Am. J. Bot.*,
91:1508–1522.
Arndt H., Dietrich D., Auer B., Cleven E. J., Gräfenhan T., Weitere M. & Mylnikov A. P. 2000.
Functional diversity of heterotrophic flagellates in aquatic ecosystems. *In*: Leadbeater B &
Green J (ed.), The Flagellates. Taylor and Francis, London. p. 240–268.
Boenigk J. & Arndt H. 2000. Particle handling during interception feeding by four species of
heterotrophic nanoflagellates. *J. Eukaryot. Microbiol.*, 47:350–8.
del Campo J., Not F., Forn I., Sieracki M. E. & Massana R. 2013. Taming the smallest predators
of the oceans. *ISME J.*, 7:351–358.
Capella-Gutierrez S., Silla-Martinez J. M. & Gabaldon T. 2009. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*, 25:1972–1973.
Cavalier-Smith T. 1986. The kingdom Chromista: origin and systematics. *In*: Round F. E. &
Chapman D. J. (eds.), Progress in Phycological Research. Bristol: Biopress Ltd. p. 309–347.
Cavalier-Smith T. 2018. Kingdom Chromista and its eight phyla: a new synthesis emphasising
periplastid protein targeting, cytoskeletal and periplastid evolution, and ancient divergences.
Protoplasma, 255:297–357.
Cavalier-Smith T. & Chao E. E. Y. 2006. Phylogeny and Megasystematics of Phagotrophic
Heterokonts (Kingdom Chromista). *J. Mol. Evol.*, 62:388–420.
Christaki U., Vazquez-Dominguez E., Courties C. & Lebaron P. 2005. Grazing impact of
different heterotrophic nanoflagellates on eukaryotic (Ostreococcus tauri) and prokaryotic
picoautotrophs (Prochlorococcus and Synechococcus). *Environ. Microbiol.*, 7:1200–1210.
De Corte D., Paredes G., Yokokawa T., Sintes E. & Herndl G. J. 2019. Differential Response of
Cafeteria roenbergensis to Different Bacterial and Archaeal Prey Characteristics. *Microb.*
Ecol., 78:1–5.
Dang H. & Lovell C. R. 2016. Microbial Surface Colonization and Biofilm Development in
Marine Environments. *Microbiol. Mol. Biol. Rev.*, 80:91–138.
Derelle R., López-García P., Timpano H. & Moreira D. 2016. A Phylogenomic Framework to
Study the Diversity and Evolution of Stramenopiles (=Heterokonts). *Mol. Biol. Evol.*,
33:2890–2898.
Díez B., Pedrós-Alió C. & Massana R. 2001. Study of Genetic Diversity of Eukaryotic
Picoplankton in Different Oceanic Regions by Small-Subunit rRNA Gene Cloning and

Sequencing. *Appl. Environ. Microbiol.*, 67:2932–2941.

Eccleston-Parry J. D. & Leadbeater B. S. C. 1994. A comparison of the growth kinetics of six marine heterotrophic nanoflagellates fed with one bacterial species. *Mar. Ecol. Prog. Ser.*, 105:167–177.

Frost B. W. 1972. Effects of size and concentration of food particles on the feeding behavior of the marine planktonic copepod Calanus pacificus. *Limnol. Oceanogr.*, 17:805–815.

Garcés E., Fernandez M., Penna A., Van Lenning K., Gutierrez A., Camp J. & Zapata M. 2006. Characterization of NW Mediterranean Karlodinium spp. (Dinophyceae) strains using morphological, molecular, chemical, and physiological methodologies. *J. Phycol.*, 42:1096–1112.

Giner C. R., Balagué V., Krabberød A. K., Ferrera I., Reñé A., Garcés E., Gasol J. M., Logares R. & Massana R. 2019. Quantifying long‐term recurrence in planktonic microbial eukaryotes. *Mol. Ecol.*, 28:923–935.

Goldman J. C. & Caron D. A. 1985. Experimental studies on an omnivorous microflagellate: implications for grazing and nutrient regeneration in the marine microbial food chain. *Deep Sea Res.*, 32:899–915.

Hansen B., Bjornsen P. K. & Hansen P. J. 1994. The size ratio between planktonic predators and their prey. *Limnol. Oceanogr.*, 39:395–403.

Heywood J. L., Sieracki M. E., Bellows W., Poulton N. J. & Stepanauskas R. 2011. Capturing diversity of marine heterotrophic protists: one cell at a time. *ISME J.*, 5:674–84.

Jürgens K. & Massana R. 2008. Protistan Grazing on Marine Bacterioplankton. *In*: Kirchman D. L. (ed.), Microbial Ecology of the Oceans. 2nd ed. Hoboken, NJ, USA, John Wiley & Sons, Inc. p. 383–441.

Katoh K. & Standley D. M. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.*, 30:772–80.

Leipe D. D., Tong S. M., Goggin C. L., Slemenda S. B., Pieniazek N. J. & Sogin M. L. 1996. 16S-like rDNA sequences from Developayella elegans, Labyrinthuloides haliotidis, and Proteromonas lacertae confirm that the stramenopiles are a primarily heterotrophic group. *Eur. J. Protistol.*, 32:449–458.

Leonard G., Labarre A., Milner D. S., Monier A., Soanes D., Wideman J. G., Maguire F., Stevens S., Sain D., Grau-Bové X., Sebé-Pedrós A., Stajich J. E., Paszkiewicz K., Brown M. W., Hall N., Wickstead B. & Richards T. A. 2018. Comparative genomic analysis of the "pseudofungus" Hyphochytrium catenoides. *Open Biol.*, 8:170184.

Massana R., del Campo J., Sieracki M. E., Audic S. & Logares R. 2014. Exploring the uncultured microeukaryote majority in the oceans: reevaluation of ribogroups within stramenopiles. *ISME J.*, 8:854–866.

Massana R., Castresana J., Balagué V., Guillou L., Romari K., Groisillier A., Valentin K. & Pedrós-Alió C. 2004. Phylogenetic and Ecological Analysis of Novel Marine Stramenopiles. *Appl. Environ. Microbiol.*, 70:3528–3534.

Massana R., Unrein F., Rodríguez-Martínez R., Forn I., Lefort T., Pinhassi J. & Not F. 2009. Grazing rates and functional diversity of uncultured heterotrophic flagellates. *ISME J.*, 3:588–596.

Rose J., Caron D., Sieracki M. & Poulton N. 2004. Counting heterotrophic nanoplanktonic protists in cultures and aquatic communities by flow cytometry. *Aquat. Microb. Ecol.*, 34:263–277.

Sherr E. B. & Sherr B. F. 2002. Significance of predation by protists in aquatic microbial food

webs. *Antonie van Leeuwenhoek, Int. J. Gen. Mol. Microbiol.*, 81:293–308.

Simon M., Jardillier L., Deschamps P., Moreira D., Restoux G., Bertolino P. & López-García P. 2015. Complex communities of small protists and unexpected occurrence of typical marine lineages in shallow freshwater systems. *Environ. Microbiol.*, 17:3610–3627.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*, 30:1312–1313.

Stocker R., Seymour J. R., Samadani A., Hunt D. E. & Polz M. F. 2008. Rapid chemotactic response enables marine bacteria to exploit ephemeral microscale nutrient patches. *Proc. Natl. Acad. Sci. U. S. A.*, 105:4209–4214.

Strassert J. F. H., Jamy M., Mylnikov A. P., Tikhonenkov D. V. & Burki F. 2019. New Phylogenomic Analysis of the Enigmatic Phylum Telonemia Further Resolves the Eukaryote Tree of Life.Shapiro B. (ed.). *Mol. Biol. Evol.*, 36:757–765.

Takishita K., Kakizoe N., Yoshida T. & Maruyama T. 2010. Molecular Evidence that Phylogenetically Diverged Ciliates Are Active in Microbial Mats of Deep-Sea Cold-Seep Sediment. *J. Eukaryot. Microbiol.*, 57:76–86.

Tan K. S. W. 2008. New insights on classification, identification, and clinical relevance of Blastocystis spp. *Clin. Microbiol. Rev.*, 21:639–665.

Tong S. M. 1995. Developayella elegans nov. gen., nov. spec., a new type of heterotrophic flagellate from marine plankton. *Eur. J. Protistol.*, 31:24–31.

Vázquez-Domínguez E., Vaqué D. & Gasol J. 2012. Temperature effects on the heterotrophic bacteria, heterotrophic nanoflagellates, and microbial top predators of the NW Mediterranean. *Aquat. Microb. Ecol.*, 67:107–121.

Weisse T., Anderson R., Arndt H., Calbet A., Hansen P. J. & Montagnes D. J. S. 2016. Functional ecology of aquatic phagotrophic protists – Concepts, limitations, and perspectives. *Eur. J. Protistol.*, 55:50–74.

Yubuki N., Pánek T., Yabuki A., Čepička I., Takishita K., Inagaki Y. & Leander B. S. 2015. Morphological Identities of Two Different Marine Stramenopile Environmental Sequence Clades: Bicosoeca kenaiensis (Hilliard, 1971) and Cantina marsupialis (Larsen and Patterson, 1990) gen. nov., comb. nov. *J. Eukaryot. Microbiol.*, 62:532–542.

FIGURE TITLES

Figure 1: SEM images of *M. mediterraneus* cells at varying angles (**A-D**), showing an ovoid body morphology and two flagella protruding from a surface depression ventrally located on the anterior portion of the cell. Anterior flagellum typically held in an arc possessing tubular mastigonemes. Scale bars measure 3 μm.

Figure 2: Maximum likelihood phylogenetic tree with partial (~1700 bp) 18S rRNA gene sequences showing the clustering of *M. mediterraneus* with the Developea group within the Gyrista clade. ML tree based off 1,000 iterations and only bootstrap values > 50 are shown. The symbol (•) represents bootstrap values of 100.

Figure 3: Numerical and functional responses of *M. mediterraneus*. **A.** showing the specific grazing rate (h^{-1}) on the y-axis as a response to increased bacterial abundance (cells/mL), with I_{max} indicating the theoretical maximum grazing rate and the resultant k_{50}. **B.** showing the specific growth rate (d^{-1}) on the y-axis, as a response to increased bacterial abundance (cells/mL), with R_{max} indicating the theoretical maximum growth rate and resultant k_{50}.

Supplementary Figure S1: Cell abundance dynamics of *M. mediterraneus* (black) and bacterial
prey (grey) in batch culture experiments (1 to 4). Experiments 1 & 2 have two replicates
indicated by A & B.

Table 1: Growth and bacterial grazing characteristics of *M. mediterraneus* by samples (1-4).
Samples 1 & 2 have replicates denoted by A & B.

Sample	Date	Initial Bacterial Conc. (cell/mL)	Growth Rate (d-1)	Doubling Time (days)	Grazing Rate (h-1)	Yield (bac/fla)	Growth Efficiency (Yield) (%)
1A	Oct. 2009	9.86E+06	0.68	1.02	1.99	70.90	35.31
1B	Oct. 2010	5.62E+07	1.37	0.51	4.38	82.94	30.19
2A	Nov. 2009	4.33E+07	1.64	0.42	5.38	83.04	30.15
2B*	Nov. 2009	3.92E+07	1.62	0.43	3.37	54.81	45.68
3	May 2011	1.24E+07	1.04	0.67	4.01	72.78	34.40
4*	Dec. 2011	4.22E+07	1.83	0.38	5.24	84.06	29.79

* with natural viruses

Table 2: Comparison of heterotrophic nanoflagellate (HNF) grazing rates ranked lowest to
highest.

HNF	Grazing Rate (h^{-1})	Reference
Cafeteria sp.	0.1	(Christaki et al. 2005)
MAST-4	1.5	(Massana et al. 2009)
Cafeteria roenbergensis	1.5	(De Corte et al. 2019)
Paraphysomonas imperforate	2.4	(Goldman and Caron 1985)
MAST-1C	3.6	(Massana et al. 2009)
Mediocremonas mediteranneus	4.1	Present study
Jakoba libera	5	(Eccleston-Parry and Leadbeater 1994)
Minorisa minuta	7.0	(del Campo et al. 2013)
Cafeteria roenbergensis	14.1	(Boenigk and Arndt 2000)
Bodo saltans	34.3	(Boenigk and Arndt 2000)
Codosiga gracilis	36	(Eccleston-Parry and Leadbeater 1994)
Spumella sp.	36.7	(Boenigk and Arndt 2000)
Stephanoexa diplocostata	37	(Eccleston-Parry and Leadbeater 1994)
Ochromonas sp.	63	(Boenigk and Arndt 2000)
Paraphysomonas imperforata	63	(Eccleston-Parry and Leadbeater 1994)
Bodo designis	160	(Eccleston-Parry and Leadbeater 1994)
Ciliophrys infusionum	259	(Eccleston-Parry and Leadbeater 1994)