Static All-Atom Energetic Mappings of the SARS-Cov-2 Spike Protein with Potential Latch Identification of the Down State Protomer

Michael H. Peters* *Virginia Commonwealth University*, Department of Chemical and Life Science Engineering, 601 West Main St, Richmond, Virginia, United States of America

Oscar Bastidas *University of Minnesota*, College of Biological Sciences, University of Minnesota, Minneapolis, Minnesota, United States of America

* To whom correspondence should be addressed. Email: (M.H.P.) mpeters@vcu.edu

Running Title: Energetic Mappings of the SARS-Cov-2 Spike Protein with Potential Latch Identification

Abstract

The SARS-Cov-2 virion responsible for the current world-wide pandemic Covid-19 has a characteristic Spike protein (S) on its surface that embellishes both a prefusion state and fusion state. The prefusion Spike protein (S) is a large trimeric protein where each protomer may be in a so-called Up state or Down state, depending on the configuration of its receptor binding domain (RBD). The Up state is believed to allow binding of the virion to ACE-2 receptors on human epithelial cells, whereas the Down state is believed to be relatively inactive or reduced in its binding behavior. We have performed detailed all-atom, dominant energy landscape mappings for noncovalent interactions (charge, partial charge, and van der Waals) of the SARS-Cov-2 Spike protein in its static prefusion state based on recent structural information. We included both interchain interactions and intrachain (domain) interactions in our mappings in order to determine any telling differences (different so-called "glue" points or "hot spots") between residues in the Up and Down state protomers. In general, the S2 or fusion machinery domain of S is relatively rigid with strong noncovalent interactions facilitated by helical secondary structures, whereas the S1 domain, which contains the RBD and N-terminal domain (NTD), is relatively more flexible and characterized by beta strand structural motifs. The S2 domain demonstrated no appreciable energetic differences between Up and Down protomers, including interchain as well as each protomer's intrachain, S1-S2 interactions. However, the S1 domain interactions across neighboring protomers, which include the RBD-NTD cross chain interactions, showed energetic differences between Up-Down and Down-Down neighboring protomers. Surprisingly, the Up-Down, RBD-NTD interactions were overall stronger and more numerous than the Down-Down cross chain interactions, including the appearance of the three residue sequence ALA520-PRO521-ALA522 associated with a turn structure in the RBD of the Up state. Additionally, our *intra*chain dominant energy mappings within each protomer, identified a significant "glue" point or possible latch for the Down state protomer between the S1 subdomain, SD1 and the RBD domain of the same protomer that was completely missing in the Up state protomer analysis. Ironically, this dominant energetic interaction in the Down state protomer involved the backbone atoms of the same three residue sequence ALA520-PRO521-ALA522 of the RBD with the R-group of GLN564 in the SD1 domain. Thus, this same three residue sequence acts as a stabilizer of the RBD in the Up conformation through its interactions with its neighboring NTD chain and a kind

of latch in the Down state conformation through its interactions with its own SD1 domain. The dominant interaction energy residues identified here are also conserved across reported variations of SARS-Cov-2, as well as the closely related virions SARS-Cov and the bat corona virus RatG13. However, these results are all based on static structure analysis and large scale dynamic simulations will be necessary to further investigate these findings. Nonetheless, static dominant energy landscape mappings represents a starting point for more detailed dynamic analyses and hopefully an improved understanding at the structure-function relationship of this highly complex protein.

Key words: SARS-Cov-2, spike protein, energetic mappings, contact plot, latch, COVID-19

1. Introduction

SARS-Cov-2 is an RNA virus responsible for the disease Covid-19 which was recently declared a world-wide pandemic by the WHO. SARS-Cov-2 virion enters human epithelial cells by attachment of it's prefusion-form Spike protein (S) with a specific cell surface receptor ACE2 (Angiotensin Converting Enzyme 2) (1-3). The Spike protein is a large, trimeric protein whose Receptor Binding Domain (RBD) undergoes somewhat unusual dynamic transformations sometimes called "breathing". From a protein engineering perspective, so-called "breathing" reflects the inherent flexibility and/or localized mobility associated with the Receptor Binding Domain (RBD) of the Spike Protein. In the so-called "Up-state" of the RBD, the (prefusion) protein is able to bind to ACE2 (Angiotensin Converting Enzyme 2) and infect (via a transformation to its fusion state) human epithelial cells (Type I and II pneumocytes; also, alveolar macrophage and nasal mucosal cells), but in the "Down-state" the Spike protein is believed to be inactive to ACE2 binding and to cellular infection. We note that the S1 domain of the Spike protein is shed in the transition from the prefusion state to the fusion state of this virion; those transformational aspects are not considered here.

The exact mechanism and specific structural details associated with the flexibility or local mobility of the RBD in the Up and Down states in SARS-Cov-2 remain unanswered. For example, it is not known whether these states exist simply randomly or by deterministic changes orchestrated by the virion or its environment. Recently unpublished long time Molecular Dynamics (MD) studies ($10\mu s$) of an isolated Spike Protein by the Shaw Group (4) noted that the protomers tended to persist in their initial states, i.e, Down states remain Down and Up states remain Up. However, the Up state protomer demonstrated further distal displacement and mobility from its initial state that was given by experimental structural data.

In order to better understand the differences between the Up and Down protomer states, we conducted an all-atom interacting energy landscape mapping of the entire Spike protein from its *.pdb (Protein Data Bank) structure file (6vsb.pdb) in order to identify interaction energy "hotspots" or "glue" points associated with relatively strong non-covalent atom-atom interactions between residues, which may be responsible for specific persistent domains of this complex trimeric protein. In doing so, we were able to identify some unique and potentially critical differences between the Up and Down protomers within the overall trimeric structure, including a possible molecular latch that helps to maintain the RBD in the down state conformation. The latch residues are conserved across the closely related virions SARS-Cov-1 and the bat corona virus

RatG13, as well as known variations of the novel corona virus. Comparative analyses between Up and Down state protomers, such as those given here, may lead to potentially new therapeutic targets aimed at preventing the attachment of the Spike protein to its host cell receptors.

2. Materials and Methods

Following our recently published study on $A\beta42$ amyloid fibrils (5), we analyzed the recently published trimeric structure of SARS-Cov-2 (PDB ID: 6vsb) according to the Coulombic (charge and partial atomic charge) and Lennard-Jones (Born and van der Waals forces) atom-atom interaction forces as laid out in the open-source energy mapping algorithm developed by Krall et al (6). This mapping algorithm efficiently parses the strongest non-covalent atom-atom interactions and their inter-atomic distances from structure file data according to empirically established criteria based on the $AMBER03$ force field model to ensure that all dominant interactions are accounted for. These energetic mappings, for example, allowed us to discern important structural differences leading to greater adhesive strengths of $A\beta42$ oligomers versus their $A\beta40$ counterparts. Following our previous studies, the parsing criteria were taken as the upper limit of $-0.1kT$ units for Lennard-Jones (van der Waals) criteria and $-0.3kT$ units for Coulombic interactions, although lower values can also be specified in the analysis part of the mappings in order to further refine the results (6).

The SARS-Cov-2 Spike protein structure consists of three chains or protomers (A, B, and C chains) of which the chain A is given in the so-called "Up" state of its RBD (6vsb.pdb), and chains B and C are in their "Down" state. We energetically mapped the interchain interactions "Up-Down" and "Down-Down" and specific domain interactions (*intra*chain interactions) for the Up and Down state protomers, including S1 and S2 domain interactions and sub domains of S1 that include the RBD domain.

3. Results

Interchain Interactions

Mappings of the dominant atom-atom interactions among the residues for the Up-Down (A-B) and Down-Down (B-C) states are illustrated in Figs. 1 and 2, respectively (Table Suppl.1 A-B and Table Suppl.2 B-C). As expected, the majority of the dominant interactions are within the S2 domain, and the entire three chain structure is greatly stabilized by this feature. There are a number of stabilizing interactions involving the SD1/SD2 domain of any protomer with the S2 and NTD domain of neighboring protomers (Supplementary Figs. Suppl.1 and Suppl.2). The pattern of the S1-S1 interchain interactions involving the RBD of any chain to the N-terminal domain (NTD) of its adjacent neighbor is potentially important as shown in Figs. 1 and 2 for clarity. We specifically examined the interchain interactions between the RBD with the NTD of its neighbor protomer for both the Up-Down (A-B) and Down-Down (B-C) interactions, shown in Figs. 3 and 4, respectively. Somewhat surprisingly, the Up-Down interactions were stronger and more numerous overall than the Down-Down interactions of these close neighbor domains. Additionally, the energetically dominant Up-Down RBD-NTD interactions involve three additional residues (ALA520-PRO521-ALA522) not seen in the Down-Down cross chain interactions. These

Figure 1: Dominant Energy Landscape for the Interaction of Chain A (Up; red) with Chain B (Down; blue). The dominant atom-atom interactions are shown by ball and sticks. Also shown is the overall chain interaction configuration looking at the trimer from the top view.

three residues, however, will be shown below to play a stabilizing *intra*chain role for the Down state protomer.

Intrachain Interactions

We next looked in detail at *intra*chain interactions for both the Up and Down protomers in order to attempt to discern any telling differences in these two states.

S1-S2 Intrachain Interactions

We first energetically mapped the S1 domain (residues 1-677; not including missing residues) to the S2 domain (residues 682-1273; not including missing residues) of both the Up protomer and Down protomer separately, as shown in Tables Suppl.3 and Suppl.4, respectively. Overall, we found no discernable differences in the energetic interactions between the Up and Down states, which again may not be surprising considering the RBD of S1 (residues 319-527) is distal to the S2 domain whether in the Up or Down protomer state. Additionally, the proximal S1 domain beyond the RBD, SD1/SD2 domains (residues 533-677), also showed no discernable differences between the Up and Down states in it's interaction with S2. We, therefore, next compared the SD1/SD2 domains to RBD domain interactions within each protomer in the Up or Down state, since these states have clear differences in separation distances owing to the more distal RBD in the Up state.

Figure 2: Dominant Energy Landscape for the Interaction of Chain B (Down; blue) with Chain C (Down; green). The dominant atom-atom interactions are shown by ball and sticks.

Figure 3: Dominant Energy Landscape for the Interaction of the RBD Chain A (Up;red) with NTD Chain B (Down;blue); oblique, top view. Only A Chain RBD residue/atoms are shown.

Figure 4: Dominant Energy Landscape for the Interaction of the RBD of Chain B (Down;blue) with the NTD of Chain C (Down;green); oblique, top view. Only B Chain residue/atoms are shown.

SD1/SD2-RBD Intrachain Interactions

Overall the energetic mappings between Up and Down states for these two S1 domains (RBD-SD1/SD2) appear very similar, as shown, for example, by the dominant partial charge interactions (Figs. 5 and 6; Tables Suppl.5 and Suppl.6) with one important difference as shown in the zoom illustration of Fig 7.

For the Down state protomer, we observe conspicuous, strong energetic interactions between the R group of GLN564 in the SD1 domain with backbone atoms of the three sequential residues ALA520-PRO521-ALA522 in the RBD domain, which are missing in the Up state mappings due to the more distal state of the RBD. These specific atom-atom interactions are shown in more detail in Figs. 8 and 9. Note the PRO521 interactions are dominated by van der Waals interactions in both instances.

Interestingly, these are exactly the same three residues of the Up-RBD that participate in the set of dominant energetic interactions with the Down-NTD of its neighbor protomer along the Up-Down interface (Fig.3 previously). Thus, these three dominant energy residues seem to play different roles depending on the Up or Down state of the protomer, which was somewhat surprising.

Figure 5: Dominant Energy Landscape for partial charge interactions between SD1/SD2 and RBD of Chain A (Up).

Figure 6: Dominant Energy Landscape for partial charge interactions between SD1/SD2 and RBD of Chain B (Down).

Figure 7: Zoomed Dominant Energy Landscape for partial charge interactions between SD1/SD2 and RBD of Chain B showing the conspicuously unique residue interactions. Note that dominant van der Waals interactions invloving PRO 521 backbone atoms are not shown here; see Tables Suppl.5 and Suppl.6.

Figure 8: Dominant Energy Landscape for the partial charge interactions between SD1 and RBD of of Chains A and B showing a possible latch for the B Chain.

Figure 9: Zoomed View of possible molecular latch between the RBD and SD1 domain: ALA520 PRO 521 ALA 522 with GLN 564.

GLN564 --- ALA520 PRO521 ALA522
interactions off

Control --- all interactions on

Figure 10: Initial dynamic test of possible latch over a period of approximatley 1 nsec. Blue-initital state; Red-final state. With latch residues "off", SD1 movement begins the detachment process within 1 nsec.

Initial Dynamic Test of Latch

In order to determine if these strong, but limited residue interactions could possibly function as a molecular latch for the Down state protomer, we conducted some preliminary all-atom implicit solvent simulations, as described in detail previously (7) of the S1 domain for the Down state protomer under two conditions. In the first condition (Control) simulations were done for the full set of residues in the S1 domain without modifications. We used the Charmm-Gui (8) to insert all missing residues from the S1 domain from 6vsb.pdb structure file. For the second simulation we used the same initial structure as Control, but we artificially turned off the partial charge and Lennard-Jones interactions between *and only between* the four participating latch residues. Preliminary results for a short time period demonstrate that the SD1 domain begins to separate from the RBD domain (Fig. 10), as expected from the energetic mapping identification of this "glue" point. Clearly, however, longer time dynamic studies are needed to fully address the functions of this possible latch as well as its dynamic transformations.

4. Discussion

The all-atom energetic mappings are shown to be useful in attempting to understand the overall functionality of the Spike protein of the novel corona virus. The S2, fusion machinery domain, for example, also acts as a rigid base for the more flexible S1, prefusion domain, where flexibility is an important part of its function. The RBD is tethered to the SD1/SD2 domain by two peptide strings: one from the NTD (approximately residues 327-335) and the other connected to the SD1/SD2 domains (approximately residues 527-535). These peptide "strings" are stabilized or "tied down" by beta strand structures near the S1/S2 interface. The RBD domain in the Down state protomer is partially stabilized by the dominant energetic interactions of the backbone atoms of three of its residues (in sequence) with the R-group of a single residue from the SD1 domain, which we have called a possible latch. Ironically, these same three residues are involved in helping to stabilize the Up state protomer by interactions with the NTD of its Down state neighbor. This study raises a number of new questions about the origins of the Up and Down state protomers, including the reasons behind the occurrence of the Up state protomer, since as we have shown here, the Down state is partially stabilized by not only the potential latch, but also neighboring chain interactions between the RBD and it's NTD neighbor. It seems reasonable, however, that the process of "Down to Up" is irreversible in the absence of deliberate outside interference. We further note that the possible three residue latch of the RBD Down state (ALA520 PRO521 ALA522) along with GLN654 of SD1 are conserved across reported variants for SARS-Cov-2 (1). All latch residues are also present in the closely related virions: SARS-Cov-1 and the bat corona virus RatG13; the latter being 96 % homologous to SARS-Cov-2 (1). It is important to point out, however, that these findings are based on given, experimental static structure data and may not accurately reflect the dynamic states of this complex protein. More complete biocomputational dynamic and in-vitro biological studies (via single site mutagenesis, for example) are necessary to develop a complete picture of the mechanisms surrounding the Up and Down protomer states, including the effects of the entire trimeric structure.

5. Acknowledgments M.H.P. would like to thank the Office of the Vice President for Research, the Department of Chemical and Life Science Engineering, and the Center for High Performance Computing at Virginia Commonwealth University for partial support of our overall research efforts surrounding Covid-19. O.B. was supported by the National Institutes of Health Institutional Research and Academic Career Development Award K12GM119955. M.H.P and O.B. would like to acknowledge future resources services, and support provided via the COVID-19 HPC Consortium (https://covid19-hpc-consortium.org/), which is a unique private-public effort to bring together government, industry, and academic leaders who are volunteering free compute time and resources, in support of COVID-19 research. Specifically, resources supporting this continued work are provided by the NASA High-End Computing (HEC) Program through the NASA Advanced Supercomputing (NAS) Division at Ames Research Center.

Open Source Software. OpenContact is freely available under the Third-Party Software Tools listings of the Protein Data Bank: https://www.rcsb.org

References

1. Wrapp, D., Wang, N., Corbett, K.S., Goldsmith, J.A., Hsieh, C.-L., Abiona, O., Graham, B.S., and McLellan, J.S. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation, *Science*, 367, 1260-1263, 2020 DOI 10.1126/science.abb2507

2. Walls, A.C., Park,. Y.-J., Alejandra Tortorici, M., Wall, A., McGuire, A.T.,Vessler, D., Structure, function and antigenicity of the SARS-Cov-2 spike glycoprotein, BioRxiv, 2020 https://doi.org/10.1101/202

3. Yan, R., Zhang, Y., Li, Y., Xia, L., Guo, Y., and Zhou, Q., Structural basis for the recognition of the SARS-CoV-2 by full-length human ACE2, *Science*, 367, 1260-1263, 2020 DOI 10.1126/science.abb2762

4. D. E. Shaw Research, "Molecular Dynamics Simulations Related to SARS-CoV-2," D. E. Shaw Research Technical Data, 2020. `http://www.deshawresearch.com/resources_sarscov2.html`

5. Bastidas,O.H., Green, B., Sprague, M., and Peters, M.H, Few Ramachandran Angle Changes Provide Interaction Strength Increase in $A\beta 42$ versus $A\beta 40$ Amyloid Fibrils, *Sci. Rep.*, 6, 36499, 2016. DOI: 10.1038/srep36499.

6. Krall,A., Brunn, J., Kankanala, S., and Peters, M.H. A Simple Contact Mapping Algorithm for Identifying Peptide Mimetics in Protein-Protein Interaction Partners. *Proteins. Structure, Function and Bioinformatics*, 82, 2253-2262, 2014. DOI:10.1002/prot.24592.

7. Peters M. H. Langevin dynamics for the transport of flexible biological macromolecules in confined geometries. *The Journal of Chemical Physics*, Jan 14;134(2):01B615, 2011.

8. S. Jo, T. Kim, V.G. Iyer, and W. Im CHARMM-GUI: A Web-based Graphical User Interface for CHARMM. J. Comput. Chem. 29:1859-1865, 2008.

Supplementary Information

Fig. Suppl.1

Fig. Suppl.2

Tables Suppl.1 - 6