Social hierarchy is established and maintained with distinct acts of aggression in male *Drosophila*

Jasper C. Simon[*] & Ulrike Heberlein

Janelia Research Campus, HHMI, Ashburn, VA 20147, USA

[*]Correspondence: *simonj10@janelia.hhmi.org*

ABSTRACT

Social interactions pivot on an animal's experiences, internal states, and feedback from others. This complexity drives the need for precise descriptions of behavior to dissect the fine detail of its genetic and neural circuit bases. In laboratory contests, *Drosophila* males reliably exhibit aggression, conventionally scored using lunges as a proxy, allowing its study with the many available genetic tools. Here, we use an explicit approach to identify the onset and reversals in hierarchical relationships and observe that distinct aggressive acts precede, concur, and follow dominance. We find that lunges are inadequate in establishing social dominance. Lunges rather appear to reflect the dominant state of a male and help in maintaining his social status. Lastly, we characterize the recurring structure of aggression that emerges through subsequent reversals in dominance. Collectively, this work provides a framework for studying the complexity of agonistic interactions in flies enabling its neurogenetic basis to be understood with precision.

INTRODUCTION

Agonistic interactions important for establishing social hierarchies over food, territory, or

mates often progress as a sequence of stereotyped acts. Like other goal-driven

behaviors [1], the steps along this sequences are triggered by context and modified by

past experience [2,3], through largely unknown mechanisms that are coupled to an

animal's internal physiological state [4]. Determining these mechanisms remains

difficult, however, because social exchanges proceed as a dynamic, reciprocal, and

continual feedback loop of interactions between two (or more) decision-making

individuals that progresses over both moment-by-moment and protracted times scales

[5]. Intra-male adversarial contests in the fruit fly *Drosophila melanogaster* seem well

suited for studying such complex social phenotypes, due to its established study in a

laboratory settling [6,7] and the multitude of techniques available for manipulating and

recording gene and neuron function [8]. Nonetheless, efforts to explain which acts lead

to dominance hierarchies [9,10] have suffered from a lack of consensus on methods

and results for studies of aggression (as discussed within [11,12]).

Here we use an explicit approach to characterize the temporal relationship among

various aggressive acts and the establishment of social hierarchy between pairs of male

flies in arenas previously developed for high-throughput phenotyping [13,14]. Using this

approach, we observe that the acts of fencing, boxing, and tussling appear well

positioned in time to be related to the establishment of social hierarchy, whereas

lunging and chasing its maintenance. Unexpectedly, the commonly studied aggressive

act of lunging, near-universally, follows the onset of social hierarchies and thus is

unlikely involved in its establishment. We carried out a simple social manipulation and

observed that males in contests between mixed pairs of genotypically high versus low

lunging lines became dominant at chance levels. This result suggests that the total

number of lunges executed by an individual is inadequate in explaining his social

outcome. We further observed that lunging persists as dominant males confront

unfamiliar opponents in unexplored settings. Lastly, we report that a faction of the

aggressive acts surveyed also intensify through subsequent reversals in social

dominance adding to a fuller appreciation of this complex, recurrent social exchange.

Together, this work provides a framework for untangling which aggressive acts causally

relate to the establishment of social dominance and those that are a consequence

RESULTS

Identifying hierarchical relationships

Upon introduction to experimental arenas, pairs of males whom had never fought

(henceforth, "naïve males") display independent and haphazard movement that may

include one or both males jumping, taking flight, and climbing the wall as they attempt to

escape. With time males settle into a period of exploration during which they eventually

meet, and commonly this initial contact leads to bouts of short, alternating pursuits that

are characteristically restricted to the floor (which is entirely made up of food; see

Methods). Predictably this engagement then progresses to a situation where one male

consistently pursues the other with the retreating male leaving the floor (resource)

presumably attempting to flee by climbing the wall. It is this explicit characteristic

sequence – of one male pursuing and a second attempting to climb the arena wall –

that we use for identifying when a hierarchical relationship is established (see **Figure 1**

and **Movie 1**).

Distinct aggressive acts precede, concur, and follow establishment

We observed adversarial contests in experimental arenas that have been widely used to

study aggression [13,14]. These arenas increase the frequency of flies' interactions by

restricting their movements near a food resource and yet retain an uncluttered

background (no food cup, no decapitated female, nor region of the chamber that is

blocked from camera view [15]). This arrangement permits methods based on computer

vision to help automatically annotate various aggressive acts [13], followed by direct

inspection for correcting erroneous annotations and identifying the onset of dominance.

Together this complimentary workflow allows a robust, high-resolution portrayal of the

aggressive acts displayed during the progression of adversarial contests (see **Figure**

S1 and **Table S1** for details of annotation and software used).

Naïve males were introduced simultaneously on food as pairs. This lessened the

disparity in which individuals discovered the contested resource and allowed

uninterrupted observation of their interactions leading up to and following the onset of

social dominance. We considered "Wing flick" and the following acts reported to span

the range of aggressive intensities (from low to high): "Fence," "Chase," "Lunge," "Box,"

and "Tussle" [6] (for descriptions of each act see **Table S1**). At times we found boxing

and tussling difficult to separate, and thus for ease of scoring they were combined into

the single category, "Box, Tussle" (as done by others [6]).

From both individual contests and aggregate data, we observed reproducible patterns of

behavior (see **Figure 2**). Males consistently displayed the highest levels of the head-to-

head interaction, "Fence," from the beginning of contests until the onset of dominance.

Thereafter, we observed only sporadic displays of fencing with no clear temporal

structure. In contrast, the other head-to-head interactions, "Box, Tussle," appeared

abruptly, peaked, and then sharply decreased immediately prior to the onset of

dominance (see **Figure 2D**), and on occasion also transiently reappeared preceding

subsequent reversals in social status. The wing display, "Wing flick" was observed more

broadly, frequent at a low level from the beginning of contests and making a salient

uptick near the onset of dominance with males continuing this escalated flicking

throughout the remainder of the contest. In contrast, displays of "Lunge" were rarely

observed before the establishment of dominance, and began and continually increased

in intensity thereafter. Lastly, occurrences of "Chase" were never recorded prior to

dominance and emerged well after establishment (as reported previously [6]). Together,

by aligning recordings from individual contests to the onset of social dominance, we

expose which acts may causally relate to its establishment and those apparently more

related to its maintenance.

Lunges are inadequate for explaining hierarchical outcomes

The act of lunging is commonly used for studying aggression (see reviews [11,16]), with

several studies suggesting they play a pivotal role in establishing dominance [9,17,18].

We, however, observed that lunging emerged after the establishment of dominance (as

seen in **Figure 2**). To confirm this finding we ran a second, independent group of

contests and carefully reviewed the incidences of lunging (both direct inspection with

slow playback to correct for false negative displays and semi-automatically for false

positives; see **Methods** for details). Again males lunged after the establishment of

dominance, with the majority of the lunges executed by the dominant male (see **Figure**

S3).

From the observation that lunging occurs after the onset of dominance we reasoned

that they were unlikely to play a role in its establishment. If this were true executing

more (or less) lunges should not influence whether an individual emerges dominant. To

test this, we set up contests with mixed pairs of males from genotypically high and low

lunging lines and measured the total number of lunges executed by those males that

emerged dominant. We saw no influence of total number of lunges on dominance

outcomes, even when males from mixed pairs executed dramatically lopsided amounts

of lunging (see fourth columns in **Figure 3A, B**; and **Figure 3 – Supplement 1**,

Methods, and **Table S2** for additional examples and details regarding the various fly

lines). It was therefore unsurprising to observe that males from both the high and low

lunging genotypes also executed lunges following the establishment of dominance (see

Figure S4). Collectively, these results provide evidence that lunges do not establish

dominance, and more likely play a role in its maintenance.

Males assert dominance by lunging at familiar and unfamiliar opponents

For the majority of contests, we observed that only a single, stable hierarchical

relationship formed between pairs of naïve males (as shown in **Figure 2** and **Figure**

S3). In approximately one in five contests, however, after the establishment of

dominance there were subsequent reversals in social status, with lunges (sometimes by

both males) clustering near each reversal. To better understand how these lunges

relate to reversals in status, we determined the identities of males executing each lunge

(see **Methods**). The majority of the lunges preceding a reversal in status were

displayed by the currently dominant male; whereas, dominant males executed nearly all

of the lunges following reversals (see **Figure S3**). This analysis provides evidence that

lunging reflects current and likely past dominance status, and further support the

interpretation that lunges play a role in maintaining a male's social status.

Both increased total time holding dominance and also the occurrence of reversals in

social status contributed to a persistent high level of lunging by individuals against

familiar opponents. Moreover, it has been reported that "winner" males lunged and

became dominant earlier in contests against naïve opponents [19]; and also, that males

appeared to change their fighting tactics as a consequence of winning or losing [20].

Therefore, to test if lunging functions as a general tactic for holding on to dominance

irrespective of opponent or location, we paired the emergent dominant males (from a

first contest between two naïve males; see **Figure 4A**) with naïve males in unexplored

arenas (see **Figure 4B**). The most salient feature of these second contests was that

unlike for pairs of naïve males, that rarely lunged before the establishment of

dominance, the majority of dominate males (although not their paired naïve opponents)

lunged prior to establishing dominance (see **Figure 4C**). The temporal sequence for all

other aggressive acts studied (including lunging after the initial establishment of

dominance) appeared comparable to those observed in contests between two naïve

males (data not shown). Finally, to examine if the precocious lunging helped dominant

males reassert dominance against unfamiliar males in unexplored arenas, we quantified

the number of contests in which dominant males that had lunged early were successful

in reasserting their dominance status. For the majority of these contests, previously

dominant males reasserted their dominance (see **Figure 4D**).

DISCUSSION

Social dominance forms when individuals yield to the agonistic advances of others,

often in the context of conflict over resources such as access to food, territories, and

reproductive opportunities [21]. However, the relationship between specific aggressive

acts and the establishment of social hierarchies remains contentious largely for they

appear so highly correlated, even with disagreements over which drives the other [22].

In this work, we use an explicit approach to characterize with high temporal resolution

how a range of known acts of aggression relate to the establishment, maintenance, and

subsequent reversals of social hierarchy.

The approach introduced within this work has several advantages. It is easy to describe

and straightforward to implement. It accommodates that adversaries compete over

resources and includes an operationally defined 'escape,' both which are important

ethological considerations [6,23]. Further, it provides the means to align and thus

compare the onset of dominance, which varies in time across contests due to

differences in various influences (e.g., inherited factors, past social histories, current

physiological states, handing by the experimenter, environmental conditions, and

feedback between adversaries). Lastly, it can be used to update the classification of

dominance if and when reversals in status occur after an initial hierarchical relationship

has been established.

A significant challenge to studying complex social behaviors such as those related to

the establishment of social hierarchy is that the numerous interactions possibly

regulating them are often highly correlated with outcomes. By using criteria to identify

the establishment of dominance that are independent from the particular acts of

aggression under study, we avoid a circular definition of social hierarchy, and thus allow

for stronger claims of causality between specific acts and its emergence (for a graphical

summary see **Figure 5**).

Unexpectedly, we show that near-universally only once males hold (or have held)

dominance do they lunge, and moreover they continue lunging against familiar

opponents even after a dominance hierarchy is settled (as has been described

previously [9,24]). From this and our observation that dominant males precociously

lunge and successfully reassert dominance over unfamiliar males in novel settings, we

propose that lunging is likely uninvolved in establishing dominance, but rather

symptomatic of a changed internal state [25] and plays a role in maintaining an

individual's social status.

The complexity of reversals in dominance requires more study. From a preliminary

characterization of a larger number of contests with reversals, we observed that lunging

appears as just one of several interacting aggressive acts likely involved in the process

(see **Figure S5**). Taken together, it is clear that further consideration is needed for a

more complete understanding of these dynamic, reversing, and escalating exchanges.

Collectively, this work advances our understanding for the causal relationships within

this complex behavioral phenotype and should prove helpful for both refining the roles

of known [6,26,27] and yet to be discovered acts of aggression. Finally, the approach

described, when used in conjunction with high-throughput screens, provides a

framework for helping to identify and clarify the genetic and neural mechanisms

underlying the coordination of behaviors involved in establishing hierarchical social

relationships in insects.

FIGURE LEDGENDS

Figure 1. Identifying social hierarchy within arenas used for studies of

aggression. (A) Behavior independent of specific aggressive acts used to determine

when social hierarchy forms between pairs of males. Pursuing males (red, solid arrow),

that often proceed to guarding the floor (red, asterisk), are classified as dominant and

those then attempting to escape by climbing the wall of the arena as subordinate (black,

dashed arrow). (B) Twenty-minute contests ordered as rows from early to late

establishment. Using the "Pursue-to-Climb" criteria clear hierarchies form in the majority

of pairings. Periods of unestablished social standing (light yellow bars) precede clear

social hierarchy (gray bars). Contests in which hierarchies formed prior to observation

(full gray bar, top row, n=12) and those in which hierarchies never formed (full yellow

bar, bottom row, n=6) were excluded from further analysis.

Movie 1. Established hierarchical relationship. Example movie from a 20-minute

contest showing an established hierarchical relationship, wherein a subordinate male

repeatedly attempts to climb the wall of the arena while a dominate male (clipped wing)

guards the floor.

Figure 2. Fence, box, and tussle interactions precede the establishment of social

hierarchy, whereas the majority of wing flicks, lunges, and chases follow. (A)

Raster plot from a single contest illustrates the relationship between the observed

reproducible sequence of aggressive acts and the establishment of social hierarchy.

Each row and color corresponds to a different act. (B) Raster plots show the temporal

structure for the various acts (discrete act = black tick). Individual contests (shown as

rows) between pairs of naïve males are ordered top-to-bottom by latency until

establishment (red ticks). After establishment males exhibit clear hierarchical

relationships (gray shading), with occasional reversals in dominance (blue ticks). (C)

Peri-event plots with contests aligned by establishment (vertical, red line). Aggressive

acts from each contest are binned into one-minute intervals and displayed above plots

with corresponding collective medians (black line) and interquartile ranges (gray

envelope). (D) Peri-event plots aligned to include the minute immediately preceding and

following establishment (vertical, red line). Discrete acts are displayed above associated

collective medians (black line) and interquartile ranges (gray envelope). (C, D)

Statistical comparisons for the frequency of acts within five- (C) and one-minute (D)

windows prior to and following establishment are noted above plots (horizontal, green

lines). In all cases the Wilcoxon signed rank test was used to assess whether

differences in the observed acts exist between paired, equal-length windows of time

before and after the establishment of social hierarchy. See **Figure S2** for details

regarding data exclusion. Common nomenclature for statistical significance used

throughout study: not significant (n.s.), *p<.05* (*), *p<.01* (**), *p<.001* (***), *p<.0001* (****),

p<.00001 (*****).

Figure 3. Number of lunges executed by dominant individuals is inadequate for

explaining hierarchical outcomes. (A) Focal males (F, above) and opponents (O,

below) were paired and the percentage of contests in which males became dominant

are shown within each stacked bar chart. Focal males from genotypically low (L, blue),

standard (S, green), and high (H, light red) aggressive lines became dominant at

chance levels when paired with same-genotype opponents. The tested pairing between

mixed high (H, light red) and low (L, blue) aggressive males also became dominant in

comparable amounts (shaded in green). In contrast, for contests in which high-

aggressive focal males (H, light red) were paired with socialized opponents (group

housed) from the same genotype, focal males became dominant 100% of the time

(compared to chance levels; Fisher exact, *p=.0005*; right-most bar). In all cases males

were socially isolated unless noted. (B) Medians with interquartile ranges for the total

number of lunges executed by dominant individuals from the contests reported in "A"

above. The number of lunges executed by dominant focal males (F) from low (blue),

standard (green), and high (light red) aggressive lines were no different than when

those of paired opponents (O) from the same genotypes became dominant. Focal

males from a high-aggressive line (F, light red) displayed a comparable, high level of

lunging when paired with all opponents (from left to right): males from the same

genotype (O', light red), genetically low-aggressive males (O'', blue), or socialized

males (O''', light red) from the same genotype (Kruskal-Wallis, H(3)=5.92, *p=.1142*).

However, in tested pairings the levels of lunges executed by the dominant, high-

aggressive focal males (F, light red) were significantly greater than those of dominant,

low-aggressive opponents (O'', blue; Wilcoxon Rank-Sum, Z=3.9244, *p=.00009*; shaded

in green) and socialized opponents from the same high-aggressive line (O''', light red;

Wilcoxon Sign, *p<.00001*).

Figure 3 – Supplement 1. Independently isolated genotypically low, standard, and

high aggressive lines identified from a P-element screen; additional genotypically

high and low aggressive lines corroborate findings in Figure 3. (A) The distribution

of P-element lines screened before outcrossing is skewed right (black, stair-step line)

with the median (vertical, white line) and interquartile range (gray shading) centered

around 67 total lunges, as calculated by summing the total lunges executed by pairs of

same-genotype males from 20-minute contests. The median number of lunges

executed by the genotypic standard (green) used throughout the study and also the low

(blue, light blue) and high (light red, red) lines are shown after outcross. (B) Focal males

(F, above) and opponents (O, below) were paired in various combinations and the

percentage of contests in which males became dominant are shown within each

stacked bar chart (adversaries are coded by letter and color as indicated in "A" above).

Focal males became dominant at chance levels in all contests. (C) Lunging executed by

dominant males in both same- and mixed- genotype pairings were consistent with those

reported in **Figure 3**. Dominant focal or opponent males arising from same-genotype

pairs displayed equivalent amounts of lunges (F vs. O, light blue and red). Focal males

from an additional high-aggressive line (F, red) displayed a comparable, high level of

lunging regardless of opponent (Kruskal-Wallis test; H (1)=0.0039, *p=0.9503*). However,

in tested pairings the levels of lunges executed by both dominant, high-aggressive focal

males (F, light red; F, red) were significantly greater than an additional dominant, low-

aggressive opponent (vs. O', light blue; Wilcoxon Rank-Sum, Z=3.8884, *p=.00010* and

versus O'', light blue; Wilcoxon Rank-Sum, Z=3.8111, *p=.00014*).

Figure 4. Lunging persists as dominant males confront unfamiliar opponents in

unexplored arenas. (A, B) Raster plots showing the temporal structure of lunges

executed by naïve (black ticks) and previously-dominate (red ticks) males in relation to

establishing social hierarchy. Individual contests (shown as rows) between pairs of

males are ordered top-to-bottom by latency until establishment. Periods preceding

establishment (yellow shading), and the prevailing dominance status for naïve (gray

shading) and previously-dominant (light red shading) males are noted as observed

within 20-minute contests. (A) Pairs of naïve males. (B) Dominate males paired with

naïve opponents in a second, unexplored arena. (C) The number of contests in which

the previously-dominant males (red bar; "Dm") lunged prior to reasserting dominance

over naïve males was significantly greater than those in which randomly selected

individuals from pairs of naïve males (black bar; "N") lunged prior to establishing

dominance (Chi-Square: X^2 (1, N=56) = 16.07, *p=.00006*). (D) The number of contests

where previously-dominant males (light red bar; "Dm") lunged prior to reasserting

dominant was also significantly greater than those in which they lunged and failed (gray

bar; "N") and became subdominant (Chi-Square: X^2 (1, N=25) = 4.89, *p=.02701*).

Figure 5. Sequence among aggressive acts in relation to establishing social

hierarchy. Depiction of the reproducible sequence among distinct aggressive acts as

observed in relationship to the emergence of social hierarchy. After adjusting to the

experimental arena, pairs of males typically recognize each other as opponents.

Successively they then exhibit fencing (red line) and then display low levels of wing

flicks (purple line). As contests escalate (graded yellow box) the frequency of fencing

bouts lessens and is replaced by boxing and tussling (orange line) concomitantly with

an increasing number of wing flicks. Transiently the amount of boxing and tussling

peaks and then sharply drops immediately preceding the establishment of hierarchical

relationships (time = 0). Lunging (blue line) follows establishment, is the principal

aggressive act during the maintenance of dominance (gray box), and is accompanied

by a steady display of wing flicks. Finally, increased displays of chase (green line)

observed after some time appear to connote stabilized hierarchical outcomes.

Figure S1. Details for including and classifying fence versus box, tussle

interactions. (A) Illustration drawn from a top viewing angle to clarify the criteria for

inclusion. For an interaction to be included the focal male must touch the front legs or

head of the opponent (blue dashed box). (B) Side views showing the postural difference

between "Fence" and "Box, Tussle." Interactions that included touch from a prone

stance were classified as "Fence." Any touching when one or both males exhibited an

erect posture with both front legs lifted off the floor were classified as "Box, Tussle."

Interactions in which males disengaged, directly executed a 360-degree rotational turn,

and then reengaged were annotated as a single continuous act.

Figure S2. Example data to show the requisite exclusions for balanced statistical

analyses. An example dataset shows the data used for statistical analyses. In order to

apply paired statistical tests, only contests allowing five minutes or more (always equal)

windows of time before and after the establishment of social hierarchy were used unless

otherwise noted. In the example shown only contests with five minutes of recorded data

before and after establishment were used (blue shaded box, above; median frequency

of behavior for corresponding period denoted as bold, blue line, below). In this example,

data recorded earlier than five minutes preceding and following establishment and also

the data from the entire first five and last two contests were excluded. The exclusions

occasionally decreased the sample size, and thereby the statistical significance, yet

never qualitatively changed results.

Figure S3. Males assert dominance by lunging. (A) Raster plot displaying the

temporal structure of lunges executed by transiently dominant males (red ticks) and

those observed dominant at end (black ticks) in relation to the establishment and

reversals in social hierarchy. Individual contests (shown as rows) between pairs of

males are ordered top-to-bottom by latency until the establishment. Periods preceding

establishment (yellow shading), ultimately transient (light red shading), and of final

dominance status (gray shading) are noted as observed within 20-minute contests.

Terminal reversals in social status for each contest are indicated with blue ticks (ticks

for intermediate reversals were excluded for clarity). (B) Lunging reflects dominance.

Uncommonly lunges were observed before establishment (bar plots highlighted by

yellow shading). Prior to terminal reversals (vertical, blue line), the currently dominant

male executed the majority of the lunges executed. Thereafter, near-universally, all

lunges came from dominant males (D, black bar), and subordinate males (t, red bar)

displayed almost none (Wilcoxon rank sum, Z=7.1057, *p<.00001*). This figure includes

the identities for which male lunged and also denotes any reversals in social status

between the pairs of naïve males as observed within the contests reported in **Figure**

4A.

Figure S4. Lunges are executed after establishing dominance irrespective of

genotype. (A) Raster plots show the temporal structure for lunges executed by males

from genotypically high (top) and low (bottom) aggressive lines (lunge = black tick).

Individual contests (shown as rows) between pairs of naïve males from the same

genotype (high vs. high and low vs. low) are ordered top-to-bottom by latency until

establishment (red ticks). After establishment males exhibit clear hierarchical

relationships (gray shading). (B) Peri-event plots with contests aligned by establishment

(vertical, red line). Lunges from each contest are binned into one-minute intervals and

displayed above plots with corresponding collective medians (black line) and

interquartile ranges (gray envelope). (C) Peri-event plots aligned to include the minute

immediately preceding and following establishment (vertical, red line). Lunges are

displayed above associated collective medians (black line) and interquartile ranges

(gray envelope). (B, C) Statistical comparisons for the frequency of lunges within five-

(B) and one-minute (C) windows prior to and following establishment are noted above

plots (horizontal, green lines). In all cases the Wilcoxon signed rank test was used as

described in **Figure S2**.

Figure S5. Recurring aggressive acts intensify through subsequent reversals in

dominance. (A) Schematic illustrating the establishment and subsequent reversals in

social dominance in contests between pairs of naïve males. The gray, switching line

indicates the state of dominance with colored arrows indicating the onset (white dot) for

the establishment (Initial, black), subsequent reversals (Intermediate, blue), and the

final hierarchical outcome (Terminal, red). (B) Idealized results for the observed

changes of aggressive acts among establishment (black line) and intermediate (blue

line) and terminal (red line) reversals of social dominance (gray shaded box). A value of

"1" indicates the highest level for any particular act to help clarify its relative increase or

decrease; drawings are based on the data in **Figure 2**, those reported here in: "C-E,"

and observations of contests using males from the genotypically high and low

aggressive lines (data not shown). Depicted in the top panel of "B," early on males

exhibit the highest levels of "Fence," which then drops during the minute immediately

preceding the initial establishment (black line). From then on only intermittent and low

levels of fencing were observed through subsequent reversals (Intermediate, blue and

Terminal, red lines). In contrast, as shown in the next panel down, the amount of "Box,

Tussle" peaks during the minute immediately preceding the initial establishment (black

line), and then peaks again and again, increasing and broadening through subsequent

reversals (Intermediate, blue and Terminal, red lines). (C-E) Peri-event plots centered

on the minutes immediately preceding and following the initial establishment (C; vertical,

black line), the intermediate subsequent reversals (D; vertical, blue line), and final

outcome (E; vertical, red line) of social hierarchy. Discrete acts are displayed above

associated collective medians (black line) and interquartile ranges (gray envelope). (C-

E) Statistical comparisons for the frequency of aggressive acts within one-minute

windows prior to and following establishment are noted above plots (horizontal, green

lines; with ±10-second window comparisons denoted within parentheses). In all cases

the Wilcoxon signed rank test was used as described in **Figure S2**. Lunges increased,

formed clusters, and the peaks of clusters shifted forward in time relative to each

subsequent reversal (C-E; shaded, green box). Lunging increased in a scalable manner

(One-way ANOVA: df(2), F(22.5470), *p<.00001* and followed by post hoc comparisons:

Initial versus Intermediate, *p=.0135*, Intermediate versus Terminal, p=.*0475*, and Initial

versus Terminal, *p<.00001*; all with Bonferroni correction). The shift in peaks was

quantified as a procession index; PI = (Lunges *minute preceding* – Lunges *minute*

following) / (Lunges *minute preceding* + Lunges *minute following*). One-way ANOVA:

df(2), F(6.1421), *p=.0049* and followed by post hoc comparisons: initial versus

Intermediate, *p=n.s.*, Intermediate versus Terminal, *p=n.s.*, and Initial versus Terminal,

p=.0046; all with Bonferroni correction). The intensity for several of the aggressive acts

increased alongside that of lunges. Collectively, the characterization exposes a

recurring structure that emerges during protracted engagements – one in which both

males reciprocally escalate fights just prior to reversals in status – seemingly in attempt

to maintain or reclaim social dominance. Data within this figure come from contests

containing reversals in dominance (n=25) identified from 146 adversarial contests.

TABLES

Table S1. Details for studying the aggressive acts reported within manuscript.

Descriptions and process for generating the data used to characterize the temporal

structure of the various aggressive acts reported within manuscript.

Act	Description	Method	Annotation	Software
Fence	Extending one or both forelegs and pushing against the forelegs or head of opponent from a prone position.	Manual	See **Figure S1**.	VCode [28]
Box, Tussle	Rearing up on hindlegs while striking, pushing, or pulling opponent with forelegs.	Manual	See **Figure S1**.	VCode [28]
Wing flick	Brief spreading followed by furling of one or both wings.	Automatic	Left and right wings flicks from both focal and opponent males outputted by CADABRA were combined (using Matlab) as follows: wingFlick = sort([wing.fli.r.obj1.t wing.fli.l.obj1.t wing.fli.r.obj2.t wing.fli.l.obj2.t])	CADABRA [13]
Lunge	Forward thrust of body with outstretched forelegs while snapping down on opponent.	Semi-Automatic	Assigning identification to time-stamped lunges classified by CADABRA (see Supplementary Tables 2,3 therein for details) provided information for which male executed a particular lunge.	CADABRA [13] followed by scoreMoviesLunges [29]
Chase	Running after opponent.	Automatic	First male following another at a speed greater than 5mm/s for ≥1s (see Supplementary Table 9 within [13] for details).	CADABRA [13]

Table S2. Behavioral measures for the out-crossed P-element lines used within

manuscript. P-element lines ordered by row from fewest to greatest number of lunges

(mean ±quartile) per 20-minute contest as identified within screen after outcrossing.

Line names indicate batch followed by trial number and come from the labelling scheme

of our laboratory. Color and intensity may be used to associate lines with results from

Figure 3 and **Figure 3 – Supplement 1**. In addition to its high levels of lunge displays,

the line 9.3 was chosen for its high amounts of chase (see below). Consequently, to

support the notion that chase events follow the establishment of social hierarchy as

observed in **Figure 2**, we analyzed chase events in contests between pairs of males

from 9.3 and also in mixed contests between males from 9.3 and the those from the low

aggressive line 11.261. In both cases, we observed that all chase events occurred

following the establishment of hierarchy (compared to equal numbers of chase events

before and after establishment; 9.3 versus 9.3, Fisher exact, n=18; *p=.0010*; 9.3 versus

11.261, Fisher exact, n=11; *p=.0351*). Together, combining the above results with those

from **Figure 2**, we report that we have only observed chases following the

establishment social hierarchies, Fisher exact, n=34, *p<.00001*.

Line	Color	Intensity	Sample size (n=contests)	Tussle	Wing flick	Lunge	Chase
11.261	Blue	Low	62	1 (0/4.3)	66.5 (12/128)	12.5 (1/26)	0 (0/1)
15.46	Light blue	Low	64	1 (0/4)	66 (24.5/213.8)	16 (6.5/37.5)	0 (0/1)
5.116	Green	Standard	301	3 (0/8)	95 (57/149)	48 (23.5/104)	0 (0/3)
9.3	Light red	High	24	2 (0.9/7.6)	138 (73/212.6)	184 (98.5/236)	4 (0/16)
11.27	Red	High	52	3 (1/8.3)	138 (85.9/214.9)	277 (186/391)	0 (0/2)

MATERIALS AND METHODS

Fly lines

All *Drosophila* lines originate from a P-element collection generated by the Heberlein

laboratory that were then outcrossed for at least six generations to w^{1118} Berlin.

Animal rearing

To lessen developmental heterogeneity, males used in experiments were collected from

lines maintained with controlled densities (by seeding vials with 5 males crossed to 10

females and removing these parental animals after 3 days), reared in customary 8-dram

plastic vials on standard media (cornmeal/yeast/molasses/agar), maintained at 25°C

and 65% relative humidity, and entrained on a 12:12-hour light-dark cycle. The lights-on

phase started at 1pm EST. Transitions between dark and light were immediate.

Animal handling

To model a uniform, probable 'ecological baseline' amount of social experience

including to have had mated, males for experiments were collected after 7 hours, yet

within the first 24 hours following eclosion using CO_2. For experiments in which the

identity of individuals were required, during collection, a small section of wing from one

of an eventual pair was removed, a procedure shown inconsequential to the outcomes

of adversarial contests [29].

Animal housing

In order to increase or suppress the level of aggression, males were housed individually in 10-mm diameter x 75-mm tall glass tubes (Fisher, Waltham, MA) or as groups of 15 males in 8-dram vials [30,31]. For both housing conditions flies had *ad libitum* access to standard media and were incubated in the same conditions as they were reared.

Animal observation

Experiments were performed on 4~6-day old males. Trials began after a 30-minute adjustment period to "lights on" and the environmental conditions of the observation room. All runs were completed within the first 4 hours of "day." Replicate experimental and control trials were intercalated and duplicate batches of trials were run over several days at least twice, weeks to months apart.

Aggression assay

Unless noted, aggression assays were performed similar to those previously described [25] modified from [14]. Briefly, pairs of males were aspirated simultaneously into a 16-mm diameter x 10-mm tall enclosed arena staged in an environmentally-control room held at 25°C and 45% humidity and their behavior was recorded for 20 minutes. To encourage consistent aggressive interactions, the entire floor of the arena was composed of a thin layer of apple juice-sugar agar made as 10g sucrose and 9g agar boiled in 400mL 100% apple juice (Mott's, Plano, TX). To keep the quality of the agar floor consistent, it was used either immediately after a 2-hour setting period or air-dried for 1 hour, wrapped inside of plastic (Saran, Racine, WI), and held at room temperature

until the following day. To impede flies from climbing, the wall of the arena was made

slippery by coating it with Fluon (BioQuip, Rancho Dominguez, CA). Similarly, to limit

flies from hanging from the ceiling, the lid of the arena was brushed with Sigmacote

(Sigma-Aldrich, St. Louis, MO), a transparent silicone paint allowing an unobstructed

view for recording from above. Both coatings were left to dry for at least 24 hours before

running experiments.

Aggression screen

The genetically high- and low- aggressive males used came from a P-element screen

(Mark Eddison, J.C.S, U.H.; unpublished data; see **Figure 3 – Supplement 1A** for the

distribution of median total lunges for all lines observed in screen). From this screen

only lines that maintained a stable phenotype after outcrossing, had normative levels of

activity as estimated by measuring their total distance travelled and total number of

jumps, and appeared otherwise healthy were further used. A line, 5.116, exhibiting

normative levels of aggression and activity was also identified in the screen. This

"standard" was used for all principal experiments reported within this work. **Table S2**

includes measures of the aggressive acts from the various lines used within the

manuscript.

Data acquisition

Intra-male interactions were recorded at 30 frames per second using a Basler A622f

camera (Edmund Optics, Barrington, NJ), with movies from an array of arenas saved

simultaneously with gVision video acquisition software [32] for further analysis.

Experimental arenas were lighted from behind with a flicker-free, uniform, white

backlight (Coherent, Santa Clara, CA), making recordings suitable for machine vision

tracking and behavioral classification methods. For the principal experiments reported in

this work, recordings started within 2 minutes after introduction. For experiments

analyzed from the P-element screen, recordings started 5-7 minutes after introductions,

allowing sufficient time for loading and 5 minutes for flies to settle.

Behavior classification

To catalog the changes in social hierarchy and also the aggressive acts, we used a

combination of manual and automatic classification methods (see **Figure S1** and **Table**

S1 for methods, descriptions, and software used). In summary, the establishment and

reversals of social dominance and also the aggressive acts, fence, box, and tussle,

were manually annotated with VCode [28] and we used CADABRA [13] to automatically

classify wing flick, lunge, and chase. To assign specific lunges to identified males we

used a semi-automated method previously reported [29]. This software application uses

as inputs both original full-length movies and corresponding time-stamped behavioral

events (generated with CADABRA [13] in our case) and iteratively makes short movies

inclusive of each event, allowing users to accurately assess if or which male executed a

particular lunge. For the P-element screen, pairs of males were tracked and

automatically scored for total number of wing flicks, tussles, lunges, chases, and jumps

using CADABRA [13].

Statistical analysis

Data processing, plotting, and statistical analyses were all conducted in Matlab

(MathWorks, Natick, MA). Details for analyses are explained and reported within each

figure legend. See **Figure S2** for specifics regarding data exclusion.

Code availability

Documentation and code used for validating the identification of which male lunged is

available at: https://github.com/JasperCSimon/scoresMovie.git.

Acknowledgments

We thank Erik Hoopfer for gifting us his experimental rig and sharing code to run

CADABRA [13] on the Janelia computer cluster; Mehrab Modi, Lisha Shao, Mark

Eddison, and Kit Longden for insightful suggestions; and past members of Heberlein

laboratory for feedback and support. This work was funded by HHMI.

Authors Contributions

J.C.S. proposed project, designed, conducted, analyzed experiments, and wrote the

manuscript; U.H. supported, advised project, and helped to write manuscript.

Declaration of interests

The authors declare no competing interests.

REFERENCES

1. Berridge, K.C. (2004). Motivation concepts in behavioral neuroscience. Physiol. Behav. *81*, 179–209.

2. Kravitz, E.A., and Huber, R. (2003). Aggression in invertebrates. Curr. Opin. Neurobiol. *13*, 736–743.

3. Freudenberg, F., Carreño Gutierrez, H., Post, A.M., Reif, A., and Norton, W.H.J. (2015). Aggression in non-human vertebrates: Genetic mechanisms and molecular pathways. Am. J. Med. Genet. Part B Neuropsychiatr. Genet. *171*, 603–640.

4. Anderson, D.J. (2016). Circuit modules linking internal states and social behaviour in flies and mice. Nat. Rev. Neurosci. *17*, 692–704.

5. Chen, P., and Hong, W. (2018). Neural circuit mechanisms of social behavior. Neuron *98*, 16–30.

6. Chen, S., Lee, A.Y., Bowens, N.M., Huber, R., and Kravitz, E.A. (2002). Fighting fruit flies: A model system for the study of aggression. PNAS *99*, 5664–5668.

7. Baier, A., Wittek, B., and Brembs, B. (2002). *Drosophila* as a new model organism for the neurobiology of aggression? J. Exp. Biol. *205*, 1233–1240.

8. Sherer, L.M., and Certel, S.J. (2019). The fight to understand fighting: neurogenetic approaches to the study of aggression in insects. Curr. Opin. Insect Sci. *36*, 18–24.

9. Yurkovic, A., Wang, O., Basu, A.C., and Kravitz, E.A. (2006). Learning and memory associated with aggression in *Drosophila melanogaster*. PNAS *103*, 17519–17524.

603   10.   Vrontou, E., Nilsen, S.P., Demir, E., Kravitz, E.A., and Dickson, B.J. (2006).

604   fruitless regulates aggression and dominance in *Drosophila*. Nat. Neurosci. *9*,

605   1469–1471.

606   11.   Kravitz, E.A., and Fernandez, M. de la P. (2015). Aggression in *Drosophila*.

607   Behav. Neurosci. *129*, 549–563.

608   12.   Asahina, K. (2017). Neuromodulation and strategic action choice in *Drosophila*

609   aggression. Annu. Rev. Neurosci. *40*, 51–75.

610   13.   Dankert, H., Wang, L., Hoopfer, E.D., Anderson, D.J., and Perona, P. (2009).

611   Automated monitoring and analysis of social behavior in *Drosophila*. Nat Methods

612   *6*, 297–303.

613   14.   Dierick, H.A. (2007). A method for quantifying aggression in male *Drosophila*

614   *melanogaster*. Nat. Protoc. *2*, 2712–2718.

615   15.   Certel, S.J., and Kravitz, E.A. (2012). Scoring and analyzing aggression in

616   *Drosophila*. Cold Spring Harb. Protoc. *2012*, 319–25.

617   16.   Hoopfer, E.D. (2016). Neural control of aggression in *Drosophila*. Curr. Opin.

618   Neurobiol. *38*, 109–118.

619   17.   Penn, J.K.M., Zito, M.F., and Kravitz, E.A. (2010). A single social defeat reduces

620   aggression in a highly aggressive strain of *Drosophila*. PNAS *107*, 12682–12686.

621   18.   Alekseyenko, O. V., Lee, C., and Kravitz, E.A. (2010). Targeted manipulation of

622   serotonergic neurotransmission affects the escalation of aggression in adult male

623   *Drosophila melanogaster*. PLoS One *5*, e10806.

624   19.   Trannoy, S., Penn, J., Lucey, K., Popovic, D., and Kravitz, E.A. (2016). Short and

625   long-lasting behavioral consequences of agonistic encounters between male

Drosophila melanogaster. PNAS *113*, 4818–4823.

20. Trannoy, S., and Kravitz, E.A. (2017). Strategy changes in subsequent fights as

consequences of winning and losing in fruit fly fights. Fly (Austin). *11*, 129–138.

21. Drews, C. (1993). The concept and definition of dominance in animal behaviour.

Behaviour *125*, 283–313.

22. Francis, R.C. (1988). On the relationship between aggression and social

dominance. Ethology *78*, 223–237.

23. Hoffmann, A.A. (1987). A laboratory study of male territoriality in the sibling

species *Drosophila melanogaster* and *D. simulans.* Anim. Behav. *35*, 807–818.

24. Chouhan, N.S., Mohan, K., and Ghose, A. (2017). cAMP signaling mediates

behavioral flexibility and consolidation of social status in *Drosophila* aggression. J.

Exp. Biol. *220*, 4502–4514.

25. Hoopfer, E.D., Jung, Y., Inagaki, H.K., Rubin, G.M., and Anderson, D.J. (2015).

P1 interneurons promote a persistent internal state that enhances inter-male

aggression in *Drosophila.* Elife *4*, 1–27.

26. Jonsson, T., Kravitz, E.A., and Heinrich, R. (2011). Sound production during

agonistic behavior of male *Drosophila melanogaster*. Fly (Austin). *5*, 29–38.

27. Duistermars, B.J., Pfeiffer, B.D., Hoopfer, E.D., and Anderson, D.J. (2018). A

brain module for scalable control of complex, multi-motor threat displays. Neuron

100, 1474–1490.

28. Haegorn, J., Hailpern, J., and Karahalios, K.G. (2008). VCode and VData:

illustrating a new framework for supporting the video annotation workflow. In

Extended Abstracts of AVI Available at:

http://social.cs.uiuc.edu/projects/vcode.html.

29. Kim, Y.-K., Saver, M., Simon, J., Kent, C.F., Shao, L., Eddison, M., Agrawal, P.,

Texada, M., Truman, J.W., and Heberlein, U. (2018). Repetitive aggressive

encounters generate a long-lasting internal state in *Drosophila melanogaster*

males. PNAS *115*, 1099–1104.

30. Hoffmann, A.A. (1990). The influence of age and experience with conspecifics on

territorial behavior in *Drosophila melanogaster*. J. Insect Behav. *3*, 1–12.

31. Wang, L., Dankert, H., Perona, P., and Anderson, D.J. (2008). A common genetic

target for environmental and heritable influences on aggressiveness in

Drosophila. PNAS *105*, 5657–5663.

32. Gus K. Lott (2010). gVision. Available at: http://gvision-hhmi.sourceforge.net.

Figure 1. Identifying social hierarchy within arenas used for studies of aggression.

Figure 2. Fence, box, and tussle interactions precede the establishment of social hierarchy, whereas the majority of wing flicks, lunges, and chases follow.

Figure 3. Number of lunges executed by dominant individuals is inadequate for explaining hierarchical outcomes.

A

Focal male (F)

Contests dominant (%)

Low (L)   Standard (S)   High (H)   H   H

n=28   n=31   n=23   n=22   n=21

n.s.   n.s.   n.s.

Chance level

n.s.

***

L   S   H   L   Group housed

Opponent (O)

B

Lunges (#/contest)

n.s.

****   *****

n.s.

n.s.

F O   F O   F O'   F O''   F O'''

Dominant male

Figure 3 – Supplement 1. Independently isolated genotypically low, standard, and high aggressive lines identified from a P-element screen; additional genotypically high and low aggressive lines corroborate findings in Figure 3.

A

B

C

Figure 4. Lunging persists as dominant males confront
unfamiliar opponents in unexplored arenas.

Figure 5. Sequence among aggressive acts
in relation to establishing social hierarchy.

Figure S1. Details for including and classifying
fence versus box, tussle interactions.

A

Include

B

Fence

Box,Tussle

Floor

Figure S2. Example data to show the requisite
exclusions for balanced statistical analyses.

Figure S3. Males assert dominance by lunging.

A

Lunge (transiently dominant)
Lunge (dominant at end)

Contests

Unestablished
hierarchy

Terminal reversal

Time (min)

B

Terminal reversal

Lunges (mean+/-std)

Unestablished
hierarchy

n.s.

*****

Male (final dominance status)

Figure S4. Lunges are executed after establishing dominance irrespective of genotype.

Figure S5. Recurring aggressive acts intensifies through subsequent reversals in dominance