Reactivation of latent infections with migration shapes population-level disease dynamics

Daniel J. Becker[1,2*], Ellen D. Ketterson[1,3], Richard J. Hall[2,4,5]

[1]Department of Biology, Indiana University, Bloomington, IN

[2]Center for the Ecology of Infectious Disease, University of Georgia, Athens, GA

[3]Environmental Resilience Institute, Indiana University, Bloomington, IN

[4]Odum School of Ecology, University of Georgia, Athens, GA

[5]Department of Infectious Diseases, College of Veterinary Medicine, University of Georgia,

Athens, GA

[*]danbeck@iu.edu

Running head: Migratory relapse and infectious disease

Keywords: animal migration; mechanistic models; recrudescence; migratory relapse; infectious

disease

Abstract

Annual migration is common across animal taxa and can dramatically shape the spatial and temporal patterns of infectious disease. Although migration can decrease infection prevalence in some contexts, these energetically costly long-distance movements can also have immunosuppressive effects that may interact with transmission processes in complex ways. Here we develop a mechanistic model for the reactivation of latent infections driven by physiological changes associated with migration (i.e., "migratory relapse") and its consequences for infectious disease dynamics. We determine conditions under which migratory relapse can amplify or reduce infection prevalence across host and pathogen traits and transmission phenologies. We show that migratory relapse can dramatically increase prevalence across the annual cycle and may be crucial for maintaining pathogens with low transmissibility and short infectious periods in migratory populations. Conversely, migratory relapse can reduce the prevalence of highly virulent pathogens by amplifying culling of infected hosts during costly migration, especially for highly transmissible pathogens and those transmitted during migration or the breeding season. Our study provides a mechanistic foundation for understanding the spatiotemporal patterns of relapsing infections in migratory hosts, with implications for zoonotic surveillance and understanding how infection patterns will responds to shifts in migratory propensity associated with environmental change. Furthermore, our work suggests that incorporating within-host processes into population-level models of pathogen transmission may be crucial for reconciling the range of migration–infection relationships observed across migratory host species.

Introduction

Long-distance migration is increasingly recognized to shape the spatial and temporal patterns of

infectious disease [1,2]. As these seasonal movements between breeding and wintering grounds

occur across animals [3], migration can facilitate the geographic spread of zoonotic pathogens

such as filoviruses and West Nile virus [4,5]. Pathogens can also threaten migratory hosts, as

observed with sockeye salmon and some waterfowl [6,7]. Accordingly, quantifying the

conditions under which migration enhances or dampens pathogen transmission is important to

identify when and where infection risks in such species are greatest and to predict the

epidemiological consequences of shifting migrations with changing land-use and climate [8,9].

Empirical studies have suggested several ecological mechanisms by which animal

migration could increase or decrease infection risks. Seasonal movement from breeding to

wintering grounds could expose hosts to a greater number of infected individuals, arthropod

vectors, or environmental infectious stages across habitats (i.e., migratory exposure [10–12]).

However, because long-distance movement is energetically demanding, migration could also

reduce infection prevalence by causing high mortality of infected hosts (i.e., migratory culling

[13]). Migration could also allow hosts to escape habitats with high infection risk (i.e., migratory

escape [14]) or temporally separate from groups of particularly infectious conspecifics (i.e.,

migratory allopatry [15]). Additionally, certain habitats experienced during migration could also

directly cull pathogens sensitive to environmental conditions (i.e., migratory recovery [16]).

Despite theoretical support [17,18] and grounding of transmission-reducing mechanisms

in some empirical systems [13,15,19], recent data syntheses suggest animal migration does not

uniformly reduce wildlife disease risks. For example, a meta-analysis across host taxa found

relatively weak negative effects of infection status and intensity on survival during migration

[20], and a comparative study across migrant, resident, and nomadic ungulates also demonstrated

little evidence for migratory escape or culling [21]. Additionally, similar analyses have suggested

mixed support for migratory exposure to explain greater infection risks with long-distance

movement. Although habitat diversity predicts parasite richness across ungulates, migratory or

nomadic species do not sample more diverse habitats than resident counterparts [21]. Further,

whereas bird species with greater habitat diversity likewise have higher helminth richness,

migratory and resident hosts do not systematically differ in habitat use [22]. Such work suggests

greater pathogen exposure of migrants may be insufficient to explain observed infection patterns

and that other factors related to host migration more likely drive increased infection risks.

One underexplored mechanism focuses on how energetic costs of migration could drive

within-host infection processes [23]. Long-distance movement requires substantial energy [24],

with some songbirds and temperate bats investing 25–50% of their body mass in fat reserves for

flight [25]. Such energy demands and physiological changes can negatively affect migrant

immunity [26,27]. Impaired immunity prior to and during migration could make hosts more

susceptible to new infections [28]. However, greater susceptibility would likely increase

pathogen transmission only when exposure occurs at these stages of the annual cycle (e.g.,

breeding grounds before fall migration, wintering grounds in early spring, and stopover sites

[11,29]). As an alternative mechanism, immunosuppression associated with migration could

cause latent infections (e.g., obtained in a previous season) to reactivate. In a rare experimental

test, modified photoperiod of redwings caused the relapse of latent *Borrelia burgdorferi*

infections upon initiation of migratory restlessness [30]. Pathogen reactivation caused by

physiological preparations for long-distance movement (i.e., "migratory relapse") could thus

facilitate migratory hosts arriving at their breeding grounds and their wintering grounds

particularly primed to infect susceptible conspecifics, arthropod vectors, and spillover hosts.

Pathogens that exhibit cycles of latency and reactivation (i.e., susceptible–infected–

latent–infected dynamics) are relatively common, particularly for viruses, bacteria, and some

protozoa [31–33], and infect migratory hosts ranging from birds to ungulates to bats (Table 1).

Immunosuppression from stressors broadly drives the reactivation of pathogens such as

herpesviruses in ungulates [34,35], haemosporidians and flaviviruses in songbirds [36,37], and

henipaviruses in flying foxes [38,39]. Similarly, migration has been implicated in the relapse of

not only *Borrelia burgdorferi* in thrushes but also West Nile virus in white storks [40] and

haemosporidians in rusty blackbirds [41], among other examples (Table 1). Recent comparative

analyses across birds also support migration-induced relapse, as temperate bird species with a

broader range of migratory movements from their wintering grounds (where vector exposure is

unlikely) to their breeding grounds are more capable of infecting susceptible ticks [42]. Such

findings more generally underscore that animal migration could play an underrecognized role in

the spatial and temporal dynamics of relapsing pathogens across diverse mobile host species.

Mechanistic models of relapsing infections in closed populations broadly suggest that

reactivation is an important determinant of pathogen persistence [39], especially when

transmission is seasonal (e.g., many vector-borne diseases [43,44]). For migratory species,

although pathogen transmission often occurs at the breeding grounds [17], exposure

opportunities can take place at other or multiple stages of the annual cycle [11,29,45,46]. If

transmission occurs in only the breeding or wintering grounds, relapse during spring and fall

migration could facilitate pathogen persistence, akin to biannual birth pulses [47], and elevate

prevalence through increasing the force of infection. However, migratory relapse may have

strong impacts only when enough hosts are latently infected, suggesting that pathogen traits

could moderate when and where relapse increases prevalence. Short infectious periods relative to

migration duration could limit how strongly migrants contribute to pathogen transmission upon

arrival at the breeding or wintering grounds. Similarly, relapse of particularly virulent pathogens

could rapidly cull infected hosts and reduce prevalence [1]. Integrating relapse into general

theory for infection dynamics in a migratory host population could accordingly help to

disentangle when animal migration is most likely to magnify infectious disease risks.

We here develop a mechanistic model of relapsing infections in a migratory host to

explore the conditions under which migration-induced pathogen reactivation amplifies or

dampens infection risks. We expand prior modeling frameworks for a host undergoing a two-

way annual migration between the breeding and wintering grounds [17,18]. By assuming the

start of spring and fall migration causes some fraction of latently infected hosts to relapse, we

explore how the frequency of reactivation affects infection dynamics across the annual cycle of a

migratory species. Additionally, we identify pathogen traits for which migratory relapse most

influences disease risk and assess how the phenology of pathogen transmission modifies

infection outcomes. Using models to explore these scenarios could help establish the kinds of

host–pathogen systems for which reactivation driven by animal migration can amplify or reduce

infection prevalence and thus better guide future empirical tests of model predictions [48].

Methods

Model structure

We developed a differential equation model describing population and infection dynamics during

breeding, migration, and overwintering periods. We coarsely parameterized our model around a

migratory songbird, but our model structure remains general enough to apply to a range of

migratory host and pathogen systems in which latent infections reactivate.

<u>Host demography and migration</u>

In the absence of infection, our model tracks the host population size, $N(Y, \tau)$, in year Y and

within-year time τ, where τ takes values between 0 and 1 corresponding to the start and end of

the calendar year, respectively. The population dynamics of the migratory host are described by

ordinary differential equations across each stage of the annual cycle [17]. To avoid offspring

contributing to reproduction in their hatch year, fecundity is proportional to the number of

migrants returning at the start of the breeding season ($N_{breed}(Y) = N(Y, \tau_0)$) [49]. Per-capita

offspring production takes the density-dependent form $b_0 - b_1 N$, where b_0 and b_1 are constants.

The per-capita mortality rate is assumed to be independent of host density but varies across the

annual cycle (μ_j, where $j = b, m, w$ denotes breeding, migration, and wintering, respectively).

The probability of survival at each stage is given by the following equation:

$\sigma_j = e^{-\mu_j T_j}$

Here, T_j ($j = b, m, w$) represents the proportion of the year occupied per stage of the annual

cycle. I the breeding season (i.e., $\tau_0 \leq \tau < \tau_0 + T_b$), population dynamics are as follows:

$\frac{dN}{d\tau} = (b_0 - b_1 N) N_{breed} - \mu_b N$

During spring and fall migrations, hosts travel between breeding and wintering grounds each for

a duration T_m and experience a per-capita mortality rate μ_m. Hosts spend the remainder of the

annual cycle at the wintering grounds (i.e., $T_w = 1 - T_b - 2T_m$) with a per-capita mortality rate

μ_w. During migration and winter, population dynamics are given by the following equations:

$\frac{dN}{d\tau} = -\mu_m N$

and

$\frac{dN}{d\tau} = -\mu_w N$

<u>Seasonal infection dynamics</u>

To represent the dynamics of relapsing infections, we categorize hosts by their current infection

status (i.e., susceptible [S], infected [I], and latent [L]), following the SILI model framework

[33,39]. Transmission from infectious to susceptible hosts in each stage of the annual cycle

occurs at the density-dependent rate β_j (see scenarios below), with infected animals becoming

latent at rate ρ (i.e., $1/\rho$ is the duration of non-lethal acute infections). We model stage-specific

costs of infection as proportional reductions in host fecundity (c_f) and survival (c_j), with each

ranging from 0 (i.e., no change) to 1 (i.e., infected hosts have a 100% reduction in fecundity or

lifespan). During the breeding season, we model infection dynamics as follows:

$\frac{dS}{d\tau} = \left(b_0 - b_1\left(S + I\left(1 - c_f\right) + L\right)\right) N_{breed} - \mu_b S - \beta_b SI$

$\frac{dI}{d\tau} = \beta_b SI - \frac{\mu_b}{1-c_b}I - \rho I$

$\frac{dL}{d\tau} = \rho I - \mu_b L$

Migratory relapse is modeled by the parameter ε, representing the fraction of latent hosts that

return to the infectious class. Because latent infections can reactivate shortly after the initiation

of migratory restlessness [30], we assume relapse occurs instantaneously at the start of both

spring and fall migrations. The starting values for latent and infected individuals at the initiation

of each migration are reset accordingly:

$L = L(1 - \varepsilon)$

and

$I = I + L\varepsilon$

Owing to immunosuppression during migration, we assume acute infections do not subside

during migration (i.e., $\rho = 0$). Infection dynamics are thus given by the following equations:

$\frac{dS}{d\tau} = -\mu_m S - \beta_m SI$

$\frac{dI}{d\tau} = \beta_m SI - \frac{\mu_m}{1-c_m} I$

$\frac{dL}{d\tau} = -\mu_m L$

Infection dynamics are modeled in a similar manner at the overwintering grounds, although

infections can here become latent owing to relatively weaker physiological tradeoffs:

$\frac{dS}{d\tau} = -\mu_w S - \beta_w SI$

$$\frac{dI}{d\tau} = -\frac{\mu_w}{1-c_w}I - \rho I$$

$$\frac{dL}{d\tau} = \rho I - \mu_w L$$

Parameterization

We parameterized our model using the dark-eyed junco (*Junco hyemalis*), a temperate songbird

with diverse migratory strategies throughout North America [50]. Juncos breed across Alaska,

Canada, and the northern United States, migrating south in autumn and wintering throughout the

central and eastern United States. The timing of the junco annual cycle is well-characterized,

with migrants beginning to depart their wintering grounds in early March, migrations spanning

four to eight weeks, and reproduction starting at the breeding grounds in May [51,52]. Migrants

depart their breeding grounds in early October and arrive at the wintering grounds in November.

Juncos lay an average of four eggs per clutch across 1–2 broods, resulting in a maximum per-

capita fecundity of eight juveniles reared over the five-month breeding season [50]. The survival

probability at the breeding grounds is high, whereas that during spring and fall migration is

lower. As the survival probability during winter is slightly higher than during migration [51], we

assume $\sigma_m < \sigma_w < \sigma_b$ to characterize temperate migrants. However, we also assess sensitivity

to this assumption by modeling equal winter and breeding survival probabilities ($\sigma_w = \sigma_b$), as

likely occurs for many Neotropical migrants [53]. Although migratory relapse has not been

observed in juncos, this species can be infected with relapsing pathogens (e.g., *B. burgdorferi*

and haemosporidians) and displays physiological costs of long-distance migration [19,54,55].

Model analysis and infection scenarios

Because our main aim was to assess how migratory relapse affects long-term infection

prevalence (I/N), we varied the proportion of hosts that relapse at the start of migration (ε) from

0 (i.e., no reactivation of infection) to 1 (all latent infections reactivate at the onset of migration).

In our baseline scenario, we assumed pathogen transmission occurs only at the breeding grounds

[17] (i.e., $\beta_b = \beta$, $\beta_m = \beta_w = 0$). This assumption holds for systems where only the breeding

season produces enough susceptible hosts to enable transmission or where exposure is driven by

breeding behavior [56]. Limiting transmission to the breeding grounds can also coarsely

approximate vector-borne diseases given the phenology of many arthropod vectors [31].

To identify pathogen traits that might shape the degree to which migratory relapse affects

infection prevalence, we next systematically varied the duration of acute infection ($1/\rho$) and

transmission rate (β) alongside our relapse parameter (ε). Because past work on juncos supports

migratory culling in this particular host system [19], and because particularly virulent pathogens

could rapidly cull latent hosts that become infected with migratory relapse [1], we also varied the

cost of infection for survival during migration (c_m). We assumed that the costs of infection to

fecundity (c_f) and survival during both breeding (c_b) and wintering (c_w) stages were low.

Lastly, because host aggregations (e.g., at stopover sites [29]) or particular environmental

conditions (e.g., tropical habitats supporting winter activity of vectors [11]) can facilitate some

pathogens being transmitted outside of only the breeding season, we explored model behavior

across three additional transmission phenologies: (*i*) winter only ($\beta_w = \beta$, $\beta_b = \beta_m = 0$), (*ii*)

migration only ($\beta_m = \beta$, $\beta_b = \beta_w = 0$) or (*iii*) year-round ($\beta_b = \beta_m = \beta_w = \beta$).

Across all parameterizations, we ran our model for at least five years and allowed

simulations to continue until the maximum mean difference in prevalence per timestep in the

previous five years was below 0.0001, representing a steady state annual cycle. Most iterations

reached steady state in 40 years. Figure 1 shows our model structure and example population

sizes. All simulations were conducted in R using the *deSolve* package [57]. For each unique set

of model parameters, we recorded the maximum infection prevalence across the steady state

annual cycle. Maximum prevalence is notably important for understanding the epidemiological

consequences of migratory relapse for both pathogen spillover and wildlife conservation [49].

Results

Migratory relapse and host–pathogen dynamics

In the absence of pathogen relapse ($\varepsilon = 0$) and when transmission occurs only in the breeding

season, infection prevalence in the migratory population is relatively low for our baseline

parameterization. Prevalence increases early in the breeding season, attains its maximum prior to

fall migration, and decreases over the winter through lack of transmission (Fig. 2). In contrast,

migratory relapse generally amplifies prevalence across the annual cycle, especially for low-

virulence pathogens (i.e., causing a modest increase in migratory mortality); at its most extreme

($\varepsilon = 1$), reactivation of all latent hosts upon initiation of migration triggers dramatic pulses of

infection back into the host population (Fig. 2A). Prevalence then declines across the breeding

season, as relapsed hosts transition to latency. As more hosts acquire infection, migratory relapse

reduces population size, with the largest reductions occurring for pathogens of intermediate

virulence (Fig. 2A). Intermediate infection costs decrease the mean and variance in prevalence

with migratory relapse across the annual cycle (Fig. 2B). However, when infection poses strong

costs for migrant survival and the fraction of reactivation is high, migratory relapse converts

hosts from latency to actively infected and in turn enhances the effect of migratory culling on

infected individuals (i.e., mortality), thereby reducing population-level prevalence (Fig. 2C).

Sensitivity to pathogen traits and host overwinter survival

To understand broader contexts where migratory relapse amplifies or minimizes infection

prevalence, we systematically covaried the cost of infection to migratory survival (c_m), the

fraction of relapsing hosts upon migration (ε), pathogen transmissibility (β), and the duration of

non-lethal acute infection ($1/\rho$). We explored patterns for two host migration scenarios: that of a

temperate migrant with winter survival lower than breeding survival ($\sigma_w < \sigma_b$) and that of a

Neotropical migrant where winter survival is equivalent to breeding survival ($\sigma_w = \sigma_b$).

For the baseline (i.e., temperate migrant), migratory relapse broadly increases the

maximum prevalence when costs of infection are low (Fig. 3). For pathogens with low

transmissibility ($\beta = 0.05$) and short acute infections (i.e., one month relative to the six-week

migration), migratory relapse allows pathogens to persist that otherwise would be unable to

invade the host population (Fig. 3). As expected, pathogens with higher transmissibility and

longer acute infections attain greater peak prevalence when costs of infection are low to

intermediate. However, migratory relapse can decrease prevalence relative to models assuming

no relapse ($\varepsilon = 0$) when costs of infection and transmission rates are both high (Fig. 3).

These patterns were generally consistent when winter survival reflected Neotropical

migrant systems ($\sigma_w = \sigma_b$). Maximum prevalence was overall greater, with the pathogen

typically able to persist without relapse (Fig. S1). With low costs of infection for migrant

survival, migratory relapse produced greater increases in prevalence, particularly with higher

transmission rates and longer infectious periods. At higher infection costs, the difference in

maximum prevalence between models without relapse ($\varepsilon = 0$) and with complete relapse ($\varepsilon = 1$) was minimized so that migratory relapse only dampened infection by virulent pathogens.

Sensitivity to transmission phenology

We lastly considered how the timing of pathogen transmission within the migratory host annual

cycle affects model outcomes. The amplifying effects of migratory relapse on prevalence are

most pronounced when transmission rates are low, irrespective of transmission phenology (Fig.

4). Yet when transmission rates are high, complete migratory relapse ($\varepsilon = 1$) can reduce

prevalence relative to a model without relapse ($\varepsilon = 0$) especially when transmission occurs

during migration and throughout the annual cycle. For the former, this infection-dampening

effect of relapse is particularly pronounced for pathogens with short infectious periods (Fig. S2).

These patterns were similar for the Neotropical migrant scenario (i.e. high winter survival),

although fewer regions of parameter space reduced infection prevalence with relapse (Fig. S3).

Discussion

Determining the conditions under which migration amplifies or dampens pathogen transmission

is important to identify when and where infection risks are greatest in highly mobile species.

Migratory species host various pathogens with cycles of latency and reactivation, including

several with zoonotic potential [30,39,40], but theory to date on migratory host–pathogen

interactions does not account for migration-induced pathogen reactivation and its population-

level consequences. Using a mathematical model, we found that migratory relapse can increase

or decrease infection prevalence throughout the annual cycle, dependent upon pathogen traits

and transmission phenology. For relatively low-virulence pathogens, gains to the infectious class

through relapse outpace losses of infected individuals through migratory culling to increase

prevalence. In contrast, migratory relapse can exacerbate migratory culling to reduce prevalence

for pathogens that are especially virulent, highly transmissible, and can be spread during the

breeding and migratory stages of the annual cycle. Past theory on host–pathogen dynamics

generally concludes that migration reduces infection prevalence by reducing transmission

opportunity (i.e., migratory escape) and infected host survival (i.e., migratory culling) [16–18].

Our work instead suggests that incorporating physiological processes into host–pathogen models

(i.e., energetic demands of migration that lead to immunosuppression and relapse [24,26,27])

could account for the wider range of migration–infection relationships observed in wildlife.

For an annual cycle typical of migratory songbirds, and a range of biologically plausible

pathogen traits, our model generated dramatic increases in infection prevalence with an increase

in the proportion of hosts experiencing migratory relapse. In some cases, relapse was necessary

to maintain persistence of less-transmissible pathogens with shorter acute infections across the

annual cycle. Although prior theory has not examined relapse induced by physiological changes

with migration, past models of relapsing pathogens have demonstrated that reactivation can

facilitate pathogen persistence. In the absence of immigration, recurrent cycles of acute infection

and latency were necessary to explain henipavirus dynamics in straw-colored fruit bats [39], and

more frequent relapse optimized pathogen invasion potential for human malaria (*Plasmodium*

vivax) [44]. Our model suggests migratory relapse can generate pronounced biannual peaks in

prevalence that maintain more infection throughout the annual cycle than by direct transmission

alone. This pattern differs in its seasonal timing and magnitude from those generated by seasonal

birth pulses or relapse induced by immune tradeoffs with reproduction [36,38,58,59], especially

in systems where offspring benefit from prolonged maternal immunity [60]. Because this pattern

was most evident when infection had low costs for migrant survival, our model predictions may

be especially applicable to migratory reservoir hosts that themselves do not experience high

morbidity or mortality (e.g., henipaviruses in flying foxes and *Borrelia burgdorferi* in songbirds

[30,39]). Given the high prevalence of actively infectious hosts during migration, migratory

stopover sites and arrival periods of migrants returning to breeding or wintering grounds could

be prioritized as important targets for surveillance and interventions to limit spillover risk [61].

Our model also identified contexts in which migratory relapse can decrease prevalence,

or even cause pathogen extinction, by removing more actively infectious hosts with migratory

culling. Reductions in prevalence were most likely to be seen in highly transmissible pathogens,

those with long infectious periods, and when transmission occurred at the breeding grounds or

during migration. Systems for which such conditions are met include rhabdoviruses in salmonid

fish as well as haemosporidian parasites and flaviviruses in some songbirds [7,19,62,63]. High

relapse frequency and high mortality of infected migrants might be more common for species

undertaking strenuous migrations, such as those involving prolonged periods of powered flight.

This supports prior predictions that migratory culling is more likely to operate in long-distance

migrations with fewer stopovers than shorter migrations or where migratory species move and

forage in a stable environmental window (e.g., ungulates "surfing the green wave") [17,64].

Given the sensitivity of our model to the degree of relapse and the costs of infection for

migrant survival, estimation of these parameters will be important for applying this framework to

empirical systems. Detecting pathogen relapse requires sampling hosts across their annual cycle

[33], which can be logistically challenging for highly mobile species [65]. However,

advancements in tracking technologies (e.g., lightweight geolocators, stable isotopes) are

improving inference into migratory networks, which could facilitate temporal sampling across

breeding, wintering, and stopover habitats [66]. Additionally, multiple diagnostic efforts are

necessary to differentiate compartments within the SILI framework. For example, serology,

PCR, and microscopy can distinguish uninfected (i.e., susceptible), acutely infected, and latent

hosts for haemosporidian parasites [67], and pathogen genotyping could disentangle whether

infections are more likely to indicate reactivation rather than novel transmission events [33].

Integrating longitudinal sampling and these diagnostics with mark–recapture could help to

estimate infection- and stage-specific survival probabilities and infer pathogen costs [68].

To examine conditions under which relapse during migration increases or decreases

infection risks, we focused our model on a simple system with a single interaction between a

migratory host and its pathogen. For tractability, we ignored alternative ecological interactions

such as multiple host species (and thus a more generalist pathogen) or explicit arthropod vectors.

However, high prevalence in relapsing migrants suggests migrants could introduce pulses of

infection into seasonally sympatric co-occurring species. This could be particularly important for

pathogen maintenance in partially migratory species such as juncos, where long-distance

migrants form mixed flocks with residents at shared wintering grounds [50]. For vector-borne

pathogens with cycles of latency and reactivation, such as *Borrelia burgdorferi* and some

flaviviruses [30,69], migrants undergoing relapse could also have disproportionate influence on

infecting vectors upon their arrival at the breeding grounds. Future extensions to our model could

assess how phenological overlap between migrants and vectors shapes disease outcomes [49,70].

Because many species are shifting the timing and extent of their migratory movements in

response to climate and anthropogenic change [9,71], associated changes to host physiology and

survival could alter how migratory relapse influences infection patterns. Deterioration of

resources at breeding or wintering sites could increase the proportion of relapsing hosts,

increasing prevalence of low-virulence pathogens or preventing persistence of high-virulence

pathogens if migratory mortality is also elevated. Additionally, many migratory species are

increasingly overwintering in anthropogenic habitats. For example, Anna's hummingbirds and

European blackcaps both spend more of their annual cycle in urbanized regions owing to

abundant and predictable anthropogenic food resources [72,73]. Such resources could dampen or

amplify the likelihood of pathogen relapse at spring departure depending on their nutritional

quality or effects on host density and crowding [74]. Other stressors associated with urban

habitats, such as artificial light at night, could further amplify the likelihood of relapse [75,76].

These consequences of urban habituation could be especially relevant for human health in the

context of wildlife reservoirs of relapsing zoonoses, such as flying foxes and henipaviruses [77].

Many pathogens of concern to human, domestic animal, and wildlife health exhibit cycles

of latency and reactivation. Our model provides a generalizable and mechanistic framework for

understanding the dynamics of such infections in migratory hosts. We demonstrate theoretical

support for migratory relapse increasing infectious disease risks but also emphasize the context

dependence of these patterns on host and pathogen traits. Empirical estimates of relapse and

survival across the annual cycle will be important for linking such frameworks with empirical

systems, which will be critical for predicting how anthropogenic changes in both climate and

land use alter animal migration, reactivation of latent infections, and pathogen spillover risks.

Data accessibility

R code for reproducing the main analyses is available in the Dryad Digital Repository:

https://datadryad.org/stash/share/eT4DN6oJ0BlGmv5fqjg16cm3HThgXmvgllWemPpInYY.

Author contributions

DJB and RJH designed the model, DJB and EDK parameterized the model, and DJB performed

the analyses and wrote the first draft of the manuscript. All authors contributed to revisions.

Acknowledgements

We thank members of the Ketterson lab for constructive feedback on this manuscript.

Funding

DJB was supported by an appointment to the Intelligence Community Postdoctoral Research

Fellowship Program, administered by Oak Ridge Institute for Science and Education through an

interagency agreement between the U.S. Department of Energy and the Office of the Director of

National Intelligence. EDK was supported by the Environmental Resilience Institute, funded by

Indiana University's Prepared for Environmental Change Grand Challenge Initiative. RJH was

supported by the National Science Foundation (DEB-1754392 and DEB-1911925).

Competing interests

We declare no competing interests.

Tables

Table 1. Examples of pathogens that exhibit cycles of latency and reactivation, their detection in migratory or nomadic host species, and likely drivers of observed relapse events.

Pathogen	Host taxon	Behavior	Likely driver of relapse	Reference
Borrelia burgdorferi	Redwing (*Turdus iliacus*)	Migrant	Migration	[30]
Eastern equine encephalitis virus	Passeriformes	Migrant	Migration, reproduction	[69]
Plasmodium spp.	Passeriformes	Migrant	Migration, reproduction	[31]
West Nile virus	White storks (*Ciconia ciconia*)	Migrant	Migration	[40]
Haemosporida	Rusty blackbird (*Euphagus carolin*)	Migrant	Migration	[41]
Infectious hematopoietic necrosis virus	Sockeye salmon (*Oncorhynchus nerka*)	Migrant	Migration	[63]
Brucella abortus	Elk (*Cervus canadensis*)	Migrant	Reproduction	[58]
Equine herpesviruses	Grévy's zebras (*Equus grevyi*)	Migrant	Translocation	[35]
Henipaviruses	Flying foxes (*Pteropus* spp.)	Nomad	Reproduction, nutritional stress	[38]

419   Table 2. Model parameters, their definitions, and default values. All rates are given in units of

420   years^{-1}. Parameterization for host demography and migration is based on juncos [50–52].

Parameter	Definition	Value
Annual cycle		
T_b	Proportion of year spent at breeding grounds	0.42 (~5 months)
T_m	Proportion of year spent in each migration	0.12 (1.5 months)
T_w	Proportion of year spent at wintering grounds	0.34 (~4 months)
τ_0	Start of the breeding season per annual cycle	0.33 (~early May)
Host demography		
b_0	Density-independent reproduction rate	3.3
b_1	Density-dependent reproduction rate	0.003
σ_b	Breeding survival probability	0.95
σ_m	Migration survival probability (one-way)	0.79
σ_w	Winter survival probability	0.85 (temperate), 0.95 (Neotropical)
Pathogen traits		
β	Density-dependent transmission rate	0.05 (low), 0.5 (high)
ρ	Infected-to-latent rate	4 (1 month), 12 (3 months)
ε	Fraction of relapsing hosts at the start of migration	0–1
c_f	Cost of infection for fecundity	0.2
c_b	Cost of infection for breeding survival	0.2
c_m	Cost of infection for migratory survival	0.2–0.8
c_w	Cost of infection for winter survival	0.2

Figures and legends

Figure 1. (A) Model schematic of relapsing infections in a migratory host through the annual

cycle, for a pathogen transmitted at the host breeding grounds. Filled circles depict the number

susceptible (S), infectious (I) and latently infected (L) hosts, and arrows represent gain or losses

to each class through demographic or infection processes. Arrows between panels illustrate that a

fraction (ε) of latently infected hosts relapse at the start of migration. (B) Example dynamics

over the steady state annual cycle. Solid lines depict numbers of susceptible (S, light gray),

infectious (I, pink), and latent (L, dark gray) hosts, and vertical dashed lines delineate the spring

and fall migratory periods. Infection parameters used are the transmission rate ($\beta_b = 0.05$), cost

of infection for migrant survival ($c_m = 0.5$), rate of transition from infectious to latent ($\rho = 4$,

corresponding to a three-month duration of acute infection), and the probability of relapse at the

onset of migration ($\varepsilon = 0.75$); demographic parameters are the default values listed in Table 2.

Figure 2. Steady state population size (top row) and infection prevalence (bottom row) across the

annual cycle as a function of the fraction of relapse at migration onset (ε, colored lines) and low,

medium, and high infection costs for migratory survival (c_m, columns). Transmission occurs at

the breeding grounds ($\beta_b = 0.5$), infection is acute for three months ($\rho = 4$), and winter survival

represents a temperate migrant ($\sigma_w = 0.85$). All other parameter values are listed in Table 2.

Figure 3. Sensitivity of the relationship between peak infection prevalence and virulence (i.e.,

cost of infection for migratory survival, c_m) to pathogen traits. Transmission occurs only at the

breeding grounds for moderate ($\beta_b = 0.05$, top) and high ($\beta_b = 0.5$, bottom) transmissibility.

Lines show the fraction of relapsing hosts at the start of migration (ε), and columns indicate an

increasing duration of acute infection ($1/\rho$) from 1 to 3 months. Winter survival represents a

temperate migrant ($\sigma_w = 0.85$), and all other parameter values are listed in Table 2.

Figure 4. Effects of transmission phenology and pathogen traits on how infection prevalence

responds to migratory relapse. We calculate the response of infection prevalence to relapse

(vertical axis) as the difference in maximum prevalence between models where all individuals

relapse ($\varepsilon = 1$) and no individuals relapse ($\varepsilon = 0$); values above or below the dashed line

indicate that relapse increases or decreases peak prevalence, respectively. Columns represent

different transmission phenologies (breeding only, migration only, winter only, or year-round),

Rows represent moderate ($\beta = 0.05$, top) and high ($\beta = 0.5$, bottom) transmissibility, line width

represent short versus long infectious periods ($1/\rho$), and the horizontal axis represents costs of

infection for migrant survival (c_m). Winter survival is representative of temperate migrants

($\sigma_w = 0.85$), and all other parameter values are provided in Table 2.

References

1. Altizer S, Bartel R, Han BA. 2011 Animal migration and infectious disease risk. *science* **331**, 296–302.

2. Fritzsche McKay A, Hoye BJ. 2016 Are Migratory Animals Superspreaders of Infection? *Integr. Comp. Biol.* **56**, 260–267. (doi:10.1093/icb/icw054)

3. Dingle H. 2014 *Migration: the biology of life on the move*. Oxford University Press, USA.

4. Owen J, Moore F, Panella N, Edwards E, Bru R, Hughes M, Komar N. 2006 Migrating Birds as Dispersal Vehicles for West Nile Virus. *EcoHealth* **3**, 79. (doi:10.1007/s10393-006-0025-9)

5. Ogawa H *et al.* 2015 Seroepidemiological Prevalence of Multiple Species of Filoviruses in Fruit Bats (Eidolon helvum) Migrating in Africa. *J. Infect. Dis.* **212**, S101–S108. (doi:10.1093/infdis/jiv063)

6. Kirby JS, Stattersfield AJ, Butchart SH, Evans MI, Grimmett RF, Jones VR, O'Sullivan J, Tucker GM, Newton I. 2008 Key conservation issues for migratory land-and waterbird species on the world's major flyways. *Bird Conserv. Int.* **18**, S49–S73.

7. Jeffries KM *et al.* 2014 Immune response genes and pathogen presence predict migration survival in wild salmon smolts. *Mol. Ecol.* **23**, 5803–5815.

8. Wilcove DS, Wikelski M. 2008 Going, going, gone: is animal migration disappearing. *PLoS Biol.* **6**, e188.

9. Satterfield DA, Marra PP, Sillett TS, Altizer SM. 2018 Responses of migratory species and their pathogens to supplemental feeding. *Philos. Trans. R. Soc. B Biol. Sci.*

10. Figuerola J, Green AJ. 2000 Haematozoan parasites and migratory behaviour in waterfowl. *Evol. Ecol.* **14**, 143–153.

11. Waldenström J, Bensch S, Kiboi S, Hasselquist D, Ottosson U. 2002 Cross-species infection of blood parasites between resident and migratory songbirds in Africa. *Mol. Ecol.* **11**, 1545–1554.

12. Morgan ER, Medley GF, Torgerson PR, Shaikenov BS, Milner-Gulland EJ. 2007 Parasite transmission in a migratory multiple host system. *Ecol. Model.* **200**, 511–520.

13. Bradley CA, Altizer S. 2005 Parasites hinder monarch butterfly flight: implications for disease spread in migratory hosts. *Ecol. Lett.* **8**, 290–300.

14. Folstad I, Nilssen AC, Halvorsen O, Andersen J. 1991 Parasite avoidance: the cause of post-calving migrations in Rangifer? *Can. J. Zool.* **69**, 2423–2429.

15. Krkošek M, Gottesfeld A, Proctor B, Rolston D, Carr-Harris C, Lewis MA. 2007 Effects of
host migration, diversity and aquaculture on sea lice threats to Pacific salmon populations.
Proc. R. Soc. Lond. B Biol. Sci. **274**, 3141–3149.

16. Shaw AK, Binning SA. 2016 Migratory recovery from infection as a selective pressure for
the evolution of migration. *Am. Nat.* **187**, 491–501.

17. Hall RJ, Altizer S, Bartel RA. 2014 Greater migratory propensity in hosts lowers pathogen
transmission and impacts. *J. Anim. Ecol.* **83**, 1068–1077. (doi:10.1111/1365-2656.12204)

18. Johns S, Shaw AK. 2015 Theoretical insight into three disease-related benefits of migration.
Popul. Ecol. **58**, 213–221. (doi:10.1007/s10144-015-0518-x)

19. Slowinski SP, Fudickar AM, Hughes AM, Mettler RD, Gorbatenko OV, Spellman GM,
Ketterson ED, Atwell JW. 2018 Sedentary songbirds maintain higher prevalence of
haemosporidian parasite infections than migratory conspecifics during seasonal sympatry.
PLOS ONE **13**, e0201563. (doi:10.1371/journal.pone.0201563)

20. Risely A, Klaassen M, Hoye BJ. 2018 Migratory animals feel the cost of getting sick: A
meta-analysis across species. *J. Anim. Ecol.* **87**, 301–314.

21. Teitelbaum CS, Huang S, Hall RJ, Altizer S. 2018 Migratory behaviour predicts greater
parasite diversity in ungulates. *Proc R Soc B* **285**, 20180089. (doi:10.1098/rspb.2018.0089)

22. Leung TL, Koprivnikar J. 2016 Nematode parasite diversity in birds: the role of host
ecology, life history and migration. *J. Anim. Ecol.* **85**, 1471–1480.

23. Daversa DR, Fenton A, Dell AI, Garner TWJ, Manica A. 2017 Infections on the move: how
transient phases of host movement influence disease spread. *Proc R Soc B* **284**, 20171807.
(doi:10.1098/rspb.2017.1807)

24. Wikelski M, Tarlow EM, Raim A, Diehl RH, Larkin RP, Visser GH. 2003 Costs of
migration in free-flying songbirds. *Nature* **423**, 704. (doi:10.1038/423704a)

25. Guglielmo CG. 2018 Obese super athletes: fat-fueled migration in birds and bats. *J. Exp.*
Biol. **221**, jeb165753. (doi:10.1242/jeb.165753)

26. Owen JC, Moore FR. 2008 Swainson's thrushes in migratory disposition exhibit reduced
immune function. *J. Ethol.* **26**, 383–388. (doi:10.1007/s10164-008-0092-1)

27. Eikenaar C, Hegemann A. 2016 Migratory common blackbirds have lower innate immune
function during autumn migration than resident conspecifics. *Biol. Lett.* **12**, 20160078.
(doi:10.1098/rsbl.2016.0078)

28. Van Dijk JG, Hoye BJ, Verhagen JH, Nolet BA, Fouchier RA, Klaassen M. 2014 Juveniles
and migrants as drivers for seasonal epizootics of avian influenza virus. *J. Anim. Ecol.* **83**,
266–275.

29. Krauss S, Stallknecht DE, Negovetich NJ, Niles LJ, Webby RJ, Webster RG. 2010
Coincident ruddy turnstone migration and horseshoe crab spawning creates an ecological
'hot spot' for influenza viruses. *Proc. R. Soc. B Biol. Sci.* **277**, 3373–3379.
(doi:10.1098/rspb.2010.1090)

30. Gylfe A, Bergström S, Lundström J, Olsen B. 2000 Reactivation of Borrelia infection in
birds. *Nature* **403**, 724–725. (doi:10.1038/35001663)

31. Beaudoin RL, Applegate JE, Davis DE, McLean RG. 1971 A model for the ecology of avian
malaria. *J. Wildl. Dis.* **7**, 5.

32. Virgin HW, Wherry EJ, Ahmed R. 2009 Redefining chronic viral infection. *Cell* **138**, 30–50.
(doi:10.1016/j.cell.2009.06.036)

33. Plowright RK, Peel AJ, Streicker DG, Gilbert AT, McCallum H, Wood J, Baker ML, Restif
O. 2016 Transmission or Within-Host Dynamics Driving Pulses of Zoonotic Viruses in
Reservoir–Host Populations. *PLoS Negl. Trop. Dis.* **10**, e0004796.
(doi:10.1371/journal.pntd.0004796)

34. Lamontagne L, Sadi L, Joyal R. 1989 Serological evidence of bovine herpesvirus 1-related
virus infection in the white-tailed deer population on Anticosti Island, Quebec. *J. Wildl. Dis.*
25, 202–205.

35. Seeber PA, Quintard B, Sicks F, Dehnhard M, Greenwood AD, Franz M. 2018
Environmental stressors may cause equine herpesvirus reactivation in captive Grévy's zebras
(Equus grevyi). *PeerJ* **6**, e5422. (doi:10.7717/peerj.5422)

36. Applegate JE. 1970 Population Changes in Latent Avian Malaria Infections Associated with
Season and Corticosterone Treatment. *J. Parasitol.* **56**, 439–443. (doi:10.2307/3277599)

37. Reisen WK, Chiles RE, Green EN, Fang Y, Mahmood F, Martinez VM, Laver T. 2003
Effects of immunosuppression on encephalitis virus infection in the house finch, Carpodacus
mexicanus. *J. Med. Entomol.* **40**, 206–214.

38. Plowright RK, Field HE, Smith C, Divljan A, Palmer C, Tabor G, Daszak P, Foley JE. 2008
Reproduction and nutritional stress are risk factors for Hendra virus infection in little red
flying foxes (Pteropus scapulatus). *Proc. R. Soc. B* **275**, 861–869.
(doi:10.1098/rspb.2007.1260)

39. Glennon EE *et al.* 2019 What is stirring in the reservoir? Modelling mechanisms of
henipavirus circulation in fruit bat hosts. *Philos. Trans. R. Soc. B Biol. Sci.* **374**, 20190021.
(doi:10.1098/rstb.2019.0021)

40. Malkinson M, Banet C, Weisman Y, Pokamunski S, King R, Deubel V. 2002 Introduction of
West Nile Virus in the Middle East by Migrating White Storks. *Emerg. Infect. Dis.* **8**, 392–
397. (doi:10.3201/eid0804.010217)

41. Barnard WH, Mettke-Hofmann C, Matsuoka SM. 2010 Prevalence of hematozoa infections among breeding and wintering rusty blackbirds. *The Condor* **112**, 849–853.

42. Becker DJ, Han BA. 2020 The macroecology and evolution of avian competence for Borrelia burgdorferi. *bioRxiv* , 2020.04.15.040352. (doi:10.1101/2020.04.15.040352)

43. Murdock CC, Foufopoulos J, Simon CP. 2013 A Transmission Model for the Ecology of an Avian Blood Parasite in a Temperate Ecosystem. *PLOS ONE* **8**, e76126. (doi:10.1371/journal.pone.0076126)

44. White MT, Shirreff G, Karl S, Ghani AC, Mueller I. 2016 Variation in relapse frequency and the transmission potential of Plasmodium vivax malaria. *Proc. R. Soc. B Biol. Sci.* **283**, 20160048.

45. Hasselquist D, Östman Ö, Waldenström J, Bensch S. 2007 Temporal patterns of occurrence and transmission of the blood parasite Haemoproteus payevskyi in the great reed warbler Acrocephalus arundinaceus. *J. Ornithol.* **148**, 401–409. (doi:10.1007/s10336-007-0144-2)

46. Miller-Butterworth CM, Vonhof MJ, Rosenstern J, Turner GG, Russell AL. 2014 Genetic Structure of Little Brown Bats (Myotis lucifugus) Corresponds with Spread of White-Nose Syndrome among Hibernacula. *J. Hered.* **105**, 354–364. (doi:10.1093/jhered/esu012)

47. Hayman DTS. 2015 Biannual birth pulses allow filoviruses to persist in bat populations. *Proc. R. Soc. Lond. B Biol. Sci.* **282**, 20142591. (doi:10.1098/rspb.2014.2591)

48. Restif O *et al.* 2012 Model-guided fieldwork: practical guidelines for multidisciplinary research on wildlife ecological and epidemiological dynamics. *Ecol. Lett.* **15**, 1083–1094. (doi:10.1111/j.1461-0248.2012.01836.x)

49. Brown LM, Hall RJ. 2018 Consequences of resource supplementation for disease risk in a partially migratory population. *Phil Trans R Soc B* **373**, 20170095. (doi:10.1098/rstb.2017.0095)

50. Nolan V, Ketterson ED, Cristol DA, Rogers CM, Clotfelter ED, Titus RC, Schoech SJ, Snajdr E. 2002 *Dark-eyed junco (Junco hyemalis).* American Ornithologists' Union.

51. Ketterson ED, Nolan Jr V. 1982 The role of migration and winter mortality in the life history of a temperate-zone migrant, the Dark-eyed Junco, as determined from demographic analyses of winter populations. *The Auk* **99**, 243–259.

52. Ketterson ED, Nolan Jr V. 1985 Intraspecific variation in avian migration: evolutionary and regulatory aspects. *Migr. Mech. Adapt. Significance* **27**, 553–579.

53. Sillett TS, Holmes RT. 2002 Variation in survivorship of a migratory songbird throughout its annual cycle. *J. Anim. Ecol.* **71**, 296–308. (doi:10.1046/j.1365-2656.2002.00599.x)

54. Wright SA, Tucker JR, Donohue AM, Castro MB, Kelley KL, Novak MG, Macedo PA. 2011 Avian Hosts of Ixodes pacificus (Acari: Ixodidae) and the Detection of Borrelia

burgdorferi in Larvae Feeding on the Oregon Junco. *J. Med. Entomol.* **48**, 852–859. (doi:10.1603/ME11001)

55. Becker DJ, Talbott KM, Smiley TM, Clark KL, Sauer PE, Ketterson ED. 2019 Leukocyte profiles vary with breeding latitude and infection status in a seasonally sympatric songbird. *Anim. Migr.* **6**, 28–40. (doi:10.1515/ami-2019-0004)

56. Altizer S, Dobson A, Hosseini P, Hudson P, Pascual M, Rohani P. 2006 Seasonality and the dynamics of infectious diseases. *Ecol. Lett.* **9**, 467–484.

57. Soetaert KER, Petzoldt T, Setzer RW. 2010 Solving differential equations in R: package deSolve. *J. Stat. Softw.* **33**.

58. Thorne ET, Morton JK. 1978 Brucellosis in elk. II. Clinical effects and means of transmission as determined through artificial infections. *J. Wildl. Dis.* **14**, 280.

59. Peel AJ, Pulliam JRC, Luis AD, Plowright RK, O'Shea TJ, Hayman DTS, Wood JLN, Webb CT, Restif O. 2014 The effect of seasonal birth pulses on pathogen persistence in wild mammal populations. *Proc. R. Soc. Lond. B Biol. Sci.* **281**, 20132962. (doi:10.1098/rspb.2013.2962)

60. Peel AJ, Baker KS, Hayman DTS, Broder CC, Cunningham AA, Fooks AR, Garnier R, Wood JLN, Restif O. 2018 Support for viral persistence in bats from age-specific serology and models of maternal immunity. *Sci. Rep.* **8**, 3859. (doi:10.1038/s41598-018-22236-6)

61. Plowright RK, Becker DJ, McCallum H, Manlove KR. 2019 Sampling to elucidate the dynamics of infections in reservoir hosts. *Philos. Trans. R. Soc. B* (doi:10.1098/rstb.2018.0336)

62. George TL, Harrigan RJ, LaManna JA, DeSante DF, Saracco JF, Smith TB. 2015 Persistent impacts of West Nile virus on North American bird populations. *Proc. Natl. Acad. Sci.* **112**, 14290–14294. (doi:10.1073/pnas.1507747112)

63. Mulcahy D, Jenes CK, Pascho R. 1984 Appearance and quantification of infectious hematopoietic necrosis virus in female sockeye salmon (Oncorhynchus nerka) during their spawning migration. *Arch. Virol.* **80**, 171–181.

64. Aikens EO, Kauffman MJ, Merkle JA, Dwinnell SP, Fralick GL, Monteith KL. 2017 The greenscape shapes surfing of resource waves in a large migratory herbivore. *Ecol. Lett.* **20**, 741–750.

65. Becker DJ, Crowley DE, Washburne AD, Plowright RK. 2019 Temporal and spatial limitations in global surveillance for bat filoviruses and henipaviruses. *Biol. Lett.* **15**, 20190423. (doi:10.1098/rsbl.2019.0423)

66. Fudickar AM, Ketterson ED. 2018 Genomes to space stations: the need for the integrative study of migration for avian conservation. *Biol. Lett.* **14**, 20170741.

67. Jarvi SI, Schultz JJ, Atkinson CT. 2002 PCR diagnostics underestimate the prevalence of
avian malaria (Plasmodium relictum) in experimentally-infected passerines. *J. Parasitol.* **88**,
153–158.

68. Rushing CS. 2019 Estimability of migration survival rates from integrated breeding and
winter capture–recapture data. *Ecol. Evol.* **9**, 849–858. (doi:10.1002/ece3.4826)

69. Crans WJ, Caccamise DF, McNelly JR. 1994 Eastern Equine Eneephalomyelitis Virus in
Relation to the Avian Community of a Coastal Cedar Swamp. *J. Med. Entomol.* **31**, 711–728.
(doi:10.1093/jmedent/31.5.711)

70. Hall RJ, Brown LM, Altizer S. 2016 Modeling vector-borne disease risk in migratory
animals under climate change. *Integr. Comp. Biol.* **56**, 353–364. (doi:10.1093/icb/icw049)

71. Howard C, Stephens PA, Tobias JA, Sheard C, Butchart SHM, Willis SG. 2018 Flight range,
fuel load and the impact of climate change on the journeys of migrant birds. *Proc. R. Soc. B
Biol. Sci.* **285**, 20172329. (doi:10.1098/rspb.2017.2329)

72. Plummer KE, Siriwardena GM, Conway GJ, Risely K, Toms MP. 2015 Is supplementary
feeding in gardens a driver of evolutionary change in a migratory bird species? *Glob.
Change Biol.* **21**, 4353–4363. (doi:10.1111/gcb.13070)

73. Greig EI, Wood EM, Bonter DN. 2017 Winter range expansion of a hummingbird is
associated with urbanization and supplementary feeding. *Proc R Soc B* **284**, 20170256.
(doi:10.1098/rspb.2017.0256)

74. Becker DJ, Streicker DG, Altizer S. 2015 Linking anthropogenic resources to wildlife–
pathogen dynamics: a review and meta-analysis. *Ecol. Lett.* **18**, 483–495.
(doi:10.1111/ele.12428)

75. Kernbach ME *et al.* 2019 Light pollution increases West Nile virus competence of a
ubiquitous passerine reservoir species. *Proc. R. Soc. B* **286**, 20191051.

76. Ouyang JQ, de Jong M, van Grunsven RH, Matson KD, Haussmann MF, Meerlo P, Visser
ME, Spoelstra K. 2017 Restless roosts: Light pollution affects behavior, sleep, and
physiology in a free-living songbird. *Glob. Change Biol.* **23**, 4987–4994.

77. Kessler MK *et al.* 2018 Changing resource landscapes and spillover of henipaviruses. *Ann.
N. Y. Acad. Sci.* **1429**, 78–99. (doi:10.1111/nyas.13910)