The role of conidia in the dispersal of *Ascochyta rabiei*

Ihsanul Khaliq[1,*], Joshua Fanning[2], Paul Melloy[1], Jean Galloway[3], Kevin Moore[4], Daniel Burrell[1], Adam H Sparks[1]

[1]University of Southern Queensland, Centre for Crop Health, Toowoomba, Queensland 4350, Australia

[2]Agriculture Victoria, Horsham, Victoria, 3401, Australia

[3]Department of Primary Industries and Regional Development (DPIRD), Northam, WA, 6401, Australia

[4]New South Wales Department of Primary Industries, 4 Marsden Park Rd, Tamworth, NSW, 2340, Australia

* **Corresponding author**:
E-mail address: Ihsan.Khaliq@usq.edu.au
Phone: +61 7 4631 1877

Abstract

Ascochyta rabiei asexual spores (conidia) were assumed to spread over short distances

(~10 m) in a combination of rain and strong wind. The potential distance of conidial spread

was investigated in three rainfall and three sprinkler irrigation events. Chickpea trap plants

were distributed at the distances of 0, 10, 25, 50 and 75 m from infected chickpea plots

before scheduled irrigation and forecast rainfall events. Trap plants were transferred to a

controlled temperature room (20 °C) for 48 h (100% humidity) after being exposed in the field

for 2–6 days for rainfall events, and for one day for irrigation events. After a 48 h incubation

period, trap plants were transferred to a glasshouse (20 °C) to allow lesion development.

Lesions on all plant parts were counted after two weeks, which gave an estimate of the

number of conidia released and the distance travelled. Trap plants at all distances were

infected in all sprinkler irrigation and rainfall events. The highest number of lesions on trap

plants were recorded closest to the infected plots – the numbers decreased as the distance

from the infected plots increased. There was a positive relationship between the amount of

rainfall and the number of lesions recorded. A generalised additive model was developed that

efficiently described spatial patterns of conidial spread. With further development, the model

can be used to predict the spread of *A. rabiei*. This is the first systematic study to show that

conidia distribute *A. rabiei* over longer distances than previously reported.

61   **Keywords:** Ascochyta blight, epidemiology, conidial spread, wind-driven rain, chickpea,
62   sprinkler irrigation.

Introduction

Chickpea (*Cicer arietinum* L.) is the second most important legume crop globally and is the most widely grown legume grain crop in Australia with >1,060,000 ha harvested in 2018 (FAOSTAT 2020). Ascochyta blight caused by *Ascochyta rabiei* (syn. *Phoma rabiei*) is one of the most devastating chickpea diseases worldwide (Pande et al. 2005). With the exception of the Ord region in northern Western Australia, *A. rabiei* is the major biotic constraint to chickpea production in Australia with almost all chickpea growing areas affected (Bretag et al. 2008). *Ascochyta rabiei* survives on infected seed, volunteer chickpea plants and infested stubble, forming conidia that initiate primary infection. *Ascochyta rabiei* survives for up to two years on infested stubble at the ground surface but loses viability in up to five months when infested stubble are 5–40 cm deep in soil (W. J. Kaiser 1973; Navas‐Cortés et al. 1995). Conidia are hyaline, oval to oblong in shape, none to one septate, slightly bent at one or both ends and occasionally bicelled with size range from 8.2 to 10 x 4.2 to 4.5 µm (Nene 1982). The optimum range for the fungus growth, pycnidial production and conidial germination is 20 °C ; temperatures below 10 °C and above 30 °C have been found unfavourable for the fungus (Nene 1982; W. J. Kaiser 1973). Although there is no information on the effect of natural ultra violet light on *A. rabiei* conidia survival (Coventry 2012), exposure of cultures to wavelengths in the near ultra violet and blue range increased the rate of mycelial growth, pycnidial formation and conidial production in all isolates (W. J. Kaiser 1973).

Infected seed gives rise to infected seedlings through transmission from germinating seeds. Conidia are spread by rain splash or wind driven rain. Pycnidial formation, conidial production, host infection and disease development are favoured by temperatures between 5 and 30 °C (optimum 20 °C), relative humidity > 95 % (Nene 1982), and wetness period of 10 h or more (Khan 1999). A longer wetness period is required for spore germination at sub-optimal temperatures, and infection is rare in hot and dry conditions (Jhorar et al. 1998). Relative humidity is more important than temperature, thus *A. rabiei* development is very limited at lower humidity (e.g. 86 %) irrespective of temperature (Navas‐Cortés et al. 1998). Symptoms develop within 5–6 days after infection in cool and moist (20 °C and over 90 % humidity) conditions (Khan 1999). Pycnidia develop on the infected tissues, producing conidia that are responsible for the secondary spread of the pathogen (Pande et al. 2005). Conidia are the only spore type to have been reported in Australia. Putative ascospores have been reported only once in Australia (Galloway and MacLeod 2003), however, only one mating type has been detected and detailed molecular studies have shown Australian *A. rabiei* populations to be asexually reproducing (Leo et al. 2016; Bruvo et al. 2004; Mehmood et al. 2017), thus both primary and secondary infections are attributed to conidial infection.

Rain is needed for *A. rabiei* to infect and spread within a crop; dew alone will not
result in significant spread (Moore et al. 2016). During a rain event, pycnidia imbibe water
and hydrostatic pressure forces conidia in the pycnidia to exude through the ostiole in a
cirrhus. When a raindrop strikes the cirrhus, the kinetic energy breaks up the spore mass and
disperses the conidia to nearby healthy plants, thus spreading the disease (W. Kaiser 1992;
Nene and Reddy 1987; Coventry 2012). *Ascochyta rabiei* conidia have been reported to
spread over 1 m by splash dispersal (Kimber 2002). The amount of kinetic energy in
raindrops varies with the rainfall intensity, duration and size distribution with small drops
having less kinetic energy and thus remove fewer spores (Sache 2000). Splash dispersal by
large raindrops is considered more destructive because they tend to carry more spores
(Sache 2000). Spores can also be washed from upper leaves to lower leaves (Sache 2000).
Thunderstorms cause rapid spore dispersal from pycnidia and can result in diminution of
inoculum (Coventry 2012). Low intensity, intermittent rains combined with high wind have
been found to disperse spores more efficiently without diminishing the spore supply
(Coventry 2012).

Irrigation is the principal mechanism for increasing grain yield, supplementing rainfall
in drought seasons, or in areas receiving low rainfalls. Compared with furrow or flood
irrigation, sprinkler irrigation results in the wetting of the entire crop canopy and constitutes a
major source of moisture on all plant parts during dry conditions, thus increasing the risk of
foliar diseases (Rotem and Palti 1969). Sprinkler irrigation plays an important role in the
epidemiology of splash dispersed pathogens because spores are dispersed to their host
under moist conditions and their germination is practically ensured (Lomas 1991). Unlike
other forms of irrigation, water droplets generated by overhead sprinkler irrigation have the
impact energy to dislodge spores (Lomas 1991). This makes it likely to play an important role
in conidial spread for Ascochyta blight, *as A. rabiei* requires the impact energy of falling drops
to disperse its conidia from pycnidia. Despite this, there is no study on the distance and
quantity of *A. rabiei* conidia disperse during sprinkler irrigation events. Moreover, an
assumption exists that conidia disperse over short distances during sprinkler irrigation
events. Kimber et al. (2007) studied the spread of *A. rabiei* from primary infection foci
(infected seedlings) in small plots (15 m x 9 m) in Israel. Sprinkler irrigation was used to
irrigate plots, but only during dry periods, to provide conditions conducive to disease spread.
Moreover, the number of sprinkler irrigation events and amount of water applied during each
irrigation event were not recorded; therefore, any quantifiable relationship between irrigation,
disease and the pathogen spread (via conidia) was not established.

It has been reported that *A. rabiei* conidia spread over short distances (~1 m) via
splash dispersal, and up to 10 m in the presence of rain and strong wind (in the direction of

wind) (Kimber 2002; Kimber et al. 2007). However, the actual distance and number of conidia

spread via wind driven rain have not been determined systematically (Coventry 2012).

Kimber et al. (2007) showed Ascochyta blight spread over short distances (< 10 m) from

primary infection foci (infected seedlings) in the direction of the prevailing winds. However,

this was spread from a rather limited quantity of inoculum (a single infected seedling). The

potential spread is likely to be greater when inoculum is plentiful. Previous studies have

considered the spread of *A. rabiei* in biological rather than epidemiological terms, focusing on

disease development and spread in the paddock and not on the spread of the pathogen via

conidia (Coventry 2012). However, the presence of a pathogen does not automatically lead

to disease development. Disease occurs due to complex interactions between a virulent

pathogen, a susceptible host(s) and an environment favorable for long enough for the

pathogen to cause disease (Agrios 2005).

Trapping spores on 'trap plants' distributed at various distances from the source of

infection for a short period of time is an excellent way to investigate the distance of conidial

spread. Trap plants are distributed at various distances from a source of infection for a

relatively short period of time to trap spores, and then transferred to a glasshouse to allow

lesion development (Salam et al. 2011). The numbers of lesions give an estimate of the

number of spores released and the distance travelled. Lesion development indicates that the

dispersed conidia are viable. The actual distance *A. rabiei* conidia travel from a standing crop

has not been determined in such a systematic manner on a large scale. Based on

preliminary work aimed to determine the distance of *A. rabiei* conidia dispersal from previous

year's infested stubble (Galloway, unpublished), we propose the pathogen is likely to spread

over longer distances, than reported in the literature, in wind driven rain. Experiments

conducted on plant pathogens, in a wind tunnel and rain tower, have shown that spores

released by rain splash and then transported by wind are likely to be important in long-

distance dispersal of pathogens (Fitt et al. 1989). Improving our understanding on how *A.*

rabiei may spread in wind driven rain can help in predicting spatial patterns of conidial spread

during rainfall events, and subsequent disease epidemics through modelling techniques

(Coventry 2012). Models are required to predict disease risk and spread for varying

environmental conditions, beyond their domain of study, as it is impractical to investigate

pathogens' epidemiology in all growing environments (Diggle et al. 2002).

To investigate *A. rabiei* conidial spread in epidemiological rather than biological terms,

focusing on the pathogen spread via conidia rather than the disease development and

subsequent spread in the paddock, the present study asked the following questions: (1) can

conidia spread over longer distances than previously reported, and (2) do conidia spread

over the same distances in rainfall and sprinkler irrigation events?

 Methods

 Study areas description

This study was conducted at three different locations, Horsham (-36.728879, 142.157311), a

paddock just outside Horsham (-36.745871, 142.115969) and Curyo (-35.779312,

142.778332), in Victoria, Australia. All three sites had not had chickpea planted on them for

at least 4 years. Trials established by Agriculture Victoria to assess the efficacy of fungicides

against Ascochyta blight in chickpea were used to determine the distance of conidial spread

in rainfall and sprinkler irrigation events. Genetically pure chickpea seed (variety PBA Striker)

was sourced from the previous year's disease free seed-increase plots. Chickpea plots were

sown to dimension 10 m x 2 m on 13 May 2019. A handful (~300 mL) of naturally infested

stubble inoculum was evenly scattered across each plot at three to four nodes plant growth

stage on 26 June 2019 to initiate infection. Infested residues were collected, after passing

through the back of the harvester, from the previous year's trial of similar nature. A domestic

mulcher was used to break down residues into smaller pieces, and inoculum was mixed

thoroughly before inoculating plots. Disease incidence for each plot was recorded as the

percentage of plot with stem infections (with 0 % being no stem infections; 50 % being half

the plot having stem infections; and 100 % being plot completely dead). Control plots (that

received no fungicide treatments) had 90–100 % plants dead when conidia spread

experiments were conducted.

Experimental design

A method used by Salam et al. (2011) was modified for this study. At each location, 2–3

weeks old chickpea trap plants (variety PBA striker) were distributed at the distances of 0,

10, 25, 50 and 75 m from the control plots downwind of the prevailing wind direction.

Downwind directions for the control plots were determined based on previous years' weather

data. Trap plants were placed at the distances of 0–75 m along radial transects (there were

no other chickpea crops downwind of the control plots where chickpea trap plants were

distributed). There were 10 transects separated by 10-degree increments for a total of 90

degrees between the first and the last transect. There were four trap plant stations (trap

plants units) at both 0 and 10 m, and 10 trap plant stations at the distances of 25, 50 and

75 m (Table 1, Fig. S1). At the distances 0–25 m, each trap plant station consisted of 10

chickpea seedlings in two pots (five seedlings per pot). At the distances 50 and 75 m, each

trap plant station consisted of three chickpea seedlings in a single pot. The distances

between individual trap plant stations at each distance (0, 10, 25, 50 and 75 m) were

determined using the Law of Cosine (Table 1). As there were only 4 trap plant stations at the

distances of 0 and 10 m, they were placed along transects 1, 4, 7 and 10, with 30 degrees

difference (space) between each trap plant stations, to increase the chances of trapping

conidia. At distances of 25–75 m: all 10-trap plant stations were placed along 10 transects

with 10 degrees space between the individual trap plant stations (Table 1). The locations of

trap plant stations across transects were marked with flags to ensure trap plants were always

exchanged in the same spot for the entire duration of the experiment (Table S1). No

fungicides were applied over the duration trap plants were deployed in the paddock (Table 2).

Spore trapping

The distance of conidial spread was determined from September through to

December 2019. Spores were trapped in two rainfall events (event one ~ 4. 6 mm, and event

two ~ 18.6 mm) in Horsham, and a single rainfall event of 0.8 mm in Curyo (Table 2). A

sprinkler irrigation system (Hunter ®: Model MP300090210) located in a paddock just outside

Horsham was used to determine the distance of conidial spread for three irrigation events.

Sprinklers were set 110 cm above ground and were operated at 3.5 bar pressure. Each

sprinkler irrigation event was equivalent to 10–12 mm of simulated rain over approximately

90 minutes (a small rain shower of ~0.6 mm occurred during sprinkler irrigation event 3)

(Table 2).

In order to determine the distance of conidial spread in rainfall and sprinkler irrigation

events, trap plants were distributed in the paddock at the distances of 0–75 m from control

plots before forecast rainfall and scheduled irrigation events, respectively. Trap plants were

transferred to a controlled temperature room set at 20 °C for 48 hours (100 % humidity) after

being exposed in the field for 2–6 days for rainfall events, and for one day for irrigation

events (Table 2). The exposure period in paddock for the irrigation events was shorter for two

reasons. Firstly, each irrigation event was a single continuous controlled event as opposed to

natural rainfall events that fell in multiple showers over several days. Secondly, only the

infected plot was being irrigated (trap plants at the distances of 10–75 m were not getting

wet), therefore longer exposure period may have led to conidial desiccation on trap plants at

the distances of 10–75 m. After 48 hours incubation period, the trap plants were transferred

to a glasshouse (20 °C) to allow lesion development. Plants were watered daily via trays to

prevent splash dispersal to nearby plants. The numbers of lesions were counted on all plant

parts after two weeks giving an estimate of the number of conidia released and the distance

travelled.

Weather data was recorded by an automated weather station (Measurement

Engineering Australia, Adelaide, Australia). The data collected composed of date and time,

air temperature (° C), dew point (˚ C), rainfall (mm), relative humidity (%), wind speed (ms^{-1}),

and average wind direction (˚) recorded at 10-minutes intervals. The wind vane was not

properly calibrated for the weather station at Curyo. Therefore, Birchip Cropping Group's

(BCG) agriculture farm (located 10 km away from the experimental location) weather

station's (Davis Australia) wind direction data was used for the interpretation of results (wind

direction data was not included in the analysis, as wind direction is highly unstable and

constantly changes). Wind direction data, recorded by Australia Bureau of Meteorology, at

two locations (Hopetoun Airport 39 km, and Charlton 75 km; located in the same area as

BCG and Curyo) were retrieved using the R programming language (R Core Development

Team 2019) contributed packages 'bomrang' v0.7.0 (Sparks et al. 2017) and 'stationaRy'

v0.5.1 (Iannone 2020) and compared with the BCG's weather station data to ensure the

BCG's weather station wind direction data was valid (Fig. S2).

Development of a model to describe spatial patterns of conidial spread

A preliminary linear mixed model analysis was conducted in the R programming language,

using the lmer() function from the contributed package 'lme4' v1.1-1.23 (Bates et al. 2015), to

investigate the influence of 'spread event' (as a random variable: any spread event

regardless of sprinkler irrigation or rainfall events) and 'experimental location' on the number

of lesions recorded. A reductive approach was used to investigate the significance of the

random variables 'spread event' and 'experimental location' before modelling spatial patterns

of conidial spread. Preliminary analysis showed the influence of 'experimental location' to be

non- significant implying conidial spread patterns were consistent between experimental

locations. This result allowed generalised additive model (GAM) to compare the influence of

weather variables without the need to include experimental location as a dependant factor.

Generalised additive models were used because they have been reported to efficiently

describe the effect of various environmental variables on plant pathogen interactions (Sparks

et al. 2011; Yee and Mitchell 1991). The GAM for conidial dispersion, based on the pooled

data across all experimental locations for all six spread events, was fitted using the R

programming language via the gam() function from the contributed package 'mgcv' v1.8-31

(Wood 2011). Standard information criterion, such as Akaike's Information Criterion (AIC)

(Akaike 1974), the Bayesian Information Criterion (BIC) (Gideon 1978), as well as adjusted

R-squared values were compared across models to select the best fitting model.

Letting the mean number of lesions for pot i (pooled across all experimental locations

for all six spread events) be the response variable Y_i, and $\boldsymbol{x_i}$ be the associated covariate

vector with the distance of pot i from the infected plot being x_{i1}, the mean wind speed being

x_{i2} and the amount of irrigation or rainfall being x_{i3}, during spread events, the GAM model

that achieved best fit was of the form:

$$Y_i \sim \mathsf{Tw}_p(\mu_i, \phi), \qquad \log \mu_i = \alpha + \sum_{j=1}^{3} f_j(x_{ij})$$

where the f_j are independent "thin-plate splines" which are the default smoother used by the

gam() function. For details about the mathematical specification of these and other

smoothing splines, see Wood (2017). The use of the Tweedie distribution family to model the

response data is now standard in the case of interval scale measurements with substantial

zero-inflation, i.e., a positive probability mass at zero will indicate no lesions were observed

(Dunn 2004; Hasan and Dunn 2010). As it is implemented in the tw() function passed to the

family parameter in the gam() function, the power parameter p – which relates the Tweedie

variance to its mean – is estimated from the data (Wood 2011). The chosen link function for

this family was the logarithm link; the canonical or natural link function for Tweedie

distributions, and the default supplied to the gam() function. The gam() function also allowed

the dimension of the basis for smoothed terms to be manually selected via the parameter k,

which controls the "wiggliness" of the resultant curve. The value of k was chosen to be 5

($i.e. k = 5$) for all three predictors due to rather limited data.

Data cleaning and visualisation

Data were processed and visualised in R using the contributed packages 'tidyverse' v1.3.0

(Wickham et al. 2019), 'lubridate' v1.7.8 (Grolemund and Wickham 2011) and` 'clifro' (Seers

and Shears 2015) for wind roses. The best fitting GAM was visualised using a custom

function created for this work that uses the contributed package 'mgcViz' v0.1.6 (Fasiolo et

al. 2019). Mean wind speed was calculated using a circular averaging function from

'SDMTools' v1.1-221.2 (VanDerWal et al. 2014).

Results

Model fit

The linear mixed model analysis indicated that experimental location had no significant effect

on the number of conidia trapped and distance travelled during each spread event, thus

experimental location was not used when fitting the GAMs. Thirteen GAM models were

evaluated to describe spatial patterns of conidial spread. The model selected as the best

fitting had an adjusted R-squared value of 0.674, AIC value of 663 and a BIC value of 709.

The smooth terms for distance (D) and precipitation (P) were significant terms (p < 0.05). The

smooth term for wind speed (W) was not significant. However, including this term only as a

linear predictor reduced the fitness of the model, R-squared value of 0.657, AIC value of 689

and BIC value of 729, so the smoothed term was included in the final model (Fig. 1).

 Dispersal in rainfall events

Trap plants at all distances (0–75 m) were infected in all 3-rainfall events, indicating conidia

spread to 75 m in each rainfall event. There was a strong relationship ($R^2 = 0.657$) between

mean wind speed, rainfall intensity, distance from the infected plot, and the number of

conidia trapped (counted as lesions). The highest number of lesions on trap plants were

recorded closest to the control plot at 0 m in each rainfall event - the numbers decreased as

the distance from the control plots increased (Fig. 2). The number of lesions recorded

increased with increasing rainfall. At Horsham, a total of 159 lesions were recorded at all

distances in rainfall event 1 (~ 4.8 mm), and 275 lesions were recorded in rainfall event 2

(~18.6 mm) (Table 2). In rainfall event 1 (Horsham Rain 1), a uniform conidial spread pattern

across transects was observed at the distances of 0–25 m (i.e., all trap plant stations at the

distances of 0–25 m were infected), whereas a non-uniform conidial spread pattern across

transects was observed at the distances of 50 and 75 m (Fig. 3). Uniform conidial spread

pattern across transects at all distances was observed in rainfall event 2 (Horsham Rain 2).

Mean wind speed and direction varied during rainfall events 1 and 2 (Fig. 4). Mean wind

direction was predominantly in the direction of trap plants (NE, SE) during both spread

events (Fig. 4)

At Curyo, a total of 193 lesions were recorded in a single rainfall event (Curyo Rain 1) of 0.8

mm (Table 2). Conidial spread pattern was almost uniform across transects at the distances

of 0–50 m, while a non-uniform conidial spread pattern across transects was observed at 75

m (Fig. 3). Mean wind speed and direction varied during the spread event (Fig. S2). Mean

wind direction was not in the direction of trap plants (Fig. S2) but, compared to rainfall

events-1 and 2, wind direction appeared to have limited influence on the number of lesions

recorded (Table 2), and the spatial patterns of conidial spread at Curyo (Fig. 3).

Dispersal with sprinkler irrigation

Trap plants at all distances (0–75 m) were infected in all three irrigation events. There was a

strong correlation between mean wind speed, irrigation, distance from the infested plot and

the number of conidia trapped (counted as lesions). The highest numbers of lesions on trap

plants were recorded closest to the control plot in each irrigation event - the numbers

decreased as the distance from the control plot increased (Fig. 2). The total number of

lesions recorded on trap plants varied greatly between all three irrigation events, with a total

of 250 lesions recorded in event 1 (~10–12 mm), 368 lesions in event 2 (~10–12 mm) and

477 lesions in event 3 (~10–12 mm irrigation event and a small rain shower of 0.6 mm)

(Table 2). In event 1 and 3 (Horsham Irrg1 and Horsham Mxd1), a uniform conidial spread

pattern across almost all transects was observed at the distances of 0–50 m, whereas

conidial spread pattern across transects was not uniform at 75 m (Fig. 3). In event 2: conidial

spread pattern across transects was uniform at the distances of 0–25 m, whereas a non-

uniform conidial spread pattern across transects was observed at the distances of 50 and 75

m (Fig. 3). Mean wind speed and direction varied between each spread event (Fig. 4).

Maximum mean wind speed (6.5 ms^{-1}) (Table 2) and maximum gust (11.2 ms^{-1}) were

recorded was recorded during irrigation event 3. Mean wind direction was predominantly in

the direction of trap plants (NE, SE) during all spread events (Fig. 4).

Discussion

Conidia were shown to distribute *A. rabiei* over longer distances than previously reported.

Conidia dispersed 75 m from infected plots in all sprinkler irrigation and rainfall events. The

mean number of lesions recorded on trap plants decreased as the distance from infected

plots increased. There was a strong relationship (R^2 = 0.657) between mean wind speed, the

amount of precipitation (rainfall and/or sprinkler irrigation) and the number of conidia trapped.

The experimental location had no significant effect on the number of conidia trapped and

distance travelled during each spread event (P>0.01). A model was developed that efficiently

described spatial patterns of conidial spread under field conditions. Given more data and

further development, this model can be used to predict the spread of *A. rabiei*.

357   The highest mean number of lesions on trap plants were recorded closest to the

control plots in all spread events – the numbers decreased as the distance from the control

plots increased. These results support the findings from previous studies (Coventry 2012;

Kimber et al. 2007) who showed *A. rabiei* inoculum density decreased as the distance from

infected foci increased. A similar dispersal pattern has been observed for other splash

dispersed pathogens, such as *Colletotrichum acutatum* (Yang et al. 1990),

Parastagonospora nodorum (previously known as *Septoria nodorum*) (Griffiths and Ao 1976),

and *Ascochyta fabae* (Pedersen et al. 1994). For splash dispersed pathogens, plants close

to the source of infection are more exposed to spore load due to localised splash dispersal

and become infected quickly, resulting in more lesions (Coventry 2012). More lesions were

recorded on trap plants with increased rainfall and in an event with a combination of a small

rain shower and a sprinkler irrigation event compared to a sprinkler irrigation event alone (in

case of sprinkler irrigation event 3, i.e., Horsham Mixd), suggesting that increased

precipitation dislodged more conidia, which were dispersed by wind driven rain. The amount

of rainfall has been reported to have a major influence on Ascochyta blight severity and the

distance disease spreads (Coventry 2012).

373   Wind speed and direction coinciding with increased rainfall influenced the spatial

patterns of conidial spread. Uniform conidial spread across transects at all distances was

observed in rainfall event 2 (Horsham Rain 2), which indicates strong winds from NW
directions coincided with increased rainfall and dispersed conidia across all transects
downwind (SE, NE) from the infected plot. Although trap plants were located downwind in
rainfall event 1 (Horsham Rain 1), uniform conidial spread across all transects was not
observed at 50–75 m, which reflects the influence of reduced rainfall and rather low wind
speed (compared to rainfall event 2) during the spread event. Compared to rainfall event 1 at
Horsham, the amount of rainfall and wind direction appeared to have limited influence on the
number of conidia trapped and the spatial patterns of conidial spread for rainfall event 3
(Curyo Rain 1) at Curyo. More lesions (193 lesions) were recorded in a single rainfall event
of 0.8 mm at Curyo, although mean wind direction was not in the direction of trap plants. In
addition, conidial spread pattern was almost uniform at the distances of 0–50 m. Conversely,
fewer lesions (159) were recorded in a rainfall event of 4.8 mm at Horsham when trap plants
were located downwind; and conidial spread pattern was uniform at the distances of 0–25 m
only. However, a direct comparison between these two rainfall events is not possible due to a
location difference and rather limited data. Furthermore, our model does not consider
continuously changing wind direction and dynamics associated with turbulence, which may
assist in explaining variation in spatial patterns of conidial spread. Conidia dispersed 75 m
and spatial pattern of conidial spread was uniform across transects at the distances of 0–50
m, when the amount of rainfall was as low as 0.8 mm (Curyo Rain 1 event with mean wind
speed of 3.6 ms^{-1}); and mean wind direction was not predominantly in the direction of trap
plants. This suggests that only a small amount of rain is required to dislodge a large number
of conidia. A similar dispersal pattern has been reported for the sexual spores (ascospores)
of *A. rabiei*, where ascospores discharge depended more on the occurrence of rain than the
amount (Trapero-Casas et al. 1996).

More conidia (368) were trapped in sprinkler irrigation event 2, compared to sprinkler
irrigation event 1, but conidial spread pattern across transects was not uniform at the
distances of 50–75 m. Conversely, fewer conidia (250) were trapped in sprinkler irrigation
event 1, but conidial spread pattern was almost uniform across transects at all distances.
The lower number of spores and non-uniform spatial spread pattern for sprinkler irrigation
event 2 is rather surprising because an equal amount (~10–12 mm) of irrigation was applied
in both sprinkler irrigation events. Moreover, stronger mean wind speed was recorded for
sprinkler irrigation event 2 and the non-infected trap plants were located downwind (the
mean wind direction was from the west, therefore trap plants placed at the distances of 50–
75 m east of the infected plot were expected to be uniformly infected). The reasons for this
dispersal pattern are not clear, and our analysis is limited by rather limited data. Detailed

studies are needed to get a more complete picture of spatial patterns of conidial spread
during sprinkler irrigation events.

Above canopy irrigation reduces mean ambient air temperature by 7–9 °C and
increases humidity up to 30 % during the irrigation period (Rotem and Palti 1969). This has
implications for Ascochyta blight development within an irrigated chickpea paddock, as *A.*
rabiei only requires 3–10 h leaf wetness for host penetration and infection at the optimum
temperature (Khan 1999; Moore et al. 2016). Therefore, extended periods of sprinkler
irrigation events may not be required for disease development, especially in a cool and moist
season. In the laboratory; more than 50 % of conidia, incubated on membrane filters,
germinated after four days at relative humidity as low as 12.5 % (temperature 20 °C)
(Coventry 2012). *Ascochyta rabiei* conidia have been shown to survive and remain infective,
when inoculated on chickpea seedlings in a growth chamber, during dry periods lasting 24 h
(Armstrong-Cho et al. 2004). However, conidia are unlikely to withstand dry conditions for
extended periods of time under field conditions, especially at sub-optimal temperatures. For
conidial spread with sprinkler irrigation, this implies that even though conidia may spread to
equal distances (75 m in this case) during sprinkler irrigation and rainfall events, leaf wetness
provided by dew and/or humidity over 90% to facilitate conidial penetration and infection will
be important in establishing disease on chickpea outside the irrigated paddock in the
absence of rain (if temperature requirement is met). Conversely; rainfall usually occurs over
a large area, and only a few mm of rain later in the day or at night satisfies the leaf wetness
requirement (Moore et al. 2016). Therefore, leaf wetness provided by dew and/or high
relative humidity is not necessary to facilitate conidial penetration and infection at longer
distances for conidia dispersed in rainfall events. This further implies that rainfall might be
more important than sprinkler irrigation for conidial spread and subsequent disease
development at farther distances. Future work determining the viability of conidia after
exposure to different duration of dry periods, under field conditions, is necessary to
understand the duration of dry period over which *A. rabiei* conidia can survive when
dispersed outside the irrigated paddock. To the best of our knowledge, this is the first
systematic study to determine conidia dispersal with sprinkler irrigation.

The finding that conidia travelled 75 m during all rainfall and sprinkler irrigation events
suggests that conidia can potentially play a role in the long-distance dispersal of *A. rabiei.*
Conidia dispersed 50 m from naturally infested stubble from the previous year (there were no
trap plants farther than 50 m to test long-distance dispersal) (Galloway, unpublished). The
sexual spores of the fungus (ascospores) have been reported to travel up to hundreds of
metres to several kilometres from the source of infection (Trapero-Casas et al. 1996; W.
Kaiser and Küsmenoglu 1997; Trapero-Casas and Kaiser 1992). The present study did not

have trap plants farther than 75 m from the infected plot and wind speed during spread
events was rather low with a maximum mean wind speed of 6.5 ms^{-1} and maximum gust of
11.2 ms^{-1}. It is likely that conidia could dispersed farther than 75 m, especially when wind
speed is high and there are strong wind gusts. It has been hypothesized that if conidial
dispersal is aided by turbulent winds and conidia enter the turbulent boundary layer, they
could be dispersed to distances that are associated with ascospores only (McCartney and
West 2007; Coventry 2012). While the role of conidia in long-distance dispersal has not been
recognized, it has been hypothesized that liquid suspensions in air (aerosols) can
incorporate a large number of conidia in heavy rain that can be transported over longer
distances by wind (Coventry 2012). A severe Ascochyta blight epidemic developed across
the southern region of Australia in 1998. It is speculated that aerosols developed during
heavy rains, which travelled over hundreds of meters to kilometers and contributed to the
epidemic development (Coventry 2012). This suggests that role of conidia in the long-
distance dispersal has been underestimated. Further work distributing trap plants at
distances farther than 75 m is necessary to determine the potential distance conidia can
travel.

When fitting observed data to logistic models, the known biology of pathogens needs
to be considered rather than simply considering the shape of the curve (Sparks et al. 2008).
Generalised additive models are data-driven not model-driven, as they allow the data to
determine the shape of the response curves rather than being limited by the shapes
attainable through some parametric specification. The inability to control environmental
factors, such as wind speed, amount of rainfall, temperature and humidity makes it difficult to
use parametric statistical models e.g., generalised linear models (GLMs) to make inferential
comparisons across sites without a high risk of the presence of confounding variables.
Generalised additive models offer a flexible modelling approach that provides a good
balance between highly interpretable linear models that have undesirable predictive
capabilities and black-box machine learning approaches, such as neural networks, which
have excellent predictive performance but low interpretability (Hastie and Tibsbirani 1990).
This capacity to achieve interpretable separation of covariate contributions in a strong
descriptive model is the main reason that GAMs were chosen to model spatial patterns of
conidial spread. A model was developed that efficiently described spatial patterns of conidial
spread under field conditions. The model will be useful for chickpea growers in identifying the
scale of potential damage and could be used to predict the spread of *A. rabiei* with further
development. When further developed, this model can also be used to inform chickpea
growers how far to grow chickpea from the neighbouring paddocks.

A comparison of conidial spatial dispersal patterns in identical amount of irrigation
and rainfall events would have been beneficial, but constantly changing wind speed and
direction makes a direct comparison difficult under field conditions. Conidial spread was
investigated in the late stage of plant growth with 90–100 % plants in plots being infected.
Detailed studies at different plant growth stages and disease incidence levels are now
needed to provide a more complete picture of distance conidia disperse and their spatial
dispersal patterns. It should be noted that counting lesions on trap plants gave an estimate of
only viable conidia dispersed (not the total number of conidia dispersed), which could cause
disease under field conditions if conditions are favourable for disease development. This
study was primarily conducted to get an estimate of the distance conidia travel and remain
viable. Future studies should consider using trap plants, spore traps and molecular
approaches (e.g. trapping spores sticky tapes and subsequent spore count using quantitative
PCR) to get an accurate estimate on the number of conidia dispersed. Spore trapping
coupled with quantitative PCR has been demonstrated to be a useful tool for epidemiological
studies and biosecurity surveillance (Vogelzang 2012). More work is required to establish the
equivalency of number of lesions to the number of conidia. Many other factors are involved in
the spore dispersal, including changing wind directions, turbulence and topographic factors
(Coventry 2012), which add complexity to the model and were not considered in the present
model. These factors need to be addressed in future models.

Acknowledgements

Grains Research and Development Corporation (GRDC), Australia, provided financial
assistance in this work through the project USQ-1903-003RTX. Agriculture Victoria and
GRDC provided financial assistance in this work through their co-investment project
DAV00150, and DJP1097-001RTX.

We thankfully acknowledge Jason Brand and his team for trial sites management. We thank
Andrew Hallet for technical support and assistance with field work.

Conflict of interest

We confirm that we have no conflict to interest to disclose.

Ethical approval

This study did not involve working with animals or humans

Consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

The raw data are documented and available from https://doi.org/10.5281/zenodo.3842293.

All raw and generated data and further associated materials have been made further

available as a part of a research compendium available from

https://doi.org/10.5281/zenodo.3810826.

Code availability

All code used in the analyses and data visualisation have been made available as

a research compendium, available from https://doi.org/10.5281/zenodo.3810826.

References

Agrios, G. (2005). *Plant Pathology*. San Diego, CA: Elsevier academic press.

Akaike, H. (1974). A new look at the statistical model identification. *IEEE Transactions on*
Automatic Control, 19(6), 716-723.

Armstrong-Cho, C., Gossen, B., & Chongo, G. (2004). Impact of continuous or interrupted
leaf wetness on infection of chickpea by *Ascochyta rabiei*. *Canadian Journal of*
Plant Pathology, 26(2), 134-141.

Bretag, T., MacLeod, W., Kimber, R., Moore, K., Knights, E., & Davidson, J. (2008).
Management of Ascochyta blight in chickpeas in Australia. *Australasian Plant*
Pathology, 37(5), 486-497.

Bruvo, R., Michiels, N. K., D'SOUZA, T. G., & Schulenburg, H. (2004). A simple method for
the calculation of microsatellite genotype distances irrespective of ploidy level.
Molecular Ecology, 13(7), 2101-2106.

Coventry, S. A. (2012). Factors affecting short and long distance dispersal of fungal
pathogens: chickpea Ascochyta blight as a model. PhD dissertation. The
University of Adelaide, Australia.

Diggle, A. J., Salam, M. U., Thomas, G. J., Yang, H., O'Connell, M., & Sweetingham, M. (2002).
AnthracnoseTracer: a spatiotemporal model for simulating the spread of
anthracnose in a lupin field. *Phytopathology, 92*(10), 1110-1121.

Dunn, P. K. (2004). Occurrence and quantity of precipitation can be modelled
simultaneously. *International Journal of Climatology: A Journal of the Royal*
Meteorological Society, 24(10), 1231-1239.

FAOSTAT (2020). FAOSTAT database. Available at http://www.fao.org/faostat/en/ -
data/QC.

Fasiolo, M., Nedellec, R., Goude, Y., & Wood, S. N. (2019). Scalable visualization methods
for modern generalized additive models. *Journal of computational and Graphical*
Statistics, DOI: 10.1080/10618600.2019.1629942.

Fitt, B. D., McCartney, H., & Walklate, P. (1989). The role of rain in dispersal of pathogen
inoculum. *Annual Review of Phytopathology, 27*(1), 241-270.

Galloway, J., & MacLeod, W. (2003). Didymella rabiei, the teleomorph of *Ascochyta rabiei*,
found on chickpea stubble in Western Australia. *Australasian Plant Pathology,*
32(1), 127-128.

Gideon, S. (1978). Estimating the dimension of a model. *The Annals of Statistics, 6*(2),
461-464.

Griffiths, E., & Ao, H. C. (1976). Dispersal of *Septoria nodorum* spores and spread of
glume blotch of wheat in the field. *Transactions of the British Mycological Society,*
67(3), 413-418.

Grolemund, G., & Wickham, H. (2011). Dates and times made easy with lubridate. *Journal*
of Statistical Software, 40(3), 1-25.

Hasan, M. M., & Dunn, P. K. (2010). A simple Poisson–gamma model for modelling rainfall
occurrence and amount simultaneously. *Agricultural and Forest Meteorology,*
150(10), 1319-1330.

Hastie, T. J., & Tibsbirani, R. (1990). *Generalized Additive Models*. New York: Chapman and
Hall.

Iannone, R. (2020). stationaRy: Detailed Meteorological Data from Stations All Over the
World. R package version 0.5.1. https://CRAN.R-project.org/package=stationaRy.

Jhorar, O., Butler, D., & Mathauda, S. (1998). Effects of leaf wetness duration, relative
humidity, light and dark on infection and sporulation by *Didymella rabiei* on
chickpea. *Plant Pathology, 47*(5), 586-594.
Kaiser, W. (1992). Epidemiology of Ascochyta rabiei. In K. Singh, & M. Saxena (Eds.),
Disease resistance breeding in chickpea (pp. 117-134). Aleppo, Syria: ICARDA.
Kaiser, W., & Küsmenoglu, I. (1997). Distribution of mating types and the teleomorph of
Ascochyta rabiei on chickpea in Turkey. *Plant Disease, 81*(11), 1284-1287.
Kaiser, W. J. (1973). Factors affecting growth, sporulation, pathogenicity, and survival of
Ascochyta rabiei. Mycologia, 65(2), 444-457.
Khan, M. S. A. (1999). Epidemiology of Ascochyta blight of chickpea in Australia. PhD
dissertation. The University of Adelaide,
Kimber, R. (2002). Survival and transmission of *Ascochyta rabiei* (Ascochyta blight) of
chickpea. Honours thesis, University of Adelaide, Australia.,
Kimber, R., Shtienberg, D., Ramsey, M., & Scott, E. (2007). The role of seedling infection in
epiphytotics of Ascochyta blight on chickpea. *European Journal of Plant Pathology,*
117(2), 141-152.
Leo, A. E., Linde, C. C., & Ford, R. (2016). Defence gene expression profiling to *Ascochyta*
rabiei aggressiveness in chickpea. *Theoretical and Applied Genetics, 129*(7), 1333-
1345.
Lomas, J. (1991). Sprinkler irrigation and plant disease under semi‐arid climatic
conditions 1. *EPPO Bulletin, 21*(3), 365-370.
McCartney, A., & West, J. (2007). Dispersal of fungal spores through the air. In J.
Dijksterhuis, & R. A. Samson (Eds.), *Food Mycology - A Multifaceted Approach to*
Fungi and Food (pp. 79-96): CRC press.
Mehmood, Y., Sambasivam, P., Kaur, S., Davidson, J., Leo, A. E., Hobson, K., et al. (2017).
Evidence and consequence of a highly adapted clonal haplotype within the
Australian Ascochyta rabiei population. *Frontiers in plant science, 8*, 1029.
Moore, K., Hobson, K., Sambasivam, P., Ford, R., Harden, S., Nash, P., et al. (2016).
Chickpea ascochyta - is the pathogen changing and what are the implications for
management. Available at https://grdc.com.au/resources-and-publications/grdc-
update-papers/tab-content/grdc-update-papers/2015/02/chickpea-ascochyta-
is-the-pathogen-changing-and-what-are-the-implications-for-management.
Navas‐Cortés, J., Trapero‐Casas, A., & Jiménez‐Díaz, R. (1995). Survival of *Didymella*
rabiei in chickpea straw debris in Spain. *Plant Pathology, 44*(2), 332-339.
Navas‐Cortés, J., Trapero‐Casas, A., & Jiménez‐Díaz, R. (1998). Influence of relative
humidity and temperature on development of Didymella rabiei on chickpea
debris. *Plant Pathology, 47*(1), 57-66.
Nene, Y. (1982). A review of Ascochyta blight of chickpea. *International Journal of Pest*
Management, 28(1), 61-70.
Nene, Y., & Reddy, M. (1987). Chickpea diseases and their control. In M. Saxena, & R.
Singh (Eds.), *The Chickpea* (pp. 233–270). Wallingford, UK: CAB International.
Pande, S., Siddique, K., Kishore, G., Bayaa, B., Gaur, P., Gowda, C., et al. (2005). Ascochyta
blight of chickpea (*Cicer arietinum* L.): a review of biology, pathogenicity, and
disease management. *Australian Journal of Agricultural Research, 56*(4), 317-332.
Pedersen, E., Morrall, R., McCartney, H., & Fitt, B. D. (1994). Dispersal of conidia of
Ascochyta fabae f. sp. lentis from infected lentil plants by simulated wind and rain.
Plant Pathology, 43(1), 50-55.
R Core Development Team (2019). R: A language and environment for statistical
computing. Vienna, Austria: R Foundation for Statistical Computing.

Rotem, J., & Palti, J. (1969). Irrigation and plant diseases. *Annual Review of*
Phytopathology, 7(1), 267-288.
Sache, I. (2000). Short-distance dispersal of wheat rust spores. *EDP Sciences 20*, 757-767.
Salam, M. U., Galloway, J., Diggle, A. J., MacLeod, W. J., & Maling, T. (2011). Predicting
regional-scale spread of ascospores of *Didymella pinodes* causing ascochyta blight
disease on field pea. *Australasian Plant Pathology, 40*(6), 640-647.
Seers, B. M., & Shears, N. T. (2015). New Zealand's Climate Data in R—An Introduction to
clifro. The University of Auckland, New Zealand. URL:
http://stattech.wordpress.fos.auckland.ac.nz/2015-02-new-zealands-climate-
data-in-r-an-introduction-to-clifro/.

.
Sparks, A. H., Esker, P., Bates, M., Dall'Acqua, W., Guo, Z., Segovia, V., et al. (2008). Ecology
and epidemiology in R: disease progress over time. The Plant Health Instructor
DOI:101094/PHI-A-2008-0129-02.

.
Sparks, A. H., Forbes, G. A., Hijmans, R., & Garrett, K. A. (2011). A metamodeling
framework for extending the application domain of process‐based ecological
models. *Ecosphere, 2*(8), 1-14.
Sparks, A. H., Padgham, M., Parsonage, H., & Pembleton, K. (2017). bomrang: fetch
Australian Government Bureau of Meteorology data in R. *The Journal of Open*
Source Software, 2, 17.
Trapero-Casas, A., & Kaiser, W. (1992). Development of *Didymella rabiei*, the teleomorph
of *Ascochyta rabiei*, on chickpea straw. *Phytopathology, 82*(11), 1261-1266.
Trapero-Casas, A., Navas-Cortés, J., & Jimenez-Diaz, R. (1996). Airborne ascospores of
Didymella rabiei as a major primary inoculum for Ascochyta blight epidemics in
chickpea crops in southern Spain. *European Journal of Plant Pathology, 102*(3),
237-245.
VanDerWal, J., Falconi, L., Januchowski, S., Shoo, L., & Storlie, C. (2014). Species
Distribution Modelling Tools, Tools for processing data associated with species
distribution modelling exercises. *CRAN. R-project. org/package= SDMTools*.
Vogelzang, B. (2012). Spore trapping coupled with quantitative PCR: potential for
application to biosecurity surveillance. PhD dissertation. The University of
Adelaide, Australia.
Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., et al. (2019).
Welcome to the Tidyverse. *Journal of Open Source Software, 4*(43), 1686.
Wood, S. N. (2011). Fast stable restricted maximum likelihood and marginal likelihood
estimation of semiparametric generalized linear models. *Journal of the Royal*
Statistical Society (B), 73(1), 3-36.
Yang, X., Madden, L., Wilson, L., & Ellis, M. (1990). Effects of surface topography and rain
intensity on splash dispersal of *Colletotrichum acutatum*. *Phytopathology, 80*(10),
1115-1120.
Yee, T. W., & Mitchell, N. D. (1991). Generalized additive models in plant ecology. *Journal*
of Vegetation Science, 2(5), 587-602.

 Figure captions

Fig. 1 Generalised additive model fit for smoothed terms of three predictors: distance, mean
windspeed (mws) and total precipitation (sum_rain)

Fig. 2 Mean number of conidia dispersed (counted as lesions) during each spread event at
each distance, where 'n' (indicated by the point size) represents the number of pots that
shared the same mean number of lesions counted per trap plant per pot. The mean count
values are shown on the y-axis and distance conidia dispersed is shown on the x-axis. Red
dots indicate median lesions counted per trap plant. Conidia dispersed 75 m in all irrigation
and rainfall event

Fig. 3 Heatmaps showing spatial patterns of conidial spread during each spread event.
Spatial spread patterns were determined by counting lesions on trap plants (placed along
radial transects at the distances of 0–75 m from infected plots), where 'n' represents mean
number of lesions per trap plant per pot. Concentric circles show distance of trap plants from
the infected plots: the inner single solid dot represents 0 m distance and the outer most
dotted line represents 75 m distance from the infected plot

Fig. 4 Wind roses, for all spread events except Curyo Rain 1, showing wind speed and
direction recorded at 10 minutes interval