Conserved anti-inflammatory effects and sensing of butyrate in zebrafish

Pradeep Manuneedhi Cholan [1], Alvin Han [2], Brad R Woodie [1,2], Angela RM Kurz [3], Warwick

J Britton [1,4,5], Lihua Ye [2], Zachary C Holmes [2], Jessica R McCann [2], Lawrence A David [2],

John F Rawls [2], Stefan H Oehlers [1,4]

[1] Tuberculosis Research Program at the Centenary Institute, The University of Sydney,

Camperdown NSW 2050, Australia

[2] Department of Molecular Genetics and Microbiology, Duke Microbiome Center, Duke

University School of Medicine, Durham NC 27710, USA

[3] Centenary Imaging and Sydney Cytometry at the Centenary Institute, The University of

Sydney, Camperdown NSW 2050, Australia

[4] The University of Sydney, Discipline of Infectious Diseases and Immunology, Faculty of

Medicine and Health, and Marie Bashir Institute, Camperdown NSW 2050, Australia

[5] Department of Clinical Immunology, Royal Prince Alfred Hospital, Camperdown, NSW

2050 Australia

Corresponding author: Dr Stefan Oehlers stefan.oehlers@sydney.edu.au

ORCID: 0000-0003-0260-672X Twitter: @oehlerslab

Abstract

Short chain fatty acids (SCFAs) are produced by microbial fermentation of dietary fiber in the

gut. Butyrate is a particularly important SCFA with anti-inflammatory properties and is

generally present at lower levels in inflammatory diseases associated with gut microbiota

dysbiosis in mammals. We aimed to determine if SCFAs are produced by the zebrafish

microbiome and if SCFAs exert conserved effects on zebrafish immunity as an example of the

non-mammalian vertebrate immune system. We demonstrate that bacterial communities from

adult zebrafish intestines synthesize all three main SCFA *in vitro*, although SCFA were below

our detectable limits in zebrafish intestines *in vivo*. Immersion in butyrate, but not acetate or

propionate, reduced the recruitment of neutrophils and M1-type pro-inflammatory

macrophages to wounds. We found conservation of butyrate sensing by neutrophils via

orthologs of the *hydroxycarboxylic acid receptor 1* (*hcar1*) gene. Neutrophils from Hcar1-

depeleted embryos were no longer responsive to the anti-inflammatory effects of butyrate,

while macrophage sensitivity to butyrate was independent of Hcar1. Our data demonstrate

conservation of anti-inflammatory butyrate effects and identify the presence of a conserved

molecular receptor in fish.

Keywords: zebrafish, butyrate, short chain fatty acid, inflammation, neutrophil, macrophage,

tumor necrosis factor

Introduction

Short chain fatty acids (SCFAs) are microbial metabolites produced in the gut by the

anaerobic fermentation of dietary fiber and protein in the large intestine [1]. The most abundant

SCFAs are acetate, butyrate, and propionate. In addition to providing the host with an energy

source, microbially-derived SCFAs exert anti-inflammatory effects through inhibition of

histone deacetylates (HDAC) and activation of G-protein coupled receptors (GPCRs) [2]. Most

research on SCFAs has reported their anti-inflammatory properties in mammals [3-5]. However,

the anti-inflammatory mechanism responsible for the anti-inflammatory effects of SCFA

administration has not been reported in fish species to date.

Zebrafish are an important model of vertebrate gut physiology with key experimental

advantages including high fecundity, transparency, and well-developed gut digestive function

by 6 days post fertilization (dpf) [6]. There is a high degree of intestinal immune conservation

across vertebrates, including the sensitivity of zebrafish intestinal epithelial cell progenitors to butyrate [7-9]. However, SCFA production has not been previously observed in the intestines of zebrafish, and it is unclear if the intestinal lumen of the zebrafish intestine provides a suitable niche for SCFA production [9]. Nevertheless, detectable amounts of acetate, butyrate, and propionate have been measured in several species of teleosts [10-14].

We have used conservation between mammalian intestinal function and immunity to create zebrafish models of human intestinal inflammation [15, 16]. Key pattern recognition molecule families, such as the Toll-like receptors and Nod-like receptors, are evolutionarily ancient, and have conserved roles in zebrafish intestinal immunity [7, 8, 17, 18]. However, conservation of host molecules responsible for sensing SCFAs has not been explored in teleosts. Mammals utilize a wide range of molecules to sense SCFAs including G protein-coupled receptors (GPRs) GPR81 (also known as HCAR1) which is primarily present on immune cells and GPR109A which is expressed on intestinal epithelial cells. Microbially-derived SCFAs also exert direct effects on host physiology through histone deacetylase (HDAC) inhibition [19-21].

In this study, we investigated whether SCFAs are produced in the zebrafish intestine, and if the anti-inflammatory effects and sensing of SCFAs are conserved in zebrafish. We find that the pattern of SCFA production by the zebrafish intestinal microbiota is different from that seen in mammals, but that the anti-inflammatory effects mechanisms of butyrate are conserved across vertebrate species and development regardless of the ability of their endogenous microbiota to produce measurable butyrate.

Methods

2.1 Zebrafish handling

Adult zebrafish were housed at the Centenary Institute (Sydney Local Health District AEC

Approval 17-036) and Duke University School of Medicine ((Duke Institutional Animal Care

and Use Committee Protocol Approval A115-16-05). Adult zebrafish experimentation was

approved by the Institutional Animal Care and Use Committees of Duke University approval

A115-16-05. Zebrafish adults were reared, housed, and fed as previously described [22]. All

zebrafish embryo research experiments and procedures were completed in accordance with

Sydney Local Health District animal ethics guidelines under approval 17-036. Zebrafish

embryos were obtained by natural spawning and embryos were maintained and raised in E3

media at 28ºC.

2.2 SCFA quantification from adult zebrafish

Adult zebrafish were euthanized with 200 – 300 mg/L of ethyl 3-aminobenzoate

methanesulfonate (tricaine) (Sigma, E10521) prior to dissection. For each sample, intestines

dissected from five adult (90+ dpf) EK WT zebrafish males (roughly 0.2 g total) were pooled

and homogenized using a Precellys 24 High-Powered Bead Homogenizer at 5500 rpm for 3

cycles at 20 seconds per cycle with a 10 second delay between cycles. Samples were then

acidified with HCl to pH 2, pelleted by centrifugation, and the supernatant was harvested.

Filtered supernatant was stored at -80°C until quantification.

SCFA quantification as carried out on an Agilent 7890B GC FID, with an HP-FFAP capillary

column (25 m length, ID 0.2 mm, film thickness 0.33 µm). Concentrations were determined

using a linear model fit of a standard curve that encompasses the sample concentration range.

Standardized concentrations used for each C2-C5 SCFA were as follows: 0.2, 0.5, 1, 2, 4, and

8 mM.

2.3 In vitro synthesis of SCFA by zebrafish gut commensals

Freshly dissected intestines from four adult (6-month-old) EK WT zebrafish males and frozen

mouse fecal pellets were homogenized under reducing conditions to preserve the anaerobes.

Samples were handled in a Coy anaerobic chamber and used to inoculate tubes containing

brain-heart infusion (BHI) media (Thermo Scientific, OXOID) or Gifu anaerobic media

(Sigma), both supplemented with deoxygenated hemin and vitamin K to a final concentration

of 12.5 mg/L of hemin and 2.5 mg/L of vitamin K.

Tubes were incubated in a sealed anaerobic jar with a Gas-Pak (Becton and Dickinson) to

maintain anaerobic conditions at 28°C for 24 hours. Samples were then acidified to pH 2 with

HCl, pelleted by centrifugation, and the supernatant was harvested. Filtered supernatant was

stored at -80°C until quantification with methods identical to those listed above in Section

2.2.

2.4 Drug treatments

Embryos were treated with 30 mM sodium acetate (Sigma; S2889), 30 mM sodium butyrate

(Sigma; B5887), 30 mM sodium propionate (Sigma; P1880), 50 µg/mL dexamethasone

(Sigma; D4902), or 100 mM 6-aminocaproic acid (Sigma; A2504). Drug stocks were

dissolved in DMSO or PBS and added to E3.

2.5 Tail wounding experiment

Caudal fin amputation was performed on 5 dpf embryos unless otherwise indicated. Zebrafish

embryos were anesthetized with tricaine. Embryos were cut posterior to the notochord using a

sterile scalpel. Embryos were then recovered to fresh E3 and kept at 28ºC.

2.6 Imaging

Live zebrafish embryos were anesthetized using tricaine, mounted on 3% methylcellulose

(Sigma, M0512), and imaged using a Leica M205FA. ImageJ software was used to quantify

the fluorescent pixel count within 100 µm of the wound site.

Additional high resolution and time-lapse microscopy was carried out on anesthetized

embryos embedded in 1% low melt agarose in a 96 well-plate with a Leica SP8 confocal

microscope or Deltavision Elite microscope.

2.7 Neutrophil tracking

Time-lapse images were processed and analyzed using ImageJ. Neutrophils were tracked

using the Trackmate plugin in ImageJ software and further quantified using Chemokine and

Migration tool software (Ibidi).

2.8 Germ-free derivation and microdissection of embryos

Germ-free zebrafish were created and maintained as previously described [23]. The gut and

body of 5 dpf embryos were separated using a 25-gauge needle and added to Trizol LS

(Invitrogen; 10296010) for RNA extraction.

2.9 RNA extraction, cDNA synthesis and quantitative PCR (qPCR)

10 - 20 zebrafish embryos were pooled and lysed using a 25-gauge needle in Trizol LS for

RNA extraction. cDNA was synthesized using a High-capacity reverse transcription kit

(ThermoFisher Scientific, 4368814). qPCR was carried out using Power UP SYBR green

master mix (ThermoFisher Scientific, 4385610) on a CFX96 Real-Time system (BioRad).

Primer pairs (5`- 3`); *18s* TCGCTAGTTGGCATCGTTTATG and

CGGAGGTTCGAAGACGATCA; *hcar1* CATCGTCATCTACTGCTCCAC and

GCTAACACAAACCGCACA.

2.10 gRNA synthesis and CRISPR injections

gRNA templates for *hcar1-2* (5`- 3`): Target 1

TAATACGACTCACTATAGGTACCGGCGGCTCGATTGGGTTTTAGAGCTAGAAATA

GC, Target 2

TAATACGACTCACTATAGGAGCAACTCTCGCTTCACTGTTTTAGAGCTAGAAATA

GC , Target 3

TAATACGACTCACTATAGGGATTCGAGAGATGTTACTGTTTTAGAGCTAGAAATA

GC. gRNA was synthesized as previously described [24].

A 1:1 solution of gRNA and 500 µg/mL of Cas9 nuclease V3 (Integrated DNA Technology)

was prepared with phenol red dye (Sigma, P0290). Freshly laid eggs were collected from

breeding tanks and the solution was injected in the yolk sac of the egg before the emergence

of the first cell with a FemtoJet 4i (Eppendorf).

2.11 Statistics

All statistical analyses (t-tests and ANOVA where appropriate) were performed using

GraphPad Prism8. Outliers were removed using ROUT, with Q = 1%. All data are

representative of at least 2 biological replicates.

Results

3.1 Zebrafish gut commensals are capable of producing SCFA **ex vivo**

We initially tried to detect SCFAs in whole intestines and their contents dissected from adult

zebrafish using gas chromatography but levels of acetate, butyrate, or propionate were below

our limit of detection.

We next sought to determine if the conventional zebrafish gut microbiota had the capacity to

produce SCFAs using *ex vivo* culture on two rich medias, BHI and Gifu. We found that

microbial communities cultured from conventionally-reared adult zebrafish intestines were

able to synthesize acetate under both aerobic and anaerobic conditions (**Figure 1**). Butyrate

and propionate were only detected under anaerobic conditions. The highest concentrations of

SCFA were detected under anaerobic conditions in BHI where acetate, propionate, and

butyrate were present in a roughly 90:5:5 ratio.

3.2 Butyrate reduces the recruitment of zebrafish neutrophils to a wound

We first observed the effect of SCFAs (acetate, butyrate and propionate) on neutrophil

migration following a tail wound injury using $Tg(lyzC:DsRed)^{nz50}$ and $Tg(lyzC:GFP)^{nz117}$

transgenic zebrafish lines where neutrophils are fluorescently labelled (**Figure 2A**). We

observed a significant reduction in the number of recruited neutrophils at 6 hours post

wounding (hpw) in embryos exposed to butyrate by immersion **(Figure 2B)**. There were no

changes seen with acetate or propionate, but dexamethasone, a corticosteroid anti-

inflammatory used as a positive control, reduced neutrophil recruitment as expected.

We next assessed the quality of neutrophil recruitment by intravital imaging. We observed

reduced neutrophil velocity (**Figure 2C**), and increased meandering index (total distance

traveled / Euclidean distance) in butyrate-treated embryos (**Figure 2D**).

We next sought to determine if butyrate sensitivity is dependent on intestinal maturity by

repeating the tail wound experiment using 2 dpf embryos, a developmental stage prior to

significant intestinal morphogenesis [25], and found neutrophil recruitment was overall reduced

compared to 5 dpf, but still further inhibited by butyrate immersion (**Figure 2E**).

3.3 Butyrate reduces the proportion of TNF positive macrophages at the wound site

Next, we examined the effect of SCFAs on macrophage recruitment and polarization

following wounding. *Tg(mfap4:tdTomato)^{xt12}* transgenic zebrafish were used to visualize

macrophage numbers [26]. Relative to control embryos, macrophage recruitment was reduced

by butyrate and increased by propionate treatment at 6 hpw (**Figure 3A**). Consistent with a

lack of effect on neutrophil recruitment, acetate treatment did not affect the number of

recruited macrophages, and as the positive control anti-inflammatory dexamethasone

significantly reduced macrophage recruitment. These changes were maintained at 24 hpw

when inflammation is in the resolution phase (**Figure 3B**).

We next used the *TgBAC(tnfa:gfp)^{pd1026}* line to monitor inflammatory gene expression [27]. As

expected, since macrophages are the primary producers of TNF in zebrafish embryos, total

TNF promotor fluorescence area at 24 hpw reflected the trend of macrophages recruitment in

each treatment condition with butyrate, but not acetate or propionate, reducing TNF promoter

expression　(**Figure**　**3C**).　We　next　crossed　the　*TgBAC(tnfa:gfp)^{pd1026}*　and

Tg(mfap4:tdTomato)^{xt12} lines to monitor macrophage inflammatory polarization as defined by

inflammatory TNF promotor expression (**Figure 3D**) [28]. Butyrate treatment reduced the

percentage of *TgBAC(tnfa:gfp)^{pd1026}* positive macrophages (**Figure 3E**).

3.4 Butyrate does not have toxic effects as measured by hemostatic indices

Butyrate has been previously shown to reduce the proliferation of zebrafish intestinal

epithelial cells [9]. This raises the possibility that inhibition of leukocyte recruitment by

butyrate immersion was due to toxicity. Changes to zebrafish hemostasis have been observed

in models of toxicity and inflammation [29, 30].

We used *Tg(fabp10a:fgb-EGFP)^(mi4001)*, where fibrin clots are visualized by GFP deposition,

and *Tg(-6.0itga2b:eGFP)^(la2)*, where thrombocytes are GFP-labeled, transgenic zebrafish lines

to monitor hemostasis following transection of the dorsal aorta and posterior cardinal vein [31,

32]. We stabilized clots with aminocaproic acid as a positive control [33]. Fibrinogen

accumulation in *Tg(fabp10a:fgb-EGFP)^(mi4001)* embryos was unchanged at the wound site in

response to butyrate treatment, however we noted that propionate treatment caused increased

fibrinogen accumulation **(Supplementary Figure 1A)**. No changes were observed in

thrombocytes accumulation in the *Tg(-6.0itga2b:eGFP)^(la2)* line following any of the SCFA

treatments **(Supplementary Figure 1B)**.

3.5 Characterization of the zebrafish hydrocarboxylic acid receptor 1

Hydrocarboxylic acid receptor 1 (*HCAR1*) is an important receptor for butyrate in mammals

[19]. We identified a conserved region of zebrafish chromosome 10, human chromosome 12,

and mouse chromosome 5 containing *density regulated re-initiation and release factor (denr)*,

coiled-coil domain-containing 62 (*ccdc62*), *huntingtin interacting protein 1 related a (hip1ra)*

loci and the single exon *hcar* family **(Figure 4A)**. Two copies of the putative zebrafish *hcar1*

(with 93% identity and lacking sufficient divergence to differentiate by PCR) were identified

as annotated by the single-entry NM_001163295.1 (Danio rerio hydroxycarboxylic acid

receptor 1-2 (hcar1-2), mRNA) suggesting the possibility of a tandem duplication event

(Supplementary File). Kuei *et al*. previously annotated what we refer as *hcar1a* as *gpr81-2*

and *hcar1b* as *gpr81-1* [34]. The predicted 933/936 bp transcript of *hcar1a/hcar1b* is expected

to give rise to a 310/311 amino acid protein with 89% amino acid identity. The

Hcar1a/Hcar1b hypothetical proteins contain 7 predicted transmembrane helix domains

characteristic of a GPCR and an E-value of $7.05^{-157}/3.21^{-166}$ for the HCAR subfamily [35]. The

predicted zebrafish proteins have approximately 43% identity to the human HCAR1, HCAR2,

and HCAR3; and mouse HCAR1 and HCAR2 proteins.

Mammalian *HCAR1* is highly expressed by intestinal epithelial cells and neutrophils [19].

Compared to other immune cells, *hcar1* expression is enriched in zebrafish neutrophils [36, 37].

We sought to characterize expression of *hcar1* in the intestines of zebrafish embryos using

microdissection of 5 dpf embryos. We found increased *hcar1* expression in dissected guts

compared to the rest of the embryo by RT-qPCR analysis (**Figure 4B**). Interestingly, there

were no significant changes observed with the absence of microbial colonization in germ-free

embryos (**Supplementary Figure 2**).

3.6 The anti-inflammatory effects of butyrate on neutrophils, but not macrophages, are

dependent on Hcar1.

To determine if butyrate acts through the Hcar1 receptor, we next used CRISPR/Cas9

technology to knockdown *hcar1* expression in zebrafish embryos. We utilized three target

sites in *hcar1a*, two of which had strong homology to sequences in *hcar1b* (Supplementary

File). We confirmed ~50% transcript depletion by RT-qPCR for a shared *hcar1a/hcar1b*

sequence (**Figure 4C**). Embryo development was morphologically normal compared to

embryos injected with control scrambled guide RNA-Cas9 complex (**Figure 4D**).

We performed tail wound injury on *hcar1* knockdown embryos. Knockdown of *hcar1*

abrogated the effect of butyrate treatment on neutrophil recruitment (**Figure 4E**). As expected

from the lack of expression on macrophages, *hcar1* knockdown did not affect the reduced

macrophage recruitment **(Figure 4F)** and TNF promotor expression **(Figure 4G)** induced by

butyrate immersion.

Discussion

This study shows for the first time that commensal microbiota residing in the zebrafish

intestine are capable of producing SCFAs. Experimentally, we demonstrate that the effects,

and sensing, of butyrate are conserved between zebrafish and mammals. Out of the three main

SCFAs, only the anti-inflammatory effect of butyrate was found to be conserved in zebrafish

embryos. We applied the commonly used tail wounding model to demonstrate an anti-

inflammatory effect of butyrate on zebrafish neutrophils and macrophages. Using Crispr-Cas9

targeted mutagenesis, we also identified conserved butyrate responsiveness of the zebrafish

Hcar1 receptor.

The anti-inflammatory effects of butyrate have been established in numerous *in vivo* and *in*

vitro studies of mammalian hosts but not in fish [38]. Our zebrafish tail wound model

demonstrates conservation of this property in a bony fish. Bony fish diverge from mammals

approximate 420 million years ago suggesting that the sensing of microbially-derived SCFAs

has been conserved from a common ancestor.

Our finding that immune cells are responsive to butyrate even before intestinal lumen

formation in early embryonic development is surprising as 2 dpf embryos are usually

contained within relatively impervious chorions that prevent microbial colonization of the

embryo. This suggests that the ability to sense xenobiotic SCFAs is programmed alongside the ability to sense more traditional microbially-associated molecular patterns via pattern recognition molecules.

The HCAR1/GPR81 butyrate receptor is expressed by many mammalian innate immune cells [19]. Expression in zebrafish is strongest in granulocytes [36, 37]. Our knockdown experiments further demonstrate Hcar1 is necessary for the butyrate sensitivity of neutrophils but not macrophages. Thus, the butyrate-Hcar1-neutrophil behavior axis is evolutionarily ancient.

Human macrophages in the presence of butyrate have been shown to differentiate into an M2 phenotype which exhibits anti-microbial and tissue reparative properties [39, 40]. These effects are independent of HCAR1 signal transduction and are believed to be an effect of butyrate acting as a histone deacetylase inhibitor [41, 42]. SCFAs can permeate cell membranes through passive diffusion or through specific transporters such as the proton-coupled monocarboxylate-transporter 1 (MCT1) and sodium-coupled monocarboxylate-transporter 1 (SMCT1) [43-46]. Consistent with this literature, we show butyrate reduces the expression of pro-inflammatory TNF by zebrafish macrophages independent of Hcar1 expression. Our data suggests the HCAR1-independent immunosuppressive actions of butyrate may be conserved across vertebrate evolution.

Our data demonstrate that, under nutrient-rich *in vitro* conditions, gut commensal microbiota from adult zebrafish are capable of synthesizing the three most important SCFAs: acetate, propionate, and butyrate. However, the ratio of acetate, propionate, and butyrate produced under anaerobic conditions in BHI media (90:5:5) differed from the ratio typically observed in mammalian colons (60:20:20) [1]. This may be due to the differing bacterial communities

found in zebrafish and mammalian intestines. The most abundant bacterial phyla in the adult

zebrafish intestine are Proteobacteria and Fusobacteria, whereas mouse and human intestines

are dominated by members of phyla Bacteroidetes and Firmicutes [47]. Considering the SCFA

production we observed *in vitro*, our inability to detect SCFA *in vivo* was surprising. We

anticipate this could be due to rapid host or microbial metabolism of SCFA produced within

the zebrafish gut, or the composition of the diet fed to the zebrafish tested in this experiment.

Zebrafish are omnivores and were fed protein rich diets in the Duke aquaculture facility. A

diet with more SCFA substrates such as carbohydrates and fiber may yield detectable SCFA

production *in situ*.

Interestingly, we observed increased macrophage and fibrinogen clot accumulation at the

wound site following propionate treatment, indicative of a pro-inflammatory effect. Although

this is at odds with anti-inflammatory effects of propionate in mammals [48-50], it is consistent

with evidence of an immunostimulatory effect of propionate in teleosts [51, 52].

Overall this manuscript provides further evidence of conserved mechanisms of host-microbe

interaction within vertebrates. We present evidence that immunological sensitivity to butyrate

is conserved across vertebrates. Furthermore, there is conservation of the molecular

machinery that senses butyrate even down to the responsiveness of individual leukocyte

lineages.

Acknowledgements

Funding: Australian National Health and Medical Research Council Project Grant

APP1099912; The University of Sydney Fellowship G197581; NSW Ministry of Health

under the NSW Health Early-Mid Career Fellowships Scheme H18/31086 to SHO.

The authors acknowledge the facilities and the scientific and technical assistance of the

BioImaging Facility and Sydney Cytometry at Centenary Institute.

The authors declare no conflicts of interest.

References

1.   den Besten G, van Eunen K, Groen AK, Venema K, Reijngoud DJ, Bakker BM. The
role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy
metabolism. J Lipid Res 2013; 54:2325-40.
2.   Tan J, McKenzie C, Potamitis M, Thorburn AN, Mackay CR, Macia L. The role of
short-chain fatty acids in health and disease. Adv Immunol 2014; 121:91-119.
3.   Tedelind S, Westberg F, Kjerrulf M, Vidal A. Anti-inflammatory properties of the
short-chain fatty acids acetate and propionate: a study with relevance to inflammatory bowel
disease. World J Gastroenterol 2007; 13:2826-32.
4.   Zheng N, Gao Y, Zhu W, Meng D, Walker WA. Short chain fatty acids produced by
colonizing intestinal commensal bacterial interaction with expressed breast milk are anti-
inflammatory in human immature enterocytes. PLoS One 2020; 15:e0229283.
5.   Baxter NT, Schmidt AW, Venkataraman A, Kim KS, Waldron C, Schmidt TM.
Dynamics of Human Gut Microbiota and Short-Chain Fatty Acids in Response to Dietary
Interventions with Three Fermentable Fibers. mBio 2019; 10.
6.   Matty MA, Oehlers SH, Tobin DM. Live Imaging of Host-Pathogen Interactions in
Zebrafish Larvae. Methods Mol Biol 2016; 1451:207-23.
7.   Oehlers SH, Flores MV, Chen T, Hall CJ, Crosier KE, Crosier PS. Topographical
distribution of antimicrobial genes in the zebrafish intestine. Dev Comp Immunol 2011;
35:385-91.
8.   Oehlers SH, Flores MV, Hall CJ, Swift S, Crosier KE, Crosier PS. The inflammatory
bowel disease (IBD) susceptibility genes NOD1 and NOD2 have conserved anti-bacterial
roles in zebrafish. Dis Model Mech 2011; 4:832-41.
9.   Kaiko GE, Ryu SH, Koues OI, Collins PL, Solnica-Krezel L, Pearce EJ, et al. The
Colonic Crypt Protects Stem Cells from Microbiota-Derived Metabolites. Cell 2016;
165:1708-20.
10.   Romano N, Simon W, Ebrahimi M, Fadel AHI, Chong CM, Kamarudin MS. Dietary
sodium citrate improved oxidative stability in red hybrid tilapia (Oreochromis sp.) but
reduced growth, health status, intestinal short chain fatty acids and induced liver damage.
Aquaculture 2016; 458:170-6.
11.   Titus E, Ahearn GA. Short-chain fatty acid transport in the intestine of a herbivorous
teleost. J Exp Biol 1988; 135:77-94.
12.   Mountfort DO, Campbell J, Clements KD. Hindgut fermentation in three species of
marine herbivorous fish. Appl Environ Microbiol 2002; 68:1374-80.
13.   Hao YT, Wu SG, Jakovlić I, Zou H, Li WX, Wang GT. Impacts of diet on hindgut
microbiota and short-chain fatty acids in grass carp (Ctenopharyngodon idellus). Aquaculture
Research 2017; 48:5595-605.
14.   Ebrahimi M, Daeman NH, Chong CM, Karami A, Kumar V, Hoseinifar SH, et al.
Comparing the effects of different dietary organic acids on the growth, intestinal short-chain
fatty acids, and liver histopathology of red hybrid tilapia (Oreochromis sp.) and potential use
of these as preservatives. Fish Physiol Biochem 2017; 43:1195-207.

15.   Oehlers SH, Flores MV, Hall CJ, Wang L, Ko DC, Crosier KE, et al. A whole animal
chemical screen approach to identify modifiers of intestinal neutrophilic inflammation. FEBS
J 2017; 284:402-13.
16.   Oehlers SH, Flores MV, Hall CJ, Crosier KE, Crosier PS. Retinoic acid suppresses
intestinal mucus production and exacerbates experimental enterocolitis. Dis Model Mech
2012; 5:457-67.
17.   Koch BEV, Yang S, Lamers G, Stougaard J, Spaink HP. Intestinal microbiome adjusts
the innate immune setpoint during colonization through negative regulation of MyD88. Nat
Commun 2018; 9:4099.
18.   Cheesman SE, Neal JT, Mittge E, Seredick BM, Guillemin K. Epithelial cell
proliferation in the developing zebrafish intestine is regulated by the Wnt pathway and
microbial signaling via Myd88. Proc Natl Acad Sci U S A 2011; 108 Suppl 1:4570-7.
19.   Ranganathan P, Shanmugam A, Swafford D, Suryawanshi A, Bhattacharjee P,
Hussein MS, et al. GPR81, a Cell-Surface Receptor for Lactate, Regulates Intestinal
Homeostasis and Protects Mice from Experimental Colitis. J Immunol 2018; 200:1781-9.
20.   Arpaia N, Campbell C, Fan X, Dikiy S, van der Veeken J, deRoos P, et al. Metabolites
produced by commensal bacteria promote peripheral regulatory T-cell generation. Nature
2013; 504:451-5.
21.   Furusawa Y, Obata Y, Fukuda S, Endo TA, Nakato G, Takahashi D, et al. Commensal
microbe-derived butyrate induces the differentiation of colonic regulatory T cells. Nature
2013; 504:446-50.
22.   Murdoch CC, Espenschied ST, Matty MA, Mueller O, Tobin DM, Rawls JF. Intestinal
Serum amyloid A suppresses systemic neutrophil activation and bactericidal activity in
response to microbiota colonization. PLoS Pathog 2019; 15:e1007381.
23.   Melancon E, Gomez De La Torre Canny S, Sichel S, Kelly M, Wiles TJ, Rawls JF, et
al. Best practices for germ-free derivation and gnotobiotic zebrafish husbandry. Methods Cell
Biol 2017; 138:61-100.
24.   Wu RS, Lam, II, Clay H, Duong DN, Deo RC, Coughlin SR. A Rapid Method for
Directed Gene Knockout for Screening in G0 Zebrafish. Dev Cell 2018; 46:112-25 e4.
25.   Ng AN, de Jong-Curtain TA, Mawdsley DJ, White SJ, Shin J, Appel B, et al.
Formation of the digestive system in zebrafish: III. Intestinal epithelium morphogenesis. Dev
Biol 2005; 286:114-35.
26.   Walton EM, Cronan MR, Beerman RW, Tobin DM. The Macrophage-Specific
Promoter mfap4 Allows Live, Long-Term Analysis of Macrophage Behavior during
Mycobacterial Infection in Zebrafish. PLoS ONE 2015; 10:e0138949.
27.   Marjoram L, Alvers A, Deerhake ME, Bagwell J, Mankiewicz J, Cocchiaro JL, et al.
Epigenetic control of intestinal barrier function and inflammation in zebrafish. Proc Natl
Acad Sci U S A 2015; 112:2770-5.
28.   Nguyen-Chi M, Laplace-Builhe B, Travnickova J, Luz-Crawford P, Tejedor G, Phan
QT, et al. Identification of polarized macrophage subsets in zebrafish. Elife 2015; 4:e07288.
29.   McLeish JA, Chico TJ, Taylor HB, Tucker C, Donaldson K, Brown SB. Skin
exposure to micro- and nano-particles can cause haemostasis in zebrafish larvae. Thromb
Haemost 2010; 103:797-807.
30.   Hortle E, Johnson KE, Johansen MD, Nguyen T, Shavit JA, Britton WJ, et al.
Thrombocyte Inhibition Restores Protective Immunity to Mycobacterial Infection in
Zebrafish. J Infect Dis 2019; 220:524-34.
31.   Vo AH, Swaroop A, Liu Y, Norris ZG, Shavit JA. Loss of fibrinogen in zebrafish
results in symptoms consistent with human hypofibrinogenemia. PLoS ONE 2013; 8:e74682.
32.   Lin HF, Traver D, Zhu H, Dooley K, Paw BH, Zon LI, et al. Analysis of thrombocyte
development in CD41-GFP transgenic zebrafish. Blood 2005; 106:3803-10.

33. Liu Y, Kretz CA, Maeder ML, Richter CE, Tsao P, Vo AH, et al. Targeted
mutagenesis of zebrafish antithrombin III triggers disseminated intravascular coagulation and
thrombosis, revealing insight into function. Blood 2014; 124:142-50.
34. Kuei C, Yu J, Zhu J, Wu J, Zhang L, Shih A, et al. Study of GPR81, the lactate
receptor, from distant species identifies residues and motifs critical for GPR81 functions. Mol
Pharmacol 2011; 80:848-58.
35. Marchler-Bauer A, Bo Y, Han L, He J, Lanczycki CJ, Lu S, et al. CDD/SPARCLE:
functional classification of proteins via subfamily domain architectures. Nucleic Acids Res
2017; 45:D200-D3.
36. Rougeot J, Torraca V, Zakrzewska A, Kanwal Z, Jansen HJ, Sommer F, et al. RNAseq
Profiling of Leukocyte Populations in Zebrafish Larvae Reveals a cxcl11 Chemokine Gene as
a Marker of Macrophage Polarization During Mycobacterial Infection. Frontiers in
immunology 2019; 10:832.
37. Athanasiadis EI, Botthof JG, Andres H, Ferreira L, Lio P, Cvejic A. Single-cell RNA-
sequencing uncovers transcriptional states and fate decisions in haematopoiesis. Nat Commun
2017; 8:2045.
38. Inan MS, Rasoulpour RJ, Yin L, Hubbard AK, Rosenberg DW, Giardina C. The
luminal short-chain fatty acid butyrate modulates NF-kappaB activity in a human colonic
epithelial cell line. Gastroenterology 2000; 118:724-34.
39. Ji J, Shu D, Zheng M, Wang J, Luo C, Wang Y, et al. Microbial metabolite butyrate
facilitates M2 macrophage polarization and function. Sci Rep 2016; 6:24838.
40. Schulthess J, Pandey S, Capitani M, Rue-Albrecht KC, Arnold I, Franchini F, et al.
The Short Chain Fatty Acid Butyrate Imprints an Antimicrobial Program in Macrophages.
Immunity 2019; 50:432-45 e7.
41. Davie JR. Inhibition of histone deacetylase activity by butyrate. J Nutr 2003;
133:2485S-93S.
42. Chang PV, Hao L, Offermanns S, Medzhitov R. The microbial metabolite butyrate
regulates intestinal macrophage function via histone deacetylase inhibition. Proc Natl Acad
Sci U S A 2014; 111:2247-52.
43. Miyauchi S, Gopal E, Fei YJ, Ganapathy V. Functional identification of SLC5A8, a
tumor suppressor down-regulated in colon cancer, as a Na(+)-coupled transporter for short-
chain fatty acids. J Biol Chem 2004; 279:13293-6.
44. Ritzhaupt A, Wood IS, Ellis A, Hosie KB, Shirazi-Beechey SP. Identification and
characterization of a monocarboxylate transporter (MCT1) in pig and human colon: its
potential to transport L-lactate as well as butyrate. J Physiol 1998; 513 (Pt 3):719-32.
45. Borthakur A, Priyamvada S, Kumar A, Natarajan AA, Gill RK, Alrefai WA, et al. A
novel nutrient sensing mechanism underlies substrate-induced regulation of monocarboxylate
transporter-1. Am J Physiol Gastrointest Liver Physiol 2012; 303:G1126-33.
46. Alarcon P, Manosalva C, Conejeros I, Carretta MD, Munoz-Caro T, Silva LMR, et al.
d(-) Lactic Acid-Induced Adhesion of Bovine Neutrophils onto Endothelial Cells Is
Dependent on Neutrophils Extracellular Traps Formation and CD11b Expression. Front
Immunol 2017; 8:975.
47. Rawls JF, Mahowald MA, Ley RE, Gordon JI. Reciprocal gut microbiota transplants
from zebrafish and mice to germ-free recipients reveal host habitat selection. Cell 2006;
127:423-33.
48. Provost KA, Smith M, Miller-Larsson A, Gudleski GD, Sethi S. Bacterial regulation
of macrophage bacterial recognition receptors in COPD are differentially modified by
budesonide and fluticasone propionate. PLoS One 2019; 14:e0207675.

49.   Ciarlo E, Heinonen T, Herderschee J, Fenwick C, Mombelli M, Le Roy D, et al.
Impact of the microbial derived short chain fatty acid propionate on host susceptibility to
bacterial and fungal infections in vivo. Sci Rep 2016; 6:37944.
50.   Park JW, Kim HY, Kim MG, Jeong S, Yun CH, Han SH. Short-chain Fatty Acids
Inhibit Staphylococcal Lipoprotein-induced Nitric Oxide Production in Murine Macrophages.
Immune Netw 2019; 19:e9.
51.   Hoseinifar SH, Zoheiri F, Caipang CM. Dietary sodium propionate improved
performance, mucosal and humoral immune responses in Caspian white fish (Rutilus frisii
kutum) fry. Fish Shellfish Immunol 2016; 55:523-8.
52.   Safari R, Hoseinifar SH, Kavandi M. Modulation of antioxidant defense and immune
response in zebra fish (Danio rerio) using dietary sodium propionate. Fish Physiol Biochem
2016; 42:1733-9.
504

Figure Legends

Figure 1: In vitro synthesis of SCFA by zebrafish gut microbiota

Concentrations of short-chain fatty acids (SCFA) synthesized by conventional microbiota

harvested from adult zebrafish intestines. SCFA content of nutrient media used to culture

microbes is provided under "Negative". Asterisks indicate measurements within the range of

standards. Error bars are shown as mean ± SE, n = 2. Bars without an asterisk indicate

concentrations that were outside the standard range but were detectable.

Figure 2: Butyrate reduces the recruitment of zebrafish neutrophils to a wound

(A) Cartoon describing the standard cut site transecting the dorsal aorta and cardinal vein of a

5 dpf zebrafish embryo, and the fin cut site used for live imaging studies. (B) Neutrophil

counts at 6 hpw. Each dot represents a single embryo. (C) Velocity of wound-recruited

neutrophils calculated from live imaging studies. Each dot represents the average of 10

neutrophils from a single embryo. (C) Meandering index (Total distance/Euclidean distance)

of wound-recruited neutrophils calculated from live imaging studies. Each dot represents the

average of 10 neutrophils from a single embryo. (D) Neutrophil count at 6 hpw in 2 dpf

zebrafish.

Figure 3: Butyrate reduces macrophage recruitment to the wound site and pro-

inflammatory differentiation.

(A) Macrophage fluorescent area at 6 hpw. (B) Macrophage fluorescent area at 24 hpw. (C)

Total TNF promotor fluorescent area at the wound site after 24 hpw. (D) Representative

images of double transgenic red macrophage, green TNF promoter activity embryos tail

wounds at 24 hpw. Scale bar represents 100 µm. (D) Quantification of wound site TNF

expressing macrophages at 24 hpw.

Figure 4: Characterization of zebrafish hydrocarboxylic acid receptor 1 and

responsiveness to butyrate.

(A) Synteny diagram illustrating *HCAR1* in a conserved region of human chromosome 12,

mouse chromosome 5, and zebrafish chromosome 10. (B) Quantification of *hcar1* expression

in dissected gut and body of zebrafish embryos. Each dot represents a biological replicate of

at least 10 embryos. (C) Quantification of *hcar1* expression in 5 dpf embryos injected with

hcar1-targeting Crispr-Cas9 complexes. Each dot represents a biological replicate of at least

10 embryos. (D) Morphology of the control and crispant embryos. Scale bar represents 100

µm. (E) Quantification of neutrophil area at 6 hpw in control and crispant embryos exposed to

butyrate by immersion. (E) Quantification of macrophage area at 6 hpw in control and

crispant embryos exposed to butyrate at 6 hpw. (F) Total TNF promotor-driven fluorescent

area at the wound site at 24 hpw.

Supplementary Figure 1: Effects of SCFA administration on zebrafish hemostasis.

(A) Clotting at 2 hpw following a tail wound in 5 dpf zebrafish. (B) Thrombosis at 3 hpw

following a tail wound in 5 dpf zebrafish.

Supplementary Figure 2: Expression of *hcar1* in germ-free embryos

Quantification of *hcar1* expression in guts and bodies dissected from conventionally raised

and germ-free embryos.