Convergent Antibody Responses to SARS-CoV-2 Infection in Convalescent Individuals

Davide F. Robbiani[1][*][#], Christian Gaebler[1][*], Frauke Muecksch[2][*], Julio C. C. Lorenzi[1][*], Zijun Wang[1][*], Alice Cho[1][*], Marianna Agudelo[1][*], Christopher O. Barnes[6][*], Anna Gazumyan[1][*], Shlomo Finkin[1][*], Thomas Hagglof[1][*], Thiago Y. Oliveira[1][*], Charlotte Viant[1][*], Arlene Hurley[4], Hans-Heinrich Hoffmann[3], Katrina G. Millard[1], Rhonda G. Kost[5], Melissa Cipolla[1], Kristie Gordon[1], Filippo Bianchini[1], Spencer T. Chen[1], Victor Ramos[1], Roshni Patel[1], Juan Dizon[1], Irina Shimeliovich[1], Pilar Mendoza[1], Harald Hartweger[1], Lilian Nogueira[1], Maggi Pack[1], Jill Horowitz[1], Fabian Schmidt[2], Yiska Weisblum[2], Eleftherios Michailidis[3], Alison W. Ashbrook[3], Eric Waltari[7], John E. Pak[7], Kathryn E. Huey-Tubman[6], Nicholas Koranda[6], Pauline R. Hoffman[6], Anthony P. West, Jr.[6], Charles M. Rice[3], Theodora Hatziioannou[2], Pamela J. Bjorkman[6], Paul D. Bieniasz[2,8,#], Marina Caskey[1,#], Michel C. Nussenzweig[1,8,#]

[1]Laboratory of Molecular Immunology, The Rockefeller University, New York, NY 10065, USA
[2]Laboratory of Retrovirology, The Rockefeller University, New York, NY 10065, USA
[3]Laboratory of Virology and Infectious Disease, The Rockefeller University, New York, NY 10065, USA
[4]Hospital Program Direction, The Rockefeller University, New York, NY 10065, USA
[5]Hospital Clinical Research Office, The Rockefeller University, New York, NY 10065, USA
[6]Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA 91125, USA
[7]Chan Zuckerberg Biohub, 499 Illinois Street, San Francisco, CA 94158, USA
[8]Howard Hughes Medical Institute
[*]Equal contribution
[#]Send correspondence to Paul D. Bieniasz: pbieniasz@rockefeller.edu, Marina Caskey: mcaskey@rockefeller.edu, Michel C. Nussenzweig: nussen@rockefeller.edu, or Davide F. Robbiani: drobbiani@rockefeller.edu

Abstract

During the COVID-19 pandemic, SARS-CoV-2 infected millions of people and claimed hundreds of thousands of lives. Virus entry into cells depends on the receptor binding domain (RBD) of the SARS-CoV-2 spike protein (S). Although there is no vaccine, it is likely that antibodies will be essential for protection. However, little is known about the human antibody response to SARS-CoV-2[1-5]. Here we report on 149 COVID-19 convalescent individuals. Plasmas collected an average of 39 days after the onset of symptoms had variable half-maximal neutralizing titers ranging from undetectable in 33% to below 1:1000 in 79%, while only 1% showed titers >1:5000. Antibody cloning revealed expanded clones of RBD-specific memory B cells expressing closely related antibodies in different individuals. Despite low plasma titers, antibodies to three distinct epitopes on RBD neutralized at half-maximal inhibitory concentrations (IC_{50}s) as low as single digit ng/mL. Thus, most convalescent plasmas obtained from individuals who recover from COVID-19 do not contain high levels of neutralizing activity. Nevertheless, rare but recurring RBD-specific antibodies with potent antiviral activity were found in all individuals tested, suggesting that a vaccine designed to elicit such antibodies could be broadly effective.

Between April 1 and May 8, 2020, 157 eligible participants enrolled in the study. Of these, 111

(70.7%) were individuals diagnosed with SARS-CoV-2 infection by RT-PCR (cases), and 46

(29.3%) were close contacts of individuals diagnosed with SARS-CoV-2 infection (contacts).

While inclusion criteria allowed for enrollment of asymptomatic participants, 8 contacts that did

not develop symptoms were excluded from further analyses. The 149 cases and contacts were free

of symptoms suggestive of COVID-19 for at least 14 days at the time of sample collection.

Participant demographics and clinical characteristics are shown in Table 1 and Extended Data

Tables 1 and 2. Only one individual who tested positive for SARS-CoV-2 infection by RT-PCR

remained asymptomatic. The other 148 participants reported symptoms suggestive of COVID-19

with an average onset of approximately 39 days (range 17 to 67 days) before sample collection. In

this cohort, symptoms lasted for an average of 12 days (0-35 days), and 11 (7%) of the participants

were hospitalized. The most common symptoms were fever (83.9%), fatigue (71.1%), cough

(62.4%) and myalgia (61.7%) while baseline comorbidities were infrequent (10.7%) (Table 1 and

Extended Data Tables 1 and 2). There were no significant differences in duration or severity (see

Methods) of symptoms, or in time from onset of symptoms to sample collection between genders

or between cases and contacts. There was no age difference between females and males in our

cohort (Extended Data Fig. 1).

Plasma samples were tested for binding to the SARS-CoV-2 RBD and trimeric spike (S) proteins

by ELISA using anti-IgG or -IgM secondary antibodies for detection (Fig. 1, Extended Data Table

1 and Extended Data Figs. 2 and 3). Eight independent negative controls and the plasma sample

from participant 21 (COV21) were included for normalization of the area under the curve (AUC).

Overall, 78% and 70% of the plasma samples tested showed anti-RBD and anti-S IgG AUCs that

were at least 2 standard deviations above the control (Fig. 1 a, b). In contrast, only 15% and 34%

of the plasma samples showed IgM responses to anti-RBD and anti-S that were at least 2 standard

deviations above control, respectively (Fig. 1 c, d). There was no positive correlation between anti-

RBD or -S IgG or IgM levels and duration of symptoms or the timing of sample collection relative

to onset of symptoms (Fig. 1e, and Extended Data Figs. 3 a-c and 3 g-j). On the contrary, as might

be expected, anti-RBD IgM titers were negatively correlated with duration of symptoms and the

timing of sample collection (Fig. 1e and Extended Data Fig. 3h). Anti-RBD IgG levels were

modestly correlated to age, and the severity of symptoms including hospitalization (Fig. 1 f, g and

Extended Data Fig. 3k). Interestingly, females had lower anti-RBD and -S IgG titers than males

(Fig. 1h, Extended Data Fig. 2f).

To measure the neutralizing activity in convalescent plasmas we used HIV-1-based virions

carrying a nanoluc luciferase reporter that were pseudotyped with the SARS-CoV-2 spike (SARS-

CoV-2 pseudovirus, see Methods, Fig. 2 and Extended Data Fig. 4). The overall level of

neutralizing activity in the cohort, as measured by the half-maximal neutralizing titer (NT_{50}) was

generally low, with 33% undetectable and 79% below 1,000 (Fig. 2 a, b). The geometric mean

NT_{50} was 121 (arithmetic mean = 714), and only 2 individuals reached NT_{50}s above 5,000 (Fig. 2

a, b and Extended Data Table 1).

Notably, levels of anti-RBD- and -S IgG antibodies correlated strongly with NT_{50} (Fig. 2 c, d).

Neutralizing activity also correlated with age, duration of symptoms and symptom severity

(Extended Data Fig. 5). Consistent with this observation, hospitalized individuals with longer

symptom duration showed slightly higher average levels of neutralizing activity than non-

hospitalized individuals (p=0.0495, Fig. 2e). Finally, we observed a significant difference in

neutralizing activity between males and females (p=0.0031, Fig. 2f). The difference between males

and females was consistent with higher anti-RBD and -S IgG titers in males, and could not be

attributed to age, severity, timing of sample collection relative to onset of symptoms or duration

of symptoms (Fig. 1h, Extended Data Fig. 1 a-d and 2f).

To determine the nature of the antibodies elicited by SARS-CoV-2 infection we used flow

cytometry to isolate individual B lymphocytes with receptors that bound to RBD from the blood

of 6 selected individuals including the 2 top and 4 high to intermediate neutralizers (Fig. 3). The

frequency of antigen-specific B cells, identified by their ability to bind to both Phycoerythrin (PE)-

and AF647-labeled RBD, ranged from 0.07 to 0.005% of all circulating B cells in COVID-19

convalescents but they were undetectable in pre-COVID-19 controls (Fig. 3a and Extended Data

Fig. 6). We obtained 534 paired IgG heavy and light chain (IGH and IGL) sequences by reverse

transcription and subsequent PCR from individual RBD-binding B cells from the 6 convalescent

individuals (see Methods and Extended Data Table 3). When compared to the human antibody

repertoire, several IGHV and IGLV genes were significantly over-represented (Extended Data Fig.

7). The average number of V genes nucleotide mutations for IGH and IGL was 4.2 and 2.8,

respectively (Extended Data Fig. 8), which is lower than in antibodies cloned from individuals

suffering from chronic infections such as Hepatitis B or HIV-1, and similar to antibodies derived

from primary malaria infection or non-antigen-enriched circulating IgG memory cells[6-8] (Wang et

al, in press, https://www.biorxiv.org/content/10.1101/2020.03.04.976159v1.full). Among other

antibody features, IGH CDR3 length was indistinguishable from the reported norm and

hydrophobicity was below average (Extended Data Fig. 8)[9].

As is the case with other human pathogens, there were expanded clones of viral antigen binding B

cells in all COVID-19 individuals tested (see Methods and Fig. 3 b,c). Overall, 32.2% of the

recovered IGH and IGL sequences were from clonally expanded B cells (range 21.8-57.4% across

individuals, Fig. 3b). Antibodies that shared specific combinations of IGHV and IGLV genes in

different individuals comprised 14% of all the clonal sequences (colored pie slices in Fig. 3 b,c).

Remarkably, the amino acid sequences of some antibodies found in different individuals were

nearly identical (Fig. 3 e,d). For example, antibodies expressed by clonally expanded B cells with

IGHV1-58/IGKV3-20 and IGHV3-30-3/IGKV1-39 found repeatedly in different individuals had

amino acid sequence identities of up to 99% and 92%, respectively (Fig. 3d and Extended Data

Table 4). We conclude that the IgG memory response to the SARS-CoV-2 RBD is rich in recurrent

and clonally expanded antibody sequences.

To examine the binding properties of anti-SARS-CoV-2 antibodies, we expressed 84

representative antibodies, 56 from clones and 28 from singlets (Extended Data Table 5). ELISA

assays showed that 94% (79 out of 84) of the antibodies tested including clonal and unique

sequences bound to the SARS-CoV-2 RBD with an average half-maximal effective concentration

(EC_{50}) of 6.1 ng/mL (Fig. 4a and Extended Data Fig. 9a). A fraction of those (7 out of 79, or 9%)

cross-reacted with the RBD of SARS-CoV with a mean EC_{50} of 120.1 ng/mL (Extended Data Fig.

9b and c). No significant cross-reactivity was noted to the RBDs of MERS, HCoV-OC43, HCoV-

229E or HCoV-NL63.

To determine whether the monoclonal antibodies have neutralizing activity, we tested them against

the SARS-CoV-2 pseudovirus (Fig. 4 and Extended Data Table 6). Among 79 RBD binding

antibodies tested, we found 40 that neutralized SARS-CoV-2 pseudovirus with nanogram per

milliliter half-maximal inhibitory concentrations (IC_{50}s) ranging from 3 to 709 (Fig. 4 b,c and e,

Extended Data Table 6). A subset of the most potent of these antibodies were also tested against

authentic SARS-CoV-2 and neutralized with IC_{50}s of less than 5 ng/ml (Fig. 4 d,e).

Potent neutralizing antibodies were found in individuals irrespective of their plasma NT_{50}s. For

example, C121, C144, and C135 with IC_{50}s of 1.64, 2.55 and 2.98 ng/mL against authentic SARS-

CoV-2, respectively, were obtained from individuals COV107, COV47, and COV72 whose

plasma NT_{50} values were of 297, 10,433 and 3,138, respectively (Figs. 2b and 4). Finally, clones

of antibodies with shared IGHV and IGLV genes were among the best neutralizers, e.g., antibody

C002 composed of IGHV3-30/IGKV1-39 is shared by the 2 donors with the best plasma

neutralizing activity (red pie slice in Fig. 3b and Fig. 4). We conclude that even individuals with

modest plasma neutralizing activity harbor rare IgG memory B cells that produce potent SARS-

CoV-2 neutralizing antibodies.

To determine whether human anti-SARS-CoV-2 monoclonal antibodies with neutralizing activity

can bind to distinct domains on the RBD, we performed bilayer interferometry experiments in

which a preformed antibody-RBD immune complex was exposed to a second monoclonal. The

antibodies tested comprised 3 groups, all of which differ in their binding properties from CR3022,

an antibody that neutralizes SARS-CoV and binds to, but does not neutralize SARS-CoV-2[10,11].

Representatives of each of the 3 groups include: C144 and C101 in Group 1; C121 and C009 in

Group 2; C135 in Group 3. All of these antibodies can bind after CR3022. Groups 1 and 2 also

bind after Group 3, and Groups 1 and 2 differ in that Group 1 can bind after Group 2 but not vice

versa (Fig. 4 f-n). We conclude that similar to SARS-CoV, there are multiple distinct neutralizing

epitopes on the RBD of SARS-CoV-2.

To further define the binding characteristics of Groups 1 and 2 antibodies, we imaged SARS-CoV-

2 S–Fab complexes by negative stain electron microscopy (nsEM) using C002 (Group 1, an

IGHV3-30/IGKV1-39 antibody, which is clonally expanded in 2 donors), C119 and C121 (both in

Group 2) Fabs (Fig. 4 f-r and Extended Data Fig. 10). Consistent with the conformational

flexibility of the RBD, 2D class averages showed heterogeneity in both occupancy and

conformation of bound Fabs for both groups (Fig. 4o-q). The 3D reconstructions of S-Fab

complexes revealed both a fully-occupied RBD and partial density for a second bound Fab,

suggesting that Fabs from both groups are able to recognize "up" and "down" states of the RBD

as previously described for some of the antibodies targeting this epitope[12,13]. The 3D

reconstructions are also consistent with BLI measurements indicating that Groups 1 and 2

antibodies bind a RBD epitope distinct from antibody CR3022, which is only accessible on the

RBD "up" conformational state[11], and bind the RBD with different angles of approach, with Group

1 antibodies most similar to the approach angle of the SARS-CoV antibody S230 (Fig. 4r)[14].

Human monoclonal antibodies with neutralizing activity against pathogens ranging from viruses

to parasites have been obtained from naturally infected individuals by single cell antibody cloning.

Several have been shown to be effective in protection and therapy in model organisms and in early

phase clinical studies, but only one antiviral monoclonal is currently in clinical use[15]. Antibodies

are relatively expensive and more difficult to produce than small molecule drugs. However, they

differ from drugs in that they can engage the host immune system through their constant domains

that bind to Fc gamma receptors on host immune cells[16]. These interactions can enhance immunity

and help clear the pathogen or infected cells, but they can also lead to disease enhancement during

Dengue[17] and possibly coronavirus infections[18]. This problem has impeded Dengue vaccine

development but would not interfere with the clinical use of potent neutralizing antibodies that can

be modified to prevent Fc gamma receptor interactions and remain protective against viral

pathogens[19].

Antibodies are essential elements of most vaccines and will likely be crucial component of an

effective vaccine against SARS-CoV-2[20](PMID:32434945; PMID:32434946). Recurrent

antibodies have been observed in other infectious diseases and vaccinal responses [21-24](Wang et

al, in press, https://www.biorxiv.org/content/10.1101/2020.03.04.976159v1.full). The observation

that plasma neutralizing activity is low in most convalescent individuals, but that recurrent anti-

SARS-CoV-2 RBD antibodies with potent neutralizing activity can be found in individuals with

unexceptional plasma neutralizing activity suggests that humans are intrinsically capable of

generating anti-RBD antibodies that potently neutralize SARS-CoV-2. Thus, vaccines that

selectively and efficiently induce antibodies targeting the SARS-CoV-2 RBD may be especially

effective.

Table

Table 1. Cohort characteristics

Gender	n	Average age	Case/Contact	Average duration		Average Sx Severity (0-10)	ELISA binding (AUC)				Neutralization (NT50)
				Sx total	Sx onset to visit		RBD		S		
							IgG	IgM	IgG	IgM	
Male	83	45 (19-76)	65/18	12 (0-31)	39 (21-63)	5.8 (0-10)	2.44	1.61	4.65	1.62	867
Female	66	42 (19-75)	46/20	12 (1-35)	38 (17-67)	5.4 (1-9)	1.99	1.58	4.36	1.86	522

Sx=symptoms

 Figures

Figure 1. Plasma antibodies against SARS-CoV-2. a-d, Graphs show results of ELISAs

measuring plasma reactivity to RBD (**a, b**) and S protein (**c, d**). Left shows optical density units at

450 nm (OD, Y axis) and reciprocal plasma dilutions (X axis). Negative controls in black;

individuals 21, and 47 in blue and red lines and arrowheads, respectively. Right shows normalized

area under the curve (AUC) for controls and each of 149 individuals in the cohort. **e**, Symptom

(Sx) onset to time of sample collection in days (X axis) plotted against normalized AUC for IgM

binding to RBD (Y axis) r=0.5517 and p=<0.0001. **f**, Participant age in years (X axis) plotted

against normalized AUC for IgG binding to RBD (Y axis) r=0.1827 and p=0.0258. The r and p

values for the correlations in **e** and **f** were determined by two-tailed Spearman's. **g,** IgG anti-RBD

normalized AUC for outpatients and hospitalized individuals p=0.0178. **h,** IgG anti-RBD

normalized AUC for males and females p=0.0063. For **g** and **h** horizontal bars indicate median

values. Statistical significance was determined using two-tailed Mann-Whitney U test.

Figure 2. Neutralization of SARS-CoV-2 pseudovirus by plasma. **a,** Graph shows normalized

relative luminescence values (RLU, Y axis) in cell lysates of 293T$_{ACE2}$ cells 48 hours after

infection with nanoluc-expressing SARS-CoV-2 pseudovirus in the presence of increasing

concentrations of plasma (X axis) derived from 149 participants (grey, except individuals 47 and

21 in red, and blue lines, bars and arrowheads, respectively) and 3 negative controls (black lines).

Standard deviations of duplicates of one representative experiment are shown. **b,** Ranked average

half-maximal inhibitory plasma neutralizing titer (NT_{50}) for the 59 of 149 individuals with NT_{50}s

>500 and individual 107. See also Extended Data Table 1. Asterisks indicate donors from which

antibody sequences were derived. **c,** AUC for anti-RBD IgG ELISA (X axis) plotted against NT_{50}

(Y axis) r=0.6432, p=<0.0001. **d,** AUC for anti-S IgG ELISA (X axis) plotted against NT_{50} (Y

axis) r=0.6721, p=<0.0001. **e,** NT_{50} for outpatients and hospitalized individuals p=0.0495. **f,** NT_{50}

for all males and females in the cohort p=0.0031. Dotted line in c to f (NT_{50}=5) represents lower

limit of detection (LLOD). Samples with undetectable neutralizing titers were plotted at LLOD.

Correlations in **c** and **d** were determined by two-tailed Spearman's. Statistical significance in **e** and

f was determined using two-tailed Mann-Whitney U test. Horizontal bars indicate median values.

Figure 3. Anti-SARS-CoV-2 RBD antibodies. a. Representative flow cytometry plots showing dual AF647- and PE-RBD binding B cells in control and 6 study individuals (for gating strategy see Extended Data Fig. 6). Percentages of antigen specific B cells are indicated. Control is a healthy control sample obtained pre-COVID-19. **b,** Pie charts depicting the distribution of antibody sequences from 6 individuals. The number in the inner circle indicates the number of sequences analyzed for the individual denoted above the circle. White indicates sequences isolated only once, and grey or colored pie slices are proportional to the number of clonally related sequences. Red, blue, orange and yellow pie slices indicate clones that share the same IGHV and

IGLV genes. **c,** Circos plot shows sequences from all 6 individuals with clonal relationships

depicted as in **b**. Interconnecting lines indicate the relationship between antibodies that share V

and J gene segment sequences at both IGH and IGL. Purple, green and gray lines connect related

clones, clones and singles, and singles to each other, respectively. **d,** Sample sequence alignment

for antibodies originating from different individuals that display highly similar IGH V(D)J and

IGL VJ sequences including CDR3s. Amino acid differences in CDR3s to the bolded reference

sequence above are indicated in red and dots represent identities.

Figure 4. Anti-SARS-CoV-2 RBD antibody reactivity. a, Graph show results of ELISA assays measuring monoclonal antibody reactivity to RBD. Optical density units at 450 nm (OD, Y axis) vs. antibody concentrations (X axis). C121, C135 C144 and isotype control in red, green, purple, and black respectively, in all panels. **b,** Graph shows normalized relative luminescence values (RLU, Y axis) in cell lysates of $293T_{ACE2}$ cells 48 hours after infection with SARS-CoV-2 pseudovirus in the presence of increasing concentrations of monoclonal antibodies (X axis). **c,** RLU for SARS-CoV-2 pseudovirus assay (Y axis) vs. titration of monoclonal antibodies C121, C135 and C144 in one of two independent experiments (see Extended Data Table 6). **d,** SARS-

CoV-2 real virus neutralization assay. Infected cells (Y axis) vs. titration of monoclonal antibodies

C121, C135 and C144 in two independent experiments. For a and b panels, isotype control

antibody in black. **e,** IC50s for antibodies assayed in **b** and **d**. **f,** Diagrammatic representation of

biolayer interferometry experiment. **g,** Graph shows binding of C144, C101, C121, C009, C135,

and CR3022[10,11] to RBD. **h-m**, Secondary antibody binding to preformed IgG-RBD complexes

(Ab1). The table displays the shift in nanometers after second antibody (Ab2) binding to the

antigen in the presence of the first antibody (Ab1). Values are normalized by the subtraction of the

autologous antibody control. **o-q,** Representative 2D-class averages and 3D reconstructed

volumes for SARS-CoV-S 2P trimers complexed with C002, C119, and C121 Fabs. 2D-class

averages with observable Fab density are boxed. **r**, Overlay of S-Fab complexes with fully-

occupied C002 (blue), C121 (magenta) and C119 (orange) Fabs aligned on the RBD "up"

conformational state. The SARS-CoV-2 S model with 1 "up" RBD state (PDB 6VYB) was fit into

the density and the SARS-CoV antibody S230 (PDB 6NB6) shown as reference (green ribbon).

Methods

Study participants. Study participants were recruited at the Rockefeller University Hospital in New York from April 1 through May 8, 2020. Eligible participants were adults aged 18-76 years who were either diagnosed with SARS-CoV-2 infection by RT-PCR and were free of symptoms of COVID-19 for at least 14 days (cases), or who were close contacts (e.g., household, co-workers, members of same religious community) with someone who had been diagnosed with SARS-CoV-2 infection by RT-PCR and were free of symptoms suggestive of COVID-19 for at least 14 days (contacts). Exclusion criteria included presence of symptoms suggestive of active SARS-CoV-2 infection, or hemoglobin < 12 g/dL for males and < 11 g/dL for females.

Most study participants were residents of the Greater New York City tri-state region and were enrolled sequentially according to eligibility criteria. Participants were first interviewed by phone to collect information on their clinical presentation, and subsequently presented to the Rockefeller University Hospital for a single blood sample collection. Participants were asked to rate the highest severity of their symptoms on a numeric rating scale ranging from 0 to 10. The score was adapted from the pain scale chart, where 0 was the lack of symptoms, 4 were distressing symptoms (e.g. fatigue, myalgia, fever, cough, shortness of breath) that interfered with daily living activities, 7 were disabling symptoms that prevented the performance of daily living activities, and 10 was unimaginable/unspeakable discomfort (in this case, distress due to shortness of breath). All participants provided written informed consent before participation in the study and the study was conducted in accordance with Good Clinical Practice.

Blood samples processing and storage. Peripheral Blood Mononuclear Cells (PBMCs) were obtained by gradient centrifugation and stored in liquid nitrogen in the presence of FCS and

DMSO. Heparinized plasma and serum samples were aliquoted and stored at -20°C or less. Prior

to experiments, aliquots of plasma samples were heat-inactivated (56C for 1 hour) and then stored

at 4C.

Cloning, expression and purification of recombinant coronavirus proteins. Codon-optimized

nucleotide sequences encoding the SARS-CoV-2 S ectodomain (residues 16-1206) and receptor

binding domain (RBD; residues 331-524) were synthesized and subcloned into the mammalian

expression pTwist-CMV BetaGlobin vector by Twist Bioscience Technologies based on an early

SARS-CoV-2 sequence isolate (GenBank MN985325.1). The SARS-CoV-2 RBD construct

included an N-terminal human IL-2 signal peptide and dual C-terminal tags ((GGGGS)$_2$-

HHHHHHHH (octa-histidine), and GLNDIFEAQKIEWHE (AviTag)). In addition, the

corresponding S1^B or receptor binding domains for SARS-CoV S (residues 318-510; GenBank

AAP13441.1), MERS-CoV S (residues 367-588; GenBank JX869059.2), HCoV-NL63 (residues

481-614; GenBank AAS58177.1), HCoV-OC43 (residues 324-632; GenBank AAT84362.1), and

HCoV-229E (residues 286-434; GenBank AAK32191.1) were synthesized with the same N- and

C-terminal extensions as the SARS-CoV-2 RBD construct and subcloned into the mammalian

expression pTwist-CMV BetaGlobin vector (Twist Bioscience Technologies). The SARS-CoV-2

S ectodomain was modified as previously described [4]. Briefly, the S ectodomain construct included

an N-terminal mu-phosphatase signal peptide, 2P stabilizing mutations (K986P and V987P),

mutations to remove the S1/S2 furin cleavage site ($_{682}$RRAR$_{685}$ to GSAS), a C-terminal extension

(IKGSG-RENLYFQG (TEV protease site), GGGSG-YIPEAPRDGQAYVRKDGEWVLLSTFL

(foldon trimerization motif), G-HHHHHHHH (octa-histidine tag), and GLNDIFEAQKIEWHE

(AviTag)). The SARS-CoV-2 S 2P ectodomain and RBD constructs were produced by transient

transfection of 500 mL of Expi293 cells (Thermo Fisher) and purified from clarified transfected

cell supernatants four days post-transfection using Ni^{2+}-NTA affinity chromatography (GE Life

Sciences). Affinity-purified proteins were concentrated and further purified by size-exclusion

chromatography (SEC) using a Superdex200 16/60 column (GE Life Sciences) running in 1x TBS

(20 mM Tris-HCl pH 8.0, 150 mM NaCl, and 0.02% NaN_3). Peak fractions were analyzed by SDS-

PAGE, and fractions corresponding to soluble S 2P trimers or monomeric RBD proteins were

pooled and stored at 4˚C.

ELISAs. ELISAs to evaluate antibodies binding to SARS-CoV-2 RBD and trimeric spike proteins

were performed by coating of high binding 96 half well plates (Corning #3690) with 50 µL per

well of a 1µg/mL protein solution in PBS overnight at 4°C. Plates were washed 6 times with

washing buffer (1xPBS with 0.05% Tween 20 (Sigma-Aldrich)) and incubated with 170 µL per

well blocking buffer (1xPBS with 2% BSA and 0.05% Tween20 (Sigma)) for 1 hour at room

temperature (RT). Immediately after blocking, monoclonal antibodies or plasma samples were

added in PBS and incubated for 1 hr at RT. Plasma samples were assayed at a 1:200 starting

dilution and seven additional 3-fold serial dilutions. Monoclonal antibodies were tested at 10µg/ml

starting concentration and 10 additional 4-fold serial dilutions. Plates were washed 6 times with

washing buffer and then incubated with anti-human IgG or IgM secondary antibody conjugated to

horseradish peroxidase (HRP) (Jackson Immuno Research 109-036-088 and 109-035-129) in

blocking buffer at a 1:5000 dilution. Plates were developed by addition of the HRP substrate, TMB

(ThermoFisher) for 10 minutes, then the developing reaction was stopped by adding 50µl 1M

H_2SO_4 and absorbance was measured at 450nm with an ELISA microplate reader (FluoStar

Omega, BMG Labtech). For plasma samples, a positive control (plasma from patient COV21,

diluted 200-fold in PBS) was added in duplicate to every assay plate. The average of its signal was

used for normalization of all the other values on the same plate with Excel software prior to

calculating the area under the curve using Prism 8 (GraphPad). For monoclonal antibodies, the

half-maximal effective concentration (EC_{50}) was determined using 4-parameter nonlinear

regression (GraphPad Prism).

$293T_{ACE2}$ cells. For constitutive expression of ACE2 in 293T cells, a cDNA encoding ACE2,

carrying two inactivating mutations in the catalytic site (H374N & H378N), was inserted into CSIB

3' to the SFFV promoter[25]. $293T_{ACE2}$ cells were generated by transduction with CSIB based virus

followed by selection with 5 μg/ml Blasticidin.

SARS-CoV-2 and SARS-CoV pseudotyped reporter viruses. A plasmid expressing a C-

terminally truncated SARS-CoV-2 S protein (pSARS-CoV2-S_{trunc}) was generated by insertion of

a human-codon optimized cDNA encoding SARS-CoV-2 S lacking the C-terminal 19 codons

(Geneart) into pCR3.1. The S ORF was taken from "Wuhan seafood market pneumonia virus

isolate Wuhan-Hu-1" (NC_045512). For expression of full-length SARS-CoV S protein, "Human

SARS coronavirus Spike glycoprotein Gene ORF cDNA clone expression plasmid (Codon

Optimized)" (here referred to as pSARS-CoV-S) was obtained from SinoBiological (Cat:

VG40150-G-N). An *env*-inactivated HIV-1 reporter construct (pNL4-3ΔEnv-nanoluc) was

generated from pNL4-3[26] by introducing a 940 bp deletion 3' to the *vpu* stop-codon, resulting in a

frameshift in *env*. The human codon-optimized nanoluc Luciferase reporter gene (*Nluc,* Promega)

was inserted in place of nucleotides 1-100 of the *nef*-gene. To generate pseudotyped viral stocks,

293T cells were transfected with pNL4-3ΔEnv-nanoluc and pSARS-CoV2-S_{trunc} or pSARS-CoV-

S using polyethyleneimine. Co-transfection of pNL4-3ΔEnv-nanoluc and S-expression plasmids

leads to production of HIV-1-based virions carrying either the SARS-CoV-2 or SARS-CoV spike

protein on the surface. Eight hours after transfection, cells were washed twice with PBS and fresh

media was added. Supernatants containing virions were harvested 48 hours post transfection,

filtered and stored at -80°C. Infectivity of virions was determined by titration on $293T_{ACE2}$ cells.

Pseudotyped virus neutralization assay. Five-fold serially diluted plasma from COVID-19

convalescent individuals and healthy donors or four-fold serially diluted monoclonal antibodies

were incubated with the SARS-CoV-2 or SARS-CoV pseudotyped virus for 1 hour at 37°C

degrees. The mixture was subsequently incubated with $293T_{ACE2}$ cells for 48 hours after which

cells were washed twice with PBS and lysed with Luciferase Cell Culture Lysis 5x reagent

(Promega). Nanoluc Luciferase activity in lysates was measured using the Nano-Glo Luciferase

Assay System (Promega). Relative luminescence units obtained were normalized to those derived

from cells infected with SARS-CoV-2 or SARS-CoV pseudotyped virus in the absence of plasma

or monoclonal antibodies. The half-maximal inhibitory concentration for plasma (NT_{50}) or

monoclonal antibodies (IC_{50}) was determined using 4-parameter nonlinear regression (GraphPad

Prism).

Cell lines, virus and virus titration. VeroE6 kidney epithelial cells (*Chlorocebus sabaeus*) and

Huh-7.5 hepatoma cells (*H. sapiens*) were cultured in Dulbecco's Modified Eagle Medium

(DMEM) supplemented with 1% nonessential amino acids (NEAA) and 10% fetal bovine serum

(FBS) at 37°C and 5% CO_2. All cell lines have been tested negative for contamination with

mycoplasma and were obtained from the ATCC (with the exception for Huh-7.5).

SARS-CoV-2, strain USA-WA1/2020, was obtained from BEI Resources and amplified in VeroE6

cells at 33°C. Viral titers were measured on Huh-7.5 cells by standard plaque assay (PA). Briefly,

500 μL of serial 10-fold virus dilutions in Opti-MEM were used to infect 400,000 cells seeded the

day prior in a 6-well plate format. After 90 min adsorption, the virus inoculum was removed, and

cells were overlayed with DMEM containing 10% FBS with 1.2% microcrystalline cellulose

(Avicel). Cells were incubated for five days at 33°C, followed by fixation with 3.5% formaldehyde

and crystal violet staining for plaque enumeration. All experiments were performed in a biosafety

level 3 laboratory.

Microscopy-based neutralization assay of authentic SARS-CoV-2. The day prior to infection

VeroE6 cells were seeded at 12,500 cells/well into 96-well plates. Antibodies were serially diluted

in BA-1, mixed with a constant amount of SARS-CoV-2 (grown in VeroE6) and incubated for 60

min at 37°C. The antibody-virus-mix was then directly applied to VeroE6 cells (MOI of ~0.1

PFU/cell). Cells were fixed 18 hours post infection by adding an equal volume of 7% formaldehyde

to the wells, followed by permeabilization with 0.1% Triton X-100 for 10 min. After extensive

washing, cells were incubated for 1 hour at room temperature with blocking solution of 5% goat

serum in PBS (catalog no. 005–000-121; Jackson ImmunoResearch). A rabbit polyclonal anti-

SARS-CoV-2 nucleocapsid antibody (catalog no. GTX135357; GeneTex) was added to the cells

at 1:500 dilution in blocking solution and incubated at 4°C overnight. A goat anti-rabbit

AlexaFluor 594 (catalog no. A-11012; Life Technologies) at a dilution of 1:2,000 was used as a

secondary antibody. Nuclei were stained with Hoechst 33342 (catalog no. 62249; Thermo

Scientific) at a 1:1,000 dilution. Images were acquired with a fluorescence microscope and

analyzed using ImageXpress Micro XLS (Molecular Devices, Sunnyvale, CA). All statistical

analyses were done using Prism 8 software (GraphPad).

Biotinylation of viral protein for use in flow cytometry. Purified and Avi-tagged SARS-CoV-2

RBD was biotinylated using the Biotin-Protein Ligase-BIRA kit according to manufacturer's

instructions (Avidity). Ovalbumin (Sigma, A5503-1G) was biotinylated using the EZ-Link Sulfo-

NHS-LC-Biotinylation kit according to the manufacturer's instructions (Thermo Scientific).

Biotinylated Ovalbumin was conjugated to streptavidin-BV711 (BD biosciences, 563262) and

RBD to streptavidin-PE (BD biosciences, 554061) and streptavidin-Alexa Fluor 647 (AF647,

Biolegend, 405237) respectively[27].

Single cell sorting by flow cytometry. PBMCs were enriched for B cells by negative selection

using a pan B cell isolation kit according to the manufacturer's instructions (Miltenyi Biotec, 130-

101-638). The enriched B cells were incubated in FACS buffer (1 X Phosphate-buffered Saline

(PBS), 2% calf serum, 1 mM EDTA) with the following anti-human antibodies: anti-CD20-PECy7

(BD Biosciences, 335793), anti-CD3-APC-eFluro 780 (Invitrogen, 47-0037-41), anti-CD8-APC-

eFluro 780 (Invitrogen, 47-0086-42), anti-CD16-APC-eFluro 780 (Invitrogen, 47-0168-41), anti-

CD14-APC-eFluro 780 (Invitrogen, 47-0149-42), as well as Zombie NIR (BioLegend, 423105),

and fluorophore-labeled RBD and Ovalbumin for 30 minutes on ice[27]. Single CD3$^-$CD8$^-$CD16$^-$

CD20$^+$Ova$^-$RBD-PE$^+$RBD-AF647$^+$ B cells were sorted into individual wells of a 96-well plates

containing 4 μl of lysis buffer (0.5 X PBS, 10mM DTT, 3000 units/mL RNasin Ribonuclease

Inhibitors (Promega, N2615) per well using a FACS Aria III (Becton Dickinson). The sorted cells

were frozen on dry ice, and then stored at −80°C or immediately used for subsequent RNA reverse

transcription. Although cells were not stained for IgG expression, they are memory B cells based

on the fact that they are CD20+ (a marker absent in plasmablasts) and they express IgG (since

antibodies were amplified from these cells using IgG-specific primers).

Antibody sequencing, cloning and expression. Antibodies were identified and sequenced as described previously[22,28,29]. Briefly, RNA from single cells was reverse-transcribed (SuperScript III Reverse Transcriptase, Invitrogen, 18080-044) and the cDNA stored at -20°C or used for subsequent amplification of the variable IGH, IGL and IGK genes by nested PCR and Sanger sequencing[28]. Amplicons from the first PCR reaction were used as templates for Sequence- and Ligation-Independent Cloning (SLIC) into antibody expression vectors. Recombinant monoclonal antibodies and Fabs were produced and purified as previously described[30,31].

Biolayer interferometry.

BLI assays were performed on the Octet Red instrument (ForteBio) at 30°C with shaking at 1,000 r.p.m. Epitope binding assays were performed with protein A biosensor (ForteBio 18-5010), following the manufacture protocol "classical sandwich assay". (1) Sensor check: sensors immersed 30 sec in buffer alone (buffer ForteBio 18-1105). (2) Capture 1st Ab: sensors immersed 10min with Ab1 at 40 μg/mL. (3) Baseline: sensors immersed 30sec in buffer alone. (4) Blocking: sensors immersed 5 min with IgG isotype control at 50 μg/mL. (6) Antigen association: sensors immersed 5 min with RBD at 100 μg/mL. (7) Baseline: sensors immersed 30 sec in buffer alone. (8) Association Ab2: sensors immersed 5min with Ab2 at 40 μg/mL. Curve fitting was performed using the Data analysis software (ForteBio).

Computational analyses of antibody sequences. Antibody sequences were trimmed based on quality and annotated using Igblastn v1.14.0[32] with IMGT domain delineation system. Annotation was performed systematically using Change-O toolkit v.0.4.5[33]. Heavy and light chains derived

from the same cell were paired, and clonotypes were assigned based on their V and J genes using

in-house R and Perl scripts (Fig. 3 b,c). All scripts and the data used to process antibody sequences

are publicly available on GitHub (https://github.com/stratust/igpipeline).

The frequency distributions of human V genes in anti-SARS-CoV-2 antibodies from this study

was compared to Sequence Read Archive SRP010970[34]. The V(D)J assignments were done using

IMGT/High V-Quest and the frequencies of heavy and light chain V genes were calculated for 14

and 13 individuals, respectively, using sequences with unique CDR3s. The two-tailed t test with

unequal variances was used to determine statistical significance (Extended Data Fig. 7).

Nucleotide somatic hypermutation and CDR3 length were determined using in-house R and Perl

scripts. For somatic hypermutations, IGHV and IGLV nucleotide sequences were aligned against

their closest germlines using IgBlast and the number of differences were considered nucleotide

mutations. The average mutations for V genes was calculated by dividing the sum of all nucleotide

mutations across all patients by the number of sequences used for the analysis. To calculate the

GRAVY scores of hydrophobicity[35] we used Guy H.R. Hydrophobicity scale based on free energy

of transfer (kcal/mole)[36] implemented by the R package Peptides available in the Comprehensive

R Archive Network repository (https://journal.r-project.org/archive/2015/RJ-2015-001/RJ-2015-

001.pdf). We used 534 heavy chain CDR3 amino acid sequences from this study and 22,654,256

IGH CDR3 sequences from the public database of memory B-cell receptor sequences[37]. The

Shapiro-Wilk test was used to determine whether the GRAVY scores are normally distributed.

The GRAVY scores from all 533 IGH CDR3 amino acid sequences from this study

(sequence COV047_P4_IgG_51-P1369 lacks CDR3 amino acid sequence) were used to perform

the test and 5000 GRAVY scores of the sequences from the public database were randomly

selected. The Shapiro-Wilk p-values were 6.896 x 10^{-3} and 2.217 x 10^{-6} for sequences from this

study and the public database, respectively, indicating the data are not normally distributed.

Therefore, we used the Wilcoxon non-parametric test to compare the samples, which indicated a

difference in hydrophobicity distribution ($p = 5 \times 10^{-6}$; Extended Data Fig. 8).

Negative-stain EM Data Collection and Processing

Purified Fabs (C002, C119, and C121) were complexed with SARS-CoV-2 S trimer at a 2-fold

molar excess for 1 min and diluted to 40 μg/mL in TBS immediately before adding 3 μL to a

freshly-glow discharged ultrathin, 400 mesh carbon-coated copper grid (Ted Pella, Inc.). Samples

were blotted after a 1 min incubation period and stained with 1% uranyl formate for an additional

minute before imaging. Micrographs were recorded on a Thermo Fisher Talos Arctica transmission

electron microscope operating at 200 keV using a K3 direct electron detector (Gatan, Inc) and

SerialEM automated image acquisition software[38]. Images were acquired at a nominal

magnification of 28,000x (1.44 Å/pixel size) and a -1.5 to -2.0 μm defocus range. Images were

processed in cryoSPARC v2.14, and reference-free particle picking was completed using a

gaussian blob picker[39]. Reference-free 2D class averages and *ab initio* volumes were generated in

cryoSPARC, and subsequently 3D-classified to identify classes of S-Fab complexes, that were

then homogenously refined. Figures were prepared using UCSF Chimera[40].

Acknowledgements: We thank all study participants who devoted time to our research; Drs. Barry Coller and Sarah Schlesinger, the Rockefeller University Hospital Clinical Research Support Office and nursing staff. Dr. Joseph L. DeRisi for facilitating interactions with the Chan Zuckerberg BioHub. All members of the M.C.N. laboratory for helpful discussions, Drs. Amelia Escolano, Gaëlle Breton and Bernardo Reis, and Maša Jankovic for laboratory support. This work was supported by NIH grant P01-AI138398-S1 (M.C.N., C.M.R., P.J.B.) and 2U19AI111825 (M.C.N. and C.M.R).; the Caltech Merkin Institute for Translational Research and P50 AI150464 (P.J.B.), George Mason University Fast Grant and the European ATAC consortium (EC 101003650) to D.F.R.; 3 R01-AI091707-10S1 to C.M.R.; R37-AI64003 to P.D.B.; R01AI78788 to T.H.; The G. Harold and Leila Y. Mathers Charitable Foundation to C.M.R.. Electron microscopy was performed in the Caltech Beckman Institute Resource Center for Transmission Electron Microscpy (Drs. Songye Chen and Andrey Malyutin, Directors). C.G. was supported by the Robert S. Wennett Post-Doctoral Fellowship, in part by the National Center for Advancing Translational Sciences (National Institutes of Health Clinical and Translational Science Award program, grant UL1 TR001866), and by the Shapiro-Silverberg Fund for the Advancement of Translational Research. P.D.B. and M.C.N. are Howard Hughes Medical Institute Investigators.

Contributions: D.F.R., P.D.B., P.J.B., T.H., C.M.R. and M.C.N. conceived, designed and analyzed the experiments. D.F.R., M.C. and C.G. designed clinical protocols. F.M., J.C.C.L., Z.W., A.C., M.A., C.O.B., S.F., T.H., C.V., K.G., F.B., S.T.C., P.M., H.H., L.N., F.S., Y.W., H.-H.H., E.M., A.W.A., K.E.H.T., N.K. and P.R.H. carried out all experiments. A.G. and M.C. produced antibodies. C.O.B., J.P. and E.W. produced SARS-CoV-2 proteins. A.H., R.K., J.H., K.G.M., C.G. and M.C. recruited participants and executed clinical protocols. R.P., J.D., M.P. and

519   I.S. processed clinical samples. T.Y.O., A.P.W. and V.R. performed bioinformatic analysis.

520   D.F.R., P.D.B., P.J.B., T.H., C.M.R. and M.C.N. wrote the manuscript with input from all co-

521   authors.

523   **Declaration of conflict:** In connection with this work The Rockefeller University has filed a

524   provisional patent application on which D.F.R. and M.C.N. are inventors.

Extended Data Tables

Extended Data Table 1. Individual participant demographics and clinical characteristics

ID	Age	Gender	Race	Ethnicity	Case/Contact	Duration (days) Sx total	Duration (days) Sx onset to visit	Sx Severity (0-10)	ELISA binding (AUC) RBD IgG	ELISA binding (AUC) RBD IgM	ELISA binding (AUC) S IgG	ELISA binding (AUC) S IgM	Neutralization (NT50)
5	43	M	White	Non-Hispanic	Case	9	41	9	2.52	2.35	5.51	2.43	5.0
7	40	M	White	Non-Hispanic	Case	11	30	6	2.92	2.54	7.39	2.37	2730.4
8	37	M	White	Non-Hispanic	Case	3	57	5	2.11	0.81	4.46	0.88	5.0
9	35	F	White	Non-Hispanic	Case	12	54	5	2.90	1.07	4.44	2.80	5.0
12	27	F	White	Non-Hispanic	Contact	7	24	3	1.47	1.71	4.31	1.23	5.0
13	28	M	White	Non-Hispanic	Case	5	25	3	1.97	4.10	3.95	1.38	173.2
18	55	M	White	Non-Hispanic	Case	16	28	6	2.57	1.75	3.92	1.31	410.3
20	26	F	White	Non-Hispanic	Case	2	17	5	1.80	2.00	4.49	1.68	5.0
21	54	M	White	Hispanic	Contact	11	27	7	5.60	2.88	7.60	2.47	5052.7
24	34	M	White	Non-Hispanic	Case	15	30	4	2.29	1.97	4.36	1.45	280.7
26	66	M	White	Non-Hispanic	Case	2	35	3	1.67	1.86	4.00	1.58	276.1
27	28	M	White	Non-Hispanic	Case	9	32	4	1.79	1.64	4.41	1.30	739.3
28	26	M	White	Non-Hispanic	Case	7	21	4	2.39	2.25	4.93	1.66	888.9
29	26	F	White	Non-Hispanic	Contact	5	35	4	1.69	1.54	3.93	1.97	5.0
30	30	M	White	Non-Hispanic	Contact	3	35	4	3.07	2.55	4.86	1.34	5.0
31	51	M	White	Non-Hispanic	Case	9	33	3	1.29	1.60	4.69	1.20	192.3
32	46	F	White	Non-Hispanic	Case	8	32	3	1.30	2.18	3.79	0.83	47.4
37	27	M	White	Non-Hispanic	Case	6	28	6	2.13	1.52	4.11	1.04	286.3
38	57	F	White	Non-Hispanic	Case	10	38	4	1.87	2.02	4.30	1.12	518.9
40	44	M	White	Non-Hispanic	Case	7	23	5	1.72	1.50	3.81	1.47	42.1
41	35	M	White	Non-Hispanic	Contact	10	29	8	2.13	1.67	4.15	0.98	302.5
42	40	M	White	Non-Hispanic	Contact	20	36	8	2.37	1.50	5.82	1.43	627.1
45	55	F	White	Non-Hispanic	Case	7	48	6	1.95	0.97	3.53	0.75	5.0
46	39	M	White	Non-Hispanic	Case	8	30	2	2.68	1.45	4.14	1.29	59.2
47	43	F	White	Non-Hispanic	Case	11	33	5	3.02	2.30	6.23	2.66	10433.3
48	37	F	White	Non-Hispanic	Case	7	21	5	1.51	1.85	3.47	1.51	173.4
50	27	F	White	Non-Hispanic	Contact	7	28	4	1.64	2.32	5.42	1.45	924.7
51	21	M	White	Non-Hispanic	Contact	8	31	5	1.76	1.69	3.94	1.03	1499.2
54	40	F	White	Non-Hispanic	Contact	3	24	3	1.80	2.10	4.99	1.42	5.0
55	36	M	White	Non-Hispanic	Case	3	49	2	1.43	1.23	3.90	2.09	5.0
56	75	F	White	Non-Hispanic	Case	22	40	3	1.79	1.81	5.89	1.47	1388.4
57	66	M	White	Non-Hispanic	Case	6	21	5	1.54	2.10	4.33	1.00	2048.9
58	64	F	White	Non-Hispanic	Contact	1	32	2	1.20	1.71	3.95	1.21	5.0
64	28	F	White	Non-Hispanic	Contact	11	32	6	2.36	2.06	4.48	1.66	776.7
67	19	F	N/A	Hispanic	Case	5	29	6	2.56	1.98	5.48	1.32	2052.9
71	45	F	White	Non-Hispanic	Case	12	48	7	1.55	2.04	3.86	1.53	33.3
72	42	M	White	Non-Hispanic	Case	16	35	8	4.05	3.58	6.05	2.59	3138.2
75	46	F	White	Non-Hispanic	Case	10	36	4	1.64	2.37	3.98	1.20	271.5
76	49	F	White	Non-Hispanic	Case	28	34	4	1.88	1.65	5.17	0.97	219.8
77	37	M	White	Non-Hispanic	Contact	6	33	4	1.33	1.83	3.56	1.16	5.0
81	44	F	White	Non-Hispanic	Contact	3	35	2	1.58	1.73	3.82	1.10	5.0
82	46	M	N/A	Non-Hispanic	Case	0		0	2.19	1.92	5.41	1.81	130.7
88	41	M	White	Non-Hispanic	Case	7	23	4	1.82	3.32	4.97	1.37	424.7
95	44	M	White	Non-Hispanic	Case	9	36	6	2.61	1.62	6.03	1.62	961.9
96	48	F	White	Non-Hispanic	Case	9	30	3	3.93	1.93	6.25	2.26	927.7
97	39	M	White	Non-Hispanic	Case	9	31	3	1.58	1.61	4.03	2.33	202.7
98	35	F	White	Non-Hispanic	Case	2	24	4	1.78	1.47	5.38	1.22	249.0
99	36	F	White	Non-Hispanic	Case	13	29	5	2.50	3.27	4.38	2.49	1127.6
107	53	F	White	Non-Hispanic	Contact	10	29	4	1.74	1.41	4.66	0.90	297.5
108	75	M	White	Non-Hispanic	Case	16	41	7	1.37	0.88	3.47	1.36	557.5
110	27	M	White	Non-Hispanic	Case	1	25	1	1.30	1.60	4.00	0.93	5.0
114	30	F	White	Non-Hispanic	Case	15	36	7	1.65	1.92	3.53	1.55	110.9
115	65	F	White	Non-Hispanic	Contact	20	41	6	2.10	3.28	4.32	3.27	1127.7
119	56	M	White	Non-Hispanic	Case	13	48	3	1.36	1.26	4.41	1.59	650.3
120	56	F	White	Non-Hispanic	Case	26	48	6	0.99	0.97	3.65	1.21	100.6
121	19	M	White	Non-Hispanic	Contact	3	42	2	0.85	0.87	3.56	1.26	5.0
122	21	F	White	Non-Hispanic	Contact	3	38	1	1.44	0.94	3.39	1.22	5.0
123	26	M	White	Non-Hispanic	Contact	12	34	6	0.94	0.95	3.39	1.40	5.0
124	63	F	Asian	Non-Hispanic	Contact	4	37	3	1.58	2.06	3.49	1.32	5.0
125	51	F	White	Non-Hispanic	Case	10	26	3	1.92	3.49	3.86	1.24	126.5
127	24	F	White	Non-Hispanic	Case	10	43	6	1.80	2.50	4.37	2.41	883.5
130	39	M	White	Non-Hispanic	Contact	7	28	5	1.24	1.72	3.91	1.34	5.0
131	39	M	White	Non-Hispanic	Case	5	25	4	1.46	1.38	4.44	1.03	7.8
132	36	M	White	Non-Hispanic	Contact	10	50	6	2.15	1.76	4.84	1.97	5.0
134	27	F	White	Non-Hispanic	Contact	16	22	5	2.51	2.18	6.84	1.94	2700.6
135	62	F	White	Non-Hispanic	Case	8	31	6	2.20	2.02	3.80	1.13	350.0
140	63	F	White	Non-Hispanic	Case	28	47	1	1.05	1.24	3.58	1.28	52.4
149	41	M	White	Non-Hispanic	Contact	17	28	6	1.68	2.02	3.67	1.09	494.9
150	50	F	White	Non-Hispanic	Contact	12	45	7	1.15	0.43	3.18	1.82	5.0
154	68	M	Asian	Non-Hispanic	Case	16	30	9	3.19	2.19	4.85	1.29	928.2
157	50	M	White	Non-Hispanic	Case	10	32	8	2.40	2.86	3.90	2.06	741.7
166	28	F	White	Non-Hispanic	Case	13	45	2	1.27	0.66	3.45	0.94	5.0
167	50	F	White	Non-Hispanic	Contact	11	41	6	1.43	3.71	3.93	8.74	5.0
172	38	F	White	Non-Hispanic	Case	8	22	9	1.71	2.58	4.29	1.20	301.1
173	47	M	White	Non-Hispanic	Case	5	47	7	2.57	4.14	4.78	4.48	646.9
178	26	F	White	Non-Hispanic	Case	6	24	4	1.54	1.59	3.66	1.02	5.0
179	39	M	White	Non-Hispanic	Contact	10	37	3	1.89	2.25	3.83	1.73	370.1
182	44	F	White	Non-Hispanic	Contact	10	38	6	3.80	1.77	5.36	8.05	1503.7
183	43	F	White	Non-Hispanic	Case	13	44	8	1.56	1.02	3.88	1.55	240.1
185	54	M	White	Non-Hispanic	Case	11	44	8	3.51	1.39	5.55	2.11	1806.8
186	38	F	N/A	N/A	Case	8	26	2	1.73	2.47	4.37	1.23	296.9
195	24	M	White	Non-Hispanic	Case	18	42	5	2.74	2.55	5.01	2.33	1315.1
200	60	F	White	Non-Hispanic	Case	17	39	7	2.40	1.00	4.38	1.51	1014.4
201	50	M	White	Non-Hispanic	Contact	15	33	6	4.37	2.57	6.15	1.57	3897.4
202	57	M	White	Non-Hispanic	Case	21	34	7	2.10	2.08	5.07	1.17	257.9
205	64	M	White	Non-Hispanic	Case	7	36	4	4.51	0.70	6.12	1.69	924.4
222	28	M	Asian	Non-Hispanic	Case	11	29	7	1.28	0.69	3.94	3.46	5.0
229	45	M	White	Non-Hispanic	Case	10	63	4	2.92	1.42	4.90	1.58	1272.9
230	50	M	White	Non-Hispanic	Case	18	33	7	3.80	0.47	3.48	0.88	5.0
232	38	F	White	Non-Hispanic	Case	13	43	7	1.57	0.70	4.24	5.70	94.3
233	55	M	White	Non-Hispanic	Case	20	41	3	2.07	2.11	4.51	1.07	173.2
241	36	M	White	Non-Hispanic	Case	12	30	7	2.27	2.66	4.54	1.46	923.1

ID	Age	Sex	Race	Ethnicity	Case/Contact								
	30	M	White	Non-Hispanic	Case	10	52	8	2.40	2.86	5.50	2.50	741.7
166	28	F	White	Non-Hispanic	Case	13	45	2	1.27	0.66	3.45	0.94	5.0
167	50	F	White	Non-Hispanic	Contact	11	41	6	1.43	3.71	3.93	8.74	5.0
172	38	F	White	Non-Hispanic	Case	8	22	9	1.71	2.58	4.29	1.20	301.1
173	47	M	White	Non-Hispanic	Case	5	47	7	2.57	4.14	4.78	4.48	646.9
178	26	F	White	Non-Hispanic	Case	6	24	4	1.54	1.59	3.66	1.02	5.0
179	39	M	White	Non-Hispanic	Contact	10	37	3	1.89	2.25	3.83	1.73	370.1
182	44	F	White	Non-Hispanic	Contact	10	38	6	3.80	1.77	5.36	8.05	1503.7
183	43	F	White	Non-Hispanic	Case	13	44	8	1.56	1.02	3.88	1.55	240.1
185	54	M	White	Non-Hispanic	Case	11	44	8	3.51	1.39	5.55	2.11	1806.8
186	38	F	N/A	N/A	Case	8	26	2	1.73	2.47	4.37	1.23	296.9
190	54	F	White	Non-Hispanic	Case	18*	63	9	3.24	1.24	7.38	1.42	598.1
195	24	M	White	Non-Hispanic	Case	18	42	5	2.74	2.55	5.01	2.33	1315.1
200	60	F	White	Non-Hispanic	Case	17	39	7	2.40	1.00	4.38	1.51	1014.4
201	50	M	White	Non-Hispanic	Contact	15	33	6	4.37	2.57	6.15	1.57	3897.4
202	57	M	White	Non-Hispanic	Case	21	34	7	2.10	2.08	5.07	1.17	257.9
205	64	M	White	Non-Hispanic	Case	7	36	4	4.51	0.70	6.12	1.69	924.4
222	28	M	Asian	Non-Hispanic	Case	11	29	7	1.28	0.69	3.94	3.46	5.0
229	45	M	White	Non-Hispanic	Case	10	63	4	2.92	1.42	4.90	1.58	1272.9
230	50	M	White	Non-Hispanic	Case	18	33	7	3.80	0.47	3.48	0.88	5.0
232	38	F	White	Non-Hispanic	Case	13	43	7	1.57	0.70	4.24	5.70	94.3
233	55	M	White	Non-Hispanic	Case	20	41	3	2.07	2.11	4.51	1.07	173.2
241	36	M	White	Non-Hispanic	Case	12	30	7	2.27	2.66	4.54	1.46	923.1
242	59	M	White	Non-Hispanic	Case	10	42	6	4.91	1.94	4.81	2.16	1353.0
243	30	F	Asian	Non-Hispanic	Case	6	26	5	2.92	2.57	5.06	1.14	1300.2
246	44	F	White	Non-Hispanic	Case	10	38	7	2.05	2.79	6.09	1.32	566.0
255	33	M	White	Non-Hispanic	Case	14	44	6	2.14	0.70	4.20	1.24	172.5
256	63	F	White	Non-Hispanic	Case	27	42	6	1.72	1.96	4.26	7.79	141.6
258	52	M	White	Non-Hispanic	Contact	14	48	6	2.64	1.20	4.52	1.85	4145.9
279	41	M	White	Non-Hispanic	Case	7	38	8	1.68	2.13	3.77	1.90	308.9
280	59	M	White	Non-Hispanic	Case	6	32	7	2.53	3.07	4.61	1.19	1072.1
302	47	F	White	Non-Hispanic	Case	35*	49	7	1.48	0.97	4.06	1.26	5.0
310	34	F	White	Non-Hispanic	Case	17	35	5	3.95	1.24	9.44	3.07	485.5
314	46	M	White	Non-Hispanic	Case	11	38	7	2.12	0.88	4.56	1.51	667.1
315	29	F	White	Non-Hispanic	Case	15	42	8	3.02	0.69	3.98	1.03	376.5
319	50	M	White	Non-Hispanic	Case	5	38	6	3.71	2.28	3.79	1.05	5.0
323	39	F	White	Non-Hispanic	Case	7	45	7	1.05	1.03	3.53	1.42	5.0
325	52	M	White	Non-Hispanic	Case	16	38	8	2.25	1.47	4.83	2.28	1603.3
343	21	F	White	Non-Hispanic	Case	16	49	5	1.63	0.94	3.37	1.70	5.0
352	44	M	White	Non-Hispanic	Case	16	43	4	3.54	0.92	4.50	1.07	519.2
353	60	M	White	Non-Hispanic	Case	14	49	6	5.38	1.12	5.69	1.05	855.5
356	22	F	White	Non-Hispanic	Contact	16	38	3	1.37	0.61	2.98	1.09	5.0
357	27	F	White	Non-Hispanic	Contact	34	56	5	7.49	1.10	2.77	1.13	5.0
364	29	M	White	Non-Hispanic	Contact	14	49	6	0.97	0.58	2.90	0.89	5.0
366	41	F	White	Non-Hispanic	Contact	9	34	7	0.98	0.52	3.51	1.19	5.0
373	35	F	White	Non-Hispanic	Case	12	51	7	1.69	1.26	5.14	1.69	5.0
388	47	F	White	Non-Hispanic	Contact	14	41	9	1.57	1.06	3.61	1.69	5.0
393	69	M	White	Non-Hispanic	Case	23*	54	9	1.28	1.74	3.81	2.65	715.4
394	48	F	Multiple	Hispanic	Case	7	67	4	2.05	0.87	4.34	2.02	1281.5
397	52	M	White	Non-Hispanic	Case	22	45	8	3.32	0.59	5.01	0.87	1516.9
403	52	M	Asian	Non-Hispanic	Case	18*	39	10	5.36	1.09	10.01	1.36	3887.8
406	65	M	White	Non-Hispanic	Case	20	56	8	4.69	0.90	7.51	1.15	1288.7
410	34	M	White	Non-Hispanic	Case	12	46	8	1.06	0.56	3.95	0.76	5.0
421	62	F	White	Non-Hispanic	Contact	12	43	9	0.95	1.07	3.34	1.35	5.0
426	65	M	White	Non-Hispanic	Case	18	51	6	2.07	0.55	3.95	1.49	804.8
437	43	F	Asian	Non-Hispanic	Case	14	34	7	2.54	0.47	4.30	1.44	698.8
460	36	M	White	Non-Hispanic	Case	11	39	6	2.94	3.18	5.51	2.80	1906.7
461	49	M	White	Non-Hispanic	Case	7	39	5	3.38	0.94	4.67	2.02	1076.6
462	28	F	White	Non-Hispanic	Case	16	45	5	1.36	0.38	3.07	1.11	5.0
470	28	F	White	Non-Hispanic	Case	17	51	4	1.26	0.86	3.97	1.50	5.0
478	31	M	White	Non-Hispanic	Case	16	52	4	1.43	0.93	3.70	1.97	263.2
481	28	F	Asian	Non-Hispanic	Case	15	43	8	1.70	0.39	3.46	1.24	5.0
486	64	F	White	Non-Hispanic	Case	11	41	10	1.70	1.00	3.68	1.29	5.0
500	46	M	White	Non-Hispanic	Case	12	53	5	1.10	0.82	3.49	1.34	5.0
501	32	M	Asian	Non-Hispanic	Case	18*	53	10	2.62	0.65	4.51	1.21	718.8
502	52	M	White	Non-Hispanic	Case	16*	53	9	5.10	0.61	5.10	1.62	2171.8
506	46	M	White	Non-Hispanic	Case	12	59	9	0.84	0.81	3.13	1.21	5.0
507	39	M	White	Non-Hispanic	Case	15	60	8	1.92	0.96	4.56	1.52	5.0
509	36	M	White	Non-Hispanic	Case	11	50	5	1.99	1.01	3.99	1.45	5.0
526	49	M	Asian	Non-Hispanic	Case	11	34	7	3.36	1.45	5.88	1.57	4193.3
537	52	M	White	Non-Hispanic	Case	15	45	6	1.47	0.95	3.65	1.58	923.3
539	73	F	White	Non-Hispanic	Case	19*	54	10	2.82	0.63	4.46	1.45	487.9
547	59	M	White	Non-Hispanic	Case	15*	36	9	2.97	1.53	5.08	2.59	2900.6
587	54	M	PI	N/A	Case	17*	51	8	3.22	0.60	4.01	1.49	473.1
632	38	M	White	Non-Hispanic	Contact	10	43	6	2.49	0.86	4.50	1.63	572.3
633	39	M	White	Non-Hispanic	Contact	8	57	4	1.25	1.04	3.38	1.73	5.0
652	76	M	White	Non-Hispanic	Case	18*	56	10	4.75	1.46	8.96	3.80	2324.0
664	45	F	White	Non-Hispanic	Case	17*	42	10	1.68	0.43	3.93	1.32	5.0
675	47	M	White	Non-Hispanic	Contact	31	47	5	0.79	0.93	2.94	1.38	5.0

*hospitalized, Sx=symptoms

Extended Data Table 2. Frequency of symptoms and comorbidities reported by participants

Symptom	Participants (n=149)	%
Fever	125	83.9
Fatigue	106	71.1
Cough	93	62.4
Myalgia	92	61.7
Shortness of breath	66	44.3
Headache	63	42.3
Loss of smell/taste	50	33.3
Sore throat	38	25.3
Diarrhea	32	21.3
Presence of comorbidities (HTN, CAD, DM, COPD, asthma, cancer)	16	10.7

HTN (hypertension), CAD (coronary artery disease), DM (diabetes mellitus), COPD (chronic obstructive pulmonary disease)

Extended Data Tables 3 and 4 are provided as separate Excel files.

Extended data table 5. Sequences of cloned recombinant antibodies

Antibody ID	IGH VDJ (nt)	IGH VDJ (aa)	IGL VJ (nt)	IGL VJ (aa)
C002	GAGGTGCAGC	EVQLVESGGGV	GACATCCAG	DIQLTQSPSSL
C003	GAGGTGCAGC	EVQLVESGGGL	GAAATTGTGT	EIVLTQSPGTLS
C004	CAGGTGCAGC	QVQLVQSGAEV	GCCATCCGG	AIRMTQSPSSL
C005	CAGGTGCAGC	QVQLVQSGPEV	GAAATTGTGT	EIVLTQSPGTLS
C006	CAGGTGCAGC	QVQLVESGGGL	CAGTCTGTG	QSVLTQPPSAS
C008	GAGGTGCAGC	EVQLVESGGGV	GACATCCAG	DIQMTQSPSTL
C009	CAGGTGCAGC	QVQLVQSGAEV	CAGTCTGCC	QSALTQPPSAS
C010	GAGGTGCAGC	EVQLVESGGGV	GACATCCAG	DIQLTQSPSSL
C013	CAGGTGCAGC	QVQLVQSGAEV	GAAATTGTGT	EIVLTQSPATLS
C016	GAGGTGCAGC	EVQLVESGGGV	GCCATCCGG	AIRMTQSPSSL
C017	GAAGTGCAGC	EVQLVESGGGL	GAAATTGTGT	EIVLTQSPATLS
C018	GAGGTGCAGC	EVQLVESGGGV	GACATCCAG	DIQLTQSPSSL
C019	CAGGTGCAGC	QVQLVQSGAEV	TCCTATGAGC	SYELTQPPSVS
C021	CAGGTGCAGC	QVQLQESGPGL	GATATTGTGA	DIVMTQSPLSLI
C022	CAGGTGCAGT	QVQLQESGPGL	GACATCCAG	DIQMTQSPSTL
C027	GAGGTGCAGC	EVQLVESGGGV	GACATCCAG	DIQMTQSPSTL
C029	CAGGTGCAGC	QVQLQESGPGL	GATATTGTGA	DIVMTQSPLSLI
C030	GAGGTGCAGC	EVQLVESGGGV	GACATCCAG	DIQMTQSPSTL
C031	GAGGTGCAGC	EVQLVESGGGL	GACATCCAG	DIQMTQSPSSL
C101	CAGGTGCAGC	QVQLVESGGGL	GAAATTGTGT	EIVLTQSPGTLS
C102	CAGGTGCAGC	QVQLVESGGGL	GAAATTGTGT	EIVLTQSPGTLS
C103	CAGGTGCAGC	QVQLQQWGAGL	GAAATTGTGT	EIVLTQSPGTLS
C104	CAGGTGCAGC	QVQLQQWGAGL	GAAATTGTGT	EIVLTQSPGTVS
C105	CAGGTGCAGC	QVQLVESGGGL	CAGTCTGCC	QSALTQPPSAS
C106	CAGCTGCAGC	QLQLQESGPGL	TCCTATGAGC	SYELTQPPSVS
C107	CAGGTTCAGC	QVQLVQSGAEV	CAGTCTGTG	QSVLTQPPSAS
C108	CAGGTGCAGC	QVQLQESGPGL	CAGTCTGCC	QSALTQPASVS
C109	GAAGTGCAGC	EVQLVESGGGL	CAGTCTGCC	QSALTQPPSAS
C110	CAGGTACAGC	QVQLQQSGAEV	GACATCCAG	DIQMTQSPSTL
C111	CAGCTGCAGC	QLQLQESGPGL	CAGTCTGTG	QSVLTQPPSVS
C112	CAGGTGCAGC	QVQLVESGGGV	CAGTCTGCC	QSALTQPASVS
C113	CAGGTGCAGC	QVQLVESGGGV	GACATCCAG	DIQMTQSPSTL
C114	CAGGTGCAGC	QVQLVESGGGL	CAGTCTGTG	QSVLTQPPSVS
C115	CAGGTGCAGC	QVQLVESGGGL	GATATTGTGA	DIVMTQSPLSL
C116	CAGGTGCAGC	QVQLVESGGGV	AATTTTATGC	NFMLTQPHSVS
C117	CAGGTGCAGC	QVQLVESGGGV	CAGTCTGTG	QSVLTQPPSVS
C118	CAGGTGCAGC	QVQLVESGGGV	CAGCCTGTG	QPVLTQPSPAS
C119	CAGGTCCAGC	QVQLVQSGAEV	CAGTCTGCC	QSALTQPASVS
C120	GAGGTGCAGC	EVQLVESGGGL	GCCATCCGG	AIRMTQSPSSL
C121	CAGGTGCAGC	QVQLVQSGAEV	CAGTCTGCC	QSALTQPASVS
C122	GAGGTGCAGC	EVQLVESGGGL	GACATCCAG	DIQLTQSPSFLS
C123	GAGGTGCAGC	EVQLVESGGGL	GACATCCAG	DIQLTQSPSFLS
C124	GAGGTGCAGC	EVQLVESGGGL	GAAATTGTGT	EIVLTQSPATLS
C125	CAGGTGCAGC	QVQLVQSGPEV	GAAATTGTGT	EIVLTQSPGTLS
C126	CAGGTGCAGC	QVQLQESGPGL	AATTTTATGC	NFMLTQPHSVS
C127	CAGGTGCAGC	QVQLVQSGAEV	CAGTCTGTG	QSVLTQPPSAS
C128	GAGGTGCAGC	EVQLLESGGGL	GAAATTGTGT	EIVLTQSPGTLS
C130	CAGGTGCAGC	QVQLVQSGAEV	TCCTATGAGC	SYELTQPPSVS
C131	CAGGTGCAGC	QVQLVQSGSEV	GAAATAGTGA	EIVMTQSPATLS
C132	CAGGTGCAGC	QVQLQESGPGL	CAGTCTGCC	QSALTQPASVS
C133	CAGGTGCAGC	QVQLVESGGGV	GACATCCAG	DIQMTQSPSSL
C134	GAGGTGCAGC	EVQLLESGGGL	TCCTATGAGC	SYELTQPPSVS
C135	CAGGTGCAGC	QVQLVESGGGV	GACATCCAG	DIQMTQSPSTL
C138	GAGGTGCAGC	EVQLVESGGGL	AATTTTATGC	NFMLTQPHSVS
C139	GAGGTGCAGC	EVQLVESGGGV	GACATCCAG	DIQMTQSPSSL
C140	GAGGTGCAGC	EVQLVESGGGL	GACATCCAG	DIQLTQSPSFLS
C141	GAGGTGCAGC	EVQLVESGGGV	AATTTTATGC	NFMLTQPHSVS
C143	GAGGTGCAGC	EVQLVESGGGL	CAGTCTGCC	QSALTQPASVS
C144	GAGGTGCAGC	EVQLVESGGGL	CAGTCTGCC	QSALTQPASVS
C145	GAGGTGCAGC	EVQLVESGGGL	CAGTCTGCC	QSALTQPASVS
C146	GAGGTGCAGC	EVQLVESGGGL	CAGTCTGCC	QSALTQPASVS
C147	GAGGTGCAGC	EVQLVQSGAEV	CAGGCTGTG	QAVVTQEPSLT
C148	GAGGTGCAGC	EVQLVESGGGL	GAAATAGTGA	EIVMTQSPATLS
C150	GAGGTGCAGC	EVQLVESGGGL	CAGTCTGCC	QSALTQPASVS
C151	GAGGTGCAGC	EVQLVESGGGL	AATTTTATGC	NFMLTQPHSVS
C153	GAGGTGCAGC	EVQLVESGGGL	CAGTCTGCC	QSALTQPASVS
C154	CAGGTGCAGC	QVQLVESGGGV	GACATCCAG	DIQMTQSPSSL
C155	GAGGTGCAGC	EVQLVESGGGL	GAAATAGTGA	EIVMTQSPATLS
C156	CAGGTGCAGC	QVQLVESGGGV	TCCTATGAGC	SYELTQPPSVS
C164	GAGGTGCAGC	EVQLVESGGGL	CAGTCTGCC	QSALTQPASVS
C165	CAGGTGCAGC	QVQLVQSGAEV	GAAATTGTGT	EIVLTQSPGTLS
C201	GAAGTGCAGC	EVQLVESGGGL	CATCCGGAT	IRMTQSPSSVS
C202	GAGGTGCAGC	EVQLVESGGGL	GACATCCAG	DIQLTQSPSSL
C204	GAGGTGCAGC	EVQLLESGGGL	GACATCCAG	DIQLTQSPSSL
C205	CAGGTGCAGC	QVQLVQSGAEV	GAAATTGTGT	EIVLTQSPGTLS
C207	GAGGTGCAGC	EVQLLESGGGL	GAAATTGTGT	EIVLTQSPATLS
C208	GAGGTGCAGC	EVQLVQSGAEV	GAAATTGTGT	EIVLTQSPGTLS
C209	CAGGTGCAGC	QVQLVQSGAEV	TCCTATGAGC	SYELTQPPSVS
C210	GAGGTGCAGC	EVQLVESGGGL	GACATCCAG	DIQLTQSPSFLS
C211	GAGGTGCAGC	EVQLVESGGGL	GAAATAGTGA	EIVMTQSPATLS
C212	CAGGTGCAGC	QVQLVQSGAEV	CTGACTCAG	LTQPASVSGSP
C214	CAGGTGCAGC	QVQLVESGGGV	GACATCCAG	DIQLTQSPSSL
C215	GAGGTGCAGC	EVQLLESGGGL	GACATCCAG	DIQLTQSPSSL
C216	GAGGTGCAGC	EVQLVESGGGL	GACATCCAG	DIQLTQSPSSL

Extended Data Table 6. Effective and inhibitory concentrations of the monoclonal antibodies

Participant ID	Antibody ID	SARS-CoV-2 IC50 ng/ml	SARS-CoV-2 IC80 ng/ml	SARS-CoV-2 IC90 ng/ml	SARS-CoV-2 RBD EC50 ng/ml	SARS-CoV RBD EC50 ng/ml
COV21	C002	8.88	21.95	37.61	3.14	>1000
COV21	C003	313.79	992.62	>1000	6.37	>1000
COV21	C004	10.67	41.08	91.71	2.39	>1000
COV21	C005	60.49	130.65	205.20	4.41	>1000
COV21	C006	321.51	>1000	>1000	1.81	>1000
COV21	C008	625.46	>1000	>1000	4.63	>1000
COV21	C009	4.82	14.54	29.34	1.80	>1000
COV21	C010	>1000	>1000	>1000	5.44	>1000
COV21	C013	42.48	360.59	>1000	2.59	>1000
COV21	C016	>1000	>1000	>1000	7.41	>1000
COV21	C017	72.67	256.18	543.87	1.63	>1000
COV21	C018	>1000	>1000	>1000	1.53	>1000
COV21	C019	>1000	>1000	>1000	11.85	>1000
COV21	C021	>1000	>1000	>1000	1.25	>1000
COV21	C022	73.57	314.71	736.87	2.40	5.99
COV21	C027	>1000	>1000	>1000	2.65	696.05
COV21	C029	>1000	>1000	>1000	4.13	>1000
COV21	C030	>1000	>1000	>1000	2.89	>1000
COV21	C031	>1000	>1000	>1000	22.69	>1000
COV107	C101	8.20	30.15	65.30	1.51	>1000
COV107	C102	34.03	84.21	143.23	4.54	>1000
COV107	C103	4.38	12.58	23.59	3.77	>1000
COV107	C104	23.31	72.12	140.28	8.31	>1000
COV107	C105	26.09	72.24	133.70	5.20	>1000
COV107	C106	>1000	>1000	>1000	19.03	106.75
COV107	C107	>1000	>1000	>1000	11.55	>1000
COV107	C108	480.69	>1000	>1000	5.32	>1000
COV107	C110	18.44	45.11	77.28	7.29	>1000
COV107	C112	111.79	701.99	>1000	3.38	>1000
COV107	C113	>1000	>1000	>1000	6.93	>1000
COV107	C114	>1000	>1000	>1000	9.51	>1000
COV107	C115	198.33	958.18	>1000	3.40	>1000
COV107	C116	>1000	>1000	>1000	37.56	>1000
COV107	C117	348.00	>1000	>1000	5.38	>1000
COV107	C118	103.69	417.76	>1000	3.45	3.817
COV107	C119	9.12	39.45	97.78	3.57	>1000
COV107	C120	13.26	26.73	40.30	1.41	>1000
COV107	C121	6.73	14.31	22.33	2.85	>1000
COV107	C122	22.80	57.77	100.12	2.67	>1000
COV107	C123	149.22	355.47	595.51	1.92	>1000
COV107	C124	341.82	937.26	>1000	2.23	>1000
COV107	C125	43.32	92.54	144.26	1.87	>1000
COV107	C126	>1000	>1000	>1000	4.78	>1000
COV107	C127	68.74	190.96	347.31	2.62	>1000
COV072	C128	101.22	263.35	460.73	1.95	>1000
COV072	C130	>1000	>1000	>1000	2.05	>1000
COV072	C131	30.52	178.90	759.11	1.67	>1000
COV072	C132	708.67	>1000	>1000	2.92	>1000
COV072	C133	>1000	>1000	>1000	1.98	>1000
COV072	C134	>1000	>1000	>1000	1.57	>1000
COV072	C135	16.61	32.81	48.90	1.80	>1000
COV072	C136	>1000	>1000	>1000	33.61	>1000
COV072	C137	>1000	>1000	>1000	25.22	>1000
COV072	C138	>1000	>1000	>1000	2.94	>1000
COV072	C139	>1000	>1000	>1000	1.89	>1000
COV072	C140	23.88	66.24	120.69	2.19	>1000
COV072	C141	>1000	>1000	>1000	1.71	>1000
COV047	C143	>1000	>1000	>1000	3.66	>1000
COV047	C144	6.91	17.28	29.66	3.24	>1000
COV047	C145	3.04	14.51	36.79	3.86	>1000
COV047	C146	>1000	>1000	>1000	>1000	>1000
COV047	C147	>1000	>1000	>1000	>1000	>1000
COV047	C148	>1000	>1000	>1000	64.69	>1000
COV047	C150	>1000	>1000	>1000	8.11	>1000
COV047	C151	31.79	363.97	>1000	4.30	>1000
COV047	C153	70.71	490.08	>1000	3.17	>1000
COV047	C154	435.50	>1000	>1000	2.92	10.65
COV047	C155	11.00	35.75	77.01	3.30	>1000
COV047	C156	>1000	>1000	>1000	3.32	>1000
COV047	C164	239.15	865.40	>1000	2.06	>1000
COV072	C165	40.81	138.66	297.38	4.25	>1000
COV96	C201	>1000	>1000	>1000	2.98	>1000
COV96	C202	>1000	>1000	>1000	3.40	>1000
COV96	C204	>1000	>1000	>1000	3.73	9.41
COV96	C205	>1000	>1000	>1000	>1000	>1000
COV96	C207	158.52	960.39	>1000	1.87	>1000
COV96	C208	>1000	>1000	>1000	>1000	>1000
COV96	C209	>1000	>1000	>1000	3.79	>1000
COV96	C210	50.73	155.24	298.90	2.83	>1000
COV96	C211	12.79	34.87	62.89	2.82	>1000
COV96	C212	>1000	>1000	>1000	>1000	>1000
COV96	C214	>1000	>1000	>1000	5.75	>1000
COV96	C215	>1000	>1000	>1000	5.33	17.944
COV96	C216	>1000	>1000	>1000	9.53	>1000

Extended Data Figures

Extended Data Figure 1. Clinical correlates. a, Age distribution (Y axis) for all males and

females in the cohort p=0.2074. **b,** Duration of symptoms in days (Y axis) for all males and females

in the cohort p=0.8704. **c,** Time between symptom onset and plasma collection (Y axis) for all

males and females in the cohort p=0.5514. **d,** Subjective symptom severity on a scale of 0-10 (Y

axis) for all males and females in the cohort p=0.1888. **e,** Age distribution (Y axis) for all cases

and contacts in the cohort p=0.0305. **f,** Duration of symptoms in days (Y axis) for all cases and

contacts in the cohort p=0.1241. **g,** Time between symptom onset and plasma collection in days

(Y axis) for all cases and contacts in the cohort p=0.1589. **h,** Symptom severity (Y axis) for all

cases and contacts in the cohort p=0.0550. **i,** Age distribution (Y axis) for all outpatient and

hospitalized participants p=0.0024. **j,** Duration of symptoms in days (Y axis) for all outpatient and

hospitalized participants in the cohort p=<0.0001. **k,** Time between symptom onset and plasma

collection in days (Y axis) for all outpatient and hospitalized participants in the cohort p=0.0001.

l, Symptom severity (Y axis) for all outpatient and hospitalized participants in the cohort

p=<0.0001. Horizontal bars indicate median values. Statistical significance was determined using

two-tailed Mann-Whitney U test.

Extended Data Figure 2. Clinical correlates of plasma antibody titers. a, AUC for IgG anti-RBD (Y axis) for all cases and contacts in the cohort p=0.0107. **b,** AUC for IgM anti-RBD (Y axis) for all cases and contacts in the cohort p=0.5371. **c,** AUC for IgG anti-S (Y axis) for all cases and contacts in the cohort p=0.0135. **d,** AUC for IgM anti-S (Y axis) for all cases and contacts in the cohort p=0.7838. **e,** AUC for IgM anti-RBD (Y axis) for all males and females in the cohort p=0.9597. **f,** AUC for IgG anti-S (Y axis) for all males and females in the cohort p=0.0275. **g,** AUC for IgM anti-S (Y axis) for all males and females in the cohort p=0.5363. **h,** AUC for IgM anti-RBD (Y axis) for all outpatient and hospitalized participants in the cohort p=0.0059. **i,** AUC for IgG anti-S (Y axis) for all outpatient and hospitalized participants in the cohort p=0.0623. **j,** AUC for IgM anti-S (Y axis) for all outpatient and hospitalized participants in the cohort p=0.2976. Horizontal bars indicate median values. Statistical significance was determined using two-tailed Mann-Whitney U test.

Extended Data Figure 3. Additional clinical correlates of plasma antibody titers. a, Time between symptom onset and plasma collection in days (X axis) plotted against AUC for IgG anti-RBD (Y axis) r=-0.0261 p=0.7533. **b,** Time between symptom onset and plasma collection in days (X axis) plotted against AUC for IgG anti-S (Y axis) r=-0.1495 p=0.0697. **c,** Time between symptom onset and plasma collection in days (X axis) plotted against AUC for IgM anti-S (Y axis) r=0.1496 p=0.0695. **d,** Age (X axis) plotted against AUC for IgM anti-RBD (Y axis) r=0.0172 p=0.8355. **e,** Age (X axis) plotted against AUC for IgG anti-S (Y axis) r=0.1523 p=0.0638. **f,** Age (X axis) plotted against AUC for IgM anti-S (Y axis) r=0.0565 p=0.4934. **g,** Duration of symptoms in days (X axis) plotted against AUC for IgG anti-RBD (Y axis) r=0.1525, p=0.0633. **h,** Duration of symptoms in days (X axis) plotted against AUC for IgM anti-RBD (Y axis) r=-0.3187,

p=<0.0001. **i,** Duration of symptoms in days (X axis) plotted against AUC for IgG anti-S (Y axis)

r=0.0329, p=0.6904. **j,** Duration of symptoms in days (X axis) plotted against AUC for IgM anti-

S (Y axis) r=0.0824, p=0.3177. **k,** Severity of symptoms (X axis) plotted against AUC for IgG

anti-RBD (Y axis) r=0.2679 p=0.0010. **l,** Severity of symptoms (X axis) plotted against AUC for

IgM anti-RBD (Y axis) r=-0.1943 p=0.0176. **m,** Severity of symptoms (X axis) plotted against

AUC for IgG anti-S (Y axis) r=0.1187 p=0.1492. **n,** Severity of symptoms (X axis) plotted against

AUC for IgM anti-S (Y axis) r=0.1597 p=0.0517. All correlations were analyzed by two-tailed

Spearman's.

Extended Data Figure 4. Diagrammatic representation of the SARS-CoV2 pseudovirus luciferase assay. a, Co-transfection of pNL4-3ΔEnv-nanoluc and pSARS-CoV-2 spike vectors into 293T cells leads to production of SARS-CoV-2 Spike-pseudotyped HIV-1 particles (SARS-CoV-2 pseudovirus) carrying the *Nanoluc* gene. **b,** SARS-CoV-2 pseudovirus is incubated for 1 h at 37°C with plasma or monoclonal antibody dilutions. The virus-antibody mixture is used to infect ACE2-expressing 293T cells, which will express nanoluc Luciferase upon infection. **c,** Relative luminescence units (RLU) reads from lysates of ACE2-expressing 293T cells infected with increasing amounts of SARS-CoV-2 pseudovirus.

Extended Data Figure 5. Clinical correlates of neutralization. a, Anti-RBD IgM AUC (X axis) plotted against NT_{50} (Y axis) r=0.3119, p=0.0001. **b,** Anti-S IgM AUC (X axis) plotted against NT_{50} (Y axis) r=0.3211, p=<0.0001. **c,** Duration of symptoms in days (X axis) plotted against NT_{50} (Y axis) r=0.1997, p=0.0146. **d,** Time between symptom onset and plasma collection in days (X axis) plotted against NT_{50} (Y axis) r=-0.1344, p=0.1033. **e,** Symptom severity (X axis) plotted

597 against NT_{50} (Y axis) r=0.2234, p=0.0062. **f,** Age (X axis) plotted against NT_{50} (Y axis) r=0.3005,

598 p=0.0002. All correlations were analyzed by two-tailed Spearman's. Dotted line (NT_{50}=5)

599 represents lower limit of detection (LLOD) of pseudovirus neutralization assay. Samples with

600 undetectable neutralizing titers were plotted at LLOD.

603 **Extended Data Figure 6. Flow cytometry.** Gating strategy used for cell sorting. Gating was on

604 singlets that were CD20[+] and CD3[-]CD8[-]CD16[-]Ova[-]. Sorted cells were RBD-PE[+] and RBD-

605 AF647[+].

Extended Data Figure 7. Frequency distributions of human V genes. The two-tailed t test with unequal variance was used to compare the frequency distributions of human V genes of anti-SARS-CoV-2 antibodies from this study to Sequence Read Archive SRP010970[34].

Extended Data Figure 8. Analysis of antibody somatic hypermutation and CDR3 length. a,
For each individual, the number of somatic nucleotide mutations (Y axis) at the IGVH and IGVL
are shown on the left panel, and the amino acid length of the CDR3s (Y axis) are shown on the
right panel. The horizontal bar indicated the mean. **b,** same as in a but for all antibodies combined.
c, Distribution of the hydrophobicity GRAVY scores at the IGH CDR3 in antibody sequences
from this study compared to a public database (see Methods).

Extended Data Figure 9. Binding of the monoclonal antibodies to the RBD of SARS-CoV-2

and SARS-CoV. a, EC_{50} values for binding to the RBD of SARS-CoV-2. **b and c,** Binding curves

and EC_{50} values for binding to the RBD of SARS-CoV.

Extended Data Figure 10. Biolayer interferometry experiment. showing binding of antibodies

C144, C101, C002, C121, C009, C019 (see also main text Fig. 4). Graphs show secondary

antibody binding to preformed C121 IgG-RBD complexes. The table displays the shift in

nanometers after second antibody (Ab2) binding to the antigen in the presence of the first antibody

(Ab1). Values are normalized by the subtraction of the autologous antibody control.

References

630 1   Graham, R. L., Donaldson, E. F. & Baric, R. S. A decade after SARS: strategies for
631   controlling emerging coronaviruses. *Nat Rev Microbiol* **11**, 836-848,
632   doi:10.1038/nrmicro3143 (2013).
633 2   Gralinski, L. E. & Baric, R. S. Molecular pathology of emerging coronavirus infections. *J*
634   *Pathol* **235**, 185-195, doi:10.1002/path.4454 (2015).
635 3   Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is
636   Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280 e278,
637   doi:10.1016/j.cell.2020.02.052 (2020).
638 4   Walls, A. C. *et al.* Structure, Function, and Antigenicity of the SARS-CoV-2 Spike
639   Glycoprotein. *Cell* **181**, 281-292 e286, doi:10.1016/j.cell.2020.02.058 (2020).
640 5   Jiang, S., Hillyer, C. & Du, L. Neutralizing Antibodies against SARS-CoV-2 and Other
641   Human Coronaviruses. *Trends Immunol*, doi:10.1016/j.it.2020.03.007 (2020).
642 6   Scheid, J. F. *et al.* Broad diversity of neutralizing antibodies isolated from memory B
643   cells in HIV-infected individuals. *Nature* **458**, 636-640, doi:10.1038/nature07930 (2009).
644 7   Tiller, T. *et al.* Autoreactivity in human IgG+ memory B cells. *Immunity* **26**, 205-213,
645   doi:10.1016/j.immuni.2007.01.009 (2007).
646 8   Murugan, R. *et al.* Clonal selection drives protective memory B cell responses in
647   controlled human malaria infection. *Sci Immunol* **3**, doi:10.1126/sciimmunol.aap8029
648   (2018).
649 9   Briney, B., Inderbitzin, A., Joyce, C. & Burton, D. R. Commonality despite exceptional
650   diversity in the baseline human antibody repertoire. *Nature* **566**, 393-397,
651   doi:10.1038/s41586-019-0879-y (2019).
652 10   ter Meulen, J. *et al.* Human monoclonal antibody combination against SARS coronavirus:
653   synergy and coverage of escape mutants. *PLoS Med* **3**, e237,
654   doi:10.1371/journal.pmed.0030237 (2006).
655 11   Yuan, M. *et al.* A highly conserved cryptic epitope in the receptor binding domains of
656   SARS-CoV-2 and SARS-CoV. *Science* **368**, 630-633, doi:10.1126/science.abb7269
657   (2020).
658 12   Walls, A. C. *et al.* Unexpected Receptor Functional Mimicry Elucidates Activation of
659   Coronavirus Fusion. *Cell* **176**, 1026-1039 e1015, doi:10.1016/j.cell.2018.12.028 (2019).
660 13   Pinto, D. *et al.* Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV
661   antibody. *Nature*, doi:10.1038/s41586-020-2349-y (2020).
662 14   Zhu, Z. *et al.* Potent cross-reactive neutralization of SARS coronavirus isolates by human
663   monoclonal antibodies. *Proc Natl Acad Sci U S A* **104**, 12123-12128,
664   doi:10.1073/pnas.0701000104 (2007).
665 15   Salazar, G., Zhang, N., Fu, T. M. & An, Z. Antibody therapies for the prevention and
666   treatment of viral infections. *NPJ Vaccines* **2**, 19, doi:10.1038/s41541-017-0019-3
667   (2017).
668 16   Bournazos, S. & Ravetch, J. V. Anti-retroviral antibody FcgammaR-mediated effector
669   functions. *Immunol Rev* **275**, 285-295, doi:10.1111/imr.12482 (2017).
670 17   Feinberg, M. B. & Ahmed, R. Advancing dengue vaccine development. *Science* **358**,
671   865-866, doi:10.1126/science.aaq0215 (2017).
672 18   Iwasaki, A. & Yang, Y. The potential danger of suboptimal antibody responses in
673   COVID-19. *Nat Rev Immunol*, doi:10.1038/s41577-020-0321-6 (2020).

19 Van Rompay, K. K. A. *et al.* A combination of two human monoclonal antibodies limits
fetal damage by Zika virus in macaques. *Proc Natl Acad Sci U S A* **117**, 7981-7989,
doi:10.1073/pnas.2000414117 (2020).
20 Plotkin, S. A. Correlates of protection induced by vaccination. *Clin Vaccine Immunol* **17**,
1055-1065, doi:10.1128/CVI.00131-10 (2010).
21 Scheid, J. F. *et al.* Sequence and structural convergence of broad and potent HIV
antibodies that mimic CD4 binding. *Science* **333**, 1633-1637,
doi:10.1126/science.1207227 (2011).
22 Robbiani, D. F. *et al.* Recurrent Potent Human Neutralizing Antibodies to Zika Virus in
Brazil and Mexico. *Cell* **169**, 597-609 e511, doi:10.1016/j.cell.2017.04.024 (2017).
23 Ehrhardt, S. A. *et al.* Polyclonal and convergent antibody response to Ebola virus vaccine
rVSV-ZEBOV. *Nat Med* **25**, 1589-1600, doi:10.1038/s41591-019-0602-4 (2019).
24 Pappas, L. *et al.* Rapid development of broadly influenza neutralizing antibodies through
redundant mutations. *Nature* **516**, 418-422, doi:10.1038/nature13764 (2014).
25 Kane, M. *et al.* Identification of Interferon-Stimulated Genes with Antiretroviral Activity.
Cell Host Microbe **20**, 392-405, doi:10.1016/j.chom.2016.08.005 (2016).
26 Adachi, A. *et al.* Production of acquired immunodeficiency syndrome-associated
retrovirus in human and nonhuman cells transfected with an infectious molecular clone. *J*
Virol **59**, 284-291 (1986).
27 Wang, Z. *et al.* Isolation of single HIV-1 Envelope specific B cells and antibody cloning
from immunized rhesus macaques. *J Immunol Methods* **478**, 112734,
doi:10.1016/j.jim.2019.112734 (2020).
28 Tiller, T. *et al.* Efficient generation of monoclonal antibodies from single human B cells
by single cell RT-PCR and expression vector cloning. *J Immunol Methods* **329**, 112-124,
doi:10.1016/j.jim.2007.09.017 (2008).
29 von Boehmer, L. *et al.* Sequencing and cloning of antigen-specific antibodies from
mouse memory B cells. *Nat Protoc* **11**, 1908-1923, doi:10.1038/nprot.2016.102 (2016).
30 Klein, F. *et al.* Enhanced HIV-1 immunotherapy by commonly arising antibodies that
target virus escape variants. *J Exp Med* **211**, 2361-2372, doi:10.1084/jem.20141050
(2014).
31 Schoofs, T. *et al.* Broad and Potent Neutralizing Antibodies Recognize the Silent Face of
the HIV Envelope. *Immunity* **50**, 1513-1529 e1519, doi:10.1016/j.immuni.2019.04.014
(2019).
32 Ye, J., Ma, N., Madden, T. L. & Ostell, J. M. IgBLAST: an immunoglobulin variable
domain sequence analysis tool. *Nucleic Acids Res* **41**, W34-40, doi:10.1093/nar/gkt382
(2013).
33 Gupta, N. T. *et al.* Change-O: a toolkit for analyzing large-scale B cell immunoglobulin
repertoire sequencing data. *Bioinformatics* **31**, 3356-3358,
doi:10.1093/bioinformatics/btv359 (2015).
34 Rubelt, F. *et al.* Onset of immune senescence defined by unbiased pyrosequencing of
human immunoglobulin mRNA repertoires. *PLoS One* **7**, e49774,
doi:10.1371/journal.pone.0049774 (2012).
35 Kyte, J. & Doolittle, R. F. A simple method for displaying the hydropathic character of a
protein. *J Mol Biol* **157**, 105-132, doi:10.1016/0022-2836(82)90515-0 (1982).
36 Guy, H. R. Amino acid side-chain partition energies and distribution of residues in
soluble proteins. *Biophys J* **47**, 61-70, doi:10.1016/S0006-3495(85)83877-7 (1985).

37   DeWitt, W. S. *et al.* A Public Database of Memory and Naive B-Cell Receptor
Sequences. *PLoS One* **11**, e0160853, doi:10.1371/journal.pone.0160853 (2016).
38   Mastronarde, D. N. Automated electron microscope tomography using robust prediction
of specimen movements. *J Struct Biol* **152**, 36-51, doi:10.1016/j.jsb.2005.07.007 (2005).
39   Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for
rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296,
doi:10.1038/nmeth.4169 (2017).
40   Goddard, T. D., Huang, C. C. & Ferrin, T. E. Visualizing density maps with UCSF
Chimera. *J Struct Biol* **157**, 281-287, doi:10.1016/j.jsb.2006.06.010 (2007).
729