Shifts from pulled to pushed range expansions caused by reduction of landscape connectivity

Authors

Maxime Dahirel[1], Aline Bertin[1], Marjorie Haond[1], Aurélie Blin[1], Eric Lombaert[1], Vincent Calcagno[1], Simon Fellous[2], Ludovic Mailleret[1,3], Thibaut Malausa[1], Elodie Vercken[1]

[1]Université Côte d'Azur, INRAE, CNRS, ISA, France

[2]CBGP, INRAE, CIRAD, IRD, Montpellier SupAgro, Univ Montpellier, Montpellier, France

[3]Université Côte d'Azur, INRIA, INRAE, CNRS, Sorbonne Université, BIOCORE, Sophia Antipolis, France

Corresponding author: MD, maxime.dahirel@yahoo.fr

Abstract

Range expansions are key processes shaping the distribution of species; their ecological and evolutionary dynamics have become especially relevant today, as human influence reshapes ecosystems worldwide. Many attempts to explain and predict range expansions assume, explicitly or implicitly, so-called "pulled" expansion dynamics, in which the low-density edge populations provide most of the "fuel" for the species advance. Some expansions, however, exhibit very different dynamics, with high-density populations behind the front "pushing" the expansion forward. These two types of expansions are predicted to have different effects on e.g. genetic diversity and habitat quality sensitivity. However, empirical studies are lacking due to the challenge of generating reliably pushed vs. pulled expansions in the laboratory, or discriminating them in the field. We here propose that manipulating the degree of connectivity among populations may prove a more generalizable way to create pushed expansions. We demonstrate this with individual-based simulations as well as replicated experimental range expansions (using the parasitoid wasp *Trichogramma brassicae* as model). By analysing expansion velocities and neutral genetic diversity, we showed that reducing connectivity led to pushed dynamics. Low connectivity alone, i.e. without density-dependent dispersal, can only lead to "weakly pushed" expansions, where invasion speeds conforms to pushed expectations but not genetic diversity. In empirical expansions however, low connectivity may in some cases also lead to adjustments to the dispersal-density function, recreating "classical" pushed expansions. In the current context of habitat loss and fragmentation, we need to better account for this relationship between connectivity and expansion regimes to successfully predict the ecological and evolutionary consequences of range expansions.

Keywords biological control, biological invasions, density-dependent dispersal, individual-based model, range shifts, *Trichogramma*

Introduction

Range expansions and range shifts into novel habitats and landscapes are key ecological processes shaping the abundance and distribution of species (Sexton, McIntyre, Angert, & Rice, 2009). Understanding their ecological and evolutionary drivers and consequences has become especially relevant in the context of more frequent biological invasions (Miller et al., 2020; Renault, Laparie, McCauley, & Bonte, 2018) or increasing impacts of climate change (Hill, Griffiths, & Thomas, 2011; Lenoir et al., 2020).

Range expansions are usually modelled and analysed in a framework based on the Fisher-KPP partial differential equation and its numerous declinations (see e.g. Hastings et al., 2005; Lewis, Petrovskii, & Potts, 2016). In this framework, range dynamics generally converge to solutions of constant profiles moving in space at a fixed velocity, called travelling waves. Travelling waves are considered "pulled" when spread is driven mostly, if not only, by the dynamics at the leading edge of the expansion. In biological range expansions, this happens when growth and dispersal rates are maximal at low densities; the velocity of the wave then only depends on low-density dispersal and growth (e.g. Birzu, Matin, Hallatschek, & Korolev, 2019). While growth and dispersal functions that are expected to generate pulled expansions certainly happen in nature (Harman, Goddard, Shivaji, & Cronin, 2020; Matthysen, 2005; Williams & Levine, 2018), they are not the only ones possible. For instance, growth rates can be reduced at low densities compared to intermediate ones, a phenomenon known as the Allee effect (Berec, Angulo, & Courchamp, 2007). Additionally, dispersal is often maximal at high densities (positive density-dependence) as it provides a mechanism to escape increased competition (Harman et al., 2020; Matthysen, 2005). In both cases, we would expect the advance of the expansion to be driven not by the low-density front populations, but by the population dynamics in a region located behind the front, where growth and/or dispersal are maximal. These "pushed" waves (Stokes, 1976) behave very differently than pulled waves (see e.g. Lewis et al., 2016). They will typically advance faster than expected based solely on growth and dispersal rates observed at low densities (i.e. in edge patches; Gandhi, Korolev, & Gore, 2019). The ratio of the actual expansion velocity to the velocity expected for the corresponding pulled expansion has been proposed as a

quantitative indicator discriminating pulled, "semi-pushed" and pushed expansions (e.g. Birzu,

Hallatschek, & Korolev, 2018). This ratio has been connected to other key metrics expected to

vary along the pushed/pulled gradient (Birzu et al., 2018). In the context of biological range

expansions or shifts, this includes the rate at which the edge or front of the expansion loses

genetic diversity due to drift and successive founding events as it advances. Indeed, models and

experiments show that neutral genetic diversity is lost at the expanding edge much more slowly

in pushed than in pulled expansions (Birzu et al., 2018, 2019; Gandhi et al., 2019; Hallatschek &

Nelson, 2008; Roques, Garnier, Hamel, & Klein, 2012), with estimated times to allele fixation

differing by up to several orders of magnitude (Gandhi et al., 2019).

A key challenge for empirically studying the ecological and evolutionary consequences of

pushed and pulled expansions (which, besides neutral genetic diversity, remain poorly known;

see discussion in Birzu et al., 2019) is to obtain comparable expansions known to be either

pushed or pulled. One could contrast strains, populations or closely related species naturally

differing in density-dependent dispersal and/or Allee effects (Harman et al., 2020; Jacob,

Chaine, Huet, Clobert, & Legrand, 2019; Matthysen, 2005; Walter, Grayson, Blackburn, Tobin, &

Johnson, 2019). However, they are likely to also differ in other key traits (Jacob et al., 2019),

making direct comparisons difficult. A better experimental solution would be to manipulate the

focal species/population/strain's environment to change the presence or strength of Allee

effects and/or density-dependent dispersal. For instance, Gandhi et al. (2019; 2016) managed to

reliably produce pushed expansions by changing the substrate on which *Saccharomyces*

cerevisiae yeasts were grown: while yeasts grow best at low densities on galactose, they present

an Allee effect when they have to use sucrose. Externally manipulating density-dependent

dispersal may be more difficult, but could conceivably also be done by manipulating resource

quality (Endriss et al., 2019; Van Allen & Bhavsar, 2014). Adding or removing conspecific cues

independently of actual population size may also be an option (De Meester & Bonte, 2010), but

not all organisms have resources or cues that are easy and useful to manipulate experimentally

(Fellous, Duncan, Coulon, & Kaltz, 2012).

Rather than manipulating the within-habitat conditions, another solution that may be easier to

generalize would be to reduce the level of connectivity among habitats. In a spatially structured

environment, this could be done experimentally by e.g. altering the number, length or quality of

the physical links between patches/populations. We hypothesize such experimental

manipulations could lead to more pushed expansions because of dispersal stochasticity. Indeed,

at low population sizes, reducing the dispersal rate sharply increases the risk the smaller

populations at the edge of the expansion will fail to send any disperser, due to stochasticity

alone (**Fig. 1**). Reduced dispersal rate could thus increase the influence of population density on

dispersal success, leading to pushed expansions. This effect would occur when populations at

the very edge of the expansions are of few individuals, typically less than a hundred (**Fig. 1**).

Edge population densities low enough to trigger the effect described **Fig. 1** are probably not

uncommon in some taxonomic groups, where core population sizes are in the 100-1000 range

(Krauss, Steffan-Dewenter, & Tscharntke, 2003; Santini, Isaac, & Ficetola, 2018). These are

however much lower densities than those considered in most pushed expansion models and

experiments (e.g. Birzu et al., 2018; Gandhi et al., 2019). The idea that expansions that should

be pulled at asymptotically large carrying capacities can nonetheless appear pushed at lower

densities, at least in terms of velocity dynamics, is not new. It was previously raised by Panja &

van Saarloos (2002) and reviewed/expanded on in Panja (2004), which called these "weakly

pushed" expansions (not to be confused with "semi-pushed" expansions *sensu* Birzu et al.,

2018). "Weakly pushed" expansions arise when we stop assuming continuous fluid-like

population densities can approximate the stochastic dynamics of discrete individuals in

discrete/discretized space. Revisiting earlier range expansion studies including discreteness, we

can indeed find hints that added dispersal stochasticity or decreased connectivity leads to

population patterns resembling those seen in pushed expansions (Morel-Journel et al., 2016;

Reluga, 2016; Williams & Levine, 2018). However, there is still no actual confirmation that

reduced connectivity can produce pushed expansions (even in the "weakly pushed" sense), or

that these expansions would share all key characteristics of pushed expansions that interest

ecologists and evolutionary biologists (such as the dynamics of genetic diversity).

In the present study, we combined simulations and experimental approaches to study the effect

of reduced connectivity on the pushed vs. pulled status of expansions, at population sizes that

are realistic for many "macroscopic" organisms. In individual-based simulations, we examined

expansion velocities and the dynamics of neutral genetic diversity to (a) confirm Allee effects

and density-dependent dispersal (Birzu et al., 2019) still lead to pushed expansions at "low"

equilibrium population sizes K, and (b) show that reducing connectivity can also generate

(weakly) pushed fronts even in the absence of the other two mechanisms. We complemented

this approach by using minute wasps of the genus *Trichogramma* (Haond et al., 2018; Morel-

Journel et al., 2016) in replicated experimental landscapes (Larsen & Hargreaves, 2020), to

investigate whether reduced connectivity influenced the velocity and neutral genetic dynamics

of "real" range expansions.

Figure 1. (A) Probability that at least one individual disperses from a patch ($\Pr(n > 0)$) as a function of its population size N pre-dispersal and average dispersal rate D. (B) Coefficient of variation of the the numbers of dispersers. Dispersal stochasticity is obtained by assuming the number of dispersers n from a patch is drawn from a Binomial distribution $n \sim \mathrm{Binom}(N, D)$. Note the log scale on the x-axes.

Methods

Simulations

To determine whether the new mechanism we propose, reduced connectivity, can actually generate pushed expansions, we used an individual-based model (IBM) approach. The model is in discrete time and space and simulates the dynamics of a haploid clonal species with non-overlapping generations, expanding in one direction in a one-dimensional landscape. Conceptually, the model is inspired by previous models by Birzu et al. (2018, 2019) and Haond et al. (2018), as well as Gandhi et al. (2019, 2016)'s experiments. Range expansions unfold during 100 generations in a landscape long enough that they never run out of empty patches. In practice, given the expansions advance in only one direction and individuals can only disperse at most one patch per generation (see below), this means any landscape length higher than the number of generations. All patches are of equal and constant quality, determined by the equilibrium population density K.

At the start of the expansion, K adult individuals that have not yet reproduced are introduced in one patch at one of the two extremities of the landscape (coordinate x = 0). To be able to later study genetic diversity, all individuals are randomly assigned one of two allele values (coded 0 and 1) at a neutral genetic locus L. Each generation until the end of a run, the life cycle then happens as follows:

(1) Adult individuals disperse with a probability $D = \text{logit}^{-1}(\text{logit}(D_0) + \beta_{density} \times N/K)$, where N is the patch population size immediately before the dispersal phase, D_0 is the (hypothetical) dispersal rate at N = 0, and $\beta_{density}$ the slope of the dispersal-density relationship on the logit scale. This function is based on the way empirical dispersal-density data are usually analysed (through generalized linear models; see e.g. De Meester & Bonte, 2010; Van Allen & Bhavsar, 2014). Dispersers randomly move to one of the nearest neighbouring patches; that is, the maximal dispersal distance is of 1 patch.

169 (2) Reproduction occurs post-dispersal; the number of offspring F each individual produces is drawn from a $\text{Poisson}(\mu)$ distribution. The mean fecundity μ is based on a Ricker equation modified to allow potential Allee effects (Morel-Journel et al., 2016): $\mu = e^{r_0(1-N/K)(1-A/N)}$. In this equation, r_0 is the hypothetical population growth rate at $N = 0$ in the absence of Allee effects, and A an Allee threshold such that $A = 0$ leads to no Allee effects, $0 < A \leq 1$ leads to weak Allee effects (*sensu* e.g. Berec et al., 2007, i.e. where positive density-dependent growth never leads to negative growth rates), and $A > 1$ leads to strong Allee effects (where growth rates are negative for $N < A$). All new individuals inherit the value at the neutral locus L from their parent with no mutation.

(3) All adults die; juveniles then become adults.

Figure 2. Dispersal (A) and fecundity (B) functions used in simulations. The functions differ depending on equilibrium population size K because they are defined in terms of N/K, rather than N. Dots mark the values of dispersal and fecundity used in the definition of the reference velocity v_F.

The model was written in Netlogo (Wilensky, 1999), version 6.1.1, and set up using the *nlrx* R

package (Salecker, Sciaini, Meyer, & Wiegand, 2019). We tested 5 scenarios (× 2 possible values

of K; see below and **Fig. 2**). The "reference" scenario had no Allee effects, no density-

dependent dispersal ($\beta_{density}$ = 0) and a dispersal rate D_0 set to 0.2, a biologically "typical" rate

according to experimental (e.g. Fronhofer et al., 2018) or natural observations (Marjamäki,

Contasti, Coulson, & McLoughlin, 2013; Stevens et al., 2013), spanning many taxa. Three other

scenarios, which we expected to lead to pushed expansions, each differed from the reference

by one parameter: either a weak Allee effect was present (A = 0.95), there was positive-density-

dependent dispersal ($\beta_{density}$ = 1) or connectivity was reduced by half (D_0 = 0.1). The fifth

scenario was a combination of reduced connectivity and density-dependent dispersal ($\beta_{density}$ =

1 and D_0 = 0.1). In all cases r_0 was set to log(5), which is well within the range of plausible values

for insects (Hassell, Lawton, & May, 1976). We tested each scenario for two values of the

equilibrium density K, 225 and 450 individuals per patch, as the relationship between K and the

expansion velocity v is expected to differ between pushed and pulled expansions (Haond et al.,

2018). These densities are within the range of K used in Haond et al. (2018)'s simulations and

experiments, and correspond respectively to 50% and 100% of the largest possible population in

our own experimental expansions (see below). These values of K are also within one order of

magnitude of those seen for some species of butterflies (Krauss et al., 2003) or herbaceous

plants (Dauber et al., 2010) in patchy environments, or in land molluscs (Kappes et al., 2009)

assuming a typical home range of a few m² at most (Bailey, 1989). Each scenario × K

combination was replicated 100 times.

Experimental range expansions

We used laboratory strains of the haplo-diploid egg parasitoid *Trichogramma brassicae*

Bezdenko, 1968 (Hymenoptera: Trichogrammatidae) as our model in experiments

(**Supplementary Material 1**). *T. brassicae* wasps are raised routinely in the lab using eggs of the

Mediterranean flour moth *Ephestia kuehniella* Zeller 1879 (Lepidoptera: Pyralidae) as

substitution host. *E. kuehniella* eggs are irradiated before use, which prevents their larval

development while still allowing *Trichogramma* wasps to use them as hosts (St-Onge, Cormier,

Todorova, & Lucas, 2014). To be able to better generalize our results, we used three

independent genetic mixes of *T. brassicae* (**Supplementary Material 1**). Note that the effect of

genetic background itself is not the object of the present manuscript, and is thus not analyzed

directly in any of our statistical models, for simplicity (as in Van Petegem et al., 2018).

Figure 3. A representative subset of the experimental landscapes, showing the "patches"

connected by 1 or 3 tubes, depending on treatment, as well as clusters of host eggs on paper

strips for easy manipulation. Picture by Aline Bertin. Inset: *Trichogramma brassicae* on *Ephestia*

kuehniella eggs. Picture by Géraldine Groussier, used with authorisation.

To experimentally test the effects of reduced connectivity on range expansions, we set up a

series of one-dimensional artificial landscapes (**Fig. 3**), in which we monitored *T. brassicae*

wasps for 14 non-overlapping generations (including initially released adults, i.e. generation 0).

The landscapes were made of closed plastic vials (5 cm diameter, 10 cm height) connected to

each of their nearest neighbours by either three 20 cm flexible tubes ("reference"-level

connectivity) or one 40 cm tube (reduced connectivity) (tube internal diameter: 5 mm).

Landscapes were initially 5 patches long (including the release patch), to allow dispersal kernels

with movements > 1 patch, and extended as needed to ensure there were always at least 2

available empty patches beyond the current front. Each treatment was replicated 12 times (4

per genetic mix), for a total of 24 experimental landscapes. Landscapes were initiated by

introducing ≈ 300 adult wasps (range: 261 − 324) in one extremity patch (x = 0), so the

expansion could only advance in one direction in each landscape. Landscapes were kept under

standardized conditions (23°C, 70% relative humidity, 16:8 L:D). Each generation then unfolded

along the following steps:

- (a) We provided approximately 450 new eggs of *Ephestia kuehniella* per patch for wasps to lay

eggs in. *E. kuehniella* eggs were presented on paper strips to facilitate handling.

- (b) Adults (and old egg strips) were removed after 48 hours to enforce non-overlapping

generations and standardized generation times.

- (c) When parasitoid larval development was advanced enough to identify signs of parasitism

(host eggs darkening after ≈ 7 days), we temporarily removed egg strips from patches and

assessed the presence/ absence of parasitized eggs by eye. We photographed all patches with

parasitized eggs for semi-automatic estimations of population sizes (Nikon D750 60mm f/2.8D

macro lens, 24.3 Mpix). Population sizes were estimated (as % of host eggs parasitized) using

ImageJ and the Codicount plugin (Abramoff, Magalhães, & Ram, 2004; Perez, Burte, Baron, &

Calcagno, 2017). We used four different macros each trained on a different set of pictures, to

account for "observer" effects; each macro tended to give consistently biased estimates, but

combining them led to on average unbiased estimates (**Supplementary Material 2**). These

population size estimates should be relatively robust to superparasitism, as in the vast majority

of cases only 1 wasp per host survives to this stage, even when multiple eggs have been laid in

the host (Corrigan, Laing, & Zubricky, 1995). Eggs were then replaced in their patches until the

emergence of the adults, when we started a new cycle from step (a).

We confirmed the experimental design led to reduced connectivity by comparing the average

egg-laying distances (in patches from release sites) between treatments at the start of the

experiment, i.e. the only time point when the source patch of all egg-laying individuals was

known. As dispersal is defined as "movement of individuals or propagules with potential

consequences for gene flow across space" (Ronce, 2007), this metric is in effect a measure of

average (effective) dispersal distance. The average distance was reduced by ≈ 20% in "reduced

connectivity" landscapes compared to "reference" ones (**Supplementary Material 3**).

To determine how genetic diversity evolved during experimental range expansions and whether

it was influenced by connectivity, we kept and genotyped adult female wasps after their

removal from the experimental landscapes. In each landscape, we sampled initially released

individuals (i.e. generation 0; hereafter "origin" wasps) as well as wasps found in both the

release patch ("core" wasps) and the most advanced population ("edge" wasps) at generations

4, 8 and 12. In each patch × time combination, we aimed to sample about 20 individuals; the

actual number was sometimes lower especially in edge populations with fewer individuals

(average ± SD: 18.10 ± 2.63 wasps; range: 4-28; total: 3043). Wasps were genotyped at 19

microsatellite loci; microsatellite characteristics, as well as the DNA extraction and amplification

protocols, are detailed in **Supplementary Material 4.**

Statistical analyses

All data (experimental and simulated) were analyzed in a Bayesian framework, using R, version

4.0.3 (R Core Team, 2020) and the *brms* package version 2.14.4 (Bürkner, 2017) as frontends to

the Stan language (RStan version 2.19.3; Carpenter et al., 2017; Stan Development Team, 2018).

We used (non)linear multilevel/mixed models. The model descriptions below are summaries

aiming for clarity rather than completeness; formal and complete write-ups for all statistical

models are given in **Supplementary Material 5**. We use general-purpose "weakly informative"

priors based on McElreath (2020) for all parameters except one where there is strong

theoretical prior knowledge (see **Supplementary Material 5**). For each model, we ran four

chains with enough iterations to ensure effective sizes were satisfactory for all fixed effect,

random effect and distributional parameters (both bulk- and tail-ESS *sensu* Vehtari, Gelman,

Simpson, Carpenter, & Bürkner, 2020 > 1000). In addition to graphical posterior checks (Gabry,

Simpson, Vehtari, Betancourt, & Gelman, 2019; Vehtari et al., 2020), we checked chain

convergence using the improved $\hat{R}$ statistic by Vehtari et al. (2020). All credible intervals given in

text and figures are 95% Higher posterior Density Intervals (HDIs). Data handling and figure

generation were mainly done using the various *tidyverse* packages (Wickham et al., 2019), as

well as the *tidybayes* (Kay, 2019) and *patchwork* (Pedersen, 2019) packages.

Expansion velocity (simulations and experimental data)

In the long run, the position X_t of the front (here using the distance between the farthest populated patch and the release site) is expected to increase linearly with time t, whether expansions are pushed or pulled (e.g. Lewis et al., 2016):

$X_t = vt,$

where v is the asymptotic expansion velocity. However, expansions only settle on the velocity v after a transition period; the above equation is likely ill-suited for estimating front positions when expansions are followed during their early stages and population sizes are low (so stochasticity is high). In these cases, which correspond to the present study, we propose to use the following model:

$X_t = v_t t,$

$v_t = v + (v_1 - v) \times e^{-\lambda_{[v]}(t-1)},$

where v_1 is the initial velocity after 1 generation of expansion (we use v_1 as our "starting point", and thus $t - 1$, because estimating velocities at $t = 0$ leads to convergence problems), and $\lambda_{[v]}$ the rate of exponential convergence to the asymptotic velocity v.

We fitted this model to both simulated and experimental front locations, assuming a lognormal distribution to account for the fact distances travelled are always strictly positive. We note that one could assume a power-law decay, rather than an exponential one (Panja, 2004). We found however that a power-law model performed either much worse (simulated data) or similarly (experimental data) than the exponential one, and thus used the exponential decay in both cases (see **Supplementary Material 6**, as well as the companion script for detailed model comparisons, **Data availability**).

For experimental data, the submodels for $\log(v_1)$, $\log(v)$ and $\log_{10}(\lambda_{[v]})$ included fixed effects for treatment as well as random effects of replicate identity. For simulated data, we made three slight adjustments. First, we only used data from every fifth generation, to reduce computation time (by about an order of magnitude, based on preliminary tests) without significant impact on

predictive success. Second, the submodel for v used logit(v)instead of log(v), as velocities

were by design ≤ 1 (due to nearest neighbour dispersal). Finally, the model was simplified by

setting v_1 to 1; given the initial population size at t = 0, all simulated landscapes were all but

guaranteed to send at least one individual to the next patch during the first generation, (see **Fig.**

1), leading to a velocity ≈ 1 patch/generation at t = 1.

In simulations, the growth and dispersal functions are fully known (**Fig. 2**), so we were also able

to directly compare the estimated v to their respective v_F, the predicted velocity for a pulled

wave with the same D_0 and r_0 (Birzu et al., 2019; Lewis et al., 2016):

$v_F = 2\sqrt{d_0 r_0},$

where d_0 = 0.5 D_0 for a one-dimensional landscape with nearest neighbour dispersal (see

e.g. Haond et al., 2018). Pulled expansions are expected to have $\frac{v}{v_F}$ = 1, fully pushed expansions

to have $\frac{v}{v_F} > \frac{3}{2\sqrt{2}}$, and so-called semi-pushed expansions to be in-between (Birzu et al., 2018,

2019). Because the formula we use for Allee effects leads to a fecundity of 0 at density = 0, and

because growth and dispersal at density = 0 are not meaningful anyway, we used r_1 and D_1 at N

= 1 to estimate v_F rather than the "true" r_0 and D_0. Note that the formula we use for v_F is

based on the continuum assumption, and should not be exactly valid for discrete systems with

stochasticity in growth, like ours. In fact, we would expect $v_{F[\text{true}]} < v_{F[\text{deterministic}]}$ for such

systems (e.g. Hallatschek & Korolev, 2009). However, some of us previously showed that for a

simulated system like ours, using the simpler continuum formula for v_F is good enough, with an

expected bias of ≈ 1% for 200 ≲ K ≲ 500 (Haond et al., 2018). This is much less than the

smallest difference in speeds between pulled and pushed expansions (≈ 6%, because $\frac{v}{v_F} \geq \frac{3}{2\sqrt{2}}$),

and should therefore not affect our decision to classify a type of expansion as pushed or pulled.

Relationship between equilibrium population density K and velocity v

It has been demonstrated that, over the range of densities we consider in our experiments and

simulations, the correlation between K and the asymptotic velocity v could be another

indicator of pushed expansions (Haond et al., 2018). Pushed expansions would show a positive

correlation between K and v, while pulled expansions should in theory show no correlation.

However, this indicator cannot be considered alone, as demographic stochasticity can also

create such a correlation (Brunet & Derrida, 1997; Hallatschek & Korolev, 2009). The main

difference seems to be that K-v correlations created by demographic stochasticity alone are

associated with $\frac{v}{v_F} < 1$ (Brunet & Derrida, 1997; Hallatschek & Korolev, 2009), while $\frac{v}{v_F} > 1$ for

pushed expansions (see above).

In simulations, we simply compared the posterior distributions of v at K = 225 and K = 450, as

K was fixed.

In experiments, we used the fact that K and v spontaneously varied among replicates, and

analysed among-replicate correlations using a bivariate multilevel model. This allowed us to

estimate correlations accurately while accounting for various sources of uncertainty, including

observer/macro error for population sizes (**Supplementary Material 5**). We first started by

designing univariate models. The univariate model for velocities v is the one described in the

previous section. For K, we used estimated population sizes in the starting patch (x = 0), as this

was the only patch we expected to be at equilibrium from the start, based on release densities

(Morel-Journel et al., 2016). We initially assumed the estimated percentage of hosts identified

as "parasitized" could be analysed using a Beta model, with fixed effects of experimental

treatment and random effects of replicate landscape and computer macro, the latter to account

for consistent macro-level bias (**Supplementary Material 5**). However, we found using posterior

checks (Gabry et al., 2019) that this model failed to accurately represent data distribution and

variability. We then used instead a Student t model on logit-transformed percentages, with the

same fixed and random effects; this performed better (see companion code, **Data availability**).

Following this, we fitted a bivariate model using the two selected models for front position and

population sizes. Replicate-level random effects for front parameters and for K were combined

in the same variance-covariance matrix; this covariance matrix was estimated separately for

each treatment. This allowed us to obtain treatment-specific posteriors for the replicate-level

correlation between K and v.

Genetic diversity, experimental data

When expected heterozygosity (Nei, 1973) H is used as a measure of genetic diversity, theory predicts the genetic diversity both in core patches and at the edge of a range expansion should decay exponentially with time t (Birzu et al., 2018; Hallatschek & Nelson, 2008):

$$H_t = H_0 \times e^{-\lambda_{[g]} \times t},$$

where H_0 is the initial heterozygosity, and $\lambda_{[g]}$ the rate of decay of genetic diversity. The decay rate $\lambda_{[g]}$ is proportional to the inverse of the effective population size N_e, with the exact relationship depending on ploidy (see e.g. Coop, 2020). For experimental data, we used this equation directly in a non-linear model to estimate whether the dynamics of genetic diversity varied between our two treatments and between core and edge patches. Multilocus expected heterozygosity was first calculated for each location (core/edge) × time combination using microsatellite data and the *adegenet* package (Jombart, 2008). Our submodel for $\lambda_{[g]}$ included fixed effects for treatment, location (initial release patch for "core"/ most advanced patch for "edge") and their interactions, as well as a random effect of replicate identity. The submodel for the initial diversity H_0 only included the random effect of replicate, as we did not expect differences between treatments beyond random fluctuations, and core/edge patches are the same at $t = 0$. Location was included in the $\lambda_{[g]}$ submodel as a centred dummy variable (-0.5 for "core", 0.5 for "edge") set to 0 at $t = 0$ as, again, core/edge patches are the same at $t = 0$. The submodels were estimated on logit(H_0) and $\log_{10}(\lambda_{[g]})$ to keep them within proper bounds; expected heterozygosities are proportions, and the decay rate $\lambda_{[g]}$ must be positive. We fitted the overall model on logit(H_t) using a Student t distribution, rather than on H_t using a Beta distribution (and a logit link). This is because the former is likely to be more robust to rare outliers (Kruschke, 2015), like those caused by sampling effects before genotyping (see companion script for detailed model comparisons, **Data availability**).

Genetic diversity, simulated data

With simulated data, we cannot use the equation above as our basis to fit a model. This is because the way we simulate neutral genetic diversity (one locus with only two alleles in a

haploid species) means values of H_t = 0 are very frequent (especially as t increases), and our

previous model cannot handle data containing zeroes. We instead used the fact that, with two

alleles and if a given treatment is replicated a large number of times, a version of $\lambda_{[g]}$ can also

be recovered from the way among-replicate variance in allelic frequencies V (for either allele)

changes with time (e.g. Gandhi et al., 2019):

$V = V_{max}(1 - e^{-\lambda_{[g]} \times t})$,

where V_{max} is the asymptotic variance reached when all replicates have fixed one of the alleles,

which is equal to the product of initial allelic frequencies. As for experimental data, our

submodel for $\log_{10}(\lambda_{[g]})$ included fixed effects for treatment, location (core/edge) and their

interactions. The submodel for $\text{logit}(V_{max})$ included only a constant intercept, as V_{max} should

be identical between all cases (and ≈ 0.25) but for random sampling fluctuations. We fitted this

model on all data with t > 0 using a Beta distribution, as the issues raised for experimental data

above regarding outliers did not apply (each data point here being the summarized outcome of

100 independent populations).

Results

Expansion velocity v and correlation with K

In simulations, absolute asymptotic expansion velocity v differed between treatments. Density-

dependent dispersal, when alone, led to higher velocities than in reference expansions, while

Allee effects and reduced connectivity (with or without density-dependent dispersal) led to

slower expansions (**Fig. 4**, **Supplementary Materials 6,7**). Combining reduced connectivity and

density-dependent dispersal led to velocities closer to the reference than reduced connectivity

alone (**Fig. 4**, **Supplementary Materials 6,7**). Relative velocities $\frac{v}{v_F}$ in the reference landscapes

were very close to those expected for pulled expansions (1.01 and 0.99 for K = 450 and 225

respectively, **Fig. 4**). The other four treatments all had higher velocity ratios and were firmly in

the range corresponding to true pushed expansions based on Birzu et al. (2018) (average

velocity ratios $\frac{v}{v_F} \geq \frac{3}{2\sqrt{2}}$, **Fig. 4**). Expansions were faster when K = 450 than when K = 225 for all

five treatments (**Fig. 4, Supplementary Materials 6,7**).

Figure 4. Posterior distribution of the average asymptotic expansion velocity (v) in simulations,

depending on simulation scenario and equilibrium population size K (black dots: posterior

means, The posterior is in all cases narrower than the dot used to depict the mean). The

horizontal lines mark the range between $\frac{v}{v_F} = 1$ (i.e. pulled expansions) and $\frac{v}{v_F} = \frac{3}{2\sqrt{2}}$ (limit

between semi-pushed and fully pushed expansions; Birzu et al., 2018, 2019). Light grey dots are

observed speeds at the end of each simulation run (t = 100).

In experiments, absolute estimated asymptotic velocities were virtually indistinguishable

between reference and "reduced connectivity" landscapes (**Fig. 5**). There was no correlation

between replicate-level estimates of K and v in "reference" landscapes (r = 0.04 [-0.47; 0.57],

Fig. 6), while there was a positive correlation when connectivity was reduced (r = 0.51 [0.02;

0.92], **Fig. 6**). One must note however that credible intervals are wide, meaning we cannot say

with certainty that the two correlations are different (Δ = 0.47 [-0.22; 1.20]). There was also no

clear evidence that mean population sizes in the "core" starting patch differed between

treatments, with 59.6% [44.6; 72.7] of hosts successfully parasitised on average in "reference"

landscapes vs 63.7% [50.4; 77.0] in "reduced connectivity" landscapes (based on univariate

model; Δ = -4.07% [-9.36; 1.39])(**Fig. 6A**).

Figure 5. (A) Front locations as a function of the number of generations since release and experimental landscape type. Posterior means and 95% credible bands are displayed along observed trajectories for each landscape. (B) Posterior average asymptotic velocity v as a function of landscape type. Dots are posterior means, vertical bars 95% credible intervals.

Figure 6. Correlation between population size in the core K and asymptotic velocity v in experiments. (A) Replicate-level predictions for K and v as a function of treatment (see text for comparison of means). (B) Posterior distribution of the K-v correlation coefficient (latent-scale) as a function of treatment. In both subplots dots are posterior means, bars 95% credible intervals.

Shifts in genetic diversity

In simulations, genetic diversity declined faster, as measured by the decay rate λ, in edge patches than in core patches for all treatments (**Fig. 7**, **Supplementary Materials 6,7**). In edge patches, the addition of density-dependent dispersal and Allee effects led to a slower rate of diversity loss compared to the reference (**Fig. 7**, **Supplementary Materials 6,7**). By contrast, reducing connectivity led to a faster rate of diversity loss (**Fig. 7**, **Supplementary Materials 6,7**), but only when there was no density-dependent dispersal at the same time. Core patch dynamics were much less variable: however, when a treatment did deviate from the reference,

it mostly did so in the same direction as in edge patches (see e.g. "density-dependent dispersal"

or "reduced connectivity" at K = 225, or "reduced connectivity + density dependent dispersal"

at K = 450; **Fig. 7**, **Supplementary Materials 6,7**). In all five treatments and whether we looked

at core or edge patches, genetic diversity was lost faster when K = 225 than when K = 450 (**Fig.**

7, **Supplementary Materials 6,7**).

Figure 7. Posterior distribution of the average decay rate of genetic diversity with time (λ) in

simulations, depending on simulation scenario, equilibrium population size K and patch location

(either the most advanced patch at the time of measure (edge, A), or the original release site

(core, B)). Dots are posterior means, vertical bars 95% credible intervals. Please note that

posteriors for core and edge patches are displayed on different scales on the y-axis, for

readability.

Figure 8. (A) Observed (points) and predicted (lines and bands) genetic diversity as a function of generations since release and landscape type. For predictions, lines are the posterior means and bands correspond to the 95% credible intervals. (B) Posterior distribution of the average genetic diversity decay rate λ as a function of landscape type and patch location (either the original release site (core) or the most advanced patch at the time of measure (edge)). Dots are posterior means, vertical bars 95% credible intervals.

In the experimental landscapes, genetic diversity also decayed on average in all tested contexts of our experiment (**Fig. 8**), and we also found differences between edge and core patches and between both treatments. The rate of genetic diversity loss did not differ clearly between treatments in core patches (mean $\Delta_{\text{reference}-\text{reduced}}$ = - 6.5 × 10^{-3}, 95% CI : [-15.5 × 10^{-3}, 2.1 × 10^{-3}]), but diversity was lost more rapidly in reference edge patches than in "reduced connectivity" ones (mean $\Delta_{\text{reference}-\text{reduced}}$ = 18.3 × 10^{-3}, 95% CI : [1.8 × 10^{-3}, 35.1 × 10^{-3}]) (**Fig. 8**, **Supplementary Material 6,7**). In addition, while edge patches lost diversity faster than core patches in reference landscapes, (mean $\Delta_{\text{edge}-\text{core}}$ = 40.6 × 10^{-3}, 95% CI : [27.7 × 10^{-3}, 53.7 × 10^{-3}]), this difference was still present but reduced when connectivity was reduced (mean $\Delta_{\text{edge}-\text{core}}$ = 15.8 × 10^{-3}, 95% CI : [8.0 × 10^{-3}, 24.3 × 10^{-3}]) (**Fig. 8**).

Discussion

By combining an individual-based model and replicated range expansions in experimental landscapes, we showed that reducing connectivity can lead to more pushed expansions. Discrepancies between the simulation and experimental results, as well as comparisons between the different simulation treatments, potentially shed light on the mechanisms at play in each context.

Expansion velocity as a quantitative and qualitative indicator of pushed expansions

First, our simulation results largely confirm previous theoretical and empirical results regarding the effects of density-dependent dispersal and Allee effects (Birzu et al., 2018, 2019; Gandhi et al., 2019, 2016; Haond et al., 2018; Roques, Garnier, Hamel, & Klein, 2012). We show that both mechanisms lead to expansions that can be classified as "pushed" based on the velocity ratio $\frac{v}{v_F}$ (**Fig. 2**), i.e. advancing faster than expected based on dispersal and growth at low densities. We also show that in the reference scenario, v can be lower than v_F, which is expected as a result of demographic stochasticity (Brunet & Derrida, 1997; Hallatschek & Korolev, 2009), although the difference is in our case minimal (Haond et al., 2018).

In simulations, reducing connectivity led to slower expansions (lower absolute speed v) that

were also pushed (higher velocity ratio $\frac{v}{v_F}$) (**Fig. 4**), confirming our main prediction. Although it

was not presented in connectivity terms, a similar result was recently found using a model

inspired by phage-bacteria interactions (Hunter, Liu, Möbius, & Fusco, 2020), where reduced

connectivity was *de facto* achieved using increased densities of phage-resistant bacteria as a

barrier. In our own study, these pushed expansions are characterised by a mean dispersal rate

independent of population density; this means they would likely converge to a pulled wave as K

tends to infinity and the effects of discreteness and stochasticity become negligible. As such,

they should be considered a type of "weakly" pushed expansions (sensu Panja, 2004; Panja &

van Saarloos, 2002). Nonetheless, they still advance faster than one would expect if they were

pulled; further work is needed to understand the consequences of ignoring this when

attempting to forecast biological invasions or climate-induced range shifts in fragmented

environments.

We also confirm that, over the range of K we studied, increasing the equilibrium population size

K leads to faster range expansions, as previously shown by Haond et al. (2018) for pushed

expansions. Interestingly, this K-v relationship is seen in all simulated treatments, even the

reference one. This may indicate that reference dispersal rates are already low enough to cause

weakly pushed expansions. More likely, given the $\frac{v}{v_F}$ ratio of reference expansions is ≈1 or even

<1 (**Fig. 4**), this may indicate that, in their case, the K-v relationship stems here from

demographic (Hallatschek & Korolev, 2009), rather than dispersal stochasticity.

In experimental landscapes, contrary to simulations, expansions advanced just as fast when

connectivity was reduced (**Fig. 5**). Importantly, it does not mean that reducing connectivity does

not lead to more pushed expansions, on the contrary. What differentiates between pushed and

pulled expansions is not absolute velocity v (which can be either higher or lower in pushed

expansions, **Fig. 4** ; Gandhi et al., 2019), but the ratio to the "equivalent" pulled wave's velocity,

so $\frac{v}{v_F}$. While we do not have direct quantitative information on the v_F of our experimental

landscapes, we have some partial and indirect qualitative information. Indeed, a reduction of

connectivity and thus dispersal implies by definition a reduction of v_F, growth being equal (see

also **Supplementary Material 3**). If $v_{[\text{reference}]} \simeq v_{[\text{reduced}]}$ (**Fig. 5**) and $v_{F[\text{reference}]} >$

$v_{F[\text{reduced}]}$, then it follows that $\frac{v_{[\text{reference}]}}{v_{F[\text{reference}]}} < \frac{v_{[\text{reduced}]}}{v_{F[\text{reduced}]}}$. This would mean that "reduced

connectivity" expansions are indeed more pushed than reference expansions in our experiment.

We acknowledge that this conclusion hinges on growth rates being similar between connectivity

treatments; they should be, as wasps came from the same stock populations, but evolution

during expansions cannot be excluded *a priori* (Van Petegem et al., 2018). Our interpretation

that reduced connectivity leads to pushed expansions in experiments is further supported by

the presence of a K-v correlation that is absent in the "reference" treatment (**Fig. 6**)(Haond et

al., 2018) (but see simulation results above). More studies are needed to confirm these results

as both our velocities and our correlations are estimated with non-negligible uncertainty, due to

limited sample sizes. Nonetheless, our data show that it is possible, in principle, to make

qualitative assessments of expansion "pushiness" relative to a reference, even without

knowledge of the underlying dispersal or growth functions. This is very interesting for the study

of biological invasions and range shifts in the wild, especially in (larger) organisms that are not

easily reared and maintained in laboratory conditions. More work is needed to determine

whether more quantitative insights can also be obtained from incomplete life-history and

population dynamics data typical of "natural" study systems.

Mismatches to genetic diversity predictions show pushed

expansion are diverse

A key theoretical prediction is that pushed expansions lose genetic diversity at the edge more

slowly than comparable pulled expansions (Birzu et al., 2019; Roques et al., 2012). While our

simulations with density-dependent dispersal or Allee effects conform to this prediction (**Fig. 7**),

this is not the case for the pushed expansions we generated by reducing connectivity alone:

these expansions lost genetic diversity slightly faster than the pulled reference (**Fig. 7**) despite

being clearly "pushed" based on velocity ratios. On the one hand, this is unsurprising, as we

should expect a negative relationship between genetic diversity at expansion edges and

landscape fragmentation (e.g. Hill, Hughes, Dytham, & Searle, 2006; Mona, Ray, Arenas, &

Excoffier, 2014); less connectivity means new populations are founded by fewer individuals,

leading to stronger founder effects and drift. On the other hand, this creates an apparent

conflict between the genetic and velocity-based definitions of pushed expansions, which were

implied to be intrinsically correlated (e.g. Birzu et al., 2018). We suggest that, although "weakly"

pushed expansions can behave like "classical" pushed expansions with respect to speed, they

are actually qualitatively different, and we should not expect all theoretical predictions based

on "classical" pushed expansions to apply to them. Further research is thus needed to better

pinpoint where "classical" and "weakly" pushed expansions diverge or converge in their

dynamics. Setups using natural inter-individual variation in dispersal (Saastamoinen et al., 2018;

Schreiber & Beckman, 2020; Stevens, Pavoine, & Baguette, 2010) to decouple the effects of

reduced connectivity itself from those of increased dispersal variability might be particularly

useful here.

In experimental landscapes, reducing connectivity did slow the decay of genetic diversity at the

edge of the expansion (**Fig. 8**). Given population genetics models predict a faster loss of diversity

with increased fragmentation when expansions are not pushed (Mona et al., 2014), this result is

strong evidence that our experiments did generate (more) pushed expansions, this time in the

"classical" sense. On the other hand, experiments are here in direct contradiction with our

simulation results. This mismatch may indicate increased dispersal stochasticity is not the only

mechanism at play here.

Based on simulation results (**Fig. 7**), we must first consider the possibility that experimental

landscapes with reduced connectivity lose genetic diversity more slowly simply because they

exhibit larger core populations, rather than due to a shift in regime from (more) pulled to

(more) pushed expansions. There is however no clear evidence that changes in connectivity led

to changes in equilibrium population sizes (see Results), so we do not believe this is the main

driver of our genetic diversity results. In addition, even if there were evidence of changes in K,

this would have led to differences in genetic diversity not only in edge but also in core

populations, which we did not observe (**Fig. 8**).

From a dispersal ecology standpoint, one of the most obvious differences between our

simulations and experiments is that we did not allow simulated individuals to adjust their

dispersal-density decision rules as a function of their environment. Dispersal decisions are

multicausal, with many factors interacting to shape the benefits-cost balance of dispersal

(Matthysen, 2012). For instance, emigration by *Notonecta undulata* backswimmers is density-

dependent in parasite-free ponds, but density-independent in mite-infected ponds (Baines,

Diab, & McCauley, 2020). In *Erigone atra* spiders, dispersal by rappelling depends on the

interaction of sex, density and local sex-ratios (De Meester & Bonte, 2010). In *Tribolium*

castaneum beetles, dispersal is less density-dependent out of lower-quality habitat (Endriss et

al., 2019; Van Allen & Bhavsar, 2014). In our context, one might wonder how reduced

connectivity and density-dependent dispersal might interact. Theory predicts that density-

dependent emigration should be more advantageous when connectivity is reduced (more

precisely, when costs of dispersal are higher; Travis, Murrell, & Dytham, 1999; Rodrigues &

Johnstone, 2014). In line with these predictions, dispersal was density-independent in *Tribolium*

castaneum metapopulations with low connectivity, density-dependent in metapopulations with

high connectivity (Govindan, Feng, DeWoody, & Swihart, 2015). If we want to see whether

simulations and experiments actually agree, these previous studies suggest we should actually

compare our experimental treatments to the "reference" vs "reduced connectivity + density-

dependent dispersal" pair of treatments in our simulations, rather than the "reference" vs "low

connectivity alone" pair. Confirming this hypothesis, the "reduced connectivity + density-

dependent dispersal" simulated expansions are a better match to the experiments in that (i)

while pushed, their absolute speed v is closer to reference expansions, and (ii) they do lose

genetic diversity slower than reference expansions. For this mechanism to plausibly apply to our

experimental system, *Trichogramma* wasps need to be able to detect when density and

competition increase locally, and adjust their spatial behaviour in response, which evidence

suggests they do (Wajnberg, Fauvergue, & Pons, 2000). Importantly, the hypothesis that low

connectivity drives changes in the density-dependence of dispersal itself, leading to "true" and

not simply "weakly" pushed expansions, does not imply these changes result from evolution

during the expansion. Indeed, in at least four out of the five experiments cited above, the

context-specific adjustments to the dispersal-density function involved plastic responses (Baines

et al., 2020; De Meester & Bonte, 2010; Endriss et al., 2019; Van Allen & Bhavsar, 2014). Further

experiments under standardised densities and connectivity are needed to determine whether our results are explained by plastic adjustments or evolutionary changes.

Conclusion

In this study, we demonstrated a new mechanism for generating pushed(-like) expansions (**Fig. 1**). We showed that the genetic and velocity aspects of the pushed/pulled distinction, which have been assumed to be tightly linked (Birzu et al., 2018) can be decoupled when (weakly) pushed expansions are caused solely by reduced connectivity. We also confirm pushed expansions can be detected and analysed in systems with (relatively) low population sizes, even though theory has mostly been developed on much higher population sizes. This means the pushed-pulled framework may be valuable to help understand and better predict range expansions and shifts in a broad range of taxa and contexts (including under natural conditions). As ecologists and evolutionary biologists are becoming more aware and more interested in the potential effects of pushed expansions (Miller et al., 2020; Williams, Hufbauer, & Miller, 2019), our results suggest we must be careful about which questions we ask, and whether the dimensions of "pushiness" we collect data on are the appropriate ones. For instance, assuming an expansion loses less genetic diversity simply because its measured ratio is higher may not always be appropriate (**Figs 4, 7**). This increased focus on ecological pushed expansions is nonetheless welcome, as our results highlight more studies are needed to better understand how their impacts depend on the underlying causal mechanisms.

Our experiment shows that, in some cases, reducing connectivity may limit the loss of genetic diversity without impeding spread rates. Founder effects and low genetic diversity are associated with lower adaptive potential and lower chances of population persistence (e.g. Szűcs, Melbourne, Tuff, Weiss-Lehman, & Hufbauer, 2017). Our results thus lead to the somewhat counterintuitive conclusion that reducing connectivity might, in some cases, help expanding species. This conclusion has strong implications for the management of invasive species and the conservation of species undergoing climate-induced range shifts, as one of the main causes of pushed expansions, positive density-dependent dispersal, is frequent in nature (Harman et al., 2020; Matthysen, 2005). Finally, density-dependent dispersal (Fronhofer, Gut, &

Altermatt, 2017; Travis, Mustin, Benton, & Dytham, 2009) and Allee effects (Datta, Korolev, Cvijovic, Dudley, & Gore, 2013; Erm & Phillips, 2020) themselves may evolve during range expansions, and the effects of habitat fragmentation on dispersal ecology and evolution are abundantly documented (Cote et al., 2017; Jacob, Laurent, Morel-Journel, & Schtickzelle, 2020). For these reasons, we call for more systematic eco-evolutionary studies of context-dependent dynamics during range expansions and shifts.

Acknowledgments

We thank participants to the 2019 conference of the British Ecological Society for their helpful comments, as well as Géraldine Groussier for letting us use her picture of *Trichogramma brassicae*.

Data availability

Data, Netlogo model code and R scripts to reproduce all analyses presented in this manuscript are available on GitHub (https://github.com/mdahirel/pushed-pulled-2020-dynamics) and Zenodo (doi: 10.5281/zenodo.3969988).

Funding

This work was funded by the French Agence Nationale de la Recherche (TriPTIC, ANR-14-CE18-0002; PushToiDeLa, ANR-18-CE32-0008), and received funding from the European Union Seventh Framework Programme FP7 (grant agreement FP7-IAPP #324475 "COLBICS").

Conflicts of interest disclosure

The authors declare they have no financial conflict of interest in relation with the content of this article. Several authors are recommenders for PCI (PCI Evol Biol: EL, VC, SF, TM, EV; PCI Ecology and PCI Zoology: EL, VC, EV).

References

Abramoff, M. D., Magalhães, P. J., & Ram, S. J. (2004). Image processing with ImageJ. *Biophotonics International*, *11*(7), 36–42. Retrieved from http://dspace.library.uu.nl/handle/1874/204900

Bailey, S. E. R. (1989). Foraging behaviour of terrestrial gastropods: integrating field and laboratory studies. *Journal of Molluscan Studies*, *55*(2), 263–272. doi: 10.1093/mollus/55.2.263

Baines, C. B., Diab, S., & McCauley, S. J. (2020). Parasitism risk and infection alter host dispersal. *The American Naturalist*, *196*(2), 119–131. doi: 10.1086/709163

Berec, L., Angulo, E., & Courchamp, F. (2007). Multiple Allee effects and population management. *Trends in Ecology & Evolution*, *22*(4), 185–191. doi: 10.1016/j.tree.2006.12.002

Birzu, G., Hallatschek, O., & Korolev, K. S. (2018). Fluctuations uncover a distinct class of traveling waves. *Proceedings of the National Academy of Sciences*, *115*(16), E3645–E3654. doi: 10.1073/pnas.1715737115

Birzu, G., Matin, S., Hallatschek, O., & Korolev, K. S. (2019). Genetic drift in range expansions is very sensitive to density dependence in dispersal and growth. *Ecology Letters*, *22*(11), 1817–1827. doi: 10.1111/ele.13364

Brunet, E., & Derrida, B. (1997). Shift in the velocity of a front due to a cutoff. *Physical Review E*, *56*(3), 2597–2604. doi: 10.1103/PhysRevE.56.2597

Bürkner, P.-C. (2017). brms: an R package for Bayesian multilevel models using Stan. *Journal of Statistical Software*, *80*(1), 1–28. doi: 10.18637/jss.v080.i01

Carpenter, B., Gelman, A., Hoffman, M. D., Lee, D., Goodrich, B., Betancourt, M., … Riddell, A. (2017).
Stan: a probabilistic programming language. *Journal of Statistical Software*, *76*(1), 1–32. doi:
10.18637/jss.v076.i01

Coop,   G.   (2020).   *Population   and   quantitative   genetics*.   Retrieved   from
https://github.com/cooplab/popgen-notes

Corrigan, J. E., Laing, J. E., & Zubricky, J. S. (1995). Effects of parasitoid to host ratio and time of day of
parasitism on development and emergence of *Trichogramma minutum* (Hymenoptera:
Trichogrammatidae) parasitizing eggs of *Ephestia kuehniella* (Lepidoptera: Pyralidae). *Annals of*
the Entomological Society of America, *88*(6), 773–780. doi: 10.1093/aesa/88.6.773

Cote, J., Bestion, E., Jacob, S., Travis, J., Legrand, D., & Baguette, M. (2017). Evolution of dispersal
strategies and dispersal syndromes in fragmented landscapes. *Ecography*, *40*(1), 56–73. doi:
10.1111/ecog.02538

Datta, M. S., Korolev, K. S., Cvijovic, I., Dudley, C., & Gore, J. (2013). Range expansion promotes
cooperation in an experimental microbial metapopulation. *Proceedings of the National Academy*
of Sciences, *110*(18), 7354–7359. doi: 10.1073/pnas.1217517110

Dauber, J., Biesmeijer, J. C., Gabriel, D., Kunin, W. E., Lamborn, E., Meyer, B., … Petanidou, T. (2010).
Effects of patch size and density on flower visitation and seed set of wild plants: a pan-European
approach. *Journal of Ecology*, *98*(1), 188–196. doi: https://doi.org/10.1111/j.1365-
2745.2009.01590.x

De Meester, N., & Bonte, D. (2010). Information use and density-dependent emigration in an agrobiont
spider. *Behavioral Ecology*, *21*(5), 992–998. doi: 10.1093/beheco/arq088

Endriss, S. B., Vahsen, M. L., Bitume, E. V., Monroe, J. G., Turner, K. G., Norton, A. P., & Hufbauer, R. A.
(2019). The importance of growing up: juvenile environment influences dispersal of individuals
and their neighbours. *Ecology Letters*, *22*(1), 45–55. doi: 10.1111/ele.13166

Erm, P., & Phillips, B. L. (2020). Evolution transforms pushed waves into pulled waves. *The American*
Naturalist, *195*(3), E87–E99. doi: 10.1086/707324

Fellous, S., Duncan, A., Coulon, A., & Kaltz, O. (2012). Quorum sensing and density-dependent dispersal
in an aquatic model system. *PLoS ONE*, *7*(11), e48436. doi: 10.1371/journal.pone.0048436

Fronhofer, E. A., Gut, S., & Altermatt, F. (2017). Evolution of density-dependent movement during
experimental range expansions. *Journal of Evolutionary Biology*, *30*(12), 2165–2176. doi:
10.1111/jeb.13182

Fronhofer, E. A., Legrand, D., Altermatt, F., Ansart, A., Blanchet, S., Bonte, D., … Cote, J. (2018). Bottom-
up and top-down control of dispersal across major organismal groups. *Nature Ecology &*
Evolution, *2*(12), 1859–1863. doi: 10.1038/s41559-018-0686-0

Gabry, J., Simpson, D., Vehtari, A., Betancourt, M., & Gelman, A. (2019). Visualization in Bayesian
workflow. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*, *182*(2), 389–402.
doi: 10.1111/rssa.12378

Gandhi, S. R., Korolev, K. S., & Gore, J. (2019). Cooperation mitigates diversity loss in a spatially
expanding microbial population. *Proceedings of the National Academy of Sciences*, *116*(47),
23582–23587. doi: 10.1073/pnas.1910075116

Gandhi, S. R., Yurtsev, E. A., Korolev, K. S., & Gore, J. (2016). Range expansions transition from pulled to
pushed waves as growth becomes more cooperative in an experimental microbial population.
Proceedings of the National Academy of Sciences, *113*(25), 6922–6927. doi:
10.1073/pnas.1521056113

Govindan, B. N., Feng, Z., DeWoody, Y. D., & Swihart, R. K. (2015). Intermediate disturbance in
experimental landscapes improves persistence of beetle metapopulations. *Ecology*, *96*(3), 728–
736. doi: 10.1890/14-0044.1

Hallatschek, O., & Korolev, K. S. (2009). Fisher waves in the strong noise limit. *Physical Review Letters*,
103(10), 108103. doi: 10.1103/PhysRevLett.103.108103

Hallatschek, O., & Nelson, D. R. (2008). Gene surfing in expanding populations. *Theoretical Population*
Biology, *73*(1), 158–170. doi: 10.1016/j.tpb.2007.08.008

Haond, M., Morel-Journel, T., Lombaert, E., Vercken, E., Mailleret, L., & Roques, L. (2018). When higher
carrying capacities lead to faster propagation. *bioRxiv (Peer-Reviewed and Recommended by*
Peer Community in Ecology), 307322. doi: 10.1101/307322

Harman, R. R., Goddard, J., Shivaji, R., & Cronin, J. T. (2020). Frequency of occurrence and population-
dynamic consequences of different forms of density-dependent emigration. *The American*
Naturalist, *195*(5), 851–867. doi: 10.1086/708156

Hassell, M. P., Lawton, J. H., & May, R. M. (1976). Patterns of dynamical behaviour in single-species
populations. *Journal of Animal Ecology*, *45*(2), 471–486. doi: 10.2307/3886

Hastings, A., Cuddington, K., Davies, K. F., Dugaw, C. J., Elmendorf, S., Freestone, A., … Thomson, D.
(2005). The spatial spread of invasions: new developments in theory and evidence. *Ecology*
Letters, *8*(1), 91–101. doi: 10.1111/j.1461-0248.2004.00687.x

Hill, J. K., Griffiths, H. M., & Thomas, C. D. (2011). Climate change and evolutionary adaptations at
species' range margins. *Annual Review of Entomology*, *56*(1), 143–159. doi: 10.1146/annurev-
ento-120709-144746

Hill, J. K., Hughes, C. L., Dytham, C., & Searle, J. B. (2006). Genetic diversity in butterflies: interactive
effects of habitat fragmentation and climate-driven range expansion. *Biology Letters*, *2*(1), 152–
154. doi: 10.1098/rsbl.2005.0401

Hunter, M., Liu, T., Möbius, W., & Fusco, D. (2020). Virus-host interactions shape viral dispersal giving
rise to distinct classes of travelling waves in spatial expansions. *arXiv:2009.11357 [Physics]*.
Retrieved from http://arxiv.org/abs/2009.11357

Jacob, S., Chaine, A. S., Huet, M., Clobert, J., & Legrand, D. (2019). Variability in dispersal syndromes is a
key driver of metapopulation dynamics in experimental microcosms. *The American Naturalist*,
194(5), 613–626. doi: 10.1086/705410

Jacob, S., Laurent, E., Morel-Journel, T., & Schtickzelle, N. (2020). Fragmentation and the context-
dependence of dispersal syndromes: matrix harshness modifies resident-disperser phenotypic
differences in microcosms. *Oikos*, *129*(2), 158–169. doi: 10.1111/oik.06857

Jombart, T. (2008). adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics, *24*(11), 1403–1405. doi: 10.1093/bioinformatics/btn129

Kappes, H., Jordaens, K., Hendrickx, F., Maelfait, J.-P., Lens, L., & Backeljau, T. (2009). Response of snails
and slugs to fragmentation of lowland forests in NW Germany. *Landscape Ecology*, *24*(5), 685–
697. doi: 10.1007/s10980-009-9342-z

Kay, M. (2019). *tidybayes: tidy data and geoms for Bayesian models*. doi: 10.5281/zenodo.1308151

Krauss, J., Steffan-Dewenter, I., & Tscharntke, T. (2003). How does landscape context contribute to
effects of habitat fragmentation on diversity and population density of butterflies? *Journal of*
Biogeography, *30*(6), 889–900. doi: 10.1046/j.1365-2699.2003.00878.x

Kruschke, J. (2015). *Doing Bayesian data analysis: a tutorial with R, JAGS, and Stan* (2nd ed.). Boston:
Academic Press.

Larsen, C. D., & Hargreaves, A. L. (2020). Miniaturizing landscapes to understand species distributions.
Ecography, *43*(n/a). doi: 10.1111/ecog.04959

Lenoir, J., Bertrand, R., Comte, L., Bourgeaud, L., Hattab, T., Murienne, J., & Grenouillet, G. (2020).
Species better track climate warming in the oceans than on land. *Nature Ecology & Evolution*.
doi: 10.1038/s41559-020-1198-2

Lewis, M., Petrovskii, S. V., & Potts, J. (2016). *The mathematics behind biological invasions*. Springer
International Publishing. doi: 10.1007/978-3-319-32043-4

Marjamäki, P. H., Contasti, A. L., Coulson, T. N., & McLoughlin, P. D. (2013). Local density and group size
interacts with age and sex to determine direction and rate of social dispersal in a polygynous
mammal. *Ecology and Evolution*, *3*(9), 3073–3082. doi: 10.1002/ece3.694

Matthysen, E. (2005). Density-dependent dispersal in birds and mammals. *Ecography*, *28*(3), 403–416.
Retrieved from http://onlinelibrary.wiley.com/doi/10.1111/j.0906-7590.2005.04073.x/pdf

Matthysen, E. (2012). Multicausality of dispersal: a review. In J. Clobert, M. Baguette, T. G. Benton, & J.
M. Bullock (Eds.), *Dispersal ecology and evolution* (pp. 3–18). Oxford, UK: Oxford University
Press.

McElreath, R. (2020). *Statistical rethinking: a Bayesian course with examples in R and Stan* (2nd edition).
Boca Raton, USA: Chapman and Hall/CRC.

Miller, T. E. X., Angert, A. L., Brown, C. D., Lee-Yaw, J. A., Lewis, M., Lutscher, F., … Williams, J. L. (2020).
Eco-evolutionary dynamics of range expansion. *Ecology*, *n/a*(n/a), e03139. doi: 10.1002/ecy.3139

Mona, S., Ray, N., Arenas, M., & Excoffier, L. (2014). Genetic consequences of habitat fragmentation
during a range expansion. *Heredity*, *112*(3), 291–299. doi: 10.1038/hdy.2013.105

Morel-Journel, T., Girod, P., Mailleret, L., Auguste, A., Blin, A., & Vercken, E. (2016). The highs and lows of
dispersal: how connectivity and initial population size jointly shape establishment dynamics in
discrete landscapes. *Oikos*, *125*(6), 769–777. doi: 10.1111/oik.02718

Nei, M. (1973). Analysis of gene diversity in subdivided populations. *Proceedings of the National*
Academy of Sciences, *70*(12), 3321–3323. doi: 10.1073/pnas.70.12.3321

Panja, D. (2004). Effects of fluctuations on propagating fronts. *Physics Reports*, *393*(2), 87–174. doi:
10.1016/j.physrep.2003.12.001

Panja, D., & van Saarloos, W. (2002). Fronts with a growth cutoff but with speed higher than the linear
spreading speed. *Physical Review E*, *66*(1), 015206. doi: 10.1103/PhysRevE.66.015206

Pedersen, T. L. (2019). *patchwork: the composer of plots*. Retrieved from https://CRAN.R-
project.org/package=patchwork

Perez, G., Burte, V., Baron, O., & Calcagno, V. (2017). Une méthode d'analyse d'image automatique pour
quantifier rapidement les nombres d'œufs et les taux de parasitisme chez *Trichogramma* sp. In
Cahiers techniques de l'INRA. Innovations entomologiques : du laboratoire au champ ! (pp. 135–
142).

R Core Team. (2020). R: a language and environment for statistical computing (Version 4.0.0). Vienna,
Austria: R Foundation for Statistical Computing. Retrieved from https://www.R-project.org/

Reluga, T. C. (2016). The importance of being atomic: ecological invasions as random walks instead of
waves. *Theoretical Population Biology*, *112*, 157–169. doi: 10.1016/j.tpb.2016.06.002

Renault, D., Laparie, M., McCauley, S. J., & Bonte, D. (2018). Environmental adaptations, ecological
filtering, and dispersal central to insect invasions. *Annual Review of Entomology*, *63*(1), 345–368.
doi: 10.1146/annurev-ento-020117-043315

Rodrigues, A. M. M., & Johnstone, R. A. (2014). Evolution of positive and negative density-dependent
dispersal. *Proceedings of the Royal Society of London B: Biological Sciences*, *281*(1791),
20141226. doi: 10.1098/rspb.2014.1226

Ronce, O. (2007). How does it feel to be like a rolling stone? Ten questions about dispersal evolution.
Annual Review of Ecology, Evolution, and Systematics, *38*, 231–253. Retrieved from
http://www.jstor.org/stable/30033859

Roques, L., Garnier, J., Hamel, F., & Klein, E. K. (2012). Allee effect promotes diversity in traveling waves
of colonization. *Proceedings of the National Academy of Sciences*, *109*(23), 8828–8833. doi:
10.1073/pnas.1201695109

Saastamoinen, M., Bocedi, G., Cote, J., Legrand, D., Guillaume, F., Wheat, C. W., … Delgado, M. del M.
(2018). Genetics of dispersal. *Biological Reviews*, *93*(1), 574–599. doi: 10.1111/brv.12356

Salecker, J., Sciaini, M., Meyer, K. M., & Wiegand, K. (2019). The nlrx R package: a next-generation
framework for reproducible NetLogo model analyses. *Methods in Ecology and Evolution*, *10*(11),
1854–1863. doi: 10.1111/2041-210X.13286

Santini, L., Isaac, N. J. B., & Ficetola, G. F. (2018). TetraDENSITY: A database of population density
estimates in terrestrial vertebrates. *Global Ecology and Biogeography*, *27*(7), 787–791. doi:
10.1111/geb.12756

Schreiber, S. J., & Beckman, N. G. (2020). Individual variation in dispersal and fecundity increases rates of
spatial spread. *AoB PLANTS*, *12*(3). doi: 10.1093/aobpla/plaa001

Sexton, J. P., McIntyre, P. J., Angert, A. L., & Rice, K. J. (2009). Evolution and ecology of species range
limits. *Annual Review of Ecology, Evolution, and Systematics*, *40*(1), 415–436. doi:
10.1146/annurev.ecolsys.110308.120317

Stan Development Team. (2018). RStan: the R interface to Stan (Version 2.17.3). Retrieved from
http://mc-stan.org

Stevens, V. M., Pavoine, S., & Baguette, M. (2010). Variation within and between closely related species
uncovers high intra-specific variability in dispersal. *PLoS ONE*, *5*(6), e11123. doi:
10.1371/journal.pone.0011123

Stevens, V. M., Trochet, A., Blanchet, S., Moulherat, S., Clobert, J., & Baguette, M. (2013). Dispersal
syndromes and the use of life-histories to predict dispersal. *Evolutionary Applications*, *6*(4), 630–
642. doi: 10.1111/eva.12049

Stokes, A. N. (1976). On two types of moving front in quasilinear diffusion. *Mathematical Biosciences*, *31*(3), 307–315. doi: 10.1016/0025-5564(76)90087-0

St-Onge, M., Cormier, D., Todorova, S., & Lucas, É. (2014). Comparison of *Ephestia* kuehniella eggs sterilization methods for *Trichogramma* rearing. *Biological Control*, *70*, 73–77. doi: 10.1016/j.biocontrol.2013.12.006

Szűcs, M., Melbourne, B. A., Tuff, T., Weiss-Lehman, C., & Hufbauer, R. A. (2017). Genetic and demographic founder effects have long-term fitness consequences for colonising populations. *Ecology Letters*, *20*(4), 436–444. doi: 10.1111/ele.12743

Travis, J. M. J., Murrell, D. J., & Dytham, C. (1999). The evolution of density–dependent dispersal. *Proceedings of the Royal Society of London B: Biological Sciences*, *266*(1431), 1837–1842. doi: 10.1098/rspb.1999.0854

Travis, J. M. J., Mustin, K., Benton, T. G., & Dytham, C. (2009). Accelerating invasion rates result from the evolution of density-dependent dispersal. *Journal of Theoretical Biology*, *259*(1), 151–158. doi: 10.1016/j.jtbi.2009.03.008

Van Allen, B. G., & Bhavsar, P. (2014). Natal habitat effects drive density—dependent scaling of dispersal decisions. *Oikos*, *123*(6), 699–704. doi: 10.1111/oik.01240

Van Petegem, K., Moerman, F., Dahirel, M., Fronhofer, E. A., Vandegehuchte, M. L., Van Leeuwen, T., … Bonte, D. (2018). Kin competition accelerates experimental range expansion in an arthropod herbivore. *Ecology Letters*, *21*(2), 225–234. doi: 10.1111/ele.12887

Vehtari, A., Gelman, A., Simpson, D., Carpenter, B., & Bürkner, P.-C. (2020). Rank-normalization, folding, and localization: an improved \widehat{R} for assessing convergence of MCMC. *Bayesian Analysis*. doi: 10.1214/20-BA1221

Wajnberg, E., Fauvergue, X., & Pons, O. (2000). Patch leaving decision rules and the Marginal Value Theorem: an experimental analysis and a simulation model. *Behavioral Ecology*, *11*(6), 577–586. doi: 10.1093/beheco/11.6.577

Walter, J. A., Grayson, K. L., Blackburn, L. M., Tobin, P. C., & Johnson, D. M. (2019). Spatiotemporal variability in Allee effects of invading gypsy moth populations. *Biological Invasions*. doi: 10.1007/s10530-019-02096-5

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., … Yutani, H. (2019). Welcome

to the Tidyverse. *Journal of Open Source Software*, *4*(43), 1686. doi: 10.21105/joss.01686

Wilensky, U. (1999). *NetLogo*. Evanston, IL, USA: Center for Connected Learning and Computer-Based

Modeling, Northwestern University. Retrieved from http://ccl.northwestern.edu/netlogo/

Williams, J. L., Hufbauer, R. A., & Miller, T. E. X. (2019). How evolution modifies the variability of range

expansion. *Trends in Ecology & Evolution*, *34*(10), 903–913. doi: 10.1016/j.tree.2019.05.012

Williams, J. L., & Levine, J. (2018). Experimental evidence that density dependence strongly influences

plant invasions through fragmented landscapes. *Ecology*, *99*(4), 876–884. doi: 10.1002/ecy.2156