Deciphering a marine bone degrading microbiome reveals a complex community effort

Erik Borchert[a,#], Antonio García-Moyano[b], Sergio Sanchez-Carrillo[c], Thomas G. Dahlgren[b,d],

Beate M. Slaby[a], Gro Elin Kjæreng Bjerga[b], Manuel Ferrer[c], Sören Franzenburg[e] and Ute

Hentschel[a,f]

[a]GEOMAR Helmholtz Centre for Ocean Research Kiel, RD3 Research Unit Marine Symbioses,

Kiel, Germany

[b]NORCE Norwegian Research Centre, Bergen, Norway

[c]CSIC, Institute of Catalysis, Madrid, Spain

[d]Department of Marine Sciences, University of Gothenburg, Gothenburg, Sweden

[e]IKMB, Institute of Clinical Molecular Biology, University of Kiel, Kiel, Germany

[f]Christian-Albrechts University of Kiel, Kiel, Germany

Running Head: Marine bone degrading microbiome

[#]Address correspondence to Erik Borchert, eborchert@geomar.de

Abstract word count: 229

Text word count: 4908 (excluding Abstract, Importance, Materials and Methods)

Abstract

The marine bone biome is a complex assemblage of macro- and microorganisms, however the enzymatic repertoire to access bone-derived nutrients remains unknown. The bone matrix is a composite material made up mainly of organic collagen and inorganic hydroxyapatite. We conducted field experiments to study microbial assemblages that can use organic bone components as nutrient source. Bovine and turkey bones were deposited at 69 m depth in a Norwegian fjord (Byfjorden, Bergen). Metagenomic sequence analysis was used to assess the functional potential of microbial assemblages from bone surface and the bone eating worm *Osedax mucofloris,* which is a frequent colonizer of whale falls and known to degrade bone. The bone microbiome displayed a surprising taxonomic diversity revealed by the examination of 59 high-quality metagenome assembled genomes from at least 23 bacterial families. Over 700 genes encoding enzymes from twelve relevant enzymatic families pertaining to collagenases, peptidases, glycosidases putatively involved in bone degradation were identified. Metagenome assembled genomes (MAGs) of the class Bacteroidia contained the most diverse gene repertoires. We postulate that demineralization of inorganic bone components is achieved by a timely succession of a closed sulfur biogeochemical cycle between sulfur-oxidizing and sulfur-reducing bacteria, causing a drop in pH and subsequent enzymatic processing of organic components in the bone surface communities. An unusually large and novel collagen utilization gene cluster was retrieved from one genome belonging to the gammaproteobacterial genus *Colwellia*.

Importance

Bones are an underexploited, yet potentially profitable feedstock for biotechnological advances

and value chains, due to the sheer amounts of residues produced by the modern meat and poultry

processing industry. In this metagenomic study we decipher the microbial pathways and

enzymes that we postulate to be involved in bone degradation marine environment. We herein

demonstrate the interplay between different bacterial community members, each supplying

different enzymatic functions with the potential to cover an array of reactions relating to the

degradation of bone matrix components. We identify and describe a novel gene cluster for

collagen utilization, which is a key function in this unique environment. We propose that the

interplay between the different microbial taxa is necessary to achieve the complex task of bone

degradation in the marine environment.

Introduction

The marine environment is a treasure trove for novel microbial assemblages and organic catalysts (enzymes) (1-3). The oceans cover approximately 70% of the Earth surface with an estimated volume of about 2×10^{18} m^3 and due to its incredible environmental variability (e.g. temperature, pressure, salinity, light availability), it has sparked the evolution of an unprecedented range of different microbes and hence enzymatic activities (4-6). Genome sequencing of individual microbial isolates of complex communities has allowed us to get a glimpse of their diversity and their potential functions in multiple environmental contexts. The lack of cultivable microbes has further driven the development of functional and sequence-driven metagenomic analyses, and enabled us to decipher complex interactions in entire microbial consortia (7-9).

The deep-sea was for a long time seen as an almost lifeless environment, as no one could imagine life to be possible under conditions vastly different from shallower ocean waters in respect to nutrient and energy resources. Nowadays we know that even the deep-sea is teeming with life; hydrothermal vents, sponge grounds and coral gardens are recognized as examples of unique and complex habitats (10-12). Nonetheless, the deep-sea is also a harsh environment with limited nutrient sources. In this respect sudden events like a whale fall, create a locally defined but significant nutrition source for deep-sea life, that can last for years or even decades (13). These whale carcasses are rapidly stripped of their soft tissue by scavengers (i.e., hagfish, sleeper sharks, rat tail fish, crabs), but the energy-rich bones and cartilage remain as a recalcitrant nutrient source. More than 15 years ago *Osedax* was described, a genus of bone-eating annelid worms (14), and has since then been investigated for its diversity, ecology and how it accesses the organic compounds of whale bones (14-17). These worms are gutless and rather bore cavities

into bones and develop a root tissue in these cavities for food intake. Furthermore, this

evolutionary novel and specialized organ was shown to harbor endosymbionts, typically

affiliated to Oceanospirillales (14, 18-21). *Osedax* species are known for their ability to acidify

their environment via elevated expression levels of vacuolar-H^+-ATPase (VHA) specifically in

their root tissue and of carbonic anhydrase throughout their body, to dissolve hydroxyapatite and

access collagen and lipids from the bone matrix (16). Miyamoto *et al.* found a high number of

matrix metalloproteinases in the genome of *Osedax japonicus* compared to other invertebrates,

potentially assisting in digestion of collagen and other proteins derived from bones (15). The

species can thus be regarded as a member of the bone biome and an important facilitator in this

degradation process. In the northern North Atlantic *Osedax mucofloris* was described in 2005

and has been shown to consistently colonize bone material on the sea floor below a depth of 30

m (22, 23).

Bone is a recalcitrant and heterogeneous composite material made of a mineral phase, an organic

phase and water. Hydroxyapatite crystals in the mineral phase contribute to the structural

strength in bones. The organic phase includes proteins, such as collagen and structural

glycoproteins (e.g. proteins decorated with sugars such as mannose, galactose, glucosamine,

galactosamine, N-acetylglucosamine, N-acetylgalactosamine, rhamnose, sialic acid and fucose),

lipids and cholesterol composed of various triglycerides (24-26). Up to 90% of the protein

content in mature bone is made of type I collagen, a triple helical molecule rich in glycine,

hydroxyproline and proline that assembles into fibrils with a high number of hydrogen bonds,

hydrophobic interactions and covalent cross-linking, which together confer high structural

stability to collagen fibrils (27). Due to this structural and chemical complexity, it is expected

that the degradation of the recalcitrant bone matrix will require a synergistic multi-enzyme

system and require a microbial community effort. Similar multi-enzyme systems are well

described for example for the degradation of lignocellulose, another important organic polymer

(28, 29). This system will likely include essential enzymes in the breakdown of the organic

matrix, namely collagenases that break the peptide bonds in collagen and other

proteases/peptidases (endo- and exopeptidases) that attack the glycoproteins. Furthermore,

neuraminidases (sialidases), α-mannosidases, α-/β-galactosidases, α-fucosidase, α-rhamnosidase

and α/β-N-acetylhexosaminidase (glucose and galactose-like), all glycoside hydrolase enzymes

are likely involved in cleavage of glycosidic linkages. Finally, in the digestion of the cholesterol-

containing marrow, cholesterol oxidases may be involved.

To date only a few studies have been published that focus on microbial communities to

understand the necessary complex interactions in bone degradation, mainly relying on 16S rRNA

gene sequencing data (30-33) and one metagenomic study of a whale fall (34). We here provide

a first comprehensive overview and identify putative key functions involved in bone degradation

of the marine bone microbiome retrieved from deployed bone material, including microbial

communities from the gutless worm *Osedax mucofloris* and free-living microbial assemblages

developing on the bone surface.

Results

Recovery of artificially deployed bone for bone microbiome metagenomic analysis

Turkey and bovine bones were deployed at 69 m depths in Byfjorden, a fjord outside Bergen,

Norway. After nine months of incubation, underwater images taken by a remotely operated

vehicle (ROV) showed microbial colonization of the bone surfaces (supplementary figure S1A).

Osedax mucofloris worms were observed, especially on the joints, and in some cases forming

small colonies of several individuals inside a single cavity under the external bone tissue.

Although not subject of this study, a larger diversity of invertebrate fauna including

Ophryotrocha, *Vigtorniella* and *Capitella* worms were also observed. Dense microbial mats

developed asymmetrically with preference for the joint adjacent sections (epiphysis), which also

appeared in aquaria settings (Supplementary figure S1B).

Two sets of samples, an *Osedax*-associated bone microbiome (OB), and a bone surface-

associated biofilm (BB) were collected, each consisting of four individual metagenomes (Table

1). The co-assemblies of each sample set comprised of >300 000 contigs and 342.7 Mb for the

OB-metagenomes and >1 000 000 contigs and 1.22 Gb for the BB-metagenomes respectively,

considering only contigs >500 bp (Supplementary table S1). The individual metagenomes were

profiled separately with Kaiju (35), a database aided metagenomic read taxonomy classifier. The

OB-metagenomes comprised of 45-76% bacterial, 15-34% eukaryotic and 0.4-0.5% archaeal

reads and the BB-metagenomes are made up of 92-95% bacterial, 2-4% eukaryotic and 0.4-0.7%

archaeal reads (Supplementary table S1).

Metagenome assembled genomes (MAGs) from the marine bone microbiome

display taxonomic diversity and novelty

59 high-quality MAGs (>90% completion and <10% redundancy) were extracted from the co-

assembled metagenomes (Supplementary Table S3 for MAG sequence statistics). The MAGs

span 11 phyla, 14 classes, 19 orders and at least 23 families. About 63% of the MAGs (37/59)

are taxonomically novel as determined by their relative evolutionary divergence (RED) (36) to

their closest common ancestor (Supplementary table S2). One MAG could be only identified up

to phylum level, seven to class level, seven to order level, 18 up to family level and four up to

genus level. The taxonomy of most MAGs was fully resolved based on 120 marker genes. The

three best represented phyla were Proteobacteria (22 MAGs), Campylobacterota (14 MAGs) and

Bacteroidota (8 MAGs). However, the percental distribution of the most abundant classes differs

between the two metagenome sets. The OB-MAGs were dominated by the classes

Gammaproteobacteria (27%), Campylobacteria (27%) and Alphaproteobacteria (20%), while the

BB-MAGs were mainly affiliated with Gammaproteobacteria (30%), Campylobacteria (23%)

and Bacteroidia (16%).

Sulfur cycling in the marine bone microbiome

All MAGs were investigated using multigenomic entropy based score pipeline (MEBS) (37) for

their ability to utilize sulfur as energy source via the abundance of selected marker genes for

related pathways (Figure 2). Sulfur cycling is of relevance to bone degradation due to the

generation of free protons by sulfur and sulfur compound oxidation processes (thiotrophy),

which leads to an acidification.

MAGs affiliated to the order Campylobacterales encode almost complete Sox (*sox*XYZABCD)

and Sor enzyme systems for sulfur oxidation. Furthermore, Gammaproteobacteria affiliated to

Beggiatoaceae and other unclassified Gammaproteobacteria, are all potentially capable of

thiotrophy via utilization of reduced sulfur compounds as electron donors (flavocytochrome c

sulfide dehydrogenase (*fcc*B) and adenosine-5'-phospho-sulfate reductase (*apr*AB)) and partial

predicted Sox sulfur/thiosulfate oxidation pathway. MAGs identified as Desulfuromonadia,

Desulfobacteria, Desulfobulbia and Desulfovibrionia possess marker genes for sulfur reduction

(*qmo*ABC, *hyd*ACD, *sre*ABC) and lack Sox and Sor pathway genes, all of which belong to the

new proposed phylum of Deltaproteobacteria (38). In all Gammaproteobacteria (except BB32),

Desulfobacteria, Desulfobulbia and Desulfovibrionia, genes for dissimilatory sulfite reductase

(*dsr*ABC) are present. Müller *et al.* (2015) described that gammaproteobacterial *dsr*AB-type

genes are commonly involved in oxidative reactions, whereas *dsr*AB in Desulfobacterota are

reductive type *dsr*AB (39). All MAGs contain at least partial pathways for dissimilatory

tetrathionate reduction (*ttr*ABC), thiosulfate disproportionation (*phs*ABC and rhodanase),

dimethylsulphide (DMS) degradation (*ddh*ABC) and contain also genes for sulfoacetaldehyde

degradation (*isf*D, xsc and *saf*D). In addition, the phenotypic trait of H_2S production was

identified in 10 MAGs (Traitar analysis (40)), two of which were Marinifiliaceae, two

Krumholzibacteria, two *Sulfurospirillum*, one Spirochaetaceae, two Desulfobacteraceae, and one

Pseudodesulfovibrio.

The anticipated thiotrophy has the potential to contribute massively to the acidification of the

environment via the oxidation of reduced sulfur compounds leading to production of sulfuric

acid (41). This requires a close interaction between sulfur-reducing bacteria (SRB) producing

hydrogen sulfide and sulfur-oxidizing bacteria (SOB) utilizing the hydrogen sulfide, while

releasing protons (Figure 3). The Traitar analysis identified 10 MAGs potentially able to produce

hydrogen sulfide, including known SRB like Desulfobacteraceae, *Pseudodesulfovibrio*, and

others like *Sulfurospirillum* (42-44). The bone microbiome is especially enriched in taxa

containing known SOB, like the large filamentous bacteria Beggiatoales (5 MAGs) (45) and

Campylobacterales (10 MAGs) (46). Furthermore, one MAG identified as Desulfobulbaceae was

found in the bone associated metagenomes. Member of this group are known to be able to

perform sulfur oxidation and sulfur reduction (47, 48).

Acidification by carbonic anhydrases

Carbonic anhydrases were identified in 51 of 59 MAGs. Nineteen out of 94 carbonic anhydrases

contained a signal peptide for extracellular export (16 MAGs). 15 were predicted to contain a

Sec signal peptide (SPI) and four to encode lipoprotein (SPII) signal peptides. Four out of five

Beggiatoales MAGs were predicted to contain carbonic anhydrases with a SPI signal peptide

(BB2, BB3 and BB20) or a SPII signal peptide (BB16). The remaining three SPII signal peptides

were found in carbonic anhydrases from Campylobacterales (BB8, BB10 and BB11)

Interestingly BB8 contains at least three carbonic anhydrases, one with a SPI, one with a SPII

and one where a signal peptide was not predicted. Three SPI signal peptides were found in

carbonic anhydrases from unclassified γ-proteobacteria (BB4, BB34 and BB36). The remaining

SPI including carbonic anhydrases were found in five Campylobacterales MAGs (BB14, BB26

(three carbonic anhydrases and two containing SPI signal peptides), BB30, BB41 and OB7) and

one Desulfobulbaceae MAG (BB13). Based on phylogenetic relationship to known carbonic

anhydrases described in Capasso *et al.*, 2015 (49), 18 out of 19 carbonic anhydrases belong to

the α-carbonic anhydrase family and one to the β-family, with no γ-family carbonic anhydrases

found (Supplementary figure S4).

Gelatin hydrolysis

With respect to microbial bone degradation, the phenotypic feature of gelatin hydrolysis was

analyzed using the Traitar software (40), which   provides genome-informed phenotype

predictions. 22 MAGs showed capacity of gelatin hydrolysis (19 MAGs in the bone surface

community (BB) and three in the *Osedax* associated communities (OB)). With gelatin being a

primarily bone collagen derived compound, we consider gelatin hydrolysis a key trait for the

microbial community studied herein. All eight Bacteroidia affiliated MAGs possess the gelatin

hydrolysis trait, seven Gammaproteobacteria MAGs, one tentative Planctomycetota MAG, one

Spirochaetia   MAG,   two   Krumholzibacteria   MAGs,   one   Thiovulaceae,   one

Geopsychrobacteraceae and one Fermentibacteria MAG (Figure 3).

Enzymes involved in bone degradation

Based on the structure and composition of mature vertebrate bone tissue, we hypothesized that

12 different COGs and peptidase/collagenase families were relevant for the enzymatic attack of

the bone organic matrix. This "bone-degradome" comprised the peptidase families S1

(COG0265), S8/S53 (and Pfam00082), U32 (COG0826) and M9 collagenase (including

Pfam01752), mannosidases (COG0383), sialidases (COG4409), glucuronidases (COG3250),

glucosaminidases (COG1472), galactosaminidases (COG0673), α-galactosidases (Pfam16499),

cholesterol oxidases (COG2303) and fucosidases (COG3669) (choice of enzymes is justified in

the Materials and Methods section). We constructed HMM profiles that were used to screen the

abundance of each enzyme family in all MAGs (Figure 4). In total 722 enzymes belonging to the

12 investigated enzyme families were identified in the 59 MAGs. The glycosidase families of

mannosidases, galactoaminidases, glucosaminidases and the peptidase families S1, S8/S53 and

U32 were widespread in the MAGs (Figure 4). M9 collagenases and α-galactosidases

(Pfam16499) were only found in three MAGs. The M9 collagenases where solely found in

Enterobacterales (BB5, BB44 and OB12). Pfam16499 was only identified in Bacteroidales

(BB22, BB24 and OB13). The most abundant group of enzymes were the S1 peptidases (141

hits), followed by galactosaminidases (COG0673) (116 hits) and U32 peptidases (99 hits)

(Figure 4), constituting 20%, 16% and 14% of all identified bone degrading enzymes,

respectively. In general, Bacteroidales (BB17, BB22, BB24, BB29, BB42 and OB13) displayed

the most diverse set of enzyme families related to bone degradation, as they contained genomic

evidence of all enzymes besides M9 collagenases. MAGs belonging to the orders

Desulfuromonadia, Desulfobulbia, Desulfobacteria, Desulfovibrio, Campylobacteria (all of them

driving the sulfur biogeochemical cycle), as well as some undefined α-proteobacteria and γ-

proteobacteria appear to have no or few mannosidases (COG0383), glucuronidases (COG3250),

fucosidases (COG3669), sialidases (COG4409) and α-galactosidases (Pfam16499).

Collagen degradation

We investigated the genomic context of each M9 collagenase for potential links to metabolic

pathways, such as proline utilization (Supplementary figure S4). *Colwellia* MAG BB5 possessed

an approximately 21 kbp-long gene cluster presumably devoted to collagen utilization, which is

unique in the dataset and in the public databases. The functional cluster spans at least 15

different genes (Figure 5A), featuring a secreted Zn-dependent M9 collagenase, a secreted

peptidyl-prolyl cis-trans isomerase (cyclophilin-type PPIase), a secreted unknown protein and an

unknown Zn/Fe chelating domain-containing protein. Additionally, one putative transporter

(MFS family), a TonB-dependent receptor and several genes involved in the catabolism of

proline and hydroxyproline, e.g. prolyl-aminopeptidase YpdF, intracellular peptidyl-prolyl cis-

trans isomerase (rotamase), pyrroline reductase, hydroxyproline dipeptidase, 4-hydroxyproline

epimerase and others. Moreover, genes involved in transcription regulation such as PutR and the

stringent starvation protein A and B were identified in the same putative gene cluster.

To explore the conservation of this gene cluster, we retrieved fourteen representative *Colwellia*

genomes of marine origin from the NCBI repository (Supplementary table S5) (50-63). To

minimize methodological bias, the nucleotide sequences of these genomes were likewise

annotated with RAST (rapid annotation using subsystem technology) and screened for M9

collagenase using the previously established HMM profile. 22 annotated M9 collagenases were

identified in seven out of fourteen genomes. In the genomes of *Colwellia piezophila* ATCC

BAA-637 (57) and *Colwellia psychrerythraea* GAB14E (59) a gene cluster comparable to the

one in MAG BB5 was identified (Figure 6) and found to be largely conserved between the three

species. The conserved core constitutes of the M9 collagenase, a D-hydroxyproline

dehydrogenase, an epimerase, a secreted cyclophilin-type peptidyl-prolyl isomerase, a

ketoglutaric acid dehydrogenase, a MFS transporter, an aminopeptidase, a bifunctional 1-

pyrroline-5-carboxylate reductase/ornithine cyclodeaminase and a hydroxyproline dipeptidase.

BB5 additionally contains several other relevant genes, such as a PutR regulator, stringent

starvation proteins A and B, a TonB dependent receptor, Zn/Fe binding domain protein, 1-

pyrroline-4-hydroxy-2-carboxylate deaminase and an intracellular peptidyl-prolyl cis-trans

isomerase (rotamase, PpiD).

Specificity of the bone microbiome

To investigate the potential specificity of the bone microbiome in respect to its taxonomic

composition and function the generated MAGs were compared to 832 seawater MAGs generated

from the Tara Ocean datasets, comprising 93 metagenomes from various locations (64). The

bone microbiome was rebinned to retrieve all MAGs with >70% completion to match the Tara

Ocean threshold for high-quality MAGs, resulting in 86 MAGs. The taxonomic class abundances

were compared between the two datasets and Gammaproteobacteria and Bacteroidia appeared in

similar abundances (bone microbiome 23.26% and 15.12% and Tara Oceans 23.56% and 13.34%

respectively), but other than those high taxonomic ranks, the datasets were different. The bone

microbiome is dominated by Campylobacteria, accounting for 26.74% of the MAGs, whereas

only 0.12% Campylobacteria MAGs are found in the Tara Oceans dataset. In contrast

Alphaproteobacteria are better represented in the Tara Oceans dataset, than in the bone

microbiome, 21.88% to 5.81% respectively. The bone microbiome furthermore also contains a

number of bacterial classes that are represented at low levels (1-5% of the MAGs), which are

either not represented in the Tara Oceans dataset or only at miniscule levels as low as 0.1-0.5%

(Supplementary figure S4A). The functional repertoire was compared via screening the Tara

Oceans dataset with the previously generated HMM profiles for enzymes potentially involved in

bone degradation and calculated as ratio of enzymes per MAG. The ratios for 9 out of 12 profiled

enzyme families were higher in the bone microbiome, ratios for α-N-acetylgalactosaminidases

(COG0383) and cholesterol oxidases (COG2303) were higher in the Tara Oceans dataset and the

ratio for S1 peptidases was equal between the two MAG sets (Supplemetary figure S4B).

Discussion

In this study, 59 high quality MAGs were reconstructed from microbes colonizing bone surfaces

and from symbionts of the bone-eating worm *Osedax mucofloris*. Metabolic reconstruction

revealed a complex, diverse and specialized community. Our MAGs span at least 23 bacterial

families and uncover a large potential for taxonomic novelty (over 50% according to genome-

based taxonomy) from species up to class level in the bone microbiome. Interestingly, only

genomes of gram-negative bacteria were reconstructed and despite gram-positive bacteria being

widespread in the marine environment, they make up only minor portions of the metagenomes

(4-5% of the reads affiliated to Actinobacteria and Firmicutes in the *Osedax* metagenomes and 5-

9% in the Biofilm metagenomes respectively) (65). This is remarkable since they are known to

carry out potentially relevant metabolic processes (thiotrophy, sulfidogenesis) (66, 67), are

capable of dealing with low pH conditions which are likely encountered during bone dissolution

(68), and they possess high capacity for the secretion of hydrolytic enzymes (69). Despite this

underrepresentation of gram-positive taxa, this study reveals the existence of a specialized bone-

degrading microbiome in the marine environment and starts to explore the enzymatic activities

involved in the complete demineralization of bone material. The bone microbiome is different

from seawater communities and from other specialized habitats (recolonized volcanic eruption

site) in its microbial composition as well as functional makeup (Supplementary figure 4).

The role of *Osedax* endosymbionts in bone utilization

Two distinct bacterial endosymbiont genomes belonging to the order Oceanospirillales have

previously been sequenced, but their role in bone degradation in the marine environment

remained unclear (18). The bacterial fraction of the herein sequenced *Osedax mucofloris*

metagenome is made up of 7% to 22% Oceanospirillales affiliated reads, whereas the bone

surface metagenome only contains 1% to 4% reads of this order according to the performed

Kaiju analysis. This relative difference confirms that the methodological approach to minimize

cross-contamination was successful and that the OB-MAGs affiliated to Oceanospiralalles likely

represent the symbiotic community of *Osedax mucofloris* worms. Two MAGs belonging to the

Oceanospirillales were identified in the *Osedax*-associated metagenome, belonging to the genera

Neptunomonas (OB1) and *Amphritea* (OB2). Both genera are known to have an aerobic

organotrophic metabolism and also able to thrive as free-living bacteria (70). In fact, the scarce

representation of Oceanospirillales in the bone surface has been reported before (30, 31) and

contrasts with their character as common dwellers of the marine environment (71, 72). Although

we cannot rule out that Oceanospirillales preferentially colonize bone surfaces in earlier or later

stages, their preference for a symbiont life supports the notion of a casual and facultative

association to *Osedax* worms, triggered by the common benefit from a sudden nutrient bonanza

as previously hypothesized (18, 21).

The degradative functions within the bone microbiome

Acidification via a closed sulfur biogeochemical cycle

Free-living microbial communities, must deal with similar challenges as the *Osedax* holobiont to

access the nutrient-rich, collagen-made organic bone matrix and eventually the lipid-rich bone

marrow by dissolving the hydroxyapatite. The association in large specialized microbial

consortia may be a beneficial strategy for achieving this task. We hypothesize that sulfur-driven

geomicrobiology (sulfate/thiosulfate/tetrathionate reduction and sulfide/sulfur/thiosulfate

oxidation) is the major responsible factor for bone dissolution in the marine environment by free-

living bacterial communities. Campylobacterales are one of the most abundant bacterial orders in the herein investigated metagenomes, in both the *Osedax*-associated metagenomes (OB) and the bone surface biofilms (BB). Campylobacterales represent the most abundant group in terms of absolute read number, although it is the second largest taxon with reconstructed MAGs (Figure 1). Members of the Campylobacterales have previously been found to be associated with *Osedax*, albeit not as endosymbionts (21). The majority of retrieved Campylobacterales MAGs (14 in total) belong to different families of aerobic and facultative anaerobic (nitrate, manganese) sulfur-oxidizing bacteria (73, 74) (Thiovulaceae, Sulfurovaceae and Arcobacteraceae). Other aerobic/facultatively anaerobic (nitrate) sulfur oxidizing bacteria are also well represented in the order Beggiatoales (Gammaproteobacteria, 5 MAGs). *Beggiatoa*-like bacterial mats are commonly associated with whale falls (75) indicating a potential indifference regarding the bone type they dwell on. Sulfide oxidation produces elemental sulfur or sulfate (41, 45) while releasing protons and thereby causing a drop in pH. This acidification mechanism has been linked to bone demineralization. The dissolution of the hydroxyapatite mineral exposes the organic matrix to enzymatic degradation (30, 31). Besides thiotrophy, that seems to be a major acid-producing mechanism in the microbial community, other mechanisms might also contribute significantly. In this respect, a number of carbonic anhydrases (CA) were annotated, normally housekeeping genes involved in internal pH homeostasis and other processes (76), but known to play a role in environmental acidification by *Osedax* (15). Here, the CAs were found to contain signal peptides for extracellular export (19 out of 94) and therefore could also be involved in acidification. Interestingly, 18 out of 19 identified and potentially secreted CAs belong to the α-CA family and only one member of the β-CA family was found (Supplementary figure S4). The

α-CA family is only found in gram negative bacteria, which is also the case here, and it is

evolutionarily the youngest of the three bacterial CA families (49).

Besides a large number of SOB, eight MAGs related to SRB were identified that are affiliated to

the families Desulfobulbaceae (also SOB), Desulfobacteraceae, Geopsychrobacteraceae and

Desulfovibrionacaeae. Moreover, they are prevalently associated with the free-living community

attached to the bone surface in this study. Sulfate, tetrathionate or thiosulfate can serve as

electron acceptors and/or donors and gene markers for all pathways are present in the

metagenomes (Figure 2). Microbial sulfidogenesis on the bone surface or the surrounding

sediments can feed the thiotrophic community and therefore accelerate the demineralization

process. The generated sulfide is known to quickly react with iron, blackening the bone surfaces

with insoluble iron sulfide (77). In our incubation experiments, blackening occurs preferentially

on the epiphysis, which is also where complex white/pink microbial mats are forming over time

(Supplementary figure S1B). However, from our analysis SRB seem unable to degrade large

complex molecules. This is supported by the lack of bone-degrading enzymes herein

investigated, such as: S8/S53 peptidases, mannosidases, sialidases, fucosidases and α-

galactosidases. SRB are likely dependent on the generation of simple organic compounds

produced as metabolites by fermenters or aerobic organotrophic bacteria of the wider bone

microbiome. The bone dissolution driven by sulfur geomicrobiology relies on other specialized

members of the community to degrade the organic matrix and to fuel the acid generation.

Degradation of organic compounds via peptidases, glucosidases and oxidases

Once the inorganic hydroxyapatite is removed, an array of different enzymes is required to digest

the various organic bone components. Bacteroidia appear to be especially remarkable in this

respect and represent the third most abundant taxon. Eight high quality MAGs could be

reconstructed, seven of them from the bone surface metagenome. Bacteroidia, and especially the

family of Flavobacteriaceae, are known to be versatile degraders of polysaccharides like agar

(78), chitin (79), ulvan (80), alginate (81), carrageen (82), cellulose and xylanose (83) and

polypeptides like elastin (84), spongin (85) and others. The recently described Marinifilaceae

family (86) includes isolates that are reported to present xylanase activity (87). Despite the

discrepancy between abundance versus reconstructed genomes, the Bacteroidia MAGs appear to

be the most versatile order of the investigated MAGs in respect to their richness in bone

degrading enzymes (Figure 4), and all were predicted to possess the gelatin hydrolysis trait

(Figure 3). They were also the only MAGs containing sialidases (COG4409) and α-

galactosidases (Pfam16499) (Figure 4). Due to this taxa-specific trait and their presence limited

to the bone-surface associated microbiome, we hypothesize that Bacteroidia play a pivotal and

specialized role in the free-living community via the degradation of specific organic bone

components.

Differential microbial colonization of the spongy cancellous bone tissue over the cortical

compact bone has also been observed in the terrestrial environment and has been related to easier

access to the red marrow (88) although a priming effect linked to the differential composition of

the bone cannot be ruled out. Complex microbial mats form preferentially on the epiphysis of the

long bones and this area is normally covered with hyaline cartilage (89) which was not removed

before deployment. Cartilage is a related connective tissue made of nonfibrous type II collagen

and a sulfated-proteoglycan matrix rich in N-acetyl-galactosamine and glucuronic acid residues.

This would explain the abundance of alpha-galactosidases, N-acetyl glucosaminidase and

glucuronidases. Moreover, other groups such as Kiritimatiellales (PVC superphylum) are known

marine anaerobic saccharolytic microbes specialized in degrading sulfated polymers that we find

in this environment (90).

Collagen degradation by Gammaproteobacteria

Peptidases and especially M9 collagenases are of special interest for the degradation of the

proteinogenic compounds within bone, as they are able to degrade collagen, the main source of

carbon in this environment. The class γ-proteobacteria is comparatively enriched in these

enzymes and it is the best represented class in the dataset, with 17 MAGs. Of particular interest

are the MAGs affiliated to the order Enterobacterales (two MAGs of the families Kangiellaceae

and one Alteromondaceae). They possess the gelatin hydrolysis trait (Figure 3, MAGs BB5,

BB44 and OB12), have a high number of S1 and U32 peptidases, and are the only MAGs with

M9 collagenases. The *Colwellia* MAG BB5 is particularly remarkable as it contains an entire

gene cluster dedicated to collagen utilization (Figure 5A). The collagen degradation gene cluster

comprises at least 15 different genes, including a M9 collagenase, a PepQ proline dipeptidase, an

aminopeptidase YpdF, several transporters, epimerase, isomerases and others. The gene cluster

encodes nearly the entire pathway necessary to unwind and hydrolyze triple-helical collagen,

transport and uptake of collagen oligopeptides into the cell and utilization of its main

components, mainly hydroxyproline and proline, for energy production via the TCA cycle and/or

the urea cycle or for polyamine biosynthesis (Figure 5B). Accessory genes for an alternative

catabolic route of proline to glutamate are also encoded elsewhere in the genome (P5CDH). This

kind of functional condensation for collagen utilization has not been described before in

Colwellia or elsewhere. Interestingly, *Colwellia* bacteria are also one partner in a dual

extracellular symbiosis with sulfur-oxidizing bacteria in the mussel *Terua* sp. 'Guadelope',

retrieved from a whale fall in the Antilles arc and supposedly involved in the utilization and

uptake of bone components (91). A cluster of functionally related genes was found in the

publicly available genomes of *Colwellia piezophila* and *Colwellia psychrerythraea.* However,

the gene cluster described for MAG BB5 contains several supplementary features, like a

rotamase, a 1-pyrroline-4-hydroxy-2-carboxylate deaminase and a Zn/Fe binding domain protein

potentially attributed to collagen utilization, which are absent in the published genomes (Figure

6). Moreover, the gene cluster contains regulatory elements like the PutR regulator and stringent

starvation proteins known to be activated under acid stress or amino acid starvation conditions in

Escherichia coli (92). This supports our hypothesis that other members of the microbial

community need to dissolve the bone calcium phosphate via acid secretion, before collagen and

other organic bone compounds can be accessed. In summary the publicly available gene clusters

lack regulatory elements to switch on the collagen utilization pathway under 'bone

degrading'/acidified conditions and misses key enzymes to exploit collagens key components

proline (rotamase missing to transverse D-proline to L-proline) and hydroxyproline (pyrroline-4-

hydroxy-2-carboxylate deaminase missing that breaks down 1-pyrroline-4-hydroxy-2-

carboxylate to alpha-ketoglutarate semialdehyde).

Bone degradation – a complex microbial community effort

The marine bone microbiome is a complex assemblage of various bacterial classes that requires

the synergistic action of many different interwoven enzymatic reactions to access the recalcitrant

bone material for its nutritional resources. A scenario how we envision the orchestration of this

complex process is depicted in Figure 7. The primary requirement in utilizing organic bone

compounds is likely the dissolution of mineralized calcium phosphate (hydroxyapatite) by

acidification, which can potentially be performed via proton release by a versatile community of

sulfur-oxidizing (SOB) γ-proteobacteria (mainly *Beggiatoa*-like), Campylobacterales

(*Sulfurimonas*, *Sufurospirillum*, *Sulfurovum*), Desulfobubales and α-proteobacteria (Figure 7-I).

This acidification via thiotrophy may be fueled by sulfur-reducing bacteria (SRB), like

Desulfobacteraceae, Geopsychrobacteraceae, *Pseudodesulfovibrio*, creating a sulfur

biogeochemical loop between SRB and SOB (Figure 7-II). Once the organic compounds

(collagen, fatty acids, proteins, peptidoglycans) are accessible, the Bacteroidia

(Flavobacteriaceae and Marinifiliaceae) and γ-proteobacteria (Alteromonadaceae and

Kangiellaceae) may become the main protagonists (Figure 7 -III and IV). These Bacteroidia are

especially rich in bone degrading enzymes, but importantly the γ-proteobacteria are the only

members identified with M9 collagenases and *Colwellia* contains an entire gene cluster dedicated

to collagen degradation (Figure 5). Herein we disentangled the potential functional roles of

specialized members of the bone-degrading microbial community, which together make bone-

derived nutrients accessible – not only to themselves, but also to generalists within the bone

microbiome. We posit that Flavobacteriales and Enterobacterales are the most promising

candidates for novel enzyme discovery, as they display the most versatile sets of bone degrading

enzymes.

Materials and Methods

Sample collection

Bone material after manual meat deboning was kindly provided by a local slaughterhouse operated by Norilia (Norway). Since deboning does not completely eliminate the animal tissue attached to the bone, some remains were still present. Therefore, in order to avoid bacterial colonization and decomposition, all bone material was kept at -20 C until deployment. Four sets of turkey thigh bones and one bovine lower leg bone were placed in a crab trap and deposited at the bottom of Byfjorden (60,397093N; 5,301293E) close to Bergen, Norway, at a depth of 69 m and approximately 150 m off shore in May 2016, incubated for nine months and retrieved using a small ROV (Table 1). The material was transported to the lab in styrofoam boxes for either processing within two hours (bone surfaces) or for prolonged incubation in seawater aquaria and subsequent dissection of *Osedax* worms. The meatless bone surfaces were scraped with a sterile scalpel for microorganisms and *Osedax mucofloris* specimens were extracted from the bone using sterile scissors and forceps. Their root tissue was dissected from the body, rinsed in sterile 70% (v/v) sea water and preserved in storage solution (700 g/l ammonium sulfate, 20 mM sodium citrate and 25 mM EDTA, pH 5.2) and stored at -70 °C until further processing.

DNA extraction and sequencing

DNA was extracted from 10 to 50 mg sample of either scrapped biofilm or *Osedax* root tissue, using the Qiagen AllPrep DNA/RNA Mini Kit according to the manufacturer's instructions with cell lysis by a bead beating step in Lysing Matrix E tubes (MP Biomedicals) in a FastPrep homogenizer (MP Biomedicals) with a single cycle of 30s at a speed of 5500 rpm. The obtained

DNA was quantified and quality controlled using a NanoDrop2000 (ThermoFisher Scientific)

and a Qubit fluorometer 3.0 (ThermoFisher Scientific). The obtained DNA concentrations

ranged from 11,9 ng/μl to 166 ng/μl according to Qubit readings. The DNA libraries were

prepared using the Nextera DNA library Prep Kit according to the manufacturer's guidelines. 50

ng DNA was used for the preparations. In brief, the DNA was fragmented by the Nextera

transposome at 55°C for 5 min and barcoded adapters were added in a 5 cycle PCR

amplification. The resulting libraries, including ~140 bp of adapter had an average fragment size

of 436 bp (± 112 bp) and were sequenced on an Illumina HiSeq4000 platform (150-bp paired-

end reads) at the Institute of Clinical Molecular Biology (IKMB), Kiel University, Germany.

Metagenomic read profiling

Illumina raw reads were quality trimmed and adapters were removed with Trimmomatic version

0.36 (93). The quality filtered reads were used individually and combined with respect to their

sample source (either *Osedax*-associated or bone surface biofilms) to profile the taxonomic

origin of the reads with Kaiju (35).

Metagenomic assembly, binning, taxonomic identification, ORF prediction and annotation

For each sample type (*Osedax mucofloris* and bone surface biofilm communities), the quality

filtered metagenomic reads (Trimmomatic version 0.36) were co-assembled with SPAdes v3.12

(94) for kmers 21, 33, 55, 77 and 99, with the metaSPAdes-assembler option enabled (see

Supplementary table S1 for read counts). Binning was conducted on the resulting assemblies

using the MetaWRAP pipeline (version 1.0.1) (95). This pipeline combines three different

implemented binning methods, CONCOCT (96), MaxBin2.0 (97) and metaBAT2 (98), to

retrieve high quality MAGs. We only considered high-quality MAGs with >90% completeness

and <10% redundancy for further analyses. CheckM was used for quality assessment of the

assembled genomes (99), and GTDB-Tk version 0.1.3 (36) was used for taxonomic

identification, coupled with an estimate of relative evolutionary divergence (RED) to their next

common ancestor. RED is a normalization method to assign taxonomic ranks according to

lineage-specific rates of evolution, based on branch lengths and internal nodes in relation to the

last common ancestor calculated by GTDBTk. Open reading frames (ORF) of the obtained

MAGs were predicted with Prodigal version 2.6.3 (100). Predicted ORFs were annotated using

eggNOG-mapper v1 (101) with eggNOG orthology data version 4.5 (102). Additionally, the

MAGs were annotated and metabolic models were calculated using the RAST (rapid annotation

using subsystem technology) server (103, 104). The MAGs were further investigated for the

presence or absence of major metabolic pathways and phenotypic microbial traits based on their

genomic sequences using MEBS (multigenomic entropy based score, version 1.2) (37) and

Traitar (40). MEBS is a software package used here to detect genes related to sulfur metabolism,

this was done by providing protein fasta files of the high-quality MAGs, that are annotated by

MEBS with Interproscan (105) and are then searched with HMM profiles for genes related to

sulfur metabolism. The sulfur metabolism related genes investigated by MEBS are based

primarily on the MetaCyc database (106). Traitar is a software package that can predict 67

phenotypic traits from a genome sequence. In brief, the analysis is based on known phenotypic

traits of 234 bacterial species and infers from their genome Pfam families that are either present

or absent in a specific trait. In this manuscript the traits gelatin hydrolysis and H_2S production

are of interest and these are based the presence of 70 and 43 and absence of 51 and 22 Pfam

families respectively. Phylogenomic trees were calculated with FastTree (107) as maximum

likelihood trees and visualized with iTOL (108, 109) and heatmaps were visualized with

Heatmapper (110). Gene cluster maps were drawn with Gene Graphics (111). Signal peptides

were predicted with SignalP-5.0 server using nucleotide sequences to predict the presence of

Sec/SPI, Tat/SPI and Sec/SPII signal peptides in a given sequence (112).

Enzyme profiling

Based on the organic composition of bone matrix, we hypothesized twelve enzyme families to be

necessary for its degradation. Accordingly, the following enzymes were selected for in-depth

studies: (i) M9 collagenases (pfam01752), S1 peptidases (COG0265), S8/S53 peptidases

(pfam00082) and U32 proteases (COG0826) which hydrolyze peptide bonds in collagen and

glycoproteins; (ii) sialidases (COG4409), β-D-glucuronidases (COG3250), β-N-acetyl-D-

glucosaminidases (COG1472), α-N-acetylgalactosaminidases (COG0673), α-galactosidases

(pfam16499), fucosidases (COG3669) and mannosidases (COG0383) which cleave glycosidic

lingakes, and (iii) cholesterol oxidases (COG2303) which degrade lipids such as cholesterol. One

reference database for each of these families was generated using the NCBI repository, based on

sequences from 287 M9 collagenases, 4453 S1 peptidases, 3237 S8/S53 peptidases, 3653 U32

proteases, 267 COG4409, 873 COG3250, 1274 COG1472, 6140 COG0673, 279 COG3669, 206

COG0383 and 1119 COG2303. The databases included the closest protein homologs of all

protein families of interest for bone-degradation, and at least one representative sequence from

all taxonomic groups (containing such enzymes) was represented. The reference databases were

used to generate Hidden Markov Model (HMM) profiles for each enzyme family with HMMER

version 3.1b1 (113) using the *hmmbuild* option after an alignment of each sequence set was built

with Clustal W version 2.1 (114). The MAGs were screened for the twelve enzyme families of

interest using the generated HMM profiles using HMMer version 3.1b1 with the *hmmsearch*

option and a bitscore threshold of 100.

Tara Oceans comparison

This comparison is based on the work of Meren *et al.* (2018), who binned 93 metagenomes

generated from the Tara Oceans project (115). The metagenomes represent 61 surface water

samples and 32 samples from the deep chlorophyll maximum layer of the water column. They

generated 957 non-redundant high-quality bacterial, archaeal and eukaryotic genomes. The 957

MAGs were here reanalyzed with GTDB-Tk and the generated HMM profiles as previously

described. 832 MAGs were identified as of bacterial origin and included in the comparison. The

Tara Oceans MAGs were generated with a quality threshold of >70% completion, therefore the

bone metagenomes were rebinned (>70% completion, <10% redundancy) for better comparison

to avoid bias due to the higher threshold used for functional analysis in other parts of this

manuscript.

Data availability

The raw sequencing reads have been deposited in the sequence read archive (SRA) of NCBI

under the BioProject ID PRJNA606180 and with the BioSample accession numbers

SAMN14086998 (A5), SAMN14087000 (A9), SAMN14087001 (A9n), SAMN14087003 (B4),

SAMN14087005 (D1), SAMN14087006 (D2), SAMN14087007 (I1) and SAMN14087008 (I3).

The 59 high-quality MAGs analyzed in this study were deposited in the NCBI database as well,

as part of the BioProject ID PRJNA606180, the BioSample accession numbers are

SAMN16086327-SAMN16086385 (Biofilm MAGs 1-44 SAMN16086327-SAMN16086370 and

Osedax MAGs 1-15 SAMN16086371-SAMN16086385).

Authors contributions

T.D., E.B., A.G.M. and G.E.K.B. developed the conceptual idea, conducted the sampling of the source material, contributed to content, reviewed and edited the manuscript. S.S.C., E.B. and M.F. identified enzymes of interest, constructed the reference databases and aided in result interpretation. E.B., B.S., S.F. and A.G.M. conducted bioinformatic analyses and E.B., B.S. and U.H. wrote the manuscript.

Acknowledgements

To the memory of Prof. Hans Tore Rapp and his effort on characterizing *Osedax mucofloris*. We acknowledge financial support of ERA-NET Marine Biotechnology (GA no.: 604814) funded under the FP7 ERA-NET scheme and nationally managed from the German Federal Ministry of Education and Research and Norwegian Research Council. M.F. acknowledges the support of the grants PCIN-2017-078 (within the Marine Biotechnology ERA-NET) and BIO2017-85522-R from the Ministerio de Economía y Competitividad, Ministerio de Ciencia, Innovación y Universidades (MCIU), Agencia Estatal de Investigación (AEI), Fondo Europeo de Desarrollo Regional (FEDER) and Competiveness. Additional funding was received from the Norwegian Biodiversity Information Centre (PA 809116 knr. 47-14). We thank Hans T. Kleivdal for early developments of the concept. We thank Norilia AS for supplying bone residue material for the field work and ROV AS for providing underwater services during the deployment and sampling campaign. The authors acknowledge Kira S. Makarova at the National Center for Biotechnology Information (NCBI, Bethesda MD, USA) for help during the design of the reference database of enzyme families relevant to bone degradation.

References

<u>References</u>

1.   Debashish G, Malay S, Barindra S, Joydeep M. 2005. Marine enzymes. Adv Biochem
Eng Biotechnol 96:189-218.

2.   Imhoff JF, Labes A, Wiese J. 2011. Bio-mining the microbial treasures of the ocean: new
natural products. Biotechnol Adv 29:468-82.

3.   Trincone A. 2011. Marine biocatalysts: enzymatic features and applications. Mar Drugs
9:478-99.

4.   Adams MW, Perler FB, Kelly RM. 1995. Extremozymes: expanding the limits of
biocatalysis. Biotechnology (N Y) 13:662-8.

5.   Aertsen A, Meersman F, Hendrickx ME, Vogel RF, Michiels CW. 2009. Biotechnology
under high pressure: applications and implications. Trends Biotechnol 27:434-41.

6.   Feller G, Gerday C. 2003. Psychrophilic enzymes: hot topics in cold adaptation. Nat Rev
Microbiol 1:200-8.

7.   Alcaide M, Tchigvintsev A, Martínez-Martínez M, Popovic A, Reva ON, Lafraya Á,
Bargiela R, Nechitaylo TY, Matesanz R, Cambon-Bonavita MA, Jebbar M, Yakimov
MM, Savchenko A, Golyshina OV, Yakunin AF, Golyshin PN, Ferrer M, Consortium M.
2015. Identification and characterization of carboxyl esterases of gill chamber-associated
microbiota in the deep-sea shrimp *Rimicaris exoculata* by using functional
metagenomics. Appl Environ Microbiol 81:2125-36.

8.   Kennedy J, O'Leary ND, Kiran GS, Morrissey JP, O'Gara F, Selvin J, Dobson AD. 2011.
Functional metagenomic strategies for the discovery of novel enzymes and biosurfactants
with biotechnological applications from marine ecosystems. J Appl Microbiol 111:787-
99.

9.   Popovic A, Tchigvintsev A, Tran H, Chernikova TN, Golyshina OV, Yakimov MM,

Golyshin PN, Yakunin AF. 2015. Metagenomics as a tool for enzyme discovery:

hydrolytic enzymes from marine-related metagenomes. Adv Exp Med Biol 883:1-20.

10.   Jørgensen BB, Boetius A. 2007. Feast and famine--microbial life in the deep-sea bed. Nat

Rev Microbiol 5:770-81.

11.   Naganuma T. 2000. Microbes on the edge of global biosphere. Biol Sci Space 14:323-31.

12.   Sogin ML, Morrison HG, Huber JA, Mark Welch D, Huse SM, Neal PR, Arrieta JM,

Herndl GJ. 2006. Microbial diversity in the deep sea and the underexplored "rare

biosphere". Proc Natl Acad Sci U S A 103:12115-20.

13.   Smith C, Baco A. 2003. Ecology of whale falls at the deep-sea floor, p 311-354. *In*

Gibson R, Atkinson R (ed), Oceanography and Marine Biology: an Annual Review, vol

41. Taylor&Francis.

14.   Rouse GW, Goffredi SK, Vrijenhoek RC. 2004. *Osedax*: bone-eating marine worms with

dwarf males. Science 305:668-71.

15.   Miyamoto N, Yoshida MA, Koga H, Fujiwara Y. 2017. Genetic mechanisms of bone

digestion and nutrient absorption in the bone-eating worm *Osedax japonicus* inferred

from transcriptome and gene expression analyses. BMC Evol Biol 17:17.

16.   Tresguerres M, Katz S, Rouse GW. 2013. How to get into bones: proton pump and

carbonic anhydrase in *Osedax* boneworms. Proc Biol Sci 280:20130625.

17.   Higgs ND, Glover AG, Dahlgren TG, Little CT. 2011. Bone-boring worms:

characterizing the morphology, rate, and method of bioerosion by *Osedax mucofloris*

(Annelida, Siboglinidae). Biol Bull 221:307-16.

18. Goffredi SK, Yi H, Zhang Q, Klann JE, Struve IA, Vrijenhoek RC, Brown CT. 2014.

Genomic versatility and functional variation between two dominant heterotrophic

symbionts of deep-sea *Osedax* worms. ISME J 8:908-24.

19. Goffredi SK, Orphan VJ, Rouse GW, Jahnke L, Embaye T, Turk K, Lee R, Vrijenhoek

RC. 2005. Evolutionary innovation: a bone-eating marine symbiosis. Environ Microbiol

7:1369-78.

20. Goffredi SK, Johnson SB, Vrijenhoek RC. 2007. Genetic diversity and potential function

of microbial symbionts associated with newly discovered species of *Osedax* polychaete

worms. Appl Environ Microbiol 73:2314-23.

21. Verna C, Ramette A, Wiklund H, Dahlgren TG, Glover AG, Gaill F, Dubilier N. 2010.

High symbiont diversity in the bone-eating worm *Osedax mucofloris* from shallow

whale-falls in the North Atlantic. Environ Microbiol 12:2355-70.

22. Glover AG, Källström B, Smith CR, Dahlgren TG. 2005. World-wide whale worms? A

new species of *Osedax* from the shallow north Atlantic. Proc Biol Sci 272:2587-92.

23. Dahlgren T, Wiklund H, Källström B, Lundälv T, Smith C, Glover A. 2006. A shallow-

water whale-fall experiment in the North Atlantic. Cahiers De Biologie Marine 47:385-

389.

24. Robey PG. 2008. Noncollagenous bone matrix proteins, p 335-349. *In* Bilezikian JP,

Raisz LG, Martin JT (ed), Principles of bone biology, Third edition ed, vol Volume I.

Academic press.

25. Sato S, Rahemtulla F, Prince CW, Tomana M, Butler WT. 2009. Acidic glycoproteins

from bovine compact bone. Connective Tissue Research 14:51-64.

26.   Kuboki Y, Watanabe T, Tazaki M, Taktta H. 1991. Comparative biochemistry of bone

matrix proteins in bovine and fish. *In* Suga S, Nakahara H (ed), Mechanisms and

phylogeny of mineralization in biological systems doi:https://doi.org/10.1007/978-4-431-

68132-8_79. Springer, Tokyo.

27.   Shoulders MD, Raines RT. 2009. Collagen structure and stability. Annu Rev Biochem

78:929-58.

28.   de Gonzalo G, Colpa DI, Habib MH, Fraaije MW. 2016. Bacterial enzymes involved in

lignin degradation. J Biotechnol 236:110-9.

29.   Puentes-Tellez PE, Salles JF. 2018. Construction of effective minimal active microbial

consortia for lignocellulose degradation. Microbial Ecology 76:419-429.

30.   Vietti L, Bailey J, Ricci E. 2014. Insights into the microbial degradation of bone in

marine environments from rRNA gene sequencing of biofilms on lab-simulated carcass-

falls. The Paleontological Society Special Publications

doi:doi:10.1017/S2475262200012582:120-121.

31.   Vietti LA. 2014. Insights into the microbial degradation of bones from the marine

vertebrate fossil record: an experimental approach using interdisciplinary analyses. Ph.D.

dissertation. University of Minnesota, University of Minnesota Digital Conservancy.

32.   Taboada S, Bas M, Avila C, Riesgo A. 2020. Phylogenetic characterization of marine

microbial biofilms associated with mammal bones in temperate and polar areas. Marine

Biodiversity 50.

33.   Hewitt OH, Diez-Vives C, Taboada S. 2020. Microbial insights from Antarctic and

Mediterranean shallow-water bone-eating worms. Polar Biology 43:1605-1621.

34.   Freitas RC, Marques HIF, Silva MACD, Cavalett A, Odisi EJ, Silva BLD, Montemor JE,

Toyofuku T, Kato C, Fujikura K, Kitazato H, Lima AOS. 2019. Evidence of selective

pressure in whale fall microbiome proteins and its potential application to industry. Mar

Genomics 45:21-27.

35.   Menzel P, Ng KL, Krogh A. 2016. Fast and sensitive taxonomic classification for

metagenomics with Kaiju. Nature Communications 7.

36.   Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil PA,

Hugenholtz P. 2018. A standardized bacterial taxonomy based on genome phylogeny

substantially revises the tree of life. Nat Biotechnol 36:996-1004.

37.   De Anda V, Zapata-Peñasco I, Poot-Hernandez AC, Eguiarte LE, Contreras-Moreira B,

Souza V. 2017. MEBS, a software platform to evaluate large (meta)genomic collections

according to their metabolic machinery: unraveling the sulfur cycle. Gigascience 6:1-17.

38.   Waite DW, Chuvochina M, Pelikan C, Parks DH, Yilmaz P, Wagner M, Loy A,

Naganuma T, Nakai R, Whitman WB, Hahn MW, Kuever J, Hugenholtz P. 2020.

Proposal to reclassify the proteobacterial classes Deltaproteobacteria and Oligoflexia, and

the phylum Thermodesulfobacteria into four phyla reflecting major functional

capabilities. Int J Syst Evol Microbiol doi:10.1099/ijsem.0.004213.

39.   Müller AL, Kjeldsen KU, Rattei T, Pester M, Loy A. 2015. Phylogenetic and

environmental diversity of DsrAB-type dissimilatory (bi)sulfite reductases. ISME J

9:1152-65.

40.   Weimann A, Mooren K, Frank J, Pope PB, Bremges A, McHardy AC. 2016. From

genomes to phenotypes: Traitar, the Microbial Trait Analyzer. mSystems 1.

41.   Yamamoto M, Takai K. 2011. Sulfur metabolisms in epsilon- and gamma-proteobacteria

in deep-sea hydrothermal fields. Front Microbiol 2:192.

42.   Muyzer G, Stams AJ. 2008. The ecology and biotechnology of sulphate-reducing

bacteria. Nat Rev Microbiol 6:441-54.

43.   Barton LL, Fauque GD. 2009. Biochemistry, physiology and biotechnology of sulfate-

reducing bacteria. Adv Appl Microbiol 68:41-98.

44.   Kruse S, Goris T, Westermann M, Adrian L, Diekert G. 2018. Hydrogen production by

Sulfurospirillum species enables syntrophic interactions of Epsilonproteobacteria. Nat

Commun 9:4872.

45.   Schulz HN, Jorgensen BB. 2001. Big bacteria. Annu Rev Microbiol 55:105-37.

46.   Gevertz D, Telang AJ, Voordouw G, Jenneman GE. 2000. Isolation and characterization

of strains CVO and FWKO B, two novel nitrate-reducing, sulfide-oxidizing bacteria

isolated from oil field brine. Appl Environ Microbiol 66:2491-501.

47.   Trojan D, Schreiber L, Bjerg JT, Bøggild A, Yang T, Kjeldsen KU, Schramm A. 2016. A

taxonomic framework for cable bacteria and proposal of the candidate genera

Electrothrix and *Electronema*. Syst Appl Microbiol 39:297-306.

48.   Pfeffer C, Larsen S, Song J, Dong M, Besenbacher F, Meyer RL, Kjeldsen KU, Schreiber

L, Gorby YA, El-Naggar MY, Leung KM, Schramm A, Risgaard-Petersen N, Nielsen

LP. 2012. Filamentous bacteria transport electrons over centimetre distances. Nature

491:218-21.

49.   Capasso C, Supuran CT. 2015. An overview of the alpha-, beta- and gamma-carbonic

anhydrases from Bacteria: can bacterial carbonic anhydrases shed new light on evolution

of bacteria? J Enzyme Inhib Med Chem 30:325-32.

50. Bowman JP, Gosnik JJ, McCammon SA, Lewis TE, Nichols DS, Nichols PD, Skerratt H,

Staley JT, McMeekin TA. 1998. *Colwellia demingiae* sp. nov., *Colwellia hornerae* sp.

nov., *Colwellia rossensis* sp. nov. and *Colwellia psychrotropica* sp. nov.: psychrophilic

Antarctic species with the ability to synthesize docosahexaenoic acid (22:ω63).

International Journal of Systematic and Evolutionary Microbiology 48:1171-1180.

51. Bowman JP, McCammon SA, Brown MV, Nichols DS, McMeekin TA. 1997. Diversity

and association of psychrophilic bacteria in Antarctic sea ice. Appl Environ Microbiol

63:3068-78.

52. Campeao ME, Swings J, Silva BS, Otsuki K, Thompson FL, Thompson CC. 2019.

"*Candidatus Colwellia aromaticivorans*" sp. nov., "*Candidatus Halocyntiibacter*

alkanivorans" sp. nov., and "*Candidatus Ulvibacter alkanivorans*" sp. nov. Genome

Sequences. Microbiol Resour Announc 8.

53. Christiansen L, Bech PK, Schultz-Johansen M, Martens HJ, Stougaard P. 2018. *Colwellia*

echini sp. nov., an agar- and carrageenan-solubilizing bacterium isolated from sea urchin.

Int J Syst Evol Microbiol 68:687-691.

54. Jung SY, Oh TK, Yoon JH. 2006. *Colwellia aestuarii* sp. nov., isolated from a tidal flat

sediment in Korea. Int J Syst Evol Microbiol 56:33-7.

55. Kim YO, Park IS, Park S, Nam BH, Jung YT, Kim DG, Yoon JH. 2017. *Colwellia mytili*

sp. nov., isolated from mussel *Mytilus edulis*. Int J Syst Evol Microbiol 67:31-36.

56. Kusube M, Kyaw TS, Tanikawa K, Chastain RA, Hardy KM, Cameron J, Bartlett DH.

2017. *Colwellia marinimaniae* sp. nov., a hyperpiezophilic species isolated from an

amphipod within the Challenger Deep, Mariana Trench. Int J Syst Evol Microbiol

67:824-831.

57.   Nogi Y, Hosoya S, Kato C, Horikoshi K. 2004. *Colwellia piezophila* sp. nov., a novel

767   piezophilic species from deep-sea sediments of the Japan Trench. Int J Syst Evol

768   Microbiol 54:1627-1631.

769   58.   Park S, Jung YT, Yoon JH. 2016. *Colwellia sediminilitoris* sp. nov., isolated from a tidal

770   flat. Int J Syst Evol Microbiol 66:3258-3263.

771   59.   Techtmann SM, Fitzgerald KS, Stelling SC, Joyner DC, Uttukar SM, Harris AP, Alshibli

772   NK, Brown SD, Hazen TC. 2016. *Colwellia psychrerythraea* strains from distant deep

773   sea basins show adaptation to local conditions. Frontiers in Environmental Science 4.

774   60.   Xu ZX, Zhang HX, Han JR, Dunlap CA, Rooney AP, Mu DS, Du ZJ. 2017. *Colwellia*

775   *agarivorans* sp. nov., an agar-digesting marine bacterium isolated from coastal seawater.

776   Int J Syst Evol Microbiol 67:1969-1974.

777   61.   Yu Y, Li HR, Zeng YX. 2011. *Colwellia chukchiensis* sp. nov., a psychrotolerant

778   bacterium isolated from the Arctic Ocean. Int J Syst Evol Microbiol 61:850-853.

779   62.   Zhang C, Guo W, Wang Y, Chen X. 2017. *Colwellia beringensis* sp. nov., a

780   psychrophilic bacterium isolated from the Bering Sea. Int J Syst Evol Microbiol 67:5102-

781   5107.

782   63.   Zhang DC, Yu Y, Xin YH, Liu HC, Zhou PJ, Zhou YG. 2008. *Colwellia polaris* sp. nov.,

783   a psychrotolerant bacterium isolated from Arctic sea ice. Int J Syst Evol Microbiol

784   58:1931-4.

785   64.   Delmont TO, Quince C, Shaiber A, Esen OC, Lee STM, Rappe MS, McLellan SL,

786   Lucker S, Eren AM. 2018. Nitrogen-fixing populations of Planctomycetes and

787   Proteobacteria are abundant in surface ocean metagenomes (vol 3, pg 804, 2018). Nature

788   Microbiology 3.

789   65.   Gontang EA, Fenical W, Jensen PR. 2007. Phylogenetic diversity of gram-positive

790   bacteria cultured from marine sediments. Appl Environ Microbiol 73:3272-82.

791   66.   Teske A, Brinkhoff T, Muyzer G, Moser DP, Rethmeier J, Jannasch HW. 2000. Diversity

792   of thiosulfate-oxidizing bacteria from marine sediments and hydrothermal vents. Appl

793   Environ Microbiol 66:3125-33.

794   67.   Leloup J, Fossing H, Kohls K, Holmkvist L, Borowski C, Jørgensen BB. 2009. Sulfate-

795   reducing bacteria in marine sediment (Aarhus Bay, Denmark): abundance and diversity

796   related to geochemical zonation. Environ Microbiol 11:1278-91.

797   68.   Cotter PD, Hill C. 2003. Surviving the acid test: responses of gram-positive bacteria to

798   low pH. Microbiol Mol Biol Rev 67:429-53, table of contents.

799   69.   Anné J, Economou A, Bernaerts K. 2017. Protein secretion in gram-positive bacteria:

800   From multiple pathways to biotechnology. Curr Top Microbiol Immunol 404:267-308.

801   70.   Jang H, Yang SH, Seo HS, Lee JH, Kim SJ, Kwon KK. 2015. *Amphritea spongicola* sp

802   nov., isolated from a marine sponge, and emended description of the genus *Amphritea*.

803   International Journal of Systematic and Evolutionary Microbiology 65:1866-1870.

804   71.   Wang JH, Lu Y, Nawaz MZ, Xu J. 2018. Comparative genomics reveals evidence of

805   genome reduction and high extracellular protein degradation potential in *Kangiella*.

806   Frontiers in Microbiology 9.

807   72.   Cao SN, He JF, Zhang F, Lin L, Gao Y, Zhou QM. 2019. Diversity and community

808   structure of bacterioplankton in surface waters off the northern tip of the Antarctic

809   Peninsula. Polar Research 38.

73.   Lastovica A, On S, Zhang L. 2014. The family *Campylobacteraceae*, p 307-335. *In*

Rosenburg E, DeLong E, Lory S, Stackebrandt E, Thompson F (ed), The Prokaryotes

doi:https://doi.org/10.1007/978-3-642-39044-9_274. Springer.

74.   Takai K, Campbell BJ, Cary SC, Suzuki M, Oida H, Nunoura T, Hirayama H, Nakagawa

S, Suzuki Y, Inagaki F, Horikoshi K. 2005. Enzymatic and genetic characterization of

carbon and energy metabolisms by deep-sea hydrothermal chemolithoautotrophic isolates

of Epsilonproteobacteria. Appl Environ Microbiol 71:7310-20.

75.   Deming JW, Reysenbach AL, Macko SA, Smith CR. 1997. Evidence for the microbial

basis of a chemoautotrophic invertebrate community at a whale fall on the deep seafloor:

bone-colonizing bacteria and invertebrate endosymbionts. Microsc Res Tech 37:162-70.

76.   Supuran CT, Capasso C. 2017. An overview of the bacterial carbonic anhydrases.

Metabolites 7.

77.   Vietti L, Bailey J, Fox D, Rogers R. 2015. Rapid formation of framboidal sulfides on

bone surfaces from a simulated marine carcass fall. PALAIOS 30:327-334.

78.   Naretto A, Fanuel M, Ropartz D, Rogniaux H, Larocque R, Czjzek M, Tellier C, Michel

G. 2019. The agar-specific hydrolase. J Biol Chem 294:6923-6939.

79.   Nedashkovskaya OI, Kim SG, Stenkova AM, Kukhlevskiy AD, Zhukova NV, Mikhailov

VV. 2018. *Aquimarina algiphila* sp. nov., a chitin degrading bacterium isolated from the

red alga *Tichocarpus crinitu*s. Int J Syst Evol Microbiol 68:892-898.

80.   Konasani VR, Jin C, Karlsson NG, Albers E. 2018. A novel ulvan lyase family with

broad-spectrum activity from the ulvan utilisation loci of *Formosa agariphila* KMM

3901. Sci Rep 8:14713.

81.   Li S, Wang L, Chen X, Zhao W, Sun M, Han Y. 2018. Cloning, expression, and

833   biochemical characterization of two new oligoalginate lyases with synergistic

834   degradation capability. Mar Biotechnol (NY) 20:75-86.

835   82.   Shen J, Chang Y, Chen F, Dong S. 2018. Expression and characterization of a κ-

836   carrageenase from marine bacterium *Wenyingzhuangia aestuarii* OF219: A

837   biotechnological tool for the depolymerization of κ-carrageenan. Int J Biol Macromol

838   112:93-100.

839   83.   Tan H, Miao R, Liu T, Yang L, Yang Y, Chen C, Lei J, Li Y, He J, Sun Q, Peng W, Gan

840   B, Huang Z. 2018. A bifunctional cellulase-xylanase of a new *Chryseobacterium* strain

841   isolated from the dung of a straw-fed cattle. Microb Biotechnol 11:381-398.

842   84.   Rochat T, Pérez-Pascual D, Nilsen H, Carpentier M, Bridel S, Bernardet JF, Duchaud E.

843   2019. Identification of a novel elastin-degrading enzyme from the fish pathogen. Appl

844   Environ Microbiol 85.

845   85.   Choi KD, Lee GE, Park JS. 2018. *Aquimarina spongiicola* sp. nov., isolated from

846   spongin. Int J Syst Evol Microbiol 68:990-994.

847   86.   Iino T, Mori K, Itoh T, Kudo T, Suzuki K, Ohkuma M. 2014. Description of

848   *Mariniphaga anaerophila* gen. nov., sp. nov., a facultatively aerobic marine bacterium

849   isolated from tidal flat sediment, reclassification of the Draconibacteriaceae as a later

850   heterotypic synonym of the Prolixibacteraceae and description of the family

851   Marinifilaceae fam. nov. Int J Syst Evol Microbiol 64:3660-7.

852   87.   Han Z, Shang-Guan F, Yang J. 2019. Molecular and biochemical characterization of a

853   bimodular xylanase From. Front Microbiol 10:1507.

88. Emmons A, Mundorff A, Keenan S, Davoren J, Andronowski J, Carter D, DeBruyn J.

2019. Patterns of microbial colonization of human bone from surface-decomposed

remains. bioRxiv 664482.

89. Robey P. 2002. Bone matrix proteoglycans and glycoproteins. *In* Bilezikian J, Raisz L,

Rodan G (ed), Principles of bone biology, 2nd Edition ed

doi:https://doi.org/10.1016/B978-012098652-1.50116-5. Elsevier.

90. van Vliet DM, Palakawong Na Ayudthaya S, Diop S, Villanueva L, Stams AJM,

Sánchez-Andrea I. 2019. Anaerobic degradation of sulfated polysaccharides by two

novel. Front Microbiol 10:253.

91. S D, O G. 2016. *Colwellia* and sulfur-oxidizing bacteria: An unusual dual symbiosis in a

Terua mussel (Mytilidae: Bathymodiolinae) from whale falls in the Antilles arc. Deep

Sea Research Part I 115:112-122.

92. Hansen AM, Qiu Y, Yeh N, Blattner FR, Durfee T, Jin DJ. 2005. SspA is required for

acid resistance in stationary phase by downregulation of H-NS in *Escherichia coli*. Mol

Microbiol 56:719-34.

93. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics 30:2114-20.

94. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,

Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G,

Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its

applications to single-cell sequencing. J Comput Biol 19:455-77.

95. Uritskiy GV, DiRuggiero J, Taylor J. 2018. MetaWRAP-a flexible pipeline for genome-

resolved metagenomic data analysis. Microbiome 6:158.

96. Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, Ijaz UZ, Lahti L, Loman

NJ, Andersson AF, Quince C. 2014. Binning metagenomic contigs by coverage and

composition. Nat Methods 11:1144-6.

97. Wu YW, Simmons BA, Singer SW. 2016. MaxBin 2.0: an automated binning algorithm

to recover genomes from multiple metagenomic datasets. Bioinformatics 32:605-7.

98. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, Wang Z. 2019. MetaBAT 2: an

adaptive binning algorithm for robust and efficient genome reconstruction from

metagenome assemblies. PeerJ 7:e7359.

99. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM:

assessing the quality of microbial genomes recovered from isolates, single cells, and

metagenomes. Genome Res 25:1043-55.

100. Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal:

prokaryotic gene recognition and translation initiation site identification. BMC

Bioinformatics 11:119.

101. Huerta-Cepas J, Forslund K, Coelho LP, Szklarczyk D, Jensen LJ, von Mering C, Bork P.

2017. Fast genome-wide functional annotation through orthology assignment by

eggNOG-Mapper. Mol Biol Evol 34:2115-2122.

102. Huerta-Cepas J, Szklarczyk D, Forslund K, Cook H, Heller D, Walter MC, Rattei T,

Mende DR, Sunagawa S, Kuhn M, Jensen LJ, von Mering C, Bork P. 2016. eggNOG 4.5:

a hierarchical orthology framework with improved functional annotations for eukaryotic,

prokaryotic and viral sequences. Nucleic Acids Res 44:D286-93.

103. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S,

Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil

LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O,

Vonstein V, Wilke A, Zagnitko O. 2008. The RAST Server: rapid annotations using

subsystems technology. BMC Genomics 9:75.

104.   Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S,

Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and

the Rapid Annotation of microbial genomes using Subsystems Technology (RAST).

Nucleic Acids Res 42:D206-14.

105.   Quevillon E, Silventoinen V, Pillai S, Harte N, Mulder N, Apweiler R, Lopez R. 2005.

InterProScan: protein domains identifier. Nucleic Acids Research 33:W116-W120.

106.   Caspi R, Billington R, Ferrer L, Foerster H, Fulcher CA, Keseler IM, Kothari A,

Krummenacker M, Latendresse M, Mueller LA, Ong Q, Paley S, Subhraveti P, Weaver

DS, Karp PD. 2016. The MetaCyc database of metabolic pathways and enzymes and the

BioCyc collection of pathway/genome databases. Nucleic Acids Research 44:D471-

D480.

107.   Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-likelihood

trees for large alignments. PLoS One 5:e9490.

108.   Letunic I, Bork P. 2007. Interactive Tree Of Life (iTOL): an online tool for phylogenetic

tree display and annotation. Bioinformatics 23:127-8.

109.   Letunic I, Bork P. 2019. Interactive Tree Of Life (iTOL) v4: recent updates and new

developments. Nucleic Acids Res 47:W256-W259.

110.   Babicki S, Arndt D, Marcu A, Liang Y, Grant JR, Maciejewski A, Wishart DS. 2016.

Heatmapper: web-enabled heat mapping for all. Nucleic Acids Res 44:W147-53.

111.   Harrison KJ, Crécy-Lagard V, Zallot R. 2018. Gene Graphics: a genomic neighborhood

923   data visualization web application. Bioinformatics 34:1406-1408.

924   112.   Almagro Armenteros JJ, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak S,

925   von Heijne G, Nielsen H. 2019. SignalP 5.0 improves signal peptide predictions using

926   deep neural networks. Nat Biotechnol 37:420-423.

927   113.   Eddy SR. 2011. Accelerated Profile HMM Searches. PLoS Comput Biol 7:e1002195.

928   114.   Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H,

929   Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG.

930   2007. Clustal W and Clustal X version 2.0. Bioinformatics 23:2947-8.

931   115.   Delmont TO, Quince C, Shaiber A, Esen OC, Lee STM, Rappe MS, MacLellan SL,

932   Lucker S, Eren AM. 2018. Nitrogen-fixing populations of Planctomycetes and

933   Proteobacteria are abundant in surface ocean metagenomes. Nature Microbiology 3:804-

934   +.

943   **Tables:**

944   **Table 1:** Metagenome sampling information and number of retrieved metagenome assembled

945   genomes (MAG).

Sample	Sample type	Bone type, organism	Collection dates	Sampling location (GPS)	High quality MAGs
A5 A9 A9n B4	*Osedax mucofloris*	Femur, turkey (*Meleagris gallopavo*)	08.01.2017 08.02.2017 08.02.2017 14.04.2017	Byfjorden, Bergen, Norway (60,397093N; 5,301293E)	15 (OB)
D1 D2 I1 I3	Bone surface biofilm communities	Tibia, cow (*Bos taurus*)	02.2017 11.12.2017 27.01.2017 11.12.2017		44 (BB)

Figures

Figure 1: Phylogenomic maximum likelihood tree of the obtained 59 high-quality MAGs from *Osedax*-associated bone microbiome (OB) and bone surface-associated biofilm (BB). Bootstrap values greater than 0.5 are displayed as circles on the branches. The five most common bacterial classes are colored (green – Bacteroidia, light purple – Desulfuromonadia, purple – Campylobacteria, orange – Alphaproteobacteria and blue – Gammaproteobacteria). Order level identifications are listed and MAGs for which only class level identification could be inferred are marked with an asterisk.

Figure 2: Whole genome metabolic pathway comparison for genes of the sulfur metabolism. Analysis was done with MEBS (37) and MAGs were phylogenetically grouped according to GTDBTk pipeline (36). The grey scale represents the completeness of a given pathway or multi-enzyme system shown in the heatmap for each MAG. The OB MAGs are highlighted in blue. The "Sox system" constitutes of the *sox*XYZABCD genes and the "Sor system" of *sor*ABDE.

Figure 3: Abundance and taxonomic affiliation of predicted gelatin hydrolysers (green) (Traitar) in *Osedax* and biofilm derived MAGs. Additionally, sulfur reducers (blue) and sulfur oxidizers (orange), (MEBS) are shown, while white indicates absence of these traits. MAGs are displayed at the deepest taxonomic classification obtained. The size of the circles reflects the number of MAGs within each clade.

Figure 4: Abundance heatmap of the 12 investigated enzyme COG classes in the 59 bone degradome MAGs. The MAGs are arranged according to their taxonomic affiliation. The absolute abundance of each enzyme COG class is depicted in the diagram on top of the heatmap. The OB MAGs are highlighted in blue.

Figure 5: Collagen utilization pathway scheme in MAG BB5. A) The gene cluster in BB5 spans

approximately 21 kb, comprising 15 genes for collagen utilization, each color-coded respective

to its functional group: orange for collagen hydrolysis, blue for uptake and transport, green for

proline (Pro) utilization, ocher for hydroxyproline (Hyp) utilization and brown for unknown

function. The purple box is indicative of a signal-peptide for secretion. B) Metabolic model for

collagen utilization in *Colwellia* BB5. Arrows and genes are color-coded in the same functional

groups as in A. Dashed arrows point to a major metabolic pathway. Metabolite abbreviations:

P4C (1-pyrroline 4-hydroxy-2-carboxylate), KGSA (alpha-ketoglutarate semialdehyde), KG

(alpha-ketoglutarate), P5C (1-pyrroline-5-carboxylate). Enzyme abbreviations: D-aa dHase (D-

hydroxyproline dehydrogenase), dAminase (pyrroline-4-hydroxy-2-carboxylate deaminase), di-

oxo dHase (KGSA dehydrogenase), P5CR/ornithine dAminase (bifunctional 1-pyrroline-5-

carboxylate reductase/ornithine cyclodeaminase), PRODH (proline dehydrogenase), P5CDH

(pyrroline-5-carboxylate dehydrogenase).

Figure 6: Conservation between M9 collagen degradation gene clusters in *Colwellia*

psychrerythraea GAB14E, *Colwellia piezophila* ATCC BAA-637 and the MAG *Colwellia* BB5

drawn at scale. dHase, dehydrogenase; PPIase, peptidyl-prolyl cis trans isomerase; Hyp, D-aa,

dAminase etc. Color coding and gene names are indicated.

Figure 7: Hypothesis of the interplay in the marine bone microbiome and degradome. Sulfur-

oxidizing bacteria (SOB, shown with a halo) convert elemental sulfur and H_2S into sulfate and

protons that lead to an acidification and therefore bone demineralization. Sulfate reducing (SRB,

green) and sulfur disproportioning bacteria produce H_2S from sulfate. Enterobacterales and other

especially Gammaproteobatceria secret collagenases to degrade collagen. Bacteroidia and other

bacteria secret gylcosidases and other enzymes to hydrolyze the organic bone components

(glycosides, esters, lipids). This exemplifies a bone demineralization loop that fuels itself as long

as sulfur is available and degrades the organic bone components in the process.

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017 **Supplementary material**

1018 **Supplementary tables**

1019 **Supplementary table S1:** Metagenome statistics

	Reads			Individual statistics				Combined statistics			Taxonomic distribution of raw reads (Kaiju)		
	raw [million]	filtered [million]	dereplicated [million]	size [Mb]	contigs	N50	largest contig	size [Mb]	contigs	N50	Bacteria [%]	Eukaryota (fungi) [%]	Archaea [%]
A5	24.8	18.75	13.32	214.6	185.1k	1175	1645k				76	15 (4)	0.40
A9	24.9	18.71	12.8	170.9	169k	1007	156k	342.7	309.5k	1115	45	34 (9)	0.40
A9n	14.4	6.76	5.12	88.6	90.8k	926	299k				45	34 (9)	0.40
B4	19.2	12.65	8.86	120.7	132.7k	910	137k				47	34 (9)	0.40
D1	24.1	20.59	12.16	192.9	121.8k	2543	918k				94	34 (9)	0.40
D2	35.8	32.5	15.67	175.2	156.9k	1134	1234k	1220	1022k	1293	92	34 (9)	0.40
I1	35.9	31.96	26.07	534.3	440.8k	1336	1105k				95	34 (9)	0.40
I3	32.3	29.29	15.48	249.1	227.9k	1136	195k				93	34 (9)	0.40

 Supplementary table S2: Taxonomic affiliation of MAGs according to Parks *et al.* (2018) and

 taxonomic novelty identified by RED (*) (36).

MAG	Phylum	Class	Order	Family	Genus
BB1	Planctomycetota*	-	-	-	-
BB2	Proteobacteria	γ-proteobacteria	Beggiatoales*	-	-
BB3	Proteobacteria	γ-proteobacteria	Beggiatoales	Beggiatoaceae*	-
BB4	Proteobacteria	γ-proteobacteria*	-	-	-
BB5	Proteobacteria	γ-proteobacteria	Enterobacterales	Alteromonadaceae	Colwellia
BB6	Proteobacteria	α-proteobacteria	Rhizobiales*	-	-
BB7	Spirochaetota	Spirochaetia	Spirochaetales	Spirochaetaceae*	-
BB8	Campylobacterota	Camylobacteria	Campylobacterales	-	-
BB9	Campylobacterota	Camylobacteria	Campylobacterales	Sulfurovaceae	Sulfurovum
BB10	Campylobacterota	Camylobacteria	Campylobacterales	Thiovulaceae*	-
BB11	Campylobacterota	Camylobacteria	Campylobacterales	Thiovulaceae	Sulfurimonas
BB12	Fermentibacterota	Fermentibacteria	Fermentibacterales	Fermentibacteraceae*	-
BB13	Desulfobacterota	Desulfobulbia	Desulfobulbales	Desulfobulbaceae*	-
BB14	Campylobacterota	Camylobacteria	Campylobacterales	Sulfurovaceae*	-
BB15	Campylobacterota	Camylobacteria	Campylobacterales	Sulfurospirillaceae	Sulfurospirillum
BB16	Proteobacteria	γ-proteobacteria	Beggiatoales	Beggiatoaceae	Marithrix*
BB17	Bacteroidota	Bacteroidia	Flavobacteriales	Ichthyobacteriaceae*	-
BB18	Krumholzibacteriota	Krumholzibacteria*	-	-	-
BB19	Proteobacteria	γ-proteobacteria	Pseudomonadales	Hahellaceae*	-
BB20	Proteobacteria	γ-proteobacteria	Beggiatoales*	-	-
BB21	Desulfuromonadota	Desulfuromonadia	Desulfuromonadales	Geopsychrobacteraceae	Desulfuromusa
BB22	Bacteroidota	Bacteroidia	Bacteroidales*	-	-
BB23	Bacteroidota	Bacteroidia	Flavobacteriales	Flavobacteriaceae	Winogradskyella
BB24	Bacteroidota	Bacteroidia	Flavobacteriales	Flavobacteriaceae*	-
BB25	Desulfuromonadota	Desulfuromonadia	Desulfuromonadales	Geopsychrobacteraceae	Desulfuromusa
BB26	Campylobacterota	Camylobacteria	Campylobacterales	Thiovulaceae*	-
BB27	Desulfuromonadota	Desulfuromonadia	Desulfuromonadales	Geopsychrobacteraceae*	-
BB28	Campylobacterota	Camylobacteria	Campylobacterales	Arcobacteraceae*	-
BB29	Bacteroidota	Bacteroidia	Bacteroidales	Marinifilaceae	-
BB30	Campylobacterota	Camylobacteria	Campylobacterales	Arcobacteraceae	Arcobacter
BB31	Proteobacteria	γ-proteobacteria	Beggiatoales*	-	-
BB32	Proteobacteria	γ-proteobacteria	Xanthomonadales	Marinicellaceae*	-
BB33	Proteobacteria	α-proteobacteria	Rhodobacterales	Rhodobacteraceae	Lentibacter*
BB34	Proteobacteria	γ-proteobacteria*	-	-	-
BB35	Bacteroidota	Bacteroidia	Flavobacteriales	Flavobacteriaceae	-
BB36	Proteobacteria	γ-proteobacteria*			
BB37	Proteobacteria	γ-proteobacteria	Pseudomonadales	Halieaceae*	-

BB38	Chloroflexota	Anaerolineae	Anearolineales*	-	-
BB39	Krumholzibacteriota	Krumholzibacteria*	-	-	-
BB40	Desulfobacterota	Desulfobacteria	Desulfobacterales	Desulfobacteraceae	-
BB41	Campylobacterota	Camylobacteria	Campylobacterales	Sulfurovaceae*	-
BB42	Bacteroidota	Bacteroidia	Flavobacteriales	Flavobacteriaceae	Maribacter*
BB43	Verrucomicrobiota	Kiritimatiellae	Kiritimatiellales*	-	-
BB44	Proteobacteria	γ-proteobacteria	Enterobacterales	Kangiellaceae*	-
OB1	Proteobacteria	γ-proteobacteria	Pseudomonadales	Nitrincolaceae	Neptunomonas
OB2	Proteobacteria	γ-proteobacteria	Pseudomonadales	Nitrincolaceae	Amphritea
OB3	Desulfobacterota	Desulfovibrionia	Desulfovibrionales	Desulfovibrionaceae	Pseudodesulfovibrio
OB4	Proteobacteria	α-proteobacteria	Sphingomonadales	Emcibacteraceae*	-
OB5	Campylobacterota	Camylobacteria	Campylobacterales	Thiovulaceae	Sulfurimonas
OB6	Proteobacteria	γ-proteobacteria*	-	-	-
OB7	Campylobacterota	Camylobacteria	Campylobacterales	Arcobacteraceae	Arcobacter
OB8	Campylobacterota	Camylobacteria	Campylobacterales	Thiovulaceae	Sulfurimonas
OB9	Proteobacteria	α-proteobacteria	Rhodobacterales	Rhodobacteraceae	-
OB10	Verrucomicrobiota	Kiritimatiellae	Kiritimatiellales*	-	-
OB11	Campylobacterota	Camylobacteria	Campylobacterales	Sulfurospirillaceae	Sulfurospirillum
OB12	Proteobacteria	γ-proteobacteria	Enterobacterales	Kangiellaceae*	-
OB13	Bacteroidota	Bacteroidia	Bacteroidales	Marinifilaceae	Labilibaculum*
OB14	Desulfobacterota	Desulfobacteria	Desulfobacterales	Desulfobacteraceae	-
OB15	Proteobacteria	α-proteobacteria*	-	-	-

1039

Supplementary table S3: MAG sequence data. All MAGs labeled 'OB' were retrieved from

Osedax samples and all MAGs labeled with 'BB' from bone surface biofilms.

MAG	Genome size [Mbp]	Longest contig [kbp]	N50 [kbp]	No. of contigs	Predicted genes	GC content [%]	No. of bone-degrading Enzymes
BB1	4.68	155	429	38	3723	44.50	24
BB2	4.18	853	248	108	3538	44.10	13
BB3	4.65	75	23	362	4126	37.70	11
BB4	5.18	1.050	198	68	4705	38.80	13
BB5	3.26	82	13	395	3018	36.10	16
BB6	2.77	72	19	244	2783	40.10	10
BB7	4.87	78	15	489	4636	40.10	22
BB8	3.08	89	18	413	3332	32.90	11
BB9	2.16	126	36	159	2259	36.80	5
BB10	3.01	137	35	142	2950	36.20	11
BB11	2.39	21	3	809	2958	35.40	6

BB12	3.24	329	141	127	2929	45.10	10
BB13	4.24	16	3	1688	4778	48	16
BB14	2.32	74	33	103	2265	34.20	8
BB15	2.68	415	191	33	2695	30.70	5
BB16	3.09	481	237	22	2867	34.10	13
BB17	3.64	81	16	327	3171	32.30	13
BB18	4.43	267	114	176	3652	52.60	23
BB19	4.07	273	132	88	3778	41.30	13
BB20	4.49	73	19	338	3725	37.90	13
BB21	2.89	145	51	186	2812	44.40	8
BB22	5.23	124	28	294	4210	34.20	32
BB23	3.72	128	30	190	3427	23.40	9
BB24	4.64	57	16	582	4438	31.30	58
BB25	3.31	211	67	118	3069	44.30	8
BB26	3.59	77	23	242	3547	44.60	8
BB27	2.58	39	9	514	2602	43.10	9
BB28	3.02	103	27	187	3056	26.30	4
BB29	4.16	122	21	304	3375	34.40	21
BB30	1.99	35	9	292	2109	25.80	4
BB31	5.34	141	35	263	4342	45.30	15
BB32	3.62	326	117	52	3061	35.20	14
BB33	3.25	137	77	74	3319	56.80	9
BB34	4.87	115	31	302	4339	38.20	9
BB35	3.13	1.080	834	7	2773	31.80	12
BB36	4.29	149	54	138	3907	39.10	9
BB37	3.94	120	29	353	3831	51.50	11
BB38	3.62	242	62	144	3260	44.20	14
BB39	4.06	50	7	814	3832	59.10	17
BB40	6.73	296	113	106	5886	45.10	0
BB41	2.54	64	13	288	2631	32.90	7
BB42	3.59	118	30	186	3352	33.10	23
BB43	3.69	317	19	330	3767	49.10	10
BB44	4.03	21	5	985	4077	40.20	17
OB1	4.21	150	53	143	3949	43.10	13
OB2	4.15	563	111	53	3804	47.20	14
OB3	3.96	2.090	2.090	150	3665	48.70	16
OB4	2.73	66.9	15	778	2962	38.90	4
OB5	2.05	107	50	81	2113	31.90	5
OB6	3.88	180	68	100	3643	39.10	11
OB7	1.78	24	5	586	1916	26.10	2
OB8	2.92	106	17	269	3159	31.60	7

OB9	3.13	432	190	55	3181	55.10	6
OB10	3	55	14	316	2910	48.60	10
OB11	2.69	324	101	212	2676	30.50	5
OB12	4.68	59	21	361	4128	39.90	22
OB13	6.74	32	7	1266	5928	34.40	82
OB14	6.81	436	121	365	6165	44.90	17
OB15	2.25	20	4	800	2345	36.10	13

1042

1043

1044

1045   **Supplementary table S4:** *Colwellia* genomes used in this study for comparison to MAG BB5.

Name	NCBI BioProject accession number	Isolation source	Number of M9 collagenases	
Colwellia piezophila	PRJNA182419	Deep-sea sediment	2	
Colwellia psychrerythraea	PRJNA258170	*Terua* mussel	11	
Colwellia hornerae	PRJNA516280	Arctic sea ice	0	
Colwellia demingiae	PRJNA516284	Arctic sea ice	1	
Candidatus *Colwellia aromaticivorans*	PRJNA478776	Microcosm experiments with oil in seawater	0	
Colwellia echini	PRJNA420580	Sea urchin	0	
Colwellia beringensis	PRJNA378583	Marine sediment, Bering Sea	1	
Colwellia agarivorans	PRJNA371543	Coastal sea water	0	
Colwellia marinimaniae	PRJDB5767	Amphipod from Challenger Deep	4	
Colwellia sediminilitoris	PRJNA381102	Tidal flat, South Sea, South Korea	0	
Colwellia polaris	PRJNA380006	Arctic sea ice	0	
Colwellia mytili	PRJNA381102	Mussel *Mytilus edulis*	2	
Colwellia aestuarii	PRJNA371561	Tidal flat Korea	0	
Colwellia chukchiensis	PRJNA380006	Arctic ocean	1	

1046

1047

1048

Supplementary figures

Supplementary figure S1: A) ROV image of bone incubation experiment in the Byfjorden at 68 m depth, shown is a cow tibia. B) Bones retrieved from the Byfjorden after nine months of incubation, bacterial mats and blackening at the epiphysis can be seen.

Supplementary figure S2: Maximum-likelihood tree of all 94 obtained carbonic anhydrases and relevant reference sequences from Capasso *et al.*, 2015 (40).

Supplementary figure S3: Genomic context of all identified M9 collagenase in the investigated MAGs. M9 collagenases are encircled in red, all genes potentially involved in collagen/proline utilization pathways are encircled in green. The graphic was made with SnapGene software (from GSL Biotech; available at snapgene.com).

Supplementary figure S4: Comparison of the bone microbiome to seawater metagenomes. Tara Oceans seawater MAGs have been chosen for comparison. The Tara Oceans dataset comprises here 832 bacterial MAGs with >70% completion. The bone microbiome dataset has been rebinned to >70% completion and contains now 86 MAGs for this comparison. A) Percental bacterial class abundance comparison between bone microbiome MAGs (orange) and Tara Oceans MAGs (blue). Only bacterial classes present within the bone microbiome have been included in this graph, all classes only present in the Tara Oceans dataset are combined in ‚Others'. B) Enzyme abundance comparison. Shown is the ratio per MAG for each investigated enzyme class. All MAGs have been profiled with the here established HMM profiles and the total obtained number of enzymes was divided by the number of MAGs per dataset. Highlighted in red are the higher ratios and in bold red ratios at least twice as high as in the other datatset.

Figure 1: Phylogenomic maximum likelihood tree of the obtained 59 high-quality MAGs from *Osedax*-associated bone microbiome (OB) and bone surface-associated biofilm (BB). Bootstrap values greater than 0.5 are displayed as circles on the branches. The five most common bacterial classes are colored (green – Bacteroidia, light purple – Desulfuromonadia, purple – Campylobacteria, orange – Alphaproteobacteria and blue – Gammaproteobacteria). Order level identifications are listed and MAGs for which only class level identification could be inferred are marked with an asterisk.

Figure 2: Whole genome metabolic pathway comparison for genes of the sulfur metabolism. Analysis was done with MEBS (37) and MAGs were phylogenetically grouped according to GTDBTk pipeline (36). The grey scale represents the completeness of a given pathway or multi-enzyme system shown in the heatmap for each MAG. The OB MAGs are highlighted in blue. The "Sox system" constitutes of the *soxXYZABCD* genes and the "Sor system" of *sorABDE*.

Figure 3: Abundance and taxonomic affiliation of predicted gelatin hydrolysers (green) (Traitar) in *Osedax* and biofilm derived MAGs. Additionally, sulfur reducers (blue) and sulfur oxidizers (orange), (MEBS) are shown, while white indicates absence of these traits. MAGs are displayed at the deepest taxonomic classification obtained. The size of the circles reflects the number of MAGs within each clade.

Figure 4: Abundance heatmap of the 12 investigated enzyme COG classes in the 59 bone degradome MAGs. The MAGs are arranged according to their taxonomic affiliation. The absolute abundance of each enzyme COG class is depicted in the diagram on top of the heatmap. The OB MAGs are highlighted in blue.

Figure 5: Collagen utilization pathway scheme in MAG BB5. A) The gene cluster in BB5 spans approximately 21 kb, comprising 15 genes for collagen utilization, each color-coded respective to its functional group: orange for collagen hydrolysis, blue for uptake and transport, green for proline (Pro) utilization, ocher for hydroxyproline (Hyp) utilization and brown for unknown function. The purple box is indicative of a signal-peptide for secretion. B) Metabolic model for collagen utilization in *Colwellia* BB5. Arrows and genes are color-coded in the same functional groups as in A. Dashed arrows point to a major metabolic pathway. Metabolite abbreviations: P4C (1-pyrroline 4-hydroxy-2-carboxylate), KGSA (alpha-ketoglutarate semialdehyde), KG (alpha-ketoglutarate), P5C (1-pyrroline-5-carboxylate). Enzyme abbreviations: D-aa dHase (D-KGSA (alpha-ketoglutarate semialdehyde), dAminase (pyrroline-4-hydroxy-2-carboxylate deaminase), di-oxo dHase (KGSA dehydrogenase), hydroxyproline dehydrogenase), P5CR/ornithine dAminase (bifunctional 1-pyrroline-5-carboxylate reductase/ornithine cyclodeaminase), PRODH (proline dehydrogenase), P5CDH (pyrroline-5-carboxylate dehydrogenase).

Figure 6: Conservation between M9 collagen degradation gene clusters in *Colwellia psychrerythraea* GAB14E, *Colwellia piezophila* ATCC BAA-637 and the MAG *Colwellia* BB5 drawn at scale. dHase, dehydrogenase; PPIase, peptidyl-prolyl cis trans isomerase; Hyp, D-aa, dAminase etc. Color coding and gene names are indicated.

Material	Process	Enzymes	Organisms
Mineral matrix (hydroxyapatite)	Demineralization via acidification	Sulfur cycling processes (oxidation via Sox, Sor cycle, ...)	SOB (Beggiatoa, Thiovulaceae, Sulfurovaceae)
Collagen fibers	Enzymatic degradation	Collagenases, peptidases	Enterobacterales (Kangiellaceae, Colwellia)
Bone marrow	Utilization of glycosides, lipids, etc.	Glycosidases, cholesterol oxidases	Bacteroidia

Figure 7: Hypothesis of the interplay in the marine bone microbiome and degradome. Sulfur-oxidizing bacteria (SOB, shown with a halo) convert elemental sulfur and H_2S into sulfate and protons that lead to an acidification and therefore bone demineralization. Sulfate reducing (SRB, green) and sulfur disproportioning bacteria produce H_2S from sulfate. Enterobacterales and other especially Gammaproteobacteria secret collagenases to degrade collagen. Bacteroidia and other bacteria secret gylcosidases and other enzymes to hydrolyze the organic bone components (glycosides, esters, lipids). This exemplifies a bone demineralization loop that fuels itself as long as sulfur is available and degrades the organic bone components in the process.