Repeated global migrations on different plant hosts by the tropical pathogen

Phytophthora palmivora

Jianan Wang[1], Michael D. Coffey[2], Nicola De Maio[3], Erica M. Goss[1]*

[1] Department of Plant Pathology and Emerging Pathogens Institute, University of Florida,

Gainesville, FL, USA 32611;

[2] Department of Plant Pathology and Microbiology, University of California, Riverside, CA,

USA 92521;

[3] Nuffield Department of Medicine, University of Oxford, OX3 7BN, Oxford, United Kingdom.

*e-mail: emgoss@ufl.edu

Keywords: host tracking, host shift, Oomycete, genetic diversity, structured coalescent analysis

Running title: Host tracking by a broad host range plant pathogen

ORCID ID:

Jianan Wang: https://orcid.org/0000-0002-7704-666X
Nicola De Maio: https://orcid.org/0000-0002-1776-8564
Erica M. Goss: https://orcid.org/0000-0003-3512-2107

Abstract

The genetic structure and diversity of plant pathogen populations are the outcomes of

evolutionary interactions with hosts and local environments, and migration at different scales,

including human-enabled long-distance dispersal events. As a result, patterns of genetic variation

in present populations may elucidate the history of pathogens. *Phytophthora palmivora* is a

devastating oomycete that causes disease in a broad range of plant hosts in the tropics and

subtropics worldwide. The center of diversity of *P. palmivora* is in Southeast Asia, but it is a

destructive pathogen of hosts native to South America. Our objective was to use multilocus

sequence analysis to resolve the origin and historical migration pathways of *P. palmivora*. Our

analysis supports Southeast Asia as a center of diversity of *P. palmivora* and indicates that a

single colonization event was responsible for the global pandemic of black pod disease of cacao.

Analysis using the structured coalescent indicated that *P. palmivora* emerged on cacao and that

cacao has been the major source of migrants to populations in Asia, Africa, and Pacific Islands.

To explain these results, we hypothesize widespread introgression between the pandemic cacao

lineage and populations native to Asia and the Pacific Islands. The complex evolutionary history

of *P. palmivora* is a consequence of geographic isolation followed by long-distance movement

and host jumps that allowed global expansion with cacao, coconut and other hosts.

With continued growth of human populations and change in global climate, there is increasing

concern over plant diseases affecting food crop and economic security[1]. Epidemics of plant

disease can result in major yield losses and associated economic consequences[2]. In low- and

middle-income countries, plant diseases can have a relatively larger impact on local socio-

economic development[3,4]. Understanding the population biology and evolution of plant

pathogens can aid in disease management. For example, determining the geographic history of a

plant pathogen can help locate plant germplasm that co-evolved with the pathogen and exhibits

disease resistance, which can be employed in plant breeding programs. Knowledge of plant

pathogen migration patterns can lead to informed decision-making for disease prevention,

specifically to limit re-introductions and pathogen re-emergence[5]. Global pandemics may

originate from a very successful invasive population that itself is responsible for multiple

secondary invasions[3,6–8]. Hence, mitigation strategies may specifically target invasive

populations or genotypes[9,10].

However, the above efforts can be hampered by pathogens making dramatic host jumps.

Host jumps are pervasive in the emergence or spread of plant pathogens and have been identified

as a crucial mechanism underlying pathogen diversification and ultimately speciation[11,12].

International travel and global trade over the course of human history have exposed hosts to new

pathogens, thus, facilitating host jumps[1,3,9,13–15]. Deciphering complex interactions among plant

hosts, pathogens and human activities can elucidate major drivers behind the emergence, re-

emergence and dispersal of plant pathogens[1]. Many such studies of pathogen emergence have

targeted major crop pathogens that exhibit relatively narrow host associations[5,16–21]. When

pathogens have multiple host associations, can the influence of different hosts on pathogen

emergence and dispersal be resolved?

Phytophthora is a genus of plant-damaging Oomycetes, including more than 100

described species[22–25]. *Phytophthora* species can infect a broad range of plant hosts[26] and have

caused enormous economic losses to agro-ecosystems and ecological damage in natural

ecosystems. The relatively narrow host range pathogen *Phytophthora infestans,* causal agent of

potato and tomato late blight, was globally distributed on potato, leading to the Great Irish

Famine. In 2008, *P. infestans* spread across the Eastern United States on tomato plants for home

gardens[27]. *P. ramorum* is a broad host range pathogen, which dispersed on ornamental plants

followed by host jumps to timber and forest trees[28]. The resulting disease outbreaks in North

America and Europe have been economically costly and have changed the ecology of coastal

California forests[29]. *Phytophthora palmivora* (Butler) Butler (1919) is a destructive tropical and

subtropical plant pathogen with a very broad host range in the tropics and subtropics[30,31]. *P.*

palmivora causes problematic diseases of coconut and other palms (bud rot), cacao (black pod,

canker, and cherelle wilt), rubber (black stripe), durian (fruit rot and canker), orchids and other

ornamentals, and more[26]. Annual losses due to black pod of cacao, caused primarily by *P.*

palmivora, have been estimated at more than US$400 million per year[103] and bud rot endemics

have affected more than 70,000 ha of oil palm in Colombia[32]. However, like many other tropical

pathogens, research on *P. palmivora* population biology and genetics has lagged far behind

temperate *Phytophthora*, even though this pathogen is responsible for significant economic

losses. To advance research on *P. palmivora*, the genome of an isolate from cacao was recently

sequenced. Estimated genome size was greater than 151.23 Mb with 44 327 genes and initial

analysis of gene models indicated that *P. palmivora* experienced a genome doubling event[33].

87   The hypothesized center of origin of *P. palmivora* has changed over time. Initially,

Central or South America was suspected to be the native region of the pathogen, because of the

susceptibility of indigenous hosts and the apparent global distribution of the pathogen on cacao

(*Theobroma cacao*), which is native to South America[34]. Findings of high levels of genetic

diversity in *P. palmivora* populations in Southeast Asia, particularly among isolates from native

hosts coconut and durian, led to the proposal that Southeast Asia is the center of origin of *P.*

palmivora[30,35]. Recent advances in population genetics and coalescent model-based approaches

have revolutionized methods to identify plant pathogen centers of origin, reconstruct migration

pathways and reveal population genetic structure[36–44]. These new tools have substantially

advanced knowledge of the evolutionary history and epidemiology of major plant pathogens, but

have not been applied towards understanding the global spread of *P. palmivora*.

98   The main objective of this study was to describe the global population structure of *P.*

palmivora and historical migration pathways using Bayesian and coalescent model-based

inference approaches. We specifically addressed the following questions: (1) Are populations of

P. palmivora structured by host or geography? (2) Where is the center of origin of *P. palmivora*?

(3) What were the main migration pathways out of the center of origin? (4) Did host shifts drive

the global expansion of the *P. palmivora*?

Results

Nucleotide diversity by geographic region. We evaluated DNA sequence variation in four

genes (one mitochondrial and three nuclear) for three major geographic regions (Table 1). Higher

values of average pairwise nucleotide diversity (π) and Watterson's theta (θ_W) were observed in

the Asia-Pacific region than in Africa and the Americas. Tajima's D and Fu & Li's D* and F*

test statistics were not significantly different from zero except for the sample from the Americas

(Table 1). A significantly positive value of Tajima's D was observed for *trp1* in the Americas,

indicating more intermediate frequency alleles than expected under neutral evolution, while all

three test statistics were significantly negative for *coxII* in the Americas, indicating excessive

rare polymorphisms.

Population subdivision and structure. Population differentiation was tested by AMOVA

(Table 2). All four loci showed a significant portion of genetic variation (8-12%) distributed

among regions (Africa, Americas and Asia-Pacific). Differentiation between the two most

common hosts in the data set, coconut and cacao, explained around one quarter of the genetic

variation in the nuclear loci (23-27%).

To further explore population structure, *P. palmivora* isolates were grouped by Bayesian

clustering, using an allelic data set from the three nuclear loci (Fig. 1). When the number of

clusters (K) was set to 2, most isolates from the Asia-Pacific region were in one cluster, and the

majority of isolates from Africa and Americas were in the other. As K was increased, isolates

from Indonesia, Philippines, Malaysia, and Pacific Island formed new clusters. Based on delta K,

the optimum value of K was 2. The genetic structure within the Asia-Pacific region was greater

than in Africa and the Americas. The small sample from south Asia (India and Sri Lanka)

resembled populations in Africa and the Americas, the majority of which were isolated from

cacao. Bayesian clustering of clone-corrected data produced the same overall pattern as the full

data set (Supplementary Fig. 1).

The non-parameterized clustering method DAPC showed the first discriminant

component separating isolates collected from cacao in all geographic regions from isolates

collected from coconut and other hosts (Fig. 2). The second discriminant component separated

isolates sampled from durian and other hosts from coconut and cacao isolates. Six isolates

collected from coconut and other hosts in the Asia-Pacific and Americas clustered with isolates

from cacao, consistent with isolates shifting from cacao to other hosts.

Fig. 1. STRUCTURE assignments of *P. palmivora* isolates to populations for values of K from 2 to 6. Greater genetic structure is apparent in the Asia-Pacific region. Origins of isolates are indicated below each bar. Isolates from cacao are indicated by a blue symbol, those from coconut with a red symbol, and those from durian with an orange symbol. Isolates with purple or gray symbols are from other hosts. Region is indicated by the shape of the symbol (circle for Asia-Pacific, triangle for Africa, square for Americas).

Fig. 2. DAPC clusters of *P. palmivora* isolates from different hosts and geographic regions.
All isolates collected from cacao in the three geographic regions grouped together. Asia-Pacific
isolates sampled from coconut and durian grouped into two largely distinct clusters.

Global migration patterns. Discrete Bayesian phylogeographic analysis was used to infer the
geographic location of the root state of each locus[45,46]. The reconstructed root states supported
Indonesia and/or Philippines as the inferred evolutionary origin of our sample (Fig. 3). The
posterior probabilities for Philippines or Indonesia as the root state varied by locus
(Supplementary Table 4). The corresponding analysis with BASTA, which is not as biased by
sampling patterns, produced similar maximum clade credibility genealogies but with low
posterior probabilities for geographic root states.

Fig. 3. Root state posterior probability values inferred using discrete phylogeographic
analysis for each location (country) by locus: *β-tubulin*, *pelota*, *trp1* and *coxII*. The posterior
probabilities for an Indonesia root (red bar) and/or a Philippines root (magenta bar) are higher
than for the other locations (other colors) for each locus, inferred independently. Due to limited
space, only five locations with the five highest posterior probabilities for *β-tubulin* are shown.
For *pelota*, *trp1* and *cox II*, only the locations with posterior probabilities more than 0.01 are
shown.

The maximum clade credibility genealogies for the four loci (Fig. 4; Supplementary Figs.
2-4) indicate that lineages presently associated with Southeast Asia and the Pacific Islands
diverged much earlier than lineages associated with Central and South America and Central
Africa (mainly isolated from cacao). The mitochondrial locus, *coxII*, produced three lineages,
one of which shows early diversification and is represented by isolates from SE Asia, Australia,
Papua New Guinea, Vanuatu, French Polynesia, Hawaii, Ivory Coast and Jamaica (Fig. 4).

Across loci, isolates from Pacific Islands represented lineages that diverged prior to

diversification of cacao-associated lineages. We observed that isolates from different hosts

within given geographic regions had distinct evolutionary histories. For example, an isolate

collected from coconut in Ivory Coast had distinct ancestry from the cacao isolates collected in

Central Africa, and this was also observed for isolates from Florida and Jamaica. For the nuclear

loci, we observed that most of the isolates from cacao exhibited two distinct haplotypes while

showing little variation within each haplotype among isolates, consistent with a clonal lineage.

Isolates from Indonesia and Philippines were less likely to have two diverged alleles but they

exhibited more variation among isolates.

Fig. 4. Maximum clade credibility genealogy for the *coxII* locus. Colors of branches indicate

the most probable geographic origin of each lineage.

Host switching. To complement the phylogeographic analysis and infer patterns of historical

migration between major host groups, we conducted a multi-locus BASTA analysis by host

group and region (Americas, Asia-Pacific and Africa). Based on analyses of genetic structure

and phylogeography, we expected that migration occurred from coconut or other native hosts in

SE Asia to hosts in the Pacific, to cacao in the Americas, and then back to cacao and other hosts

in Asia. The BASTA analysis did not confirm this pattern. Using a two-host model to determine

the directionality of gene flow between cacao and coconut, we inferred cacao to be the root host

with high posterior probabilities (≥0.96 for each locus) and significant migration from cacao to

coconut (Supplementary Table 5a). We examined if the existence of unsampled populations

might have affected the analyses using a three-host model with one unsampled host. Cacao and

the unsampled host were inferred as the root hosts with nearly even posterior probabilities. Mean

rates of migration from cacao to coconut and the unsampled host to coconut were moderate to

high, but the 95% HPD of the migration rates from the unsampled host included zero

(Supplementary Table 5b), indicating uncertainty in these estimates. To incorporate geography,

we examined a three-deme model, with populations cacao-America, cacao-Asia-Pacific and

coconut-Asia. Cacao-America was inferred as the root state with high posterior probabilities

(≥0.93 for each locus). Cacao-America was the major source of migrants, while cacao-Asia-

Pacific and coconut-Asia were the major sinks (Table 3A; Supplementary Figs. 5-8).

Incorporating an unsampled population into this model produced uncertainty in the root state

with posterior probabilities of 0.63-0.68 for cacao-Asia-Pacific and 0.27-0.31 for the unsampled

population and wide 95% HPDs on immigration estimates from the unsampled population (Table

3b). To explore if all remaining hosts, other than coconuts and cacao, in the Asia-Pacific region

could represent the unsampled population, we examined a four-deme model replacing the

unsampled deme with a putative deme we designated Other-Asia-Pacific. Similar to the previous

models without an unsampled population, cacao-America was inferred as the root state with high

posterior probabilities (≥0.96 for each locus), and cacao-Asia-Pacific and coconut-Asia were the

two major sinks. Other-Asia-Pacific was inferred as a third major sink (Table 3c), which is the

opposite migration direction to the models incorporating an unsampled deme. Models that

explicitly incorporated the few cacao isolates from Africa produced results generally consistent

with the models without the cacao-Africa deme (Supplementary Table 6a, b).

Discussion

The evolutionary history of crop pathogens is as complex as the phylogeographic and

agricultural histories of their hosts and the human populations that have cultivated them. Our

multilocus sequence analysis of a global sample of *Phytophthora palmivora* isolates revealed

population subdivision among three major geographic regions (Africa, Americas, Asia-Pacific)

and between hosts, here highly sampled cacao and coconut. We found greater genetic diversity

and more complex population structure within Southeast Asia as compared to other global

regions, and reduced genetic variation among isolates collected from cacao. Therefore, our

results generally support Southeast Asia as the *P. palmivora* center of diversity[30,35]. On the other

hand, our analyses suggest that the Americas are a significant source of the genetic diversity now

observed in the Asia-Pacific region. Here, we argue that the complex evolutionary history of *P.*

palmivora reflects the history of human travel and trade in the tropics.

The greatest genetic diversity in *P. palmivora* appears to be within and among Pacific

island chains, which naturally introduce geographic isolation due to historically infrequent

movement of plant hosts, including by human migrations[47]. Genetically diverse isolates were

associated with coconut palms and a variety of other tropical hosts. However, because the

distribution of hosts sampled was not independent of geographic region, we cannot fully separate

the relative influence of host and geography. For example, American and African isolates largely

clustered together and separately from isolates from the Asia-Pacific region, but cacao was the

most-sampled host in Africa and the Americas. Indeed, the population genetic structure of *P.*

palmivora on cacao suggests that a single colonization was responsible for the global pandemic

of *P. palmivora* causing black pod of cacao. The globally successful "cacao strain" is

characterized by a single *coxII* haplotype and STRUCTURE cluster based on the genotypes of

three nuclear loci. The most frequent haplotypes for each locus are associated with cacao. Many

of the isolates from the Americas, Africa, and South Asia represented this cacao-associated

lineage. The resulting pattern of genetic variation in the Americas produced positive values of

Tajima's D test across nuclear housekeeping genes and a statistically significant negative value

for the mitochondrial *coxII* locus. These values are consistent with a recent population

bottleneck[48]. In the Americas and South Asia, the "cacao strain" was isolated from palms and

rubber, suggesting movement from cacao to other hosts. For example, two isolates obtained from

cacao and oil palm in Colombia had identical sequences at the four sequenced loci. These

isolates shared haplotypes with isolates from cacao worldwide and commercial oil palm

production only began in Colombia in 1945[102], therefore we infer that *P. palmivora* colonized oil

palm from cacao.

We conducted phylogeographic analyses to infer the potential historical processes behind

the observed patterns of genetic variation. Using discrete phylogeography, we inferred root states

of our *P. palmivora* sample to be Indonesia or the Philippines, depending on the locus. We might

expect that the pathogen has a metapopulation structure across these island nations, mediated by

sea level changes joining or separating islands, dispersal of infected fruits following sea currents,

or, more recently, trade in hosts infected by *P. palmivora*. The corresponding BASTA analysis

was not able to infer a root location, and our reduced parameter analyses that grouped isolates by

geographic region and host group inferred an American origin. BASTA is less susceptible to the

effects of sampling bias, but inference of an American origin was unexpected because of the

higher genetic diversity of the pathogen in the Asia-Pacific region. Simulations suggest that high

posterior probabilities provided by the discrete model likely underestimate posterior

uncertainty[44]. Our discrete model results reflect deep sampling in Southeast Asia, while the

BASTA results indicate that sampling across a greater diversity of hosts and locations is needed.

A long history of controversy regarding the geographic and evolutionary origin of the late blight

pathogen, *P. infestans*, has been exacerbated by limited sampling of wild relatives of potato in

the South American Andes[49,50] and the geographic origins of many other widespread

Phytophthora pathogens are unknown.

Migration estimates over the history of our sample of *P. palmivora*, inferred by BASTA,

suggests movement from cacao in the Americas to cacao in Asia and Africa, and from cacao to

other hosts in Asia. Because we observed genetic diversity on other hosts, it is unlikely that the

sole origin of the pathogen is cacao. Indeed, we could not exclude the presence of an alternative,

unsampled source population. In our inferred genealogies, we observed that haplotypes were not

segregating independently, particularly for isolates from cacao, which likely represent an

asexually reproducing lineage. The phylogeny for the *coxII* mitochondrial locus showed two

groups of haplotypes, a large group of closely related haplotypes and a second group of low

frequency, diversified haplotypes, which could represent different histories of ancestral

populations of *P. palmivora*. One explanation for the observed patterns is introgression, perhaps

repeated introgression, from the pandemic lineage on cacao to endemic populations of

Phytophthora in Southeast Asia and the Pacific Islands. We do not yet have evidence of

populations of *P. palmivora* or a closely related species on wild plants in the Americas which

could have sourced migration to Asia. However, *P. palmivora* infects important tropical cash

crops native to South America, including cacao[52], rubber[53], and neotropical orchids[54]. Genome

sequencing of an isolate of *P. palmivora* from cacao indicates that this pathogen underwent

genome doubling. We speculate that genome doubling was the consequence of a hybridization

event between strains of different geographic origin, because genome doubling is commonly

associated with interspecific hybridization[51].

284   We can draw parallels between global patterns in the genetic variation of *P. palmivora*

and historical global movement of the hosts considered here (Fig. 5). The Philippines represents

one of two independent domestications of coconut (*Cocos nucifera*)[47,55]. Historical and genetic

data have clarified the routes of coconut as it spread throughout the tropics and subtropics[55].

Various lines of evidence suggest that Pacific coconut was brought from the Philippines to

eastern Polynesia and to the Pacific coast of Latin America by pre-Columbian Austronesians.

Trade occurred in the opposite direction as well, because herbarium specimens and

anthropological studies suggest pre-historic introduction of sweet potato from the Pacific coast of

South America to Polynesia[56]. Later, the Portuguese set up plantations of the South Asian

domestication (Indo-Atlantic genotypes) of coconut in West Africa, Brazil, and the Caribbean.

Our findings of early divergence in our *P. palmivora* sample from the Pacific Islands is

consistent with early movement, possibly on Pacific coconut, whereas isolates from Ivory Coast

and Jamaica with deep nodes suggest movement on Indo-Atlantic coconut. *P. palmivora* infects

numerous other tropical fruits and ornamentals native to the Asia-Pacific region, on which the

A. Historical movement of major hosts

B. Inferred *P. palmivora* migration rates between regions and host groups

Fig. 5. Movement of major hosts of *P. palmivora* out of their centers of origin and inferred

migration of *P. palmivora* by host group. (**a**) Simplified historical movement of major *P.*

palmivora hosts. Indo-Atlantic and Pacific coconut were domesticated in South and Southeast

Asia, respectively. Cacao and rubber originated in South America. Many other tropical crops

host *P. palmivora*, including ornamentals. (**b**) Summary of BASTA inferences of *P. palmivora*

migration (solid lines, width proportional to mean migration rate) and potential migration from

an unsampled deme (dotted lines). BASTA inferred high migration rates from cacao in the

Americas to cacao in Asia-Pacific and Africa, and migration from cacao in Asia-Pacific to

coconut and other hosts (indicated by a durian fruit), as well as migration from cacao in the

Americas directly to other Asia-Pacific hosts.

pathogen could have dispersed or persisted over hundreds of years. If we assume that the

divergence in sequences of isolates from Hawaii and other Pacific Islands are associated with

human colonization of the Eastern Pacific islands around 1200 AD, we find that the colonization

of global cacao production is during the last 150-200 years, as expected. The major documented

movements of cacao to Africa and Asia was during the colonial period, and originated from

Venezuela, Trinidad and Brazil[52,57]. Rubber is another major South American export and host of

P. palmivora, with modern plantations located in South and Southeast Asia and West Africa[59,60].

The history of rubber in Asia started with the introduction of specimen trees from the Amazon

region of Brazil to Ceylon, Singapore, and Penang in 1876, and rubber was established in

plantations in Malaysia through 1898[58].

We hypothesize that the diversity and genetic divergence of *P. palmivora* isolates in the

Asia-Pacific region is an example of host-tracking. Our suggest that there may a long history of

association of *P. palmivora* with coconut. Alternatively, *P. palmivora* may have long-term

associations with other hosts native to Southeast Asia and coconut served primarily as an early

vector for long-distance dispersal. The host-tracking model of plant pathogen emergence has

been seen in other host-pathogen systems in Southeast Asia, such as the black Sigatoka pathogen

Mycosphaerella fijiensis on banana[61] and blast fungus *Magnaporthe oryzae* on rice[62]. In contrast,

the nearly monomorphic population *P. palmivora* found in Americas, South Asia, and Africa can

be explained by a pandemic lineage of *P. palmivora* on cacao, following a genetic bottleneck

likely associated with initial colonization of this host. The population of *P. palmivora* associated

with cacao resembles the pattern of emergence of globally-distributed clonal lineages of *P.*

infestans on potato[21,63–65]. Several other oomycetes have exhibited similar genetic bottlenecks

upon intercontintental spread, including *Phytophthora ramorum*[28]. These events are often

associated with a host jump, particularly colonization of plants for export.

The evolutionary and geographic history of *P. palmivora* could be clarified by

examination of additional isolates from uncultivated plants, which may represent older or more

isolated populations, or by using preserved DNA from herbarium samples. In general, the

distributions of crop pests and pathogens are highly dependent on the distribution of their hosts,

even though the specific mechanisms of global spread differ among species[66]. The idiosyncrasies

of plant movement during human history shaped the genetic variation and structure of modern

populations of cultivated plants and of their pathogens. Therefore, efforts to manage disease may

benefit from teasing apart historical dependencies that structure host and pathogen genetic

variation and that are likely to mediate the plant-pathogen interaction.

Methods

P. palmivora **isolates.** Genomic DNA was provided from 95 isolates of *P. palmivora* in the

World Oomycete Genetic Resource Collection at the University of California, Riverside, USA.

The isolates were originally collected from 23 hosts and 22 countries on five continents (Africa,

Asia, Australia, North America and South America) (Supplementary Table 1). For analysis,

isolates were assigned to one of three large geographical regions: 1) Asia-Pacific (Asia, Australia

and Pacific Islands); 2) Americas (including the Caribbean islands); and 3) Africa.

Multilocus sequence typing. Four housekeeping genes known to contain variation within

Phytophthora species were used in this study: mitochondrial gene *coxII* with the adjacent

spacer[67], and three nuclear genes, *β-tubulin*, *pelota* and *trp1*[68,69]. Primers for the nuclear genes

were modified for *P. palmivora* based on draft whole genome sequence kindly provided by S.

Schornack and S. Kamoun (Supplementary Table 2).

For PCR, a total reaction volume of 25.0 µl was prepared including 1.0 ul DNA template,

1X OneTaq master mix (NEB), 0.2uM primers and 3.0 uM $MgCl_2$. PCR amplifications were

performed using the following cycling protocol: initial denaturation at 94 °C for 3 minutes;

followed by 35 cycles of 94 °C for 45 seconds, [(67 °C for 45 seconds (*pelota*); 58 °C for 45

seconds (*ß-tubulin*); 60 °C for 45 seconds (*coxII*); 59°C for 60 seconds (*trp1*)) and 68 °C for 1

minute; then a final extension step was 68 °C for 10 minutes. PCR products were prepared for

sequencing using ExoSAP-IT (USB Corporation, Cleveland, OH, USA) and directly sequenced

in both directions at Interdisciplinary Center for Biotechnology Research (ICBR) at the

University of Florida, USA. Reads were assembled and edited using software Geneious 6.1.8

(https://www.geneious.com). For the three nuclear loci, haplotype phase was inferred using the

program PHASE[70], assuming diploidy. The settings for PHASE were as follows: MR0 (the

default model, which is the general model for recombination rate variation[71]); number of

iterations=10000 for *ß-tubulin* and *trp1*, 2000 for *pelota*; thinning interval=1; burn-in=100. For

each nuclear locus, ten isolates that exhibited more than one heterozygous site were randomly

selected and cloned using the pGEM-T Easy Vector System (Promega Corporation, Madison,

WI, USA). Sequences were aligned with Geneious 6.1.8[72]. Indels were removed for analysis.

Nucleotide diversity and neutrality tests. Number of individuals (Nind), number of sequences

(Nseq), number of haplotypes (Nhap), segregating sites (S), average pairwise nucleotide

diversity (π)[73,74], Watterson's theta (θ_w)[75], Tajima's D[76–79], number of mutations, number of

singleton mutations, Fu and Li's D* and F*[76,77,80–85] and minimum number of recombination

events (Rm)[86] were determined for each of the four genes and by region using DnaSP v5[87].

Population subdivision and structure. Population subdivision among the three major

geographic regions (Africa, Americas and Asia-Pacific) and between two major hosts (coconut,

native to Southeast Asia; and cacao, native to South America) were examined for each gene

locus by analysis of molecular variance (AMOVA) in Arlequin 3.5.2.1[88].

The population structure of *P. palmivora* was examined by model-based Bayesian

clustering carried out in STRUCTURE 2.3.4[36]. For this analysis, allelic data were used from the

three nuclear loci. The program STRUCTURE estimates parameters independently in the

posterior probability distribution of allele frequencies. Parameters are estimated under the null

model of panmixia within populations, which is characterized by Hardy-Weinberg equilibrium at

each locus and an absence of linkage disequilibrium. The algorithm is robust to some departures

from these assumptions[89–91]. Based on allele frequencies at each locus, STRUCTURE assigns Q-

values (inferred ancestry) to each individual. Isolate P3444 was excluded, due to missing data for

pelota and *trp1*. To reduce the effect of asexual reproduction on STRUCTURE inferences, two

different input files were prepared. One used the full data set of 94 isolates and a second used a

clone-corrected data set of 69 isolates. Clone correction is a method to remove bias caused by

dominant or overrepresented genotypes generated by epidemic asexual reproduction. Here, we

clone-corrected globally by including one representative isolate of each multilocus genotype. We

applied the following workflow to analyze both input files. STRUCTURE was run using the

admixture model, and cluster numbers (K) from K=1 to K=15 were evaluated using 500 000

iterations after a burn-in period of 500 000 iterations. To evaluate the stability of the results

across repeated runs, 20 independent runs were conducted. The runs for each value of K were

evaluated based on the second order rate of change of the likelihood function with respect to K[92]

using the online program *Structure Harvester* v.0.6.94[93]. Due to stochastic effects among

replicate STRUCTURE runs, assignment probabilities were compiled for multiple runs in the

program CLUMPP version 1.2, which simplifies the assessment of replicate data by calculating

medians[94]. The parameters used were M=3, W=0 and S=2, GREEDY_OPTION=2, and

REPEATS=10000. The graphical visualization of the output was produced using R base function

'*barplot*'[95].

Population subdivision, based on allelic data from the three nuclear genes, was also

examined using model-free discriminant analysis of principal components (DAPC) implemented

in the R package *adegenet* 1.4.2[41]. DAPC was run to confirm the results of Bayesian clustering

approach. DAPC does not make biological assumptions and is less computationally intensive

than STRUCTURE. This method maximizes the variation between groups while minimizing

variation within group. First, DAPC transforms the data using principal components analysis

(PCA) and then discriminant analysis (DA) assigns individuals to clusters. The data

transformation ensures that the variables inputted to DA are uncorrelated and that their number is

less than that of analyzed individuals, so as to overcome the drawbacks of direct application of

DA. The appropriate number of principal components retained in the analysis can be determined

by cross-validation[41]. Because only one isolate, P8039, was in the pre-defined coconut-Africa

group and could not be used for in both training and validation sets, we removed this isolate and

retained 93 isolates for the DAPC analyses.

Genealogies and Phylogeographic Analysis. To infer the phylogeographic history of *P.*

palmivora, genealogies were inferred by Bayesian evolutionary analysis by sampling trees

(BEAST) and location states by discrete trait analysis[37,46]. We conducted separate analyses for

each locus, each of which is expected to have a different genealogy from the others. BEAST

executes ancestral reconstruction of discrete states, here location of collection, in a Bayesian

statistical framework for evolutionary hypothesis testing using rooted, time-measured

phylogenies. In this analysis, locations associated with branch nodes were characterized by

continuous time Markov chain models, comparable to nucleotide, codon or amino acid

substitution models and Bayesian stochastic search variable selection (BSSVS) was used to

model the phylogeographic dispersion process[45]. This approach uses the geographic locations of

the samples to reconstruct the ancestral states of tree nodes and the root state. A strict molecular

clock model was used. The mutation rate was set as a constant 1.0, consequently the estimation

of branch lengths is in substitutions per site. We used jModelTest 2.1[96] to obtain the best fitting

models of nucleotide substitution for each gene alignment. A coalescent tree prior and a prior of

constant population size were assumed. Three replications of independent MCMC were run for

each locus and the first 10% of recorded states were discarded as burn-in (parameter settings in

Supplementary Table 3). The program was run until effective sample size estimates of greater

than 200 were obtained, with good mixing and convergence in independent runs, as assessed in

Tracer v1.6[97]. Maximum clade credibility (MCC) trees were summarized by TreeAnnotator

based on common ancestor height[98] and visualized using R package *rBt*[99].

We complemented the above analysis with Bayesian structured coalescent approximation

(BASTA)[44] in BEAST 2.0[100]. BASTA incorporates migration into the structured coalescent-

based model rather than modeling migration independently from the coalescent process as in the

above discrete phylogeographic analysis. BASTA phylogeographic inferences are less influenced

by sampling bias and variation in sampling intensity, and can include unsampled ghost demes.

All four loci were used in the analysis, were assumed to be unlinked, and were assigned different

nucleotide substitution models, as determined using jModeltest2 for each locus. The symmetric

BSSVS with 24 locations (shown in Fig. 4) was used to model the phylogeographic dispersion

process, which assumed a prior in which 60 migration rates are expected to be non-zero[45]. We

assumed the same population size for all demes. Two billion steps were used in the MCMC

chain for each run, and one tree was recorded every 200 000 steps. To reduce the lengthy

running time and to obtain an effective sample size estimates of greater than 200 with good

mixing and convergence, two parallel runs with different seed numbers were implemented, and

the two parallel tree files for each locus were merged. The merged tree files were assessed in

Tracer v1.6[97]. After the first 10% of recorded 'burn-in' states were discarded, MCC trees were

summarized by TreeAnnotator based on common ancestor height[98].

We also used BASTA to explore major between-host transmissions in relation to

geography by estimating migration rates between demes defined by host and region. We started

with a simple model of the two major hosts, cacao and coconut, to determine which of these two

is more likely to be the ancestral host of *P. palmivora*. We incorporated geography by splitting

cacao isolates into cacao-Asia-Pacific and cacao-America demes, resulting in three demes:

cacao-America, cacao-Asia-Pacific and coconut-Asia. We also examined the effect of adding an

unsampled (ghost) deme. Because we did not include isolates collected from hosts other than

coconuts and cacao in the Asia-Pacific region, we grouped these isolates into a deme to

determine if these hosts represented the unsampled (ghost) deme. Finally, we examined the

effect on the above models of adding a cacao-Africa deme, although the sample size from Africa

is small. The four loci were assumed to be unlinked and assigned different nucleotide

substitution models, as above. We ran migration models that allowed for asymmetric migration

between demes, assuming the same population size for all demes. Three hundred million steps

were used in the MCMC chain for each run, and one tree was recorded every 100 000 steps.

Mixing was assessed and trees examined as above. The MCC trees were visualized by

IcyTree[101]. Two replications of independent MCMC were run for each model.

Data availability

The DNA sequence data, inferred haplotypes and the analysis scripts that support the findings of

this study will be made available from the Dryad repository upon submission of this preprint to

another journal.

References:

1.   Zhan, J., Thrall, P. H., Papaïx, J., Xie, L. & Burdon, J. J. Playing on a pathogen's weakness:
Using evolution to guide sustainable plant disease control strategies. *Annu. Rev.*
Phytopathol. **53**, 19–43 (2015).

2.   Savary, S., Ficke, A., Aubertot, J.-N. & Hollier, C. Crop losses due to diseases and their
implications for global food production losses and food security. *Food Sec.* **4**, 519–537
(2012).

3.   Estoup, A. & Guillemaud, T. Reconstructing routes of invasion using genetic data: why,
how and so what? *Molecular Ecology* **19**, 4113–4130 (2010).

4.   Lawson Handley, L.-J. *et al.* Ecological genetics of invasive alien species. *BioControl* **56**,
409–428 (2011).

5.   Taole, M., Bihon, W., Wingfield, B. D., Wingfield, M. J. & Burgess, T. I. Multiple
introductions from multiple sources: invasion patterns for an important Eucalyptus leaf
pathogen. *Ecol Evol* **5**, 4210–4220 (2015).

6.   Hänfling, B., Carvalho, G. R. & Brandl, R. mt-DNA sequences and possible invasion
pathways of the Chinese mitten crab. *Marine Ecology Progress Series* **238**, 307–310
(2002).

7.   Downie, D. A. Locating the sources of an invasive pest, grape phylloxera, using a
mitochondrial DNA gene genealogy. *Mol. Ecol.* **11**, 2013–2026 (2002).

8.   Floerl, O., Inglis, G. J., Dey, K. & Smith, A. The importance of transport hubs in stepping-
stone invasions. *Journal of Applied Ecology* **46**, 37–45 (2009).

9.   Singh, R. P. *et al.* The emergence of Ug99 races of the stem rust fungus is a threat to world
wheat production. *Annual Review of Phytopathology* **49**, 465–481 (2011).

10. Cooke, D. E. L. *et al.* Genome analyses of an aggressive and invasive lineage of the Irish
potato famine pathogen. *PLOS Pathogens* **8**, e1002940 (2012).

11. Stukenbrock, E. H. Evolution, selection and isolation: a genomic view of speciation in

fungal plant pathogens. *New Phytologist* **199**, 895–907 (2013).

12. Restrepo, S., Tabima, J. F., Mideros, M. F., Grünwald, N. J. & Matute, D. R. Speciation in
fungal and oomycete plant pathogens. *Annu Rev Phytopathol* **52**, 289–316 (2014).

13. Sax, D. F. & Brown, J. H. The paradox of invasion. *Global Ecology and Biogeography* **9**,
363–371 (2000).

14. Tilman, D. *et al.* Forecasting agriculturally driven global environmental change. *Science*
292, 281–284 (2001).

15. Lockwood, J. L., Cassey, P. & Blackburn, T. The role of propagule pressure in explaining
species invasions. *Trends Ecol. Evol.* **20**, 223–228 (2005).

16. Couch, B. C. *et al.* Origins of host-specific populations of the blast pathogen *Magnaporthe*
oryzae in crop domestication with subsequent expansion of pandemic clones on rice and
weeds of rice. *Genetics* **170**, 613–630 (2005).

17. Stukenbrock, E. H., Banke, S., Javan-Nikkhah, M. & McDonald, B. A. Origin and
domestication of the fungal wheat pathogen *Mycosphaerella graminicola* via sympatric
speciation. *Mol. Biol. Evol.* **24**, 398–411 (2007).

18. Gladieux, P. *et al.* On the Origin and Spread of the Scab Disease of Apple: Out of Central
Asia. *PLOS ONE* **3**, e1455 (2008).

19. Munkacsi, A. B., Stoxen, S. & May, G. *Ustilago maydis* populations tracked maize through
domestication and cultivation in the Americas. *Proc. Biol. Sci.* **275**, 1037–1046 (2008).

20. Ali, S. *et al.* Origin, Migration routes and worldwide population genetic structure of the
wheat yellow rust pathogen *Puccinia striiformis* f.sp. *tritici. PLOS Pathogens* **10**, e1003903
(2014).

21. Goss, E. M. *et al.* The Irish potato famine pathogen *Phytophthora infestans* originated in
central Mexico rather than the Andes. *PNAS* **111**, 8791–8796 (2014).

22. Nelson, S. C. & Abad, Z. G. *Phytophthora morindae*, a new species causing black flag
disease on noni (*Morinda citrifolia* L) in Hawaii. *Mycologia* **102**, 122–134 (2010).

23. Érsek, T. & Ribeiro, O. Mini Review Article: An annotated list of new *Phytophthora*
species described post 1996. *Acta Phytopathologica et Entomologica Hungarica* **45**, 251–
266 (2010).

24. Hansen, E. M., Reeser, P. W. & Sutton, W. Phytophthora beyond agriculture. *Annual*
Review of Phytopathology **50**, 359–378 (2012).

25. Kroon, L. P. N. M., Brouwer, H., de Cock, A. W. A. M. & Govers, F. The genus
Phytophthora anno 2012. *Phytopathology* **102**, 348–364 (2012).

26. Erwin, D. C. & Ribeiro, O. K. Phytophthora diseases worldwide. American
Phytopathological Society (APS Press), St. Paul, MN, USA (1996).

27. Fry, W. E. *et al.* Five reasons to consider *Phytophthora infestans* a reemerging pathogen.
Phytopathology **105**, 966–981 (2015).

28. Grünwald, N. J., Garbelotto, M., Goss, E. M., Heungens, K. & Prospero, S. Emergence of
the sudden oak death pathogen *Phytophthora ramorum. Trends Microbiol.* **20**, 131–138

(2012).

29.   John T. Kliejunas. Sudden oak death and *Phytophthora ramorum*: a summary of the
literature. Gen. Tech. Rep. PSW-GTR-234. U.S. Department of Agriculture, Forest Service,
Pacific Southwest Research Station, Albany, CA (2010).

30.   Mchau, G. R. A. & Coffey, M. D. Isozyme diversity in *Phytophthora palmivora*: Evidence
for a southeast Asian centre of origin. *Mycological Research* **98**, 1035–1043 (1994).

31.   Lamour, K. *Phytophthora: A Global Perspective*. (CABI, 2013).

32.   Torres, G. A. *et al.* Bud Rot Caused by *Phytophthora palmivora*: A Destructive emerging
disease of oil palm. *Phytopathology* **106**, 320–329 (2016).

33.   Ali, S. S. *et al. Phytophthora megakarya* and *P. palmivora*, closely related causal agents of
cacao black pod rot, underwent increases in genome sizes and gene numbers by different
mechanisms. *Genome Biol Evol* (2017). doi:10.1093/gbe/evx021

34.   Zentmyer, G. A. Origin and distribution of four species of *Phytophthora*. *Transactions of
the British Mycological Society* **91**, 367–378 (1988).

35.   Oudemans, P. & Coffey, M. D. A revised systematics of twelve papillate *Phytophthora*
species based on isozyme analysis. *Mycological Research* **95**, 1025–1046 (1991).

36.   Pritchard, J. K., Stephens, M. & Donnelly, P. Inference of population structure using
multilocus genotype data. *Genetics* **155**, 945–959 (2000).

37.   Drummond, A. J. & Rambaut, A. BEAST: Bayesian evolutionary analysis by sampling
trees. *BMC Evol. Biol.* **7**, 214 (2007).

38.   Cornuet, J.-M. *et al.* Inferring population history with DIY ABC: a user-friendly approach
to approximate Bayesian computation. *Bioinformatics* **24**, 2713–2719 (2008).

39.   Beaumont, M. A. Approximate Bayesian computation in evolution and ecology. *Annual
Review of Ecology, Evolution, and Systematics* **41**, 379–406 (2010).

40.   Beerli, P. & Palczewski, M. Unified framework to evaluate panmixia and migration
direction among multiple sampling locations. *Genetics* **185**, 313–326 (2010).

41.   Jombart, T., Devillard, S. & Balloux, F. Discriminant analysis of principal components: A
new method for the analysis of genetically structured populations. *BMC Genetics* **11**, 94
(2010).

42.   Kamvar, Z. N., Tabima, J. F. & Grünwald, N. J. Poppr: An R package for genetic analysis
of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ* **2**, e281
(2014).

43.   Kamvar, Z. N., Brooks, J. C. & Grünwald, N. J. Novel R tools for analysis of genome-wide
population genetic data with emphasis on clonality. *Front Genet* **6**, (2015).

44.   De Maio, N., Wu, C.-H., O'Reilly, K. M. & Wilson, D. New Routes to Phylogeography: A
Bayesian structured coalescent approximation. *PLOS Genetics* **11**, e1005421 (2015).

45.   Lemey, P., Rambaut, A., Drummond, A. J. & Suchard, M. A. Bayesian phylogeography
finds its roots. *PLOS Computational Biology* **5**, e1000520 (2009).

46.   Drummond, A. J., Suchard, M. A., Xie, D. & Rambaut, A. Bayesian phylogenetics with

BEAUti and the BEAST 1.7. *Mol. Biol. Evol.* **29**, 1969–1973 (2012).

47. Gunn, B. F., Baudouin, L. & Olsen, K. M. Independent origins of cultivated coconut
(*Cocos nucifera* L.) in the Old World tropics. *PLOS ONE* **6**, e21143 (2011).

48. Fay, J. C. & Wu, C. I. A human population bottleneck can account for the discordance
between patterns of mitochondrial versus nuclear DNA variation. *Mol. Biol. Evol.* **16**,
1003–1005 (1999).

49. Adler, N. E. *et al.* Genetic Diversity of *Phytophthora infestans* sensu lato in Ecuador
provides new insight into the origin of this important plant pathogen. *Phytopathology* **94**,
154–162 (2004).

50. Goss, E. M. Genome-enabled analysis of plant-pathogen migration. *Annual Review of*
Phytopathology **53**, 121–135 (2015).

51. Buggs, R. J. A., Soltis, P. S. & Soltis, D. E. Does hybridization between divergent
progenitors drive whole-genome duplication? *Mol. Ecol.* **18**, 3334–3339 (2009).

52. Motamayor, J. C. *et al.* Cacao domestication I: the origin of the cacao cultivated by the
Mayas. *Heredity* **89**, 380–386 (2002).

53. Verheye, W. Growth and production of rubber. *Land Use, Land Cover and Soil Sciences*.
Encyclopedia of life support systems (EOLSS), UNESCO-EOLSS Publishers, Oxford, UK
(2010).

54. Hine, R. B. Pathogenicity of *Phytophthora palmivora* in the Orchidaceae. Plant Dis. Rep.
46, 643–645 (1962).

55. Baudouin, L. & Lebrun, P. Coconut (Cocos nucifera L.) DNA studies support the
hypothesis of an ancient Austronesian migration from Southeast Asia to America. *Genet*
Resour Crop Evol **56**, 257–262 (2009).

56. Roullier, C., Benoit, L., McKey, D. B. & Lebot, V. Historical collections reveal patterns of
diffusion of sweet potato in Oceania obscured by modern plant movements and
recombination. *PNAS* **110**, 2205–2210 (2013).

57. Wood, G. R. A history of early cocoa introductions. *Cocoa Growers' Bull*. **44,** 7-12 (1991).

58. Gin, O. K. *Southeast Asia : A Historical Encyclopedia, From Angkor Wat to East Timor*.
ABC-CLIO, Inc., Santa Barbara, CA, USA (2004).

59. Hallucinogenic Plants: Richard Evans Schultes, Elmer W. Smith: 9780307243621:
Amazon.com: Books. Available at: https://www.amazon.com/Hallucinogenic-Plants-
Richard-Evans-Schultes/dp/0307243621. (Accessed: 16th March 2019)

60. Frank Z. *The International Natural Rubber Market 1870-1930*. Economic History
Association (2010).

61. Robert, S., Ravigne, V., Zapater, M.-F., Abadie, C. & Carlier, J. Contrasting introduction
scenarios among continents in the worldwide invasion of the banana fungal pathogen
Mycosphaerella fijiensis. Mol. Ecol. **21**, 1098–1114 (2012).

62. Saleh, D., Milazzo, J., Adreit, H., Fournier, E. & Tharreau, D. South-East Asia is the center
of origin, diversity and dispersion of the rice blast fungus, *Magnaporthe oryzae*. *New*
Phytol. **201**, 1440–1456 (2014).

63. Grünwald, N. J. & Flier, W. G. The biology of *Phytophthora infestans* at its center of
origin. *Annu Rev Phytopathol* **43**, 171–190 (2005).

64. Gómez-Alpizar, L., Carbone, I. & Ristaino, J. B. An Andean origin of *Phytophthora*
infestans inferred from mitochondrial and nuclear gene genealogies. *PNAS* **104**, 3306–3311
(2007).

65. Wang, J. *et al.* High levels of diversity and population structure in the potato late blight
pathogen at the Mexico centre of origin. *Mol. Ecol.* **26**, 1091–1107 (2017).

66. Bebber, D. P., Holmes, T. & Gurr, S. J. The global spread of crop pests and pathogens.
Global Ecology and Biogeography **23**, 1398–1407 (2014).

67. Martin, F. N. Mitochondrial haplotype determination in the oomycete plant pathogen
Phytophthora ramorum. Curr Genet **54**, 23–34 (2008).

68. Blair, J. E., Coffey, M. D., Park, S.-Y., Geiser, D. M. & Kang, S. A multi-locus phylogeny
for *Phytophthora* utilizing markers derived from complete genome sequences. *Fungal*
Genet. Biol. **45**, 266–277 (2008).

69. Goss, E. M., Carbone, I. & Grünwald, N. J. Ancient isolation and independent evolution of
the three clonal lineages of the exotic sudden oak death pathogen *Phytophthora ramorum.*
Mol. Ecol. **18**, 1161–1174 (2009).

70. Stephens, M., Smith, N. J. & Donnelly, P. A new statistical method for haplotype
reconstruction from population data. *Am. J. Hum. Genet.* **68**, 978–989 (2001).

71. Li, N. & Stephens, M. Modeling linkage disequilibrium and identifying recombination
hotspots using single-nucleotide polymorphism data. *Genetics* **165**, 2213–2233 (2003).

72. Kearse, M. *et al.* Geneious Basic: An integrated and extendable desktop software platform
for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647–1649 (2012).

73. Nei, M. *Molecular Evolutionary Genetics*. Columbia University Press, New York, NY,
USA (1987).

74. Nei, M. & Miller, J. C. A simple method for estimating average number of nucleotide
substitutions within and between populations from restriction data. *Genetics* **125**, 873–879
(1990).

75. Watterson, G. A. On the number of segregating sites in genetical models without
recombination. *Theor Popul Biol* **7**, 256–276 (1975).

76. Kimura, M. *The Neutral Theory of Molecular Evolution*. Cambridge University Press,
Cambridge, UK (1983). doi:10.1017/CBO9780511623486

77. Tajima, F. Evolutionary relationship of DNA sequences in finite populations. *Genetics* **105**,
437–460 (1983).

78. Tajima, F. Statistical method for testing the neutral mutation hypothesis by DNA
polymorphism. *Genetics* **123**, 585–595 (1989).

79. Tajima, F. Measurement of DNA polymorphism. Takahata N, Clark AG (eds), *Mechanisms*
of Molecular Evolution., Sinauer Associates Inc, Sunderland, MA, USA, 37–59 (1993).

80. Ewens, W. J. The sampling theory of selectively neutral alleles. *Theoretical Population*

Biology **3**, 87–112 (1972).

81. Strobeck, C. Average number of nucleotide differences in a sample from a single
subpopulation: a test for population subdivision. *Genetics* **117**, 149–153 (1987).

82. Fu, Y. X. & Li, W. H. Statistical tests of neutrality of mutations. *Genetics* **133**, 693–709
(1993).

83. Simonsen, K. L., Churchill, G. A. & Aquadro, C. F. Properties of statistical tests of
neutrality for DNA polymorphism data. *Genetics* **141**, 413–429 (1995).

84. Fu, Y. X. Statistical properties of segregating sites. *Theor Popul Biol* **48**, 172–197 (1995).

85. Fu, Y. X. Statistical tests of neutrality of mutations against population growth, hitchhiking
and background selection. *Genetics* **147**, 915–925 (1997).

86. Hudson, R. R. & Kaplan, N. L. Statistical properties of the number of recombination events
in the history of a sample of DNA sequences. *Genetics* **111**, 147–164 (1985).

87. Librado, P. & Rozas, J. DnaSP v5: A software for comprehensive analysis of DNA
polymorphism data. *Bioinformatics* **25**, 1451–1452 (2009).

88. Excoffier, L. & Lischer, H. E. L. Arlequin suite ver 3.5: A new series of programs to
perform population genetics analyses under Linux and Windows. *Mol Ecol Resour* **10**, 564–
567 (2010).

89. Falush, D., Stephens, M. & Pritchard, J. K. Inference of population structure using
multilocus genotype data: Linked loci and correlated allele frequencies. *Genetics* **164**,
1567–1587 (2003).

90. Halkett, F., Simon, J.-C. & Balloux, F. Tackling the population genetics of clonal and
partially clonal organisms. *Trends Ecol. Evol. (Amst.)* **20**, 194–201 (2005).

91. Kaeuffer, R., Réale, D., Coltman, D. W. & Pontier, D. Detecting population structure using
STRUCTURE software: effect of background linkage disequilibrium. *Heredity (Edinb)* **99**,
374–380 (2007).

92. Evanno, G., Regnaut, S. & Goudet, J. Detecting the number of clusters of individuals using
the software STRUCTURE: a simulation study. *Mol. Ecol.* **14**, 2611–2620 (2005).

93. Earl, D. A. & vonHoldt, B. M. STRUCTURE HARVESTER: a website and program for
visualizing STRUCTURE output and implementing the Evanno method. *Conservation*
Genet Resour **4**, 359–361 (2012).

94. Jakobsson, M. & Rosenberg, N. A. CLUMPP: a cluster matching and permutation program
for dealing with label switching and multimodality in analysis of population structure.
Bioinformatics **23**, 1801–1806 (2007).

95. Team RC. R: A language and environment for statistical computing. (2013).

96. Darriba, D., Taboada, G. L., Doallo, R. & Posada, D. jModelTest 2: more models, new
heuristics and parallel computing. *Nat. Methods* **9**, 772 (2012).

97. Rambaut, A., Suchard, M., Xie, D. & Drummond, A. *Tracer v1.6*. Available from
http://beast.bio.ed.ac.uk/Tracer (2014).

98. Heled, J. & Bouckaert, R. R. Looking for trees in the forest: Summary tree from posterior

samples. *BMC Evolutionary Biology* **13**, 221 (2013).

99. Sánchez-Ramírez, S. *rBt: Handy functions for dealing with BEAST trees in R*. GitHub
repository: https://github.com/santiagosnchez/rBt (2018).

100. Bouckaert, R. *et al.* BEAST 2: A software platform for Bayesian evolutionary analysis.
PLOS Computational Biology **10**, e1003537 (2014).

101. Vaughan, T. G. IcyTree: rapid browser-based visualization for phylogenetic trees and
networks. *Bioinformatics* **33**, 2392–2394 (2017).

102. Marin-Burgos, V., Clancy, J. S. Understanding the expansion of energy crops beyond the
global biofuel boom: evidence from oil palm expansion in Colombia. *Energ. Sustain.*
Soc. **7,** 21 (2017). https://doi.org/10.1186/s13705-017-0123-2

103. Drenth, A. & Sendall, B. Economic impact of *Phytophthora* diseases in Southeast Asia.
Drenth, A. & Guest, D.I., eds. Diversity and management of *Phytophthora* in Southeast
Asia. Australian Centre for International Agricultural Research (ACIAR), Canberra,
Australia, ACIAR Monograph No. 114, 10–28 (2004).

Acknowledgements

714 We thank Jordan McLendon for assistance in the laboratory, Sebastian Schornack and Sophien

715 Kamoun for sharing *P. palmivora* genome data for primer design, and Santiago Sánchez-

716 Ramírez for assistance with *rBt*. This work was supported by the University of Florida

717 Department of Plant Pathology and Emerging Pathogens Institute.

Author contributions

720 JW and EMG designed the experiments. MDC contributed the DNA samples of *Phytophthora*

721 *palmivora* from the World Oomycete Genetic Resource Collection. JW performed the data

722 analyses with contributions from EMG and NDM. JW and EMG wrote the manuscript with

723 contribution from NDM and MDC. All authors reviewed and approved the manuscript.

Competing interests

726 The authors declare no competing financial interests.

TABLES

Table 1. Nucleotide variation by locus and geographic region. Statistics given are number of individuals (Nind), number of sequences (Nseq), number of haplotypes (Nhap), segregating sites (S), average pairwise nucleotide diversity (π), Watterson's theta (θ_W), Tajima's D, number of mutations, number of singleton mutations, Fu and Li's D* and F*, and minimum number of recombination events (Rm).

Locus	Geographic region	Nind	Nseq	Nhap	S	Π	θ_W	Tajima's D	Number of mutations	Number of singleton mutations	Fu & Li's D*	Fu & Li's F*	Rm
β-tubulin	Africa	6	12	4	23	0.0076	7.6	1.53	23	6	0.44	0.82	1
	America	18	36	7	30	0.0072	7.2	1.20	30	10	-0.61	<-0.01	6
	Asia-Pacific	66	132	45	55	0.0088	10.1	0.56	55	11	-0.19	0.15	14
pelota	Africa	6	12	6	14	0.0097	4.6	0.96	15	5	0.12	0.38	0
	America	18	36	8	20	0.0091	4.8	0.63	20	8	-1.01	-0.56	3
	Asia-Pacific	68	136	38	36	0.0108	6.6	-0.06	38	13	-1.82	-1.31	8
trp1	Africa	5	10	5	9	0.0053	3.2	1.42	10	2	0.71	1.00	0
	America	16	32	6	10	0.0048	2.5	**2.19**	10	2	0.31	1.05	2
	Asia-Pacific	69	138	24	35	0.0045	6.4	-1.24	37	10	-1.00	-1.32	6
coxII and spacer	Africa	5	5	2	10	0.0068	4.8	-1.19	10	10	-1.19	-1.26	0
	America	17	17	4	7	0.0018	2.1	**-2.00**	8	7	**-2.41**	**-2.65**	0
	Asia-Pacific	62	62	28	24	0.0083	5.1	-0.25	25	3	0.80	0.50	7

Bold: $P < 0.05$.

Table 2. Analysis of molecular variance among three major geographic regions and two major hosts.

Loci	Source of variation	Percentage of variation	F_{ST}*
β-tubulin	Among regions	7.77	0.08
pelota		10.01	0.10
trp1		11.54	0.12
coxII and spacer		8.16	0.08
β-tubulin	Among hosts	22.64	0.23
pelota		27.38	0.27
trp1		23.97	0.24
coxII and spacer		11.90	0.12

*All values of F_{ST} were statistically significant with $P < 0.05$.

Table 3. Estimates of migration rates between demes defined by hosts and geographical regions.

a. Three-deme model*

Source deme	Root state probability	Mean estimated immigration rate [95% HPD]:		
		Cacao-America	Cacao-AsiaPacific	Coconut-Asia
Cacao-America	0.93-0.94	-	**0.447** **[0.158-0.785]**	0.108 [0-0.271]
Cacao-AsiaPacific	<0.1	0.069 [0-0.188]	-	**0.306** **[0.076-0.575]**
Coconut-Asia	<0.1	0.031 [0-0.091]	0.043 [0-0.130]	-

* *P.palmivora* isolates from coconuts in America were not present in our sample
Bold: lower boundary of 95% HPD > 0.001

b. Four-deme model, including an unsampled deme

Source deme	Root state probability	Mean estimated immigration rate [95% HPD]:			
		Cacao-America	Cacao-AsiaPacific	Coconut-Asia	Unsampled
Cacao-America	0.63-0.68	-	**0.476** **[0.171-0.827]**	0.11 [0-0.273]	0.069 [0-0.212]
Cacao-AsiaPacific	<0.1	0.063 [0-0.180]	-	**0.314** **[0.053-0.604]**	0.082 [0-0.248]
Coconut-Asia	<0.1	0.031 [0-0.092]	0.048 [0-0.146]	-	0.085 [0-0.252]
Unsampled	0.27-0.31	0.076 [0-0.192]	0.113 [0-0.321]	0.272 [0-0.606]	-

Bold: lower boundary of 95% HPD > 0.001

c. Four-deme model, including a putative Others-AsiaPacific deme

Source deme	Root state probability	Cacao-America	Cacao-AsiaPacific	Coconut-Asia	Others-AsiaPacific
			Mean estimated immigration rate [95% HPD]:		
Cacao-America	0.95-0.97	-	**0.403** **[0.149-0.698]**	0.104 [0-0.228]	**0.222** **[0.034-0.433]**
Cacao-AsiaPacific	<0.1	0.037 [0-0.110]	-	**0.195** **[0.004-0.395]**	**0.316** **[0.004-0.599]**
Coconut-Asia	<0.1	0.018 [0-0.054]	0.047 [0-0.139]	-	**0.359** **[0.148-0.599]**
Others-AsiaPacific	<0.1	0.028 [0-0.081]	0.050 [0-0.153]	0.052 [0-0.152]	-

Bold: lower boundary of 95% HPD > 0.001