1
2
3
4
5 **FGF signaling in development beyond canonical pathways**

11 Ayan T. Ray, Pierre Mazot[1], J. Richard Brewer[2], Catarina Catela[3], Colin J. Dinsmore, and
12 Philippe Soriano*
13
14
15 Department of Cell, Developmental, and Regenerative Biology
16 Icahn School of Medicine at Mount Sinai
17 New York, NY 10029
23 *Corresponding author: philippe.soriano@mssm.edu
24
25
26 Present address: [1]INSERM Transfert, Paris, France
27   [2]Department of Immunobiology, Yale University, New Haven, CT 06520
28   [3]Department of Neurobiology, University of Chicago, Chicago, IL 60637

29
30
31

32 Keywords: FGF; ERK1/2; Craniofacial development; Neural crest; Cell adhesion

33
34

FGFs are key developmental regulators which engage a signal transduction cascade through receptor tyrosine kinases, typically involving ERK1/2, PI3K/AKT, and other effectors. However, it remains unknown if all FGF activities depend on kinase activity or these canonical signal transduction cascades. To address these questions, we generated allelic series of knock-in *Fgfr1* and *Fgfr2* mouse strains, carrying point mutations that disrupt binding of signaling effectors to the receptors, alone or in combination. We also produced a kinase dead allele of *Fgfr2* which broadly phenocopies the null mutant. When interrogated in cranial neural crest cells, point mutations in either receptor revealed discrete functions for signaling pathways in specific craniofacial contexts, but failed to recapitulate the single or double null mutant phenotypes even in their most extensive combination. Furthermore, we found that together these signaling mutations abrogated the established FGF-induced signal transduction pathways, yet certain FGF functions such as cell-matrix and cell-cell adhesion remained unaffected. Our studies establish combinatorial roles of both *Fgfr1* and *Fgfr2* in development and identify novel kinase-dependent cell adhesion properties of FGF receptors, independent of well-established roles in intracellular signaling.

Classic models of receptor tyrosine kinase activation involve ligand binding, receptor dimerization, transactivation of the kinase domain, phosphorylation of intracellular tyrosines, and binding of effectors that orchestrate activation of downstream signaling pathways [1]. Different thresholds in dimer strength and stability may also come into play to regulate signaling dynamics [2-4]. For FGFRs, where downstream pathways have been particularly well studied, numerous lines of evidence point to ERK1/2 as the main effector of FGF signaling [5,6]. Although characterization of effector binding to RTKs provides critical insights on signaling specificity, assessing relative pathway significance requires *in vivo* validation. A previous analysis showed that knock-in *Fgfr1* point mutations disrupting binding of multiple signaling effectors, alone or in combination, did not recapitulate the *Fgfr1*[-/-] phenotype despite eliminating ERK1/2 outputs [7]. These results, although suggestive of additional signaling pathways, were difficult to interpret due to expression of another receptor, FGFR2. Using cranial neural crest (cNCC)-derived mesenchyme to understand FGF signaling, we now investigate the relative contributions of both FGF receptors and their signaling outputs to craniofacial development. We find that signaling mutations for each receptor disrupt classical signal transduction pathways in the absence of the other receptor, but do not recapitulate the null phenotypes. We furthermore show that FGF signaling is kinase-dependent and identify non-canonical functions of FGF signaling that help reconcile the gap in our phenotypic analyses.

Fgfr1 and *Fgfr2* orchestrate craniofacial development by regulating cell survival

To interrogate FGFR1/2 functions, we initially investigated how loss of both receptors in cNCCs

influences craniofacial development. We found both receptors extensively co-expressed in the

cNCC-derived mesenchyme and overlying epithelia, using fluorescent *Fgfr1* and *Fgfr2* reporter

alleles (Extended Data Fig. 1A). Next, conditional null alleles (henceforth denoted *cKO*) of *Fgfr1*

and *Fgfr2* were combined with *Wnt1Cre* drivers active in NCCs [8,9]. At embryonic day (E) 18.5,

Fgfr1$^{cKO/cKO}$ embryos displayed a fully penetrant facial cleft, *Fgfr2*$^{cKO/cKO}$ mutants had no overt

phenotype, and *Fgfr1*$^{cKO/cKO}$*; Fgfr2*$^{cKO/cKO}$ double mutants exhibited severe agenesis of most NCC

derived craniofacial structures including the frontal and nasal bones, nasal cartilage, maxilla, and

mandible (Fig. 1A and Extended Data Fig. 1B). Both *Fgfr1*$^{cKO/cKO}$*; Fgfr2*$^{cKO/+}$ and *Fgfr1*$^{cKO/cKO}$*;*

Fgfr2$^{cKO/cKO}$ embryos exhibited defects in mandible development with hypomorphic proximal

structures including angular and coronoid processes (Extended Data Fig. 1C). Overall, double

mutants showed reduced or no ossification of cNCC-derived craniofacial skeletal structures

(Extended Data Fig. 1B-D and Supplementary Information Fig. S1A-D). At E9.5 and E10.5,

Fgfr1$^{cKO/cKO}$*; Fgfr2*$^{cKO/cKO}$*; ROSA26*$^{mT/mG}$ double mutants developed hypoplastic pharyngeal arches

with reduced NCC lineage GFP$^+$ cells (Fig. 1B and Extended Data Fig. 2A). We observed reduced

expression of facial prominence markers along with midline morphogenesis defects, but ectoderm

gene expression remained unaffected, suggesting reduced NCC numbers rather than patterning

defects (Extended Data Fig. 2B). FGFR1 and FGFR2 are thought to function predominantly in

mesenchymal or epithelial contexts, respectively, since *Fgfr1c* mutants recapitulate many aspects

of the *Fgfr1*$^{-/-}$ phenotype [10], and *Fgfr2b* mutants are reminiscent of *Fgfr2*$^{-/-}$ embryos [11]. Here, we

show that both receptors function coordinately within the NCC-derived mesenchyme, as

combined loss of *Fgfr1* and *Fgfr2* leads to significantly more severe midface and mandibular

defects than loss of either receptor alone.

A reduction in NCC numbers in the midface might occur due to reduced proliferation,

increased cell death, or both. Cell proliferation remained unaffected (Extended Data Fig. 3),

however, we observed enhanced cell death in *Fgfr1*$^{cKO/cKO}$*; Fgfr2*$^{cKO/cKO}$; *ROSA26*$^{mT/mG}$ E10.5

mutants in the craniofacial mesenchyme, notably in the Lateral Nasal Process (LNP) relative to

the Medial Nasal Process (MNP; Fig. 1C, D). We next asked to what extent this observation could

explain the overall morphological defects. The BH3-only protein BIM plays a critical role in

initiating apoptotic pathway in multiple cell types by binding and repressing the function of several

pro-survival BCL-2 family members [12-14], and is involved in craniofacial development [15]. Consistent

with a role for cell survival in the etiology of craniofacial defects, we found that *Fgfr1*$^{cKO/cKO}$*;*

Fgfr2$^{cKO/+}$; Bim$^{+/-}$ E17.5 mutants exhibited a 55% reduction in midline separation along with a
partial rescue of medial skeletal structures, including the anterior nasal cartilage, palatine process,
premaxilla, primary and secondary palate, pterygoid process and basisphenoid bone, compared
to control embryos (Fig. 1E). The reduction in *Bim* levels significantly rescued defects particularly
at the level of the nasal mesenchyme and the midface (Fig. 1F and Supplementary Information
Fig. 1E). BIM is a known target of phosphorylation by several MAP kinases, particularly ERK1/2
which targets it for ubiquitination and proteasomal degradation [16]. JNK and PI3K/AKT activation
are also known to affect BIM levels [17]. Cell survival through BIM may therefore be regulated by
FGF since several of these signaling pathways are engaged by FGFR1 and FGFR2 [5]. Taken
together, these results underscore the importance of cell death in the conditional double null
mutant phenotype.
114
Combined *Fgfr1/2* signaling mutations do not recapitulate the null phenotypes
To interrogate signaling mechanisms *in vivo*, we generated an allelic series of knock-in
point mutations at the *Fgfr2* locus preventing binding of effectors to the receptor (Fig. 2A-B;
Extended Data Fig. 4), similar to previous *Fgfr1* mutations [7]. The *Fgfr2^{F}*, *Fgfr2^{C}*, and *Fgfr2^{PG}*
mutations were designed to disrupt binding of FRS2, CRK-L and PLCγ/GRB14, respectively. We
also generated compound *Fgfr2^{CPG}* and *Fgfr2^{FCPG}* signaling mutants by combining multiple
signaling mutations. Co-immunoprecipitation and western blot analysis showed that each
signaling mutation disrupted effector binding (Fig. 2B). *Fgfr2* signaling mutant alleles were
evaluated for their ability to partially or completely recapitulate the *Fgfr2$^{-/-}$* E10.5 placenta and limb
phenotype [18-20]. Surprisingly, all signaling allele mutants were at least partially viable and fertile
as homozygotes (Supplementary Information Table S1 and Supplementary Information Fig. S2).
Fgfr2$^{F/F}$ and *Fgfr2$^{FCPG/FCPG}$* homozygous mutants also exhibited defects in lacrimal gland
branching morphogenesis, a process regulated by FGF signaling [21,22] (Extended Data Fig. 5).
IRS2 has been found to bind FGFR2 [23], but no genetic interaction was observed between *Irs2$^{-/-}$*
and *Fgfr2$^{FCPG/FCPG}$* mutants.
Because *Fgfr1* and *Fgfr2* are co-expressed in cNCCs, we reasoned that discrete functions
of signaling pathways downstream of one receptor could be masked by the presence of the other,
wild-type receptor. To test this hypothesis, we analyzed compound conditional hemizygous *Fgfr1^{F}*,
Fgfr1^{FCPG} and *Fgfr2^{F}* mutations over the *Fgfr1^{cKO}* and *Fgfr2^{cKO}* conditional null alleles. *Fgfr1$^{F/cKO}$*;
Fgfr2$^{cKO/cKO}$ and *Fgfr1$^{FCPG/cKO}$; Fgfr2$^{cKO/cKO}$* conditional mutants developed severe agenesis of
midface structures, nasal cartilage, and mandible but the phenotype was less severe than in
Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$ mutants. The nasal cartilage and mandible defects were further

exacerbated in $Fgfr1^{FCPG/cKO}$; $Fgfr2^{cKO/cKO}$ mutants (Fig. 2C, Extended Data Fig. 6A, red arrows,

and Supplementary Information Table S2). $Fgfr1^{cKO/cKO}$; $Fgfr2^{F/cKO}$ embryos developed more

severe midline fusion and mandible defects than $Fgfr1^{cKO/cKO}$; $Fgfr2^{cKO/+}$ controls. Similar to

$Fgfr1^{cKO/cKO}$; $Fgfr2^{cKO/cKO}$ mutants, morphological defects in $Fgfr1^{F/cKO}$; $Fgfr2^{cKO/cKO}$ and

$Fgfr1^{FCPG/cKO}$; $Fgfr2^{cKO/cKO}$ embryos arose as early as E10.5 and were accompanied by cell death

in the LNP (Extended Data Fig. 6B-E).

We next examined the importance of coordinate engagement of signaling pathways by

both receptors upon ligand activation. $Fgfr1^{C/C}$; $Fgfr2^{C/C}$ and $Fgfr1^{CPG/CPG}$; $Fgfr2^{CPG/CPG}$ double

mutants were recovered in normal numbers, fertile, and did not exhibit a craniofacial phenotype.

Compound $Fgfr1^{F/F}$; $Fgfr2^{+/F}$ mutants displayed hypoplastic nasal prominences and midline fusion

defects at E15.5 (Fig. 2D) with no defect in mandibular development (Extended Data Fig. 6F), but

$Fgfr1^{F/F}$; $Fgfr2^{F/F}$ mutant embryos were not recovered at E10.5. Similar hypoplastic nasal and

mandibular prominence defects were observed in $Fgfr1^{FCPG/FCPG}$; $Fgfr2^{+/FCPG}$ compound mutants

(Extended Data Fig. 6G). Interestingly, $Fgfr1^{FCPG/FCPG}$; $Fgfr2^{FCPG/FCPG}$ mutants survived to E7.5 and

still formed mesoderm, as evidenced by $Eomes$ staining (Fig. 2E), in contrast to $Fgfr1^{-/-}$; $Fgfr2^{-/-}$

double mutants which fail at implantation on the same genetic background [24]. These results

indicate that signaling mutations in both $Fgfr1$ and $Fgfr2$ interact genetically during development,

but that the combination of the most severe signaling mutations fails to recapitulate the double

null mutant phenotype.

Fgfr1/2 signaling mutations abrogate signal transduction cascades

FGFs activate numerous signaling pathways upon ligand stimulation [5]. To evaluate

intracellular pathway activation downstream of wild-type FGFR2, FGFR2^F, FGFR2^{CPG}, and

FGFR2^{FCPG}, we generated immortalized E10.5 frontonasal prominence cell lines (iFNPs) by

crossing to *Ink4a/Arf* mutants [25]. These cells express predominantly *Fgfr1* and to a lesser extent

Fgfr2, but no *Fgfr3* or *Fgfr4*. We then eliminated FGFR1 expression by CRISPR/Cas9

mutagenesis leaving FGFR2 as the sole receptor, and interrogated activation of six pivotal FGF-

engaged pathways [5] by FGF1, which gave more robust responses than FGF8b (Extended Data

Fig. 7) . For ERK1/2, but also for PI3K/AKT and PLCγ, robust activation seen in *wt*-iFNPs was

diminished in both $Fgfr2^{F/F}$ or $Fgfr2^{CPG/CPG}$-iFNPs, but only eliminated in $Fgfr2^{FCPG/FCPG}$ mutant cells

(Fig. 2F and Extended Data Fig. 8). Our data thus indicates that both FRS2 and CRKL/PLCγ

binding is necessary for activation of these pathways, as for *Fgfr1* [7]. P38, STAT3 and JNK

activation were also abrogated in $Fgfr2^{FCPG/FCPG}$-iFNPs, with variable effects in $Fgfr2^{F/F}$-iFNPs and

$Fgfr2^{CPG/CPG}$-iFNPs (Extended Data Fig. 8). Last, we revisited cell signaling pathways in

Fgfr1^{FCPG/FCPG} mutants [7]. We used CRISPR/Cas9 to create *Fgfr1*^{FCPG/FCPG}; *Fgfr2*^{CRISPR-KO}-iFNP

cells, and found that both ERK1/2 and pAKT activation were eliminated in these cells (Extended

Data Fig. 8F-G). Together, these results indicate that the most severe signaling mutation

combinations in *Fgfr1* and *Fgfr2* effectively abrogate classic signal transduction pathways for each

receptor.

Fgfr2 function requires its kinase activity

The lack of more severe phenotypes in *Fgfr2* signaling mutants raised the possibility that

FGFR2 acts independent of kinase activity. To address this question, we generated a K517A

kinase dead (KD) mutation in the ATP binding site at the *Fgfr2* locus [26,27] (Fig. 2A and Extended

Data Fig. 4A, C). *Fgfr2*^{KD/+} heterozygous embryos (27/60) showed no obvious defects at E10.5.

Fgfr2^{KD/+} heterozygotes appeared normal during later developmental stages up to P0, but fewer

than expected (24/78) were recovered after weaning. They exhibited no limb or craniofacial

abnormalities, but significantly more severe semi-dominant defects in lacrimal gland development

than *Fgfr2*^{F/F} and *Fgfr2*^{FCPG/FCPG} mutants by P15 (Extended Data Fig. 5A-C).

No *Fgfr2*^{KD/KD} homozygotes were recovered at birth (Supplementary Information Table S1).

Morphological examination of *Fgfr2*^{KD/KD} embryos at E10.5 showed characteristic *Fgfr2*^{-/-}

phenotypes, with absence of limb buds, incomplete connection of the allantois to the ectoplacental

cone, and dilated pericardium (Fig. 3A), suggesting that FGFR2 broadly operates in a kinase-

dependent fashion. Moreover, since the *Fgfr2b* constitutive mutation leads to similar limb

phenotypes but no placental insufficiency [11], and phenotypes in both tissues are observed in

Fgfr2^{-/-} or *Fgfr2*^{KD/KD} mutants, FGFR2 activity must be kinase-dependent in both mesenchymal

and epithelial contexts. However, *Fgfr2*^{KD/KD} embryos exhibited additional phenotypes including

severe posterior truncations and craniofacial defects (Fig. 3A). We tested for complementation

between *Fgfr2*^{KD} and *Fgfr2*⁻ alleles to evaluate phenotypic differences. *Fgfr2*^{KD/-} embryos showed

absence of limb buds and defects in the chorio-allantoic junction, along with dilated pericardium,

similar to *Fgfr2*^{-/-} embryos, and posterior truncations defects similar to *Fgfr2*^{KD/KD} embryos (Fig.

3A). In contrast, defects in the forebrain, medial and lateral nasal prominences, and maxillary and

mandibular prominences appeared less severe than in *Fgfr2*^{KD/KD} mutants (Fig. 3B-D).

FGFRs can form heterodimers in culture [27,28], although these have never been

demonstrated *in vivo* in the absence of over-expression. The semi-dominant effects observed in

Fgfr2^{KD/+} heterozygous mutants affecting postnatal lacrimal gland development, and the

craniofacial and mesoderm defects observed in *Fgfr2*^{KD/KD} embryos suggest that the *Fgfr2*^{KD} allele

not only inactivates *Fgfr2* but also suppresses *Fgfr1* activity through FGFR2^{KD}:FGFR1

heterodimers, wherever they are co-expressed. This would explain why *Fgfr2^KD^* mutants still
generally resemble *Fgfr2^-/-^* mutants which are lethal at a similar stage, rather than *Fgfr1^-/-^* mutants
[7,18,24]. While signaling might occur through heterodimers, this mechanism cannot fully account for
the discrepancy between FCPG and null phenotypes for either receptor, as *Fgfr1^FCPG/FCPG^;*
*Fgfr2^FCPG/FCPG^* double mutants develop until E7.5 with a significant degree of mesoderm formation
(Fig. 2E) whereas *Fgfr1; Fgfr2* double null mutants have a more severe defect at implantation [18,24].
Alternatively, the FGFR2^KD^ receptor could act as a dominant negative by titrating/sequestering
ligand away from functional receptors, if it is distributed differently in the cell, but this is unlikely
as *Fgfr2^FCPG/+^* mutants do not show dominant negative effects and both knock-in alleles should
be expressed at similar levels.

Cell matrix and cell adhesion properties are retained beyond canonical signaling

Since signaling mutant cells showed near complete inactivation of classic RTK signaling
activities, but the corresponding mutant mice failed to recapitulate the null mutant phenotype, we
reasoned that some function engaged by FGF signaling must be retained in the most severe
FCPG mutants. Previous lines of evidence have implicated FGF signaling in the control of cell-
matrix [29] or cell-cell adhesion [24,30,31]. We examined *Fgfr1; Fgfr2* dependent cell
spreading/migration on extracellular matrix across a wound, using primary *Fgfr1^cKO/+^; Fgfr2^cKO/+^,*
Fgfr1^FCPG/+^; Fgfr2^cKO/+^, or *Fgfr1^cKO/cKO^; Fgfr2^cKO/cKO^* FNP cells. Spreading of control cells over the
wound over twelve hours was comparable upon FGF, PDGF, and serum stimulated conditions.
Interestingly, *Fgfr1^FCPG/cKO^; Fgfr2^cKO/cKO^* cells showed comparable spreading capacities in all three
stimulation conditions. However, *Fgfr1^cKO/cKO^; Fgfr2^cKO/cKO^* double mutant cells failed to spread into
the wound area in response to FGF, while responses to PDGF and serum were normal (Fig. 4A).

We next investigated focal adhesion formation and found that GFP^+^ *Fgfr1^cKO/+^; Fgfr2^cKO/+^;*
*ROSA26^mT/mG^* control cells and *Fgfr1^FCPG/cKO^; Fgfr2^cKO/cKO^; ROSA26^mT/mG^* mutant cells formed
numerous Paxillin enriched focal adhesions in response to FGF, PDGF or serum. Double null
mutant cells however failed to form any Paxillin^+^ foci upon FGF treatment, but still responded
normally to PDGF and serum as expected (Fig. 4B and Extended Data Fig. 9A). Western blot
analysis showed reduced pFAK and total Paxillin levels in *Fgfr1^cKO/cKO^; Fgfr2^cKO/cKO^* cells compared
to *Fgfr1^cKO/+^; Fgfr2^cKO/+^* or *Fgfr1^FCPG/cKO^; Fgfr2^cKO/cKO^* cells, and FAK phosphorylation depended on
FGF (Extended Data Fig. 9B-D). Likewise, we analyzed *Fgfr1^CRISPR-KO^; Fgfr2^+/+^* and *Fgfr1^CRISPR-KO^;*
*Fgfr2^FCPG/FCPG^* -iFNP cells and found that they spread and formed Paxillin^+^ focal adhesions in
response to serum, PDGF, or FGF, in contrast to *Fgfr1^CRISPR-KO^; Fgfr2^-/-^* -iFNP cells (Fig. 4B and
Extended Data Fig. 9E). These observations indicate that although FGFR1^FCPG^ and FGFR2^FCPG^

lose most FGF-dependent intracellular kinase signaling outputs, they still retain functions
pertaining to cell-matrix interactions and that cell spreading and stabilization of cell-matrix
interactions are actively governed by FGF signaling to its receptors.

We also explored if *Fgfr1^{FCPG}* cells established stable cell-cell contacts comparable to
control cells. We found extensive adherens junctions among *Fgfr1$^{cKO/+}$; Fgfr2$^{cKO/+}$; ROSA26$^{mT/mG}$*
control cells and *Fgfr1$^{FCPG/cKO}$; Fgfr2$^{cKO/cKO}$; ROSA26$^{mT/mG}$* cells, marked by localized β-catenin
along cell boundaries (Fig. 5A-B and Extended Data Fig. 9F). In contrast, double null mutant cells
showed far fewer cell-cell contacts with no localized β-catenin accumulation, suggesting that
contacts are either unstable or do not mature (Fig. 5B). Last, we examined cell-cell contacts *in*
vivo, in the E11.5 LNP. GFP$^+$ cNCCs in the mesenchyme exhibited extensive cell-cell contacts
within the LNP in both control and *Fgfr1$^{FCPG/cKO}$; Fgfr2$^{cKO/cKO}$; ROSA26$^{mT/mG}$* embryos. Strikingly,
GFP$^+$ cNCCs in the double null mutant LNP were mostly isolated and interspersed, whereas cell
contacts in the MNP remained unaffected (Fig. 5C). Taken together, these results indicate that
the most severe signaling mutations in *Fgfr1* and *Fgfr2* still retain cell-matrix and cell-cell
interactions otherwise lost in the nulls, while abrogating classic signal transduction pathways.

Discussion

FGFs are best known for activating signal transduction cascades, most prominently
ERK1/2 [5,6], Surprisingly, we find that mutations in *Fgfr1* and *Fgfr2* which broadly eliminate classic
RTK signaling outputs fail to recapitulate the *Fgfr1$^{-/-}$* or *Fgfr2$^{-/-}$* phenotypes. The near complete
abrogation of multiple signal transduction outputs in the most severe *Fgfr1* and *Fgfr2* signaling
alleles indicates that we have interrogated relevant cell signaling pathways and implies the
existence of non-canonical functions not impacted by our signaling mutations. Importantly, while
the *FCPG* mutations disrupt the ability of intracellular effectors to engage classical RTK activity,
they do not abrogate the receptor's kinase activity, suggesting that FGFR1^{FCPG} and FGFR2^{FCPG}
receptors may be able to phosphorylate other targets.

A wide body of literature has implicated FGFRs in cell adhesion through interactions of
the extracellular domain, which remains untouched in any of our signaling mutations, with cell
adhesion molecules. FGF ligand binding to FGFRs is known to involve a third party, heparan
sulfate proteoglycans (HSPGs)[32,33, 34], which in turn interact with integrins, regulating cell-matrix
adhesion [35-37]. Consistent with a critical role for FGFRs, we found that cell-matrix adhesion was
unperturbed in cells carrying the most severe signaling mutations, but lost in the absence of
FGFRs following FGF stimulation. Because FGF ligand still promotes cell-matrix adhesion in cells
derived from *Fgfr1/2^{FCPG}* mutants, the activity of cell adhesion receptors must be enhanced

independent of traditional signaling outputs, either directly through their interaction with FGFRs
or indirectly by an unknown molecule/kinase. Defects in FGF-dependent adhesion could also
result in the induction of anoikis [38], a process previously shown to be regulated by *Bim* [39],
potentially linking our observed adhesion defects and increase in cell death in the LNP. Last, focal
adhesion assembly and phosphorylation defects have also been observed in *Fgfr1*$^{-/-}$*; Fgfr2*$^{-/-}$
keratinocytes [29], suggesting that engagement of FGF signaling has a broad function in regulating
cell adhesion in both mesenchymal and epithelial contexts. Other receptor tyrosine kinases, like
PDGFRs, have been found to bind integrins [40] and PDGFR signaling mutations have also failed
to recapitulate null mutant phenotypes [41,42].

In addition, both FGFR1 and FGFR2 are known to interact through their extracellular
domain acid box with various cell adhesion molecules such as cadherins [43], critical players in
cNCC migration [44]. Consistent with a role for FGFRs in mediating cell-cell adhesion through
cadherins, we observed that cells derived from the most severe cell signaling mutants made
strong β-catenin cell contacts, in contrast to double null mutant cells. We also observed fewer
contacts *in vivo*, where double null mutant cells showed very limited interactions in the LNP.
FGFR1/2 association with cadherins may thus have a dual role, one in promoting cell motility
through FGF-induced ERK1/2 signaling [43], and an opposite one in cell adhesion, as ERK1/2
signaling is abrogated in *Fgfr1*^{FCPG} and *Fgfr2*^{FCPG} mutants. FGF signaling regulates E-cadherin
localization, and in the absence of *Fgfr1*, E-cadherin polarization is affected in mural
trophectoderm [24], *Drosophila* mesoderm [31], and zebrafish cardiomyocytes [30]. FGF signaling also
regulates cadherin switching during epithelial to mesenchymal transition, including neural crest
cell delamination [45-47]. Taken together, our results indicate that FGFRs regulate cell adhesion,
and possibly other processes, beyond their classic signaling cascades (Fig. 5D). Additional
genetic, biochemical and cell biological studies may identify further non-canonical roles for these
receptors beyond their traditional activities in signal transduction.

Acknowledgments. We thank Jia Li and Chantel Dixon for technical assistance; Kevin Kelley for

stable tissue culture facilities; Elaine Fuchs for helpful insights into cell adhesion mechanisms;

Jerry Chipuk for conversations about cell death; and our laboratory colleagues, Stu Aaronson,

Rob Krauss, and Sergei Sokol for critical comments on the manuscript. We thank the NYU School

of Dentistry Micro-CT Core and the Mt. Sinai Flow Cytometry and Microscopy facilities for

assistance and advice. C.J.D was supported by F32 DE026678 from NIH/NIDCR. This work was

supported in part by the Tisch Cancer Institute at Mount Sinai (P30 CA196521 Cancer Center

Support Grant, for access to Mt. Sinai cores) and by grant RO1 DE022778 from NIH/NIDCR to

P.S.

Figures, Extended Figures and Supplementary Information

Figure 1

Figure 1: Defects in craniofacial morphogenesis in *Fgfr1/2* double mutants.

(A)   Alcian blue/ alizarin red staining of mouse skulls at E18.5 showed a facial cleft in *Fgfr1$^{cKO/cKO}$* embryos, which was exacerbated in *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/+}$* mutants. *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$* mutants exhibited the most severe defect, with agenesis of proximal structures, as well as severe reduction of the mandible.

(B)   *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$*; *ROSA26$^{mT/mG}$* mutants showed fewer GFP$^+$ cNCCs in the PA1 (yellow arrow), PA2 and migratory stream relative to control embryos at E10.5.

(C)   Apoptosis was examined at E10.5 by TUNEL. Increased TUNEL positive cells were observed *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$* mutants in the lateral nasal process (LNP) compared to controls. Fewer TUNEL positive cells were observed in the medial nasal process (MNP).

(D)   Quantitation of TUNEL positive foci across different mutant genotypes show a 40-fold increase in cell death in *Fgfr1/2* double mutants in the LNP ("*" inside the bar graph represents significance level compared to control) at E10.5.

(E)   Inferior view (mandibles removed) of alcian blue/ alizarin red stained mouse skulls in *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/+}$; Bim$^{+/+}$* and *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/+}$*; *Bim$^{+/-}$* embryos showed partial rescue of medial structures, the nasal cartilage (NC) and palatine (PL) process, premaxilla (PMX), maxilla (MX), basisphenoid (BS) and palatal shelves (PS) in *Bim$^{-/+}$* mutants. Black bar (**-**) measures midline separation. Average midline separation (mm) is reduced by 2-fold (*N*=4, P>0.0109) in *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/+}$*; *Bim$^{+/-}$* as represented in the graph.

(F)   Frontal views of control *Fgfr1$^{cKO/+}$; Fgfr2$^{cKO/+}$; Bim$^{+/+}$*, *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$; Bim$^{+/+}$*, and *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$; Bim$^{+/-}$* embryos at E17.5 showing partial phenotypic rescue. Black bar (**-**) indicates intercanthal distance.

Figure 2

Figure 2: Coordinate roles *Fgfr1* and *Fgfr2* signaling mutation in development.

(A)   Schematic representation of the *Fgfr2* allelic series. Critical effectors that bind to FGFR2 are listed on the left of the *Fgfr2^{WT}* allele (WT). Critical residues for this binding are annotated on the right. Amino acid substitutions for each allele are provided to the right of all mutant alleles generated.

(B)   Co-immunoprecipitation experiments confirmed the ability of F, C and PG mutations to disrupt FRS2, CRKL and PLCγ binding to FGFR2, respectively. Constructs expressing *Fgfr2^{WT}*, *Fgfr2^{PG}*, *Fgfr2^{CPG}*, *Fgfr2^{F}* and *Fgfr2^{FCPG}*-triple Flag-tag cDNA were overexpressed in 3T3 cells. FLAG pull-downs were then immunoblotted to show interactions of FRS2, CRKL and PLCγ to FGFR2^{WT} and mutant receptors.

(C)   Sagittal view of E16.5 alcian blue/ alizarin red stained skulls from control littermate (top) and *Fgfr1* or *Fgfr2* signaling (bottom) mutants. In *Fgfr1^{F/cKO}; Fgfr2^{cKO/cKO}* mutants, the frontal bone, nasal cartilage (NC), squamosal bone (SQ), tympanic bulla (T), maxilla (MX) and mandible (MD) were affected. *Fgfr1^{FCPG/cKO}; Fgfr2^{cKO/cKO}* mutants showed a more severe defect. *Fgfr2^{F}* signaling mutants (*Fgfr1^{cKO/cKO}; Fgfr2^{F/cKO}*) showed defects in mandible along with reduced ossification of the premaxilla (PMX) and maxilla and loss of the squamosal bone and tympanic bulla.

(D)   DAPI stained frontal facial view for *Fgfr1^{F}* and *Fgfr2^{F}* mutant embryos at E15.5. *Fgfr1^{F/F}; Fgfr2^{+/F}* embryos showed a severe facial cleft compared to other genotypes.

(E)   CDX2 and EOMES expression in *Fgfr1^{FCPG}* and *Fgfr2^{FCPG}* compound signaling mutants at E7.5. In contrast to *Fgfr1^{-/-}; Fgfr2^{-/-}* which fail to implant on the 129S4 co-isogenic background, we could recover *Fgfr1^{FCPG/FCPG}; Fgfr2^{FCPG/FCPG}* compound mutants at E7.5 but not at E10.5. *Fgfr1^{FCPG/FCPG}; Fgfr2^{FCPG/FCPG}* mutants were growth-retarded but expressed mesodermal EOMES and the trophectoderm marker CDX2.

(F)   iFNP cells (*Fgfr1^{CRISPR-KO}*) derived from the indicated genotypes, *Fgfr2^{wt/wt}* (*R2^{wt/w}*), *Fgfr2^{F/F}* (*R2^{F/F}*), *Fgfr2^{CPG/CPG}* (*R2^{CPG/CPG}*) and *Fgfr2^{FCPG/FCPG}* (*R2^{FCPG/FCPG}*), were stimulated with FGF1 for the indicated times (0-60 mins) to analyze pERK1/2 and pAKT activation. Quantification of pathway activation for pERK1/2 and pAKT, normalized to GAPDH, is reported as mean ± standard deviation with a minimum of three independent biological replicates.

Figure 3

Figure 3: A kinase dead mutation in *Fgfr2* recapitulates multiple aspects of the *Fgfr2*[^-/-]

phenotype.

(A)   Morphological examination of *Fgfr2* kinase dead phenotype at E10.5. *Fgfr2*$^{KD/-}$ and

Fgfr2$^{KD/KD}$ mutants had no limb buds (red arrows), and had defects in the allantois which was

loosely or incompletely held by the chorion to the ectoplacental cone, along with a dilated

pericardium similar to *Fgfr2*$^{-/-}$ mutants. Both *Fgfr2*$^{KD/-}$ and *Fgfr2*$^{KD/KD}$ mutants displayed severe

posterior truncations and *Fgfr2*$^{KD/KD}$ mutants showed a more severe phenotype. *Fgfr2*$^{KD/KD}$

mutants also displayed craniofacial defects with poorly developed facial prominences compared

to *Fgfr2*$^{KD/-}$ mutants.

(B)   Wholemount mRNA *in situ* hybridization for cNCC mesenchyme marker *Msx1* at E10.5.

Msx1 was expressed in the facial prominences (LNP, MNP and mandibular prominences) in *Fgfr2*$^{-}$

$^{/-}$ embryos at similar levels compared to control *Fgfr2*$^{+/-}$ embryo. *Msx1* expression was also

observed in *Fgfr2*$^{KD/-}$ mutants but was reduced in *Fgfr2*$^{KD/KD}$ mutants.

(C)   Wholemount mRNA in situ hybridization for midline marker *Nkx2.1*. *Nkx2.1* was expressed

in the midline floor plate of *Fgfr2*$^{-/-}$ mutants, but not detected in *Fgfr2*$^{KD/-}$ and *Fgfr2*$^{KD/KD}$ mutants.

(D)   *Meox1* mRNA expression in somites of E10.5 *Fgfr2*$^{+/-}$ embryos was comparable to *Fgfr2*$^{-/-}$

mutants, but severely reduced in both *Fgfr2*$^{KD/-}$ and *Fgfr2*$^{KD/KD}$ mutants (white arrows) along with

defects in axis elongation and posterior somite formation (red arrow).

Figure 4

A

Starvation (T_0) ← (T_{12}) →

Starve   FGF   PDGF   Serum

*Fgfr1^cKO/+^; Fgfr2^cKO/+^*

*Fgfr1^FCPG/cKO^; Fgfr2^cKO/cKO^*

*Fgfr1^cKO/cKO^; Fgfr2^cKO/cKO^*

B

Starvation   FGF   PDGF   Serum

GFP
Paxillin
Paxillin

*Fgfr1^cKO/+^; Fgfr2^cKO/+^*

*Fgfr1^FCPG/cKO^; Fgfr2^cKO/cKO^*

*Fgfr1^cKO/cKO^; Fgfr2^cKO/cKO^*

*Fgfr1^CR-KO^; Fgfr2^+/+^*

*Fgfr1^CR-KO^; Fgfr2^FCPG/FCPG^*

*Fgfr1^CR-KO^; Fgfr2^CR-KO^*

Paxillin/ Paxillin/ Phalloidin

Figure 4: Cell matrix adhesion properties are retained in FCPG mutants.

(A) Scratch/wound healing assay was used to study cell spreading. Control (*Fgfr1$^{cKO/+}$;*
Fgfr2$^{cKO/+}$) and *Fgfr1$^{FCPG/cKO}$; Fgfr2$^{cKO/cKO}$* primary FNP cells showed active spreading over 12
hours (T$_{12}$) in response to FGF1, PDGF-A, or serum. *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$* mutant cells showed
limited spreading upon FGF stimulation but normal spreading in response to PDGF or serum.

(B) Focal-adhesion formation was assayed by Paxillin immunostaining in *Fgfr1* and *Fgfr2*
signaling mutant primary FNP cells. GFP$^+$ *Fgfr1$^{cKO/+}$; Fgfr2$^{cKO/+}$; ROSA26$^{mT/mG}$* control,
Fgfr1$^{FCPG/cKO}$; Fgfr2$^{cKO/+}$; ROSA26$^{mT/mG}$, and *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$; ROSA26$^{mT/mG}$* mutant cells
were treated with FGF1, PDGFA or serum for 3 hours before analyzing Paxillin localization to
focal adhesions. In control cells and *Fgfr1$^{FCPG/cKO}$; Fgfr2$^{cKO/+}$*, we detected multiple Paxillin$^+$ foci
upon FGF1, PDGFA, or serum stimulation. *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$; ROSA26$^{mT/mG}$* mutant FNP
cells failed to form focal adhesions in response to FGF, in contrast to PDGF or serum (quantified
in Extended Data Fig. 9A). Cell spreading properties were also analyzed for *Fgfr2$^{WT/WT}$*
(*Fgfr1$^{CRISPR-KO}$; Fgfr2$^{WT/WT}$*), *Fgfr2$^{FCPG/FCPG}$* (*Fgfr1$^{CRISPR-KO}$; Fgfr2$^{FCPG/FCPG}$*), and *Fgfr2$^{-/-}$* (*Fgfr1$^{CRISPR-}$*
^{KO}; Fgfr2$^{CRISPR-KO}$) -iFNP cells in which *Fgfr1* was disrupted. Addition of FGF1, PDGFA or serum
resulted in robust cell spreading and formation of Paxillin$^+$ focal adhesion in *Fgfr1$^{CRISPR-KO}$;*
Fgfr2$^{FCPG/FCPG}$ and *Fgfr1$^{CRISPR-KO}$; Fgfr2$^{+/+}$* -iFNP cells. *Fgfr2$^{-/-}$* (*Fgfr1$^{CRISPR-KO}$; Fgfr2$^{CRISPR-KO}$*) -iFNP
cells showed severe defects in focal adhesion formation upon FGF1 treatment, but was
unaffected in response to PDGFA or serum (quantified in Extended Data Fig. 9E).

Figure 5

Figure 5: Cell-cell adhesion is retained in *Fgfr* signaling mutants.

(A)   GFP$^+$ primary FNP cells from control *Fgfr1$^{cKO/+}$; Fgfr2$^{cKO/+}$; ROSA26$^{mT/mG}$* and *Fgfr1$^{FCPG/cKO}$; Fgfr2$^{cKO/cKO}$; ROSA26$^{mT/mG}$* embryos formed extensive cell-cell contacts, in contrast to *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$; ROSA26$^{mT/mG}$* mutants. Quantification is provided in Extended Data Fig. 9F. β-catenin was localized along the cell contact boundaries in GFP$^+$ FNP cells from control and *Fgfr1$^{FCPG/cKO}$; Fgfr2$^{cKO/cKO}$; ROSA26$^{mT/mG}$* embryos in contrast to *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$; ROSA26$^{mT/mG}$* mutants.

(B)   β-catenin immunofluorescence was quantified and represented as mean for each genotype.

(C)   Extensive cell-cell contacts with β-catenin localization (insets) were observed *in vivo* in control *Fgfr1$^{cKO/+}$; Fgfr2$^{cKO/+}$; ROSA26$^{mT/mG}$* and *Fgfr1$^{FCPG/cKO}$; Fgfr2$^{cKO/cKO}$; ROSA26$^{mT/mG}$* cells in both developing LNP and MNP at E11.5. In contrast, *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$; ROSA26$^{mT/mG}$* double null mutants GFP$^+$ cells remained sparse in the LNP and showed no localized β-catenin.

(D)   Model of FGF-mediated cell signaling pathways. In wild-type (WT) cells, activation by FGFs engages a canonical RTK signal transduction pathway, leading for the activation of ERK1/2, PI3K/AKT, PLCγ, and additional pathways. In addition, FGFs activate non-canonically both cell-matrix as well as cell-cell adhesion, in a kinase-dependent manner, possibly facilitated through interactions of the FGF receptors through their extracellular domain with cell adhesion receptors. *Fgfr1^{FCPG}* or *Fgfr2^{FCPG}* mutant cells fail to activate a classical RTK signal transduction pathway (light grey) but can still promote cell adhesion (black), as their kinase activity has not been disrupted. In null mutant cells, neither FGF-induced cell signaling nor cell adhesion are observed (light grey), since the receptors are not expressed.

Extended Data Figure 1

A
DAPI | FGFR1-GFP | FGFR1-mCherry | Merged
E10.5

B
Fgfr1^{cKO/+}; Fgfr2^{cKO/+} | *Fgfr1^{cKO/cKO}; Fgfr2^{cKO/+}* | *Fgfr1^{cKO/cKO}; Fgfr2^{cKO/cKO}*
N F I
PMx
Mx
MD
E18.5 | MD E18.5 | E18.5

C
Fgfr1^{cKO/+}; Fgfr2^{cKO/+} | *Fgfr1^{cKO/cKO}; Fgfr2^{cKO/+}* | *Fgfr1^{cKO/cKO}; Fgfr2^{cKO/cKO}*
N F
SQ
CRP CDP
PMX
MX
MD AGP

D
Fgfr1^{cKO/+}; Fgfr2^{cKO/+} | *Fgfr1^{cKO/cKO}; Fgfr2^{cKO/+}* | *Fgfr1^{cKO/cKO}; Fgfr2^{cKO/cKO}*
E14.5

Extended Data Fig. 1: Defects in craniofacial morphogenesis in *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$* double mutants.

(A)   Spatial domain of FGFR1 and FGFR2 expression in facial prominences at E10.5. GFP and mCherry immunohistochemistry were used to detect expression from *Fgfr1-GFP* and *Fgfr2-mCherry* reporter alleles. GFP expression was primarily restricted to the mesenchyme (green arrow). Although mCherry expression was restricted to the epithelium, many cells in the mesenchyme also express mCherry (red arrow). mCherry expression is downregulated in the epithelium lining the nasal pit (yellow asterisk). FGFR1 and FGFR2 co-expression in the craniofacial skeleton is also observed E17.5 (Supplementary Information Fig. S1a).

(B)   Micro-CT analysis of ossified structures in the head at E18.5. An overall reduction or complete loss of ossification of cNCC-derived facial bones in *Fgfr1/2* conditional double mutant is seen compared to the control. Frontal (F) and nasal bones (N), pre-maxilla (PMX) and maxilla (MX) were severely affected. Mesoderm derived bones including parietal and interparietal (I) bones remained unaffected (yellow asterisk). *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/+}$* mutants showed an overall reduction in the size of the mandible (MD), with the proximal structures most severely affected. A more detailed morphometry analysis regarding changes in skull length, intercanthal distance, length of frontal bone and length of nasal cartilage at E18.5 for different genotypes is provided in Supplementary Information Fig S1B.

(C)   Sagittal view of alcian blue/ alizarin red stained mouse skull at E16.5 for different genotypes. Frontal (F) and nasal (N) bone, mandible (MD), premaxilla (PMX), maxilla (MX) and the squamous (SQ) bone was severely affected in *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$* conditional double mutants. In *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/+}$* mutants also showed partial mandibular defect where proximal condylar (CDP), coronoid (CRP) and angular process (AGP) of the mandible was affected.

(D)   Alcian blue/ alizarin red staining of mouse skulls at E14.5. Alizarin red staining mark ossified region in mandible and maxilla in control (*Fgfr1$^{cKO/+}$; Fgfr2$^{cKO/+}$*) skull. Yellow arrowhead marks the Meckel's cartilage. In *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/+}$* mutants, we observed reduced ossification (red arrow) in the mandible with a significantly smaller Meckel's cartilage (yellow arrow). Meckel's cartilage was lost in *Fgfr1/2* conditional double mutant with no observable ossification.

Extended Data Figure 2

Extended Data Fig. 2: Molecular changes and defects in craniofacial ossification in
***Fgfr1^{cKO/cKO}; Fgfr2^{cKO/cKO}* double mutants.**

(A)　　Conditional *Fgfr1-Fgfr2* null mutant embryos carrying the *ROSA26^{mT/mG}* reporter analyzed
in whole mount. cNCCs in the head express GFP (green). At E9.5 (top panel), *Fgfr1/2* conditional
double mutants (*Fgfr1^{cKO/cKO}; Fgfr2^{cKO/cKO}*; *ROSA26^{mT/mG}*) had hypoplastic pharyngeal arches PA1
and PA2 (yellow arrows), shown by reduced GFP fluorescence compared to *Fgfr1-Fgfr2*
conditional heterozygotes. *Fgfr1^{cKO/+}; Fgfr2^{cKO/cKO}* and *Fgfr1^{cKO/cKO}; Fgfr2^{cKO/+}* mutants appeared
normal at this stage. Frontal view at E10.5 (lower panels). A wide midline separation was
observed in both *Fgfr1/2* conditional double (*Fgfr1^{cKO/cKO}; Fgfr2^{cKO/cKO}*; *ROSA26^{mT/mG}*) and
Fgfr1^{cKO/cKO}; Fgfr2^{cKO/+}; *ROSA26^{mT/mG}* mutants (yellow asterisk) compared to controls and
Fgfr1^{cKO/+}; Fgfr2^{cKO/cKO}; *ROSA26^{mT/mG}* mutants.

(B)　　Whole mount mRNA *in situ* hybridization was used to compare the expression of *Alx3,*
Msx1, Six3, Nkx2.1, Fgf8 and *Shh* at E10.5 in the facial primordia for *Fgfr1^{cKO/+}; Fgfr2^{cKO/+}* control
and *Fgfr1^{cKO/cKO}; Fgfr2^{cKO/cKO}* mutants. *Msx1* expression was reduced, and *Alx3*, *Six3* and *Nkx2.1*
expression were lost in *Fgfr1/2* conditional double mutants. The expression of *Fgf8* and *Shh*
remained unaffected in the *Fgfr1^{cKO/cKO}; Fgfr2^{cKO/cKO}* mutants.

472
473

Extended Data Figure 3

A
GFP
EdU
DAPI
LNP
MNP
Fgfr1^{cKO/+}; Fgfr2^{cKO/+};
Rosa26^{mT/mG}

Fgfr1^{cKO/+}; Fgfr2^{cKO/+};
Rosa26^{mT/mG}
EdU

Fgfr1^{cKO/cKO}; Fgfr2^{cKO/cKO};
Rosa26^{mT/mG}
LNP
MNP
GFP
EdU
DAPI

Fgfr1^{cKO/cKO}; Fgfr2^{cKO/cKO};
Rosa26^{mT/mG}
EdU

B
Percentage of
proliferative cells
Percentage of EdU$^+$ cells
R1^{cKO/+}; R2^{cKO/+}
R1^{cKO/cKO}; R2^{cKO/cKO}

Extended Data Fig. 3: Cell proliferation in *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$* double mutants.

(A)   EdU incorporation was used to assay cell proliferation in *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$;*

ROSA26$^{mT/mG}$ double mutants. No change in the proliferation index of GFP$^+$ cNCC lineage cells

was observed at E10.5 in the MNP and LNP across various genotypes.

(B)   Quantification of the mean percentage of EdU$^+$ cells normalized to GFP$^+$ cNCC lineage

cells, in control versus double null mutant cells suggests cell proliferation remained unaffected in

Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$ double mutants.

Extended Data Figure 4

Extended Data Fig. 4: Allelic series of *Fgfr2* signaling mutations.

(A) Representation of *Fgfr2* WT locus, targeting vectors and the corresponding targeted loci

for C, PG and F alleles, the *Fgfr2^{FCPG}* locus, and the KD allele. Targeting vectors C, PG and F

introduced Y466F, Y769F/Y779F and L434A/R426A mutations sequentially to the *Fgfr2* locus,

respectively. The targeting vector for KD allele introduced the K517A mutation. ES cells were

targeted first with C targeting vector generating *Fgfr2^{+/C}* mutant cells. *Fgfr2^{+/C}* ES cells were

targeted with PG targeting vector generating either *Fgfr2^{C/PG}* or *Fgfr2^{+/CPG}* mutant cells. Finally,

Fgfr2^{+/CPG} was targeted with F targeting vector resulting in *Fgfr2^{F/CPG}* or *Fgfr2^{+/FCPG}* mutant ES

cells. Multiple LoxP sites were introduced at the *Fgfr2* locus during successive rounds of targeting,

and the neo cassette was removed *in vivo* or in culture before generating signaling mouse mutants.

We were unable to perform a conditional mutation for the *Fgfr2^{FCPG}* allele due to the serial

retention of multiple lox sites during the generation of this allele. Abbreviation: L, loxP site; F, FRT

site; DTA, Diptheria toxin A cassette; Neo, Neomycin resistance cassette.

(B) Southern blot confirmation for proper targeting of F, C, PG and KD mutations at the *Fgfr2*

locus. 5' probes used for each targeting events are indicated. BamHI restriction enzyme

generated 5kb and 2.6kb band for WT and F targeted loci, respectively. SphI digestion generated

10kb and 8.4 kb fragment for WT and C targeted loci, respectively. KpnI digestion generated a

5.6kb band for WT locus and a 7.6kb band for PG targeted locus. EcoRI and EcoRV restriction

digestion generated 7.7kb WT fragment and 9.7kb KD-mutant fragment.

(C) Western blot from facial prominence cells harvested from E9.5 wild-type and *Fgfr2^{KD/KD}*

embryos show full length FGFR2^{KD} protein expression.

Extended Data Figure 5

A

Fgfr2^{F/+} P21 | *Fgfr2^{F/F}* P21
Fgfr2^{FCPG/+} P21 | *Fgfr2^{FCPG/FCPG}* P21
Fgfr2^{+/+} P21 | *Fgfr2^{KD/+}* P21

B

Fgfr2^{F/+} *Fgfr2^{FCPG/+}* *Fgfr2^{+/+}*

Fgfr2^{F/F} *Fgfr2^{FCPG/FCPG}* *Fgfr2^{KD/+}*

C

No. of branch points

Area of lacrimal gland (Px)

Extended Data Fig. 5: Lacrimal gland defects in *Fgfr2* signaling mutants.

(A) Sagittal view of the left eye at P21 for the mentioned genotypes. $Fgfr2^{F/F}$, $Fgfr2^{FCPG/FCPG}$

and $Fgfr2^{KD/+}$ mice showed fully penetrant defects with onset at P15 which included laceration

around the eye. More severe defects were observed in $Fgfr2^{KD/+}$ mutants.

(B) The lacrimal gland from signaling mutants was dissected at P5 and stained with

acetocarmine. Representative images are shown here for $Fgfr2^{F/F}$, $Fgfr2^{FCPG/FCPG}$ and $Fgfr2^{KD/+}$

compared to littermate $Fgfr2^{F/+}$, $Fgfr2^{FCPG/+}$ and $Fgfr2^{+/+}$ controls.

(C) Acetocarmine stained preparation of exorbital lacrimal glands at P5 from $Fgfr2^{F/F}$,

$Fgfr2^{FCPG/FCPG}$ and $Fgfr2^{KD/+}$ mice showed reduced branching and reduced overall size of the gland.

Flat mounts of the acetocarmine stained lacrimal glands for each genotype were microscopically

analyzed and primary and secondary branching points were counted manually. The bar graph

shows the mean ± standard deviation of the relative number of branch points for each genotype

analyzed. The relative size of the lacrimal gland for each genotype was also assessed by

comparing images acquired in the same magnification as a mean of the area observed in flat

mounts. $Fgfr2^{F/F}$, $Fgfr2^{FCPG/FCPG}$ and $Fgfr2^{KD/+}$ compared to littermate $Fgfr2^{F/+}$, $Fgfr2^{FCPG/+}$ and

$Fgfr2^{+/+}$controls show reduced branching and a reduction in the overall size of the gland.

Extended Data Figure 6

Extended Data Fig. 6: Analysis of *Fgfr1* and *Fgfr2* single and compound signaling mutants.

(A)　　Micro-CT analysis of control *Fgfr1$^{cKO/+}$; Fgfr2$^{cKO/+}$* embryos at E18.5 compared to *Fgfr1$^{F/cKO}$;*

Fgfr2$^{cKO/+}$ and *Fgfr1$^{FCPG/cKO}$; Fgfr2$^{cKO/+}$* embryos at similar stages. The mandible (red arrow) was

more severely affected in *Fgfr1$^{FCPG/cKO}$; Fgfr2$^{cKO/+}$* than in *Fgfr1$^{F/cKO}$; Fgfr2$^{cKO/+}$* embryos.

(B)　　Frontal facial view of DAPI-stained E10.5 *Fgfr1$^{F/cKO}$; Fgfr2$^{cKO/+}$* embryos compared to

control *Fgfr1$^{cKO/+}$; Fgfr2$^{cKO/+}$* embryos. Midline separation (yellow asterisk) was observed and facial

prominences were hypoplastic (yellow arrow), however, no defect in the mandible (red arrow) was

observed.

(C)　　Frontal facial view of DAPI-stained E10.5 *Fgfr1$^{FCPG/cKO}$; Fgfr2$^{cKO/+}$* embryos compared to

control *Fgfr1$^{cKO/+}$; Fgfr2$^{cKO/+}$* embryos. In the mutants, midline separation (yellow asterisk) and

facial prominence (yellow arrow) defects were more severe than *Fgfr1$^{F/cKO}$; Fgfr2$^{cKO/+}$* embryos. A

defect in the mandible (red arrow) was also more severe than *Fgfr1$^{F/cKO}$; Fgfr2$^{cKO/+}$* embryos.

(D-E) Cell-death analysis by TUNEL staining in the facial prominences of (D) *Fgfr1$^{F/cKO}$;*

Fgfr2$^{cKO/+}$ and (E) *Fgfr1$^{FCPG/cKO}$; Fgfr2$^{cKO/+}$* embryos at E10.5, showing multiple dying cells in both

the MNP and the LNP compared to the *Fgfr1$^{cKO/+}$; Fgfr2$^{cKO/+}$* controls.

(F)　　Frontal facial view of DAPI-stained *Fgfr1^{F}* and *Fgfr2^{F}* compound mutant embryos at E10.5.

Fgfr1$^{F/F}$; Fgfr2$^{+/F}$ embryos had defects in the development of facial prominences, as opposed to

Fgfr1$^{F/+}$; Fgfr2$^{F/+}$ controls. Mandibular prominences were hypoplastic, however, the maxillary

prominences, FNP and MNP were more severely affected. *Fgfr1$^{F/F}$; Fgfr2$^{+/+}$* mutants, *Fgfr1$^{+/+}$;*

Fgfr2$^{F/F}$ mutants, and *Fgfr1$^{F/+}$; Fgfr2$^{F/F}$* mutants did not show obvious defects at this stage. *Fgfr1$^{F/F}$;*

Fgfr2$^{+/F}$ mutant embryos were also developmentally retarded. We did not recover *Fgfr1$^{F/F}$; Fgfr2$^{F/F}$*

mutant embryos by E9.5.

(G)　　The craniofacial phenotype at E10.5 of DAPI-stained *Fgfr1^{FCPG}* and *Fgfr2^{FCPG}* compound

mutant embryos. We observed the posterior truncation for *Fgfr1$^{FCPG/FCPG}$* mutant embryos. The

introduction of one copy of *Fgfr2^{FCPG}* allele on a *Fgfr1$^{FCPG/FCPG}$* background resulted in severe

growth retardation of the facial prominences compared to control *Fgfr1$^{+/FCPG}$; Fgfr2$^{FCPG/FCPG}$*

embryos. *Fgfr2$^{FCPG/FCPG}$* and *Fgfr1$^{+/FCPG}$; Fgfr2$^{FCPG/FCPG}$* embryos did not exhibit abnormal

phenotypes and were comparable to WT embryos.

Extended Data Figure 7

A

WT-iFNP

Relative expressiomn
FgfR1 FgfR2 FgfR3 FgfR4
■ R1-WT-R2WT ■ CRISPR

F-iFNP

Relative expressiomn
FgfR1 FgfR2 FgfR3 FgfR4
■ R1WT-R2F ■ CRISPR

CPG-iFNP

Relative expressiomn
FgfR1 FgfR2 FgfR3 FgfR4
■ R1WT-R2CPG ■ CRISPR

FCPG-iFNP

Relative expressiomn
FgfR1 FgfR2 FgfR3 FgfR4
■ R1WT-R2FCPG ■ CRISPR

B

Relative expressiomn
Alx4 Dlx5 Lhx8 FoxD1 Twist2 Msx1 Sox9 Dlx1
■ WT RTPCR ■ iFNP RTPCR

C

Extended Data Fig. 7: Characterization of *Fgfr1 null* mutant iFNP cells.

(A)   *Fgfr1, Fgfr3, Fgfr4* and *Fgfr2^{WT}* mRNA expression in wild-type and *Fgfr2^F, Fgfr2^{CPG}* and

Fgfr2^{FCPG} iFNP cells before and after *Fgfr1*-CRISPR disruption, as determined by RT-qPCR.

(B)   *Alx4, Dlx1, Dlx5, FoxD1, Twist1* and *Lhx8* were all expressed at similar levels by

quantitative real-time PCR (qRT-PCR) in both immortalized FNP (iFNP) cells and primary FNP

cells derived from wild-type embryos. Only *Msx1* expression was 1.8-fold higher in wild-type

primary FNP cells compared to iFNP cell lines. These results identify iFNP cells as a suitable cell

type to interrogate FGF signaling functions in craniofacial development.

(C)   *WT*-iFNP cells were serum starved overnight and stimulated with either 50 ng/mL FGF1

or 50 ng/mL FGF8-b and 5 μg/mL heparin for the indicated times (0-60 mins). Robust ERK1/2

activation was observed with FGF1 treatment compared to FGF8-b treatment.

Extended Data Fig. 8: Intracellular signaling upon FGF1 stimulation.

(A)   iFNP cells derived from the indicated genotypes, $Fgfr2^{wt/wt}$ ($R2^{wt/w}$), $Fgfr2^{F/F}$ ($R2^{F/F}$),

$Fgfr2^{CPG/CPG}$ ($R2^{CPG/CPG}$) and $Fgfr2^{FCPG/FCPG}$ ($R2^{FCPG/FCPG}$), were serum starved overnight and

stimulated with 50 ng/mL FGF1 and 5 µg/mL heparin for the indicated times (0-60 mins).

Phospho-blots were stripped and reblotted with GAPDH as a loading control. Activation of pP38,

pPLCγ, pJNK and pSTAT3 was investigated.

(B-E)   Quantification of pathway activation for (B) pP38, (C) pPLCγ, (D) pJNK and (E) pSTAT3

normalized to the GAPDH loading control, is reported as mean ± standard deviation with a

minimum of three independent biological replicates.

(F-G) $Fgfr1^{FCPG}$ $Fgfr2$ null iFNP ($Fgfr1^{FCPG}$; $Fgfr2^{CRISPR-KO}$) cells generated using CRISPR/Cas9.

Upon FGF1 stimulation, we observed peak activation of (F) pERK1/2 after 2 mins in $Fgfr1^{FCPG/FCPG}$

iFNP cells. We did not observe activation over background in $Fgfr1^{FCPG/FCPG}$; $Fgfr2^{CRISPR-KO}$ iFNP

cells. (G) pAKT activation was undetectable in both $Fgfr1^{FCPG/FCPG}$; $Fgfr2^{CRISPR-KO}$ in contrast to

$Fgfr1^{FCPG/FCPG}$; $Fgfr2^{+/+}$ iFNP cells.

Extended Data Figure 9

A — *Fgfr1^{cKO/+}; Fgfr2^{cKO/+}; ROSA26^{mT/mG}*, *Fgfr1^{FCPG/cKO}; Fgfr2^{cKO/+}; ROSA26^{mT/mG}*, *Fgfr1^{cKO/cKO}; Fgfr2^{cKO/cKO}; ROSA26^{mT/mG}*

B — *Fgfr1^{+/cKO}; Fgfr2^{+/cKO}*, *Fgfr1^{FCPG/cKO}; Fgfr2^{cKO/cKO}*, *Fgfr1^{cKO/cKO}; Fgfr2^{cKO/cKO}*

C — *Fgfr1^{cKO/+}; Fgfr2^{cKO/+}*, *Fgfr1^{FCPG/cKO}; Fgfr2^{cKO/+}*, *Fgfr1^{cKO/cKO}; Fgfr2^{cKO/cKO}*

D

E

F

Extended Data Fig. 9: FGFR signaling in cell-matrix and cell-cell adhesion.

(A) Quantification of Paxillin immunostaining mean fluorescence intensity in FNP cells, shown

for different genotypes in Fig. 4B. Mean fluorescence intensity in starvation media act as a

baseline. Upon serum stimulation or PDGF treatment, cells from all genotypes showed a robust

increase in focal adhesions. Upon FGF treatment, control cells and $Fgfr1^{FCPG/cKO}$; $Fgfr2^{cKO/+}$;

$ROSA26^{mT/mG}$ FNP cells showed robust paxillin accumulation, in contrast to $Fgfr1^{cKO/cKO}$;

$Fgfr2^{cKO/cKO}$; $ROSA26^{mT/mG}$ double mutant cells.

(B) Western Blot analysis of freshly harvested FNP cells from control, $Fgfr1^{FCPG/cKO}$; $Fgfr2^{cKO/+}$;

$ROSA26^{mT/mG}$ and $Fgfr1^{cKO/cKO}$; $Fgfr2^{cKO/cKO}$; $ROSA26^{mT/mG}$ double mutant FNP cells for pFAK, total

FAK, β-catenin and Paxillin. GAPDH was used as a loading control. Total FAK and β-catenin

levels remained constant across all genotypes.

(C) Bar graph shows relative levels of pFAK, total FAK and Paxillin. pFAK and Paxillin levels

were comparable in control and $Fgfr1^{FCPG/cKO}$; $Fgfr2^{cKO/+}$; $ROSA26^{mT/mG}$ FNP cells, but severely

reduced in $Fgfr1^{cKO/cKO}$; $Fgfr2^{cKO/cKO}$; $ROSA26^{mT/mG}$ FNPs. Total FAK levels remained unchanged.

(D) Freshly harvested $Fgfr1^{cKO/+}$; $Fgfr2^{cKO/+}$; $ROSA26^{mT/mG}$ control, $Fgfr1^{FCPG/cKO}$; $Fgfr2^{cKO/+}$;

$ROSA26^{mT/mG}$, and $Fgfr1^{cKO/cKO}$; $Fgfr2^{cKO/cKO}$; $ROSA26^{mT/mG}$ double mutant FNP cells were treated

with serum or FGF1 for 15 mins, before assessing levels of pFAK by western blot. $Fgfr1^{cKO/+}$;

$Fgfr2^{cKO/+}$; $ROSA26^{mT/mG}$ control and $Fgfr1^{FCPG/cKO}$; $Fgfr2^{cKO/+}$; $ROSA26^{mT/mG}$ FNP cells showed a

comparable increase in FAK activation upon FGF stimulation. In contrast, $Fgfr1^{cKO/cKO}$; $Fgfr2^{cKO/cKO}$;

$ROSA26^{mT/mG}$ double mutant FNP cells failed to activate FAK in both serum-stimulated and FGF

treated conditions.

(E) Quantification of mean fluorescence from Paxillin immunostaining in $Fgfr2^{WT/WT}$

($Fgfr1^{CRISPR-KO}$ $Fgfr2^{WT/WT}$), $Fgfr2^{FCPG/FCPG}$ ($Fgfr1^{CRISPR-KO}$ $Fgfr2^{FCPG/FCPG}$), and $Fgfr2^{-/-}$ ($Fgfr1^{CRISPR-}$

^{KO} $Fgfr2^{CRISPR-KO}$) -iFNP cells upon serum starvation and growth factor treatments shown in Fig.

4B. Upon serum stimulation or PDGF treatment, cells from all genotypes showed a robust

increase in focal adhesions. Upon FGF treatment, both control and $Fgfr1^{CRISPR-KO}$ $Fgfr2^{FCPG/FCPG}$

iFNP cells showed robust paxillin accumulation at focal adhesions, whereas $Fgfr1^{CRISPR-KO}$ $Fgfr2$

$^{CRISPR-KO}$ double null mutant iFNP cells showed poor response.

(F) Quantification of percentage of cells that form cell boundaries (n=500 contacts for each

genotype) in $Fgfr1^{cKO/+}$; $Fgfr2^{cKO/+}$; $ROSA26^{mT/mG}$ control, $Fgfr1^{FCPG/cKO}$; $Fgfr2^{cKO/+}$; $ROSA26^{mT/mG}$,

and $Fgfr1^{cKO/cKO}$; $Fgfr2^{cKO/cKO}$; $ROSA26^{mT/mG}$ double mutant FNP cells. When freshly harvested

primary FNP cells were serum stimulated for 3 hrs, both control and $Fgfr1^{FCPG/cKO}$; $Fgfr2^{cKO/+}$;

$ROSA26^{mT/mG}$ cells formed comparable numbers of cell-cell contacts (Figure 7D). In contrast, this

number was reduced to half in double mutant FNP cells.

Supplementary Information Image S1

Supplementary Information Figure S1: Craniofacial ossification and organogenesis defects in *Fgfr1/2* mutants.

(A)   Spatial domain of FGFR1 and FGFR2 expression in the frontal bone at E17.5. GFP and mCherry immunohistochemistry was used to detect expression from *Fgfr1-GFP* and *Fgfr2-mCherry* reporter alleles. Co-expression of both FGFR1-GFP and FGFR2-mCherry was observed in the cartilage and the perichondrium (yellow arrow).

(B)   Quantification of skull length, intercanthal distance, length of frontal and nasal bone for different genotypes measured in millimeters (mm).

(C)   Histological examination at E15.5 showed defective organogenesis in multiple organs, including the palate, tongue, and skeleton. In the double mutants several skeletal structures such as palate (P), nasopharyngeal lumen (N) and pre-sphenoid (PS) were affected compared to the controls. Non-neural crest derived muscles in the tongue were also affected. SMA immunofluorescence showed organized muscle fiber differentiation in controls. In the *Fgfr1/2* mutants, muscle fiber differentiation appeared random and disorganized.

(D)   Differentiation of craniofacial skeleton was assessed by studying the expression of cartilage differentiation marker, *Col2a1,* on sections. mRNA *in situ* hybridization analysis at E14.5 showed *Col2a1* expression in the chondrocranium (skull base) and mandible in the control (inset shows low magnification view of the region). In the double mutants, however, expression was restricted to a narrow band of cells in the skull base. Expression of *Col2a1* and of *Col10a1* and Runx2 (mature cartilage markers) was assessed at E17.5 on sections (inset shows a low magnification view of the region). *Col2a1* mRNA expression was restricted to the skull base, mandible and frontal bone in controls. A similar expression was also observed in conditional double mutants. At E17.5, *Col10a1* mRNA expression was much broader in the controls. In conditional double mutants, *Col10a1* expression was undetectable suggesting a block in terminal chondrogenic differentiation. We detected the expression of mature cartilage and late osteoblast marker, RUNX2, in both controls and the conditional double mutants.

(E)   Increased cell-death observed in conditional double mutants was partially rescued in *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$; Bim$^{+/-}$* mutants at E10.5. More TUNEL$^+$ cells were observed in the LNP in *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$* and *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/+}$* mutants compared to corresponding *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$; Bim$^{+/-}$* and *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/+}$; Bim$^{+/-}$* counterparts. Compared to controls, reduced TUNEL positive foci were observed in *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$; Bim$^{+/-}$* mutants.

Supplementary Information Image S2

Supplementary Information Figure S2: Growth defects in *Fgfr* signaling mutants.

(A)   Growth rate (in terms of weight gain) of homozygous $Fgfr2^{F/F}$, $Fgfr2^{C/C}$, $Fgfr2^{PG/PG}$,

$Fgfr2^{CPG/CPG}$, and $Fgfr2^{FCPG/FCPG}$ female versus control until 6 weeks shown as mean ± SEM. *

indicates p-value <0.05. We noted a decrease in growth rate for $Fgfr2^{F/F}$, $Fgfr2^{PG/PG}$, and

$Fgfr2^{FCPG/FCPG}$ mutant mice compared to control.

(B) $Fgfr2^{PG/PG}$ and $Fgfr2^{CPG/CPG}$ skeletal preparation at P0 showed fusion of the caudal

vertebrae, resulting in a kinked or curly tail phenotype in rare cases (black arrow). Hemizygote

$Fgfr2^{C/-}$, $Fgfr2^{PG/-}$ and $Fgfr2^{CPG/-}$ mice were weighed at 6 weeks. $Fgfr2^{PG/-}$ were smaller than their

control littermate.

(C)   Whole mount immunohistochemistry for neurofilament staining revealed trigeminal nerve

defects at E10.5 in $Fgfr2^{FCPG/-}$ embryo but not in $Fgfr2^{F/-}$. In $Fgfr2^{F/+}$, $Fgfr2^{FCPG/+}$ and $Fgfr2^{F/-}$

embryos, trigeminal nerve migrated to the anterior domain of the PA1 (red arrow) at E10.5.

Trigeminal nerves fail to migrate into the pharyngeal arch of $Fgfr2^{FCPG/-}$ mutants.

Supplementary Information Table 1: Recovery of *Fgfr2* homozygous and *Fgfr2* hemizygous

signaling mutants.

Recovery of *Fgfr2* signaling mutants			
P4-P10	Total	*/*	Recovery
$R2^F$	386	85	22%
$R2^C$	505	131	26%
$R2^{PG}$	475	78	16%
$R2^{CPG}$	396	70	18%
$R2^{FCPG}$	646	102	16%
$R2^{KD}$	38	0	0
E18.5	Total	*/*	Recovery
$R2^F$	97	22	23%
$R2^C$	53	14	26%
$R2^{PG}$	64	14	22%
$R2^{CPG}$	42	10	24%
$R2^{FCPG}$	87	22	25%
$R2^{KD}$	45	0	-
Recovery of *Fgfr2* hemizygous signaling mutants			
P4-P10	Total	*/-	Recovery
$R2^F$	37	0	-
$R2^C$	52	10	19%
$R2^{PG}$	47	13	27%
$R2^{CPG}$	53	12	22%
$R2^{FCPG}$	28	1	3%
P0-P1	Total	Observed	Expected
$R2^F$	33	4	12%
$R2^{FCPG}$	25	2	8%
E18.5	Total	Observed	Expected
$R2^F$	33	5	15%
$R2^{FCPG}$	31	10	32%

Fgfr2 signaling mutant heterozygotes ($Fgfr2^{F/+}$, $Fgfr2^{C/+}$, $Fgfr2^{PG/+}$, $Fgfr2^{CPG/+}$, $Fgfr2^{FCPG/+}$ and

$Fgfr2^{KD/+}$) were intercrossed. Embryos were analyzed either at E18.5 or postnatally between P4

and P10. To identify *in vivo* phenotypes associated with *Fgfr2* signaling mutations, *Fgfr2* signaling

mutant heterozygotes were crossed with $Fgfr2^{+/-}$ null heterozygotes. $Fgfr2^{C/-}$, $Fgfr2^{PG/-}$ and

$Fgfr2^{CPG/-}$ embryos were analyzed at E18.5 and postnatally at P0 and between P4 and P10.

Hemizygous $Fgfr2^{F/-}$ and $Fgfr2^{FCPG/-}$ mutant mice were recovered at expected Mendelian ratios at

E18.5, but died postnatally.

Supplementary Information Table 2: Craniofacial structures differentially affected in
conditional signaling mutants.

C/CPG<F<FCPG	*C/CPG<F=FCPG*
Nasal Cartilage	
Mandible	
Squamosal	
Maxilla	
	Premaxilla
	Frontal bone
	Zygomatic bone
	Tympanic bulla
	Basisphenoid
	Internal pterygoid process

713 Micro-CT examination of *Fgfr1$^{F/cKO}$; Fgfr2$^{cKO/cKO}$* and *Fgfr1$^{FCPG/cKO}$; Fgfr2$^{cKO/cKO}$* mutant mice skull
714 were to *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$* mutants at E18.5. *Fgfr1$^{FCPG/cKO}$; Fgfr2$^{cKO/cKO}$* mutant skull show a
715 more severe defect than *Fgfr1$^{F/cKO}$; Fgfr2$^{cKO/cKO}$* mutants. Several structures are differentially
716 affected in *Fgfr1$^{F/cKO}$; Fgfr2$^{cKO/cKO}$* and *Fgfr1$^{FCPG/cKO}$; Fgfr2$^{cKO/cKO}$* mutants. Most severe defects are
717 observed in *Fgfr1$^{cKO/cKO}$; Fgfr2$^{cKO/cKO}$* mutants.

718
719

720 **Supplementary Information Table S3: Homology arms used in targeting vectors.**

	Short homology arm (corresponding primer binding region)	Notes	Long homology arm (corresponding primer binding region)	Notes
Fgfr2^F	GCTCTTTCTCTCTATGGGTT and ATTCGACTGTAATGGGGGAC	1.7kb region between exon 9-10	TGGTATCAGCTGCTCACTGG and GCCCACCCTGAATAACCATG	5.1kb spanning exon10
Fgfr2^C	CTTTCTGCATCAAAGACACA and GGTTACATCGATTGCCCAGT	1.7kb region between exon 11-12	GGCCACAATTCAGAGGGAAG and AGCCCATTTTTCCACCTTCT	5.1kb spanning exon11
Fgfr2^{PG}	TCTTGGCTGTGGCTTGTACTGG and AGGACCTCGGTGACCCACTA	1.9kb region between exon 18-19	GTAGCAGAGTGGGCAAGCTC and AGGAACTGCAAGAGGACCAA	5.4kb spanning exon19 and 3'-UTR
Fgfr2^{KD}	GGTACCCAAGTGTGGAAAAT and GTTTGCTCCTTTTTGGCTTC	SmaI to MfeI, 1.9kb spanning exon12	GAACTGATGCAGAATCCCAG and CCATTVTAAGTCCCTTGAGC	MfeI to BclI 3.7kb spanning exon13

722 **Supplementary Information Table S4: Nucleotide and amino-acid substitutions in signaling**

723 **mutations for *Fgfr2*.**

Mutation	Nucleotide substitution	Amino acid substitution	Notes
F	CTG to GCC AGA to GCA	L424A R426A	Introduces an XmaI site
C	TAT to TTT TTG to CTC	Y466F L468L	Introduces a SacI site
PG	TAC to TTC TAT to TTT	Y769F Y779F	Introduces an EcoRI site
KD	GAAG to AGCT	K517A	Introduces an AluI site

 Supplementary Information Table S5: List of oligonucleotides used in this study.

Gene	Forward	Purpose
Fgfr2^F 5' probe F	ACACTGGAGGATGGTGAAGG	Probes for Southern blot
Fgfr2^F 5' probe R	GACATGCATGCAATTGAACA	Probes for Southern blot
Fgfr2^F 3' probe F	GTCCCAGCTCAGCAATAAGC	Probes for Southern blot
Fgfr2^F 3' probe R	AGTGGTCACCTTCGGAGAGA	Probes for Southern blot
Fgfr2^C 5' probe F	GTGGGACAGCCTGGTAGGTA	Probes for Southern blot
Fgfr2^C 5' probe R	CGGGTAGGCTATGCACTCAT	Probes for Southern blot
Fgfr2^C 3' probe F	ATACTTGACCTGGCGGTTGA	Probes for Southern blot
Fgfr2^C 3' probe R	GGCATTTATAGCGTGGCATT	Probes for Southern blot
Fgfr2^{PG} 5' probe F	GTGTGAATGCTGCTTGTGCT	Probes for Southern blot
Fgfr2^{PG} 5' probe R	GCGGAAGCCTTGTTTTATGA	Probes for Southern blot
Fgfr2^{PG} 3' probe F	GGCCACGGAGTAGTTCAGAG	Probes for Southern blot
Fgfr2^{PG} 3' probe R	GCAAGCACATTTGTTTACCTG	Probes for Southern blot
Fgfr2^{KD} 5' probe F	TGGGTATGGCCAGCTCTAGT	Probes for Southern blot
Fgfr2^{KD} 5' probe R	TGCACACGAACCCAGCTATT	Probes for Southern blot
Fgfr2^{KD} 3' probe F	ATCCGTGCCTTTCTGAGTGG	Probes for Southern blot
Fgfr2^{KD} 3' probe R	TGCCAGCCATTTACCTCAGG	Probes for Southern blot
*Fgfr2^C*Forward-Genotyping	GTAAGTACTCTCCCTCTGGG	Genotyping
*Fgfr2^C*Reverse-Genotyping	AGCTGGCCATACCCAGACTTG	Genotyping
*Fgfr2^{PG}*Forward-Genotyping	CTGTTGCTTACTCTGGAGCT	Genotyping
*Fgfr2^{PG}*Reverse-Genotyping	GATACCCCAACTCCATTCAC	Genotyping
*Fgfr2^F*Forward-Genotyping	GGACCATATGAAGGGACTTG	Genotyping
*Fgfr2^F*Reverse-Genotyping	CTCGGTTTATCACTACAGCC	Genotyping
*Fgfr2^{KD}*Forward-Genotyping	ATAGCCTGCGTGTGTATCGG	Genotyping*
*Fgfr2^{KD}*Reverse-Genotyping	CGTCCCTGCAGAGTTCACAT	Genotyping*
Alx4-forward	ACACATGGGCAGCCTGTTTG	RT-PCR
Alx4-reverse	TGCTTGAGGTCTTGCGGTCT	RT-PCR
Gapdh-forward	AGGTCGGTGTGAACGGATTTG	RT-PCR
Gapdh-reverse	TGTAGACCATGTAGTTGAGGTCA	RT-PCR
Dlx5-forward	CAGAAGAGTCCCAAGCATCC	RT-PCR
Dlx5-reverse	GGTGACTGTGGCGAGTTA	RT-PCR

Lhx8-forward	ATGTATTGGAAGAGCGATCAG	RT-PCR
Lhx8-reverse	TCATTGGATGGGGTAACAAGGGC	RT-PCR
FoxD1-forward	TGAGCACTGAGATGTCCGATG	RT-PCR
FoxD1-reverse	CACCACGTCGATGTCTGTTTC	RT-PCR
Twist2-forward	AGCGCCCAGAGCTTCGAGGA	RT-PCR
Twist2-reverse	CGGCGAAGGCCTCGTTCAGG	RT-PCR
Msx1-forward	TCTCGGCCATTTCTCAGTCG	RT-PCR
Msx1-reverse	AGAGCATCTTCTGGCAGCTTG	RT-PCR
Sox9-forward	AGGAAGTCGGTGAAGAACGG	RT-PCR
Sox9-reverse	TGGAAGGTGAATTTCTCTGGG	RT-PCR
Dlx1-forward	TGGAATCCGAACTCCTCATC	RT-PCR
Dlx1-reverse	TGCTGCATAGCTTCTTGGTG	RT-PCR
Fgfr1-forward	CACATCGAGGTGAACGGGAGTAAG	RT-PCR
Fgfr1-reverse	CGCATCCTCAAAGGAGACATTCC	RT-PCR
Fgfr2-forward	GATGTGGAGTTTGTCTGCAAGGTTT	RT-PCR
Fgfr2-reverse	GACTGGTTGGCCTGCCCTATATAAT	RT-PCR
Fgfr3-forward	TGCGGTGCCTTCACAGA	RT-PCR
Fgfr3-reverse	ACTTGGACCTCTCCGTG	RT-PCR
Fgfr4-forward	GTACCCTCGGACCGCGGCACATAC	RT-PCR
Fgfr4-reverse	GCCGAAGCTGCTGCCGTTGATG	RT-PCR
Fgfr1-Ex6 -Crispr	CACCGCTGTTAGCCACACAAAGCAC	CRISPR Fgfr1
Fgfr1-Ex6 -Crispr	AAACGTGCTTTGTGTGGCTAACAGc	CRISPR Fgfr1
Fgfr1-Ex6 -Crispr	CACCGAAGAAGGAAGAAGGGGGAGG	CRISPR Fgfr1
Fgfr1-Ex6 -Crispr	AAACCCTCCCCCTTCTTCCTTCTTc	CRISPR Fgfr1
Fgfr2-Ex5 -Crispr	CACCGCTTCCTGGGCATTTGAACCC	CRISPR Fgfr2
Fgfr2-Ex5 -Crispr	AAACGGGTTCAAATGCCCAGGAAGc	CRISPR Fgfr2
Fgfr2-Ex5 -Crispr	CACCGCCATAAGATGTCATCTGAGA	CRISPR Fgfr2
Fgfr2-Ex5 -Crispr	AAACTCTCAGATGACATCTTATGGc	CRISPR Fgfr2
Fgfr1-CRISPR-Screen	TGCAGGAAGGATTCTGAGTGT	Screening Fgfr1 CRISPR clones
Fgfr1-CRISPR-Screen	GCTGTCTGGGCTAAAGTCTG	Screening Fgfr1 CRISPR clones
Fgfr2-CRISPR-Screen	CCCAAGCCCATAAAACCTGG	Screening Fgfr2 CRISPR clones

| Fgfr2-CRISPR-Screen | CAAGGACACACGTTCACAGG | Screening Fgfr2 CRISPR clones |

727 * Note: Genotyping *Fgfr2*^{KD} is followed by AluI digestion after PCR (Supplementary Information

728 Table S6)

730 **Supplementary Information Table S6: Genotyping reactions to identify *Fgfr2* signaling**
731 **mutations.**

	WT	Mutant
Fgfr2^C	553 bp	655 bp
Fgfr2^PG	377 bp	611 bp
Fgfr2^F	379 bp	614 bp
Fgfr2^KD	350 bp After AluI digest	300 bp After AluI digest

733
734

735 **Supplementary information Table S7**

	Source	Identifier
Antibodies		
FGFR2	Abcam	#ab109372
CRKL	Santa Cruz Biotechnology	#sc-319
FRS2	Santa Cruz Biotechnology	#sc-8318
FLAG2-M2	Sigma	#F1804
phospho-p44/42 MAPK Thr202/Tyr204	Cell Signaling Technology	#9101
p44/42 MAPK	Cell Signaling Technology	#9102
GAPDH	ProteinTech	#60004-1-Ig
phospho-AKT Ser473	Cell Signaling Technology	#4060
AKT	Cell Signaling Technology	#9272
phospho-p38	Cell signaling Technology	#4511
p38	Cell signaling Technology	#9212
phospho-PLCγ1 (Y783)	Cell Signaling Technology	#2821
PLCγ1	Cell Signaling Technology	#2822
phospho-JNK	Cell Signaling Technology	#4671
STAT3α	Cell signaling Technology	#8768
phospho-STAT3 (Y705)	Cell Signaling Technology	#9145
phospho-FAK	Cell Signaling Technology	#3283
FAK	Cell Signaling Technology	#3285
Paxillin	Abcam	#ab32084
IRS2	Cell Signaling Technology	#3089
β-catenin	Cell Signaling Technology	#8480
Neurofilament antibody	DSHB	#Clone 2H3
EOMES	Abcam	#ab23345
CDX2	Biogenex	#MU392A-UC
GFP	AvesLab	#GFP-1020
SMA	Cell Signaling Technology	#19245
Peroxidase AffiniPure Goat Anti-Mouse IgG (H+L)	Jackson ImmunoResearch Laboratories	#115-035-003

Peroxidase AffiniPure Goat Anti-Rabbit IgG (H+L)	Jackson ImmunoResearch Laboratories	#111-035-003
Donkey anti-Rabbit IgG (H+L), Alexa Fluor 488	ThermoFisher Scientific	#A-21206
Donkey anti-Rabbit IgG (H+L), Alexa Fluor 546	ThermoFisher Scientific	#A-10040
Donkey anti-Rabbit IgG (H+L), Alexa Fluor 647	ThermoFisher Scientific	#A-31573
Donkey anti-Mouse IgG (H+L), Alexa Fluor 488	ThermoFisher Scientific	#A-21202
Donkey anti- Mouse IgG (H+L), Alexa Fluor 546	ThermoFisher Scientific	#A-10036
Donkey anti- Mouse IgG (H+L), Alexa Fluor 647	ThermoFisher Scientific	#A-31571
Alexa Fluor® 488 AffiniPure Donkey Anti-Chicken IgY (IgG) (H+L)	Jackson ImmunoResearch Laboratories	#703-545-155

  METHODS

Generation of knock-in mice. Four distinct targeting vectors carrying the $Fgfr2^F$, $Fgfr2^C$, $Fgfr2^{PG}$,

and $Fgfr2^{KD}$ mutations were generated. The $Fgfr2^F$ targeting vector was generated by cloning a

short homology arm (1.7kb region between exon 9-10) and a long homology arm (5.1kb, spanning

exon10) into PGKneolox2DTA.2 [48]. To allow recombineering into SW105 bacteria, the neo

cassette was subsequently replaced by PGKEm7neo flanked by FRT sites, which contains both

a eukaryotic and a prokaryotic promoter. Similarly, for the $Fgfr2^C$ targeting vector, we cloned a

long homology arm (5.3kb region spanning exon11) and a short homology arm (1.7kb region

between exon 11-12) into PGKneolox2DTA.2 and used a PGKEm7neo flanked by both FRT and

LoxP sites for recombineering. For the $Fgfr2^{PG}$ targeting vector, we cloned a short homology arm

(1.9kb region 5' of exon19) and a long homology arm (5.4kb spanning exon19 and 3'-UTR) into

PGKneolox2DTA.2 and used a PGKEm7neo flanked by FRT sites for recombineering. For $Fgfr2^{KD}$

targeting vector, we cloned short homology arm (1.9kb SmaI to MfeI, spanning exon 12) and a

3.7kb long homology arm (MfeI to BclI, spanning exon 13) into PGKneolox2DTA.2 [48]. Details of

regions corresponding to homology arms are provided in Supplementary Information Table S3.

For all four alleles, site-directed mutagenesis (SDM) was performed using Phusion

polymerase (NEB#M0530). Nucleotide substitutions introduced by SDM in exon10 for $Fgfr2^F$

allele (introduces an XmaI site), exon11 for $Fgfr2^C$ allele (introduces a SacI site), exon 19 for

$Fgfr2^{PG}$ allele (introduces an EcoRI site) and in exon12 for $Fgfr2^{KD}$ allele (introduces an AluI site)

are provided below. All introduced mutations were verified by sequencing. Details of nucleotide

substitutions in $Fgfr2$ signaling mutations are provided in Supplementary Information Table S4.

The targeting vectors for $Fgfr2^F$ (linearized with NotI), $Fgfr2^C$ (linearized with XhoI), $Fgfr2^{PG}$

(linearized with NotI) and $Fgfr2^{KD}$ (linearized with NotI) were electroporated into 129S4 AK7 ES

cells. For generating the allelic series of signaling mutations, ES cells were targeted first with the

C targeting vector generating $Fgfr2^{+/C}$ mutant cells. After verifying for correct targeting events, the

neo cassette was removed by transient transfection with PGKCrebpA, leaving a single LoxP site

behind (Figure 3C). $Fgfr2^{+/C}$ ES cells were then targeted using the PG targeting vector generating

either $Fgfr2^{+/PG}$ or $Fgfr2^{+/CPG}$ mutant cells, as determined by breeding of the chimeras to

$ROSA26^{Flpo}$ mice [49]. After verifying for correct targeting events, the neo cassette was removed by

transient transfection with PGKFlpobpA [50], leaving both an FRT site and a LoxP site behind

(Figure 3C). $Fgfr2^{+/CPG}$ neo⁻ ES cells were finally targeted with the F targeting vector, resulting in

$Fgfr2^{+/F}$ or $Fgfr2^{+/FCPG}$ mutant ES cells, as determined by breeding. After verifying for correct

targeting events, the neo cassette was removed by transient transfection with Flpe, which is less

efficient than Flpo in ES cells [50], in order to not recombine sequences between exons 10-18 due

to the retention of the FRT site during the generation of the *Fgfr2^PG* allele. We screened targeting

events initially by PCR coupled with restriction digestion to identify incorporation of nucleotide

substitutions. Proper targeting was confirmed by Southern blotting using 5′ external and 3′

external probes amplified using the following primer pairs and then an internal probe against Neo.

Primers used to generate probes for confirming targeted clones using Southern blots are

described in Supplementary Information Table S5.

ES cell chimeras were bred to *Meox2-Cre* or *ROSA26Flpo* deleter mice [49,51] maintained

on a 129S4 genetic background to remove the neomycin selection cassette and the deleter alleles

were subsequently crossed out. Two independent mouse lines were generated from independent

ES cell clones for each allele, and phenotypes were confirmed in both lines. The *Fgfr2^C, Fgfr2^PG,*

Fgfr2^F, Fgfr2^FCPG and *Fgfr2^KD* alleles were maintained on the 129S4 genetic background. *Fgfr2^C,*

Fgfr2^PG, Fgfr2^F and *Fgfr2^KD* mice were genotyped using oligonucleotides listed in Supplementary

Information Table S5 and S6, with the F, C or PG primers all being able to genotype *Fgfr2^FCPG*

mice.

Mouse strains. All animal experimentation was conducted according to protocols approved by

the Institutional Animal Care and Use Committee of the Icahn School of Medicine at Mount Sinai.

Fgfr1^cKO/cKO, Fgfr2^cKO/cKO, Fgfr1-GFP and *Fgfr2-mCherry* were previously described [18,48]. *Fgfr1*

signaling mutations [7] are referred to as *Fgfr1^C, Fgfr1^F, Fgfr1^CPG* and *Fgfr1^FCPG*. *Tg(Wnt1-cre)11Rth,*

Tg(Wnt1-cre)2Sor, Cdkn2a^+/tm1Rdp, Gt(ROSA)26Sor^tm4(ACTB-tdTomato,-EGFP)Luo, and *Bcl2l11^tm1.1Ast* are

referred to in the text as *Wnt1-Cre, Wnt1-Cre2, Ink, ROSA26^mT/mG ,* and *Bim* respectively [8,9,52-54].

All lines were maintained on a 129S4 co-isogenic background, except for *Bim* which was crossed

into the *Fgfr1/2* deficient backgrounds after only six generations of backcrossing to 129S4.

***Generation of Fgfr2^-Flag3x expression vector* and stable 3T3 expression lines.** An *Fgfr2*

isoform "c" cDNA isoform was PCR amplified from primary MEFs derived from *Fgfr2^+/+, Fgfr2^PG/PG,*

Fgfr2^CPG/CPG, Fgfr2^F/F and *Fgfr2^FCPGFCPG* and subsequently digested with HindIII and XhoI. The

fragments were cloned in the pcDNA expression vector and sequence verified. Linearized

pcDNA-FGFR2 plasmids were transfected in 3T3 cells cultured in DMEM supplemented with 10%

calf serum (HyClone Laboratories #SH30072.03) with 50U/mL each penicillin and streptomycin

(Gibco #P0781). Stable clones were selected in 500µg/mL G418 (Fischer Scientific #BP6735).

10 clones from each construct (FGFR2^wt-, FGFR2^PG-, FGFR2^CPG-, FGFR2^F-, or FGFR2^FCPG-

Flag3x) were expanded and assessed for FLAG expression by western blot. Clones expressing

high FGFR2-FLAG levels were selected for further analysis.

Coimmunoprecipitation and Western blotting. Stable 3T3 cells expressing FGFR2^{WT},

FGFR2^{PG}, FGFR2^{CPG}, FGFR2^{F}, or FGFR2^{FCPG} -Flag3x were serum-starved (0.1% calf serum

supplemented DMEM) overnight, stimulated for 15 mins with 50 ng/mL FGF1 (PeproTech, 450-

33A) or FGF8b (PeproTech 100-25B) and 5 μg/mL heparin (Sigma #H3149), and lysed in ice-

cold NP-40 lysis buffer (20 mM Tris HCL at pH 8, 137 mM NaCl, 10% glycerol, 1% Nonidet [NP-

40], 2 mM EDTA, 25 mM β glycerol phosphate, 1 mM Na$_3$VO$_4$, 10 mM NaF, 1× cOmplete, EDTA-

free Protease Inhibitor Cocktail. 800 μg cell lysates were subsequently used for

immunoprecipitation with Anti-FLAG M2 magnetic beads (Sigma #M8823) using the

manufacturer's protocol. We incubated lysates with anti-FLAG M2 magnetic beads overnight at

4°C followed by five washes with lysis buffer, and precipitated proteins were eluted in Laemmli

buffer (10% glycerol, 2% SDS, 0.002% bromophenol blue, 0.062M Tris-HCl, pH 6.8) containing

10% β-mercaptoethanol, heated for 5 min at 95°C, separated by SDS-PAGE and analyzed by

western blots.

Western blot analysis was performed according to standard protocols using horseradish

peroxidase-conjugated secondary antibodies (1:10,000 dilution) developed by chemiluminescent

HRP substrate. Primary antibodies were used at the following dilutions for Western blotting:

FGFR2 (1:500 dilution), CRKL (1:500 dilution), FRS2 (1:500 dilution), Flag2 M2 (1:500 dilution),

phospho-p44/42 MAPK (1:1,000 dilution), p44/42 MAPK (1:1,000 dilution), GAPDH (1:1000

dilution), phospho-AKT (1:1,000 dilution), AKT (1:1,000 dilution), phospho-p38 (1:500 dilution),

p38 (1:500 dilution), phospho-PLCγ1 (Y783) (1:200 dilution), PLCγ1 (1:1000 dilution), pJNK

(1:500 dilution), phospho-STAT3 (Y705) (1:500 dilution), STAT3α (1:1000 dilution), phospho-FAK

(1:1000 dilution), FAK (1:1000 dilution), β-catenin (1:1000 dilution), Paxillin (1:1000 dilution), and

IRS2 (Cell Signaling Technology, 3089). For signaling pathways analyzed, the blots were

quantified for three independent biological replicates using ImageLab6.0 analysis tool. Ratio of

average of mean intensity value of pERK1/2, pAKT, pPLCγ1, pJNK, pSTAT3α, pP38 to GAPDH

± standard deviation was plotted for each pathway.

Cell derivation and culture conditions. Primary iFNPs were generated by dissecting the

maxillary and nasal prominences of E11.5 *Fgfr2$^{+/+}$; Ink$^{-/-}$*, *Fgfr2$^{F/F}$; Ink$^{-/-}$*, *Fgfr2$^{CPG/CPG}$; Ink$^{-/-}$,* and

E9.5 or E11.5 *Fgfr2$^{FCPG/FCPG}$; Ink$^{-/-}$* embryos in PBS. The tissue was disassociated with 0.125%

Trypsin-EDTA and cultured in DMEM supplemented with 20% FBS, 50 U/mL each penicillin and

streptomycin on fibronectin coated plates (0.5μg/cm^2). Cells were subsequently split 1:5 through

for at least 5 passages before immortalized cell lines were obtained. Cells were allowed to grow

until sub-confluent. All experiments were performed between passage 15 and 25. We used PX459

V2.0 vector (Addgene plasmid # 62988) to CRISPR out either *Fgfr1* or *Fgfr2* and create *Fgfr1* null,

Fgfr2 null, or *Fgfr1*: *Fgfr2* double null cells. gRNA sequences for *Fgfr1* and *Fgfr2* were selected

using CHOPCHOP gRNA design web tool and were cloned using the oligonucleotides

(Supplementary Information Table S5), as previously described [55]. Plasmids were transfected in

respective iFNP cells cultured in DMEM (Sigma #5796) supplemented with 10% calf serum with

50 U/mL each penicillin and streptomycin. Stable clones were selected in 5μg/mL Puromycin

(Sigma #P8833). Clones were verified (homozygous deletion of exon6 for *Fgfr1* and deletion

exon5 for *Fgfr2* which also introduces a frameshift mutation) using PCR (Supplementary

Information Table S5). Primary MEFs were derived from E12.5 wild type mice embryos. Embryos

were eviscerated and after removing the head, remaining tissue was chopped into 1 cm pieces

and incubated in (0.25%) 1 mL trypsin-EDTA (ThermoFisher # 25200-072) for 30 mins with

intermittent shaking. 10 mL DMEM 10% calf serum was added and the mixture was allowed to

pass through a cell-strainer. Cells collected from each embryo was plated in 0.2% gelatin

(Sigma #G1393) coated 15cm plate.

Skeletal preparations. Embryos at E14.5, E16.5 or E18.5 embryos were skinned, eviscerated,

fixed in 95% ethanol overnight, and stained (0.015% Alcian blue, 0.005% Alizarin red, 5% glacial

acetic acid, in 70% ethanol) overnight at 37°C. Skeletons were then cleared in 1% KOH and

transferred to decreasing concentrations of KOH in increasing concentrations of glycerol until

clear.

Acetocarmine and hematoxylin and eosin staining. Freshly harvested tissue was fixed in 4%

PFA at 4°C overnight followed by dehydration in 70% ethanol. For acetocarmine staining, tissues

were incubated in 0.5% aceto-carmine (0.5 g carmine stain (Sigma #C1022) dissolved in 100 ml

boiling 45% acetic acid for 15 minutes), followed by de-staining in 70% ethanol for 1 minute and

1% acid alcohol (1% HCl in 70% ethanol) for 2 minutes and 5% acid alcohol (5% HCl in 70%

ethanol) for 1 minute. For hematoxylin and eosin staining, freshly harvested tissues were

dissected in PBS, and fixed in 4% PFA followed by dehydration through a graded ethanol series,

and embedded in paraffin. 5μm sections were cut. After deparaffinization and rehydration,

sections were stained with Harris modified hematoxylin (Sigma #HHS16) followed by a 10 second

wash in acid-alcohol (1% v/v HCl in 70% EtOH), followed by counterstaining with 1% eosinY

(Sigma #17372-87-1). Tissues were washed and mounted with Permount (ThermoFisher

Scientific #SP15100).

Scratch assays. Cells were seeded onto glass coverslips coated with 5 µg/mL human plasma

fibronectin purified protein (Millipore Sigma #FC010). At ~90–100% confluency, cells were

scratched with a P1000 pipet tip, washed with PBS and incubated in fresh medium containing

either 0.1% FBS, 10% FBS, 50 ng/mL FGF1 and 5 µg/mL heparin or 10 ng/mL PDGF-AA (R&D

Systems #1055-AA-050) supplemented DMEM for 12 hrs.

Immunofluorescence and Antibodies. For immunostaining whole mount embryos were fixed in

4% paraformaldehyde solution (PFA) in PBS overnight and washed with PBS five times,

permeabilized with 0.5% Triton X100 in PBS for 30 min and blocked in 2% blocking serum for 2h

at room temperature. Primary anti-neurofilament antibody (Clone 2H3, DSHB) was used at a 1:20

dilution in 1% donkey serum in PBST; embryos were incubated overnight at 4°C. The next day,

embryos were washed 4 times in PBST and incubated with anti-mouse HRP-conjugated

secondary antibodies at a 1:1000 dilution for 4hrs at room temperature followed by washing in

PBST 4 times and signal was developed using ImmPACTDAB kit (VectorLabs #SK-4105). For

whole-mount immunofluorescence at E7.5, embryos were fixed overnight in 4:1 methanol:DMSO.

Primary antibodies for Eomes (1:100 dilution) and Cdx2 (1:100 dilution) were used. For

immunostaining cells, cells were fixed for 10 mins in 4% PFA in PBS at room temperature. Cells/

tissues were subsequently processed for immunofluorescence analysis as detailed above using

anti-paxillin primary antibody (1:250 dilution) with Alexa647 conjugated phalloidin (1:40 dilution).

For immunofluorescence on sections, antibodies for GFP (1:100 dilution), mCherry (1:100 dilution)

and SMA (1:100 dilution) was used. Supplementary Information Table S7 provides more

information on antibodies used in this study. Embryos were stained with DAPI following fixation

as previously described [56]. Cells / tissues were photographed using an Olympus DP71 digital

camera fitted onto an Olympus BX51 fluorescence microscope, Leica SP5 confocal or a

Hamamatsu C11440 camera fitted to a Zeiss Observer Z1 microscope. Epifluorescence was

imaged in Zeiss Axioplan fitted to ProgRes CT3 camera.

In situ hybridization. Labeled antisense-RNA probes were synthesized for *Alx3, Msx1, Six3,*

Nkx2.1, Fgf8, Shh, Col2a1, Col10a1 and *Meox1*. Digoxigenin-labeled anti-sense probes were

generated as described, and mRNA in situ hybridization on paraffin sections for chromogenic

detection was performed using standard protocols.

Micro-CT imaging. Micro-CT imaging of the skulls were performed using a SkyScan 1172

scanner (Bruker, Kontich, Belgium). The mouse heads were dissected and fixed in 10% neutral

buffered formalin and washed and stored in PBS at 4 °C. The skull bones were scanned with

settings of 50 kV, 500 µA, 10 µm pixel resolution, 0.3° rotation steps, and 4 frames average
imaging with a 0.5-mm Al filter at Micro-CT Core, School of Dentistry, NYU, New York. The
acquired X-ray projections were reconstructed using the Imaris software (Oxford Instruments).

Cell proliferation assay. For EdU labeling in mice, pregnant females were injected
intraperitoneally with 100 mg/kg body weight of EdU. EdU detection was carried out as per
manufacturer's instruction for Click-iT EdU Cell Proliferation Kit.

TUNEL assay. Sections were deparaffinized and were rehydrated in PBS, followed by post-
fixation in 4% PFA. In Situ Cell Death Detection Kit, TMR red user protocol was used to detect
cell death.

RT-qPCR. Cells were lysed, and mRNA was extracted according to Qiagen RNeasy kit standard
protocol. cDNA was synthesized using a 2:1 ratio of random primers to Oligo(dT) with SuperScript
IV RT (Invitrogen). qPCR was performed with PerfeCTa SYBR Green FastMix for iQ (Quanta
Biosciences) with Bio-Rad iQ5 multicolor real-time PCR detection system and analyzed with Bio-
Rad iQ5 optical system software (version 2.0). Cycling conditions were as follows: step 1, 3 min
at 95°C; step 2, 10 sec at 95°C; step 3, 30 sec at 60°C; and repeat steps 2 and 3 for 40 cycles.
Proper amplification was confirmed using a melting curve and by running samples on a gel to
ensure that the correct size band was obtained. Graphs were made using Microsoft Excel and
Prism. Primer sequence for respective genes used for RT-qPCR analysis is listed below.

β-Catenin Quantification. Cells were stained and imaged using a Leica SP5 confocal
microscope under identical conditions. Stacks were then background subtracted using a 100px
rolling ball function in ImageJ. The average pixel intensity along cell-cell junctions was measured
in a single z-plane per junction using a 15-pixel line width in ImageJ for the EGFP and β-catenin
channels.

Quantification and statistical analysis. Statistical analysis was performed using GraphPad
Prism6.0 and Microsoft Excel. Values are presented as mean ± standard deviation. The statistical
significance was determined using a student *t*-test with Holm-Sidak method.

References

1   Lemmon, M. A. & Schlessinger, J. Cell signaling by receptor tyrosine kinases. *Cell* **141**,
1117-1134, doi:10.1016/j.cell.2010.06.011 (2010).
2   Zinkle, A. & Mohammadi, M. A threshold model for receptor tyrosine kinase signaling
specificity and cell fate determination. *F1000Res* **7**, doi:10.12688/f1000research.14143.1
(2018).
3   Li, P. & Elowitz, M. B. Communication codes in developmental signaling pathways.
Development **146**, doi:10.1242/dev.170977 (2019).
4   Vasudevan, H. N., Mazot, P., He, F. & Soriano, P. Receptor tyrosine kinases modulate
distinct transcriptional programs by differential usage of intracellular pathways. *Elife* **4**,
doi:10.7554/eLife.07186 (2015).
5   Brewer, J. R., Mazot, P. & Soriano, P. Genetic insights into the mechanisms of Fgf signaling.
Genes Dev **30**, 751-771, doi:10.1101/gad.277137.115 (2016).
6   Lanner, F. & Rossant, J. The role of FGF/Erk signaling in pluripotent cells. *Development*
137, 3351-3360 (2010).
7   Brewer, J. R., Molotkov, A., Mazot, P., Hoch, R. V. & Soriano, P. Fgfr1 regulates
development through the combinatorial use of signaling proteins. *Genes Dev* **29**, 1863-
1874, doi:10.1101/gad.264994.115 (2015).
8   Danielian, P. S., Muccino, D., Rowitch, D. H., Michael, S. K. & McMahon, A. P. Modification
of gene activity in mouse embryos in utero by a tamoxifen-inducible form of Cre
recombinase. *Curr Biol* **8**, 1323-1326, doi:10.1016/s0960-9822(07)00562-3 (1998).
9   Lewis, A. E., Vasudevan, H. N., O'Neill, A. K., Soriano, P. & Bush, J. O. The widely used
Wnt1-Cre transgene causes developmental phenotypes by ectopic activation of Wnt
signaling. *Dev Biol* **379**, 229-234, doi:10.1016/j.ydbio.2013.04.026 (2013).
10   Partanen, J., Schwartz, L. & Rossant, J. Opposite phenotypes of hypomorphic and Y766
phosphorylation site mutations reveal a function for Fgfr1 in anteroposterior patterning
of mouse embryos. *Genes Dev* **12**, 2332-2344 (1998).
11   De Moerlooze, L. *et al.* An important role for the IIIb isoform of fibroblast growth factor
receptor 2 (FGFR2) in mesenchymal-epithelial signalling during mouse organogenesis.
Development **127**, 483-492 (2000).
12   Youle, R. J. & Strasser, A. The BCL-2 protein family: opposing activities that mediate cell
death. *Nat Rev Mol Cell Biol* **9**, 47-59, doi:10.1038/nrm2308 (2008).
13   Chipuk, J. E. & Green, D. R. How do BCL-2 proteins induce mitochondrial outer membrane
permeabilization? *Trends Cell Biol* **18**, 157-164, doi:10.1016/j.tcb.2008.01.007 (2008).
14   Czabotar, P. E., Lessene, G., Strasser, A. & Adams, J. M. Control of apoptosis by the BCL-2
protein family: implications for physiology and therapy. *Nat Rev Mol Cell Biol* **15**, 49-63,
doi:10.1038/nrm3722 (2014).
15   Grabow, S. *et al.* Subtle Changes in the Levels of BCL-2 Proteins Cause Severe Craniofacial
Abnormalities. *Cell Rep* **24**, 3285-3295 e3284, doi:10.1016/j.celrep.2018.08.048 (2018).
16   Clybouw, C. *et al.* Alternative splicing of Bim and Erk-mediated Bim(EL) phosphorylation
are dispensable for hematopoietic homeostasis in vivo. *Cell Death Differ* **19**, 1060-1068,
doi:10.1038/cdd.2011.198 (2012).

17 Lei, K. & Davis, R. J. JNK phosphorylation of Bim-related members of the Bcl2 family
induces Bax-dependent apoptosis. *Proc Natl Acad Sci U S A* **100**, 2432-2437,
doi:10.1073/pnas.0438011100 (2003).

18 Molotkov, A., Mazot, P., Brewer, J. R., Cinalli, R. M. & Soriano, P. Distinct Requirements
for Fgfr1 and Fgfr2 in Primitive Endoderm Development and Exit from Pluripotency.
Developmental Cell **41**, 511-526 (2017).

19 Yu, K. *et al.* Conditional inactivation of FGF receptor 2 reveals an essential role for FGF
signaling in the regulation of osteoblast function and bone growth. *Development* **130**,
3063-3074 (2003).

20 Xu, X. *et al.* Fibroblast growth factor receptor 2 (FGFR2)-mediated reciprocal regulation
loop between FGF8 and FGF10 is essential for limb induction. *Development* **125**, 753-765
(1998).

21 Garg, A. *et al.* Alx4 relays sequential FGF signaling to induce lacrimal gland morphogenesis.
PLoS Genet **13**, e1007047, doi:10.1371/journal.pgen.1007047 (2017).

22 Steinberg, Z. *et al.* FGFR2b signaling regulates ex vivo submandibular gland epithelial cell
proliferation and branching morphogenesis. *Development* **132**, 1223-1234,
doi:10.1242/dev.01690 (2005).

23 Francavilla, C. *et al.* Functional proteomics defines the molecular switch underlying FGF
receptor trafficking and cellular outputs. *Mol Cell* **51**, 707-722,
doi:10.1016/j.molcel.2013.08.002 (2013).

24 Kurowski, A., Molotkov, A. & Soriano, P. FGFR1 regulates trophectoderm development
and facilitates blastocyst implantation. *Dev Biol* **446**, 94-101,
doi:10.1016/j.ydbio.2018.12.008 (2019).

25 Fantauzzo, K. A. & Soriano , P. Generation of an Immortalized Mouse Embryonic Palatal
Mesenchyme Cell Line. *PLoS ONE* **12**, e0179078 (2017).

26 Hanks, S. K., Quinn, A. M. & Hunter, T. The protein kinase family: conserved features and
deduced phylogeny of the catalytic domains. *Science* **241**, 42-52,
doi:10.1126/science.3291115 (1988).

27 Bellot, F. *et al.* Ligand-induced transphosphorylation between different FGF receptors.
EMBO J **10**, 2849-2854 (1991).

28 Ueno, H., Gunn, M., Dell, K., Tseng, A., Jr. & Williams, L. A truncated form of fibroblast
growth factor receptor 1 inhibits signal transduction by multiple types of fibroblast
growth factor receptor. *J Biol Chem* **267**, 1470-1476 (1992).

29 Meyer, M. *et al.* FGF receptors 1 and 2 are key regulators of keratinocyte migration in
vitro and in wounded skin. *J Cell Sci* **125**, 5690-5701, doi:10.1242/jcs.108167 (2012).

30 Rasouli, S. J. *et al.* The flow responsive transcription factor Klf2 is required for myocardial
wall integrity by modulating Fgf signaling. *Elife* **7**, doi:10.7554/eLife.38889 (2018).

31 Sun, J. & Stathopoulos, A. FGF controls epithelial-mesenchymal transitions during
gastrulation by regulating cell division and apicobasal polarity. *Development* **145**,
doi:10.1242/dev.161927 (2018).

32 Yayon, A., Klagsbrun, M., Esko, J. D., Leder, P. & Ornitz, D. M. Cell surface, heparin-like
molecules are required for binding of basic fibroblast growth factor to its high affinity
receptor. *Cell* **64**, 841-848, doi:10.1016/0092-8674(91)90512-w (1991).

33   Rapraeger, A. C., Krufka, A. & Olwin, B. B. Requirement of heparan sulfate for bFGF-
mediated fibroblast growth and myoblast differentiation. *Science* **252**, 1705-1708,
doi:10.1126/science.1646484 (1991).
34   Endo, Y., Ishiwata-Endo, H. & Yamada, K. M. Extracellular matrix protein anosmin
promotes neural crest formation and regulates FGF, BMP, and WNT activities. *Dev Cell* **23**,
305-316, doi:10.1016/j.devcel.2012.07.006 (2012).
35   Geiger, B. & Yamada, K. M. Molecular architecture and function of matrix adhesions. *Cold*
Spring Harb Perspect Biol **3**, doi:10.1101/cshperspect.a005033 (2011).
36   McQuade, K. J., Beauvais, D. M., Burbach, B. J. & Rapraeger, A. C. Syndecan-1 regulates
alphavbeta5 integrin activity in B82L fibroblasts. *J Cell Sci* **119**, 2445-2456,
doi:10.1242/jcs.02970 (2006).
37   Moser, M., Legate, K. R., Zent, R. & Fassler, R. The tail of integrins, talin, and kindlins.
Science **324**, 895-899, doi:10.1126/science.1163865 (2009).
38   Frisch, S. M. & Francis, H. Disruption of epithelial cell-matrix interactions induces
apoptosis. *J Cell Biol* **124**, 619-626, doi:10.1083/jcb.124.4.619 (1994).
39   Mailleux, A. A. *et al.* BIM regulates apoptosis during mammary ductal morphogenesis, and
its absence reveals alternative cell death mechanisms. *Dev Cell* **12**, 221-234,
doi:10.1016/j.devcel.2006.12.003 (2007).
40   Borges, E., Jan, Y. & Ruoslahti, E. Platelet-derived growth factor receptor beta and
vascular endothelial growth factor receptor 2 bind to the beta 3 integrin through its
extracellular domain. *J Biol Chem* **275**, 39867-39873, doi:10.1074/jbc.M007040200
(2000).
41   Klinghoffer, R. A., Hamilton, T. G., Hoch, R. & Soriano, P. An allelic series at the
PDGFalphaR locus indicates unequal contributions of distinct signaling pathways during
development. *Dev Cell* **2**, 103-113 (2002).
42   Tallquist, M. D., French, W. J. & Soriano, P. Additive effects of PDGF receptor beta
signaling pathways in vascular smooth muscle cell development. *PLoS Biol* **1**, E52,
doi:10.1371/journal.pbio.0000052 (2003).
43   Kon, E. *et al.* N-cadherin-regulated FGFR ubiquitination and degradation control
mammalian neocortical projection neuron migration. *Elife* **8**, doi:10.7554/eLife.47673
(2019).
44   Scarpa, E. *et al.* Cadherin Switch during EMT in Neural Crest Cells Leads to Contact
Inhibition of Locomotion via Repolarization of Forces. *Dev Cell* **34**, 421-434,
doi:10.1016/j.devcel.2015.06.012 (2015).
45   Ciruna, B. & Rossant, J. FGF signaling regulates mesoderm cell fate specification and
morphogenetic movement at the primitive streak. *Dev Cell* **1**, 37-49 (2001).
46   Sun, X., Meyers, E. N., Lewandoski, M. & Martin, G. R. Targeted disruption of Fgf8 causes
failure of cell migration in the gastrulating mouse embryo. *Genes Dev* **13**, 1834-1846,
doi:10.1101/gad.13.14.1834 (1999).
47   Nieto, M. A., Huang, R. Y., Jackson, R. A. & Thiery, J. P. Emt: 2016. *Cell* **166**, 21-45,
doi:10.1016/j.cell.2016.06.028 (2016).
48   Hoch, R. V. & Soriano, P. Context-specific requirements for Fgfr1 signaling through Frs2
and Frs3 during mouse development. *Development* **133**, 663-673, doi:10.1242/dev.02242
(2006).

49   Raymond, C. S. & Soriano, P. ROSA26Flpo deleter mice promote efficient inversion of
conditional gene traps in vivo. *Genesis* **48**, 603-606, doi:10.1002/dvg.20659 (2010).
50   Raymond, C. S. & Soriano, P. High-efficiency FLP and PhiC31 site-specific recombination
in mammalian cells. *PLoS One* **2**, e162, doi:10.1371/journal.pone.0000162 (2007).
51   Tallquist, M. D. & Soriano, P. Epiblast-restricted Cre expression in MORE mice: a tool to
distinguish embryonic vs. extra-embryonic gene function. *Genesis* **26**, 113-115 (2000).
52   Bouillet, P. *et al.* Proapoptotic Bcl-2 relative Bim required for certain apoptotic responses,
leukocyte homeostasis, and to preclude autoimmunity. *Science* **286**, 1735-1738,
doi:10.1126/science.286.5445.1735 (1999).
53   Muzumdar, M. D., Tasic, B., Miyamichi, K., Li, L. & Luo, L. A global double-fluorescent Cre
reporter mouse. *Genesis* **45**, 593-605, doi:10.1002/dvg.20335 (2007).
54   Serrano, M. *et al.* Role of the INK4a locus in tumor suppression and cell mortality. *Cell* **85**,
27-37, doi:10.1016/s0092-8674(00)81079-x (1996).
55   Ran, F. A. *et al.* Genome engineering using the CRISPR-Cas9 system. *Nat Protoc* **8**, 2281-
2308, doi:10.1038/nprot.2013.143 (2013).
56   Sandell, L. L., Kurosaka, H. & Trainor, P. A. Whole mount nuclear fluorescent imaging:
convenient documentation of embryo morphology. *Genesis* **50**, 844-850,
doi:10.1002/dvg.22344 (2012).
1075