1

A kinetic model of the central carbon metabolism for acrylic acid production in *Escherichia coli*

Alexandre Oliveira[1], Joana Rodrigues[1¶], Eugénio Ferreira[1&], Lígia Rodrigues[1&], Oscar Dias[1 ¶*]

[1]Centre of Biological Engineering, University of Minho, Braga, Portugal

* Corresponding Author:
E-mail: odias@deb.uminho.pt (OD)

¶ These authors contributed equally to this work

& These authors also contributed equally to this work

Abstract

Acrylic acid is a value-added chemical used in industry to produce diapers, coatings, paints, and adhesives, among many others. Due to its economic importance, there is currently a need for new and sustainable ways to synthesise it. Recently, the focus has been laid in the use of *Escherichia coli* to express the full bio-based pathway using 3-hydroxypropionate as an intermediary through three distinct pathways (glycerol, malonyl-CoA, and *β*-alanine). Hence, the goals of this work were to use COPASI software to assess which of the three pathways has a higher potential for industrial-scale production, from either glucose or glycerol, and identify potential targets to improve the biosynthetic pathways yields.

When compared to the available literature, the models developed during this work successfully predict the production of 3-hydroxypropionate, using glycerol as carbon source in the glycerol pathway, and using glucose as a carbon source in the malonyl-CoA and *β*-alanine pathways. Finally, this work allowed to identify four potential over-expression targets (glycerol-3-phosphate dehydrogenase (G3pD), acetyl-CoA carboxylase (AccC), aspartate aminotransferase (AspAT), and aspartate carboxylase (AspC)) that should, theoretically, result in higher AA yields.

Author summary

Acrylic acid is an economically important chemical compound due to its high market value. Nevertheless, the majority of acrylic acid consumed worldwide its produced from petroleum derivatives by a purely chemical process, which is not only expensive, but it also contributes towards environment deterioration. Hence, justifying the current need for sustainable novel production methods that allow higher profit margins. Ideally, to minimise production cust, the pathway should consist in the direct bio-based production from microbial feedstocks, such as Escherichia coli, but the current yields achieved are still to low to compete with conventional method. In this work, even though the glycerol pathway presented higher yields, we identified the malonyl-CoA route, when using glucose as carbon source, as having the most potential for industrial-scale production, since it is cheaper to implement. Furthermore, we also identified potential optimisation targets for all the tested pathways, that can help the bio-based method to compete with the conventional process.

Introduction

Acrylic acid (AA) ($C_3H_4O_2$) is an important chemical compound that is one of the key components of superabsorbent polymers [1–3]. According to the Allied Market Research, in 2015, the global market for AA was valued at 12,500 million US dollars, and is expected to reach 19,500 million US dollars until 2022 [4]. Despite its economic importance, the vast majority of AA is still produced by the oxidation of propylene or propane in a purely chemical process [1,5,6]. Ergo, the principal method for AA production was found to be expensive, with a high energy demand, thus contributing to the planet's environment decay. Hence, the development of an innovative and sustainable biological production method has been attracting the attention of the scientific community [1,2,7]. In the last decade, several semi-biological methods have emerged and were optimised. These methods consist of the bio-based production of 3-hydroxypropionate (3-HP) and its subsequent chemical conversion to AA. Despite the substantial improvements obtained with these methods, this process involves a catalytic step that increases the production costs and environmental impact due to high energy demands [1–3,5]. Hence, the AA's production method should, ideally, be a bio-based direct route as, in theory, microbial feedstocks are less expensive, allowing a higher profit margin [1]. Moreover, a more sustainable bioprocess allows to decrease non-renewable resources dependence and CO_2 emissions.

Fortunately, in recent years, it has been proven that it is possible to use engineered *Escherichia coli* to convert glucose or glycerol into AA. Like in the semi-biological methods, the bioprocess is also divided into two main parts, the production of 3-HP and its subsequent conversion to AA. This part of the pathway, from 3-HP to AA, has not been extensively studied. So far, there are only three studies that successfully converted glucose or glycerol to AA in *E. coli* [1,2,7]. However, the synthesis of 3-HP is well reported, and three distinct pathways have been identified, namely the glycerol route, the malonyl-CoA route, and the *β*-alanine route. From these pathways, it is well established that the glycerol pathway is associated with the highest yields. However, one of the reactions of this route

requires the supplementation of vitamin B_{12} (Fig 1), which is an expensive practice at an industrial-

scale production, hence a significant disadvantage of this route [8,9].

Fig 1. Biosynthetic pathways for acrylic acid (AA) production from Glucose using 3-

hydroxypropionate (3-HP) as an intermediary.

3-HP can be produced from Glucose through three distinct pathways: glycerol (red arrows), malonyl-

CoA (green arrows), and *β*-alanine (blue arrows). Furthermore, *E. coli* can also direct Glycerol towards

the central carbon metabolism, allowing it to be used as a carbon source.

The bio-based method is currently considered a promising alternative to the conventional process

as the production of 3-HP increased considerably in the last few years. Recently, studies reported

productions of up to 8.10 g/L with the glycerol pathway [1], 3.60 g/L with the malonyl-CoA pathway

[10], and 0.09 g/L with the *β*-alanine pathway [11]. However, the AA yields obtained by Tong et al.

(2016) [2] (0.0377 g/L) and Chu et al. (2015) [1] (0.12 g/L) for the glycerol pathway, and Liu and Liu

(2016) [7] (0.013 g/L) for the malonyl-CoA pathway, established that this process still needs to be

optimised to compete with the currently used methods.

Taking these considerations into account, the main goals of this work are to identify the reactions

of the known routes for AA production (glycerol, malonyl-CoA, and β-alanine pathways) and to

determine which pathway have a higher potential for industrial-scale production. *E. coli*'s central

carbon metabolism (CCM) kinetic models will be used to analyse the three pathways using either

glucose or glycerol as carbon source. Finally, novel optimisation strategies to improve the AA yields

of the three biosynthetic pathways will also be sought.

Results and Discussion

Time Course Simulations

During the simulations, several issues arose, leading to changes in parameters before the analysis

of the 3-HP and AA production. These changes are explained in detail in the S1 Appendix, section 1.1.

Regarding the production of 3-HP in the glycerol pathway, simulations with models set to use

either glucose (Glu-Gly) or glycerol as carbon source (Gly-Gly), predicted, the production of 0.19 g/L

(after three hours), and 8.30 g/L (after six-hours), respectively (Fig 3). Whereas, concerning the

production of AA, the Glu-Gly and Gly-Gly models predicted 0.16 g/L and 6.71 g/L, respectively (Fig

4). From these results, glycerol seems to be associated with higher yields, which is in good agreement

with the available literature [1]. Moreover, regarding the production of AA, the intracellular

concentration of 3-HP showed that there is no accumulation (Fig 4), meaning that most 3-HP is

converted into AA. These results are most likely associated with the use of excessive enzyme

concentration to calculate the V_{max} for the heterologous pathway, which led to a state in which the main

limiting factor in the synthesis of AA was the CCM's flux distribution. However, this is not the case

in vivo, as the studies that tested the full bio-based pathway show that 3-HP and other intermediates

indeed accumulate during this process [1,2].

Fig 2. Simulation results for 3-hydroxypropionate (3-HP) production via the glycerol pathway.

(A) Glucose (GLCx) consumption and variation of extracellular 3-HP (3-HPx) over time; **(B)** Glycerol

(GLYx) consumption and variation of 3-HPx over time.

Fig 3. Simulation results for acrylic acid (AA) production via the glycerol pathway.

(A) Glucose (GLC) consumption and variation of extracellular AA (AAx) over time; **(B)** Variation of

3-hydroxypropionate (3-HP) concentration over time when using Glucose as carbon source; **(C)**

Glycerol (GLYx) consumption and variation of extracellular AAx over time; **(D)** Variation of 3-HP

concentration over time when using Glycerol as carbon source.

When comparing the predictions of the glycerol pathway models (Table 3), it is possible to observe that the predicted 3-HP concentration, with the Gly-Gly model, is slightly different from literature reports. This difference slightly increases when increasing the initial concentration of carbon. For instance, for 40 g/L of glycerol, the predicted 3-HP production is about two times higher. Nevertheless, the model representing the 3-HP production from glycerol exhibits promising results.

Table 1. Literature review on 3-hydroxypropionate (3-HP) and acrylic acid (AA) production yields by the glycerol pathway in metabolically engineered *Escherichia coli*, and comparison with the yields predicted by the dynamic models using the same initial carbon concentration.

Reference	End Product	Carbon Source	Initial Carbon Conc. (g/L)	Titer (g/L)	Predicted Titer (g/L)
Raj et al. (2009)[12]	3-HP	Glycerol	9.20	2.80	3.59
Rathnasingh et al. (2009) [13]	3-HP	Glycerol	18.40	4.40	7.59
Chu et al. (2015) [14]	3-HP	Glycerol	40.00	7.40	17.19
Chu et al. (2015) [1]	3-HP	Glycerol	40.00	8.10	17.19
		Glucose	21.50	3.90	0.42
Tong et al. (2016) [2]	AA	Glycerol	20.00	0.037	8.30

The scenario with the Glu-Gly model is considerably distinct, as a ten-fold lower concentration of 3-HP was predicted. This difference might be associated with the production of glycerol, more specifically, in the flux through G3pD and G3pP, as according to Chu et al. (2015) [1] their strain is able to accumulate more glycerol (2.5 g/L) than this model is able to produce (0.34 g/L), for the same amount of glucose. Unfortunately, such study, which presented the highest AA concentration (0.12 g/L) reported thus far, did not disclose the amount of glucose used to obtain such production. Hence, it was not possible to directly compare the predicted titer from the Glu-Gly model.

When performing simulation using glucose concentrations of 10 g/L and 20 g/L, the model predicts the production of 0.16 g/L and 0.32 g/L of AA, respectively. These results show that the AA production predicted by our model would be in good agreement with the results reported by Chu et al. (2015) [1], if 10 g/L of glucose had been used for the carbon source. However, the model does not accumulate any intermediary compounds of the heterologous pathway; thus, most 3-HP is converted into AA, which does not correctly represent the *in vivo* results. Consequently, when assessing the results to the work of Tong et al. (2016) [2] that tested the production of AA in *E. coli*, the model fails to predict the AA yield accurately, as expected.

Regarding the malonyl-CoA pathway, the model set to use glucose as carbon source (Glu-Mcoa) predicted a titer of 1.99 g/L of 3-HP, while the malonyl-CoA model set to use glycerol as carbon source (Gly-Mcoa) predicted 1.99 g/L of 3-HP (Fig 5). Moreover, the Glu-Mcoa and Gly-Mcoa models predicted the production of 1.62 g/L and 0.17 g/L of AA, respectively (Fig 6). The behaviour analysis of the model showed a considerable intracellular accumulation of dihydroxyacetone in the Gly-Mcoa model (Fig 5C). Hence, most carbon does not reach the CCM, and therefore these results should not be considered as it is not possible to determine the best carbon source to produce AA. Although literature reports suggest a consensus towards the use of glucose as a carbon source, it should be noted that no work using glycerol was found. Thus, glycerol should not be excluded as a promising alternative carbon source.

Fig 4. Simulation results for 3-hydroxypropionate (3-HP) production via the malonyl-CoA pathway. (A) Glucose (GLCx) consumption and variation of extracellular 3-HP (3-HPx) over time; **(B)** Glycerol (GLYx) consumption and variation of 3-HPx over time; **(C)** Variation of intracellular dihydroxyacetone (DHA) concentration over time when using Glycerol.

Fig 5. Simulation results for acrylic acid (AA) production via the malonyl-CoA pathway. (A) Glucose (GLC) consumption and variation of extracellular AA (AAx) over time; **(B)** Variation of 3-

hydroxypropionate (3-HP) concentration over time when using Glucose as carbon source; **(C)** Glycerol (GLYx) consumption and variation of extracellular AAx over time; **(D)** Variation of 3-HP concentration over time when using Glycerol as carbon source.

Regarding the Glu-Mcoa models, the 3-HP production predictions are very similar to those found in the literature (Table 4). However, the models failed to predict the production of AA, as Liu and Liu, (2016) [7] reported the accumulation of 3-HP, which was not replicated by the model (Fig 6).

Table 2. Literature review on 3-hydroxypropionate (3-HP) and acrylic acid (AA) production yields by the malonyl-CoA pathway in metabolically engineered *Escherichia coli*, and comparison with the yields predicted by the dynamic models using the same initial carbon concentration.

Reference	End Product	Carbon Source	Initial Carbon Conc. (g/L)	Titer (g/L)	Predicted Titer (g/L)
Cheng et al. (2016) [16]	3-HP	Glucose	10.00	1.80	1.99
Liu et al. (2016) [10]	3-HP	Glucose	20.00	3.60	3.96
Liu and Liu (2016) [7]	AA	Glucose	20.00	0.013	3.22

Finally, regarding the β-alanine pathway, as shown in Fig 7 and Fig 8, the β-alanine model set to use glycerol as carbon source (Gly-Ba) predicted the production of 0.041 g/L of 3-HP and 0.034 g/L of AA. The model set to use glucose as a carbon source (Glu-Ba) predicted the production of 0.033 g/L of 3-HP and 0.027 g/L of AA. Again, the models did not predict the accumulation of 3-HP that, although desirable, is not realistic. Although simulation results indicate a slight advantage towards using glycerol as a carbon source, there seems to be a consensus in literature towards using glucose as carbon source, as to the best of our knowledge, no studies using the glycerol pathway have been

reported. When compared to previous pathways, both carbon sources produced a considerably lower

concentration of AA in this pathway.

Fig 6. Simulation results for 3-hydroxypropionate (3-HP) production via the *β*-alanine pathway.

(A) Glucose (GLCx) consumption and variation of extracellular 3-HP (3-HPx) over time; **(B)** Glycerol

(GLYx) consumption and variation of 3-HPx over time.

Fig 7. Simulation results for acrylic acid (AA) production via the *β*-alanine pathway.

(A) Glucose (GLC) consumption and variation of extracellular AA (AAx) over time; **(B)** Variation of

3-hydroxypropionate (3-HP) concentration over time when using Glucose as carbon source; **(C)**

Glycerol (GLYx) consumption and variation of extracellular AAx over time; **(D)** - Variation of 3-HP

concentration over time when using Glycerol as carbon source.

The *β*-alanine pathway is the least studied, with very few reports, which might be associated with

the fact that such studies reported significantly lower yields, when compared with the previous

pathways [17]. Indeed, only one study was found concerning 3-HP synthesis in *E. coli* using batch

cultures [11], while studies in which AA is produced through this route are yet to be published. Despite

that, the *β*-alanine model showed promising predictions, at least when comparing to the results

published by Song et al. (2016) [11] (Table 5).

Table 3. Literature review on 3-hydroxypropionate (3-HP) production yields by the *β*-alanine pathway in metabolically engineered *Escherichia coli*, and comparison with the yields predicted by the dynamic models using the same initial carbon concentration.

Reference	End Product	Carbon Source	Initial Carbon Conc. (g/L)	Titer (g/L)	Predicted Titer (g/L)
Song et al. (2016) [11]	3-HP	Glucose	15.00	0.09	0.049

Optimisation Strategies

Ideally, all models capable of producing AA should have been optimised. However, as mentioned before, the Gly-Mcoa model presented issues with dihydroxyacetone accumulation, not being further used in this work. Additionally, the Gly-Gly and Gly-Ba models proved to be unstable when performing the metabolic control analysis (MCA), preventing the flux control coefficient (FCC) ascertainment, due to the lack of a steady-state. Henceforth, only the models designed to use glucose as a carbon source (Glu-Gly, Glu-Mcoa, Glu-Ba) were optimised.

Starting with the glycerol model, the MCA showed that the enzyme with the highest FCC, and thus a more significant influence on AA production, was the G3pD (Fig 9A), which is responsible for converting dihydroxyacetone phosphate into glycerol-3-phosphate. The reaction catalysed by this enzyme is a potential bottleneck in the pathway, thus a target for overexpression. Using COPASI's optimisation task, Mutant Glu-Gly 1 was created. This mutant included an *in silico* overexpression of the selected enzyme, in which the V_{max} of the enzyme was set to 1.392 mM/s, representing a nearly 45-fold increase that resulted in the production of 3.11 g/L of AA (Fig 10A). A subsequent MCA was performed on Mutant Glu-Gly 1, aiming at further optimising the AA production yields. However, the

204 model could not reach a steady-state; hence, the FCCs were not available, thus terminating the

205 optimisation of this model.

Fig 8. Flux Control Coefficients (FCC) results for acrylic acid formation, where the reaction with

the most impact in the yield is highlighted in red.

(A) Results for the glycerol pathway. According to the coefficients, the reaction with the most impact

is the glycerol-3-phosphate dehydrogenase (G3pD), which due to its positive FCC is a potential target

for overexpression; **(B)** Results for the malonyl-CoA pathway. The results showed that the acetyl-CoA

carboxylase (AccC) is a potential bottleneck in the pathway due to the positive FCC; hence, another

target for overexpression; **(C)** Results for the β-alanine pathway. The highest FCC was for the aspartate

aminotransferase (AspAT) which appears to be an ideal target for an overexpression.

Fig 9. Comparison between the original acrylic acid (AA) production with the results obtained

for the mutants developed with the optimisation strategies identified.

(A) AA production using glycerol pathway. Mutant 0 represents the model with the heterologous

pathway, and Mutant 1 the same model with a 45-fold increase in the V_{max} of the glycerol-3-phosphate

dehydrogenase (G3pD) reaction; **(B)** AA production using malonyl-CoA pathway. Mutant 0 represents

the model with the heterologous pathway, and Mutant 1 the same model with a 2.5-fold increase in the

V_{max} of the acetyl-CoA carboxylase (AccC); **(C)** AA production for the β-alanine pathway. Mutant 0

represents the model with the heterologous pathway, and Mutant 1 the same model with a 50-fold

increase in the V_{max} of the aspartate aminotransferase (AspAT).

In the malonyl-CoA pathway model, the FCCs identified one potential overexpression target, the

AccC (Fig 9B), which is responsible for the conversion of acetyl-CoA to malonyl-CoA. The optimum

V_{max} for this enzyme was found to be 0.568 mM/s, corresponding to approximately a 2-fold

overexpression. Mutant Glu-Mcoa 1 was able to produce a 3.11 g/L of AA, which corresponds to a

1.5-fold higher yield than Mutant Glu-Mcoa 0 (Fig 10B). Unfortunately, once again, a second iteration

revealed that the model was unable to reach a steady-state. Thus, it was not possible to identify other

potential targets using this methodology.

Regarding the β-alanine pathway, the MCA showed that the model has several reactions affecting

the AA yield. However, the reaction with the most significant impact is catalysed by AspAT enzyme

(Fig 9C). This reaction allows converting oxaloacetate and *L*-glutamate into aspartate, which is in turn

converted to β-alanine. This reaction has a positive coefficient, which indicates that it is a potential

bottleneck impairing the downstream flux towards the heterologous pathway; thus, the optimisation

goal was to overexpress this enzyme. COPASI estimated that a 50-fold overexpression was ideal for

maximising the production yields, resulting in a V_{max} of 127.4869 mM/s. With this change, the

predicted AA production was 0.97 g/L, which corresponds to a 28-fold increase (Fig 10C). A

subsequent MCA revealed that the main limiting factor to AA production in Mutant Glu-Ba 1 was the

amount of glucose provided to the model; hence the optimisation was terminated with only one target

identified. Similarly, the AspC gene, which is responsible for the production of β-alanine, can also be

considered as a limiting factor for pathway flux, thus becoming a target for optimisation, as the V_{max}

was increased for the β-alanine model to work correctly (S1 Appendix, section 1.1).

The goal of these optimisations was to provide guidelines that can be later implemented *in vivo*

and not predict the AA production accurately. Nonetheless, the final concentrations that were obtained

with the new mutants were used to compare with previous results. As shown in Table 6, the same

concentration of AA (3.11 g/L) was produced by the glycerol and malonyl-CoA pathways. The β-

alanine pathway also showed a substantial yield increase (0.97 g/L). However, the value is still

considerably lower than the obtained with remaining pathways.

Table 4. Summarised results of acrylic acid production for the mutant strains developed for the

glycerol, malonyl-CoA, and β-alanine models, using 10 g/L of glucose as substrate.

Pathway	Strain	Predicted Titer (g/L)
Glycerol	Mutant 0	0.16
	Mutant 1	3.11
Malonyl-CoA	Mutant 0	1.62
	Mutant 1	3.11
β-alanine	Mutant 0	0.027
	Mutant 1	0.97

Conclusion

In conclusion, these models seem to be more accurate in predicting 3-HP synthesis because the

V_{max} for the heterologous enzymes has calculated in excess so that it would not limit the reaction flux.

Moreover, the models also exhibited issues associated with the assimilation of glycerol and β-alanine

production. An effort was put forward towards finding proteomics data that included the absolute

quantification of such enzymes to solve these problems. However, no quantification data was found;

therefore, it will be important in the future to seek such data or, in the lack of new data, to determine it

experimentally. Additionally, an overall best carbon source does not emerge from this work. Instead,

the answer is specific to the selected pathway. When comparing the three bio-based routes, the results

suggest that the glycerol pathway leads to the highest yields, when combined with the used of glycerol

as a carbon source. However, a relevant caveat must be recalled. This pathway involves a reaction that

relies on the presence of vitamin B_{12}, which represents a significant economic disadvantage at an

industrial-scale production [8,9]. Hence, to make this route economically viable, either the yield must

be significantly improved to overcome the cost of the vitamin supplementation, or a cheaper way to

produce B_{12} must be found. Therefore, despite producing less AA, it seems to be beneficial to use the

malonyl-CoA route, as it provided the second-highest yield, does not need any vitamin

supplementation, and there is still room for optimisation [16–18]. Finally, the model optimisation

revealed four targets for overexpression with the goal of increasing the bioavailability of the

intermediaries (glycerol, malonyl-CoA, or β-alanine). Nevertheless, these models only comprise the

CCM. Thus, other strategies to force additional flux towards the heterologous pathway may prove

effective. Finally, after the model optimisation, the final step should be the validation with *in vivo*

experiments to further confirm the results herein gathered.

Materials and Methods

Kinetic Modelling

The dynamic model developed by Millard et al. (2017) [19] was extended to include the missing

pathways associated with 3-HP and AA production. This process was divided into three steps, the

extension of the CCM to include the production of the three intermediary compounds (glycerol,

malonyl-CoA, and β-alanine), the subsequent production of 3-HP, and the production of AA.

Parameter Selection. Kinetic equations and their respective parameters were retrieved from the

available literature. Databases like BioCyc [20], BRENDA [21], Sabio-RK [22] and eQuilibrator [23]

were used to identify the kinetic mechanism of each enzyme and obtain their respective parameters.

Furthermore, instead of a single value, the average of all parameters found for each enzyme, excluding

outliers, was used to obtain a better representation. A summary of the values considered when

calculating the mean value for each reaction is presented in S2 Appendix.

Regarding the parameters required to describe a reaction, the maximal rate (V_{max}) is usually not

reported in the literature. Unfortunately, this parameter is highly dependent on the specificity of the

assay conditions. Hence, the specific activity or the turnover (a.k.a. K_{cat}) are reported instead. Two

distinct methods were used, as a workaround, to estimate values for this parameter. Method 1, adapted

from the work of Chassagnole and colleagues [24], was used for reactions belonging to *E. coli*'s native

metabolism. Initially, a steady-state flux distribution is determined for the original kinetic model. Then,

a genome-scale model of *E. coli* K-12 MG1655 (in this case iML1515) [25] is used to predict the flux

of the new reactions. For this purpose, the common reactions between the kinetic model and the

stoichiometric model are constrained to the previously determined flux distribution (±0.01 mM/s).

Then, a flux variability analysis [26] is performed to estimate the maximum flux (v). By equalising v

to the respective rate law ($V_{max} \times F(X,K)$), the following equation is obtained:

$$v = V_{max} \cdot F(X,K) \Leftrightarrow V_{max} = \frac{v}{F(X,K)} \tag{1}$$

in which X is a vector of parameters, and K a vector of steady-state concentrations for the

metabolites involved in the respective reaction. Furthermore, notice that for newly added metabolites,

the steady-state concentration was assumed to be 1 mM. An example of these calculations and the

resulting V_{max} values are presented in the S1 Appendix, section 1.2.

Method 2 was used for reactions of the heterologous pathways. In this method, the V_{max} was

estimated assuming that the total concentration of enzyme was in surplus (100 mM), thus calculating

this parameter as shown in equation 2:

$$V_{max} = K_{cat} \cdot [E]_T \tag{2}$$

307   **Extension of the Central Carbon Metabolism (CCM).** The production of glycerol, malonyl-

308   CoA, and β-alanine had to be included in the model to insert the three heterologous pathways (Fig 2).

309   The reactions catalysed by the glycerol-3-phosphate dehydrogenase (G3pD) and glycerol-3-phosphate

phosphatase (G3pP) enzymes are required to produce glycerol from dihydroxyacetone phosphate. Considering that this route has to be reversible to use glycerol as a carbon source, reactions catalysed by the glycerol kinase (GlyK), glycerol dehydrogenase (GlyD) and the dihydroxyacetone phosphate transferase (DhaPT) enzymes were included too (Fig 2). As shown in Fig 2, one reaction is required to obtain malonyl-CoA, namely the reaction catalysed by the acetyl-CoA carboxylase (AccC) enzyme. Finally, three reactions were included for the β-alanine pathway. Two of these, promoted by the aspartate aminotransferase (AspAT) and the aspartate carboxylase (AspC) enzymes, are required for the production of β-alanine. A reaction, catalysed by the *L*-glutamate dehydrogenase (GluD) enzyme, is used to produce glutamate, which is required by the AspAT to produce aspartate (Fig 2) [20,27].

Fig 10. Representation of the central carbon metabolism of *Escherichia coli* and the reactions added to the kinetic model.

The reactions depicted by the blue, orange, green and yellow arrows represent, respectively, the glycolysis, pentose-phosphate pathway, tricarboxylic acid cycle and the glyoxylate shunt, which are all present in the original model. The black arrows represent the nine reactions that were added to the model. Finally, red arrows depict the Synth reactions added to account for the presence of the newly added metabolites in other pathways

All reactions and respective stoichiometry are shown below:

G3pD: Dihydroxyacetone phosphate + NADPH + H^+ = Glycerol-3-phosphate + $NADP^+$ (3)

G3pP: Glycerol-3-phosphate + $H_2O \rightarrow$ Glycerol + P_i (4)

GlyK: Glycerol + ATP $\rightarrow$ Glycerol-3-phosphate + ADP + H^+ (5)

GlyD: Glycerol + $NAD^+ \rightarrow$ Dihydroxyacetone + NADH + H^+ (6)

DhaPT: Dihydroxyacetone + Phosphoenolpyruvate → Dihydroxyacetone phosphate + Pyruvate (7)

AccC: Acetyl-CoA + ATP + HCO_3 → Malonyl-CoA + ADP + P_i (8)

GluD: α-Ketoglutarate + NADPH + NH_4^+ = *L*-Glutamate + $NADP^+$ + H_2O (9)

AspAT: Oxaloacetate + *L*-Glutamate = *L*-Aspartate + α-Ketoglutarate   (10)

AspC: Aspartate → β-alanine + CO_2 (11)

Furthermore, the kinetic law equation and the respective parameters for each of the previously

described reactions are presented in Table 1.

An additional set of pseudo-reactions was included in the model; the Synth reactions (Fig 2).

These reactions, inspired by the work of Chassagnole et al. (2002) [24] and Machado et al. (2014) [28],

are used to represent the pathways involved in the breakdown of the newly added metabolites. Mass

action kinetics was assumed for these reactions and, using the same principle as Method 1, the sum of

all fluxes from the reactions that metabolise each metabolite in the stoichiometric model was used to

determine the *k* values for each synth reaction. The resulting *k* values are available in the S1 Appendix,

section 1.2.

Pathways for Acrylic Acid Production. The following step was to insert the three heterologous

pathways to produce AA separately into the extended CCM model. All pathways encompass two

different phases, the production of an intermediary compound, namely 3-HP, and subsequent

production of AA (Fig 1).

Table 5. Rate Law (RL) equations, kinetic parameters and the respective references for each reaction that belong to the native metabolism of

Escherichia coli.

Reaction	E.C. number	RL	Equation	Parameters	Reference
G3pD	1.1.1.94	Rapid Equilibrium Random Bi Bi;	$V_{max} \cdot \dfrac{\left(A \cdot B - \left(\frac{P \cdot Q}{K_{eq}}\right)\right)}{K_{m,a} \cdot K_{m,b}} \Big/ \left[\left(1 + \frac{A}{K_{m,a}} \cdot \frac{B}{K_{m,b}}\right) + \left(1 + \frac{P}{K_{m,p}} \cdot \frac{Q}{K_{m,q}}\right) - 1\right]$	$K_{m,a} = 0.175$ mM; $K_{m,b} = 0.0037$ mM $K_{m,p} = 0.12$ mM; $K_{m,q} = 0.165$ mM $K_{eq} = 900$	[29–31]
G3pP	3.1.3.21	Michaelis-Menten	$V_{max} \cdot \dfrac{A}{K_m + A}$	$K_m = 2.9$ mM	[32]
GlyK	2.7.1.30	Random Bi Bi	$\dfrac{V_{max} \cdot A \cdot B}{K_{d,a} \cdot K_{m,b} + K_{m,b} \cdot A + K_{m,a} \cdot B + A \cdot B}$	$K_{m,a} = 0.0084$ mM; $K_{m,b} = 0.0049$ mM $K_{d,a} = 0.086$ mM	[33–36]
AccC	2.1.3.15	Order Bi Bi	$V_{max} \cdot \dfrac{A}{K_{m,A} \cdot \left(1 + \frac{P}{K_{i,P}}\right) + A} \cdot \dfrac{B}{K_{m,B} + B}$	$K_{m,a} = 0.018$ mM; $K_{m,b} = 0.06$ mM $K_{i,p} = 0.07$ mM	[37–39]
GluD	1.4.1.4	Michaelis-Menten	$V_{max} \cdot \dfrac{A}{K_{m,a} + A} \cdot \dfrac{B}{K_{m,b} + B}$	$K_{m,a} = 0.495$ mM; $K_{m,b} = 0.037$ mM	[40–43]
AspAT	2.6.1.1	Ping-Pong Bi Bi	$V_{max} \cdot \dfrac{\left(A \cdot B - \left(\frac{P \cdot Q}{K_{eq}}\right)\right)}{K_{m,a} \cdot K_{m,b}} \Big/ \left[\left(1 + \frac{A}{K_{m,a}} \cdot \frac{Q}{K_{m,q}}\right) \cdot \left(1 + \frac{B}{K_{m,b}} \cdot \frac{P}{K_{m,p}}\right)\right]$	$K_{m,a} = 19.07$ mM; $K_{m,b} = 0.19$ mM $K_{m,p} = 0.437$ mM; $K_{m,q} = 2.94$ mM $K_{eq} = 3.2$	[44–48]
AspC	4.1.1.11	Michaelis-Menten	$V_{max} \cdot \dfrac{A}{K_m + A}$	$K_m = 0.155$ mM	[49,50]
GlyD	1.1.1.6	Hill Cooperativity	$V_{max} \cdot \dfrac{A^n}{(K_{m,a})^n + B^n} \cdot \dfrac{B^n}{(K_{m,b})^n + B^n}$	$K_{m,a} = 47.83$ mM; $K_{m,b} = 1.385$ mM $n = 0.98$	[51,52]
DhaPT	2.7.1.121	Mass Action	$k \cdot A \cdot B$	Not Found	-

The first phase involves two different enzymes in each pathway. Regarding the glycerol pathway,

such enzymes are the glycerol dehydratase (GlyDH) and the 3-hydroxypropionaldehyde

dehydrogenase (3hpaD). In the malonyl-CoA pathway, the malonyl-CoA reductase (McoaR) enzyme

is responsible for the production of malonic semialdehyde (MSA), which is then converted into 3-HP

by the malonic semialdehyde reductase (MsaR). Regarding the β-alanine pathway, the *β*-alanine

aminotransferase (BaAT) enzyme promotes the conversion of β-alanine together with *α*-ketoglutarate

into *L*-glutamate and MSA. The later is then converted into 3-HP by the MsaR.

GlyDH: Glycerol → 3-HPA + H_2O (12)

3hpaD: 3-HPA + NAD^+ + H_2O → 3-HPA + NADH + $2H^+$ (13)

McoaR: Malonyl-CoA + NADPH + H^+ → MSA + CoA + $NADP^+$ (14)

MsaR: MSA + NADPH + H^+ → 3-HP + $NADP^+$ (15)

BaAT: *β*-alanine + α-Ketoglutarate → MSA + *L*-Glutamate   (16)

The final step is to convert the newly formed 3-HP into AA. This process involves the production

of 3-hydroxypropionyl-CoA (3-HP-CoA) by the 3-hydroxypropionyl-CoA synthase (3hpcoaS), the

subsequent formation of acrylyl-CoA (AA-CoA) by the 3-hydroxypropionyl-CoA dehydratase

(3hpcoaDH), and finally, the production of AA by the acrylyl-CoA thioesterase (AcoaTioE) enzyme,

as shown in Fig 1. The stoichiometry of these reactions is likewise shown below, and the respective

kinetic parameters presented in Table 2.

3hpcoaS: 3-HP + CoA + ATP → 3-HP-CoA + $2P_i$ + AMP   (17)

3hpcoaDH: 3-HP-CoA → AA-CoA + H_2O (18)

Table 6. Rate Law (RL) equations, kinetic parameters and the respective references for each reaction of the three heterologous pathways

(glycerol, malonyl-CoA and β-alanine) required to produce acrylic acid.

Reaction	E.C. number	RL	Equation	Parameters	Reference
GlyDH	4.2.1.28	Specific Activation	$$\frac{E \cdot K_{cat} \cdot A \cdot Activator}{K_{m,a} \cdot K_a + (K_{m,a} + A) \cdot Activator}$$	$K_{cat} = 0.0621 \text{ s}^{-1}$; $K_m = 6.15 \text{ mM}$ $K_a = 0.008 \text{ mM}$	[53,54]
3hpaD	1.2.1.99	Michaelis-Menten	$$\frac{E \cdot K_{cat} \cdot A \cdot B}{K_{m,a} \cdot K_{m,b} + K_{m,b} \cdot A + K_{m,a} \cdot B + A \cdot B}$$	$K_{cat} = 16.73 \text{ s}^{-1}$; $K_{m,a} = 0.39 \text{ mM}$ $K_{m,b} = 1.3 \text{ mM}$	[55]
McoaR	1.2.1.75	Michaelis-Menten	$$\frac{E \cdot K_{cat} \cdot A \cdot B}{K_{m,a} \cdot K_{m,b} + K_{m,b} \cdot A + K_{m,a} \cdot B + A \cdot B}$$	$K_{cat} = 50 \text{ s}^{-1}$; $Km_{,a} = 0.3 \text{ mM}$ $K_{m,b} = 0.03 \text{ mM}$	[56,57]
MsaR	2.6.1.19	Michaelis-Menten	$$\frac{E \cdot K_{cat} \cdot A \cdot B}{K_{m,a} \cdot K_{m,b} + K_{m,b} \cdot A + K_{m,a} \cdot B + A \cdot B}$$	$K_{cat} = 115 \text{ s}^{-1}$; $K_{m,a} = 0.07 \text{ mM}$ $K_{m,b} = 0.07 \text{ mM}$	[58]
BaTA	1.1.1.298	Ping-Pong Bi Bi	$$\frac{E \cdot K_{cat} \cdot A \cdot B}{K_{m,b} \cdot A + K_{m,a} \cdot B \cdot \left(1 + \frac{B}{K_{i,B}}\right) + A \cdot B}$$	$K_{cat} = 47.4 \text{ s}^{-1}$; $K_{m,a} = 5.8 \text{ mM}$ $K_{m,b} = 1.07 \text{ mM}$; $Ki_{,b} = 10.2 \text{ mM}$	[59]
3hpcoaS	6.2.1.36	Michaelis-Menten	$$E \cdot K_{cat} \cdot \frac{A}{K_{m,a} + A} \cdot \frac{B}{K_{m,b} + B} \cdot \frac{C}{K_{m,c} + C}$$	$K_{cat} = 36 \text{ s}^{-1}$; $K_{m,a} = 0.015 \text{ mM}$ $K_{m,b} = 0.01 \text{ mM}$; $K_{m,c} = 0.05 \text{ mM}$	[60]
3hpcoaDH	4.2.1.116	Michaelis-Menten	$$\frac{E \cdot K_{cat} \cdot A}{K_m + A}$$	$K_{cat} = 96 \text{ s}^{-1}$; $K_m = 0.06 \text{ mM}$	[61]
AcoaTioE	3.1.2.20	Michaelis-Menten	$$\frac{E \cdot K_{cat} \cdot A}{K_m + A}$$	$K_{cat} = 0.55 \text{ s}^{-1}$; $K_m = 0.167 \text{ mM}$	[62]

The enzyme concentration (E) value used for all the reactions was 100 mM.

AcoaTioE: AA-CoA + H$_2$O → AA + CoA + H$^+$ (19)

Four models were created for each of the three pathways resulting in a total of twelve distinct

models. To be more precise, for each pathway, models to produce 3-HP or AA, from glucose or

glycerol   as   carbon   sources,   were   put   forward.   All   models   can   be   found   at

https://cutt.ly/aaKineticModels.

Time Course Simulation

Time course simulations were performed to assess 3-HP and AA production over time, using the

deterministic method (LSODA) from COPASI [63], with a duration of three or six hours, to allow the

consumption of all available carbon source. Since the available carbon sources have a different number

of carbons, the initial concentration of such molecules had to ensure that the amount of carbon provided

to the model was the same. Thus, the initial concentrations for glucose and glycerol were 55.5 mM (10

g/L) and 217.2 mM (20 g/L), respectively, which allowed comparing the three pathways for each

carbon source. The model was assessed to available literature regarding these pathways, in which the

simulations' initial concentration of the carbon source was set to replicate the initial conditions of

published results.

Optimisation Strategies

The first step was to determine the flux control coefficients (FCC), through a metabolic control

analysis (MCA). Initially, a feed and drains for glycerol, malonyl-CoA, *β*-alanine, and AA were

included to replicate a continuous model and find a valid steady-state, which is required to determine

the FCCs. These coefficients reflect the level of control that each reaction has over the formation of

AA. Nevertheless, a batch version of the model was used for the optimisation. The optimisation was

performed using the automated optimisation tool provided by COPASI [63]. Here, the algorithm

proposed new *in silico* mutant strains, in which the reaction with most influence was either over-

expressed, under-expressed, or knocked-out through a change in the V_{max}, according to the respective

coefficient. The goal of the optimisation was not to meticulously predict the final concentration of AA,

but rather to find promising targets for optimisation. Hence, the changes in the V_{max} were limited to 50

times the original value to allow overcoming the influence of such reaction *in silico*, while not

impairing the *in vivo* implementation.

The objective function was the maximisation of AA concentration in the time course task. After

creating new mutants, the process was repeated to optimise the mutant strains further. The task

eventually stopped when either the glucose feed was limiting the production of AA, the limiting

reaction was already optimised, or the system could no longer reach a stable steady-state point during

the MCA.

Acknowledgments

This study was supported by the Portuguese Foundation for Science and Technology (FCT)

under the scope of the strategic funding of UIDB/04469/2020 unit and BioTecNorte operation

(NORTE-01-0145-FEDER-000004) funded by the European Regional Development Fund under the

scope of Norte2020 - Programa Operacional Regional do Norte. This article is also a result of the

project 22231/01/SAICT/2016: "Biodata.pt – Infraestrutura Portuguesa de Dados Biológicos",

supported by Lisboa Portugal Regional Operational Programme (Lisboa2020), under the PORTUGAL

2020 Partnership Agreement, through the European Regional Development Fund (ERDF).

Author Contributions. AO performed the *in silico* simulations, analysed the results and draft

the manuscript; EF and LR provided feedback and suggestions on the manuscript; JR and OD assisted

in the study conception and design, and provided feedback and suggestions on the manuscript. All

authors have read and agreed to the published version of the manuscript.

References

1. Chu HS, Ahn J-H, Yun J, Choi IS, Nam T-W, Cho KM. Direct fermentation route for the production of acrylic acid. Metab Eng. 2015;32: 23–29. doi:10.1016/j.ymben.2015.08.005

2. Tong W, Xu Y, Xian M, Niu W, Guo J, Liu H, et al. Biosynthetic pathway for acrylic acid from glycerol in recombinant *Escherichia coli*. Appl Microbiol Biotechnol. 2016;100: 4901–4907. doi:10.1007/s00253-015-7272-z

3. Beerthuis R, Rothenberg G, Shiju NR. Catalytic routes towards acrylic acid, adipic acid and ε-caprolactam starting from biorenewables. Green Chem. 2015;17: 1341–1361. doi:10.1039/C4GC02076F

4. Allied Market Research. Acrylic Acid Market Report. 2016 [cited 20 Oct 2019]. Available: https://www.alliedmarketresearch.com/acrylic-acid-market

5. Corma A, Iborra S, Velty A. Chemical Routes for the Transformation of Biomass into Chemicals. Chem Rev. 2007;107: 2411–2502. doi:10.1021/cr050989d

6. Unverricht S, Arnold H, Tenten A, Hammon U, Hans-Peter N, Klaus H. Patent No.: US 6,998,504.: Method for the catalytic gas phase oxidation of propene into acrylic acid. 2006. pp. 1–16.

7. Liu Z, Liu T. Production of acrylic acid and propionic acid by constructing a portion of the 3-hydroxypropionate/4-hydroxybutyrate cycle from *Metallosphaera sedula* in *Escherichia coli*. J Ind Microbiol Biotechnol. 2016;43: 1659–1670. doi:10.1007/s10295-016-1843-6

8.   Sankaranarayanan M, Ashok S, Park S. Production of 3-hydroxypropionic acid from glycerol by

acid tolerant *Escherichia coli*. J Ind Microbiol Biotechnol. 2014;41: 1039–1050.

doi:10.1007/s10295-014-1451-2

9.   Ashok S, Sankaranarayanan M, Ko Y, Jae K-E, Ainala SK, Kumar V, et al. Production of 3-

hydroxypropionic acid from glycerol by recombinant *Klebsiella pneumoniae ΔdhaTΔyqhD*

which can produce vitamin B_{12} naturally. Biotechnol Bioeng. 2013;110: 511–524.

doi:10.1002/bit.24726

10.   Liu C, Ding Y, Zhang R, Liu H, Xian M, Zhao G. Functional balance between enzymes in

malonyl-CoA pathway for 3-hydroxypropionate biosynthesis. Metab Eng. 2016;34: 104–111.

doi:10.1016/j.ymben.2016.01.001

11.   Song CW, Kim JW, Cho IJ, Lee SY. Metabolic Engineering of *Escherichia coli* for the

Production of 3-Hydroxypropionic Acid and Malonic Acid through *β*-Alanine Route. ACS Synth

Biol. 2016;5: 1256–1263. doi:10.1021/acssynbio.6b00007

12.   Raj SM, Rathnasingh C, Jung W-C, Park S. Effect of process parameters on 3-hydroxypropionic

acid production from glycerol using a recombinant *Escherichia coli*. Appl Microbiol Biotechnol.

2009;84: 649–657. doi:10.1007/s00253-009-1986-8

13.   Rathnasingh C, Raj SM, Jo J-E, Park S. Development and evaluation of efficient recombinant

Escherichia coli strains for the production of 3-hydroxypropionic acid from glycerol. Biotechnol

Bioeng. 2009;104: 729–739. doi:10.1002/bit.22429

14.   Chu HS, Kim YS, Lee CM, Lee JH, Jung WS, Ahn J-H, et al. Metabolic engineering of 3-

hydroxypropionic acid biosynthesis in *Escherichia coli*. Biotechnol Bioeng. 2015;112: 356–364.

doi:10.1002/bit.25444

15.   Liu Z, Liu T. Production of acrylic acid and propionic acid by constructing a portion of the
462   3-hydroxypropionate/4-hydroxybutyrate cycle from *Metallosphaera sedula* in *Escherichia coli*.
463   J Ind Microbiol Biotechnol. 2016;43: 1659–1670.

464   16.   Cheng Z, Jiang J, Wu H, Li Z, Ye Q. Enhanced production of 3-hydroxypropionic acid from
465   glucose via malonyl-CoA pathway by engineered *Escherichia coli*. Bioresour Technol.
466   2016;200: 897–904. doi:10.1016/j.biortech.2015.10.107

467   17.   Liu C, Ding Y, Xian M, Liu M, Liu H, Ma Q, et al. Malonyl-CoA pathway: a promising route
468   for 3-hydroxypropionate biosynthesis. Crit Rev Biotechnol. 2017;37: 933–941.
469   doi:10.1080/07388551.2016.1272093

470   18.   Borodina I, Kildegaard KR, Jensen NB, Blicher TH, Maury J, Sherstyk S, et al. Establishing a
471   synthetic pathway for high-level production of 3-hydroxypropionic acid in *Saccharomyces*
472   *cerevisiae* via *β*-alanine. Metab Eng. 2015;27: 57–64. doi:10.1016/j.ymben.2014.10.003

473   19.   Millard P, Smallbone K, Mendes P. Metabolic regulation is sufficient for global and robust
474   coordination of glucose uptake, catabolism, energy production and growth in *Escherichia coli*.
475   Maranas CD, editor. PLOS Comput Biol. 2017;13: e1005396. doi:10.1371/journal.pcbi.1005396

476   20.   Caspi R, Billington R, Fulcher CA, Keseler IM, Kothari A, Krummenacker M, et al. The
477   MetaCyc database of metabolic pathways and enzymes. Nucleic Acids Res. 2018;46: D633–
478   D639. doi:10.1093/nar/gkx935

479   21.   Jeske L, Placzek S, Schomburg I, Chang A, Schomburg D. BRENDA in 2019: a European
480   ELIXIR core data resource. Nucleic Acids Res. 2019;47: D542–D549. doi:10.1093/nar/gky1048

481   22.   Wittig U, Rey M, Weidemann A, Kania R, Müller W. SABIO-RK: an updated resource for

482   manually curated biochemical reaction kinetics. Nucleic Acids Res. 2018;46: D656–D660.

483   doi:10.1093/nar/gkx1065

484   23.   Flamholz A, Noor E, Bar-Even A, Milo R. eQuilibrator—the biochemical thermodynamics

485   calculator. Nucleic Acids Res. 2012;40: D770–D775. doi:10.1093/nar/gkr874

486   24.   Chassagnole C, Noisommit-Rizzi N, Schmid JW, Mauch K, Reuss M. Dynamic modeling of the

487   central carbon metabolism of *Escherichia coli*. Biotechnol Bioeng. 2002;79: 53–73.

488   doi:10.1002/bit.10288

489   25.   Monk JM, Lloyd CJ, Brunk E, Mih N, Sastry A, King Z, et al. iML1515, a knowledgebase that

490   computes *Escherichia coli* traits. Nat Biotechnol. 2017;35: 904–908. doi:10.1038/nbt.3956

491   26.   Mahadevan R, Schilling CH. The effects of alternate optimal solutions in constraint-based

492   genome-scale   metabolic   models.   Metab   Eng.   2003;5:   264–276.

493   doi:10.1016/j.ymben.2003.09.002

494   27.   Kanehisa M, Sato Y, Furumichi M, Morishima K, Tanabe M. New approach for understanding

495   genome variations in KEGG. Nucleic Acids Res. 2019;47: D590–D595. doi:10.1093/nar/gky962

496   28.   Machado D, Rodrigues LR, Rocha I. A kinetic model for curcumin production in *Escherichia*

497   *coli*. Biosystems. 2014;125: 16–21. doi:10.1016/j.biosystems.2014.09.001

498   29.   Edgar JR, Bell RM. Biosynthesis in *Escherichia coli* of m-Glycerol-3-Phosphate, a Precursor of

499   Phospholipid. In Vitro. 1978;3: 6.

30. Edgar JR, Bell RM. Biosynthesis in *Escherichia coli* of sn-glycerol-3-phosphate, a precursor of

phospholipid. Kinetic characterisation of wild type and feedback-resistant forms of the

biosynthetic sn-glycerol-3-phosphate dehydrogenase. J Biol Chem. 1978;253: 6354–6363.

31. Kito M, Pizer LI. Purification and regulatory properties of the biosynthetic L-glycerol-3-

phosphate dehydrogenase from *Escherichia coli*. J Biol Chem. 1969;244: 3316–3323.

32. Salles IM, Forchhammer N, Croux C, Girbal L, Soucaille P. Evolution of a *Saccharomyces*

cerevisiae metabolic pathway in *Escherichia coli*. Metab Eng. 2007;9: 152–159.

doi:10.1016/j.ymben.2006.09.002

33. Pettigrew D, Yu G, Liu Y. Nucleotide Regulation of *Escherichia coli* Glycerol Kinase: Initial-

Velocity and Substrate Binding Studies. Biochemistry. 1990;29: 8620–8627.

34. Hayashi S-I, Lin ECC. Purification and properties of glycerol kinase from *Escherichia coli*. J

Biol Chem. 1967;242: 1030–1035.

35. Thorner JW, Paulus H. Catalytic and allosteric properties of glycerol kinase from *Escherichia*

coli. J Biol Chem. 1973;248: 3922–3932.

36. Applebee MK, Joyce AR, Conrad TM, Pettigrew DW, Palsson B. Functional and metabolic

effects of adaptive glycerol kinase (GLPK) mutants in *Escherichia coli*. J Biol Chem. 2011;286:

23150–23159. doi:10.1074/jbc.M110.195305

37. Soriano A, Radice A, Herbitter A, Langsdorf E, Stafford J, Chan S, et al. *Escherichia coli* acetyl-

coenzyme A carboxylase: characterisation and development of a high-throughput assay. Anal

Biochem. 2005;349: 268–276.

38.   Freiberg C, Brunner N, Schiffer G, Lampe T, Pohlmann J, Brands M, et al. Identification and

characterisation of the first class of potent bacterial acetyl-CoA carboxylase inhibitors with

antibacterial activity. J Biol Chem. 2004;279: 26066–26073.

39.   Meades Jr G, Benson BK, Grove A, Waldrop GL. A tale of two functions: enzymatic activity

and translational repression by carboxyltransferase. Nucleic Acids Res. 2010;38: 1217–1227.

40.   Sharkey M, Engel P. Apparent negative co-operativity and substrate inhibition in overexpressed

glutamate dehydrogenase from *Escherichia coli*. FEMS Microbiol Lett. 2008;281: 132–139.

41.   Sakamoto N, Kotre AM, Savageau MA. Glutamate dehydrogenase from *Escherichia coli*:

purification and properties. J Bacteriol. 1975;124: 775–783.

42.   Mäntsälä P, Zalkin H. Properties of apoglutamate synthase and comparison with glutamate

dehydrogenase. J Biol Chem. 1976;251: 3300–3305.

43.   Di Fraia R, Wilquet V, Ciardiello MA, Carratore V, Antignani A, Camardella L, et al. NADP$^+$-

dependent glutamate dehydrogenase in the Antarctic psychrotolerant bacterium *Psychrobacter*

sp. TAD1: Characterisation, protein and DNA sequence, and relationship to other glutamate

dehydrogenases. Eur J Biochem. 2000;267: 121–131.

44.   Yagi T, Kagamiyama H, Nozaki M, Soda K. Glutamate-aspartate transaminase from

microorganisms. Methods Enzymol. 1985;113: 83–89.

45.   Chow MA, McElroy KE, Corbett KD, Berger JM, Kirsch JF. Narrowing substrate specificity in

a directly evolved enzyme: the *A293D* mutant of aspartate aminotransferase. Biochemistry.

2004;43: 12780–12787.

46. Mavrides C, Orr W. Multispecific aspartate and aromatic amino acid aminotransferases in

Escherichia coli. J Biol Chem. 1975;250: 4128–4133.

47. Deu E, Koch KA, Kirsch JF. The role of the conserved Lys68*: Glu265 intersubunit salt bridge

in aspartate aminotransferase kinetics: multiple forced covariant amino acid substitutions in

natural variants. Protein Sci. 2002;11: 1062–1073.

48. Fernandez FJ, de Vries D, Peña-Soler E, Coll M, Christen P, Gehring H, et al. Structure and

mechanism of a cysteine sulfinate desulfinase engineered on the aspartate aminotransferase

scaffold. Biochim Biophys Acta (BBA)-Proteins Proteomics. 2012;1824: 339–349.

49. Ramjee M, Genschel U, Abell C, Smith A. *Escherichia coli* l-aspartate-α-decarboxylase:

preprotein processing and observation of reaction intermediates by electrospray mass

spectrometry. Biochem J. 1997;323: 661–669.

50. Williamson JM, Brown GM. Purification and properties of L-Aspartate-α-decarboxylase, an

enzyme that catalyses the formation of β-alanine in *Escherichia coli*. J Biol Chem. 1979;254:

8074–8082.

51. Piattoni C, Figueroa C, Diez M, Parcerisa I, Antuña S, Comelli R, et al. Production and

characterisation of *Escherichia coli* glycerol dehydrogenase as a tool for glycerol recycling.

Process Biochem. 2013;48: 406–412.

52. Zhang H, Lountos GT, Ching CB, Jiang R. Engineering of glycerol dehydrogenase for improved

activity towards 1,3-butanediol. Appl Microbiol Biotechnol. 2010;88: 117–124.

53. Sauvageot N, Pichereau V, Louarme L, Hartke A, Auffray Y, Laplace J-M. Purification,

characterisation and subunits identification of the diol dehydratase of *Lactobacillus collinoides*.

Eur J Biochem. 2002;269: 5731–5737.

54.   Schütz H, Radler F. Propanediol-1, 2-dehydratase and metabolism of glycerol of *Lactobacillus*

brevis. Arch Microbiol. 1984;139: 366–370.

55.   Jo J-E, Raj SM, Rathnasingh C, Selvakumar E, Jung W-C, Park S. Cloning, expression, and

characterisation of an aldehyde dehydrogenase from *Escherichia coli* K-12 that utilises 3-

Hydroxypropionaldehyde as a substrate. Appl Microbiol Biotechnol. 2008;81: 51–60.

doi:10.1007/s00253-008-1608-x

56.   Hügler M, Menendez C, Schägger H, Fuchs G. Malonyl-Coenzyme A Reductase from

Chloroflexus aurantiacus, a Key Enzyme of the 3-Hydroxypropionate Cycle for Autotrophic

CO_2 Fixation. J Bacteriol. 2002;184: 2404–2410. doi:10.1128/JB.184.9.2404-2410.2002

57.   Liu C, Wang Q, Xian M, Ding Y, Zhao G. Dissection of Malonyl-Coenzyme A Reductase of

Chloroflexus aurantiacus Results in Enzyme Activity Improvement. Gill AC, editor. PLoS One.

2013;8: e75554. doi:10.1371/journal.pone.0075554

58.   Kockelkorn D, Fuchs G. Malonic Semialdehyde Reductase, Succinic Semialdehyde Reductase,

and Succinyl-Coenzyme A Reductase from *Metallosphaera sedula*: Enzymes of the Autotrophic

3-Hydroxypropionate/4-Hydroxybutyrate Cycle in *Sulfolobales*. J Bacteriol. 2009;191: 6352–

6362. doi:10.1128/JB.00794-09

59.   Liu W, Peterson PE, Langston JA, Jin X, Zhou X, Fisher AJ, et al. Kinetic and Crystallographic

Analysis of Active Site Mutants of *Escherichia coli* γ-Aminobutyrate Aminotransferase.

Biochemistry. 2005;44: 2982–2992. doi:10.1021/bi048657a

60.   Alber BE, Fuchs G. Propionyl-Coenzyme A Synthase from *Chloroflexus aurantiacus*, a Key

Enzyme of the 3-Hydroxypropionate Cycle for Autotrophic CO_2 Fixation. J Biol Chem.

2002;277: 12137–12143. doi:10.1074/jbc.M110802200

61. Teufel R, Kung J, Kockelkorn D, Alber B, Fuchs G. 3-hydroxypropionyl-coenzyme A

dehydratase and acryloyl-coenzyme A reductase, enzymes of the autotrophic 3-

hydroxypropionate/4-hydroxybutyrate cycle in the *Sulfolobales*. J Bacteriol. 2009;191: 4572–

4581.

62. Zhuang Z, Song F, Zhao H, Li L, Cao J, Eisenstein E, et al. Divergence of Function in the Hot

Dog Fold Enzyme Superfamily: The Bacterial Thioesterase YciA †. Biochemistry. 2008;47:

2789–2796. doi:10.1021/bi702334h

63. Hoops S, Sahle S, Gauges R, Lee C, Pahle J, Simus N, et al. COPASI—a COmplex PAthway

SImulator. Bioinformatics. 2006;22: 3067–3074. doi:10.1093/bioinformatics/btl485

Supporting information

594   **S1 Appendix. Supplementary results and parameter adjustments.** Explanations behind

595 parameter adjustments adopted to circumvent simulation issues, and results for the Vmax calculation

596 using method 1 and method 2.

597   **S2 Appendix. Kinetic parameters.** This file will present all the kinetic parameters and equations

598 used to model AA production.

(A)

(B)

(A)

(B)