Hyperbolic odorant mixtures as a basis for more efficient signaling between flowering plants and bees

Majid Ghaninia[1], Yuansheng Zhou[2,3], Anina C. Knauer[4], Florian P. Schiestl[4], Tatyana O. Sharpee[2,3*], Brian H. Smith[1*]

[1] School of Life Sciences, Arizona State University, 427 E. Tyler Mall, Tempe, AZ 85287, USA

[2] The Salk Institute for Biological Studies, Computational Neurobiology Laboratory, North Torrey Pines Road, La Jolla, CA 92037, USA

[3] University of California, San Diego, La Jolla, CA 92037, USA

[4] Institute of Systematic and Evolutionary Botany University of Zurich, Zollikerstrasse 107 8008 Zurich, Switzerland

*Co-equal senior and corresponding authors

Abstract

Animals use odors in many natural contexts, for example, for finding mates or food, or signaling danger. Most analyses of natural odors search for either the most meaningful components of a natural odor mixture, or they use linear metrics to analyze the mixture compositions. However, we have recently shown that the physical space for complex mixtures is 'hyperbolic', meaning that there are certain combination of variables that have a disproportionately large impact on perception and that these variables have specific interpretation in terms of metabolic processes taking place inside the flower and fruit that produce the odors. Here we show that the statistics of odorants and odorant mixtures produced by inflorescences (*Brassica rapa*) are also better described with a hyperbolic rather than a linear metric, and that combinations of odorants in the hyperbolic space are better predictors of the nectar and pollen resources sought by bee pollinators than the standard Euclidian combinations. We also show that honey bee and bumble bee antennae can detect most components of the *B. rapa* odor space, and the strength of responses correlates with positions of odorants in the hyperbolic space. In sum, a hyperbolic representation can be used to guide investigation of how information is represented at different levels of processing in the CNS.

Introduction

Natural odors are typically mixtures of many different chemical compounds (Raguso, 2008). To better understand how these different odorant compounds might be perceived and processed by insects and mammals, it is very useful to work within a low-dimensional representation of odorants (Keller et al., 2017; Ravia et al., 2020; Snitz et al., 2013), where the dimensions define common physical or perceptual properties. At the same time, there is strong evidence in the olfactory system for nonlinear interactions between odors, which suggests that other types of spaces need to be evaluated. Recently, we showed that volatile molecules present in the natural environment produced by fruits can be organized as an approximately two-dimensional map where distances between odors are determined according to a *hyperbolic* metric (Zhou et al., 2018). Hyperbolic spaces are characterized by an exponential expansion of states and provide an approximation to hierarchical tree-like structures (Krioukov et al., 2010). Positions close to the origin

describe more central nodes, whereas those further from the origin describe derivative
products of reactions in metabolic pathways.

This hyperbolic structure makes intuitive sense for semiochemicals produced by plants.
The reason for this is that different chemical compounds often arise from the same or
linked metabolic pathways within the plants (Borghi et al., 2017). And so, positions of
volatile molecules reflects, and can be used to infer, the activity of underlying metabolic
pathways. Thus, taking into account the low-dimensional and hyperbolic aspects of the
space of natural odors makes it possible to extract more reliable messages about the state
of the plant important to pollinators compared to the case where mixtures are evaluated
according to a Euclidean metric.

We focus here on floral odors produced by *Brassica rapa*, which is pollinated by insects
such as honey bees (*Apis mellifera*) and bumble bees (*Bombus* sp) that use floral odors
to identify flowers that contain nectar and pollen. Floral perfumes of *B. rapa* contain at
least 26 different chemical compounds (Kobayashi et al., 2012). We show that bees can
detect many or all of the monomolecular odorants in the mixture, and that several can
contribute to detection of the mixture. However, the mixture gives a stronger, more reliable
signal across a broader range of doses than the individual components. We show that
dimensions of the mixtures computed within hyperbolic space had a greater predictive
power about the amount of pollen and nectar than those computed using a Euclidean
metric. Our data are consistent with a model in which components of hyperbolic
submixtures together amplify important, independent messages to bees about floral
properties (Zhou et al., 2018).

Materials and methods

Experimental model and subject details

76 In this study we used foragers of two bee species, honey bees (*Apis mellifera*) and bumble
77 bees (*Bombus terristris*). The honey bee foragers were collected at the entrance of the
78 hives located in the School of Life Sciences at Arizona State University, on Tempe
79 campus, AZ, USA. The bumble bee foragers were captured from the entrance of the
80 commercially purchased boxes (Koppert Biological Systems; koppertus.com).

Preparation and electroantennographic detection

The collected foragers were immobilized by cooling and individually restrained in a row of eight truncated pipette tips ensuring that the heads and the antennae protruded from the narrow end of the pipettes. One antenna of the restrained bee was then cut off from the pedicel and subsequently transferred to a custom-made antennal holder composed of two capillary glass (tip and reference) electrodes (World Precision Instruments, Inc., USA) filled with insect saline solution. Each electrode was positioned alongside of the length of a microscope slide (75 × 25 × 1 mm) and secured on a small ball of dental wax. The tip and the cut base of the antenna were gently mounted between the tip and the reference electrodes, respectively (Supplementary Fig S1). The holder was then placed on the stage of an Olympus BX51WIF light microscope and viewed at 60× magnification for electroantennogram (EAG) experiments.

The EAG experiments were performed by inserting silver wires into the electrodes until the electrical recording with OSNs was attained. The EAG signal was amplified using an AC/DC differential amplifier (A-M Systems, Inc., Model 3000, USA), subsequently digitized using a digitizer (1440 A DigiData, Molecular Devices, USA), visualized and saved on a PC computer, and the maximum depth of the EAG deflection was measured offline using Clampfit 10.3 software. The EAG responses were normalized (and presented in percent) to the control (hexane) to compensate for the plausible antennal fatigue caused by e.g. repeated delivery of various compounds and/or prolonged recording. For normalization, the EAG responses elicited by individual compounds at a given dose were divided by the average response obtained by the control passed over the antenna at the outset and end of each run.

Odorants

One objective of our study was to evaluate honey bee and bumble bee antennae for sensitivity to a broader array of odorants identified from *B. rapa*. We selected twelve compounds associated with *B. rapa* scent that were previously identified using chemical analyses (Schiestl, 2014; Taveira et al., 2009). This subset included two odorants – phenylacetaldehyde and α-farnesene - identified as important in behavioral studies (Knauer & Schiestl, 2015) as well as other inflorescence and plant related compounds.

This subset is not a comprehensive set of chemicals identified from *B. rapa*. But it is
sufficient to test for broad sensitivity to chemicals from different classes emitted by the
plant and its inflorescences. Odorants were 1-butene-4-isothiocyanate, α-farnesene,
methyl salicylate, indole, methyl benzoate, benzyl nitrile, *(Z)*-3-hexenyl acetate, p-
anisaldehyde, phenylacetaldehyde, decanal, nonanal, acetophenone. For the mixture we
used just a 1:1 combination of the 12 odorants rather than trying to simulate a more natural
B. rapa mixture. Again, the purpose here was to evaluate how robust the signal from a
mixture, rather than specifically the natural mixture, would be relative to the individual
odorant components. Other characteristics of the compounds are displayed in Table 1. All
compounds were diluted in hexane (Fisher Scientific, USA).

Odor delivery

A 15µl aliquot of a solution of the individual compounds was subsequently loaded onto
two pieces of circular filter papers (10 mm diameter each, Whatman, UK) located in the
tip of the Pasteur pipettes. The pipette tip was placed into a hole made along a delivery
glass tube whose one end was hooked up to a stimulus controller (CS-55, Syntech, The
Netherlands) through a Teflon tube and the other end was directed towards the
preparation approximately 1-2cm from it. The stimulus controller delivered the individual
stimuli for 0.5 sec into a constant 89ml/min humidified and purified airstream passing over
the preparation though the delivery glass tube. For every run, a new antennal preparation
was used. To prevent the previously delivered odorant molecules from lingering in the
antennal periphery, residual odors were evacualted from the arena for 30 sec after each
odor delivery, and then another 30sec was allowed to pass (1 minute total time interval)
before delivering the next compound. To evaluate whether response declines initially
observed at higher doses are due to sensory adaptation persisting over the one minute
recovery interval, we repeated the experiment at high doses using a 5 minute interval.

Dose response experiment

All compounds were applied at increasing doses in linear steps ranging from 10^{-11} g/L to
10^3 g/L. Different doses of each stimulus were tested only once (always from the lowest
to highest doses) on individual antennae. Each dose was tested at least three times and
each time a new antennal preparation was used. For the complete mixture, 50µl of each

of the twelve compounds at a given dose were combined in a vial so that, at the end, the

vial contained a total volume of 600µl (50µl of twelve compounds) at that dose. Therefore,

the composite concentration of the mixture is the same as in the individual solutions, but

the concentration of individual compounds in that mixture is only 1/12th of that in the

individual solutions. Delivery of the compounds and analysis of the responses were done

as described above.

Omission (subtraction) experiment

For the omission experiment, we removed individual compounds from the 10g/L complete

mixture. The removal of a given compound in the mixture was compensated for by adding

an additional 4.5µl of each of the 11 remaining compounds. In other words, to reach the

total volume of 600µl, 54.5µl of each of the 11 remaining compounds (i.e. complete

mixture minus compound X) were combined in a separate vial. Therefore, the composite

concentration was the same as in the individual solutions, but the concentration of any

single compound was only 1/11th of what it is in the individual solutions. In total 13

recordings from both species were made and for each recording a new antennal

preparation/individual was used. After each recording, stimuli were randomized. Delivery

of the compounds and analysis of the responses were done as described above.

Floral odor sampling and analytical procedures

Odor samples from B. rapa inflorescences used herein were collected and first published

in Knauer & Schiestl (Knauer & Schiestl, 2015), where a description of methods can be

found.

Software

Clampfit 10.3 was used to analyze the intensity of the EAG responses in mV. The software

may be available online at www.moleculardevices.com. We used R 3.3.0 for data entry

and statistical analysis.

Statistics

For the dose-response experiments we used data collected at 10g/L as a reference point

and applied both the Kruskal-Wallis test and ANOVA on both honey bees and bumble

bees to compare the EAG response intensities for each of the *Brassica* odors. We used

Tukey HSD post hoc test to confirm where the difference occurred between odorants. For

the subtraction experiments we first applied both the Kruskal-Wallis test and ANOVA to

compare the responses elicited by the incomplete mixtures (mixture with individual

compounds omitted). P-values lower than 0.05 were considered to be statistically

significant. All results are represented as mean ± SEM. The "n" represents the number of

antennae tested.

Multi-dimensional Scaling

177 There are two general categories of performing multi-dimensional scaling: a metric multi-

178 dimensional scaling and a non-metric multi-dimensional scaling (Kruskal, 1964). Metric

179 MDS attempts to approximate the geometric distances *dij* within a low dimensional

180 representation using the loss function defined as:

181
$$L = \sqrt{\frac{\sum \left(d_{ij} - \widehat{d_{ij}}\right)^2}{\sum d_{ij}^2}},$$

where $\widehat{d_{ij}}$ are distances in the low-dimensional embedding. The non-metric MDS by

comparison only reconstructs original distances up to a monotonic transformation (only

rank-ordering is preserved) (Kruskal, 1964). This confers the algorithm with extra

robustness. In this work we used a non-metric MDS while embedding points into a low-

dimensional hyperbolic space and using a hyperbolic metric to evaluate distances

between data points within the low-dimensional embedding space.

There are multiple representations of a hyperbolic space. Here we used the so-called

native representation with polar coordinates (Krioukov et al., 2010). The angular

coordinates in this space are the same as in an Euclidean polar coordinates, while the

radius R characterizes the hierarchical depth of the structure and measures the degree of

hierarchy in data. The distance between two points is evaluated as:

193
$$\cosh d_{ij} = \cosh r_i \cosh r_j - \sinh r_i \sinh r_j \, cos\Delta\theta_{ij},$$

194 Where ri and rj are the radial coordinates of the two points, and$\Delta\theta_{ij}$ is the angle between

195 them. In D-dimensional HMDS, we initialize the embedding process by uniformly sampling

196 points within radius R. The directions of points are uniformly sampled around the high-

197 dimensional sphere, and the radial coordinate is sampled according to the following

198 probability distribution:

199
$$\rho(r) \sim (\sinh r)^{D-1}$$

201 During the iteration process, we update both angular and radial coordinates according to
202 the gradient descent.
203

Results

We first evaluated whether honey bees and bumble bees can detect a broad array of
chemical compounds of *B. rapa* floral scent. The measurements were made based on
electroantennogram (EAG) responses to each of the compounds and to a complete
synthetic mixture (Ignell & Hansson, 2004; Olsson & Hansson, 2013). The EAG response
to any odorant represents summed excitatory and inhibitory activity across all of the axons
from thousands of Olfactory Sensory Neurons spaced along a bee's antenna (Supl Fig 1).
Although EAG measurements are less sensitive than single sensillum recordings, they
provide a more appropriate measure for determining whether OSNs on an antenna are
collectively capable of detecting a large number of chemical compounds, and EAG's
provide a means to assess relative sensitivity to different odorants. The use of single
sensillum recordings would be prohibitive for establishing a complete representation of
chemical compounds an antenna can detect because of the large number of recordings
that would be needed.
218

Honey bees and bumble bees can detect most chemical compounds of

the *B. rapa* mixture, which also gives the most robust response

We tested each antenna from honey bees and bumble bees with an increasing dose
series of odorants diluted in hexane (Figs 1A and B; Supl Fig 2). We observed no tangible
response at doses up to 10^{-1} g/L, at which point responses to some odorants began to
increase. By 10^{-1} or 10 g/L both species' antennae responded to all odorants, with the
exception of 1-butene-4-isothiocyanate in honey bees. At higher doses, i.e. 10^2 and 10^3
g/L, responses to most pure odorants declined. This decline was likely due to sensory
adaptation that persisted for at least the minute that lapsed between each stimulation. The
sole exception was phenylacetaldehyde in honey bees, which showed no evidence of
adaptation through 10^3 g/L. We performed further EAG experiments on honey bees to
determine how the timing between odor deliveries may have affected the responses to

the highest (10^2 and 10^3 g/L) doses. We verified that when the OSNs are allowed enough time, i.e. five minutes, their responsiveness failed to show adaptation (Fig 1A).

It is worth noting that in both species the mixture elicited the most robust responses from 10 through 10^3 g/L. The dose for the mixture refers to the sum of all 12 components. Thus the dose of individual components in the mixture was only a fraction, approximately one-twelfth, of what was presented for each pure odorant at that dose. There was little evidence of adaptation in honey bees to mixtures, even at the short interstimulus intervals. Adaptation was evident in bumble bees; nevertheless, even in this case, the mixture always elicited the highest response.

Fig 1. Honey bees and bumble bees can detect almost all components of the *Brassica* mixture, but the mixture gives the most robust response. For the mixture, the x-axis represents total dose of solutes. Electroantennographic responses of (A) honey bees (mean±SE) and (B) bumble bees (mean±SE, n=3) to the selected odorant components of *Brassica rapa* scent. Note that the olfactory responses to most stimuli at 10^2 and 10^3 g/L are lower than those at 10 g/L. The responses do not show such a decline

248 after longer inter-test intervals, suggesting that decline with the shorter test interval was
249 due to adaptation (right line-graphs in A). See Supl Fig 2 for statistical analysis.
250

Most components contribute to detection of the mixture

252 To parse out the significance of specific components to the mixture we omitted individual

253 components from the mixture at 10 g/L (see Method Details). The logic was that the EAG

254 response would decrease if a component contributes in any way – either qualitatively or

255 quantitatively via, for example, intensity - to sensory processing of the mixture. Otherwise

256 the response would be unchanged when the component is deleted.

Figure 2

257
Fig 2. Several components contribute to detection of the mixture. EAG responses of
honey bee (mean±SE, n=11) and bumble bee antennae (mean±SE, n=6) to the mixtures
of *Brassica rapa* scent with individual component compounds omitted. (A) In *Apis mellifera*
omission of phenylacetaldehyde, nonanal, decanal, acetophenone, and p-anisaldehyde
from the blend significantly decreased the responses elicited by the OSNs. (B) In *Bombus*
terrestris ommission of all compounds, except 1-butene-4-isothiocyanate and nonanal,
from the blend significantly reduced the responsiveness of the OSNs compared to the

complete mixture. Significant reductions were based on 95% Confidence Intervals not
overlapping with 1.0, which was the normalized response to the mixture.
267

We found that several components contribute to detection of the mixture (Fig. 2). In
both species, removal of one component reduced the median normalized response to the
mixture. The extent and degree of reduction differed across compounds. In honey bees
(Fig 2A), removal of phenylacetaldehyde, nonanal, decanal, acetophenone, and p-
anisaldehyde all significantly reduced responses to the mixture. In bumble bees (Fig 2B),
removal of all single components except 1-butene-4-isothiocyanate and nonanal
significantly reduced responsiveness. However, even in cases where the response did
not reach significance, median values and in a few cases the 75% range, were below the
normalized mixture response. Thus, it is possible that even in those cases there is some
contribution to the mixture, and it would be prudent to include those odors in a broader
statistical analysis.

279 Submixtures represent nectar and pollen content

Since honey bees and bumble bees can perceive most if not all odorants in the *B. rapa*
mixture, we sought to establish the statistical relationships among the odorants and how
they correlate to important variables for bees, such as nectar and pollen. Different
approaches have been used to perform this kind of comparison. Previous work by Knauer
and Schiestl (Knauer & Schiestl, 2015) identified two volatile compounds,
phenylacetaldehyde and α-farnesene, that are positively associated with nectar and
pollen. We used the same data set to investigate the possibility that submixtures may
provide more complete information about nectar and pollen in *B. rapa* florets.

To find additional odor components that are associated with the nectar and pollen
content of the flower, and which may supplement detection of those important resources
for the bees, we performed two types of analyses. The first analysis was performed with
standard Principal Component Analysis using the logarithms of odorant doses. This
analysis identified combinations of odorants – i.e. submixtures - that account for largest
variance across samples (Fig 3). The top three PC's cumulatively explained 94.3 of the
total odor variance across the *B. rapa* florets and via factor loadings (p≤0.01) collectively
accounted for all of the individual odorants. PC1 explained 76.3.% of the variance and

was positively associated with benzyl nitrile (0.96) and p-anisaldehyde (0.25). This PC was positively associated with log(pollen)-to-log(nectar) ratio. The second PC explained 17% of the variance and could be associated with the same two odorants but in different proportions and signs: p-anisaldehyde (0.94) and benzyl nitrile (-0.26). This PC component was not correlated, in a statistically significant way, with any linear combination of log(pollen) or log(nectar) content. [We will see that correlations will improve once we transition to the hyperbolic space]. The third PC accounted for 3% of the variance and was positively associated with log(nectar with p=0.02). These components had the highest loading factor: Z-3-Hexenyl acetate (0.67), followed by methyl salicylate (0.55), 1-Butene-4-isothiocyanate (0.29), α-farnesene (0.17), methyl benzoate (0.14), phenlyacetaldehyde (0.13), acetophenone (0.11) and nonanal (0.11).

Fig 3. Mixtures of monomolecular odorants that predict pollen and nectar concetrations in the PCA space. (**A**) Representation of odor mixutres produced by Brassica flowers within the space of three leading principal componds (PCs). Axes that are most correlated with florescence nectar and pollen content are shown in red for log(pollen)-to-log(nectar), blue for log(nectar), and green for log(pollen). (B) Correlation with log(pollen)-to-log(nectar) ratio; (C) with log(nectar), and (D) with log(pollen).

A hyperbolic space as a model of odor predictive power

The PCA analysis assumes a Euclidian metric of the odor space (Koulakov et al., 2011), meaning that changes in the concentration of individual monomolecular odorants have the same impact on the predicted distances between the resulting mixtures regardless of the initial concentration and/or odor identity. We recently published a report that the odor spaces for different types of plant and animal derived volatiles are better fitted using a curved 'hyperbolic' metric (Zhou et al., 2018). With this metric the relevant impact of a change in the concentration of an odorant component depends on its initial concentration in ways that cannot be fully captured simply by working with logarithms of the odorant concentration. The impact of odorant concentration change also depended on its identity. The intuition for the relevance of the hyperbolic metric for natural odorants is that the hyperbolic metric approximates activity of hierarchical tree-like processes (Krioukov et al., 2010). Metabolic networks that produce natural odor mixtures certainly have such characteristics. We therefore used hyperbolic Multi-Dimensional Scaling to embed points from *B. rapa* floral volatiles onto a 3D hyperbolic space. The curvature of the spaces was the same as in (Zhou et al., 2018) with R_{max} = 7 and R_{min} = 0.9.

Embedding odorants into this hyperbolic space produced strong correlations with nectar and pollen values (Fig 4). In Fig 4A, circles represent individual monomolecular odorants, and stars represent mixtures produced by individual inflorescences. Distances between these points in hyperbolic space reflect the correlations between odorants or mixtures across samples. Points that are more closely clustered tend to be more correlated across samples. As a first step, we compared the sensitivity of EAG responses to various monomolecular odorants with the odorant positions in the hyperbolic space (Fig 4B). Here, we observed that monomolecular odorants that evoked the strongest EAG responses in honey bees (acetophenone #1, nonanal #2, and decanal #3 cf. Figure 1) have the largest component along the axis that best discriminates between pollen and nectar content (Fig 4A). The rest of the odorants were located in the opposite hemi-sphere and covered it approximately uniformly. Overall, there was a strong correlation between the position of odorants in the hyperbolic space and the EAG responses in antennae. Thus, honey bees show enhanced sensitivity to odorants that best distinguish between

pollen and nectar content of inflorescences. The correlation of EAG amplitudes to nectar/pollen in bumble bees was less clear than for honey bees. Except, like in honey bees, the antennae were most sensitive to acetophenone, which is correlated to this reward axis.

Fig 4. Hyperbolic representation of monomolecular odorants and flower odor mixtures. **(A)** Poincare ball representation of the hyperbolic space (Zhou et al., 2018) together with axes more strongly correlated with log-nectar (blue), log-pollen (green), and their ratio (red). Circles represent individual monomolecular odorants, and stars represent odor mixtures produced by individual inflorescences. Numbers denote odorant identity according to Figure 1A (honey bees). Color denotes the corresponding ratio between log-pollen and log-nectar. For inflorescences this is based on direct measurement, and for odorants on the correlation with their concentration. See also Suppl Figure 3 for a three dimensional version of this figure. (B) Correlation between the strength of EAG response in honey bees (from Figure 1) and odorant position in the hyperbolic space. (C-H) The

361 predicted correlations between projections onto these axes and nectar/pollen
362 measurements.
363

Odorants and mixtures associated with inflorescences had predictive power for the
pollen and nectar content. In particular, we could identify three axes that were strongly
correlated with the log-nectar (red), log-pollen (blue) and the ratio between log-nectar and
log-pollen (green). The corresponding correlations between projections onto these axes
and nectar and pollen flower content are shown for components odorants in Fig. 4C-E and
for mixtures from inflorescences in Figs. 4F-H. Here again one can identify three axes that
are associated respectively with the log-nectar, log-pollen, and their ratio doses (Figs 4F-
H). The correlations are all stronger and more significant than those obtained in the
Euclidean space using the PCA analysis (Figure 5). These results indicate that hyperbolic
mixtures of monomolecular odorants are better predictors of pollen/nectar flower content
than any Euclidean combinations of odorants, which have been identified assuming
independent contributions of monomolecular odorants.

377 Discussion

Many odorants and odorant mixtures serve as communication signals within and between
species. In humans, molecular features of odorants as well as mixtures of odorants are
important for determining perceptual qualities that relate to fruit pleasantness and food
quality (Gilbert et al., 2015; Schwieterman et al., 2014; Zhou et al., 2018). In fact, there is
reason to expect that many fundamental properties of olfaction will apply across a broad
phylogenetic spectrum. For example, the honey bee and fruit fly have 163 and 63
functional receptors in their genomes, respectively.   (Robertson & Wanner, 2006;
Robertson et al., 2003). With the combinatorial nature of receptor responses to any
odorant (Hallem et al., 2004), which insects share with mammals (Nara et al., 2011), the
perceptual space for odor coding is potentially enormous (Gerkin & Castro, 2015).
Therefore, it is a reasonable hypothesis that many olfactory systems have evolved to
analyze and parse different meanings out of complex natural odorant mixtures. We
propose that understanding the relationships between natural odors and the sensory
systems that have evolved to detect them can be aided by a statistical understanding of
the structure of the odor landscape, much in the same way that similar analyses have

been important for understanding vision (Bell & Sejnowski, 1997; Ruderman & Bialek,
1994) and audition (Lewicki, 2002).

In our previous study (Zhou et al., 2018), we used published data from volatiles across
several fruit varieties and mouse urine together with data from human psychophysical
ratings (Gilbert et al., 2015; Schwieterman et al., 2014). A hyperbolic space provided
better fits for both the volatiles and psychophysical preference data. The hyperbolic nature
of the physical odor space has only been explored to date in relation to human detection
(Koulakov et al., 2011; Zhou et al., 2018), although indications of hyperbolic structure were
present also in earlier results (Ravia et al., 2020; Snitz et al., 2013).

Here we use a beneficial insect-plant system related to pollination to test whether
hypotheses that stem from this logic might be more general. *B. rapa* is pollinated naturally
by honey bees, bumble bees and hover flies. When pollinated exclusively by bumble bees
over several generations, the volatile profile of *B. rapa* is increased quantitatively relative
to when pollinated by hover flies, which use more visual than olfactory cues (Gervasi &
Schiestl, 2017; Schiestl et al., 2018). This pollinator-based change in odor profile shows
the importance of odors in insects that have well-developed olfactory systems, and it
implies that there is a significant cost to odor production, which places pressure to make
signalling more efficient. Earlier analysis of the same dataset we used here identified a
small subset of chemical components, especially phenylacetaldehyde and α-farnesene,
that were reliable and – because of the cost - *honest* signals of nectar and/or pollen
(Knauer & Schiestl, 2015). Our analyses confirm that both odorants are detectable by
honey bee and bumble bee olfactory sensory neurons in the antennae. Moreover,
phenylacetaldehyde may also be more resistant to adaptation.

In addition, our analyses show that there is potentially more information present in the
relationships between mixture components. The PCA and hyperbolic MDS analyses both
indicate that combinations of odorants can serve as indicators of the amount of nectar or
pollen in a flower. However, the two methods arrive at that conclusion via different
assumptions. PCA assumes independence among the elements – odorants in this case
– and it extracts from high to low the linear combinations that explain the most variance
in the data set. This method produces correlations with pollen and nectar, but some of the
more important odorants identified by Knauer & Schiestl (Knauer & Schiestl, 2015) only

load on the third component that accounts for only a fraction of the total variance. While

bumble bees learn to pick out whatever component might indicate pollen and/or nectar

(Knauer & Schiestl, 2015), it seems as though PCA is not the best method for analysis of

floral chemistry because it almost missed two known key components.

In contrast, the hyperbolic space provided a substantially better fit for the nectar and

pollen in inflorescences, most likely because of the underlying correlations that arise from

the metabolic pathways that produce odorants in the *B. rapa* mixture. Some mixtures and

some components were better correlated to pollen, nectar, or both. In our EAG analyses,

both the concentration responses and the omission experiment show that several or most

of the components of the mixture can be detected by the antennae of both honey bees

and bumble bees. We do not at this point know what the biologically relevant range for

these compounds might be. However, even if the biologically relevant range is below the

detectable limit with EAG or more sensitive measures, these odors could act

synergistically to enhance detection of other components of the blend, as we discuss

below. Therefore, because the antenna is sensitive to several odorants in the blend, the

use of a hyperbolic multidimensional space to analyze the odor profile of *B. rapa* may be

important for understanding how this information is represented in the CNS. In fact,

sensory processing in both species may be adapted to the information content of *B. rapa*

inflorescences. In our data, EAG responses in honey bees were correlated with the

position of the odorants in the hyperbolic space, and in both bee species, the strongest

EAG responses were for odorants that had strong projections onto the axis in the

hyperbolic space that represented the pollen nectar ratio. We also show that

phenylacetaldehyde, which is a reliable signal of nectar and pollen in *B. rapa* (Knauer &

Schiestl, 2015), elicits strong EAG responses in both species, and it results in less

adaptation in honey bees.

Both odorant components and mixtures from inflorescences correlate to nectar and

pollen in hyperbolic space. So what is more important, individual components or mixtures?

There are several reasons to favor mixtures. In our experiments the mixture of all 12

components, each at a fraction of the dose of the components, elicited the strongest

response. Furthermore, at doses above 10 g/l, antennae showed strong sensory

adaptation to components. In honey bees the mixture did not elicit adaptation. In bumble

bees the mixture elicited adaptation, but responses to the mixture were still the stronger

than components at higher doses in spite of adaptation. With a mixture, more different

receptor types are targeted, exposing each individual receptor type to lower doses of its

cognate chemical compounds. Thus, less adaptation leads to more robust signaling over

a larger range of doses. Furthermore, due to non-linear competitive binding of different

molecular components to odorant receptors, odor-evoked activity patterns are more stable

and first-spike latencies are shorter for mixtures than for pure odorants (Chan et al., 2018).

Both properties – less adaptation and enhanced stability - would enhance the accuracy

and speed of detectability of mixtures over pure odorants. Competitive binding in mixtures

can also predict more complex properties of sensory receptor responses, such as

synergy, antagonism and overshadowing (Reddy et al., 2018; Singh et al., 2019). More

studies are now needed to investigate how other components may synergize signals from

important components like phenylacetaldehyde (Knauer & Schiestl, 2015), for example.

The hyperbolic space, in contrast to PCA, predicts that mixtures can have

disproportional impact on perception. Improved predictions from the use of hyperbolic

metric that we find here for *B. rapa* inflorescences indicates nonlinear synergistic

interactions associated with the presence of multiple odorants. Put differently, in a

hyperbolic space the presence of two components will exponentially increase the strength

of the cue compared to the presence of individual components. Honey bees and bumble

bees searching for pollen, nectar or both could focus on *B. rapa* odor submixtures that

independently reflect levels of each reward. Our work also suggests further studies that

would, for example, reveal the basis for correlations among components of submixtures

– e.g. common biochemical pathways, genetic pleiotropy, epigenetic factors or linkage

disequilibrium among genes that produce odor mixture components.

In summary, we have found that hyperbolic representation of odorants reveal stronger

association with both flower resources and bee antennal responses, compared to

predictions that can be obtained using a Euclidean representation. Hyperbolic

representations have several nonlinear aspects that can be particularly useful for

characterizing olfaction and natural mixtures of odorants. First, the hyperbolic structure

makes it possible to quantify hierarchical relationships in the data. As a result odorants

that are correlated with several other odorants, or are more shared across species

(Biasazin et al., 2019; Sinding et al., 2013), will be assigned more central positions in the

space. Second, based on the properties of vector addition in a hyperbolic space, one

expects odorants with more central positions to have a stronger impact within a mixture

compared to odorants with more distal positions. From this perspective, it is interesting to

note previous reports of stronger tuning in flies for those volatiles that are shared among

the plant host species (Biasazin et al., 2019). It will now be fruitful to test the model using

other coevolved odor/animal relationships, such as pheromones, kairmones,

aposematism, mimicry, crypsis or more specialized pollinator plant relationships (Burger

et al., 2021). In addition, more studies will be needed to understand how the hyperbolic

space is represted in in olfactory processing in the CNS.

Acknowledgements

We thank Dr. Jürgen Liebig for use of his lab equipment, Dr. Kavita Sharma for her advice

regarding EAG recordings, Dr. Chris Jernigan for help with statistical analysis, and Dr.

Margaret Henderson for help with preliminary statistical analyses of odorant mixtures. We

thank Kevin L. Haight for valuable comments on the first draft of the manuscript. This

research was supported by by the Next Generation Networks for Neuroscience Program

(Award #2014217), the Aileen Andrew Foundation, the National Science Foundation

(NSF) award numbers IIS-1254123 and IOS-1556388 to TOS and 1556337 to BHS. Both

awards from IOS resulted from an NSF Ideas Lab 'Cracking the Olfactory Code'. FPS and

ACK were funded by the European Union's Seventh Framework Program ([FP7/2007-

2013] [FP7/2007-2011]) under grant agreement no. 281093.

Supplemental Figure 1

508
Supplemental Figure 1: (A) a schematic drawing of an electroantennaography setup.
The two ends of the antennal shaft, after being cut off from a bee's head capsule, were
inserted in the two capillary glasses (reference and tip electrodes) containing insect
ringer's solution. For more details about the procedure see Method Details. Sample
electroantennogram traces when a honey bee was exposed to the increasing
concentrations of (B) phenylacetaldehyde and (C) complete mixture. Note that the
olfactory responses to both stimuli at 10^2 and 10^3 g/L are lower than those at 10 g/L. The
responses are restored after enough inter-delivery time was given to the OSNs (see
Figure 1).

Supplemental Figure 2

518
Supplemental Figure 2: Antennal responses to the individual components and the
mixture are different at 10 g/L. Electroantennographic responses of (A) honey bees (n=3)
and (B) bumble bees (n=3) to the scent of *Brassica rapa* at 10 g/L. We first compared the
intensity of the responses obtained when the compounds were presented individually at
10 g/L and found that the EAG values for the different odors in both species are
significantly different (ANOVA, F_{Apis}= 9.86, F_{Bombus}= 11.16, df=12, p<0.00001). We
then compared the intensity of the responses obtained when the compounds were
presented individually at 10 g/L with that in the complete mixture. The EAG values for the
different odors in both species are significantly different from those for the mixture
(ANOVA, df=12, Tukey's HSD, p<0.05). Nonetheless, in *Apis mellifera* the response
intensity to nonanal, decanal, and acetophenone alone increased to that elicited by the
mixture (Tukey's HSD, p>0.1). In *Bombus terristris*, however, the response to
phenylacetaldehyde, acetophenone, and methyl benzoate was similar to the mixture
(Tukey's HSD, p>0.1).

533
534 **Supplemental Figure 3:** Three dimensional representation of Poincare ball in Fig. 4A.
535
536

References

Bell, A. J., & Sejnowski, T. J. (1997). The "independent components" of natural scenes are edge filters. *Vision Res*, *37*(23), 3327-3338.

Biasazin, T. D., Larsson Herrera, S., Kimbokota, F., & Dekker, T. (2019). Translating olfactomes into attractants: shared volatiles provide attractive bridges for polyphagy in fruit flies. *Ecol Lett*, *22*(1), 108-118. https://doi.org/10.1111/ele.13172

Borghi, M., Fernie, A. R., Schiestl, F. P., & Bouwmeester, H. J. (2017). The Sexual Advantage of Looking, Smelling, and Tasting Good: The Metabolic Network that Produces Signals for Pollinators. *Trends Plant Sci*, *22*(4), 338-350. https://doi.org/10.1016/j.tplants.2016.12.009

Burger, H., Marquardt, M., Babucke, K., Heuel, K. C., Ayasse, M., Dötterl, S., & Galizia, C. G. (2021). Neural and behavioural responses of the pollen-specialist bee Andrena vaga to Salix odours. *J Exp Biol*, *224*(13). https://doi.org/10.1242/jeb.242166

Chan, H. K., Hersperger, F., Marachlian, E., Smith, B. H., Locatelli, F., Szyszka, P., & Nowotny, T. (2018). Odorant mixtures elicit less variable and faster responses than pure odorants. *PLoS Comput Biol*, *14*(12), e1006536. https://doi.org/10.1371/journal.pcbi.1006536

Gerkin, R. C., & Castro, J. B. (2015). The number of olfactory stimuli that humans can discriminate is still unknown. *Elife*, *4*. https://doi.org/10.7554/eLife.08127

Gervasi, D. D., & Schiestl, F. P. (2017). Real-time divergent evolution in plants driven by pollinators. *Nat Commun*, *8*, 14691. https://doi.org/10.1038/ncomms14691

Gilbert, J. L., Guthart, M. J., Gezan, S. A., Pisaroglo de Carvalho, M., Schwieterman, M. L., Colquhoun, T. A., . . . Olmstead, J. W. (2015). Identifying Breeding Priorities for Blueberry Flavor Using Biochemical, Sensory, and Genotype by Environment Analyses. *PLoS One*, *10*(9), e0138494. https://doi.org/10.1371/journal.pone.0138494

Hallem, E. A., Ho, M. G., & Carlson, J. R. (2004). The molecular basis of odor coding in the Drosophila antenna. *Cell*, *117*(7), 965-979.

Ignell, R., & Hansson, B. S. (2004). Insect olfactory neuroethology–an electrophysiological perspective. In *Advances in Insect Sensory Neuroscience* (pp. 319-347).

Keller, A., Gerkin, R. C., Guan, Y., Dhurandhar, A., Turu, G., Szalai, B., . . . Consortium, D. O. P. (2017). Predicting human olfactory perception from chemical features of odor molecules. *Science*, *355*(6327), 820-826. https://doi.org/10.1126/science.aal2014

Knauer, A. C., & Schiestl, F. P. (2015). Bees use honest floral signals as indicators of reward when visiting flowers. *Ecol Lett*, *18*(2), 135-143. https://doi.org/10.1111/ele.12386

Kobayashi, K., Arai, M., Tanaka, A., Matsuyama, S., Honda, H., & Ohsawa, R. (2012). Variation in floral scent compounds recognized by honeybees in Brassicaceae crop species. *Breed Sci*, *62*(4), 293-302. https://doi.org/10.1270/jsbbs.62.293

Koulakov, A. A., Kolterman, B. E., Enikolopov, A. G., & Rinberg, D. (2011). In search of the structure of human olfactory space. *Front Syst Neurosci*, *5*, 65. https://doi.org/10.3389/fnsys.2011.00065

Krioukov, D., Papadopoulos, F., Kitsak, M., Vahdat, A., & Boguñá, M. (2010). Hyperbolic geometry of complex networks. *Phys Rev E Stat Nonlin Soft Matter Phys*, *82*(3 Pt 2), 036106. https://doi.org/10.1103/PhysRevE.82.036106

Kruskal, J. B. (1964). Multidimensional scaling by optimizing goodness of fit to a nonmetric hypothesis. *Psychometrika*, *29(1)*, 1-27.

Lewicki, M. S. (2002). Efficient coding of natural sounds. *Nat Neurosci*, *5*(4), 356-363.
https://doi.org/10.1038/nn831
Nara, K., Saraiva, L. R., Ye, X., & Buck, L. B. (2011). A large-scale analysis of odor coding in the
olfactory epithelium. *J Neurosci*, *31*(25), 9179-9191.
https://doi.org/10.1523/JNEUROSCI.1282-11.2011
Olsson, S. B., & Hansson, B. S. (2013). Electroantennogram and single sensillum recording in
insect antennae. In *Pheromone Signaling* (pp. 157-177). Springer.
Raguso, R. A. (2008). Wake up and smell the roses: the ecology and evolution of floral scent.
Annual Review of Ecology, Evolution, and Systematics, *39*, 549-569.
Ravia, A., Snitz, K., Honigstein, D., Finkel, M., Zirler, R., Perl, O., . . . Sobel, N. (2020). A measure
of smell enables the creation of olfactory metamers. *Nature*, *588*(7836), 118-123.
https://doi.org/10.1038/s41586-020-2891-7
Reddy, G., Zak, J. D., Vergassola, M., & Murthy, V. N. (2018). Antagonism in olfactory receptor
neurons and its implications for the perception of odor mixtures. *Elife*, *7*.
https://doi.org/10.7554/eLife.34958
Robertson, H. M., & Wanner, K. W. (2006). The chemoreceptor superfamily in the honey bee,
Apis mellifera: expansion of the odorant, but not gustatory, receptor family. *Genome
Res*, *16*(11), 1395-1403. https://doi.org/gr.5057506 [pii] 10.1101/gr.5057506
Robertson, H. M., Warr, C. G., & Carlson, J. R. (2003). Molecular evolution of the insect
chemoreceptor gene superfamily in Drosophila melanogaster. *Proc Natl Acad Sci U S A*,
100 Suppl 2, 14537-14542.
Ruderman, D. L., & Bialek, W. (1994). Statistics of natural images: Scaling in the woods. In (Vol.
73, pp. 814). Physical Review Letters.
Schiestl, F. P. (2014). Correlation analyses between volatiles and glucosinolates show no
evidence for chemical defense signaling in B*rassica rapa.* In (Vol. 2, pp. 10). Frontiers in
Ecology and Evolution.
Schiestl, F. P., Balmer, A., & Gervasi, D. D. (2018). Real-time evolution supports a unique
trajectory for generalized pollination. *Evolution*, *72*(12), 2653-2668.
https://doi.org/10.1111/evo.13611
Schwieterman, M. L., Colquhoun, T. A., Jaworski, E. A., Bartoshuk, L. M., Gilbert, J. L., Tieman, D.
M., . . . Clark, D. G. (2014). Strawberry flavor: diverse chemical compositions, a seasonal
influence, and effects on sensory perception. *PLoS One*, *9*(2), e88446.
https://doi.org/10.1371/journal.pone.0088446
Sinding, C., Thomas-Danguin, T., Chambault, A., Béno, N., Dosne, T., Chabanet, C., . . . Coureaud,
G. (2013). Rabbit neonates and human adults perceive a blending 6-component odor
mixture in a comparable manner. *PLoS One*, *8*(1), e53534.
https://doi.org/10.1371/journal.pone.0053534
Singh, V., Murphy, N. R., Balasubramanian, V., & Mainland, J. D. (2019). Competitive binding
predicts nonlinear responses of olfactory receptors to complex mixtures. *Proc Natl Acad
Sci U S A*, *116*(19), 9598-9603. https://doi.org/10.1073/pnas.1813230116
Snitz, K., Yablonka, A., Weiss, T., Frumin, I., Khan, R. M., & Sobel, N. (2013). Predicting odor
perceptual similarity from odor structure. *PLoS Comput Biol*, *9*(9), e1003184.
https://doi.org/10.1371/journal.pcbi.1003184

Taveira, M., Fernandes, F., de Pinho, P. G., Andrade, P. B., Pereira, J. A., & Valentão, P. (2009).

Evolution of B*rassica rapa* var. r*apa* L. volatile composition by HS-SPMEand GC/IT-MS. In

(Vol. 93, pp. 140-146). Microchemical Journal.

Zhou, Y., Smith, B. H., & Sharpee, T. O. (2018). Hyperbolic geometry of the olfactory space. *Sci*

Adv, *4*(8), eaaq1458. https://doi.org/10.1126/sciadv.aaq1458

TABLE 1: A comprehensive list of all materials and resources used in the
manuscript.

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Chemicals, Peptides, and Recombinant Proteins		
1-Butene-4-isothiocyanate	Combi-Blocks	BATCH:A25781
α-Farnesene	ChemCruz	LOT:L0915
Methyl salicylate	SIGMA-ALDRICH	LOT:MKBW9023V
Indole	CHEM-IMPRX INT'L INC	LOT:001481-20120820
Methyl benzoate	Fluka	LOT:1339386
Benzyl nitrile	ALDRICH	LOT:BCBJ5504V
(Z)-3-Hexenyl acetate	TCI	LOT:A5JXM-TA
p-Anisaldehyde	TCI	LOT:R7QOK-TF
Phenylacetaldehyde	Santa Cruz Biotechnology	LOT:A0616
Decanal	Alfa Aesar	LOT:101709990
Nonanal	Alfa Aesar	LOT:10179106
Acetophenone	Fluka	LOT:BCBH8667V
Hexane	Fisher Scientific	LOT:165552
Experimental Models: Organisms/Strains		
Bumble bees (*Bombus terristris*)	Koppert B.V., The Netherlands	www.koppert.com/pollination
Honey bees (*Apis mellifera carniolan*)	Maintained at Arizona State University	www.asu.edu
Software and Algorithms		
Clampfit 10.3	Molecular Devices, Inc.	https://www.moleculardevices.com/
R (Data analysis)	RStudio	http://www.R-project.org/
Other		
Pipette tip	Fisherbrand	CAT:02-681-172
Borosilicate capillary glass	World Precision Instruments	LOT:1208336
Microscope slide	Fisherfinest	CAT:22-038-103
Dental wax	Surgident	LOT:1106012
BX51WIF light microscope	Olympus, Japan	SN:1H69084
AC/DC differential amplifier	A-M Systems, Inc., Model 3000, USA	SN:61676
Stimulus controller CS-55	Syntech, Germany	http://www.ockenfels-syntech.com
Digitizer	1440 A DigiData, Molecular Devices, Inc., USA	SN:814926

| Filter paper | Whatman, UK | CAT:1004-070 |

633
634