A Chemical-genetics and Nanoparticle Enabled Approach for *in vivo* Protein Kinase Analysis

Fengqian Chen[1], Qi Liu[2,3], Terrell Hilliard[1], Tingzeng Wang[1], Hongjun Liang[4], Weimin Gao[5], Leaf Huang[3], Degeng Wang[1,*]

[1]Department of Environmental Toxicology, The Institute of Environmental and Human Health (TIEHH), Texas Tech University, Lubbock, TX 79416, USA
[2]Department of Dermatology, Johns Hopkins University School of Medicine, Cancer Research Building II, Suite 216, 1550 Orleans Street, Baltimore, MD 21231, USA
[3]Division of Pharmacoengineering and Molecular Pharmaceutics, Eshelman School of Pharmacy, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA
[4] Department of Cell Physiology and Molecular Biophysics, Texas Tech University Health Sciences Center, Lubbock, TX 79430.
[5] Department of Occupational and Environmental Health Sciences, School of Public Health, West Virginia University, Morgantown, USA
*Corresponding author. E-mail: degeng.wang@ttu.edu

Abstract

The human kinome contains >500 protein kinases, and regulates up to 30% of all human proteins. Kinase study is currently hindered by a lack of *in vivo* analysis approaches, mainly due to two factors: our inability to distinguish the kinase reaction of interest from those of other members of the kinome in live cells and the cell impermeability of the ATP analogs. Herein, we aimed to overcome this issue by combining the widely used chemical genetic method developed by Dr. Kevan Shokat and colleagues with nanoparticle-mediated intracellular delivery of the ATP analog. The critical AKT1 protein kinase, which has been successfully studied with the Shokat method, was used as our initial prototype. Briefly, following the Shokat method, enlargement of the ATP binding pocket was performed by mutating the gate-keeper Methionine residue to a Glycine, prompting the mutant AKT1 to preferentially use the bulky ATP analog N^6-Benzyl-ATP-γ-S (A*TPγS) and, thus, differentiating AKT1-catalyzed and other phosphorylation events. The LPC nanoparticle was used for efficient intracellular delivery of A*TPγS, overcoming the cell impermeability issue. We demonstrated that the mutant, but not the wild type, AKT1 was able to use the delivered A*TPγS for autophosphorylation as well as phosphorylating its substrates in live cells. Thus, an *in vivo* protein kinase analysis method has been developed. The strategy should be widely applicable to other protein kinases.

1. Introduction

Protein kinases function by cleaving the γ-phosphate group off ATP and covalently adding the group onto serine, threonine or tyrosine residues of their substrate proteins, up or down-regulating their activities in vital cellular processes such as cell cycles, growth, migration and apoptosis (1,2). Very often, the substrates themselves are protein kinases. Through this phosphorylation process, the protein kinases form many signaling cascades, *e.g.*, the MAP kinase cascades. The kinases and the cascades constitute the backbone of the cellular signaling network, which bridges extracellular and intracellular environment and interconnects subcellular compartments/domains. Many diseases exhibit aberrancies in protein kinases, *i.e.,* over-activation of disease promoting kinases and/or over-repression of disease suppressive ones. Thus, the treatments often target these kinases. For instance, the type II diabetic treatment targets the AMPK protein kinase, and cancer treatments aim to correct the aberrancies of oncogenic and/or tumor-suppressive protein kinases (3,4).

However, kinase study has been hindered by a major challenge – the lack of an *in vivo* analysis technique. Currently, kinase studies rely primarily on *in vitro* assays – incubation of ATP with labeled γ-phosphate group, un-phosphorylated peptide substrate and the kinase in the reaction buffer followed by detecting/quantifying substrate tagging by the labeled γ-phosphate group from ATP. The kinase can be generated through recombinant expression followed by purification and, in case one wishes to quantify an endogenous kinase activity, isolation from cellular extract. However, it is very much questionable whether the *in vitro* substrate tagging reliably reflects the *in vivo* phosphorylation events and their magnitudes.

In order to establish *in vivo* kinase analyses in intact live cells, we need to meet two requirements. First, it is necessary to be able to distinguish the kinase reaction of interest from those of the other protein kinases, as the human kinome contains more than 500 protein kinases. Second, we need to develop a method to generate sufficient intracellular abundance of the tagged ATP analog. The analog by itself is not cell permeable, but is required as a probe to detect and/or monitor the kinase reactions, *i.e.*, transfer of the phosphate group from ATP onto the substrates.

Fortunately, the well established chemical-genetic strategy for identifying kinase-substrates relationships developed by Dr. Kevan Shokat's lab is well suited to accomplish the first requirement. The Shokat method uses site-directed mutagenesis to enlarge a kinase's ATP binding pocket. While not affecting kinase activity, the mutation specifically allows bulky synthetic ATP analogs, such as N6-Benzyl-ATP (A*TP), to fit into the enlarged ATP binding pocket (5-9). Moreover, A*TP was further modified to create A*TP-γ-S by replacing an O with a S in the γ-phosphate group. The mutant protein kinase preferentially use A*TP-gamma-S in its kinase reaction, transferring the thiophosphate tag from A*TP-γ-S onto its substrate proteins. This tag can then be alkylated by a chemical called *p*-nitrobenzylmesylate (PNBM) to generate a thiol-ester for specific antibody recognition or affinity purification followed by LC-MS analysis (10-12). In a word, this method combines ATP-binding pocket enlargement and the bulky ATP analog A*TP-gamma-S to distinguish the kinase reaction of interest from those of the other protein kinases, enabling identification of the substrates of specific kinases (13). However, due to the cell impermeability of the A*TP-γ-S molecule, this method have been executed *in vitro* with cell lysate in test tubes. In order for the thiophosphate-tagging of the substrates to occur in live cells, we need to accomplish the second requirement, *i.e.*, a reliable method for efficient intracellular A*TP-γ-S delivery.

As for this second requirement, a tool also seems exist already. The lipid/calcium/phosphate (LCP) nanoparticle is well suited for this task. It consists of a calcium phosphate (CaP) core, which carries the chemicals to be delivered, and a lipid bilayer, which encapsulates the CaP core and enables cellular uptake of the nanoparticle through endocytosis. This nanoparticle has been used for efficient intracellular delivery of Gemcitabine Triphosphate, a GTP analog used as a chemotherapy agent (2,14). The A*TP-

gamma-S molecule is chemically similar to Gemcitabine Triphosphate. Thus, we became interested in using the LCP nanoparticle as a vehicle for intracellular A*TP-γ-S delivery and, in turn, to enable intracellular thiophosphate-tagging of protein kinase substrates via the Shokat chemical genetic approach; that is, achievement of *in vivo* protein kinase analysis via a combination of the Shokat chemical genetic method and the nanoparticle delivery of the A*TP-γ-S molecule.

To accomplish this goal, we choosed the serine/threonine protein kinase AKT1 (also known as PKB) as the initial prototype. Firstly, AKT1 is functionally and therapeutically important. It is part of the critical phosphoinositide 3-kinase (PI3K) signaling module (15). The non-receptor tyrosine kinase ACK1/TNK2 can also phosphorylate AKT1, resulting in PI-3-kinase-independent activation. AKT1 phosphorylates and regulates a wide range of metabolic and/or regulatory substrate proteins such as IKKalpha, GSK3beta, and FOXO (16,17). Thus, it participates in multiple vital cellular processes, including cell survival and cell growth. As the Akt signaling process aberrancy occurs in multiple human diseases such as diabetes and cancer, AKT1 is a significant therapeutic target for multiple human diseases (18) (19). Secondly, perhaps due to the functional importance, AKT1 is one of the earliest protein kinases studied successfully with the Shokat chemical genetic approach, enhancing the likelihood of success of this study. Moreover, in addition to phosphorylating its numerous substrates, AKT1 undergoes auto-phosphorylation (20), further facilitating this initial study.

Thus, following the Shokat method, the AKT1 ATP binding pocket was enlarged by mutating the gate-keeper Methionine residue into a Glycine. We also used the LCP nanoparticle for efficient intracellular delivery of the A*TPγS molecule, overcoming the cell impermeability issue. We demonstrated that the mutant, but not the wild type, AKT1 was able to use the delivered A*TPγS for autophosphorylation as well as phosphorylating other protein substrates in live cells. In a word, an *in vivo* AKT1 kinase analysis method has been developed, to the best of our knowledge, for the first time. The strategy should be widely applicable to other protein kinases.

2. Materials and methods

2.1 Materials
Alexa-Fluor-647-ATP was obtained from Thermo Fisher Scientific (Massachusetts, USA), and N^6-Benzyladenosine- 5'- (3- thiotriphosphate) (A*TPγS) was obtained from BIOLOG Life Science Institute (Bremen, Germany). IKK alpha monoclonal antibody was obtained from Thermo Fisher Scientific (Massachusetts, USA), anti-thiophosphate ester antibody was obtained from Abcam (Cambridge, UK), and AccuRuler RGB protein ladder and BP-Fectin mammalian cell transfection reagent were obtained from BioPioneer (California, USA). Dioleoyl phosphatydic acid (DOPA), 1,2-dimyristoyl-3-trimethylammonium-propane chloride salt (DOTAP), and 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-N-[amino(polyethylene glycol)-2000] ammonium salt (DSPE-PEG$_{2000}$) were obtained from Avanti Polar Lipids, Inc. (Alabama, USA). DSPE-PEG-anisamide (AEAA) was synthesized in Dr. Leaf Huang's lab as described previously (21). Other chemicals were purchased from Sigma-Aldrich, Inc. (Missouri, USA).

2.2 Cell culture
HCT116 human colon cancer cells, originally obtained from Dr. Michael Brattain's lab (22,23), were cultured in the McCoy's 5A medium (Invitrogen, USA) supplemented with 10% fetal bovine serum, 100 U/mL penicillin, and 100 mg/mL streptomycin (Invitrogen, USA). Cells were cultivated at 37 °C and 5 % CO_2 in an incubator.

2.3 Preparation of A*TPγS-loaded LCP nanoparticles

Briefly, 100 μL of 10 mM specified ATP analog solution (A*TPγS or Alexa-Fluor-647-ATP) in 12.5 mM Na_2HPO_4 solution and 100 μL of 2.5 M $CaCl_2$ solution were dispersed in cyclohexane/Igepal solution (70/30, v/v), respectively, to form an oil phase. After 15 min of stirring, 100 μL of 25 mM DOPA was added into the oil phase for 15 min, then 40 mL of 100 % ethanol was added into the oil phase. Later, the mixture solution was centrifuged at 10,000 g for 20 min to pellet the LCP particle core out of the supernatant solution. After washing with 50 mL ethanol twice, 100 μL of 10 mg/ml cholesterol, 100 μL of 25 mg/ml DOTAP, 100 μL of 25 mg/ml DSPE-PEG, and 10 μL of 25 mg/ml DSPE-PEG-AEAA were added into the LCP core solution. After evaporating under N_2, the residual lipids were suspended in PBS to produce the layer of LCP nanoparticle. After being sonicated for 10 min, the LCP solution was dialyzed in PBS.

2.4 Characterization of A*TPγS-loaded LCP nanoparticles
The LCP nanoparticle particle sizes was determined by Malvern DLS (Royston, UK). The concentration of A*TPγS was determined by an HPLC spectrophotometer (Shimadzu Corp., Japan). Transmission electron microscope (TEM) photos of A*TPγS-loaded LCP nanoparticles were observed under JEOL 100CX II TEM (Tokyo, Japan).

2.5 Intracellular uptake behaviors
Intracellular uptake behaviors were observed with a confocal microscopy. Briefly, the HCT116 cells were plated into confocal dishes (1×10^5 cells per well) and incubated for 24 hours. Then, the cells were incubated with Alex-Fluor-647-ATP-loaded LCP nanoparticles for 4 hours with a concentration of 5 μg/mL, and were nucleus-labeled with Hoechst 33342 dye (Thermo Fisher Scientific) for 15 min. The images were taken with a Zeiss 880 confocal microscope (Germany).

2.6 Plasmid constructs

Wild type Akti plasmid was purchased from Addgene (Addgene plasmid pcDNA3-myr-HA-AKT1 1036). Point mutations made as references (24) in the ATP gatekeeper of AKT1 (M227 → G gatekeeper mutation) were introduced by GENEWIZ (New Jersey, USA), with the method being a modified Stratgene QuikChange® site-directed mutagenesis.

2.7 Expression plasmid transfection

The HCT116 cells were plated into 100 mm^2 dishes and transfected at approximately 90% confluence with the specified plasmid (wild type pcDNA3-myr-HA-AKT1 plasmid or the mutant pcDNA3-myr-HA-AKT1 plasmid) using BP-Fectin mammalian cell transfection reagent (BioPioneer, USA) in accordance with the manufacturer's instructions. One untransected dish was used as a negative control. After 72 h, cells were washed twice with cold PBS. Transfection efficacy was confirmed by western-blot measurement of the expression of the HA-tagged fusion proteins with the HA tag monoclonal antibody (Thermo Fisher Scientific, USA).

2.8 Intracellular A*TPγS delivery and thio-phosphorylation assays

Control and transfected HCT116 cells were incubated with A*TPγS-loaded LCP nanoparticles for 4 h. The cells were washed by cold PBS twice and then lysed with RIPA buffer (which contains protease and phosphatase inhibitors) for 30 minutes at 4 °C. The cell lystaes were treated with 2.5 mM PNBM at room temperature for one hour. Then, the PNBM treated cell lysates were analyzed with a western blot assay to measure protein thio-phosphorylation with the anti-thiophosphate ester rabbit monoclonal antibodies (Abcam, USA).

2.9 Immunoprecipitation assays

The PNBM-treated cell lysates were pre-cleaned by an off-target antibody. Then the cleaned cell lysates were incubated with 10 µL anti-thiophosphate ester rabbit monoclonal antibody (Abcam, USA) at 4 °C for 12 h. 100 µL protein G beads (Thermo Fisher Scientific, USA) were mixed under rotary agitation at 4 °C for 4 h, washed twice by wash buffer, and the beads were collected. Then, proteins were captured in the beads and released by elution buffer. The elution buffer was collected and stored at -80 °C.

2.10 Western blotting assay

First, total protein concentrations of the cell lysates were determined by a BCA protein assay kit (Thermo, USA). Protein extracts were separated by SDS–PAGE, then transferred onto a nitrocellulose membrane (NC membrane). After blocking the NC membrane with 5 % non-fat milk for 1 h at room temperature, it was incubated with the specified first antibody (the HA tag monoclonal antibody, the anti-thiophosphate ester antibody or the IKK alpha monoclonal antibody) at 1:1000 dilution at 4 °C overnight. Membranes were washed five times with TBST solution, incubated with the corresponding secondary antibody (1:3000 dilution) for 1.5 h at normal room temperature, and washed five times with TBST solution. Protein bands were visualized using an enhanced chemiluminescence system according to the manufacturer's instructions (Thermo Fisher Scientific, USA).

3. Results

3.1 Characterization of A*TPγS-loaded LCP Nanoparticles

As mentioned earlier, the LCP nanoparticle is a formulation that has been used for efficient intracellular delivery of the GTP analog Gemcitabine Triphosphate as a chemotherapy agent. Herein, we tested it for intracellular A*TPγS delivery to overcome the cell impermeability issue. Figure 1 shows a schematic illustration of the preparation procedure of A*TPγS-loaded LCP nanoparticles. A*TPγS is first loaded into the CaP core of the LCP nanoparticles, which is then coated with a DOPA layer. DOPA functions to prevent the aggregation of the CaP core because of the strong interaction between the CaP core and the phosphate head of DOPA. The cationic 1,2-dioleoyl-3-trimethylammonium-propane (DOTAP) lipid is then added as an outer layer onto the DOPA layer (2). Additionally, an outside layer of neutral dioleoylphosphatidylcholine (DOPC) with DSPE-PEG with a tethered targeting ligand anisamide (AA) is added. These layers keep the nanoparticles stable in the hydrophilic tissue culture medium and also enhance cellular uptake of the nanoparticles. Upon nanoparticle cellular internalization, the CaP core is rapidly dissolved in the endosome due to its low pH, leading to increased endosome osmotic pressure. Consequently, the endosome swells and ruptures, releasing the loaded A*TPγS into the cytoplasm. This quick process also avoids lysosomal degradation after endocytosis.

Figure 2 summarized the results of our characterization of synthesized nanoparticles. The loaded LCP nanoparticle sizes are exhibited in Figure 2A in a histogram. The sizes are also summarized in Figure 2B along with those of unloaded nanoparticles. On average, A*TPγS-loaded LCP nanoparticles were 83.7±8.1 nm and, as expected, about 11% larger than unloaded nanoparticles, which has a size of 75.1±5.2 nm. A transmission electron microscope (TEM) image is also shown in Figure 2C, illustrating the monodispersed nanoparticles with spherical shapes. The final concentration of A*TPγS in 1 mL of

produced nanoparticles was 0.5 mM. Thus, about 50% of the starting A*TPγS (100 µL of 10 mM solution) was loaded into the nanoparticles.

We also monitored intracellular uptake of the LCP nanoparticle by using the fluorescent ATP analog Alexa-Fluor-647-ATP. As described in Materials and Methods, this ATP analog was loaded into the LCP nanoparticles. The loaded nanoparticle was added into the tissue culture medium and incubated with the HCT116 cells for 4 hours for intracellular uptake. The cells were then nucleus-stained and observed under a confocal microscope. The result is shown in Figure 2D. Strong Alexa-Fluor-647 red fluorescence activities were spotted inside the HCT116 cells, demonstrating the cellular internalization of loaded LCP nanoparticles. Thus, we were convinced that the LCP nanoparticles could be used as a vehicle for intracellular A*TPγS delivery.

3.2 Mutated AKT1 kinase and kinase *in vivo* biological activity

Meanwhile, we generated other components of the Shokat chemical genetic method, in order to combine it with nanoparticle-mediated intracellular A*TPγS delivery to enable *in vivo* protein kinase substrate detection.

First, as brifly mentioned earlier, the AKT1 protein kinase was chosen as our initial prototype to develop an *in vivo* method for identifying protein kinase-substrates relationship. The AKT1 mutant with enlarged ATP binding pocket has been shown to bind to the bulky ATP analogue A*TP. The mutant retained protein kinase activity and exhibited the same ectopic expression level as the wild type AKT1 kinase (24,25). It was also reported by Dr. Shokat's group that ATPγS can be utilized by AKT1 to tag its substrates (26). In a word, multiple evidences support AKT1 as the most feasible choice of initial prototype. Moreover, the functional importance of AKT1 is illustrated quantitatively in Figure 3 in terms of the number of experimentally determined AKT1 subtrates. We downloaded the PhosphoSitePlus database (www.phosphosite.org) dataset and counted the number of unique experimentally verified substrates for each human protein kinases, revealing that human AKT1 has 215 unique substrates (27). As many other cell biology parameters (28-31), the substrate count follows the so-called scale-free distribution, *i.e.*, $P_{(k)} \propto K^{-\alpha}$, with $P_{(K)}$ as the number of protein kinases with K unique substrates and α a positive constants. This is illustrated in a linear relationship in a log-log histogram/plot ($\log_2(P_{(k)})$ vs. $\log_2(K)$) in figure 3. A small number of protein kinases have extra-ordinarily high substrate counts, while the majority of the kinases have only a small number of substrates. As shown by the red * symbol in figure 3, AKT1 is one of the high-substrate-count kinases; it was ranked, as shown in the insert table in figure 3, at the 7th among all human protein kinases. Therefore, AKT1 is not only the most feasible, but also a functionally important, choice for our study.

Thus, we acquired the pCDNA3-Myr-HA-AKT1 expression plasmid from AddGene. The vector expresses the AKT1 protein fused to the HA and the Myr tags. The HA-tag enables detection of the expressed AKT1 protein with the HA-tag antibody, facilitating transfection analysis. The Myr tag renders the expressed protein myristoylated, membrane anchored and, thus, contitutively active. We applied site-directed mutagenesis to mutate the gatekeeper Methionine codon into a Glycine codon in AKT1 ATP binding pocket (24). The two vectors made it possible for us to test whether the bulky A*TPγS molecule is too large for AKT1 and other wild type protein kinases, but fits into the enlarged ATP binding pocket of mutant AKT1 and leads to specific thiophosphorylation of AKT1 substrates.

Second, to test thio-phosphate-tagging of substrate proteins in live cells, the host cells were needed. The human HCT116 cell line were chosen. This cell line has an activating point mutation in the PI3KCA gene (32,33), resulting in constitutive PI3 kinase activity and thus elevated activation of the expressed AKT1 fusion proteins, further facilitating *in vivo* detection of thio-phosphate-tagging of AKT1 substrates.

Having established these components, we next tested the expression of both wildtype and mutant AKT1 proteins upon transfection of the respective expression plasmids into the HCT116 cells and subsequent treatments. 72 hours after the tranfection, the cells were treated with the A*TPγS-loaded nanoparticle for 4 hours. The cells were then lysed, and the cell lysates were treated with the PNBM alkylation to convert the thiophosphate tag into thiophosphate ester tag. We then performed western blot analysis with the HA-tag antibody to meaure the ectopic expression of wildtype and mutant AKT1 proteins. As shown in Figure 4A, both wild type and mutant AKT1 expression vectors drove strong expression of the HA-tagged proteins in the HCT116 cells upon transfection, while the control untransfected HCT116 cells exhibited no detectable HA-tagged AKT1 expression. The nanoparticle and the PNBM treatments did not affect the expression significantly.

3.3 *In vivo* utilization of bulky A*TPγS by the mutant AKT1 protein kinase

Next, we attempted to determine whether the mutant AKT1 protein can use the nanoparticle-delivered A*TPγS in its phosphorylation reaction, *i.e.*, transferring the γ-thiophosphate group onto AKT1 substrates (Figure 5). For this purpose, we tested potential AKT1 self-tagging through its autophosphorylation activity as well as tagging of another well-known AKT1 substrate, the IKKalpha protein. Since alkylation by p-Nitrobenzyl-mesylate (PNBM) should have converted the thiophosphate tag into the thiophosphate ester tag (34), the anti-thiophosphate ester monoclonal antibody should be able to recognize the tagged proteins. Thus, the treated cell lysates were then subjected to immunoprecipitation with the thiophosphate ester tag antibody. The immunoprecipitants are then examined by western blot analysis with specific antibodies to detect potential *in vivo* tagging of the corresponding proteins. The results are shown in figure 6. The HA-tag antibody was used in this analysis to detect self-tagging of the Myr-HA-AKT1 fusion proteins. As shown in Figure 6A, the fusion protein was detected in the immunoprecipitant of the cells transfected with the mutant AKT1 expression vector. Both nanoparticle treatment of the cells and PNBM alkylation of the cell lysate were required; the bands dispeared when either treatment was not performed. However, the fusion protein was not detected in the immunoprecipitants of the cells transfected with wild type AKT1 expression vector, even though the cell lysate was also treated with PNBM. Not surprisingly, the control untransfected cells gave no band due to a lack of expression of the HA-tagged fusion protein. The same was observed for the IKKalpha protein (Fig. 6B). IKKalpha was detected with an IKKalpha antibody in the immunoprecipitant of the cells transfected with the mutant AKT1 expression vector, but not in that of the cells transfected with wild type AKT1 expression vector. As expected, the antibody was able to detect endogenous IKKalpha protein in control untransfected HCT116 cells. Thus, the mutant AKT1 protein was able to use, exclusively, the nanoparticle delivered intracellular A*TPγS for *in vivo* thiophosphate-tagging of its substrates in live HCT116 cells.

AKT1 is known to phosphorylate a large number of substrates. As discussed earlier, a search of the PhosphoSitePlus database (www.phosphosite.org) revealed that human AKT1 has 215 unique human proteins as its experimentally verified substrates and was the 7[th] human kinase with most substrates (27). Thus, we next tried to determine whether the mutant AKT1 protein would use the delivered A*TPγS to tag many other proteins in the HCT116 cells. First, we tried western blot to detect the presence of tagged proteins in the cell lysate. Both wild type and mutated AKT1 expression vector transfected cell lysates were treated with PNBM, then followed by western blotting assay with the thiophosphate ester antibody. As shown in Figure 7, only the cells that were transfected with the mutant AKT1 expression vector and treated with A*TPγS-loaded nanoparticles and whose cell lysates were PNBM alkylated exhibited bands in the western blot analysis (Figure 7). Multiple prominent bands were detected, ranging from 25 to180 kDa. This is not surprising, as AKT1 is known to phosphorylate a large number of substrates. As

expected, wild type AKT1 was not able to use the bulky A*TPγS, although the cell lysate was also treated with PNBM. Once again, for cells transfected with the mutated AKT1 expression vector, both nanoparticle treatment of the cells and PNBM alkylation of the cell lysate were required; the bands dispeared when either treatment was not performed. Second, we performed immunoprecipitation of the PNBM alkylated lysates of cells transfected with expression vectors and treated with the nanoparticle. The immunoprecipitants were subjected to SDS-PAGE electrophoresis. The gel was then comassie blue stained, in order to compare the protein abundance in the two immunoprecipitants. As expected, the thiophosphate ester antibody was able to immnoprecipitate a large number of proteins from the lysate of mutant-AKT1-expressing cells, but not from that of wild-type-AKT1-expressing cells (Figure 8). Taken together, the results of the two experiments suggests that LCP-nanoparticle delivered intracellular A*TPγS was used exclusively by the mutant AKT1 protein to thiophosphate-tag its substrates in live cells. That is, by combining nanoparticle-mediated intracellular A*TPγS delivery and the Shokat chemical genetic method, an *in vivo* approach for kinase-substrate relationship determination has been established.

4. Discussion

The protein kinase study has to heavily rely on *in vitro* experimental protocols due to the cell impermeability of key analysis probes – the ATP analogs – and our inability to distinguish kinase reactions from one another in live cells. We set out to tackle this problem in this study. Nanoparticle-mediated intracellular delivery of A*TPγS overcomed the cell impermeability issue. Using the AKT1 protein kinase as the initial prototype, delivered A*TPγS enabled *in vivo* substrate thiophosphate tagging by mutant AKT1 with enlarged ATP binding pocket, which was created according to the Shokat chemical genetic method. Consequently, we were able to detect kinase-substrate relationship in live cells and in a kinase-specific manner.

The results pave the way to unbiased mass-spectrometry (MS) based proteomic study of kinase-substrate relationship. The AKT1 protein kinase, as discussed earlier, has a large number of substrates. Thus, as shown in figure 7, the thiophosphate ester antibody was able to immunoprecipitate a large number of tagged proteins from the lysate of HCT116 cells expressing the mutant AKT1 protein. Comparative MS analysis of the immunoprecipitant with that of wildtype-AKT1-expressing cells will lead to unbiased identification of both known and new AKT1 substrates.

Our approach should be applicable to many other protein kinases. Due to the conservation of the ATP binding pocket across the kinome, the Shokat chemical genetic method is widely applicable. It has been applied to a large number of human protein kinases; just to name a few, EGF receptor, SRC, p38/MAPK14 and ERK1/2. Sometimes, a different derivative of the ATP molecular was used. Consequently, the nanoparticle-mediated intracellular delivery of the bulky ATP analogs should be applicable in enabling *in vivo* analysis of these protein kinases as well.

The Shokat chemical genetic method has also been applied to other research areas. First, it has ben used to develop more specific protein kinase inhibitors. Many kinase inhibitors act by competing with ATP for binding to the target kinases. Since the ATP binding pocket are well conserved across the kinome, such kinase inhibitors would likely target multiple kinases. This issue was alleviated by modifying the inhibitor to fit it into the enlarged binding pocket of the mutant protein kinase. This technique has been applied to generate inhibitors that have been used to probe the functions of more that 70 protein kinases. Thus, whether our approach can be used to improve this application of the Shokat chemical genetic method via more efficient intracellular delivery of the inbibitors becomes an worthy future endeavor.

Second, the principle of the Shokat method has been applied to study other ATP binding proteins (35). The DDX3 gene codes for a DEAD-box protein and is frequently mutated in human cancers. The DDX3 protein belongs to a large family of ATP-dependent RNA chaperones. The ATP binding pockets of the DEAD-box proteins in this family are very similar to one another, making it challenging to develop DDX3-specific inhibitors. It was shown that mutating the hydrophobic amino acid residues, which interact with ATP N6 position in the binding pocket, enabled the mutant DDX3 protein to bind to bulky ATP analogs (35). Our strategy of nanoparticle-mediated intracellular delivery should be able to ensure efficient delivery of the bulky ATP analog based mutant-DDX3-specific inhibitors.

Moreover, it is tempting to investigate whether our approach can use additional derivatives of the A*TP (N6-Benzyl-ATP) molecule for *in vivo* probing of other aspects of protein kinase function. A*TPγS derives from A*TP by substituting the γ-phosphate group with a thiophosphate group. The purpose of the thiophosphate group is to act as a handle for kinase substrate identification. However, there are other important aspects of kinase study, *e.g.*, ATP binding and enzymatic activities. The ATP binding pocket is usually shielded from ATP binding in an inactive protein kinase, but becomes accessible upon activation of the kinase. For instance, in monomeric inactive EGF receptor (EGFR), the activation loop is in close proximity of the pocket and blocks ATP bnding. Upon ligand binding and dimerization, the C-lobe of one EGFR molecule disrupt this autoinhibitory conformation of the other EGFR molecule, enabling ATP binding and kinase activation (36). However, current ATP binding and kinase activity analyses rely on the usage of ATP analogs, and cell impermeability of these analogs renders *in vivo* analyses impossible. Using our approach for intracellular delivery of relevant A*TP derivatives should help to alleviate this cell impermeability obstacle.

Nanoparticles have been widely used for therapeutic agent delivery in biomedical researches. For instance, the LCP nanoparticle has been used to deliver gene therapy agents such as siRNA and expression vectors (37,38). It has also been used to deliver many therapeutical chemicals such as gemcitabine triphosphate (2,14), doxorubicin and paclitaxel (39). Due the similarity between A*TPγS and gemcitabine triphosphate, we choosed to use this nanoparticle for intracellular A*TPγS delivery in this study. To the best of our knowledge, this represents a new application of nanoparticles – overcoming cell impeameability of key analysis probes to enable *in vivo* execution of previously *in vitro* experimental procedures. The principle should be applicable to a wide variety of biochemical, molecular and cellular research techniques, perhaps adopting difference types of nanoparticles for different purposes. Tremoudous previously impossible research opportunities are to be opened up.

In summary, in this study, we ventured our first step towards *in vivo* protein kinase analysis by successfully combining nanoparticle-mediated probe (A*TPγS) delivery and the Shokat chemical genetic method. The former overcomed the A*TPγS cell impermeability issue, and the latter enabled differentiation of the kinase reaction of interest from those of other protein kinases. Using the AKT1 as our initial prototype, we achieved detection of kinase-substrate relationships in live cells. It is our belief that this approach can be applied to other protein kinases and other aspects of protein kinase studies. Hopefully, the principle – using nanoparticle-mediated delivery of key analysis probes to overcome cell impermeability – will be generally applicable to many other biochemical, molecular and cellular biomedical research areas.

5. Conclusion
Identification of the *in vivo* substrates of specific protein kinases in intact cells offers a great understanding of intracellular signaling pathways and benefits therapeutic targets for human diseases (40). Herein, we have developed a strategy to exclusively tag the direct substrates of a protein kinase of interest in intact cells, combining the chemical genetics method and the nanoparticle delivery system to uniquely transfer γ-thiophosphorylation tagging to AKT1 substates in intact cells. Specifically, we confirmed our strategy worked on previously known AKT1 substrate using immunoprecipitation and western blotting

assays. Finally, this *in vivo* substrate identification strategy can identify both known and novel substrates, offers a novel route to identify substrates of various kinases, and can be applied to the signaling cascade study.

5. Acknowledgments

This work was supported by National Institutes of Health (NIH) grants R15GM122006 to DW and U54 CA198999 to LH.

6. References

REFERENCES

1. Dancey, J.E., Bedard, P.L., Onetto, N. and Hudson, T.J. (2012) The genetic basis for cancer treatment decisions. *Cell*, **148**, 409-420.
2. Zhang, Y., Peng, L., Mumper, R.J. and Huang, L. (2013) Combinational delivery of c-myc siRNA and nucleoside analogs in a single, synthetic nanocarrier for targeted cancer therapy. *Biomaterials*, **34**, 8459-8468.
3. Chang, L. and Karin, M. (2001) Mammalian MAP kinase signalling cascades. *Nature*, **410**, 37-40.
4. Zhang, H., Berezov, A., Wang, Q., Zhang, G., Drebin, J., Murali, R. and Greene, M.I. (2007) ErbB receptors: from oncogenes to targeted cancer therapies. *J Clin Invest*, **117**, 2051-2058.
5. Ulrich, S.M., Kenski, D.M. and Shokat, K.M. (2003) Engineering a "methionine clamp" into Src family kinases enhances specificity toward unnatural ATP analogues. *Biochemistry*, **42**, 7915-7921.
6. Kumar, V., Weng, Y.-C., Geldenhuys, W.J., Wang, D., Han, X., Messing, R.O. and Chou, W.-H. (2015) Generation and characterization of ATP analog-specific protein kinase Cδ. *Journal of Biological Chemistry*, **290**, 1936-1951.
7. Zeliadt, N. (2012) Profile of Kevan M. Shokat. *Proceedings of the National Academy of Sciences*, **109**, 11057-11059.
8. Ultanir, S.K., Yadav, S., Hertz, N.T., Oses-Prieto, J.A., Claxton, S., Burlingame, A.L., Shokat, K.M., Jan, L.Y. and Jan, Y.N. (2014) MST3 kinase phosphorylates TAO1/2 to enable Myosin Va function in promoting spine synapse development. *Neuron*, **84**, 968-982.
9. Allen, J.J., Lazerwith, S.E. and Shokat, K.M. (2005) Bio-orthogonal affinity purification of direct kinase substrates. *Journal of the American Chemical Society*, **127**, 5288-5289.
10. Shah, K. and Shokat, K.M. (2002) A chemical genetic screen for direct v-Src substrates reveals ordered assembly of a retrograde signaling pathway. *Chem Biol*, **9**, 35-47.
11. Hertz, N.T., Wang, B.T., Allen, J.J., Zhang, C., Dar, A.C., Burlingame, A.L. and Shokat, K.M. (2010) Chemical genetic approach for kinase-substrate mapping by covalent capture of thiophosphopeptides and analysis by mass spectrometry. *Curr Protoc Chem Biol*, **2**, 15-36.
12. Dephoure, N., Howson, R.W., Blethrow, J.D., Shokat, K.M. and O'Shea, E.K. (2005) Combining chemical genetics and proteomics to identify protein kinase substrates. *Proceedings of the National Academy of Sciences*, **102**, 17940-17945.
13. Hindley, A.D., Park, S., Wang, L., Shah, K., Wang, Y., Hu, X., Shokat, K.M., Kolch, W., Sedivy, J.M. and Yeung, K.C. (2004) Engineering the serine/threonine protein kinase Raf-1 to utilise an orthogonal analogue of ATP substituted at the N 6 position. *FEBS letters*, **556**, 26-34.
14. Zhang, Y., Kim, W.Y. and Huang, L. (2013) Systemic delivery of gemcitabine triphosphate via LCP nanoparticles for NSCLC and pancreatic cancer therapy. *Biomaterials*, **34**, 3447-3458.
15. Altomare, D.A. and Testa, J.R. (2005) Perturbations of the AKT signaling pathway in human cancer. *Oncogene*, **24**, 7455-7464.
16. Testa, J.R. and Tsichlis, P.N. (2005) AKT signaling in normal and malignant cells. *Oncogene*, **24**, 7391-7393.
17. Dan, H.C., Cooper, M.J., Cogswell, P.C., Duncan, J.A., Ting, J.P.-Y. and Baldwin, A.S. (2008) Akt-dependent regulation of NF-κB is controlled by mTOR and Raptor in association with IKK. *Genes & development*, **22**, 1490-1500.

18. Manning, B.D. and Cantley, L.C. (2007) AKT/PKB signaling: navigating downstream. *Cell*, **129**, 1261-1274.
19. Engelman, J.A., Luo, J. and Cantley, L.C. (2006) The evolution of phosphatidylinositol 3-kinases as regulators of growth and metabolism. *Nat Rev Genet*, **7**, 606-619.
20. Li, X., Lu, Y., Jin, W., Liang, K., Mills, G.B. and Fan, Z. (2006) Autophosphorylation of Akt at Threonine 72 and Serine 246 A potential mechanism of regulation of Akt kinase activity. *Journal of Biological Chemistry*, **281**, 13837-13843.
21. Li, Y., Wu, Y., Huang, L., Miao, L., Zhou, J., Satterlee, A.B. and Yao, J. (2016) Sigma receptor-mediated targeted delivery of anti-angiogenic multifunctional nanodrugs for combination tumor therapy. *Journal of Controlled Release*, **228**, 107-119.
22. Brattain, M.G., Marks, M.E., McCombs, J., Finely, W. and Brattain, D.E. (1983) Characterization of human colon carcinoma cell lines isolated from a single primary tumour. *Br J Cancer*, **47**, 373-381.
23. Howell, G.M., Humphrey, L.E., Awwad, R.A., Wang, D., Koterba, A., Periyasamy, B., Yang, J., Li, W., Willson, J.K., Ziober, B.L. *et al.* (1998) Aberrant regulation of transforming growth factor-alpha during the establishment of growth arrest and quiescence of growth factor independent cells. *J Biol Chem*, **273**, 9214-9223.
24. Okuzumi, T., Ducker, G.S., Zhang, C., Aizenstein, B., Hoffman, R. and Shokat, K.M. (2010) Synthesis and evaluation of indazole based analog sensitive Akt inhibitors. *Mol Biosyst*, **6**, 1389-1402.
25. Okuzumi, T., Fiedler, D., Zhang, C., Gray, D.C., Aizenstein, B., Hoffman, R. and Shokat, K.M. (2009) Inhibitor hijacking of Akt activation. *Nat Chem Biol*, **5**, 484-493.
26. Allen, J.J., Li, M., Brinkworth, C.S., Paulson, J.L., Wang, D., Hübner, A., Chou, W.-H., Davis, R.J., Burlingame, A.L. and Messing, R.O. (2007) A semisynthetic epitope for kinase substrates. *Nature methods*, **4**, 511.
27. Hornbeck, P.V., Zhang, B., Murray, B., Kornhauser, J.M., Latham, V. and Skrzypek, E. (2015) PhosphoSitePlus, 2014: mutations, PTMs and recalibrations. *Nucleic Acids Res*, **43**, D512-520.
28. Guo, Z., Jiang, W., Lages, N., Borcherds, W. and Wang, D. (2014) Relationship between gene duplicability and diversifiability in the topology of biochemical networks. *BMC Genomics*, **15**, 577.
29. Padawer, T., Leighty, R.E. and Wang, D. (2012) Duplicate gene enrichment and expression pattern diversification in multicellularity. *Nucleic Acids Res*, **40**, 7597-7605.
30. Wang, D., Wang, T., Gill, A., Hilliard, T., Chen, F., Karamyshev, A.L. and Zhang, F. (2020) Uncovering the cellular capacity for intensive and specific feedback self-control of the argonautes and MicroRNA targeting activity. *Nucleic Acids Res*.
31. Zhang, F. and Wang, D. (2017) The Pattern of microRNA Binding Site Distribution. *Genes (Basel)*, **8**.
32. Guo, X.N., Rajput, A., Rose, R., Hauser, J., Beko, A., Kuropatwinski, K., LeVea, C., Hoffman, R.M., Brattain, M.G. and Wang, J. (2007) Mutant PIK3CA-bearing colon cancer cells display increased metastasis in an orthotopic model. *Cancer Res*, **67**, 5851-5858.
33. Kim, A., Lee, J.E., Lee, S.S., Kim, C., Lee, S.J., Jang, W.S. and Park, S. (2013) Coexistent mutations of KRAS and PIK3CA affect the efficacy of NVP-BEZ235, a dual PI3K/MTOR inhibitor, in regulating the PI3K/MTOR pathway in colorectal cancer. *Int J Cancer*, **133**, 984-996.
34. Blethrow, J.D., Glavy, J.S., Morgan, D.O. and Shokat, K.M. (2008) Covalent capture of kinase-specific phosphopeptides reveals Cdk1-cyclin B substrates. *Proceedings of the National Academy of Sciences*, **105**, 1442-1447.
35. Floor, S.N., Barkovich, K.J., Condon, K.J., Shokat, K.M. and Doudna, J.A. (2016) Analog sensitive chemical inhibition of the DEAD-box protein DDX3. *Protein Sci*, **25**, 638-649.

36. Ferguson, K.M. (2008) Structure-based view of epidermal growth factor receptor regulation. *Annu Rev Biophys*, **37**, 353-373.
37. Liu, Q., Zhu, H., Tiruthani, K., Shen, L., Chen, F., Gao, K., Zhang, X., Hou, L., Wang, D., Liu, R. *et al.* (2018) Nanoparticle-Mediated Trapping of Wnt Family Member 5A in Tumor Microenvironments Enhances Immunotherapy for B-Raf Proto-Oncogene Mutant Melanoma. *ACS Nano*, **12**, 1250-1261.
38. Yang, Y., Li, J., Liu, F. and Huang, L. (2012) Systemic delivery of siRNA via LCP nanoparticle efficiently inhibits lung metastasis. *Mol Ther*, **20**, 609-615.
39. Wu, C., Xu, J., Hao, Y., Zhao, Y., Qiu, Y., Jiang, J., Yu, T., Ji, P. and Liu, Y. (2017) Application of a lipid-coated hollow calcium phosphate nanoparticle in synergistic co-delivery of doxorubicin and paclitaxel for the treatment of human lung cancer A549 cells. *Int J Nanomedicine*, **12**, 7979-7992.
40. Manning, G., Whyte, D.B., Martinez, R., Hunter, T. and Sudarsanam, S. (2002) The protein kinase complement of the human genome. *Science*, **298**, 1912-1934.

Figure Legends

Figure 1. Nanoparticle formulation (A) Chemical structures of DOTAP, DOPA, DSPE-PEG and cholesterol. (B) Schematic illustration of the formulation procedure of LCP loaded A*TPγS LCP nanoparticles.

Figure 2. Characterization of the LCP nanoparticles. (A) Histogram of nanoparticle size. (B) Average sizes of blank LCP nanoparticles and A*TPγS-loaded LCP nanoparticles. (C) TEM images of A*TPγS-loaded LCP nanoparticles, n=6. Scale bar=X. And (D) Internalization of Alexa 647 LCP nanoparticles in HCT116 cells, n=4. Scale bar=20.0 μm.

Figure 3. High substrate count of Akt1. A log-log histogram of substrate count depicts the scale-free distribution. The red * symbol and text illustrate the high Akt1 substrate count. The insert table list top 10 high-substrate-count kinase, with AKT1 in bolded text.

Figure 4. Western blotting analysis of transfected HCT116 cells using HA tag antibody either by wild type or mutated AKT1 kinase plasmid. (B) Transfection and western blot analyses scheme.

Figure 5. Thiophosphate recoganization scheme. (A) Wild type kinase utilizes normal ATP to phosphorylate its substrates. In contrast, the mutated kinase, with enlarged ATP bind pocket, accepts bulky A*TPγS and was able to transfer the thiophosphate group to its substrates. (B) Reaction sequence for affinity labelling kinase substrates.

Figure 6. Western blotting analysis of thiophosphorylated AKT1 kinase substrates using (A) IKKα and (B) HA tag protein. (C) Thiophosphorylated kinase substrates detecting scheme. Immunoprecipitation of thiophosphate protein and validation of previously known substrates by Western blot analysis.

Figure 7. Western blotting of thiophosphorylated proteins. (A) Either wild type or mutated AKT1 kinase plasmid were transfected into HCT116 cells, intracellular incubated with A*TPγS nanoparticles, alkylated by PNBM. The thiophosphorylated substrate proteins were visualized by western blotting using an thiophosphate ester rabbit monoclonal antibody that dected thiophosphate esters.

Figure 8. Blue silver comassie stained SDS-PAGE gel. The wells for the molecular weight markers (MW), the immunoprecipitant of wild type AKT1 expression plasmid transfected cell lysate (Wild Type) and the immunoprecipitant of mutant AKT1 expression plasmid transfected cell lysate (Mutant) are marked. The antibody (IgG) heavy and light chains are also marked.

Figure 1

Figure 2

Nanoparticles	Size （nm）
Blank LCP	75.1±5.2
A*TPγS LCP	83.7±8.1

		Size (d.n...	% Intensity:	St Dev (d.n...
Z-Average (d.nm): 82.49	**Peak 1:**	99.21	100.0	35.46
Pdl: 0.157	**Peak 2:**	0.000	0.0	0.000
Intercept: 0.965	**Peak 3:**	0.000	0.0	0.000
Result quality Good				

Figure 3

Figure 4

A

C   WT   Mutated Akt1

63 kDa

+   -   -   +   +   A*TPγS NPs
+   -   +   -   +   PNBM

B

Plasmid

Cell lysates

Western blot using HA tag antibody

Figure 5

ATP

A*TPγS

Kinase substrate — Phosphorylated kinase substrate

Wild type kinase

Kinase substrate — Thiophosphorylated kinase substrate

Mutant kinase

NPs internalization

Inner cell

phosphorylation

A*TPγS

A*TPγS NPs incubation in transfected cells

Thiophosphate ester rabbit monoclonal antibody

PNBM

Cell lysates

Figure 6

A

C WT   Mutant Akt1

100 kDa

75 kDa

| + | - | - | + | + | A*TPγS NPs |
| + | - | + | - | + | PNBM |

B

C WT   Mutant Akt1

63 kDa

| + | - | - | + | + | A*TPγS NPs |
| + | - | + | - | + | PNBM |

C

Cell lysates

Thiophosphate ester rabbit
monoclonal antibody

Immunoprecipitation of
thiophosphate protein

Wash and elution

Western blot analysis using previously known substrates

Figure 7

Figure 8