TITLE

Slowly evolving dopaminergic activity modulates the moment-to-moment probability of

movement initiation.

AUTHOR NAMES

Allison E. Hamilos[1*], Giulia Spedicato[1], Ye Hong[1], Fangmiao Sun[2], Yulong Li[2] &

John A. Assad[1,3*]

AFFILIATIONS

[1]Department of Neurobiology, Harvard Medical School, Boston, Massachusetts, 02115, USA.

[2]State Key Laboratory of Membrane Biology, Peking University School of Life Sciences,

Beijing, 100871, P.R. China.

[3]Istituto Italiano di Tecnologia, Genova, Italy.

*For correspondence: A.H. (ahamilos@g.harvard.edu) or J.A.A. (jassad@hms.harvard.edu).

ABSTRACT

Clues from human movement disorders have long suggested that the neurotransmitter dopamine

plays a key role in motor control, but how the endogenous dopaminergic system regulates

movement is unknown. Here we show dynamic dopaminergic signaling over seconds-long

timescales controls movement timing in mice. Animals were trained to initiate licking after a self-

timed interval following a start-timing cue. The movement time was variable from trial-to-trial, as

expected from previous studies. Surprisingly, dopaminergic signals ramped-up over seconds

between the start-timing cue and the self-timed movement, with variable dynamics that predicted

the movement time on single trials. Steeply rising signals preceded early lick-initiation, whereas

slowly rising signals preceded later initiation. Higher baseline signals also predicted earlier self-

timed movements. Optogenetic activation of dopamine neurons during self-timing did not trigger

immediate movements, but rather caused systematic early-shifting of movement initiation,

whereas inhibition caused late-shifting, as if modulating the probability of movement. Consistent

with this view, the dynamics of the endogenous dopaminergic signals quantitatively predicted the

moment-by-moment probability of movement initiation on single trials. These results reveal a

causal role for dynamic dopaminergic signaling unfolding over seconds in modulating the decision

of when to move.

INTRODUCTION

What makes us move? Empirically, a few hundred milliseconds before movement, thousands of

neurons in the motor system suddenly become active in concert, and this neural activity is relayed

via spinal and brainstem neurons to recruit muscle fibers that power movement (*Shenoy et al.,*

2013). Yet just before this period of intense neuronal activity, the motor system is largely quiescent.

How does the brain suddenly and profoundly rouse motor neurons into the coordinated action

needed to trigger movement?

In the case of movements made in reaction to external stimuli, activity evoked first in sensory brain

areas is presumably passed along to appropriate motor centers to trigger this coordinated neural

activity, thereby leading to movement. But humans and animals can also self-initiate movement

without overt, external input (*Deecke, 1996; Hallett, 2007; Lee and Assad, 2003; Romo et al.,*

1992). For example, while reading this page, you may decide without prompting to reach for your

coffee. In that case, the movement cannot be clearly related to an abrupt, conspicuous sensory cue.

What "went off" in your brain that made you reach for your coffee at this *particular* moment, as

opposed to a moment earlier or later?

Human movement disorders may provide clues to this mystery. Patients and animal models of

Parkinson's Disease experience difficulty self-initiating movements, exemplified by perseveration

(*Hughes et al., 2013*), trouble initiating steps when walking (*Bloxham et al., 1984*), and problems

timing movements (*Malapani et al., 1998; Meck, 1986, 2006; Mikhael and Gershman, 2019*). In

contrast to these self-generated actions, externally cued reactions are often less severely affected

in Parkinson's, a phenomenon sometimes referred to as "paradoxical kinesia" (*Barthel et al., 2018;*

Bloxham et al., 1984). For example, patients' gait can be normalized by walking aids that prompt

steps in reaction to visual cues displayed on the ground (*Barthel et al., 2018*).

Because the underlying neuropathophysiology of Parkinson's includes the loss of dopaminergic

neurons (DANs), the symptomatology of Parkinson's suggests DAN activity plays an important

role in deciding when to self-initiate movement. Indeed, pharmacological manipulations of the

neurotransmitter dopamine causally and bidirectionally influence movement timing (*Dews and*

Morse, 1958; Lustig and Meck, 2005; Meck, 1986; Mikhael and Gershman, 2019; Schuster and

Zimmerman, 1961). This can be demonstrated in the context of *self-timed* movement tasks, in

which subjects reproduce a target-timing interval by making a movement following a self-timed

delay that is referenced to a start-timing cue (*Malapani et al., 1998*). Species across the animal

kingdom, from rodents and birds to primates, can learn these tasks and produce self-timed

movements that occur, on average, at about the target time, although the exact timing exhibits

considerable variability from trial-to-trial (*Gallistel and Gibbon, 2000; Meck, 2006; Mello et al.,*

2015; Merchant et al., 2013; Rakitin et al., 1998; Remington et al., 2018; Schuster and

Zimmerman, 1961; Sohn et al., 2019; Wang et al., 2018). In such self-timed movement tasks,

decreased dopamine availability/efficacy (*e.g.*, Parkinson's, neuroleptic drugs) generally produces

late-shifted movements (*Malapani et al., 1998; Meck, 1986, 2006; Merchant et al., 2013*),

whereas high dopamine conditions (*e.g.*, amphetamines) produce early-shifting (*Dews and Morse,*

1958; Schuster and Zimmerman, 1961).

Although exogenous dopamine manipulations can influence timing behavior, it remains unknown

whether endogenous DAN activity is involved in determining when to move. DANs densely

innervate the striatum, where they modulate the activity of spiny projection neurons of the direct

and indirect pathways, which are thought to exert a push-pull influence on movement centers

(*Albin et al., 1989; DeLong, 1990; Freeze et al., 2013; Grillner and Robertson, 2016*). Moreover,

phasic bursts of dopaminergic activity have been observed just prior to movement onset (within

~500 ms) (*Coddington and Dudman, 2018, 2019; da Silva et al., 2018; Dodson et al., 2016;*

Howe and Dombeck, 2016; Wang and Tsien, 2011), and dopaminergic signals have been reported

to reflect more general encoding of movement kinematics (*Barter et al., 2015; Engelhard et al.,*

2019; Parker et al., 2016). However, optogenetic activation of dopamine neurons—within

physiological range—does not elicit immediate movements (*Coddington and Dudman, 2018,*

2019). We hypothesized that rather than overtly triggering movements, the ongoing activity of

nigrostriatal DANs could influence movement initiation over longer timescales by controlling or

modulating the moment-by-moment decision of *when* to execute a planned movement.

To test this hypothesis, we trained mice to make a movement (lick) after a self-timed interval

following a start-timing cue. The mice learned the timed interval, but, as observed in other species,

the exact timing of movement was highly variable from trial-to-trial, spanning seconds. We

exploited this inherent variability by examining how moment-to-moment nigrostriatal DAN

signals differed when animals decided to move relatively early versus late. We found that

dopaminergic signals "ramped up" during the timing interval, with variable dynamics that were

highly predictive of trial-by-trial movement timing, even seconds before the movement occurred.

Optogenetic DAN manipulation during the timing interval produced bidirectional changes in the

probability of movement timing, with activation causing a bias toward earlier self-timed

movements and suppression causing a bias toward later self-timed movements. These combined

observations suggest a novel role for the dopaminergic system in movement initiation, wherein

slowly evolving signals modulate the moment-to-moment probability of whether a movement will

occur.

RESULTS

We trained head-fixed mice to make self-timed movements to receive juice rewards (*Figure 1A*).

Animals received an audio/visual start-timing cue and then had to decide when to first-lick in the

absence of further cues. Animals only received juice if they waited a proscribed interval following

the cue before making their first-lick (>3.3 s in most experiments). As expected from previous

studies, the distribution of first-lick timing was broadly distributed over several seconds, and

exhibited the canonical scalar property of timing, as described by Weber's Law (*Figure 1B* and

Figure 1—figure supplement 1A-B; (*Gallistel and Gibbon, 2000*)). We note this variability in

timing was not imposed on the animal by training it to reproduce a variety of target intervals (*e.g.*,

2 *vs*. 5 s), but is rather a natural consequence of timing behavior, even for a single target interval.

Our main objective was to exploit the inherent variability in self-timed behavior to examine how

differences in neural activity might relate to variability in movement timing. Nonetheless, the

trained animals well-understood the timing contingencies of the task. In self-timed movement tasks

in which a *single* movement is used to assess timing, the distributions of movement times (in both

rodents and monkeys) tend to anticipate the target interval, even at the expense of reward on many

trials *(Eckard and Kyonka, 2018; Kirshenbaum et al., 2008; Lee and Assad, 2003)*. In these

paradigms, however, once a movement occurs, it removes future opportunities to move, which

creates premature "bias" in the raw timing distributions (*Anger, 1956*). To correct this bias,

movement times must be normalized by the (ever-diminishing) number of opportunities to move

at each timepoint (*Jaldow et al., 1990*). This yields the hazard function (the conditional probability

of movement given that movement has not already occurred, as a function of time), which is

equivalent to the instantaneous probability of movement. For example, on the first day of training,

our animals displayed fairly flat hazard functions, indicating a uniform instantaneous probability

of movement over time—*i.e.*, the animals did not yet understand the timing contingency

(*Figure 1C-D*). However, after training, the hazard function for our animals peaked near the target

time (either 3.3 or 5 s), suggesting an accurate latent timing process reflected in the instantaneous

movement probability (*Figure 1E*). Mice trained on a variant of the self-timed movement task

without lamp-off/on events showed no systematic differences in their timing distributions

(*Figure 1—figure supplement 1C*), suggesting that the mice referenced their timing to the start-

timing cue rather than the lamp-off event.

When mice were fully trained, we employed fiber photometry to record the activity of genetically-defined DANs expressing the calcium-sensitive fluorophore GCaMP6f (12 mice, substantia nigra pars compacta (SNc); *Figure 1—figure supplement 2*). We controlled for mechanical/optical artifacts by simultaneously recording fluorescence modulation of a co-expressed, calcium-insensitive fluorophore, tdTomato. We also recorded bodily movements with neck-muscle EMG, high-speed video, and a back-mounted accelerometer.

DAN signals ramp up slowly between the start-timing cue and self-timed movement

DAN GCaMP6f fluorescence typically exhibited brief transients following cue onset and immediately before first-lick onset (*Figure 2A*), as observed in previous studies (*Coddington and Dudman, 2018; da Silva et al., 2018; Dodson et al., 2016; Howe and Dombeck, 2016; Schultz et al., 1997*). However, during the timed interval, we observed slow "ramping up" of fluorescence over seconds, with a minimum after the cue-aligned transient and maximum just before the lick-related transient. The relatively fast intrinsic decay kinetics of GCaMP6f ($t_{1/2}$ <100 ms at 37°; (*Helassa et al., 2016*)) should not produce appreciable signal integration over the seconds-long timescales of the ramps we observed.

We asked whether this ramping differed between trials in which the animal moved relatively early or late. Strikingly, when we averaged signals pooled by movement time, we observed systematic differences in the steepness of ramping that were highly predictive of movement timing (*Figure 2B-C*). Trials with early first-licks exhibited steep ramping, whereas trials with later first-licks started from lower fluorescence levels and rose more slowly toward the time of movement.

The fluorescence ramps terminated at nearly the same amplitude, regardless of the movement time.

Ramping dynamics were not evident in control tdTomato signals (*Figure 2C*), indicating that the

ramping in the GCaMP6f signals was not an optical artifact. The quantitative relationship between

GCaMP6f dynamics and movement time will be addressed in a subsequent section of this paper.

Higher pre-cue DAN signals are correlated with earlier self-timed movements

In addition to ramping dynamics, average DAN GCaMP6f signals were correlated with first-lick

timing even before cue-onset (the "baseline" period, between lamp-off to cue event), with higher

baseline fluorescence predicting earlier first-licks (Pearson's r=-0.89, n=12 mice; *Figure 2B-C*).

Because dF/F correction methods can potentially distort baseline measurements, we rigorously

tested and validated three different dF/F methods, and we also repeated analyses with raw

fluorescence values compared between pairs of sequential trials with different movement times

(*Figure 2—figure supplement 1*; see *Methods*). All reported results, including the systematic

baseline differences, were robust to dF/F correction.

In principle, the amplitude of the baseline signal on a given trial *n* could be related to the outcome

of the previous trial or the animal's behavior during the baseline interval. To test this, we

performed 4-way ANOVA on the baseline signal (averaged for each trial between lamp-off and

the start-timing cue, n=12 mice), including factors 1) outcome of the previous (*n*-1th) trial

(rewarded or unrewarded); 2) presence or absence of spontaneous licking during the baseline

period; 3) upcoming movement time on trial *n* (categorized as <3.3 s or >3.3 s, to provide a simple

binary proxy for movement time); and 4) session number (to account for signal variability across

animals and daily sessions). Although previous trial outcome and baseline-licking were significant

predictors of baseline amplitude (p<0.01 for both), the upcoming movement time had a significant

independent (main) effect ($p<10^{-5}$). This raises the possibility of an additional source of variance

in baseline dopaminergic activity that is independent from previous trial events, but potentially

influences the upcoming movement time on that trial.

Movement timing-related ramping dynamics in other dopaminergic areas and striatal

dopamine release

We found similar ramping dynamics in SNc DAN axon terminals in the dorsolateral striatum (DLS;

Figure 2—figure supplement 2A-B) at a location involved in goal-directed licking behavior

(*Sippy et al., 2015*). Ramping was also present in GCaMP6f-expressing DAN cell bodies in the

ventral tegmental area (VTA, *Figure 2—figure supplement 2C*), reminiscent of mesolimbic

ramping signals described in goal-oriented navigation tasks (*Howe et al., 2013; Kim et al., 2019*).

To determine if these movement-timing-related signals are available to downstream targets that

may be involved in movement initiation, we monitored dopamine release in the DLS with two

complementary florescent dopamine sensors (dLight1.1 and DA_{2m}) expressed broadly in striatal

cells (*Figure 3 and Figure 2—figure supplement 2D-E*). The decay kinetics of the two

extracellular dopamine sensors differ somewhat (*Patriarchi et al., 2018; Sun et al., 2020*), which

we confirmed (dLight1.1 $t_{1/2}\sim75$ ms, DA_{2m} $t_{1/2}\sim125$ ms; *Figure 3—figure supplement 1*), yet both

revealed similar timing-related ramping dynamics on average (*Figure 3 inset*). These combined

data argue that the seconds-long dopaminergic ramping signals were not artifacts of sluggish

temporal responses of the various fluorescent sensors and were ultimately expressed as ramp-like

increases in dopamine release in the striatum.

First-lick timing-predictive DAN signals are not explained by ongoing body movements

The systematic ramping dynamics and baseline differences were not observed in the tdTomato

optical control channel nor in any of the other movement-control channels, at least on average

(*Figure 4*), making it unlikely that ramping dynamics could have arisen from optical artifacts.

Nevertheless, because DANs show transient responses to salient cues and movements

(*Coddington and Dudman, 2018; da Silva et al., 2018; Dodson et al., 2016; Howe and Dombeck,*

2016; Schultz et al., 1997), it is possible that fluorescence signals could reflect the superposition

of dopaminergic responses to multiple task events, including the cue, lick, ongoing spurious body

movements, and hidden cognitive processes like timing. For example, accelerating spurious

movements could, in principle, produce motor-related neural activity that "ramps up" during the

timed interval, perhaps even at different rates on different trials.

We thus derived a nested generalized linear encoding model of single-trial GCaMP6f signals

(*Engelhard et al., 2019; Park et al., 2014; Runyan et al., 2017*), a data-driven, statistical approach

designed to isolate and quantify the contributions of task events (timing-independent predictors)

from processes predictive of movement timing (timing-dependent predictors; *Figure 5A-B* and

Figure 5—figure supplement 1A-D). The model robustly detected task-event GCaMP6f kernels

locked to cue, lick and EMG/accelerometer events, but these timing-independent predictors alone

were insufficient to capture the rich variability of GCaMP6f signals for trials with different first-

lick times, especially the timing-dependent ramp-slope and baseline offset (n=12 mice, *Figure 5C*

and *Figure 5—figure supplement 1E-G*). In contrast, two timing-dependent predictors robustly

improved the model: 1) a baseline offset with amplitude linearly proportional to first-lick time;

and 2) a "stretch" feature representing percentages of the timed interval (*Figure 5B-C* and *Figure*

5—figure supplement 1E). The baseline offset term fit a baseline level inversely proportional to

movement time, and the temporal stretch feature predicted a ramping dynamic from the time of

the cue up to the first-lick, whose slope was inversely proportional to first-lick time. Similar results

were obtained for SNc DAN axon terminals in the DLS, VTA DAN cell bodies, and extracellular

striatal dopamine release (*Figure 5—figure supplement 1H*).

We note that the stretch feature of this GLM makes no assumptions about the underlying shape of

the dopaminergic signal; it only encodes percentages of timing intervals to allow for temporal

"expansion" or "contraction" to fit whatever shape(s) were present in the data. In particular, the

stretch feature cannot produce ramping unless ramping is present in the signal *and* temporally

scales with the length of the interval. Because this feature empirically found a ramp (although not

constrained to do so), the stretch aspect indicated that the underlying ramping process took place

at different rates for trials with different movement times, at least on average.

In contrast to the GCaMP6f model, when the same GLM was applied to the tdTomato control

signal, the timing-independent predictors (which could potentially cause optical/mechanical

artifacts—cue onset, first-lick, EMG/accelerometer) improved the model, but timing-dependent

predictors did not (*Figure 5C* and *Figure 5—figure supplement 1F-H*). In addition, separate

principal component (PC) analysis revealed ramp-like and baseline-offset-like components that

explained as much as 93% of the variance in DAN signals during the timing interval

(mean: 66%, range: 16-93%), but similar PCs were not present when tdTomato control signals

were analyzed with PCA (mean variance explained: 4%, range: 1.6-15%, *Figure 5—figure*

supplement 2).

Single-trial DAN ramping and baseline signals predict movement timing

Given that ramping and baseline-offset signals were not explained by nuisance movements or

optical artifacts, we asked whether DAN GCaMP6f fluorescence could predict first-lick timing on

single trials. Using a simple threshold-crossing decoding model (*Maimon and Assad, 2006*), we

found that single-trial GCaMP6f signals were predictive of first-lick time even for low thresholds

intersecting the "base" of the ramp, with the predictive value of the model progressively improving

for higher thresholds (n=12 mice; mean R^2 low: 0.54, mid: 0.71, high: 0.82; analysis for one mouse

shown in *Figure 6A*). We will return to this observation in more detail in the upcoming section on

single-trial dynamics (see below).

To more thoroughly determine the independent, additional predictive power of DAN baseline and

ramping signals over other task variables (*e.g.*, previous trial first-lick time and reward outcome,

etc.), we derived a nested decoding model for first-lick time (*Figure 6A*). All predictors

contributed to the predictive power of the model. However, even when we accounted for the

contributions of prior trial history, tdTomato artifacts and baseline GCaMP6f signals, GCaMP6f

threshold-crossing time robustly dominated the model and absorbed much of the variance

explained by baseline dopaminergic signals, alone explaining 10% of the variance in first-lick time

on average (range: 1-27%, *Figure 6B-D*). Alternate formulations of the decoding model produced

similar results (*Figure 6—figure supplement 1*).

Characterizing single-trial dopaminergic dynamics

Although the threshold-crossing analysis made no assumptions about the underlying dynamics of the GCaMP6f signals on single-trials, in principle, ramping dynamics in *averaged* neural signals could be produced from individual trials with a single, discrete "step" occurring at different times on different trials. Ramping has long been observed in averaged neural signals recorded during perceptual decision tasks in monkeys, and there has been considerable debate over whether single-trial responses in these experiments are better classified as "ramps" or a single "step" (*Latimer et al., 2015, 2016; Shadlen et al., 2016; Zoltowski et al., 2019; Zylberberg and Shadlen, 2016*). It has even been suggested that different sampling distributions can produce opposite model classifications in ground-truth synthetic signals (*Chandrasekaran et al., 2018*).

We attempted to classify single-trial dynamics as a discrete stepping or ramping process with hierarchical Bayesian models implemented in probabilistic programs (*Figure 6—figure supplement 2A-B*). However, like the perceptual decision-making studies, we also found ambiguous results, with about half of single-trials best classified by a linear ramp and half best classified by a discrete step dynamic (*Figure 6—figure supplement 2C*). Nonetheless, three separate lines of evidence suggest that single trials are better characterized by slowly-evolving ramps:

First, the relationship of threshold-crossing time to first lick time is different for the step *vs.* ramp models when different threshold levels are sampled (*Maimon and Assad, 2006*), as schematized in *Figure 6—figure supplement 3A*. Increasing slope of this relationship is consistent with ramps on single trials, but not with a discrete step, which would be expected to have the same threshold-

crossing time regardless of threshold level (*Figure 6—figure supplement 3B*). We found that the

slope of this relationship increased markedly as the threshold level was increased, consistent with

the ramp model (n=12 mice, mean slope low: 0.46, mid: 0.7, high: 0.82, *Figure 6—figure*

supplement 3C).

Second, if single trials involve a step change occurring at different times from trial-to-trial, then

aligning trials on that step should produce a clear step on average (rather than a ramp) (*Latimer et*

al., 2015). We thus aligned single-trial GCaMP6f signals according to that optimal step position

determined from a Bayesian step model fit for each trial and then averaged the step-aligned signals

across trials. The averaged signals did not resemble a step function, but rather yielded a sharp

transient superimposed on a "background" ramping signal (*Figure 6—figure supplement 4A*).

Step-aligned tdTomato and EMG averages showed a small inflection at the time of the step, but

neither signal showed background ramping. This suggests that the detected "steps" in the

GCaMP6f signals were likely transient movement artifacts superimposed on the slower ramping

dynamic rather than *bona fide* steps.

Third, the ideal step model holds that the step occurs at different times from trial-to-trial, producing

a ramping signal when trials are averaged together. In this view, the trial-by-trial variance of the

signal should be maximal at the time at which 50% of the steps have occurred among all trials, and

the signal should be minimal at the beginning and end of the interval (when no steps or all steps

have occurred, respectively). We thus derived the optimal step time for each trial using the

Bayesian step model, and then calculated variance as a function of time within pools of trials with

similar movement times. The signal variance showed a monotonic downward trend during the

timed interval, with a minimum variance at the time of movement rather than at the point at which

50% of steps had occurred among trials, inconsistent with the discrete step model (*Figure 6—*

figure supplement 4B).

Thus, altogether, we did not find evidence for a discrete step dynamic on single trials; on the

contrary, our observations concord with slow ramping dynamics on single trials. Regardless, our

GLM movement-time decoding approaches in *Figure 6* did not make any assumptions about

underlying single-trial dynamics.

Moment-to-moment DAN activity causally controls movement timing

Because dopaminergic ramping signals robustly predicted first-lick timing and were apparently

transmitted via dopamine release to downstream striatal neurons, ramping DAN activity may

causally determine movement timing. To test this, we optogenetically activated or inhibited DANs

(in separate experiments) on 30% of randomly-interleaved trials (*Figure 7A* and *Figure 7—figure*

supplement 1). For activation experiments, we chose light levels that elevated DAN activity within

the physiological range observed in our self-timed movement task, as assayed by simultaneous

photometry in the DLS with a fluorescent sensor of released dopamine (dLight1.1, *Figure 7—*

figure supplement 2). DAN activation significantly early-shifted the distribution of self-timed

movements on stimulated trials compared to unstimulated trials (12 mice, $p < 2.8*10^{-26}$, 2-sample

Kolmogorov-Smirnov (KS) Test), whereas inhibition produced significant late-shifting compared

to unstimulated trials (4 mice, $p < 0.0004$, 2-sided KS Test) (*Figure 7B and Figure 7—figure*

supplement 3A). Stimulation of mice expressing no opsin produced no consistent effect on timing

(5 mice, $p = 0.62$, 2-sided KS Test). The direction of these effects was consistent across all animals

tested in each category (*Figure 7B*). Complementary analysis methods revealed consistent effects

(bootstrapped difference in mean first-lick times between categories: p<0.05, *Figure 7C-D*;

bootstrapped comparison of difference in area under the cdf curves: p<0.05, *Figure 7—figure*

supplement 3B; bootstrapped difference in median first-lick times between categories: p<0.05,

Figure 7—figure supplement 3C). Similar effects were obtained with activation of SNc DAN

axon terminals in the DLS (2 mice, *Figure 7—figure supplement 3A-B*).

Recent studies have shown that physiological ranges of optogenetic DAN activation (as assayed

by simultaneous recordings from DANs) fail to elicit overt movements (*Coddington and Dudman,*

2018). We likewise found that optogenetic DAN activation did not elicit immediate licking outside

the context of the task (*Figure 7—figure supplement 4A*). Additionally, optogenetic DAN

inhibition did not reduce the rate of spontaneous licking outside the context of the task (*Figure 7—*

figure supplement 4B). In both cases, we used the same light levels that had elicited the robust

shifts in timing behavior during the self-timed movement task. In other control experiments, we

purposefully drove neurons into non-physiological activity regimes during the task by applying

higher activation light levels. Over-stimulation caused large, immediate, sustained increases in

DLS dopamine (*Figure 7—figure supplement 2*), comparable in amplitude to the typical reward-

related dopamine transients on interleaved, unstimulated trials. These non-physiological

manipulations resulted in rapid, nonpurposive body movements and disrupted performance of the

task. Together, these results suggest that the optogenetic effects on timing in *Figure 7* did not

result from direct, immediate triggering or suppression of movement, nor from non-physiological

dopamine release due to over-stimulation.

Linking endogenous DAN signals to the moment-to-moment probability of movement initiation

Optogenetic manipulations of DAN activity in the physiological range appeared to modulate the *probability* of initiating the pre-potent, self-timed movement. Given that endogenous DAN signals increased during the timing interval of the self-timed movement task, we reasoned that the probability of movement should likewise increase over the course of the timed interval. We thus derived a nested probabilistic movement-state decoding model to explore the link between DAN signals and movement propensity (*Figure 8A*). We applied a GLM based on logistic regression, in which we classified each moment of time as either a non-movement (0) or movement (1) state (*Figure 8A-B*), and we examined how well various parameters could predict the probability of transitioning from the non-movement state to the movement state. Unlike the decoding model in *Figure 6*, which considers a single threshold-crossing time, the probabilistic approach takes into account continuous DAN signals. Initial model selection included previous trial history (movement time and reward outcome history) in addition to the DAN GCaMP6f signal, but Bayesian Information Criterion (BIC) analysis indicated that the instantaneous GCaMP6f signal alone was a robustly significant predictor of movement state, whereas previous trial outcomes were insignificant contributors and did not further improve the model (*Figure 8—figure supplement 1*). We thus only considered the DAN GCaMP6f signal as a predictor in subsequent analyses.

The continuous DAN GCaMP6f signal was indeed predictive of current movement state at any time t, and it served as a significant predictor of movement state, up to at least 2 seconds in the past (*Figure 8C*). However, the signals became progressively more predictive of the current movement state as time approached t. That is, the dopaminergic signal levels closer to time t tended

to absorb the behavioral variance explained by more distant, previous signal levels (*Figure 8C*),

reminiscent of how threshold crossing time absorbed the variance explained by the baseline

dopaminergic signal in the movement-timing decoding model (*Figure 6B-C*). This observation is

consistent with a diffusion-like ramping process on single trials, in which the most recent

measurement gives the best estimate of whether there will be a transition to the movement state

(but is difficult to reconcile with a discrete step process on single trials, consistent with the results

in *Figure 6—figure supplements 3-4*).

We applied the fitted instantaneous probabilities of transitioning to the movement state to derive

a fitted hazard function for each behavioral session (*Figure 8D*). The DAN GCaMP6f signals were

remarkably predictive of the hazard function, both for individual sessions and on average,

explaining 65% of the variance on average (n=12 mice). Conversely, when the model was fit on

the same data in which the timepoint identifiers were shuffled, this predictive power was

essentially abolished, explaining only 5% of the variance on average (*Figure 8E*).

Together, these results demonstrate that slowly-evolving dopaminergic signals are predictive of

the moment-to-moment probability of movement initiation. When combined with the optogenetics

results, they argue that dopaminergic signals causally modulate the moment-to-moment

probability of the pre-potent movement. In this view, trial-by-trial variability in the DAN signal

gives rise to trial-by-trial differences in movement timing in the self-timed movement task.

DISCUSSION

A role for slowly ramping DAN activity in the timing of self-initiated movements

We found that both baseline and slowly ramping DAN signals were predictive of the timing of self-initiated movements. A number of studies have reported short-latency (<500 ms) increases in DAN activity following sensory cues and immediately preceding self-initiated movements (*Coddington and Dudman, 2018; da Silva et al., 2018; Dodson et al., 2016; Howe and Dombeck, 2016; Schultz et al., 1997*), similar to the sensory- and motor-related transients we observed within ~500 ms of the cue and first-lick. However, the ramping DAN signals we observed during self-timing were markedly different. The ramping signal unfolded over *seconds*, preceding the first-lick by as long as 10 s. Furthermore, variations in both ramping dynamics and baseline amplitude predicted the trial-by-trial timing of the first-lick, whether signals were recorded from SNc cell bodies, SNc axon terminals in the DLS, or VTA cell bodies. DAN signals were also reflected in the dynamics of dopamine release in the DLS, indicating availability of this information to downstream striatal effectors.

Optogenetic augmentation and suppression of DAN activity during the timing interval causally altered movement timing. Importantly, optogenetic DAN activation within the physiological range did not evoke immediate movements, consistent with prior work (*Coddington and Dudman, 2018; Lee et al., 2020*). This suggests that, rather than serving as direct drivers of movement, DANs influence *when* movements occur by modulating the moment-by-moment probability of movement. Applying this probabilistic view to the endogenous dopaminergic signals, we found that the ramping dynamics were highly predictive of the moment-by-moment probability of movement (as

captured by the hazard function), with DAN signals became progressively better predictors as the

time of movement onset approached. These findings suggest that variations in slow DAN

dynamics affect movement timing by influencing the moment-to-moment probability of

generating a movement.

This view of dopaminergic modulation could be related to classic findings from extrapyramidal

movement disorders, in which dysfunction of the nigrostriatal pathway produces aberrations in

movement initiation rather than paralysis or paresis (*Bloxham et al., 1984; Fahn, 2011; Hallett*

and Khoshbin, 1980). That is, movements do occur in extrapyramidal disorders, but at

inappropriate times, either too little/late (*e.g.*, Parkinson's), or too often (*e.g.*, dyskinesias).

Moreover, based on the deficits observed in Parkinsonian states (*e.g.*, perseveration), this role may

extend to behavioral transitions more generally, *e.g.*, starting new movements *or* stopping ongoing

movements.

Dopaminergic ramping in other contexts

Previous studies have reported slow ramping dopaminergic signals in certain behavioral contexts,

including goal-directed navigation (*Howe et al., 2013*); multi-step tasks culminating in reward

(*Hamid et al., 2016; Howard et al., 2017; Mohebi et al., 2019*); and passive observation of

dynamic visual cues indicating proximity to reward (*Kim et al., 2019*). It has been proposed that

slowly ramping mesolimbic DAN signals could encode increasing value or reward anticipation as

animals approach reward (*Hamid et al., 2016; Mohebi et al., 2019*), or alternatively could reflect

"ongoing" reward-prediction errors (RPE) (*Kim et al., 2019; Mikhael et al., 2019*). The origin of

the ramping signals we observed in the nigrostriatal system are consistent with either value or RPE

interpretations. (In a companion theoretical paper (*Hamilos and Assad, 2020*), we examine the

possible origins of these dopaminergic signals in value and reward-prediction error computational

frameworks (*Kim et al., 2019; Mikhael and Gershman, 2019; Mikhael et al., 2019*), which offers

a reconciliation of apparently contradictory DAN signals reported in the context of a perceptual

timing task *(Soares et al., 2016)*)).

Regardless of their origin, it has been unclear how the brain *employs* slowly ramping DAN signals

in behavior. Our study moves beyond previous work by finding that trial-by-trial variability in

ramping dynamics explains the precise timing of a behavioral output—the self-timed lick—and

that optogenetically manipulating SNc DAN activity causally alters the timing of that output. Thus,

SNc ramping may not merely encode progress toward a behavioral goal, but could play a causal

role in advancing that progress. In this view, ramping DAN signals could be related to, or even

drive, ramping signals that have been observed in the motor system in anticipation of self-initiated

movements, *e.g.*, readiness potentials in human EEG recordings (Deecke, 1996; Libet et al., 1983).

This view raises the question of whether the slow-timescale DAN signals we observed are unique

to the timing requirement of our task or are present before any self-initiated movement. When we

averaged DAN signals aligned to "spontaneous" licks during the ITI, we also observed noisy, slow

ramping signals building over seconds to the time of the next lick, with a time course related to

the duration of the inter-*lick* interval (*Figure 8—figure supplement 2*). This observation raises the

possibility that slowly evolving DAN signals may be integral to the generation of self-initiated

movements more generally—although our highly trained animals may have also been "rehearsing"

timed movements between trials. It would be interesting to see whether slow ramping dynamics

predictive of movement timing could be detected in previously published datasets if DAN signals

were similarly averaged according to the interval between self-initiated movement bouts.

Relationship to setpoint and stretching dynamics in other networks during self-timed

movement

We found that DAN signals predict movement timing via two low-dimensional signals: a baseline

offset and a ramping dynamic that "stretches" depending on trial-by-trial movement timing.

Intriguingly, similar stretching of neural responses has been observed before self-timed movement

in other brain areas in rats and primates, including the dorsal striatum (*Emmons et al., 2017; Mello*

et al., 2015; Wang et al., 2018), lateral interparietal cortex (*Maimon and Assad, 2006*),

presupplementary and supplementary motor areas (*Mita et al., 2009*), and dorsomedial frontal

cortex (DMFC) (*Remington et al., 2018; Sohn et al., 2019; Wang et al., 2018; Xu et al., 2014*).

In the case of DMFC, applying dimensionality reduction to the population responses revealed two

lower-dimensional characteristics that resembled our findings in DANs: 1) the speed at which the

population dynamics unfolded was scaled ("stretched") to the length of the produced timing

interval (*Wang et al., 2018*), and 2) the population state at the beginning of the self-timed

movement interval ("setpoint") was correlated with the timed interval (*Remington et al., 2018;*

Sohn et al., 2019). Recurrent neural network models suggest variation in stretching and setpoint

states could be controlled by (unknown) tonic or monotonically-ramping inputs to the cortico-

striatal system (*Remington et al., 2018; Sohn et al., 2019; Wang et al., 2018*). We found that

DANs exhibit both baseline (*e.g.*, "setpoint") signals related to timing, as well as monotonically-

ramping input during the timing interval. Thus, through their role as diffusely-projecting

modulators, DANs could potentially orchestrate variations in cortico-striatal dynamics observed

during timing behavior.

Possible relationship to motivational/movement vigor

In operant tasks in which difficulty is systematically varied over blocks of trials, increased inter-

trial dopamine in the nucleus accumbens has been associated with higher average reward rate and

with decreased latency to engage in a new trial, suggesting a link between dopamine and

"motivational vigor," the propensity to invest effort in work (*Hamid et al., 2016; Mohebi et al.,*

2019). Intriguingly, we observed the *opposite* relationship in the self-timed movement task:

periods with higher average reward rates had *lower* average baseline dopaminergic signals and

later first-lick times. Moreover, for a given first-lick time (*e.g.*, 3.5-3.75 s), we did not detect

differences in baseline (or ramping) signals during periods with different average reward rates,

such as near the beginning or end of a session. This difference between the two tasks may be due

to their opposing strategic constraints: in the aforementioned experiments, faster trial initiation

increased the number of opportunities to obtain reward, whereas earlier first-licks tended to

decrease reward acquisition in our self-timed movement task.

The basal ganglia have also been implicated in controlling "movement vigor," generally referring

to the speed, force or frequency of movements (*Bartholomew et al., 2016; Dudman and Krakauer,*

2016; Panigrahi et al., 2015; Turner and Desmurget, 2010; Yttri and Dudman, 2016). The

activity of nigrostriatal DANs has been shown to correlate with these parameters during movement

bouts and could promote more vigorous movement via push-pull interactions with the direct and

indirect pathways (*Barter et al., 2015; da Silva et al., 2018; Mazzoni et al., 2007; Panigrahi et*

al., 2015). Movement vigor might also entail earlier self-timed movements, mediated by moment-

to-moment increases in dopaminergic activity.

If moving earlier is a signature of greater movement vigor, then earlier self-timed movements

might also be executed with greater force/speed. We looked for movement-related vigor signals,

examining both the amplitude of lick-related EMG signals and the latency between lick initiation

and lick-tube contact. We detected no consistent differences in these force- or speed-related

parameters as a function of movement time; on the contrary, the EMG signals were highly

stereotyped irrespective of the first-lick time (data not shown). It is possible that vigor might affect

movement timing without affecting movement kinematics/dynamics—but, if so, the distinction

between "timing" and "vigor" would seem largely semantical.

Overall view

Lesion and pharmacological studies have long suggested roles for the SNc and dopamine in timing

(*Meck, 2006; Merchant et al., 2013*). Broadly speaking, conditions that increase dopamine

availability affect timing as if speeding an internal "pacemaker" (*Dews and Morse, 1958; Mikhael*

and Gershman, 2019; Schuster and Zimmerman, 1961), whereas conditions that decrease

dopamine availability/efficacy generally have the opposite effect (*Malapani et al., 1998; Meck,*

1986, 2006; Merchant et al., 2013). The dopaminergic ramping signals we observed bear some

resemblance to Pacemaker-Accumulator models of neural timing, a longstanding conceptual

framework for timing behavior in which a hypothetical accumulator signals that an interval has

elapsed when it reaches a threshold level (*Gallistel and Gibbon, 2000; Lustig and Meck, 2005;*

Meck, 2006).

However, we would suggest a more nuanced view of the role of DANs in self-timed movements.

Our optogenetic manipulations indicate that, rather than abruptly triggering movement when

threshold is attained, DAN activity modulates the probability of the pre-potent movement. In this

view, as DAN activity ramps up, the probability of movement likewise increases, and thus different

rates of increase in DAN activity equate to shorter or longer elapsed intervals before movement,

on average. This framework leaves open the question of what makes movements "probabilistic."

One possibility is that recurrent cortical-basal ganglia–thalamic circuits could act to generate

movements "on their own," without direct external triggers. By providing crucial modulation of

these circuits, DANs could tune the propensity to make self-initiated movements—and

pathological loss of DANs could reduce the production of such movements. Future experiments

should address how dynamic dopaminergic input influences downstream motor circuits involved

in self-initiated movements.

562
Figure 1. Self-timed movement task. (**A**) Task schematic (3.3 s version shown). (**B**) First-lick
timing distributions generated by the same mouse exhibit the scalar property of timing
(Weber's Law). Red: 3.3 s target time (4 sessions); Blue: 5 s target time (4 sessions). For all
mice, see *Figure 1—figure supplement 1B*. (**C-E**) Hazard-function analysis. Time=0 is the
start-timing cue; dashed vertical lines are target times. (**C**) Uniform instantaneous probability
of movement over time is equivalent to a flat hazard rate (bottom) and produces an exponential
first-lick timing distribution (top). (**D**) Before Training: First day of exposure to the self-timed
movement task. Top: average first-lick timing distribution across mice; bottom: corresponding
hazard functions. Gray traces: single session data. Red traces: average among all sessions, with
shading indicating 95% confidence interval produced by 10,000x bootstrap procedure. (**E**)
Trained Behavior: Hazard functions (bottom) computed from the first-lick timing distributions
for the 3.3 s- and 5 s tasks (top) reveal peaks at the target times. Right: average first-lick timing
distribution and hazard functions for all 12 GCaMP6f photometry animals. See also *Figure*
1—figure supplements 1-2. Source data: *Figure 1—source data*.

577
578

Figure 2. SNc DAN signals preceding self-timed movement. (**A**) Left: surgical strategy for GCaMP6f/tdTomato fiber photometry. Right: average SNc DAN GCaMP6f response for first-licks between 3-3.25 s (12 mice). Data aligned separately to both cue-onset (left) and first-lick (right), with the break in the time axis indicating the change in plot alignment. (**B**) Average SNc DAN GCaMP6f responses for different first-lick times (indicated by dashed vertical lines). (**C**) Comparison of average DAN GCaMP6f and tdTomato responses on expanded vertical scale. Traces plotted up to 150 ms before first-lick. See also *Figure 2—figure supplements 1-3*. Source data: *Figure 2—source data*.

Figure 3. Striatal dopamine release during the self-timed movement task. Photometry signals averaged together from DA$_{2m}$ signals (n=4 mice) and dLight1.1 signals (n=5 mice) recorded in DLS. Axis break and plot alignment as in *Figure 2*. Dashed lines: first-lick times. Inset, left: surgical strategy. Inset, right: Comparison of dLight1.1 and DA$_{2m}$ dynamics. Expanded vertical scale to show ramping in the average signals for DA$_{2m}$ (solid trace) and dLight1.1 (dashed trace) up until the time of the first-lick (first-lick occurred between 2-3 s after the cue for this subset of the data). See also: *Figure 3—figure supplement 1*. Source data: *Figure 3—source data*.

Figure 4. Movement controls reliably detected movements, but there were no systematic differences in movement during the timing interval. **(A)** Schematic of movement-control measurements**. (B)** First-lick-aligned average movement signals on rewarded (red) and unrewarded (blue) trials. Pre-lick traces begin at the nearest cue-time (dashed red, dashed blue). Left: one session; Right: all sessions. Dashed grey line: time of earliest-detected movement on most sessions (150 ms before first-lick). Average first-lick-aligned tdTomato optical artifacts showed inconsistent excursion directions (up/down) even within the same session; signals for each artifact direction shown in *Figure 4—figure supplement 1*. Source data: *Figure 4— source data*.

Figure 5. Contribution of optical artifacts, task variables and nuisance bodily movements to SNc GCaMP6f signals. (**A**) Nested encoding model comparing the contribution of timing-independent predictors (TI) to the contribution of timing-dependent predictors (TD). (**B**) Predicted dF/F signal for one session plotted up to time of first-lick. Model error simulated 300x (shading). (**C**) Nested encoding model for one session showing the actual recorded signal (1st panel), the timing-independent model (2nd panel), and the full, timing-dependent model with all predictors (3rd panel). Top: GCaMP6f; Bottom: tdTomato (tdt). Right: relative loss improvement by timing-dependent predictors (grey dots: single sessions, line: median, box: lower/upper quartiles, whiskers: 1.5x IQR). See also *Figure 5—figure supplement 1*. Source data: *Figure 5—source data*.

Figure 6. Single-trial DAN signals predict first-lick timing. **(A)** Schematic of nested decoding model. Categories for n-1th trial predictors: 2) reaction, 3) early, 4) reward, 5) ITI first-lick (see *Methods*). Bottom: single-trial cue-aligned SNc DAN GCaMP6f signals from one session (6 trials shown for clarity). Traces plotted up to first-lick. Right: threshold-crossing model. Low/Mid/High label indicates threshold amplitude. Dots: single trials. **(B)** Model weights. Error bars: 95% CI, *: $p<0.05$, 2-sided t-test. Numbers indicate nesting-order. **(C)** Variance explained by each model nest. Grey lines: single sessions; thick black line: average. For model selection, see *Figure 6—figure supplement 1C*. **(D)** Predicted *vs*. actual first-lick time, same session as *6A*. See also *Figure 6— figure supplements 1-4*. Source data: *Figure 6—source data*.

Figure 7. Optogenetic DAN manipulation systematically and bidirectionally shifts the timing of self-timed movements. (**A**) Strategy for optogenetic DAN activation or inhibition. Mice were stimulated from cue-onset until first-lick or 7 s. (**B**) Empirical continuous probability distribution functions (cdf) of first-lick times for stimulated (blue line) versus unstimulated (grey line) trials. Arrow and shading show direction of effect. P-values calculated by Kolmogorov-Smirnov test (for other metrics, see *Figure 7—figure supplements 1-2*). (**C**) Mean 1,000,000x bootstrapped difference in first-lick time, stimulated-minus-unstimulated trials. Box: upper/lower quartile; line: median; whiskers: 1.5x IQR; dots: single sessions. (**D**) Comparison of mean first-lick time difference across all sessions. Error bars: 95% confidence interval (*: p<0.05, 1,000,000 bootstrapped mean difference in first-lick time between sessions of different stimulation categories). See also *Figure 7—figure supplements 1-4*. Source data: *Figure 7—source data*.

Figure 8. Single-trial dynamic dopaminergic signals predict the moment-to-moment probability of movement initiation. (**A**) Probabilistic movement-state model schematic. (**B**) Single-trial DAN GCaMP6f signals at SNc from one session. First-lick time truncated 150 ms before movement detection to exclude peri-movement signals. Bottom: Movement states for the trials shown as a function of time. Diagram on the right schematizes the model predictors relative to an example time=t on a single trial. (**C**) Nested model fitted coefficients. (**D**) Decoded hazard functions from full model (with all 10 predictors). Thick line=mean. n=12 mice. (**E**) Hazard function fitting with shuffled datasets abolished the predictive power of the model (same 12 mice). See also *Figure 8— figure supplements 1-2*. Source data: *Figure 8—source data*.

MATERIALS AND METHODS

Animals

Adult male and female hemizygous DAT-cre mice (*Backman et al., 2006*) (B6.SJL-Slc6a3[tm1.1(cre)Bkmm]/J, RRID:IMSR_JAX:020080; The Jackson Laboratory, Bar Harbor, ME) or *wt* C57/b6 mice were used in all experiments (> 2 months old at the time of surgery; median body weight 23.8g, range 17.3-31.9 g). Mice were housed in standard cages in a temperature and humidity-controlled colony facility on a reversed night/day cycle (12 h dark/12 h light), and behavioral sessions occurred during the dark cycle. Animals were housed with enrichment objects provided by the Harvard Center for Comparative Medicine (IACUC-approved plastic toys/shelters, e.g., Bio-Huts, Mouse Tunnels, Nest Sheets, *etc*.) and were housed socially whenever possible (1-5 mice per cage). All experiments and protocols were approved by the Harvard Institutional Animal Care and Use Committee (IACUC protocol #05098, Animal Welfare Assurance Number #A3431-01) and were conducted in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals.

Surgery

Surgeries were conducted under aseptic conditions and every effort was taken to minimize suffering. Mice were anesthetized with isoflurane (0.5-2% at 0.8 L/min). Analgesia was provided by *s.c.* 5 mg/kg ketoprofen injection during surgery and once daily for 3 d postoperatively (Ketofen, Parsippany, NJ). Virus was injected (50 nL/min) and the pipet remained in place for 10 min before removal. 200 μm, 0.53 NA blunt fiber optic cannulae (Doric Lenses, Quebec, Canada) or tapered fiber optic cannulae (200 μm, 0.60 NA, 2 mm tapered shank, OptogeniX, Lecce, Italy) were positioned at SNc, VTA or DLS and secured to the skull with dental cement (C&B Metabond,

Parkell, Edgewood, NY). Neck EMG electrodes were constructed from two Teflon-insulated 32G

stainless steel pacemaker wires attached to a custom socket mounted in the dental cement. Sub-

occipital neck muscles were exposed by blunt dissection and electrode tips embedded bilaterally.

Stereotaxic coordinates (from bregma and brain surface)

Viral Injection:

SNc: 3.16 mm posterior, +/- 1.4 mm lateral, 4.2 mm ventral

VTA: 3.1 mm posterior, +/-0.6 mm lateral, 4.2 mm ventral

DLS: 0 mm anterior, +/- 2.6 mm lateral, 2.5 mm ventral.

Fiber Optic Tips:

SNc/VTA: 4.0 mm ventral (photometry) or 3.9 mm ventral (optogenetics).

DLS: 2.311 mm ventral (blunt fiber) or 4.0 mm ventral (tapered fiber)

Virus

Photometry:

tdTomato ("tdt"): AAV1-CAG-FLEX-tdT (UNC Vector Core, Chapel Hill, NC), 100 nL

694   used alone or in mixture with other fluorophores (below), working concentration

695               $5.3*10^{12}$ gc/mL

696   gCaMP6f (at SNc or VTA): 100 nL AAV1.Syn.Flex.GCaMP6f.WPRE.SV40

697   ($2.5*10^{13}$ gc/mL, Penn Vector Core, Philadelphia, PA). Virus was mixed in a 1:3 ratio

698   with tdt (200 nL total)

699   DA_{2m} (at DLS): 200-300 nL AAV9-hSyn-DA4.4(DA2m) (working concentration:

700   *ca*. $3*10^{12}$ gc/mL, Vigene, Rockville, MD) + 100 nL tdt

dLight1.1 (at DLS): 300 nL AAV9.hSyn.dLight1.1.wPRE bilaterally at DLS (*ca.*

9.6*10^{12} gc/mL, Children's Hospital Boston Viral Core, Boston, MA) + 100 nL

AAV1.CB7.CI.TurboRFP.WPRE.rBG (*ca*. 1.01*10^{12} gc/mL, Penn Vector Core)

Optogenetic stimulation/inhibition (all bilateral at SNc):

ChR2: 1000 nL AAV5-EF1a-DIO-hChR2(H134R)-EYFP-WPRE-pA (3.2*10^{13} gc/mL,

UNC Vector Core, Chapel Hill, NC)

ChrimsonR +/- dLight1.1: 700 nL AAV1-hSyn-FLEX-ChrimsonR-tdT (4.1*10^{12} gc/mL,

UNC Vector Core, Chapel Hill, NC) +/- 400-550 nL AAV9-hSyn-dLight1.1

bilaterally at DLS (*ca*. 10^{13} gc/mL, Lin Tian Lab, Los Angeles, CA)

stGtACR2: 300 nL 1:10 AAV2/8-hSyn1-SIO-stGtACR2-FusionRed (working

concentration 4.7*10^{11} gc/mL, Addgene/Janelia Viral Core, Ashburn, VA)

Water-deprivation and acclimation

Animals recovered for 1 week postoperatively before water deprivation. Mice received daily water

supplementation to maintain ≥80% initial body weight and fed *ad libitum*. Mice were habituated

to the experimenter and their health was monitored carefully following guidelines reported

previously (*Guo et al., 2014*). Training commenced when mice reached the target weight (~8-9 d

post-surgery).

Histology

Mice were anesthetized with >400 mg/kg pentobarbital (Somnasol, Henry Schein Inc, Melville,

NY) and perfused with 10 mL 0.9% sodium chloride followed by 50mL ice-cold 4%

paraformaldehyde in 0.1 M phosphate buffer. Brains were fixed in 4% paraformaldehyde at 4°C

for >24 hr before being transferred to 30% sucrose in 0.1 M phosphate buffer for >48 hr. Brains

were sliced in 50 μm coronal sections by freezing microtome, and fluorophore expression was

assessed by light microscopy. The sites of viral injections and fiber optic placement were mapped

with an Allen Mouse Brain Atlas.

Behavioral rig, data acquisition and analysis

A custom rig provided sensory cues, recorded events and delivered juice rewards under the control

of a Teensy 3.2 microprocessor running a custom Arduino state-system behavioral program with

MATLAB serial interface. Digital and analog signals were acquired with a CED Power 1400 data

acquisition system/Spike2 software (Cambridge Electronic Design Ltd, Cambridge, England).

Photometry and behavioral events were acquired at 1,000 Hz; movement channels were acquired

at 2,000 Hz. Video was acquired with FlyCap2 or Spinnaker at 30 fps (FLIR Systems, Wilsonville,

OR). Data were analyzed with custom MATLAB statistics packages.

Self-timed movement task

Mice were head-fixed with a juice tube positioned in front of the tongue. The spout was placed as

far away from the mouth as possible so that the tongue could still reach it to discourage compulsive

licking (*Guo et al., 2014*), ~1.5 mm ventral and ~1.5 mm anterior to the mouth. During periods

when rewards were not available, a houselamp was illuminated. At trial start, the houselamp turned

off, and a random delay ensued (0.4-1.5 s) before a cue (simultaneous LED flash and 3300 Hz

tone, 100 ms) indicated start of the timing interval. The timing interval was divided into two

windows, early (0-3.333 s in most experiments; 0-4.95 s in others) and reward (3.333-7 s; 4.95-10

s), followed by the intertrial interval (ITI, 7-17 s; 10-20 s). The window in which the mouse first

licked determined the trial outcome (early, reward, or no-lick). An early first-lick caused an error

tone (440 Hz, 200 ms) and houselamp illumination, and the mouse had to wait until the full timing

interval had elapsed before beginning the ITI. Thus there was no advantage to the mouse of licking

early. A first-lick during the reward window caused a reward tone (5050 Hz, 200 ms) and juice

delivery, and the houselamp remained off until the end of the trial interval. If the timing interval

elapsed with no lick, a time-out error tone played (131 Hz, 2 s), the houselamp turned on, and ITI

commenced. During the ITI and pre-cue delay ("lamp-off interval"), there was no penalty for

licking.

Mice learned the task in 3 stages (*Figure 1—figure supplement 1A*). On the first 1-4 days of

training, mice learned a beginner-level task, which was modified in two ways: 1) to encourage

participation, if mice did not lick before 5 s post-cue, they received a juice reward at 5 s; and 2)

mice were not penalized for licking in reaction to the cue (within 500 ms). When the mouse began

self-triggering ≥50% of rewards (day 2-6 of training), the mouse advanced to the intermediate-

level task, in which the training reward at 5 s was omitted, and the mouse had to self-trigger all

rewards. After completing >250 trials/day on the intermediate task (usually day 4-7 of training),

mice advanced to the mature task, with no reaction licks permitted. All animals learned the mature

task and worked for ~400-1,500 trials/session.

Hazard function correction of survival bias in the timing distribution

The raw frequency of a particular response time in the self-timed movement task is "distorted" by

how often the animal has the chance to respond at that time (*Anger, 1956*). This bias was corrected

by calculating the hazard function, which takes into account the number of response opportunities

the animal had at each timepoint. The hazard function is defined as the conditional probability of

moving at a time, t, given that the movement has not yet occurred (referred to as "IRT/Op" analysis

in the old Differential Reinforcement of Low Rates (DRL) literature). The hazard function was

computed by dividing the number of first-movements in each 250 ms bin of the first-lick timing

histogram by the total number of first-movements occurring at that bin-time or later—the total

remaining "opportunities."

Online movement monitoring

Movements were recorded simultaneously during behavior with four movement-control

measurements: neck EMG (band-pass filtered 50-2,000 Hz, 60 Hz notch, amplified 100-1,000x),

back-mounted accelerometer (SparkFun Electronics, Boulder, CO), high-speed camera

(30 Hz, FLIR Systems, Wilsonville, OR), and tdTomato photometry. All control signals contained

similar information, and thus only a subset of controls was used in some sessions.

Photometry

Fiber optics were illuminated with 475 nm blue LED light (Plexon, Dallas, TX) (SNc/VTA: 50 µW,

DLS: 35 µW) measured at patch cable tip with a light-power meter (Thorlabs, Newton, NJ). Green

fluorescence was collected via a custom dichroic mirror (Doric Lenses, Quebec, Canada) and

detected with a Newport 1401 Photodiode (Newport Corporation, Irvine, CA). Fluorescence was

allowed to recover ≥1 d between recording sessions. To avoid crosstalk in animals with red control

fluorophore expression, the red channel was recorded at one of the 3 sites (SNc, VTA, or DLS,

550 nm lime LED, Plexon, Dallas, TX) while GCaMP6f, dLight1.1 or DA_{2m} was recorded

simultaneously only at the other implanted sites.

dF/F

795 Raw fluorescence for each session was pre-processed by removing rare singularities (single

796 points >15 STD from the mean) by interpolation to obtain F(t). To correct photometry signals for

797 bleaching, dF/F was calculated as:

$$\frac{dF}{F}(t) = \frac{F(t) - F_0(t)}{F_0(t)}$$

where $F_0(t)$ is the 200 s moving average of F(t) (*Figure 2—figure supplement 1A*). We tested

several other complementary methods for calculating dF/F and all reported results were robust to

dF/F method (see *Methods: dF/F method characterization and validation*). To ensure dF/F signal

processing did not introduce artifactual scaling or baseline shifts, we also tested several

complementary techniques to isolate undistorted F(t) signals where possible and quantified the

amount of signal distortion when perfect isolation was not possible (see *Methods: dF/F method*

characterization and validation, below, and *Figure 2—figure supplement 1C*).

dF/F method characterization and validation

dF/F calculations are intended to reduce the contribution of slow fluorescence bleaching to fiber

photometry signals, and many such methods have been described (*Kim et al., 2019; Mohebi et al.,*

2019; Soares et al., 2016). However, dF/F methods have the potential to introduce artifactual

distortion when the wrong method is applied in the wrong setting. Thus, to derive an appropriate

dF/F method for use in the context of the self-timed movement task, we characterized and

quantified artifacts produced by 4 candidate dF/F techniques.

Detailed description of complementary dF/F methods.

1. Normalized baseline: a commonly used dF/F technique in which each trial's fluorescence is normalized to the mean fluorescence during the 5 s preceding the trial.

2. Low-pass digital filter: F_0 is the low-pass, digital infinite impulse response (IIR)-filtered raw fluorescence for the whole session (implemented in MATLAB with the built-in function *lowpass* with $f_c = 5 \cdot 10^{-5}$ Hz, steepness=0.95).

3. Multiple baseline: a variation of Method 1, in which each trial's fluorescence is normalized by the mean fluorescence during the 5 s preceding the current trial, as well as 5 trials before the current trial and 5 trials after the current trial.

4. Moving average: F_0 is the 200 s moving average of the raw fluorescence at each point (100 s on either side of the measured timepoint).

Although *normalized baseline* (Method 1) is commonly used to correct raw fluorescence signals (F) for bleaching, this technique assumes that baseline activity has no bearing on the trial outcome; however, because the mouse decides when to move in the self-timed movement task, it is possible that baseline activity may differ systematically with the mouse's choice on a given trial. Thus, normalizing F to the baseline period would obscure potentially physiologically-relevant signals. More insidiously, if baseline activity *does* vary systematically with the mouse's timing, normalization can also introduce substantial amplitude scaling and y-axis shifting artifacts when correcting F with this method (*Figure 2—figure supplement 1C*, middle panels). Thus, Methods 2-4 were designed and optimized to isolate photometry signals minimally distorted by bleaching signals and systematic baseline differences during the self-timed movement task. Methods 2-4

produced the same results in all statistical analyses, and the moving average method is shown in

all figures.

Isolating minimally-distorted photometry signals with paired trial analyses of raw fluorescence.

Although slow bleaching prevents comparison of raw photometry signals (F) at one time in a

behavioral session with those at another time, the time-course of appreciable bleaching was slow

enough in the reported behavioral sessions that minimal bleaching occurred over the course of

3 trials (~1 min, *Figure 2—figure supplement 1A*). Thus, to observe the most minimally-distorted

photometry signals possible, we compared F between *pairs of consecutive* trials (*Figure 2—figure*

supplement 1B-C). We compared F baseline signals between all paired trials in which an early

trial (unrewarded first-lick between 0.7-2.9 s; abbreviated as "E") was followed by a rewarded trial

(first-lick between 3.4-7 s; abbreviated as "R"); this two-trial sequence is thus referred to as an

"ER" comparison. To ensure systematic differences did not result from subtle bleaching in the

paired-trial interval, we reversed the ordering contingency and also compared all Rewarded trials

preceding Early trials ("RE" comparison). The same systematic relationship between baseline

signals and first-lick time was found for paired trials analyzed by raw F (*Figure 2—figure*

supplement 1C, left panels).

Quantification of artifactual amplitude scaling/baseline shifts introduced by dF/F processing.

Each Candidate dF/F Method was applied to the same Paired Trial datasets described above. The

resulting paired-fluorescence datasets were normalized after processing (minimum dF/F=0,

maximum=1). The amount of distortion introduced by dF/F was quantified with a Distortion Index

(DI), which was calculated as:

$$\text{Distortion Index, DI}(t) = \text{abs}(F(t) - dF/F(t))$$

where $F(t)$ and $dF/F(t)$ are the normalized, paired-trial raw fluorescence signal or dF/F signal at

time t, respectively. t spanned from the beginning of the n-1th trial (-20 s) to the end of the nth trial

(20 s), aligned to the cue of the nth trial (*Figure 2—figure supplement 1C, bottom panels*). The

DI shown in plots has been smoothed with a 200 ms moving average kernel for clarity.

As expected, normalizing fluorescence to the baseline period (*normalized baseline*) erased the

correlation of baseline dF/F signals with first-lick time (*Figure 2—figure supplement 1C*, middle

panels). More insidiously, this also resulted in distortion of GCaMP6f dynamics *during* the timing

interval, evident in the diminished difference between E-signals compared to R-signals relative to

the shapes observed in the raw fluorescence paired-trial comparison (*Figure 2—figure*

supplement 1C, middle-bottom panel). However, dF/F Methods 2-4 visually and quantitatively

recapitulated the dynamics observed in the raw fluorescence comparison (*Figure 2—figure*

supplement 1C, right panels).

These results were corroborated by time-in-session permutation tests in which datasets for single

sessions were divided into thirds (beginning of session, middle of session, and end of session). The

differences between baseline and ramping dynamics observed in whole-session averages were

present even within these shorter blocks of time within the session (*i.e.*, faster ramping and elevated

baseline signals on trials with earlier self-timed licks). Furthermore, permutation tests in which the

block identity (begin, middle, end) was shuffled showed that this pattern held when trials with

earlier first-licks from the end of the session were compared with trials with later first-licks from

the beginning of the session (and *vice versa*).

Normalized dF/F for comparing dopamine sensor signals

DA_{2m} was about twice as bright as dLight1.1, and thus generally yielded larger and less noisy dF/F

signals. To compare the two extracellular dopamine sensors in the same plot, dF/F was normalized

for each signal by the amplitude of its lick-related transient. dF/F was calculated as usual, and then

the mean baseline-to-transient peak amplitude was measured for trials with first-licks occurring

between 2-3 s. Percentage NdF/F is reported as the percentage of this amplitude.

Dopamine sensor kinetics

dLight1.1 is an extracellular dopamine sensor derived from the dopamine-1-receptor, and has fast

reported kinetics: rise $t_{1/2} = 9.5 \pm 1.1$ ms, decay $t_{1/2} = 90 \pm 11$ ms (*Patriarchi et al., 2018*). DA_{2m} is

a new extracellular dopamine indicator derived from the dopamine-2-receptor, which provides

brighter signals. DA_{2m} signals have been reported to decay slowly in slice preparations but are

much faster *in vivo*, presumably because endogenous dopamine-clearance mechanisms are

preserved: reported rise $t_{1/2}$ ~50 ms, decay $t_{1/2}$ ~360 ms in freely behaving mice; decay $t_{1/2}$ ~190 ms

in head-fixed *drosophila* (*Sun et al., 2020*). To estimate the dopamine-sensor kinetics in our head-

fixed mice, we examined the phasic fluorescence transient occurring on unrewarded first-licks

(0.5-3.3 s), which showed a stereotyped fast rise and decay with both sensors (*Figure 2—figure

supplement 2D-E*). While the transient was somewhat complex (reminiscent of phasic burst-pause

responses sometimes observed for movement-related DAN activity (*Coddington and Dudman,

2018, 2019*), we measured the time for average fluorescence to decay from the peak of the transient

to half the baseline-to-peak amplitude. We found decay $t_{1/2}$~75 ms for dLight1.1 and $t_{1/2}$~125 ms

for DA_{2m} (*Figure 3—figure supplement 1*). Given that the dopaminergic ramping signals in our

study evolved over several seconds, the kinetics of both dopamine sensors are thus fast enough

that they should not have caused appreciable distortion of the slow ramping dynamics.

Pearson's correlation of baseline signals to first-lick time.

The mean SNc GCaMP6f signal during the minimum lamp-off interval (-0.4 s to 0 s, the cue-time)

was compared to the first-lick time for pooled trials in *Figure 2C* by calculating the Pearson

correlation coefficient. There were at least 700 trials in each pooled set of trials (0.75-4 s included).

DAN signal encoding model

To test the independent contribution of each task-related input to the photometry signal and select

the best model, we employed a nested fitting approach, in which each dataset was fit multiple times

(in "nests"), with models becoming progressively more complex in subsequent nests. The nests fit

to the GCaMP6f photometry data employed the inputs $X^{(j)}$ at each j^{th} nest:

921   Null Model: $X^{(0)} = x_0$

922   Nest 1: $X^{(1)} = X^{(0)} + \text{tdTomato (tdt)}$

923   Nest 2: $X^{(2)} = X^{(1)} + \text{cue} + \text{first-lick}$

924   Nest 3: $X^{(3)} = X^{(2)} + \text{EMG/accelerometer}$

925   Nest 4: $X^{(4)} = X^{(3)} + \text{time-dependent baseline offset}$

926   Nest 5: $X^{(5)} = X^{(4)} + \text{stretch representing percentages of interval}$

Overfitting was penalized by ridge regression, and the optimal regularization parameter for each

nest was obtained by 5-fold cross-validation to derive the final model fit for each session. Model

improvement by each input was assessed by the percentage loss improvement at the nest where

the input first appeared compared to the prior nest. The loss improvement of Nest 1 was compared

to the Null Model (the average of the photometry timeseries). The nested model of tdt control

photometry signals was the same, except Nest 1 was omitted.

The GLM for each nest takes the form:

935                                    $$Y = \theta X^{(j)}$$

Where Y is the *1xn* vector of the photometry signal across an entire behavioral session (*n* is the

total number of sampled timepoints); $X^{(j)}$ is the *dxn* design matrix for nest *j*, where the rows

correspond to the d_j predictors for nest *j* and the columns correspond to each of the *n* sampled

timepoints of Y; and θ is the *dx1* vector of fit weights.

Y is the concatenated photometry timeseries taken from trial start (lights off) to the time of first

lick. Because of day-to-day/mouse-to-mouse variation (ascribable to many possible sources, *e.g.*,

different neural subpopulations, expression levels, behavioral states, *etc.*), each session was fit

separately.

The d_j design matrix predictors were each scaled (maximum amplitude 1) and grouped by input to

the model. The timing-independent inputs were: 1. Null offset (x_0, 1 predictor), 2. tdt (1 predictor),

3. cue (24 predictors), 4. first-lick (28 predictors), and 5. EMG/accelerometer (44 predictors). The

timing-dependent inputs were: 6. timing-dependent baseline offset (1 predictor), 7. stretch

(500 predictors).

To reduce the number of predictors, the cue, first-lick and EMG/accelerometer predictors

(*Figure 5—figure supplement 1C*) were composed from sets of basis kernels as described

previously (*Park et al., 2014; Runyan et al., 2017*). The cue basis kernels were spaced 0-500 ms

post-cue and first-lick basis kernels were spaced -500 ms-0 ms relative to first-lick, the typically-

observed windows of stereotypical sensory and motor-related neural responses. For nuisance

movements (EMG/accelerometer), events were first discretized by thresholding (*Figure 5—*

figure supplement 1B) and then convolved with basis kernels spanning -500 to 500 ms around the

event. This window was consistent with the mean movement-aligned optical artifact observed in

the tdt channel. The timing-dependent baseline offset was encoded as a constant offset spanning

from lamp-off until first-lick, with amplitude taken as linearly proportional to the timed interval

on the current trial. The timing-dependent stretch input was composed of 500 predictors, with each

predictor containing 1's tiling 0.05% of the cue-to-lick interval, and 0's otherwise (*Figure 5—*

figure supplement 1D). Importantly, the stretch was not constrained in any way to form ramps.

Basis sets were optimized to minimize Training Loss, as calculated by mean squared error of the

unregularized model:

$$\text{argmin}_{X^{(j)}}(\text{Training Loss}(\Theta) = 1/n * (Y - \Theta X^{(j)})^2)$$

Superfluous basis set elements that did not improve Training Loss compared to the Null Model

were not included in the final model. Goodness of the training fit was assessed by Akaike

Information Criterion (AIC), Bayesian Information Criterion (BIC), R^2, and Training Loss. The

optimal, regularized model for each nest/session was selected by 5-fold cross-validation in which

the regularization parameter, λ_j, was optimized for minimal average Test Loss:

$$\text{argmin}_{\lambda_j} (\text{Test Loss}(\Theta,\lambda_j) = 1/n * (Y - \Theta X^{(j)})^2 + \lambda_j|\Theta|^2)$$

Test Loss for each optimal model was compared across nests to select the best model for each

session. Models were refit with the optimal λ_j to obtain the final fit.

Model error was simulated 1,000 times by redrawing θ coefficients consistent with the data

following the method described by Gelman and Hill (*Gelman and Hill, 2006*), and standard errors

were propagated across sessions. The absolute value of each predictor was summed and divided

by the total number of predictors for that input to show the contribution of the input to the model

(*Figure 5—figure supplement 1G*). To simulate the modeled session's photometry signal for each

nest j, Yfit was calculated as $\theta X^{(j)}$ and binned by the time of first-lick relative to the cue. The error

in the simulation was shown by calculating $Yfit_{sim} = \theta_{sim} X^{(j)}$ for 300 simulated sets of θ_{sim}.

Principal component analysis (PCA)

Unsmoothed ramping intervals for photometry timeseries were fit with PCA and reconstructed

with the first three principal components (PCs). To derive a PCA fit matrix with ramping intervals

of the same number of samples, the length of each trial was scaled up by interpolation to the

maximum ramping interval duration:

$\qquad$ 7 s – 0.7 s cue buffer – 0.6 s first-lick buffer = 5.7 s: 5,700 sample ramping interval

Following PC-fitting, datasets were down-sampled to produce a fit of the correct time duration.

Trials where the ramping interval was <0.1 s were excluded to exclude noise from down-sampling.

First-lick time decoding model

A nested, generalized linear model was derived to predict the first-lick time on each trial in a

session and quantify the contribution of previous reward history and photometry signals to the

prediction. The model was of the form:

$$\log(y) = bx$$

where y is the first-lick time, b is a vector of fit coefficients and x is a vector of predictors. The

nested model was constructed such that predictors occurring further back in time (such as reward

history) and confounding variables (such as tdt photometry signals) were added first to determine

the additional variance explained by predictors occurring closer to the time of first-lick, which

might otherwise obscure the impact of these other variables. The predictors, in order of nesting,

were:

Nest 0:   b0 (Null model, average log-first-lick time)

Nest 1:   b1 = b0 + first-lick time on previous trial (trial "n-1")

Nest 2-5: b2 = b1 + previous trial outcome (1,0)*

Nest 6:   b3 = b2 + median photometry signal in 10s window before lamp-off ("ITI")

Nest 7:   b4 = b3 + median photometry signal from lamp-off to cue ("lamp-off interval")

Nest 8:   b5 = b4 + tdt threshold crossing time**

Nest 9:   b6 = b5 + GCaMP6f threshold crossing time**

1015

1016 where all predictors were normalized to be in the interval (0,1).

1017

1018 * Outcomes included (in order of nest): Reaction (first-lick before 0.5 s), Early (0.5-3.333 s),

1019 Reward (3.333-7 s), ITI (7-17 s). No-lick was implied by all four outcomes encoded as zeros.

1020 ** Details on threshold-crossing time and alternative models included in *Methods: Derivation of*

1021 *threshold and alternative decoding models*.

1022

1023 To exclude the sensory- and motor-related transients locked to the cue and the first-lick events in

1024 the threshold-crossing nests, the ramping interval was conservatively defined as 0.7 s post-cue up

1025 until 0.6 s before first-lick, and the minimum ramping interval for fitting was 0.1 s. Thus, for a

1026 trial to be included in the model, the first lick occurred between 1.4 s to 17 s (end of trial).

1027

1028 Initial model goodness of fit was assessed by R^2, mean-squared loss and BIC. Models were 5-fold

1029 cross-validated with ridge regression at each nest to derive the final models, as described above.

1030 95% confidence intervals on model coefficients were calculated by 2-sided t-test with standard

1031 errors propagated across sessions.

1032

Derivation of threshold and alternative decoding models

Derivation of threshold models

As a metric of the predictive power of ramping DAN signals on first-lick time, we derived a

threshold-crossing model. A threshold-crossing event was defined as the first time after the cue

when the photometry signal exceeded and remained above a threshold level up until the time of

first-lick on each trial. Importantly, while the analysis approach is reminiscent of pacemaker-

accumulator models for timing, we make no claims that the analysis is evidence for pacemaker-

accumulator models. Rather threshold-crossing times provided a convenient metric to compare the

rate of increase in signals between trials.

1042

Photometry timeseries for GCaMP6f and tdt were de-noised by smoothing with a 100 ms Gaussian

kernel (kernel was optimized by grid screen of kernels ranging between 0-200 ms to minimize

noise without signal distortion). To completely exclude the sensory- and motor-related transients

locked to the cue and the first-lick events, the ramping interval was conservatively defined as 0.7

s post-cue up until 0.6 s before the first-lick. To eliminate chance crossings due to noise, we

imposed a stiff, debounced threshold condition: to be considered a threshold crossing event, the

photometry signal had to cross the threshold from low-to-high and remain above this level until

the end of the ramping interval.

1051

To derive an unbiased threshold for each session, we tested 100 evenly-spaced candidate threshold

levels spanning the minimum-to-maximum photometry signal during the ramping interval for each

session. Depending on threshold level, some trials never crossed, *i.e.*, signal always remained

below threshold or started and ended above threshold. Thus, the lowest candidate threshold for

which there was a maximum number of trials crossing during the timing interval was selected as

the "mid-level" threshold-crossing point. This threshold was specific to each photometry signal

tested on each session. Threshold-crossing time was included in the decoding model as the

normalized time on the ramping interval $(0,1)$. If a trial never crossed threshold, it was encoded as

a zero. If no trials ever crossed threshold, the threshold predictor was encoded as a vector of ones,

thus penalizing the model for an additional predictor but providing no new information.

1062

Multi-threshold Model

An alternative model employed 3 unbiased thresholds: 1) the lowest threshold with ≥ 50 trials

crossing ("min"); 2) the lowest threshold with the most crossings ("mid," described above); and 3)

the highest threshold with ≥50 trials crossing ("max"). For tdt datasets, trials rarely met the

monotonic threshold constraint (usually the signals oscillated above and below the threshold

throughout the ramping interval, failing to meet the debouncing constraint). Thus, to include tdt

signals as conservatively as possible, we relaxed the 50-trial minimum constraint, taking the

threshold with the most trials crossing, which was usually around 10 or fewer. The addition of

more thresholds did not substantially improve the cross-validated model compared to the single-

threshold model (*Figure 6—figure supplement 1*).

1073

Principal component analysis (PCA) threshold-crossing models

In another version of the decoding model, the threshold-crossing procedures were applied to

ramping intervals fit with the first three PCs (as described in *Methods: Principal Component*

Analysis (PCA)) to derive a PCA version of the single-threshold and multi-threshold models. PCA

analysis on tdt datasets showed no consistent PCs, and thus these PCs were not included in the

decoding model. Instead, the actual tdt data was employed in the threshold model as in the other

models described.

1081

Hierarchical Bayesian Modeling of Single-trial Dynamics.

The probability of each single-trial SNc GCaMP6f signal belonging to a ramp vs. step Model Class

was determined via Hierarchical Bayesian Model fitting with probabilistic programs written in the

novel probabilistic programming language, Gen.jl, which is embedded in the Julia Programming

Language (*Cusumano-Towner et al., 2019*). The top of the model hierarchy was the model class

(linear ramp *vs*. step function) and the lower level was the respective parameterization of the two

model classes (described below).

1089

The probability of the step *vs*. ramp model class was inferred with data-driven inference. The best

fit (step or ramp and parameterization) for each trial was calculated across 20 iterations (Gen

Traces) of hierarchical modeling with 50 rounds of probabilistic refinement (computation *via* Gen

Importance Resampling) per iteration (in model testing, models typically converged to their

steady-state probability of model class within only 30 rounds of refinement, but 50 rounds were

used conservatively to reduce the likelihood of suboptimal classifications).

1096

<u>*Data-driven inference procedure*</u>: Each iteration of model fitting began at the top level of the

hierarchy with a coin toss: with 50% probability, the probabilistic program would initialize with a

model of either the Ramp or Step class. For data-driven inference, a Gen *Proposal* for the

parameterization for this model class was then probabilistically generated. Data-driven proposals

were designed to improve fitting efficiency and reduce computation time, and this allowed for

faster convergence and better model fits as determined by the fit log-likelihood. The proposal

heuristics were as follows:

<u>Ramp model</u>: A data-driven proposal was generated by dynamic noise random sample

consensus (RANSAC (***Cusumano-Towner and Mansinghka, 2018***)) with additional data-driven

constraints (see function `ransac_assisted_model_selection_proposal` in the Gen Github

files):

1. SLOPE, *a*. The maximum data-supported slope was used to set the variance of slope

sampling:

1110 $a \sim Gaussian$(RANSAC-sampled slope, *maxslope*/2).

where *maxslope* was defined as the difference of the maximum and minimum signal within the

trial dataset divided by the total duration of the trial (by definition, the largest slope supported

by the data).

2. INTERCEPT, *b*. The initial search for the intercept ("*b-max*") was calculated as the

intercept for the calculated *maxslope* parameter), and this was used to set the noise level

on sampling of the intercept parameter:

1117 $b \sim Gaussian$(RANSAC-sampled intercept, *b-max* /2)

1118   3. NOISE, σ. Parametrized noise level was sampled as:

1119 $\sigma \sim Beta(\alpha,\beta)$

1120   where α,β are the parameters of the beta distribution with mode=std(signal).

1121

1122   <u>Step model</u>**:** The data-driven proposal included two constraints/heuristics:

1123   1. STEPTIME. *Derivative constraint*: To avoid sampling all unlikely step-times, *steptimes*

1124   were sampled uniformly from the timepoints where the derivative of the signal was in the

1125   highest 5% of the signal's derivative across the trial dataset:

1126   *steptime ~ uniform*(indices of 95th percentile of derivative of the signal)

1127   2. LEFT and RIGHT SEGMENTS. Once a *steptime* was sampled, likely *left* and *right*

1128   segment amplitudes were sampled near the mean of the signal on either side of the step,

1129   *e.g.*:

1130   *left ~ Gaussian*(mean(signal left of *steptime*), std(signal left of *steptime*))

1131   *right ~ Gaussian*(mean(signal right of *steptime*), std(signal right of *steptime*))

1132   3. NOISE, σ. The noise level was sampled as in the ramp model,

1133 $\sigma \sim Beta(\alpha,\beta)$

1134   except α,β were the parameterization of a Beta distribution with mode equal to the

1135   standard deviation of the signal left of *steptime*.

1136

1137   After model initialization for each Trace, 50 rounds of Importance Resampling of the hierarchical

1138   model were then conducted, each time randomly generating ramp or step hypotheses from the

1139   proposal heuristics. On each round, the best fitting hypothesis was retained, such that each of the

1140   20 Trace iterations of model classification returned one optimized model from the 50 rounds of

1141   Importance Resampling.

1142

1143   The probability of the model class for each single-trial was then defined as the proportion of the

1144   20 Trace iterations that found the optimal model to be derived from that model class (*e.g.*, if the

1145   model returned 15 step-fits and 5 ramp-fits, the p(ramp) was 0.25). Examples of the 20 Trace

1146   iterations for two sample trials are shown in *Figure 6—figure supplement 2B*.

1147

1148   To determine whether the step model detected step-functions in the GCaMP6f dataset, the step

1149   model was inferred alone to find step-fits for every trial, and single-trial signals were realigned to

1150   the optimal *steptime* (GCaMP6f, tdTomato, EMG, *Figure 6—figure supplement 4A-B*).

1151

1152   **Single-trial dynamics analysis with geometric modeling ("Multiple threshold modeling").**

1153   The multi-threshold procedure described above was also employed to determine whether single-

1154   trial ramping dynamics were more consistent with a continuous ramp *vs*. discrete step dynamic on

1155   single-trials. The threshold-crossing time for each trial was regressed against its first-lick time, and

1156   the slope of this relationship was reported, as well as the variance explained.

1157

Single-trial variance analysis for discrete step dynamics.

1158

For discrete step single trial dynamics to produce ramping on average, the time of the step across

trials must be distributed throughout the trial interval (importantly, a peri-motor spike occurring

consistently just before first-lick *cannot* give rise to continuous ramping dynamics on average).

As such, the variance in the GCaMP6f signals across trials for similar first-lick times should be

minimal near the time of the cue (when few trials have stepped) *and* near the time of the first-lick

(when all of the trials have stepped). This predicts an inverted-U shaped relationship of signal

variance across trials *vs*. position in the timing interval.

1166

To compare variance across trials equitably, trials were first aligned to the cue and pooled by first-

lick time in pools of 1s each (1-2 s, 2-3 s, *etc*.) truncating at the earliest first-lick time within the

pool. The variance in GCaMP6f signals across trials within a pool was quantified in 10% percent

increments of time from the cue up to the earliest first-lick time in the pool (*i.e*, 1-2 s pool truncated

at 1 s, divided into 100 ms increments). Measuring variance by percent of elapsed time within pool

allowed pooling of trials across the entire session. The shape of the variance *vs*. percent of timed

interval elapsed was compared to the inverted-U shape prediction to assess for discrete step

dynamics.

1175

1176 **Optogenetics—determining the physiological range for activation experiments**

1177 To test whether optogenetic manipulations during the self-timing task were in the physiological

1178 range, we assessed the magnitude of the effect of activation on dopamine release in the DLS by

1179 simultaneous photometry recordings with optical activation (*Figure 7—figure supplement 2*). In

two DAT-cre mice, we expressed ChrimsonR bilaterally in SNc DANs and the fluorescent

dopamine indicator dLight1.1 bilaterally in DLS neurons. SNc cell bodies were illuminated

bilaterally (ChrimsonR 550 nm lime or 660 nm crimson, 0.5-5 mW) on 30% of trials (10 Hz, 10

or 20 ms up-time starting at cue onset and terminating at first-lick). dLight1.1 was recorded with

35 μW 475 nm blue LED light at DLS. To avoid cross-talk between the stimulation LED and the

photometry recording site, the brief stimulation up-times were omitted from the photometry signal

and the missing points filled by interpolation between the adjacent timepoints.

In a few preliminary sessions, we also explored whether we could evoke short-latency licking (*i.e.*,

within a few hundred ms of the stimulation) if light levels were increased above the physiological

range for DAN signals. Rather than eliciting immediate licking, higher light levels produced bouts

of rapid, nonpurposive limb and trunk movements throughout stimulation, and task execution was

disrupted. The animals appeared to have difficulty coordinating the extension of the tongue to

touch the lick spout. Simultaneous DLS dopamine detection showed large, sustained surges in

dopamine release throughout the period of stimulation, with an average amplitude comparable to

that of the reward transient (*Figure 7—figure supplement 2, right*). This extent of dopamine

release was never observed during unstimulated trials. Consequently, to avoid overstimulation in

activation experiments, we kept light levels well below those that generated limb and trunk

movements.

Optogenetics—naïve/expert control sessions.

To determine whether optogenetic stimulation directly elicited or prevented licking, licking

behavior was first tested outside the context of the self-timed movement task on separate sessions

in the same head-fixed arena but with no cues or behavioral task. Opsin-expressing mice were

tested before any exposure to the self-timed movement task ("Naïve") as well as after the last day

of behavioral recording ("Expert"). In ChR2 control sessions, stimulation (5 mW 425 nm light, 3 s

duration, 10 Hz, 20% duty cycle) was applied randomly at the same pace as in the self-timed

movement task. stGtACR2 control sessions were conducted similarly (12 mW 425 mW light, 3 s

duration, constant illumination); but to examine if inhibition could block ongoing licking, we

increased the baseline lick-rate by delivering juice rewards randomly (5% probability checked

once every 5 s).

1211

1212 **Optogenetics—self-timed movement task.**

1213 SNc DANs were optogenetically manipulated in the context of the 3.3 s self-timed movement task.

1214 To avoid overstimulation, light levels were adjusted to be subthreshold for eliciting overt

1215 movements as described above, and mice were not stimulated on consecutive days.

1216

Activation: SNc cell bodies were illuminated bilaterally (ChR2: 0.5-5 mW 425 nm blue LED light;

ChrimsonR 550 nm lime or 660 nm crimson) on 30% of trials (10 Hz, 10 or 20% duty cycle

starting at cue onset and terminating at first-lick). DAN terminals in DLS were stimulated

bilaterally via tapered fiber optics on separate sessions.

Inactivation: SNc cell bodies were illuminated bilaterally (stGtACR2: 12 mW 425 nm blue light)

on 30% of trials (constant illumination starting at cue onset and terminating at first lick).

1223

Quantification of optogenetic effects.

The difference in the distribution of trial outcomes between stimulated and unstimulated trials on

each session was quantified in four ways.

1. 2-Sample Unsigned Kolmogorov-Smirnov Test.

2. Difference in empirical continuous probability distribution function (cdf). The difference

in the integral of the stimulated and unstimulated cdf (dAUC) was calculated for each

session from 0.7-7 s. Effect size was quantified by permutation test, wherein the identity

of each trial (stimulated or unstimulated) was shuffled, and the distribution of dAUCs for

the permuted cdfs was calculated 10,000x. Results were reported for all sessions.

3. Difference in mean movement time. Movement times on stimulated and unstimulated trials

were pooled and the distribution of movement time differences was determined by non-

parametric bootstrap, in which a random stimulated and unstimulated trial were drawn from

their respective pools 1,000,000x and the difference taken. The mean of each session's

bootstrapped distribution was compared across sessions by the 1,000,000x bootstrapped

difference of the mean between sessions of different categories.

4. Difference in median movement time. Same as above but with median.

1240

Single-trial probabilistic movement state decoding model.

The probability of transitioning to a movement state, $s_t=1$, at time=t was decoded with a logistic

generalized linear model of the form:

$$p(s_t=1) = \text{logit}(bX_t)$$

where X_t is a vector of predictors for the timepoint, t, and b is the vector of fit coefficients. The

vector of predictors was comprised of the GCaMP6f signal at every timepoint (the current time, t)

as well as the signal history, represented as 200 ms-wide signal averages moving back in time from

t. Previous trial history (n-1[th] and n-2[th] first-lick times and reward/no-reward outcomes) did not

contribute significantly to the model during model selection and were thus omitted (see Model

Selection, below).

1251

Movement state, s_t, was defined as a binary variable, where state=0 represented all timepoints

between the cue up until 160 ms before the first-lick detection (to exclude any potential peri-

movement responses), and state=1 represented the timepoint 150 ms before the first-lick. Because

there were many more state=0 than state=1 samples in a session, state=0 points were randomly

down-sampled such that states were represented equally in the fit. To avoid randomly sampling a

particular model fit by chance, each dataset was fit on 100 randomly down-sampled (bootstrapped)

sets, and the average fit across these 100 sets was taken as the model fit for the session.

1259

1260   GCaMP6f signals were smoothed with a 100 ms gaussian kernel and down-sampled to 100 Hz.

1261   The GCaMP6f predictors were then nested into the model starting with those furthest in time from

1262   the current timepoint, t:

1263

1264   Nest 0:   b0 (Null model)

1265   Nest 1:   b1 = b0 + mean GCaMP6f 1.8:2.0 s before current time=t

1266   Nest 2:   b2 = b1 + mean GCaMP6f 1.6:1.79 s before current time=t

1267   Nest 3:   b3 = b2 + mean GCaMP6f 1.4:1.59 s before current time=t

1268   Nest 4:   b4 = b3 + mean GCaMP6f 1.2:1.39 s before current time=t

1269   Nest 5:   b5 = b4 + mean GCaMP6f 1.0:1.19 s before current time=t

1270   Nest 6:   b6 = b5 + mean GCaMP6f 0.8:0.99 s before current time=t

1271   Nest 7:   b7 = b6 + mean GCaMP6f 0.6:0.79 s before current time=t

1272   Nest 8:   b8 = b7 + mean GCaMP6f 0.4:0.59 s before current time=t

1273   Nest 9:   b9 = b8 + mean GCaMP6f 0.2:0.39 s before current time=t

1274   Nest 10:   b10 = b9 + GCaMP6f signal at current time=t

1275

1276   Nesting the predictors from most distant in time to most recent permitted observation of the ability

1277   of more proximal signal levels to absorb the variance contributed by more distant signal history.

1278

The fitted hazard function was then found as the average probability of being in the movement

state across all trials in the session as calculated from the average model fit. Because s_t=0 states

were significantly down-sampled during fitting, this rescaled the fit hazard. Thus, to return the fit

hazard to the scale of the hazard function calculated from the behavioral distribution, both the fit

hazard and true hazard function were normalized on the interval (0,1), and the goodness of fit was

assessed by R^2 comparison of the fit and true hazard functions. This metric was similar between

individual session fits as well as the grand-average fit across all animals and sessions.

1286

1287   To guard against overfitting, this procedure was repeated on the same datasets, except the datasets

1288   were shuffled before fitting to erase any non-chance correlations between the predictors and the

1289   predicted probability of being in the movement state.

1290

Model selection

To evaluate the contribution of task performance history to the probability of being in the

movement state at time=t, we could not observe every timepoint in the GCaMP6f trial period

timeseries as we did in the final model because the trial history for a given timepoint was the same

for all other points in the trial; hence this created bias because the movement state=1 was

represented for all trials, but the likelihood of the a trial's 0 state being represented after down-

sampling was dependent on the duration of the trial (*i.e.*, first-lick time). Consequently, model

selection was executed on a modified version of the model that ensured that each trial would only

be represented *one time at most* in the fit. Because this greatly reduced the power of the model,

model selection was conducted on sessions from the two animals with the highest S:N ratio and

most trials to ensure the best chance of detecting effects of each predictor (*Figure 8—figure*

supplement 1).

1303

The set of permutations of GCaMP6f signal and task history were fit separately, and the best model

selected by BIC (though notably AIC and AICc were in agreement with the BIC selection). Each

model was fit in "time-slices"—windows of 500 ms from the time of the cue up until the first-lick.

Only one point for each trial was fit within this window to ensure the movement state within the

window was uniquely represented. For each time-slice model, the GCaMP6f signal for each trial

was thus averaged within the time-slice window, and the movement state was 1 only if the

movement state occurred sometime within the window. The model fit for a session was taken as

the average model fit across each of the time-slices. Notably, a time-slice required a sufficient

number of trials to be present (either in the $s_t=0$ or terminating in the movement state $s_t=1$) for the

fit to converge; once the first-lick occurred for a trial, it did not contribute data to later time-slices.

The source data files for *Figure 8—figure supplement 1* contain plots of all time-slice coefficient

fits, including for models with insufficient numbers of trials to converge.

1316

1317   **Code Availability.** All custom behavioral software and analysis tools are available with sample

1318   datasets at https://github.com/harvardschoolofmouse.

1319

1320   **Data Availability.** All datasets supporting the findings of this study are publicly available

1321   https://www.dropbox.com/sh/g671nzziba405e3/AAByufom8BYs7PctECCbdiPua?dl=0

1322   (Repository to be moved to Zenodo for publication). Source data files have been provided for all

1323   figures.

1324

ACKNOWLEDGEMENTS

We thank D. Chicharro, J.G. Mikhael, S.J. Gershman, K. Reinhold, S. Panzeri, R. Bliss and

E.N. Brown for discussions on analytical methods; J. Tenenbaum, C. Wong, and A. Lew for their

instruction and advice pertaining to probabilistic programming in Gen; V. Berezovskii, J. LeBlanc,

T. LaFratta, O. Mazor, P. Gorelik, J. Markowitz, A. Lutas, K.W. Huang, L. Hou, S.J. Lee and

N.E. Ochandarena for technical assistance; and B. Sabatini, C. Harvey, S.R. Datta, M. Andermann,

L. Tian and W. Regehr for reagents. The work was supported by NIH grants UF-NS108177 and

U19-NS113201, and NIH core grant EY-12196. A.E.H. was supported by a Harvard Lefler

Predoctoral Fellowship and a Stuart H.Q. & Victoria Quan Predoctoral Fellowship at Harvard

Medical School. The funders had no role in study design, data collection and interpretation, or the

decision to submit the work for publication.

1336

AUTHOR CONTRIBUTIONS

1337

1338 A.E.H. and J.A.A conceived the project. A.E.H. performed all experiments. G.S. assisted with

1339 experiments using tapered fiber optics; Y.H. assisted with optogenetic no-opsin control

1340 experiments. F.S. and Y.L. developed the dopamine sensor, DA_{2m}. A.E.H. and J.A.A. analyzed the

1341 data and wrote the paper.

1342

COMPETING INTERESTS

1343

J.A.A. is a co-founder of OptogeniX, which produces the tapered optical fibers used in some

experiments.

Figure 1—figure supplement 1. Self-timed movement task learning and variations. (**A**) Task learning.
Histogram of first-lick times from single sessions at different stages of training (red: reaction, grey: early,
blue: operant-rewarded, yellow: Pavlovian-rewarded). Bars >50 first-licks truncated for clarity. (**B**) Mice
adjust behavior to the timing-contingencies of the task. First-lick time distributions from tasks with different
target timing intervals. Red: 3.3 s reward-boundary. Blue: 5 s reward-boundary (all sessions, all mice). (**C**)
Mice time their first-licks relative to the start cue, not the houselamp. First-lick time distributions during
behavior with (red) and without (black) houselamp events (4 mice, 4-5 sessions/mouse on each version of
the task). Source data: *Figure 1—source data*.

1355
1356 **Figure 1—figure supplement 2**. Fiber optic placement and histology**. (A)** Approximate fiber positions for
1357 all mice. **(B)** Brightfield microscopy with polarized filter on a freshly-cut brain slice showing bilateral fiber
1358 placement at SNc (from stGtACR2 experiment). **(C)** Example of co-expression of green (DA_{2m}) and red
1359 (tdTomato) fluorophores relative to fiber optic tip.

1360
Figure 2—figure supplement 1. dF/F method validation. **(A)** Left: slow, raw fluorescence bleaching
across one session. Left inset: Minimal bleaching occurs across the first 3 trials (~1 min). Right: dF/F
removes slow bleaching dynamics. Right inset: The same 3-trial window shown for dF/F signal. **(B)**
Average raw fluorescence on paired, consecutive trials from one session aligned to cue on the n^{th} trial.
Left: n-1^{th} trial was early, n^{th} trial was rewarded ("ER" condition). Right: "RE" condition (See *Methods:
dF/F method characterization and validation*). **(C)** Comparison of baseline GCaMP6f signals on paired,
consecutive trials aligned to cue. Columns: three different versions of the signal (Raw fluorescence,
Normalized baseline dF/F method, Moving average dF/F method). Top row: ER condition; middle row: RE
condition; bottom row: distortion index. Red distortion index plot shows only Normalized baseline method.
Green distortion index plot shows overlay of Moving Average, Low-Pass Filter, and Multiple Baseline dF/F
Methods because the difference in signal distortion between these methods was indistinguishable (See
Methods: dF/F method characterization and validation). Source data: *Figure 2—source data*.

Figure 2—figure supplement 2. Average photometry signals, pooled every 250 ms by first-lick time, spanning 0.5 s (purple) to 7 s (red). Signals in main panels aligned only to cue, not first-lick. **(A)** Average DAN GCaMP6f signals at SNc cell bodies (12 mice). **(B)** DAN GCaMP6f signals at axon terminals in DLS (10 mice). **(C)** Striatal dopamine detection with dLight1.1 at DLS (5 mice). **(D)** Striatal DA_{2m} signals at DLS (4 mice). **(E)** DAN GCaMP6f signals at VTA cell bodies (4 mice). **(F)** tdTomato signals. **Insets (left)**: Cue *and* lick-aligned average signals for a single time bin before first-lick to show pre-lick ramping present in all dopaminergic signals. Left of axis break: aligned to cue. Right of axis break: aligned to first-lick. Traces plotted up until 150 ms before first-lick. Source data: *Figure 2—source data*.

1373
1374
1375
1376 **Figure 3—figure supplement 1.** Comparison of dLight1.1 (dashed) and DA_{2m} (solid) kinetics surrounding
1377 peak of unrewarded transient (first-lick: 0.5-3.3 s). Red line: ½ baseline-to-peak amplitude for measuring
1378 decay $t_{1/2}$ (see *Methods*). Source data: *Figure 3—source data*.

Figure 4—figure supplement 1. Average tdTomato optical artifacts (aligned to first-lick time) showed inconsistent directions even within the same session. Averages for all three types of artifact (consistently up, "Up"; consistently down, "Down"; and not consistent "NC") shown for all sessions. Pie plots: Breakdown of average tdt artifact direction by session at each recording site. Source data: *Figure 4—source data*.

Figure 5—figure supplement 1. DAN signal encoding model parameterization and model selection.
(**A**) Schematic of photometry timeseries fit by encoding model. The lamp-off to first-lick interval was
excised from each trial in a session (top) and concatenated to produce the timeseries fit by the model
(bottom). (**B**) EMG spikes derivation: thresholding rectified EMG at 3 standard deviations (example trial).
(**C**) Optimized basis kernels to produce timing-independent features. (**D**) Schematic of Design Matrix for
timing-dependent features. (**E**) GCaMP6f model fits by nest iteration for example session. Shading: model
error simulated 300x. (**F**) Model loss by nest iteration. Green: mean loss for SNc GCaMP6f; red: mean loss
for tdTomato (tdt); grey lines: individual sessions; grey shading: timing-dependent nests. Left: full-scale
view of all datasets. Right: mean GCaMP6f and tdt loss compared on same scale. (**G**) Summary of feature
weights across SNc GCaMP6f (left) and tdt (right) models. Coefficient weights were rectified, summed,
and divided by the number of predictors per feature. 2x standard error bars (too small to see). All features
were significant in both GCaMP6f and tdt models. (**H**) Top: examples of the full timing-dependent model
(nest 5) from additional mice for all recorded dopaminergic signals. Bottom: tdt control channel fit. Model
errors simulated 300x. Some mice show downward-going movement-related spikes at SNc cell bodies
(second panel). All mice showed downward-going movement-related spikes from SNc terminals in DLS
(middle panel). Source data: *Figure 5—source data*.

1410
1411
1412

Figure 5—figure supplement 2. Principal component analysis (PCA) of the ramping interval (0.7 s up to
first-lick relative to cue). **(A)** Left: Variance explained by first 10 principal components (PC). Right: first
three principal components. Green line: mean PC, GCaMP6f recorded at SNc; Red line: mean PC,
tdTomato (tdt) recorded at SNc and VTA; Grey lines: single-session data. X-axis shown for longest-
possible interpolated trial duration; trials of shorter duration were interpolated to have the same number of
samples for PCA. **(B)** Example session data simulated with first 3 PCs. Noisy traces: actual averaged
GCaMP6f signals truncated at first-lick onset; Smooth traces: PC fits of the same trials. Source data: *Figure*
5—source data.

Figure 6—figure supplement 1. Variations of the first-lick time decoding model. *: p<0.05, error bars:
95% confidence intervals. GCaMP6f threshold crossing time dominated every version of the model; n-1[th]
trial first-lick time was consistently the second-best predictor. Source data: *Figure 6—source data*.

1432
1433
Figure 6—figure supplement 2. Analysis of single-trial dynamics: Hierarchical Bayesian Ramp *vs*. Step
Modeling. **(A)** Schematic (see *Methods: Hierarchical Bayesian Modeling of Single-trial Dynamics*). **(B)**
Example fits from hierarchical model on 2 example single trials from the same epoch in a single session.
Green: SNc GCaMP6f single-trial signal, light grey shading: noise band, dark grey lines: model fits. Note
that the top trial is more frequently classified as a ramp, and the lower trial is more frequently classified as
a step. However, both the ramp and step models return intuitive and reasonable fits to both single-trial
signals. **(C)**. Probability of model class across all trials. X axis: 0 indicates all probabilistic fits for a given
trial returned step-class models; 1 indicates all ramp-class models. Single sessions across mice showed
considerable uncertainty in model classification. Source data: *Figure 6—source data*.
1443
1444
1445
1446

1447
1448

Figure 6—figure supplement 3. Geometric analysis of single-trial dynamics with Multiple Threshold
Modeling. **(A)** Left: linear ramp model, Right: discrete step model. Step positions drawn from uniform
distribution over the cue-to-first-lick interval. Low-, Mid- and High- level thresholds shown. **(B)** Threshold-
crossing time *vs.* first-lick time ("X-ing time *vs.* first-lick time") for (from top to bottom) High-, Mid- and
Low-level thresholds. Left: simulation predictions for ramp and step models. Right: X-ing time *vs.* first-
lick time regression fit on single trials from 1 session (data from *Figure 6A*). The step model predicts X-
ing time *vs.* first-lick time does not change across threshold levels, whereas ramp model predicts the slope
of this relationship increases as threshold is raised. Single-trial GCaMP6f data exhibits increasing X-ing
time *vs.* first-lick time slope with increasing threshold level, consistent with the ramp model but inconsistent
with the step model. **(C)**. X-ing time *vs.* first-lick time across all mice. Left column: frequency of slope
relationship across sessions, right column: variance explained. Source data: *Figure 6—source data*.

1460
1461
Figure 6—figure supplement 4. Assessing single-trial dynamics. **(A)** Single-trial signals aligned to discrete step
position as found by Bayesian step model do not exhibit discrete step dynamics. To best estimate step times, the two
animals with the highest GCaMP6f S:N were examined (Mouse B5 and B6). Left: 1 session, Right: average of signals
from both mice. **(B)** Variance of GCaMP6f signals across trials. Step times were computed by Bayesian step model.
An ideal step model predicts maximal variance at the 50th percentile step, but variance declined monotonically on
average. Grey lines: single sessions; black line: average. For detailed explanation, see *Methods: Single-trial variance*
analysis for discrete step dynamics. Source data: *Figure 6—source data*.

Figure 7—figure supplement 1. Variations on measurements of optogenetic effects. (A) Strategy for
optogenetic targeting of DANs. (B) Comparison of four complementary metrics for addressing optogenetic
effects. Left: unsigned Kolmogorov-Smirnov Distance (KS-D) analysis of differences in first-lick time
distribution. Center: signed, bootstrapped comparison of difference in area under the cdf curves (dAUC).
Right: mean and median bootstrapped difference in first-lick time. Source data: *Figure 7—source data*.

1474
1475
Figure 7—figure supplement 2. Light-power calibration for optogenetic activation of DANs. In preliminary
experiments, DLS dopamine levels were monitored during the self-timed movement task, in which SNc DANs
were activated randomly on 30% of interleaved trials. Dashed vertical lines: first-lick time. **Left**: interleaved,
unstimulated trials (2 mice, 8 sessions). **Middle**: stimulated trials at the range of light levels used in the activation
experiments show slightly elevated DLS dopamine signals compared to interleaved, unstimulated trials. First-
lick timing was generally early-shifted in these sessions. **Right**: in a subset of preliminary calibration sessions,
stimulation light levels were increased to the point where rapid, nonpurposive limb/trunk movements were
observed throughout stimulation (1 mouse, 3 sessions). DLS dopamine signals show much higher, sustained
increases throughout stimulation. Ongoing body movements disrupted task participation. Source data: *Figure*
7—source data.

Figure 7—figure supplement 3. Quantification of optogenetic effects with additional metrics. **(A)** KS-D
analysis: all sessions. "A": activation sessions; "NO": no opsin sessions; "I": inhibition sessions. Filled
circles indicate significant difference between stimulated/unstimulated trials on single session (p<0.025, 2-
sided, 2-sample KS test). Standard box plot, line: median, box: upper/lower quartiles; whiskers: 1.5x IQR.
(B) Left: bootstrapped dAUC Assay: all sessions, standard box plot as in *(A)*. Filled circles: significant
difference on single session (p<0.025, 2-sided bootstrapped dAUC test, see *Methods*). Right: comparison
of dAUC in first-lick distributions across all sessions between groups. Error bars denote bootstrapped 95%
confidence interval (*: p<0.05). **(C)** Left: median bootstrapped difference in first-lick time, stimulated-
minus-unstimulated trials, standard box plot as in *(A)*. Dots: single sessions. Right: Comparison of median
difference in first-lick time across all sessions. Error bars denote bootstrapped 95% confidence interval
(*: p < 0.05). Source data: *Figure 7—source data*.

1499
Figure 7—figure supplement 4. Optogenetic DAN stimulation does not cause or prevent licking. **(A,B)**
Stimulation-aligned lick-rate during control sessions. Animals were tested in 1-3 control sessions both before
exposure to the self-timed movement task (red) and in 1-2 control sessions after the end of behavioral training
(navy). Blue bar indicates stimulation period (3 s). Left: one session, Right: all sessions. **(A)** Activation control
sessions (no cues or rewards). Animals were head-fixed on the behavioral platform and stimulated randomly at
the same pace as the standard 3.3 s self-timed movement task. Activation did not elicit immediate licking in any
session. **(B)** Inhibition-control sessions (no cues, + random rewards). Animals were head-fixed on the behavioral
platform while receiving juice rewards at random times. Inhibition did not prevent licking in any session. Source
data: *Figure 7—source data*.

1509
1510
Figure 8—figure supplement 1. Probabilistic movement time decoding model: model selection. **(A)**
Model schematic. To assess previous trial history on the same footing as dopaminergic signals, time t during
model selection was limited to a 500 ms "time-slice," with each time-slice fit separately by the model.
Dopaminergic signals were averaged within each time-slice, such that each trial provided one and only one
dopaminergic measurement, one set of trial history terms, and one movement state per time slice (see
Methods: Single-trial probabilistic movement state decoding model, model selection). **(B)** Model fit
weights. Model ID: corresponds to the predictors included from the schematic. x-axis labels: the predictor
ID from the schematic. Predictor weights averaged across time-slices. **(C)** Model selection criteria. The
model omitting the previous trial history predictors (predictors #1-4) was consistently the best model as
selected by BIC, AIC and AICc (results similar across metrics, BIC shown alone for clarity). Source data:
Figure 8—source data.

Figure 8—figure supplement 2. Average Intertrial Interval (ITI) GCaMP6f signals aligned to most recent previous lick-time. Signals plotted up to onset of next spontaneous, self-initiated lick during the ITI. (1 mouse, 5 sessions, truncated 150 ms before lick). Source data: *Figure 8—source data.*

REFERENCES

Albin, R.L., Young, A.B., and Penney, J.B. (1989). The functional anatomy of basal ganglia disorders.
Trends Neurosci *12*, 366-375.

Anger, D. (1956). The dependence of interresponse times upon the relative reinforcement of different
interresponse times. J Exp Psychol *52*, 145-161.

Backman, C.M., Malik, N., Zhang, Y., Shan, L., Grinberg, A., Hoffer, B.J., Westphal, H., and Tomac, A.C.
(2006). Characterization of a mouse strain expressing Cre recombinase from the 3' untranslated region of
the dopamine transporter locus. Genesis *44*, 383-390.

Barter, J.W., Li, S., Lu, D., Bartholomew, R.A., Rossi, M.A., Shoemaker, C.T., Salas-Meza, D., Gaidis, E.,
and Yin, H.H. (2015). Beyond reward prediction errors: the role of dopamine in movement kinematics. Front
Integr Neurosci *9*, 39.

Barthel, C., Nonnekes, J., van Helvert, M., Haan, R., Janssen, A., Delval, A., Weerdesteyn, V., Debu, B.,
van Wezel, R., Bloem, B.R.*, et al.* (2018). The laser shoes: A new ambulatory device to alleviate freezing
of gait in Parkinson disease. Neurology *90*, e164-e171.

Bartholomew, R.A., Li, H., Gaidis, E.J., Stackmann, M., Shoemaker, C.T., Rossi, M.A., and Yin, H.H. (2016).
Striatonigral control of movement velocity in mice. Eur J Neurosci *43*, 1097-1110.

Bloxham, C.A., Mindel, T.A., and Frith, C.D. (1984). Initiation and execution of predictable and
unpredictable movements in Parkinson's disease. Brain *107 (Pt 2)*, 371-384.

Chandrasekaran, C., Soldado-Magraner, J., Peixoto, D., Newsome, W.T., Shenoy, K.V., and Sahani, M.
(2018). Brittleness in model selection analysis of single neuron firing rates. bioRxiv, 430710.

Coddington, L.T., and Dudman, J.T. (2018). The timing of action determines reward prediction signals in
identified midbrain dopamine neurons. Nat Neurosci *21*, 1563-1573.

Coddington, L.T., and Dudman, J.T. (2019). Learning from Action: Reconsidering Movement Signaling in
Midbrain Dopamine Neuron Activity. Neuron *104*, 63-77.

Cusumano-Towner, M.F., and Mansinghka, V.K. (2018). Using probabilistic programs as proposals. ArXiv
abs/1801.03612.

Cusumano-Towner, M.F., Saad, F.A., Lew, A.K., and Mansinghka, V.K. (2019). Gen: A General-Purpose
Probabilistic Programming System with Programmable Inference. Proceedings of the 40th Acm Sigplan
Conference on Programming Language Design and Implementation (Pldi '19), 221-236.

da Silva, J.A., Tecuapetla, F., Paixão, V., and Costa, R.M. (2018). Dopamine neuron activity before action
initiation gates and invigorates future movements. Nature *554*, 244-248.

Deecke, L. (1996). Planning, preparation, execution, and imagery of volitional action. Brain Res Cogn Brain
Res *3*, 59-64.

DeLong, M.R. (1990). Primate models of movement disorders of basal ganglia origin. Trends Neurosci *13*,
281-285.

Dews, P.B., and Morse, W.H. (1958). Some observations on an operant in human subjects and its
modification by dextro amphetamine. J Exp Anal Behav *1*, 359-364.

Dodson, P.D., Dreyer, J.K., Jennings, K.A., Syed, E.C., Wade-Martins, R., Cragg, S.J., Bolam, J.P., and
Magill, P.J. (2016). Representation of spontaneous movement by dopaminergic neurons is cell-type
selective and disrupted in parkinsonism. Proc Natl Acad Sci U S A *113*, E2180-2188.

Dudman, J.T., and Krakauer, J.W. (2016). The basal ganglia: from motor commands to the control of vigor.
Curr Opin Neurobiol *37*, 158-166.

Eckard, M.L., and Kyonka, E.G.E. (2018). Differential reinforcement of low rates differentially decreased
timing precision. Behav Processes *151*, 111-118.

Emmons, E.B., De Corte, B.J., Kim, Y., Parker, K.L., Matell, M.S., and Narayanan, N.S. (2017). Rodent
Medial Frontal Control of Temporal Processing in the Dorsomedial Striatum. J Neurosci *37*, 8718-8733.

Engelhard, B., Finkelstein, J., Cox, J., Fleming, W., Jang, H.J., Ornelas, S., Koay, S.A., Thiberge, S.Y.,
Daw, N.D., Tank, D.W.*, et al.* (2019). Specialized coding of sensory, motor and cognitive variables in VTA
dopamine neurons. Nature *570*, 509-513.

Fahn, S. (2011). Classification of movement disorders. Mov Disord *26*, 947-957.

Freeze, B.S., Kravitz, A.V., Hammack, N., Berke, J.D., and Kreitzer, A.C. (2013). Control of basal ganglia
output by direct and indirect pathway projection neurons. J Neurosci *33*, 18531-18539.

Gallistel, C.R., and Gibbon, J. (2000). Time, rate, and conditioning. Psychol Rev *107*, 289-344.

Gelman, A., and Hill, J. (2006). Data analysis using regression and multilevel/hierarchical models
(Cambridge ; New York: Cambridge University Press).

Grillner, S., and Robertson, B. (2016). The Basal Ganglia Over 500 Million Years. Curr Biol *26*, R1088-
R1100.

Guo, Z.V., Hires, S.A., Li, N., O'Connor, D.H., Komiyama, T., Ophir, E., Huber, D., Bonardi, C., Morandell,
K., Gutnisky, D.*, et al.* (2014). Procedures for behavioral experiments in head-fixed mice. PLoS One *9*,
e88678.

Hallett, M. (2007). Volitional control of movement: the physiology of free will. Clin Neurophysiol *118*, 1179-
1192.

Hallett, M., and Khoshbin, S. (1980). A physiological mechanism of bradykinesia. Brain *103*, 301-314.

Hamid, A.A., Pettibone, J.R., Mabrouk, O.S., Hetrick, V.L., Schmidt, R., Vander Weele, C.M., Kennedy,
R.T., Aragona, B.J., and Berke, J.D. (2016). Mesolimbic dopamine signals the value of work. Nat Neurosci
19, 117-126.

Hamilos, A.E., and Assad, J.A. (2020). Application of a unifying reward-prediction error (RPE)-based
framework to explain underlying dynamic dopaminergic activity in timing tasks. bioRxiv,
2020.2006.2003.128272.

Helassa, N., Podor, B., Fine, A., and Torok, K. (2016). Design and mechanistic insight into ultrafast calcium
indicators for monitoring intracellular calcium dynamics. Sci Rep *6*, 38276.

Howard, C.D., Li, H., Geddes, C.E., and Jin, X. (2017). Dynamic Nigrostriatal Dopamine Biases Action
Selection. Neuron *93*, 1436-1450 e1438.

Howe, M.W., and Dombeck, D.A. (2016). Rapid signalling in distinct dopaminergic axons during locomotion
and reward. Nature *535*, 505-510.

Howe, M.W., Tierney, P.L., Sandberg, S.G., Phillips, P.E., and Graybiel, A.M. (2013). Prolonged dopamine
signalling in striatum signals proximity and value of distant rewards. Nature *500*, 575-579.

Hughes, L.E., Altena, E., Barker, R.A., and Rowe, J.B. (2013). Perseveration and choice in Parkinson's
disease: the impact of progressive frontostriatal dysfunction on action decisions. Cereb Cortex *23*, 1572-
1581.

Jaldow, E.J., Oakley, D.A., and Davey, G.C. (1990). Performance on two fixed-interval schedules in the
absence of neocortex in rats. Behav Neurosci *104*, 763-777.

Kim, H.R., Malik, A.N., Mikhael, J.G., Bech, P., Tsutsui-Kimura, I., Sun, F., Zhang, Y., Li, Y., Watabe-Uchida,
M., Gershman, S.J.*, et al.* (2019). A unified framework for dopamine signals across timescales. bioRxiv,
803437.

Kirshenbaum, A.P., Brown, S.J., Hughes, D.M., and Doughty, A.H. (2008). Differential-reinforcement-of-
low-rate-schedule performance and nicotine administration: a systematic investigation of dose, dose-
regimen, and schedule requirement. Behav Pharmacol *19*, 683-697.

Latimer, K.W., Yates, J.L., Meister, M.L., Huk, A.C., and Pillow, J.W. (2015). NEURONAL MODELING.
Single-trial spike trains in parietal cortex reveal discrete steps during decision-making. Science *349*, 184-
187.

Latimer, K.W., Yates, J.L., Meister, M.L., Huk, A.C., and Pillow, J.W. (2016). Response to Comment on
"Single-trial spike trains in parietal cortex reveal discrete steps during decision-making". Science *351*, 1406.

Lee, I.H., and Assad, J.A. (2003). Putaminal activity for simple reactions or self-timed movements. J
Neurophysiol *89*, 2528-2537.

Lee, K., Claar, L.D., Hachisuka, A., Bakhurin, K.I., Nguyen, J., Trott, J.M., Gill, J.L., and Masmanidis, S.C.
(2020). Temporally restricted dopaminergic control of reward-conditioned movements. Nat Neurosci *23*,
209-216.

Libet, B., Gleason, C.A., Wright, E.W., and Pearl, D.K. (1983). Time of conscious intention to act in relation
to onset of cerebral activity (readiness-potential). The unconscious initiation of a freely voluntary act. Brain
106 (Pt 3), 623-642.

Lustig, C., and Meck, W.H. (2005). Chronic treatment with haloperidol induces deficits in working memory
and feedback effects of interval timing. Brain Cogn *58*, 9-16.

Maimon, G., and Assad, J.A. (2006). A cognitive signal for the proactive timing of action in macaque LIP.
Nat Neurosci *9*, 948-955.

Malapani, C., Rakitin, B., Levy, R., Meck, W.H., Deweer, B., Dubois, B., and Gibbon, J. (1998). Coupled
temporal memories in Parkinson's disease: a dopamine-related dysfunction. J Cogn Neurosci *10*, 316-331.

Mazzoni, P., Hristova, A., and Krakauer, J.W. (2007). Why don't we move faster? Parkinson's disease,
movement vigor, and implicit motivation. J Neurosci *27*, 7105-7116.

Meck, W.H. (1986). Affinity for the dopamine D2 receptor predicts neuroleptic potency in decreasing the
speed of an internal clock. Pharmacol Biochem Behav *25*, 1185-1189.

Meck, W.H. (2006). Neuroanatomical localization of an internal clock: a functional link between mesolimbic,
nigrostriatal, and mesocortical dopaminergic systems. Brain Res *1109*, 93-107.

Mello, G.B., Soares, S., and Paton, J.J. (2015). A scalable population code for time in the striatum. Curr
Biol *25*, 1113-1122.

Merchant, H., Harrington, D.L., and Meck, W.H. (2013). Neural basis of the perception and estimation of
time. Annu Rev Neurosci *36*, 313-336.

Mikhael, J.G., and Gershman, S.J. (2019). Adapting the flow of time with dopamine. J Neurophysiol *121*,
1748-1760.

Mikhael, J.G., Kim, H.R., Uchida, N., and Gershman, S.J. (2019). Ramping and State Uncertainty in the
Dopamine Signal. bioRxiv, 805366.

Mita, A., Mushiake, H., Shima, K., Matsuzaka, Y., and Tanji, J. (2009). Interval time coding by neurons in
the presupplementary and supplementary motor areas. Nat Neurosci *12*, 502-507.

Mohebi, A., Pettibone, J.R., Hamid, A.A., Wong, J.T., Vinson, L.T., Patriarchi, T., Tian, L., Kennedy, R.T.,
and Berke, J.D. (2019). Dissociable dopamine dynamics for learning and motivation. Nature *570*, 65-70.

Panigrahi, B., Martin, K.A., Li, Y., Graves, A.R., Vollmer, A., Olson, L., Mensh, B.D., Karpova, A.Y., and
Dudman, J.T. (2015). Dopamine Is Required for the Neural Representation and Control of Movement Vigor.
Cell *162*, 1418-1430.

Park, I.M., Meister, M.L., Huk, A.C., and Pillow, J.W. (2014). Encoding and decoding in parietal cortex
during sensorimotor decision-making. Nat Neurosci *17*, 1395-1403.

Parker, N.F., Cameron, C.M., Taliaferro, J.P., Lee, J., Choi, J.Y., Davidson, T.J., Daw, N.D., and Witten,
I.B. (2016). Reward and choice encoding in terminals of midbrain dopamine neurons depends on striatal
target. Nat Neurosci *19*, 845-854.

Patriarchi, T., Cho, J.R., Merten, K., Howe, M.W., Marley, A., Xiong, W.H., Folk, R.W., Broussard, G.J.,
Liang, R., Jang, M.J.*, et al.* (2018). Ultrafast neuronal imaging of dopamine dynamics with designed
genetically encoded sensors. Science *360*.

Rakitin, B.C., Gibbon, J., Penney, T.B., Malapani, C., Hinton, S.C., and Meck, W.H. (1998). Scalar
expectancy theory and peak-interval timing in humans. J Exp Psychol Anim Behav Process *24*, 15-33.

Remington, E.D., Narain, D., Hosseini, E.A., and Jazayeri, M. (2018). Flexible Sensorimotor Computations
through Rapid Reconfiguration of Cortical Dynamics. Neuron *98*, 1005-1019 e1005.

Romo, R., Scarnati, E., and Schultz, W. (1992). Role of primate basal ganglia and frontal cortex in the
internal generation of movements. II. Movement-related activity in the anterior striatum. Exp Brain Res *91*,
385-395.

Runyan, C.A., Piasini, E., Panzeri, S., and Harvey, C.D. (2017). Distinct timescales of population coding
across cortex. Nature *548*, 92-96.

Schultz, W., Dayan, P., and Montague, P.R. (1997). A neural substrate of prediction and reward. Science
275, 1593-1599.

Schuster, C.R., and Zimmerman, J. (1961). Timing behavior during prolonged treatment with dl-
amphetamine. J Exp Anal Behav *4*, 327-330.

Shadlen, M.N., Kiani, R., Newsome, W.T., Gold, J.I., Wolpert, D.M., Zylberberg, A., Ditterich, J., de Lafuente,
V., Yang, T., and Roitman, J. (2016). Comment on "Single-trial spike trains in parietal cortex reveal discrete
steps during decision-making". Science *351*, 1406.

Shenoy, K.V., Sahani, M., and Churchland, M.M. (2013). Cortical control of arm movements: a dynamical
systems perspective. Annu Rev Neurosci *36*, 337-359.

Sippy, T., Lapray, D., Crochet, S., and Petersen, C.C. (2015). Cell-Type-Specific Sensorimotor Processing
in Striatal Projection Neurons during Goal-Directed Behavior. Neuron *88*, 298-305.

Soares, S., Atallah, B.V., and Paton, J.J. (2016). Midbrain dopamine neurons control judgment of time.
Science *354*, 1273-1277.

Sohn, H., Narain, D., Meirhaeghe, N., and Jazayeri, M. (2019). Bayesian Computation through Cortical
Latent Dynamics. Neuron *103*, 934-947 e935.

Sun, F., Zhou, J., Dai, B., Qian, T., Zeng, J., Li, X., Zhuo, Y., Zhang, Y., Tan, K., Feng, J.*, et al.* (2020).
New and improved GRAB fluorescent sensors for monitoring dopaminergic activity in vivo.
bioRxiv, 2020.2003.2028.013722.

Turner, R.S., and Desmurget, M. (2010). Basal ganglia contributions to motor control: a vigorous tutor. Curr
Opin Neurobiol *20*, 704-716.

Wang, D.V., and Tsien, J.Z. (2011). Conjunctive processing of locomotor signals by the ventral tegmental
area neuronal population. PLoS One *6*, e16528.

Wang, J., Narain, D., Hosseini, E.A., and Jazayeri, M. (2018). Flexible timing by temporal scaling of cortical
responses. Nat Neurosci *21*, 102-110.

Xu, M., Zhang, S.Y., Dan, Y., and Poo, M.M. (2014). Representation of interval timing by temporally
scalable firing patterns in rat prefrontal cortex. Proc Natl Acad Sci U S A *111*, 480-485.

Yttri, E.A., and Dudman, J.T. (2016). Opponent and bidirectional control of movement velocity in the basal
ganglia. Nature *533*, 402-406.

Zoltowski, D.M., Latimer, K.W., Yates, J.L., Huk, A.C., and Pillow, J.W. (2019). Discrete Stepping and
Nonlinear Ramping Dynamics Underlie Spiking Responses of LIP Neurons during Decision-Making.
Neuron *102*, 1249-1258 e1210.

Zylberberg, A., and Shadlen, M.N. (2016). Cause for pause before leaping to conclusions about stepping.
bioRxiv, 085886.
1706