Updated phylogeny and sequence features of MreB from the wall-

less bacteria *Spiroplasma*

Daichi Takahashi[a], Ikuko Fujiwara[a,b], Makoto Miyata[a,b,#]

[a]Graduate School of Science, Osaka City University, Osaka, Japan

[b]The OCU Advanced Research Institute for Natural Science and

Technology (OCARINA), Osaka City University, Osaka, Japan

Running Title: Sequence analyses for *Spiroplasma* MreBs

#Address correspondence to Makoto Miyata, miyata@sci.osaka-

cu.ac.jp

ABSTRACT   *Spiroplasma* are bacteria with a helical cell shape and a lack of

peptidoglycan. They belong to the phylum *Tenericutes* that evolved from

Firmicutes, which includes the bacterium *Bacillus subtilis*. *Spiroplasma* swimming

motility is unrelated to widespread bacterial motilities, such as flagellar motility, and

is caused by helicity switching with kinks traveling along the helical cell body. The

swimming force is likely generated by five classes of bacterial actin homolog

MreBs involved in the helical bone structure. We analyzed sequences of

Spiroplasma MreBs (SMreBs) to clarify their phylogeny and sequence features.

The maximum likelihood method based on around 5,000 MreB sequences showed

that the phylogenetic tree was divided into several radiations: MreBs of

conventional bacteria, MreBs of candidate phyla radiation, and radiations for

Firmicutes MreB isoforms. SMreBs formed a clade adjacent to the radiation of

MreBH, an MreB isoform of *Firmicutes*. Sequence comparisons of SMreBs and

Bacillus MreBs were also performed to clarify the features of SMreB. In SMreBs 2

and 4, amino acids involved in the interactions for inter- and intra-protofilament

were significantly different from those in *Bacillus* MreBs. Catalytic glutamic acid

and threonine were changed to aspartic acid and lysine, respectively, in SMreB3.

SMreB5 had a significantly longer C-terminal region than the other MreBs, which

possibly forms a protein-protein interaction unique to *Spiroplasma*. A membrane-

binding region was not identified in most SMreBs 1 and 4, although it is conserved

in many walled bacterial MreBs. These features may support the functions

responsible for the unique mechanism of *Spiroplasma* swimming.

IMPORTANCE   *Spiroplasma*, a small bacterial group, is parasitic to plants and

arthropods. *Spiroplasma* species are globally widespread and swim via a "helicity

switching" mechanism. This mechanism is caused by fibril, a *Spiroplasma*-specific

protein, and five classes of MreBs, bacterial actin belonging to the same family as

major muscle protein. This study clarified the evolutional and structural features of

MreBs and provided possible clues to control the industrial damage caused by

Spiroplasma and develop manageable nano-devices to mimic helicity switching.

INTRODUCTION

The phylum *Tenericutes* consists of bacteria that evolved from the phylum

Firmicutes (1, 2). Currently, *Tenericutes* is comprised only of the class *Mollicutes*,

which includes the genera *Spiroplasma* and *Mycoplasma*. Species of these genera

are characterized by small genome size, low guanine and cytosine contents, being

mostly parasitic or commensal, and absence of peptidoglycan (3). In addition, they

cannot equip conventional machineries for bacterial motility, such as flagella or

type IV pili, due to the lack of a peptidoglycan layer (4). Instead, they have

developed three unique motility systems: *mobile* type gliding, *pneumoniae* type

gliding, and *Spiroplasma* swimming (4-11).

Spiroplasma have a helical cell shape and infect multi-cellular organisms, such

as plants and arthropods (12). They swim by rotating the cell body in viscous

environments, including host tissues. To rotate the cell body, *Spiroplasma* change

the helicity handedness at the front end by forming a kink, and propagate the kink

along the cell body from front to back (Fig 1A-B refer to videos in the following

articles) (11, 13-15). The helical cell shape originates from a flat ribbon structure

extending along the innermost line of the cell, and kink propagation is likely caused

by a structural change of the ribbon structure (Fig 1C) (13, 16-18). Electron

microscopy and proteome analyses have revealed that the flat ribbon is composed

of fibril, a *Spiroplasma* specific protein that forms a helical filament, and MreB

proteins, which are bacterial actin homologs (Fig 2A) (16-18). On the other hand,

seven copies of *mreB* genes have been found on the draft genome of a

Tenericutes bacterium, *Haloplasma contractile*, which performs characteristic

motility via tentacle-like structures extending from a coccoidal body (19, 20). Based

on these findings, MreBs are thought to be employed in the motility of *Tenericutes*

(16, 19). *Candidatus* Izimaplasma and *Mesoplasma* are spherical bacteria that

belong to *Tenericutes*. *Candidatus* Izimaplasma have three and *Mesoplasma* have

one *mreB* gene/s, respectively (19, 21, 22). While their motilities have not yet been

reported, it is possible that they use MreBs for motility.

As seen in eukaryotic actin, the MreB molecule has a U-shaped three

dimensional (3D) structure that can be further divided into four domains (Fig 2A)

(23, 24). The groove of the U-shaped structure functions as a nucleotide-binding

pocket, such as for ATP and GTP (Fig S3A) (23, 25-27). A previous study reported

that, in the presence of nucleotides, especially ATP, MreB polymerizes into non-

helical filaments with juxtaposed subunits in which two protofilaments interact

antiparallelly (Fig 2B) (23). Molecular dynamics (MD) simulations suggest that the

inter- and intra-protofilament interactions are comprised of four helices, two

strands, and three loops (H3, H5, and H8 shown in Fig 2A and Fig S3C for inter-

protofilament interaction; H9, S12, S13 and loops between S6-H1, S9-S10, and

S10-S11 shown in Fig 2A and Fig S3D for intra-protofilament interaction) (28).

Unlike the eukaryotic actin filament, MreB filaments bind to the cell membrane

directly (29). A previous study reported that many MreBs have two consecutive

hydrophobic amino acids for membrane binding in the hydrophobic loops in domain

IA (Fig 2A) (29). The study also reported that some MreBs of gram-negative

bacteria have an amphipathic helix at the N-terminus as an additional factor for

membrane binding (Fig 2A).

In various phyla of rod-shaped bacteria, MreB is well-conserved in the genomes

and is responsible for cell shape maintenance (30, 31). The MreB filament binds to

the membrane to form an elongasome complex with other proteins, which precisely

locates the peptidoglycan-synthesizing enzymes to retain the rod shape of the cells

(30, 31). Many bacterial species represented by *Escherichia coli* have a single

copy of the *mreB* gene on their genome (31), whereas the genome of *Bacillus*

subtilis has three genes to code MreB isoforms (MreB, Mbl, and MreBH), all of

which are employed in cell shape maintenance (32-35). The role of MreB is similar

to that of other bacterial actin homologs, such as ParM, MamK, and FtsA, which

are also responsible for the assignment of materials such as DNA, vesicles, and

cytoskeletal rings in cells (36).

As *Spiroplasma* MreBs (SMreBs) have a role that is distinct from those of other

bacterial actin homologs, sequence analyses among SMreBs and walled-bacterial

MreBs are useful for clarifying the origin and mechanism of *Spiroplasma*

swimming. While a previous study reported that *Spiroplasma* and *Haloplasma* form

two independent phylogenetic groups, the phylogenetical relationship among

SMreBs and walled-bacterial MreBs remains unclear (19). Moreover, amino acid

sequences specific for MreB functions, including nucleotide binding and hydrolysis,

interaction regions to form double-stranded filaments, and membrane binding

regions, have not yet been characterized for SMreBs.

In this study, we analyzed the sequences of SMreBs. Phylogenetic analyses

suggested that SMreBs evolved from MreBH in *Firmicutes*. Sequence comparisons

of SMreBs and *Bacillus* MreBs clarified their sequence features for ATP hydrolysis,

interactions for filament formation, and membrane binding.

RESULTS AND DISCUSSION

Evolutional relationships between *Spiroplasma* and other MreBs

We analyzed the phylogenetic relationships among SMreBs and other MreBs.

We first obtained 5,002 amino acid sequences of MreB family proteins through a

BLAST search, including all 170 non-redundant sequences of SMreB with those of

three new species (*S.* sp. TU-14, *S.* sp. ChiS and *S.* endosymbiont of *Megaselia*

nigra) (Table S1) (Data set 1) (37-40). Based on these sequences, we constructed

a maximum likelihood phylogenetic tree (Fig 3, S7). We adopted this phylogenetic

tree because the topology difference was indistinguishable between 4,000 and

5,002 sequences. The largest radiation of the phylogenetic tree consisted of MreB

family proteins of conventional bacteria, including *Bacillus* MreBs (BMreB). Similar

to the phylogenetic tree based on ribosomal proteins (1, 2), the radiation was

further divided into gram-positive bacteria, which have an outer membrane, and

gram-negative bacteria, which do not. Previous studies of MreB family proteins

have primarily focused on proteins in these radiations (31), unlike the present

study. In our phylogenetic tree, an MreB radiation of candidate phyla radiation

(CPR) (1, 2) (also known as Patescibacteria (41)) was formed next to the MreB

radiation of conventional bacteria. The CPR MreBs were split into two clades. Four

CPR candidate species in candidate phyla Yanofskybacteria, Taylorbacteria,

Zambryskibacteria, and Woesebacteria had MreBs of both clades, suggesting that

there are two types of MreBs that possess different roles in CPR.

Firmicutes bacteria have more than one copy of a gene to code MreB isoforms

on the genomes (32), forming three clades of MreB isoforms in addition to the

MreB radiations mentioned above. One was mostly composed of class *Bacilli* MreB

family proteins related to *Bacillus* Mbl. Another clade positioned between clades of

Bacillus Mbl and CPR MreB was primarily composed of proteins in class *Clostridia*,

distantly related to *Bacillus* MreB isoforms, suggesting the existence of a new

group of MreB isoforms. The clade including *Bacillus* MreBH (BMreBH) had a root

distinct from other clades. SMreBs formed a radiation near the BMreBH clade. In

addition, a clade of MreBs from *Haloplasma* and Izimaplasma species (HMreB)

was identified between clades of SMreB and BMreBH (Fig 3, S1). These

relationships suggest that the MreBs of *Tenericutes* species, SMreB and HMreB,

evolved from BMreBH.

Development of *Spiroplasma* swimming

Based on the phylogeny results and presence of genes for MreB isoforms in

each bacterium, we assessed the evolutional background prior to *Spiroplasma*

acquiring swimming motility using a phylogenetic tree of 16S rRNA, including newly

obtained *Spiroplasma* species (Fig 4, Data set 2) (37-40). A *Firmicutes* species, the

putative ancestor of *Tenericutes*, probably possessed MreB, Mbl, and MreBH as

the *Firmicutes* species closest to *Tenericutes* had these three MreB isoforms, as

reported previously (1, 2). The ancestral bacterium then likely lost MreB and Mbl

and duplicated MreBH before *Haloplasma* and other *Tenericutes* bacteria

separated. After the duplication of MreBH, *Haloplasma* and Izimaplasma likely

acquired HMreBs. HMreB may be involved in the characteristic motility of

Haloplasma (20), while that in Izimaplasma is unknown because its motility has not

yet been reported. A previous study suggested that *Haloplasma* and *Spiroplasma*

acquired their MreBs by independent gene duplications (19). However, HMreB was

located closer to SMreB and MreBH than to the other clades in our phylogenetic

tree (Fig 3, S1). Therefore, *Spiroplasma* likely acquired SMreBs by the duplication

of MreBH, as suggested in *Haloplasma* and Izimaplasma (Fig 4). Fibril, another

component of the ribbon structure of *Spiroplasma* cells, is specific for *Spiroplasma*

(Fig 1C) (11, 16, 17, 19, 42). Therefore, fibril was likely acquired after *Spiroplasma*

and Izimaplasma separated and may function with SMreBs to allow *Spiroplasma*

swimming.

The *Spiroplasma* genus can be divided into two subgroups: Apis and Citri-

Chrysopicola-Mirum (Citri) (19, 37-39). In terms of the *Spiroplasma* genomes,

duplications of SMreBs 1, 4, or 5 and deletions of SMreBs 1 or 2 were identified in

almost all species of the Apis group, unlike the Citri group, as previously reported

(19). *Spiroplasma* swimming has been studied by using Citri species (*S. melliferum*

and *S. eriocheiris*), both of which swim via similar kink propagations (11, 15-18,

42). The duplication and deletion of SMreBs in the Apis group may suggest their

diverse swimming formats.

Sequence comparisons among SMreBs and BMreBs

We compared the amino acid sequences of SMreB, HMreB, BMreB, and

BMreBH to characterize SMreBs (Fig 2C, S6). First, the conserved amino acids of

SMreBs 1-5, HMreB, and BMreBH sequences were compared with those of 48

sequences of BMreBs (Fig 2C, S6). Each SMreB type showed specific conserved

sequences different from those of BMreBs. SMreB 1-5 conserved 128, 187, 46,

210, and 121 amino acids in approximately 330 full-length amino acid sequences,

respectively. Of these amino acids, 50, 102, 21, 122, and 53 amino acids were

specific for each SMreB, respectively, and were not conserved in BMreBs (Table

S2, Fig 2C, S6). The specificity of these sequences to SMreBs suggests the

importance of the sequences for functions unique to SMreBs. Sequences of Citri

group SMreBs were more conserved than those of the Apis group, corresponding

to the phylogenetical spread among *Spiroplasma* species represented in the 16S

rRNA phylogeny (Table S3-4, Fig 4, S2, S6).

Characterization of SMreBs based on amino acid sequences

Next, to investigate the SMreB sequences corresponding to the sequences

conserved in walled bacteria (23, 28, 29, 43, 44), we predicted the ancestral

sequences and 3D structures for SMreBs 1-5, HMreB, BMreBH, and BMreB (Fig

S2, Data set 3). First, we focused on the 19 amino acids that construct the

nucleotide binding pocket (Fig 2A, D, S3A) (44) because MreB polymerization is

initiated when a nucleotide, such as ATP or GTP, binds to this pocket. In SMreB

sequences, most of the 12 amino acids conserved in BMreBH were also

conserved, except E212 (Fig 2C black dashed line boxes, Fig 2D, S6). We also

focused on the catalytic amino acids for nucleotide hydrolysis (23). While a

glutamic acid and threonine were conserved in almost all MreBs, interestingly, in

SMreB3, these sequences were changed to an aspartic acid and lysine,

respectively (Fig 2C cyan solid line boxes, Fig 2D, S6). As mutations on the

catalytic amino acids reduce depolymerization rates of eukaryotic actin and MamK

(45, 46), SMreB3 may also have different polymerization or depolymerization rates

than other SMreBs. Generally, an antibiotic A22 inhibits MreB polymerization by

interacting with the nucleotide in the MreB molecule (43). SMreBs also conserved

these target amino acids of A22, which was predicted by MD simulations based on

the crystal structure (Fig 2A, S3B, Fig 2C red dashed line boxes, Fig S6) (43).

Next, we focused on 87 amino acids involved in the inter- and intra-protofilament

interactions (H3, H5, H8, H9, S12, and S13 and loops between S6-H1, S9-S10,

and S10-S11 in Fig 2A, Fig S3C-D) (28). In these regions, 31, 50, 14, 69, and 32

amino acids were conserved in SMreBs 1-5, respectively (Fig 2C light green and

cyan dashed line boxes, Fig 2E-F, S6). In these amino acids, 15, 26, 6, 40, and 16

sequences were not conserved in BMreBs, respectively (white characters in light

green and cyan dashed line boxes in Fig 2C). As these regions in a walled-

bacterial MreB move toward an adjacent MreB molecule to interact with each other

(28), these amino acid substitutions of SMreBs, especially SMreBs 2 and 4, may

affect the flexibility and interaction strength of these regions, allowing them to show

polymerization characteristics different to those of walled-bacterial MreB.

We also characterized regions required for membrane binding. Two consecutive

hydrophobic amino acids are conserved in many walled-bacterial MreBs (Fig 2A).

However, SMreBs 1 and 4 of the Citri group did not have hydrophobic amino acids

(Fig 2C red solid line box, Fig S6) (23, 29). Moreover, the hydrophobic amino acids

cannot be found in 46 out of 88 sequences of the Apis group SMreBs 1, 3, 4, and 5

(Fig S6). These SMreBs may not directly interact with the membrane. All other

SMreBs (Apis group SMreB2 and Citri group SMreBs 2, 3, and 5) had hydrophobic

amino acids (Fig 2C, S6). We next focused on the N-terminal amphipathic helix

found in some MreBs of gram-negative bacteria (Fig 2A). All N-terminal regions of

the Citri group SMreB3, except for *S. melliferum* strain AS576 (QCO24517.1), were

predicted to take amphipathic helices (Fig 2C light green solid line, Fig S4A-F).

Interestingly all SpeMreB3 possessed a threonine at the hydrophobic sides (Fig

S4G, S6). This suggests that Citri group SMreB3 proteins also have affinity to the

membrane through the helix.

The lengths of C-terminal sequences varied among 133 SMreBs, and over 79%

of SMreBs were eight amino acids longer than that of BMreB (Fig S5, S6, Table

S5). In particular, SMreB5 had the longest C-terminal sequence of all SMreBs

(Table S5). Such terminal extension is a feature reminiscent of a bacterial tubulin

homolog, FtsZ, featured with a long unstructured region known as C-terminal

Conserved Peptide (CCP) (47). This is the interaction region for several proteins,

including FtsA, ZipA, and ZapD, involved in Z-ring formation, which causes cell

constriction for bacterial cytokinesis (47-50). The extension of the SMreB C-termini

may play a role in the protein-protein interactions of *Spiroplasma*, as observed in

FtsZ.

CONCLUSIONS

In this study, we suggested that the five classes of SMreBs evolved from MreBH

(Fig 3) and clarified that the classes can be characterized by polymerization,

membrane binding, and protein interactions. Based on these results, we suggest

the functions for individual SMreBs. SMreBs 2, 3, and 4 may form filaments of

sufficient size to be observed via electron microscopy, as reported previously (Fig

2C-F, S6) (17, 18). Citri group SMreBs 2, 3, and 5 can bind to membranes, but Citri

group SMreBs 1 and 4 cannot (Fig 2C, S6). SMreB5 may interact with other

proteins, such as fibril protein via their C-terminal regions (Fig 2C, Table S5, S6).

These findings would allow the clarification of the origin and mechanism of

Spiroplasma swimming.

MATERIALS AND METHODS

Phylogenetical analyses

A protein BLAST search on non-redundant protein sequences was performed to

collect 5,000 amino acid sequences of MreB in ascending order of E value by

templating MreB3 of *S. eriocheiris* (NCBI accession: WP_047791951.1) on July 14,

2019. Sequences of *B. subtilis* MreB isoforms annotated to MreB (AQR87043.1)

and MreBH (BAA07047.1) were added to the 5,000 sequences as representative

sequences, although the closest sequences were identified by the BLAST search

(WP_039074652.1; 99.11% identical with AQR87043.1, WP_061522102.1; 95.52%

identical with BAA07047.1). The nucleotide sequences of 16S rRNA were identified

by independent searches in the GenBank database. The sequence alignments and

construction of phylogenetic trees were performed via the MUSCLE program and

maximum likelihood method, respectively, using MEGA-X ver. 10.1 (51). Bootstrap

supports were estimated from 100 samples of alignment. To estimate the ancestor

of MreBs in each group, maximum likelihood phylogenetic trees by each MreB

group were constructed with a *B. subtilis* MreB (an *E. coli* MreB for the ancestor

estimation of *Bacillus* MreB) as an outgroup, and the ancestral sequences were

estimated by the ancestor estimation program of MEGA-X ver. 10.1 (51).

Structure prediction and sequence analyses

All monomeric MreB structures in this study were predicted using Rosetta

comparative modeling by templating a structure of *Caulobacter crescentus* MreB

(PDB ID: 4CZJ), using Robetta software (23, 52). Sequence comparisons among

Spiroplasma, *Haloplasma*, Izimaplasma, and *Bacillus* MreB isoforms were

performed by Clustal Omega (53). Duplicated SMreBs 1, 4, and 5 in the Apis group

species were incorporated into the analyses. Seven of the 19 amino acid

sequences contained in SMreB3 of the Apis group had no annotation. Since these

proteins were coded in a position suggesting SMreB3 as an ortholog on the

genome, they were included in the analyses. SMreB3 of the *S. melliferum* strain

CH-1 (NCBI accession: AHA83319.1) was excluded from the analyses because it

was approximately 130 amino acids shorter than the other SMreB3 sequences

comprising approximately 330 amino acids. Two sequences, MreB4 of *S.*

eriocheiris strain CCTCC_M_207170 (AHF58342.1) and a protein from *S.*

culicicola, which is unannotated but can be classified to MreB3

(WP_025363622.1), were not used for the analyses because the initiation codon

was miss-assigned. The predictions of secondary structures and amphipathic

helices were performed using PSIPRED 4.0 and AmphipaSeeK software,

respectively (54-56). Sequence identity and similarity were calculated as the ratio

of amino acids with identity and strong similarity defined by the Gonnet PAM 250

matrix over the total amino acid number excluding gap regions.

ACKNOWLEDGEMENTS

We thank Yuya Sasajima and Hana Kiyama at the Graduate School of Science,

Osaka City University, Japan for helpful discussion. The electron microscope

image of *Spiroplasma* (Fig 1A) and the schematic image of the ribbon structure

(Fig 1C) were courtesy of Yuya Sasajima (Osaka City University, Japan). This

study was supported by a Grants-in-Aid for Scientific Research (A and C) (MEXT

KAKENHI, Grant Numbers JP17H01544 to MM and JP20K06591 to IF), JST

CREST (Grant Number JPMJCR19S5) to MM, Research Foundation of Opto-

Science and Technology to IF, and the Osaka City University (OCU) Strategic

Research Grant 2019 to IF. DT is a recipient of a Japanese scholarship of JEES

Kureha (Toyobo) Scholarship.

REFERENCES

1.   Castelle CJ, Banfield JF. 2018. Major new microbial groups expand diversity

and alter our understanding of the tree of life. Cell 172:1181-1197.

2.   Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ,

Butterfield CN, Hernsdorf AW, Amano Y, Ise K, Suzuki Y, Dudek N, Relman

DA, Finstad KM, Amundson R, Thomas BC, Banfield JF. 2016. A new view

of the tree of life. Nat Microbiol 1:16048.

3.   Razin S, Hayflick L. 2010. Highlights of mycoplasma research--an historical

perspective. Biologicals 38:183-90.

4.   Miyata M, Robinson RC, Uyeda TQP, Fukumori Y, Fukushima S, Haruta S,

Homma M, Inaba K, Ito M, Kaito C, Kato K, Kenri T, Kinosita Y, Kojima S,

Minamino T, Mori H, Nakamura S, Nakane D, Nakayama K, Nishiyama M,

Shibata S, Shimabukuro K, Tamakoshi M, Taoka A, Tashiro Y, Tulum I, Wada

H, Wakabayashi K. 2020. Tree of motility - A proposed history of motility

systems in the tree of life. Genes to Cells 25:6-21.

5.   Nishikawa MS, Nakane D, Toyonaga T, Kawamoto A, Kato T, Namba K,

Miyata M. 2019. Refined mechanism of *Mycoplasma mobile* gliding based

on structure, ATPase activity, and sialic acid binding of machinery. mBio

10:e02846-19.

6.   Mizutani M, Miyata M. 2019. Behaviors and energy source of *Mycoplasma*

gallisepticum gliding. J Bacteriol 201:e00397-19.

7.   Mizutani M, Tulum I, Kinosita Y, Nishizaka T, Miyata M. 2018. Detailed

analyses of stall force generation in *Mycoplasma mobile* gliding. Biophys J

114:1411-1419.

8.   Kinosita Y, Miyata M, Nishizaka T. 2018. Linear motor driven-rotary motion

of a membrane-permeabilized ghost in *Mycoplasma mobile*. Sci Rep

8:11513.

9.   Miyata M, Hamaguchi T. 2016. Prospects for the gliding mechanism of

Mycoplasma mobile. Curr Opin Microbiol 29:15-21.

10.   Miyata M, Hamaguchi T. 2016. Integrated information and prospects for

gliding mechanism of the pathogenic bacterium *Mycoplasma pneumoniae*.

Front Microbiol 7:960.

11.   Shaevitz JW, Lee JY, Fletcher DA. 2005. *Spiroplasma* swim by a processive

change in body helicity. Cell 122:941-5.

12.   Lo WS, Huang YY, Kuo CH. 2016. Winding paths to simplicity: genome

evolution in facultative insect symbionts. FEMS Microbiol Rev 40:855-874.

13.   Nakane D, Ito T, Nishizaka T. 2020. Coexistence of two chiral helices

357   produces kink translation in *Spiroplasma* swimming. J Bacteriol 202:e00735-

358   19.

359   14.   Terahara N, Tulum I, Miyata M. 2017. Transformation of crustacean

360   pathogenic bacterium *Spiroplasma eriocheiris* and expression of yellow

361   fluorescent protein. Biochem Biophys Res Commun 487:488-493.

362   15.   Wada H, Netz RR. 2009. Hydrodynamics of helical-shaped bacterial motility.

363   Phys Rev E 80:021921.

364   16.   Liu P, Zheng H, Meng Q, Terahara N, Gu W, Wang S, Zhao G, Nakane D,

365   Wang W, Miyata M. 2017. Chemotaxis without conventional two-component

366   system, based on cell polarity and aerobic conditions in helicity-switching

367   swimming of *Spiroplasma eriocheiris*. Front Microbiol 8:58.

368   17.   Trachtenberg S, Dorward LM, Speransky VV, Jaffe H, Andrews SB,

369   Leapman RD. 2008. Structure of the cytoskeleton of *Spiroplasma melliferum*

370   BC3 and its interactions with the cell membrane. J Mol Biol 378:778-89.

371   18.   Kurner J, Frangakis AS, Baumeister W. 2005. Cryo-electron tomography

372   reveals the cytoskeletal structure of *Spiroplasma melliferum*. Science

307:436-8.

19.   Ku C, Lo WS, Kuo CH. 2014. Molecular evolution of the actin-like MreB

protein gene family in wall-less bacteria. Biochem Biophys Res Commun

446:927-32.

20.   Antunes A, Alam I, El Dorry H, Siam R, Robertson A, Bajic VB, Stingl U.

2011. Genome sequence of *Haloplasma contractile*, an unusual contractile

bacterium from a deep-sea anoxic brine lake. J Bacteriol 193:4551-2.

21.   Skennerton CT, Haroon MF, Briegel A, Shi J, Jensen GJ, Tyson GW, Orphan

VJ. 2016. Phylogenomic analysis of *Candidatus* 'Izimaplasma' species: free-

living representatives from a *Tenericutes* clade found in methane seeps.

ISME J 10:2679-2692.

22.   Tully JG, Whitcomb RF, Hackett KJ, Rose DL, Henegar RB, Bove JM, Carle

P, Williamson DL, Clark TB. 1994. Taxonomic descriptions of 8 new non-

sterol-requiring *Mollicutes* assigned to the genus *Mesoplasma*. Int J Syst

Bacteriol 44:685-693.

23.   van den Ent F, Izore T, Bharat TA, Johnson CM, Löwe J. 2014. Bacterial

actin MreB forms antiparallel double filaments. eLife 3:e02634.

24.   van den Ent F, Amos LA, Löwe J. 2001. Prokaryotic origin of the actin

cytoskeleton. Nature 413:39-44.

25.   Nurse P, Marians KJ. 2013. Purification and characterization of *Escherichia*

coli MreB protein. J Biol Chem 288:3469-3475.

26.   Mayer JA, Amann KJ. 2009. Assembly properties of the *Bacillus subtilis*

actin, MreB. Cell Motility and the Cytoskeleton 66:109-118.

27.   Bean GJ, Amann KJ. 2008. Polymerization properties of the *Thermotoga*

maritima actin MreB: Roles of temperature, nucleotides, and ions.

Biochemistry 47:826-835.

28.   Awuni E, Mu Y. 2019. Effect of A22 on the conformation of bacterial actin

MreB. Int J Mol Sci 20:E1304.

29.   Salje J, van den Ent F, de Boer P, Löwe J. 2011. Direct membrane binding

by bacterial actin MreB. Mol Cell 43:478-87.

30.   Yulo PRJ, Hendrickson HL. 2019. The evolution of spherical cell shape;

progress and perspective. Biochem Soc Trans 47:1621-1634.

31.   Shi H, Bratton BP, Gitai Z, Huang KC. 2018. How to build a bacterial cell:

MreB as the foreman of *E. coli* construction. Cell 172:1294-1305.

32.   Schirner K, Errington J. 2009. Influence of heterologous MreB proteins on

408   cell morphology of *Bacillus subtilis*. Microbiology 155:3611-21.

409   33.   Kawai Y, Asai K, Errington J. 2009. Partial functional redundancy of MreB

410   isoforms, MreB, Mbl and MreBH, in cell morphogenesis of *Bacillus subtilis*.

411   Mol Microbiol 73:719-31.

412   34.   Carballido-Lopez R, Formstone A, Li Y, Ehrlich SD, Noirot P, Errington J.

413   2006. Actin homolog MreBH governs cell morphogenesis by localization of

414   the cell wall hydrolase LytE. Dev Cell 11:399-409.

415   35.   Jones LJ, Carballido-Lopez R, Errington J. 2001. Control of cell shape in

416   bacteria: helical, actin-like filaments in *Bacillus subtilis*. Cell 104:913-22.

417   36.   Wagstaff J, Löwe J. 2018. Prokaryotic cytoskeletons: protein filaments

418   organizing small cells. Nat Rev Microbiol 16:187-201.

419   37.   Ballinger MJ, Perlman SJ. 2019. The defensive *Spiroplasma*. Curr Opin

420   Insect Sci 32:36-41.

421   38.   Tsai YM, Chang A, Kuo CH. 2018. Horizontal gene acquisitions contributed

422   to genome expansion in insect-symbiotic *Spiroplasma clarkii*. Genome Biol

423   Evol 10:1526-1532.

39.   Paredes JC, Herren JK, Schupfer F, Marin R, Claverol S, Kuo CH, Lemaitre

B, Beven L. 2015. Genome sequence of the *Drosophila melanogaster* male-

killing *Spiroplasma* strain MSRO endosymbiont. mBio 6:e02437-14.

40.   Gasparich GE. 2002. Spiroplasmas: evolution, adaptation and diversity.

Front Biosci 7:d619-40.

41.   Rinke C, Schwientek P, Sczyrba A, Ivanova NN, Anderson IJ, Cheng JF,

Darling A, Malfatti S, Swan BK, Gies EA, Dodsworth JA, Hedlund BP,

Tsiamis G, Sievert SM, Liu WT, Eisen JA, Hallam SJ, Kyrpides NC,

Stepanauskas R, Rubin EM, Hugenholtz P, Woyke T. 2013. Insights into the

phylogeny and coding potential of microbial dark matter. Nature 499:431-7.

42.   Cohen-Krausz S, Cabahug PC, Trachtenberg S. 2011. The monomeric,

tetrameric, and fibrillar organization of Fib: the dynamic building block of the

bacterial linear motor of *Spiroplasma melliferum* BC3. J Mol Biol 410:194-

213.

43.   Awuni Y, Jiang S, Robinson RC, Mu Y. 2016. Exploring the A22-bacterial

actin MreB interaction through molecular dynamics simulations. J Phys

Chem B 120:9867-74.

44.   Aylett CH, Löwe J, Amos LA. 2011. New insights into the mechanisms of

442   cytomotive actin and tubulin filaments. Int Rev Cell Mol Biol 292:1-71.

443   45.   Ozyamak E, Kollman J, Agard DA, Komeili A. 2013. The bacterial actin

444   MamK: in vitro assembly behavior and filament architecture. J Biol Chem

445   288:4265-77.

446   46.   Iwasa M, Maeda K, Narita A, Maeda Y, Oda T. 2008. Dual roles of Gln(137)

447   of actin revealed by recombinant human cardiac muscle alpha-actin

448   mutants. J Biol Chem 283:21045-21053.

449   47.   Gardner KA, Moore DA, Erickson HP. 2013. The C-terminal linker of

450   *Escherichia coli* FtsZ functions as an intrinsically disordered peptide. Mol

451   Microbiol 89:264-75.

452   48.   Wang MZ, Fang C, Ma B, Luo XX, Hou Z. 2020. Regulation of cytokinesis:

453   FtsZ and its accessory proteins. Curr Genet 66:43-49.

454   49.   Huang KH, Mychack A, Tchorzewski L, Janakiraman A. 2016.

455   Characterization of the FtsZ C-Terminal variable (CTV) region in Z-Ring

456   assembly and interaction with the Z-Ring stabilizer ZapD in *E. coli*

457   cytokinesis. PLoS ONE 11:e0153337.

50.   Erickson HP, Anderson DE, Osawa M. 2010. FtsZ in bacterial cytokinesis:

cytoskeleton and force generator all in one. Microbiol Mol Biol Rev 74:504-

28.

51.   Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular

evolutionary genetics analysis across computing platforms. Mol Biol Evol

35:1547-1549.

52.   Song Y, DiMaio F, Wang RY, Kim D, Miles C, Brunette T, Thompson J, Baker

D. 2013. High-resolution comparative modeling with RosettaCM. Structure

21:1735-42.

53.   Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R,

McWilliam H, Remmert M, Soding J, Thompson JD, Higgins DG. 2011. Fast,

scalable generation of high-quality protein multiple sequence alignments

using Clustal Omega. Mol Syst Biol 7:539.

54.   Buchan DWA, Jones DT. 2019. The PSIPRED protein analysis workbench:

20 years on. Nucleic Acids Res 47:W402-W407.

55.   Sapay N, Guermeur Y, Deleage G. 2006. Prediction of amphipathic in-plane

membrane anchors in monotopic proteins using a SVM classifier. BMC

Bioinformatics 7:255.

56.   Jones DT. 1999. Protein secondary structure prediction based on position-

specific scoring matrices. J Mol Biol 292:195-202.

 FIGURE LEGENDS

Fig 1 *Spiroplasma* swimming. (A) Electron microscope image of *Spiroplasma*.

Swimming directions are indicated with yellow arrows. (B) Swimming scheme.

Spiroplasma swim via the rotation of the cell body caused by kink propagation

along the cell axis from front to back, which causes helicity shifts of right to left

(open triangle) and left to right (closed triangle). (C) Schematic image of the ribbon

structure responsible for swimming motility. The structure composed of SMreBs

and fibril protein is presented as a gold ribbon in a cell.

Fig 2 Sequence analyses of SMreBs. (A) Structure of *Bacillus subtilis* MreB

predicted from the amino acid sequence (NP_390681.2). The structure is

composed of 15 strands and 13 helices shown as S1-S15 and H1-H13,

respectively. Individual subdomains are colored blue (IA), yellow (IB), red (IIA), and

green (IIB). An N-terminal amphipathic helix carried by some gram-negative

bacterial MreBs is presented as a rectangle bi-colored in blue and yellow. The

nucleotide binding pocket, catalytic region for nucleotide hydrolysis, A22 binding

region, inter- and intra-protofilament interaction regions, hydrophobic loop,

membrane binding amino acids in the hydrophobic loop, and N-terminal

amphipathic helix are marked with a rectangle common in style to those in Fig 2C.

(B) Filamentous structure of MreB extracted from a crystal structure of *Caulobacter*

crescentus (PDB ID: 4CZJ). It forms a non-helical filament with the juxtaposed

subunits in which two protofilaments interact antiparallelly. (C) Multiple alignment of

SMreBs 1-5, HMreB, BMreB, and BMreBH performed for MreBs 1-5 of *S.*

eriocheiris, KQ51_00688 in Izimaplasma sp. HR1, *B. subtilis* MreB, and *B. subtilis*

MreBH. The conservation levels for amino acids calculated from multiple

alignments of each MreB are indicated by the color gradient of orange markers

(amino acids calculated as identical, strongly conserved, and weakly conserved

based on the Gonnet PAM 250 matrix are indicated by deep, medium, and light

orange markers, respectively. Unrelated amino acids based on the matrix are not

colored by makers). In SMreBs 1-5, HMreB, and BMreBH, the conserved amino

acids, whose corresponding amino acids in BMreB were conserved as a different

amino acid or were not conserved, are indicated by white characters. Domains IA,

IB, IIA, and IIB are identified by underbars of coordinated colors with panel (A).

Sequences important for the specific function of MreB are indicated by rectangles

with different styles. (D-F) Focused amino acids are shown as 3D structures. The

structures for SMreBs 1-5, BMreB, HMreB, and BMreBH were marked as 1-5, B,

HI, and BH, respectively. The amino acids were colored red, pink, blue, or gold,

based on the conservation levels compared to BMreBs (red: conserved amino

acids whose corresponding amino acids in BMreB were conserved as different

amino acids, pink: conserved amino acids whose corresponding amino acids in

BMreB were not conserved, blue: not conserved amino acids whose corresponding

amino acids in BMreB were conserved, gold: conserved amino acids whose

corresponding amino acids in BMreB were conserved as the same amino acid).

The number of helices and strands is indicated for colored parts. (D) Nucleotide

binding pocket. The backbone atoms of catalytic amino acids for nucleotide

hydrolysis are indicated by spheres. (E) Inter-protofilament interaction region. (F)

Intra-protofilament interaction region.

Fig 3 No rooted maximum likelihood phylogenic tree for 5,002 amino acid

sequences of MreB family proteins. Each clade is colored according to the major

phyla (or classes). Major and minor taxa in each clade are indicated without and

with parentheses, respectively. The branches of MreB isoforms in *Bacillus subtilis*

are marked by red circles. The scale bar is in units of the number of amino acid

substitutions per site.

Fig 4 Maximum likelihood phylogeny for 16S rRNA of the organisms that are

related to the evolution of *Spiroplasma*. Sequences for 16S rRNA of proteobacteria

(*E. coli*, *Caulobacter vibrioides*, and *Helicobacter pylori*) were used as the

outgroup. Bootstrap support values are indicated by circles on nodes, solid circles

for support of 89% and above, and open circles for 60% to 89%. Stars and

triangles marked for each species indicate one duplication and deletion for genes

of MreBs colored differently as yellow: SMreB1, green: SMreB2, black: SMreB4,

and red: SMreB5. The scale bar is in units of the number of amino acid

substitutions per site.

Figure 1

Figure 2

Conservation level ■ > ■ > ▢ > ☐

Figure 3

Figure 4