In utero gene transfer system for embryos before neural tube closure reveals a role

for Hmga2 in the onset of neurogenesis

Naohiro Kuwayama[1], Yusuke Kishi[1,*], Yurie Maeda[1], Yurie Nishiumi[1], Yutaka Suzuki[2],

Haruhiko Koseki[3], Yusuke Hirabayashi[4], and Yukiko Gotoh[1,5]

[1]Graduate School of Pharmaceutical Sciences, The University of Tokyo, Tokyo 113-

0033, Japan

[2]Department of Computational Biology, Graduate School of Frontier Sciences, The

University of Tokyo, Tokyo 277-8561, Japan

[3]Laboratory for Developmental Genetics, RIKEN Center for Integrative Medical

Sciences (RIKEN-IMS), Tsurumi-ku, Yokohama 230-0045, Japan

[4]Graduate School of Engineering, The University of Tokyo, Tokyo 113-0033, Japan

[5]International Research Center for Neurointelligence (WPI-IRCN), The University of

Tokyo, Tokyo 113-0033, Japan

*Correspondence: ykisi@mol.f.u-tokyo.ac.jp

Abstract

During the early stage of mammalian neural development, neuroepithelial cells (NECs) proliferate and increase their pool size before switching their fate to become neurogenic radial glial progenitors (RGPs). The timing of this expansion-to-neurogenic phase transition is strictly regulated so as to determine the proper number of progenitors and differentiated cell types that constitute the brain. The molecular mechanism underlying this switch has remained poorly understood, however, in part because of the difficulty associated with manipulation of gene expression in NECs before their transition to RGPs, which occurs before neural tube closure. We have now developed a simple and efficient method to manipulate gene expression in mouse neocortical NECs in the expansion phase by in utero injection of viral vectors at embryonic day 7.0 to 8.0. With the use of this method, we found that knockdown of the chromatin-associated protein Hmga2 in NECs inhibited the onset of the neurogenic phase in the neocortex. This effect of Hmga2 knockdown was accompanied by inhibition of the expression of a set of genes targeted by Polycomb group proteins, which repress neurogenic genes in cortical progenitors. Our study thus establishes a novel method for manipulation of gene expression in the early stage of mouse brain development as well as uncovers a key molecular player in the generation of neurogenic progenitors in the developing mouse neocortex.

Introduction

The neocortex is the seat of higher cognitive function and has undergone evolutionary expansion in mammals. It originates from a pseudostratified sheet of neuroepithelium present at the anterior end of the neural plate before formation of the neural tube. In the course of neocortical development, neural progenitors undergo sequential waves of expansion, neurogenic and gliogenic phases (Gotz and Huttner, 2005; Hirabayashi and Gotoh, 2010; Kriegstein and Alvarez-Buylla, 2009; Mukhtar and Taylor, 2018). During the expansion phase, neural progenitors, also known as neuroepithelial cells (NECs), divide symmetrically to give rise to two progenitors per division (Rakic, 1995). This symmetric division increases the progenitor pool and results in lateral expansion of the neocortex. At the onset of neurogenesis in the mouse neocortex, NECs transform into neurogenic radial glial cells (RGCs), which mainly divide asymmetrically to give rise to one progenitor and one neuron or neuron-committed intermediate progenitor cell per division (Gao et al., 2014; Gotz and Huttner, 2005; Hartfuss et al., 2001; Miyata et al., 2002; Noctor et al., 2001). The elongated basal (radial) processes of RGCs also support the migration of newborn neurons to the pial surface, which results in radial expansion of the neocortex (Rakic, 1972). Clonal analysis of the mouse neocortex has revealed that a neurogenic RGC typically produces 8~9 cortical excitatory neurons through asymmetric division and then transforms into a gliogenic RGC (Gao et al., 2014). To avoid confusion, we will hereafter use the term neural stem-progenitor cells (NPCs) for both NECs and RGCs, given that many, if not all, of these cells are neural stem cells that self-renew and eventually produce both neurons and glial cells and that various nomenclatures are currently in use in the field.

The expansion-to-neurogenic phase (E-to-N) transition of NPC fate takes place at around embryonic day (E) 10 in the mouse neocortex. The timing of this E-to-N transition is key to determination of the number of neurons generated and the size of the cortex, given that the duration of the expansion phase determines the size of the founder NPC pool for corticogenesis. With regard to the mechanism of this transition, certain secreted (extrinsic) factors have been found to play a role, with one such factor being all-*trans* retinoic acid (atRA) released from the meninges. In mice, a reticulum of meningeal cells covers the neocortex between E9 and E10 (McLone and Bondareff, 1975) and secretes atRA, which promotes the onset of neurogenesis while suppressing

neuroepithelial elongation (Siegenthaler et al., 2009). Fibroblast growth factor (FGF) 10

is transiently expressed in anterior NPCs from E9.5 to E11.5 and, together with its

receptor FGFR2, also promotes the E-to-N transition in the anterior portion of the

neocortex (Sahara and O'Leary, 2009; K. Yoon et al., 2004). Moreover, an increase in

oxygen concentration as a result of periventricular ingrowth of blood vessels from

E10.5 to E12.5 also appears to be associated with the E-to-N transition (Lange et al.,

2016). In addition to these extrinsic factors, cell-intrinsic programs likely regulate the

E-to-N transition, given that in vitro cultures of mouse cortical NPCs as well as of

human embryonic stem (ES) cell– or primate induced pluripotent stem (iPS) cell–

derived NPCs undergo a directional transition from proliferative (symmetric) to

differentiating (asymmetric) divisions (Eiraku et al., 2008; Gaspard et al., 2008; Otani et

al., 2016; Qian et al., 1998; Shen et al., 2006; Shi et al., 2012). The molecular basis of

cell-intrinsic programs that regulate the E-to-N transition of NPC fate (in particular, the

timing of this transition) has remained largely unclear, however. Only a few candidate

intracellular mechanisms, such as those involving Notch-Hes signaling and the BAF

chromatin-remodeling complex (Anthony et al., 2005; Gaiano et al., 2000; Hatakeyama

et al., 2004; Nguyen et al., 2018), have been proposed to date.

One possible reason that the molecular basis of the E-to-N transition has

remained relatively unexplored compared with that of neurogenesis or gliogenesis later

in development is the difficulty of gene manipulation at this early stage of brain

development. At late embryonic stages (from ~E12), a gene of interest can be readily

introduced into neocortical NPCs by in utero electroporation of plasmids (Tabata and

Nakajima, 2001) or by injection of recombinant viruses into the ventricles. However,

these methods cannot be applied to earlier stages before the E-to-N transition (such as at

E8.0) because neural tube closure does not take place until E8.5 to E9.5 in the mouse

forebrain (so there is no ventricle at E8.0) and the presumptive neocortex is too small to

manipulate. Ultrasound-guided injection of viruses into the mouse nervous system at

E8.5 was introduced to overcome this difficulty (Gaiano et al., 1999). However, this

method requires an ultrasound backscatter microscopy system, and retrovirus injection

with this approach at E8.5 frequently resulted in developmental defects of neural tube

closure or death of the embryo (41 out of 54). We therefore set out to establish a new

method that does not require costly equipment and which has a high survival rate

without the induction of gross developmental defects.

A hallmark of NPCs in the neurogenic phase is their neurogenic potential—

that is, they maintain an undifferentiated state but are able to differentiate into neurons

in response to appropriate cues. The chromatin state of NPCs underlies this neurogenic

potential (Kishi and Gotoh, 2018; Nakagawa et al., 2020; Tsuboi et al., 2019).

Polycomb group (PcG) proteins temporarily suppress the expression of neurogenic

genes in NPCs through histone modification during the neurogenic phase, and this

suppression is relieved on neuronal differentiation (Albert et al., 2017; Fasano et al.,

2007; Hirabayashi et al., 2009; Morimoto-Suzki et al., 2014; Pereira et al., 2010; Tsuboi

et al., 2018). At later developmental stages (during the gliogenic phase), however, NPCs

lose their neurogenic potential as a result of persistent (robust) repression of neurogenic

genes mediated in part through polymerization of PcG proteins (Hirabayashi et al.,

2009; Tsuboi et al., 2018). High mobility group A (Hmga) proteins 1 and 2 are also

chromatin modifiers that contribute to neurogenic potential by regulating the expression

of neurogenesis-related genes in NPCs during the neurogenic phase (Fujii et al., 2013;

Kishi et al., 2012; Sakai et al., 2019). The levels of Hmga proteins in NPCs are high in

the neurogenic phase but gradually decrease during later neocortical development

(Kishi et al., 2012; Nishino et al., 2008; Sanosaka et al., 2008), with this decline

accounting in part for the loss of neurogenic potential in gliogenic NPCs (Kishi et al.,

2012). However, it has remained unknown whether Hmga proteins play a role in the E-

to-N transition of NPC fate in the early embryonic mouse neocortex.

We here describe a new and simple method for the introduction of a gene of

interest into ectodermal and endodermal cells of mouse embryos as early as E7.0 or

E8.0. This method is based on in utero injection of viral vectors into amniotic fluid and

is associated with a high survival rate without inducing gross developmental

abnormalities. Moreover, as a proof of concept, we have identified a key regulator of

the E-to-N transition with the use of this new method. We found that *Hmga2* expression

in neocortical NPCs peaks at around the E-to-N transition (at E10), and that knockdown

of Hmga2 from E8.0 delayed the onset of neurogenesis and shifted the transcriptional

profile of NPCs at this transition closer to that in the expansion phase. Importantly,

knockdown of Hmga2 from E8.0 attenuated the expression of a subset of PcG target

genes in NPCs at E11.0. Our results suggest that an increase in Hmga2 expression

triggers the onset of the neurogenic phase in part through activation of neurogenesis-

related PcG target genes.

Results

Development of a simple and efficient method for manipulation of gene expression

in mouse embryos before neural tube closure

To investigate the E-to-N transition in the mouse neocortex, we set out to develop a

simple and efficient method for the introduction of a gene of interest before neural tube

closure, ideally before E8.5. To this end, instead of the ultrasound microscopy system

adopted in a previous study (Gaiano et al., 1999), we used an optical fiber to visualize

the inner structure of the uterus including the amnion and embryo (Figure 1). Optical

fiber–based visualization of the ventricles of embryos from E9.5 was previously

described for targeted DNA injection and electroporation after neural tube closure

(Matsui et al., 2011). We found that lighting with this system was sufficient to allow

injection of a dye inside the amnion, although it was not possible to directly see the

amnion within the uterus. We first injected carboxyfluorescein succinimidyl ester

(CFSE), a cell-permeable dye that becomes fluorescent after reaction with intracellular

proteins (Quah and Parish, 2010; Telley et al., 2016), into the uterus at E8.0 (Figure S1)

To minimize possible damage caused by the injection, we inserted a sharpened capillary

at the center of the uterus at an angle perpendicular to the uterine wall while it was

positioned as shown in Figure 1d. We injected a small volume (~0.5 μl) of solution at a

depth of 1 to 2 mm below the uterine wall, which reproducibly resulted in transfer of the

solution into the amniotic fluid. The time of surgery is relatively short (~20 min)

because of the simplicity of the method, which may contribute to the minimal damage

caused to embryos. After 3 days (at E11.0), fluorescence microscopy without sectioning

revealed that 77% of injected embryos were labeled with CFSE (Figure S1a and b).

About 80% of embryos survived without gross developmental malformations at E11.0.

We also injected CFSE into amniotic fluid at E7.0, which again yielded a high rate of

embryo labeling (64%) and a high rate of survival without gross developmental

malformation (89%) at E9.0 (Figure S1b).

We next performed gene introduction via retroviral vectors with this method.

We first injected a concentrated suspension of a retrovirus encoding green fluorescent

protein (GFP) into the uterus at E8.0, and we found that 89% of embryos survived

without gross abnormalities and that 74% of embryos expressed GFP at E11.0 (Figure

2a and c). Injection of the retrovirus at E7.0 also yielded high rates of GFP-expressing

embryos and surviving embryos at E9.0 (Figure 2b and c). Flow cytometric analysis of

cortical tissue obtained from E12.0 embryos that had been injected with the retrovirus

for GFP at E8.0 showed that ~70% of the cortical cells were positive for GFP in

successful cases (Figure 2d). Immunohistofluorescence analysis of sagittal sections of

E10.0 embryos that had been injected with the retrovirus encoding GFP at E8.0 revealed

that most tissues exposed to amniotic fluid expressed GFP (Figure 2e–g). The labeled

tissues thus included the central nervous system (forebrain, midbrain, hindbrain, and

spinal cord) as well as the epidermis and endodermal tissues such as those of digestive

and respiratory organs. These results suggested that our method is also suitable for gene

introduction into such ectodermal and endodermal tissues at this early stage of mouse

development.

Given that in utero electroporation of plasmids is in general superior to

retroviral infection as a means of gene introduction with regard to efficiency and

expression level, we next injected a plasmid encoding mCherry with our new method

and performed in utero electroporation with the use of electrodes (disc size, 2.5 by 4

mm) positioned on each side of the uterus at E7.0 or E8.0. The fraction of mCherry-

expressing embryos at E9.0 and E10.0 was 19% and 40% among those electroporated at

E7.0 and E8.0, respectively (Figure S2a, b). The expression pattern of mCherry differed

among embryos, likely because of differences in the orientation of the electrodes

relative to the embryo during electroporation. Most embryos expressed mCherry in

somites (Figure S2a), whereas another manifested mCherry expression in the midbrain

and hindbrain (data not shown). No embryo examined to date was found to express

mCherry in the forebrain. These results showed that it was possible to perform gene

introduction by in utero electroporation in mouse embryos at E7.0 or E8.0 with our

method for plasmid injection, although further improvements will be necessary to allow

targeting of a specific tissue with this method.

Transcriptional profiling of NPCs in the mouse telencephalon during early

development

To investigate the molecular basis of the E-to-N transition, we first examined changes in

the transcriptional profile of NPCs around this developmental stage by RNA-

sequencing (RNA-seq) analysis. Given that the E-to-N transition takes place at ~E10 in

mice, we isolated NPCs as cells expressing the NPC marker CD133 (also known as

prominin-1) at a high level (CD133^{high} cells) from the entire telencephalon at E8.0 and

E9.0 and from the neocortex at E10.0, E11.0, and E12.0 by fluorescence-activated cell

sorting (FACS) (Figure 3a). Hierarchical clustering analysis of differentially expressed

genes (defined by maSigPro of the R package) revealed five clusters based on the

pattern of the expression changes (Figure 3b–f). The genes in cluster 1 showed a decline

in expression level, most prominently from E8.0 to E10.0 (Figure 3b), whereas those in

cluster 3 showed an increase in expression level, most prominently from E10.0 to E12.0

(Figure 3d). Gene ontology (GO) analysis showed that the genes in cluster 1 were

enriched in those related to tissue morphogenesis and embryo development (Figure 3b),

whereas the genes in cluster 3 were enriched in those related to nervous system

development and generation of neurons (Figure 3d). These findings confirmed the

proper isolation of NPCs at each stage. The genes in clusters 2 and 4 were potentially

associated with the E-to-N transition (~E10), given that their expression levels

decreased and increased, respectively, predominantly from E9.0 to E11.0 (Figure 3c and

e). The genes in cluster 2 were enriched in those related to ribosome biogenesis and

RNA processing (Figure 3c), suggesting that these processes might be regulated

differentially in the expansion versus neurogenic phase. Ribosome biogenesis has

indeed been found to be attenuated in NPCs at around this stage (Chau et al., 2018),

again providing support for the proper isolation of NPCs for this analysis. The

enrichment of genes related to RNA processing in cluster 2 is of interest, given the

pivotal role of such processing in the control of NPC differentiation (K.-J. Yoon et al.,

2017; Zahr et al., 2018; Zhang et al., 2016). The genes in cluster 4 were enriched in

those related to cell projection and cell morphogenesis (Figure 3e), possibly reflecting

the dynamic morphological changes of NPCs as they transition from NECs to RGCs.

The genes in cluster 5 may be particularly relevant to the E-to-N transition,

given that their expression peaks at E10.0 (Figure 3f). We therefore hypothesized that

this cluster includes genes that promote the E-to-N transition. The top five highly

expressed genes of cluster 5 at E10.0 are shown in Figure 3g. Among them, we focused

on *Hmga2*, given that Hmga proteins have been shown to confer neurogenic potential

on mouse neocortical NPCs at late developmental stages (Kishi et al., 2012). However,

the possible role of Hmga2 at early developmental stages—in particular, at the E-to-N

transition—has remained unknown.

Hmga2 promotes the onset of neurogenesis in the mouse neocortex

240
We first confirmed the temporal expression pattern of *Hmga2* in NPCs by reverse

transcription and quantitative polymerase chain reaction (RT-qPCR) analysis, which

again showed a peak at E10.0 (Figure 3h). To investigate the relevance of this increase

in the level of *Hmga2* expression in neocortical NPCs at this particular embryonic stage,

we knocked down Hmga2 with the use of a specific short hairpin RNA (shRNA) from

E8.0 with our novel method for early embryo manipulation. We introduced retroviruses

encoding GFP and shRNAs specific either for Hmga2 or for luciferase (control) into

NPCs at E8.0 and analyzed the embryos at E10.0 and E11.0 (Figure 4a). RT-qPCR and

immunohistofluorescence analyses confirmed the depletion of *Hmga2* mRNA and

Hmga2 protein in NPCs at these stages for embryos infected with the retrovirus for

Hmga2 shRNA at E8.0 (Figure 4b and c). We then examined the effect of Hmga2

knockdown on neurogenesis by immunostaining for Tbr2, a marker of intermediate

neuronal progenitors and immature neurons (Englund et al., 2005). Whereas Hmga2

knockdown from E8.0 did not affect the fraction of Tbr2-positive cells among GFP-

positive cells at E10.0, it significantly attenuated the increase in the fraction of such

cells apparent at E11.0 (Figure 4d and e). In contrast, Hmga2 knockdown from E8.0 did

not significantly affect the fraction of cells positive for Sox2, an NPC marker (Ellis et

al., 2004), among GFP-positive cells at E11.0 (Figure 4f and g). These results suggested

that Hmga2 promotes neurogenesis in neocortical NPCs at the onset of the neurogenic

phase.

There were at least two possible explanations for the attenuation of

neurogenesis by knockdown of Hmga2: Hmga2 might promote the E-to-N transition of

NPCs, in which case Hmga2 depletion would be expected to prolong the expansion

phase and delay the start of the neurogenic phase; or Hmga2 might promote neuronal

differentiation of NPCs during the neurogenic phase without affecting the E-to-N

transition of NPCs. To discriminate between these possibilities, we examined whether

Hmga2 knockdown might delay the developmental change in the transcriptional profile

of neocortical NPCs at the E-to-N transition. We therefore introduced retroviruses

encoding GFP and either control or Hmga2 shRNAs into embryos at E8.0 and isolated

GFP-expressing CD133[high] neocortical NPCs by FACS at E10.0 or E11.0 for RNA-seq

analysis, which we performed with the Quartz-seq method for analysis of a small

number of cells (Sasagawa et al., 2013) (Figure 5a). The expression of *Hmga2* as well

as that of *Igf2bp2*, a target gene of Hmga2 in NPCs (Fujii et al., 2013), were found to be greatly reduced in NPCs of embryos infected with the retrovirus for Hmga2 shRNA (Figure 5b), thus confirming the efficient knockdown of Hmga2 by this procedure. We then performed gene set enrichment analysis (GSEA) to compare the expression levels of signature genes of NPCs in the expansion phase between control and Hmga2-depleted NPCs at E10.0. We examined genes in cluster 2 as such signature genes, given the prominent decline in their expression level from E9.0 to E11.0 (Figure 3c), and we found that their expression level was significantly higher (enriched) in Hmga2-depleted NPCs than in control NPCs (Figure 5c). This finding suggested that Hmga2 knockdown shifts the transcriptional profile of NPCs closer to that apparent in the expansion phase, providing support for the notion that the observed increase in Hmga2 expression promotes the E-to-N transition of NPC fate in the neocortex. We previously showed that Hmga2 knockdown during the late neurogenic phase did not promote premature neuronal differentiation (Kishi et al., 2012), which further supports the notion that Hmga2 promotes the E-to-N transition of NPC fate rather than neuronal differentiation of NPCs per se.

Of interest, Hmga2 knockdown from E8.0 did not significantly increase the fraction of NPCs positive for phosphorylated histone H3 at E11.0 (data not shown), suggesting that Hmga2 might not overtly contribute to the change in cell division rate associated with the E-to-N transition. Together, our results thus implicated Hmga2 in changes in some (including the acquisition of neurogenic potential) but not all properties of NPCs related to this transition.

Hmga2 activates a subset of PcG target genes in early-stage neocortical NPCs

How does Hmga2 promote the acquisition of neurogenic potential in NPCs (without promoting neuronal differentiation per se)? In the developing mouse neocortex as well as in other contexts, PcG proteins have been shown to determine neurogenic potential through transient repression of neurogenic genes (Fasano et al., 2007; Hirabayashi et al., 2009; Mikkelsen et al., 2007; Mohn et al., 2008; Morimoto-Suzki et al., 2014; Pereira et al., 2010; Tsuboi et al., 2018). We then asked whether there is any relation between Hmga2 and PcG proteins in NPCs at the early stage of neocortical development. Given that PcG proteins catalyze histone modification including trimethylation at lysine-27 of histone H3 (H3K27me3), we performed chromatin immunoprecipitation (ChIP)-seq

analysis of Hmga2 and H3K27me3 in neocortical cells at E11.0. Clustering analysis

(and a heat map with Jaccard index) showed that loci associated with H3K27me3

showed an inverse correlation with those associated with Hmga2 in these cells (Figure

6a), consistent with a previous finding in mouse ES cells (Colombo et al., 2017). This

exclusive binding pattern may be due in large part to the preferential binding of Hmga2

and PcG proteins to AT-rich and GC-rich sequences, respectively (Colombo et al., 2017;

Ku et al., 2008; H. Li et al., 2017; Winter et al., 2011), but it was also possible that

these molecules interact functionally. We therefore examined whether knockdown of

Hmga2 in NPCs might affect the expression of PcG target genes, which we defined in

this study as genes that were derepressed (up-regulated) in response to ablation of

Ring1B, an essential component of PcG protein complexes. We first examined a set of

PcG target genes that we previously identified in a primary culture of NPCs prepared

from E18 mouse neocortex (Hirabayashi et al., 2009). GSEA of our Quartz-seq data

(Figure 5) showed that these PcG target genes were expressed at a significantly higher

level in control NPCs than in Hmga2-depleted NPCs at E11.0 (Figure 6b). We next

examined another set of PcG target genes that we identified in neocortical NPCs in vivo

by conditional deletion of *Ring1b* from E15 to E18 against a background of *Ring1a*

deletion (with the use of *Ring1a$^{-/-}$;Ring1b$^{flox/flox}$;Rosa-CreERT2$^{-/-\ or\ +/-}$* mouse embryos).

Again, GSEA revealed that these PcG target genes were expressed at a significantly

higher level in control NPCs than in Hmga2-knockdown NPCs at E11.0 (Figure 6c).

These results thus suggested that Hmga2 promotes the expression of (at least some)

PcG target genes in NPCs, with this effect possibly underlying the function of Hmga2 in

conferring neurogenic potential on these cells. Finally, we performed GO analysis of

core enriched genes whose expression was derepressed by Ring1B ablation and down-

regulated by Hmga2 knockdown (Figure 6b and 6c). The top hits of this analysis

included brain development and nervous system development (Figure 6d), providing

further support for the notion that Hmga2 promotes the expression of PcG target genes

that contribute to nervous system development.

Discussion

A directional transition of NPC fate from the expansion phase to the neurogenic phase takes place during mammalian brain development, but the cell-intrinsic programs that govern this transition have remained largely unclear. We have here investigated changes in the transcriptional profile of mouse neocortical NPCs, and we identified gene sets that are specifically up- or down-regulated around the time of this E-to-N transition. Among the highly expressed genes whose expression peaks around this transition, we focused on that for the chromatin-associated protein Hmga2 and examined the effects of its knockdown with the use of a new method we developed in this study for the genetic manipulation of early-stage NPCs before neural tube closure. We thus found that forced depletion of Hmga2 in neocortical NPCs from E8.0 hindered the increase in neurogenesis apparent at the onset of the neurogenic phase (from E10.0 to E11.0). Moreover, we found that this effect of Hmga2 depletion was likely due to a delay in the E-to-N transition of NPC fate rather than to inhibition of neuronal differentiation per se within the neurogenic phase, given the delayed change in the transcriptional profile of NPCs induced by the loss of Hmga2. Importantly, the genes whose expression was attenuated in Hmga2-depleted NPCs were enriched in PcG target genes (genes up-regulated by deletion of *Ring1b*) in these cells. On the basis of our results, we propose that the increase in Hmga2 expression apparent at the E-to-N transition facilitates the expression of a subset of PcG target genes and thereby promotes this transition of NPC fate—in particular, by promoting the acquisition of neurogenic potential.

Investigation of the molecular mechanism underlying the E-to-N transition of NPC fate in any detail requires manipulation of gene expression before neural tube closure. However, existing (convenient and simple) methods such as injection into the ventricles of viruses or of plasmids for in utero electroporation (Tabata and Nakajima, 2001) can be applied only after neural tube closure. The method for gene introduction developed in the present study has several advantages over other methods that can be used before neural tube closure, such as whole-embryo genomic manipulation (gene targeting or transgene introduction) and ultrasound-guided injection of viruses into the mouse nervous system. Whereas genomic manipulation requires a specific Cre-expressing mouse line or knowledge of specific distal elements to control expression of a gene of interest in a temporally and spatially regulated manner, our new method

allows expression of a gene of interest in NPCs as well as in other ectodermal or

endodermal cells before neural tube closure without such mouse lines or knowledge.

Our method is also less costly, less labor-intensive, and less time-consuming compared

with existing methods and is thus suitable for testing and screening a large number of

genes for their specific effects. The relatively simple and rapid nature of our procedure

also appears to be beneficial in terms of minimizing developmental abnormalities and

maximizing the survival rate of manipulated embryos compared with the ultrasound-

guided injection of viruses (Gaiano et al., 1999). Given that many recent studies based

on brain organoids have implicated various molecules as potential regulators of early

brain development, a simple and efficient method for gene introduction into early-stage

mouse NPCs would be useful for rapid testing of the functions of these molecules in the

in vivo context.

Our present results suggest that Hmga2 positively regulates the expression of

PcG target genes in neocortical NPCs. PcG proteins are known to repress a specific

subset of developmental genes in a context-dependent manner, but the mechanisms

underlying this context-dependent regulation remain obscure in many cases. Our

findings now implicate Hmga2 as a factor that, directly or indirectly, regulates PcG

protein function in a locus- and developmental stage–dependent manner. It remains

unclear, however, how Hmga2 may regulate PcG target genes. Regulation of certain

genes (such as *CcnA2*, *Igf2bp2*, and *Plag1*) through direct association of Hmga2 with

their promoters has been described (Cleynen et al., 2007; Z. Li et al., 2012; Sakai et al.,

2019; Tessari et al., 2003; Vignali and Marracci, 2020), but this mechanism of

regulation seems to be relatively minor given the global exclusion of Hmga2 from gene

promoters in mouse ES cells (Colombo et al., 2017). Although Hmga2 has been

proposed to compete with histone H1 for linker DNA and thereby to open chromatin

(Catez et al., 2004), direct evidence for such a mechanism is lacking. Further

biochemical and cellular studies will be necessary to characterize the precise role of

Hmga2 in the regulation of its target genes.

Given the proposed function of Hmga2 as a trigger of the E-to-N transition,

the timing of the increase in Hmga2 abundance may play an important role in

determining that of the E-to-N transition. We indeed found that the amount of *Hmga2*

mRNA peaks at around the time of this transition (at ~E10) in mouse neocortical NPCs.

Then, what upstream pathways (mechanisms) might regulate this increase of *Hmga2*

expression? It is possible that extrinsic factors implicated in the E-to-N transition in the

mouse neocortex, such as FGF10 and atRA, contribute to the regulation of *Hmga2*

expression. Of note, the promoter of the mouse *Hmga2* gene contains putative retinoic

acid–responsive elements that are able to bind retinoic acid receptors (data not shown).

Myc proteins as well as the Lin28–*let7* microRNA axis have also been shown to

promote the expression of *Hmga2* in other cell types (Chang et al., 2009; Copley et al.,

2013; Lee and Dutta, 2007; Mayr et al., 2007). Although we did not detect an increase

in the expression of Myc-family proteins around E9 to E11, we found that the level of

Lin28b mRNA increases in neocortical NPCs before that of *Hmga2* mRNA at around

this time of development (data not shown). It will therefore be of interest to examine the

relevance of Lin28 to the regulation of *Hmga2* expression in the context of the E-to-N

transition in the mouse neocortex.

The search for the "timer" of the E-to-N transition is especially relevant to our

understanding of the evolutionary enlargement of the brain, given that the duration of

the expansion phase (or the number of symmetric NPC divisions) is expected to

positively correlate with the size of the founder NPC pool. Indeed, the enlargement of

the neocortex in primates as compared with rodents appears to be related to lengthening

of the expansion phase from a few days to several weeks (Dehay et al., 2015; Kornack

and Rakic, 1998; Rakic, 2009; 1995) in addition to the emergence of additional

proliferative NPC populations (such as outer/basal RGCs) (Betizeau et al., 2013; Fietz

et al., 2010; Hansen et al., 2010) and the continuation of proliferative NPC divisions

even after the onset of neurogenesis (Otani et al., 2016). Even among primates, the cell-

autonomous timing of the E-to-N transition in cultured NPCs is delayed in humans

compared with nonhuman primates (Otani et al., 2016). Of interest, the level of *HMGA2*

mRNA also seems to be high in neocortical NPCs at around the onset of neurogenesis

(gestation week 7 to 8) during early human development (Nowakowski et al., 2017).

Moreover, *HMGA2* was found to be one of the genes that most markedly influence the

velocity stream associated with timing of NPC fate transition at the early stage of

human neocortical development (Eze et al., 2020), suggesting the possibility that

HMGA2 plays a conserved role in the onset of the neurogenic phase in humans. If this

is the case, the difference in the regulation of Hmga2 gene expression between mouse

and human may provide insight into the timing of this transition.

The size of the neocortex is strictly regulated during mammalian

development, with disorders that result from defects in such regulation in humans, such

as macrocephaly and microcephaly, being associated with pathological or psychiatric

conditions such as epilepsy, autism spectrum disorder, and schizophrenia (Sun and

Hevner, 2014). A single-cell transcriptomic analysis of human neocortical progenitors

indicated that genes that drive the early-stage transition of NPC fate are enriched in

those associated with neurodevelopmental or psychiatric disorders (Eze et al., 2020). It

will thus be of interest to investigate the impact of Hmga2 and its related molecules on

neocortical size and, ultimately, neocortical function.

Materials and Methods

Mouse maintenance

All animals were studied according to protocols approved by the Animal Care and Use Committee of The University of Tokyo. *Ring1a*$^{-/-}$;*Ring1b*$^{flox/flox}$ mice (Calés et al., 2008; Endoh et al., 2008) were crossed with *Rosa26-CreERT2* mice (Taconic), in which expression of *CreERT2* is driven by the endogenous *Rosa26* promoter. All mice were maintained in a temperature- and relative humidity–controlled (23° ± 3°C and 50 ± 15%, respectively) environment with a normal 12-h-light, 12-h-dark cycle. Two to six mice were housed per sterile cage (Innocage, Innovive; or Micro BARRIER Systems, Edstrom Japan) containing bedding chips (PALSOFT, Oriental Yeast; or PaperClean, SLC Japan), and they were provided with irradiated food (CE-2, CLEA Japan) and filtered water ad libitum. For in utero injection before neural tube closure, mice were allowed to mate for 2 h and the female was then checked for a vaginal plug. The time of plug confirmation was defined as E0.0. For induction of CreERT2 activity, pregnant mice were injected intraperitoneally with 150 μl of tamoxifen (Sigma) dissolved in sunflower oil (Nacalai) at a concentration of 10 mg/ml.

In utero injection and electroporation before neural tube closure

Jcl:ICR (CLEA Japan) or Slc:ICR (SLC Japan) mice were used for in utero injection experiments. Pregnant mice were injected intraperitoneally with a sodium pentobarbital–based anesthetic, and the uterine horn was exposed and pinched between a flexible fiber-optic cable and a finger (Matsui et al., 2011). Lighting from the fiber-optic cable allowed visualization of the internal uterine structure but did not allow the embryo to be distinguished from the amnion. A small volume (~0.5 μl) of CFSE (1 mM, Sigma), plasmid DNA (1 μg/μl), or retroviruses, each mixed with FastGreen (Chroma Gesellschaft), was injected with a sharpened capillary at an angle perpendicular to the center of the uterus. An injection depth of 1 to 2 mm below the uterine wall allowed the reproducible introduction of the capillary contents into amniotic fluid. For electroporation, the uterus was held between the tweezer-type electrodes (CUY652P2.5x4, Nepa Gene) and electroporated with a NEPA21 system (Nepa Gene) according to the following program: three poring pulses of 30 V with a 15-ms pulse width and 450-ms pulse interval and 10% decay (+ pulse orientation), and two transfer

pulses of 10 V with a 50-ms pulse width and 450-ms pulse interval and 10% decay (±

pulse orientation). After the injection and electroporation were completed, the uterine

horn was placed back into its original location. Embryos were imaged with a

fluorescence stereo microscope (2.0× lens, Nikon SMZ18).

Plasmid construction

The cDNA for mCherry was cloned by PCR, and its coding sequence was verified by

DNA sequencing. The cDNA was subcloned into pCAGEN (a kind gift from C. L.

Cepko) to generate the pCAGEN-mCherry plasmid.

Retrovirus production and concentration

Retroviruses were produced by transfection of PLAT-GP cells (a kind gift from T.

Kitamura) with the previously validated retroviral vectors pSIREN-siLuc (for control

shRNA, Addgene #122287) or pSIREN-siHmga2 (for Hmga2 shRNA, Addgene

#122290) (Kishi et al., 2012) and with a VSVG plasmid (a kind gift from H. Song).

Conditioned medium collected 72 h after the onset of transfection was cleared of debris

by centrifugation at $2500 \times g$ for 5 min, passed through a 0.45-μm filter, and centrifuged

at $250,000 \times g$ for 90 min at 4°C and then at $20,000 \times g$ for >12 h at 4°C. The resulting

virus pellet was suspended in phosphate-buffered saline (PBS). For in utero injection,

10 ml of conditioned medium from PLAT-GP cells in a 10-cm dish was concentrated

into 10 μl of PBS.

Flow cytometry and FACS

The telencephalon or neocortex of ICR embryos was dissected and subjected to

enzymatic digestion with a papain-based solution (Sumitomo Bakelite). For

quantification of the efficiency of gene introduction, the cells were directly subjected to

flow cytometric analysis with a FACS Aria instrument (Becton Dickinson). For

isolation of NPCs, the cells were stained with allophycocyanin-conjugated antibodies to

CD133 (1:400 dilution, BioLegend 141210) and sorted with a FACS Aria instrument

(Becton Dickinson). The threshold of CD133 intensity for selection of NPCs was set as

follows: E8.0, top 80%; E9.0, top 70%; E10.0, top 60%; E11.0, top 50%; and E12.0, top

50%.

Immunohistofluorescence analysis

Immunohistofluorescence staining was performed as previously described (Morimoto-

Suzki et al., 2014), with minor modifications including fixation for 3 h without

perfusion and the use of a blocking buffer comprising 0.1% Triton X-100 and 3%

bovine serum albumin. Primary antibodies and dilutions for immunostaining were as

follows: anti-Hmga2, 1:500 (Cell Signaling); anti-Sox2, 1:500 (Cell Signaling); anti-

Tbr2, 1:500 (chick, Millipore); anti–phospho-histone H3, 1:500 (Cell Signaling); and

anti-GFP, 1:2000 (Abcam) or 1:1000 (Nakalai). Images were acquired with a laser-

scanning confocal microscope (TSC-SP5, Leica) and ImageJ software (NIH). For

quantification, the average value of multiple sections of the telencephalon along the

rostrocaudal axis was determined for each embryo.

RT-qPCR analysis

Total RNA was isolated from cells with the use of RNAiso plus (Takara), and up to 0.5

μg of the RNA was subjected to RT with the use of ReverTra Ace qPCR RT Master Mix

with gDNA remover (Toyobo). The resulting cDNA was subjected to real-time PCR

analysis in a LightCycler 480 instrument (Roche) with either KAPA SYBR FAST for

LightCycler 480 (Kapa Biosystems) or Thunderbird SYBR qPCR Mix (Toyobo). The

amount of each target mRNA was normalized by that of *Actb* mRNA. Primer sequences

were described previously (Sakai et al., 2019).

RNA-seq analysis

Libraries for RNA-seq analysis were constructed from total RNA isolated as for RT-

qPCR analysis. An Illumina TruSeq RNA sample preparation kit was used for template

preparation, which was followed by deep sequencing on the Illumina HiSeq platform to

obtain 36-base single-end reads. Approximately 20 million sequences were obtained.

Sequenced reads of NPCs isolated from wild-type embryos or from Ring1A/B knockout

embryos were mapped to the reference mouse genome mm10 or mm9 with the use of

Hisat2 (Kim et al., 2019) or ELAND (Illumina), respectively. Only uniquely mapped

and "deduplicated" reads with no base mismatch were used. Expression information

was obtained as RPKM based on the RefSeq gene models and with the use of

featureCounts (Liao et al., 2014). Gene clustering analysis based on expression patterns

was performed with maSigPro (Conesa et al., 2006). GO analysis for functional

annotation was conducted with DAVID software (Huang et al., 2009).

Quartz-seq analysis

Both cDNA synthesis and amplification were performed with total RNA from 1000

cells as described previously (Sasagawa et al., 2013). The amplified cDNA was

prepared for sequencing with the use of a Nextera XT DNA Sample Prep Kit (Illumina)

and was subjected to deep sequencing analysis on the Illumina HiSeq platform to yield

36-base single-end reads. Approximately 2 million sequences were obtained from each

sample. Sequences were mapped to the reference mouse genome (mm10) with Hisat2

(Kim et al., 2019). Only uniquely mapped and "deduplicated" reads with no base

mismatch were used. Gene expression was quantitated as RPKM on the basis of RefSeq

gene models (mm10) with the use of featureCounts (Liao et al., 2014).

GSEA

Enrichment of signature genes was assessed with GSEA software (Subramanian et al.,

2005). Each gene set was determined as follows. Expansion-phase genes: genes

expressed in cluster 2 at ≥50 RPKM at E9.0. Ring1B target genes in cultured NPCs:

Ring1B knockout (KO)/control ≥ 1.5, expression in control ≥ 3.0 RPKM (Hirabayashi

et al., 2009). Ring1B target genes in NPCs in vivo: Ring1A/B KO/Ring1A KO ≥ 3.0,

expression in control (Ring1A KO) ≥ 1.0 RPKM.

ChIP-seq analysis

ChIP analysis was performed as previously described (Sakai et al., 2019) with

antibodies to Hmga2 (Cell Signaling) and to H3K27me3 (MBL). Two million cells

directly isolated from the neocortex of embryos at E11.0 were used for ChIP-seq of

Hmga2, and 100,000 NPCs isolated as CD133^{high} cells by FACS from the neocortex of

embryos at E11.0 were used for ChIP-seq of H3K27me3. An Illumina TruSeq DNA

sample preparation kit was used for template preparation, which was followed by deep

sequencing analysis on the Illumina HiSeq platform to obtain 36-base single-end reads.

Sequences were mapped to the reference mouse genome (mm10) with bowtie software

(Langmead et al., 2009). Only uniquely mapped reads without blacklist regions

determined by the ENCODE project (Amemiya et al., 2019) were used. Peaks for

Hmga2 and H3K27me3 were called with the use of F-seq software (Boyle et al., 2008).

Jaccard indices for sample comparisons were determined with bedtools (Quinlan and

Hall, 2010). R was used for clustering and heat map construction.

Statistical analysis

Data presented as means + SEM were compared with the two-tailed Student's paired or

Welch's *t* test or by one-way analysis of variance (ANOVA) followed by the

Bonferroni-Holm multiple-comparison test. A *p* value of <0.05 was considered

statistically significant.

Acknowledgments

We thank T. Shimogori (RIKEN) as well as M. Miura, Y. Yamaguchi, K. Hashimoto, and H. Miyazawa (The University of Tokyo) for technical advice regarding in utero injection before neural tube closure; I. Nikaido (RIKEN) for advice on Quartz-seq analysis; K. Imamura, T. Horiuchi, and S. Sugano (The University of Tokyo) for assistance with high-throughput sequencing analysis; Y. Koseki (RIKEN) for providing *Ring1* mutant mice; H. Song (Johns Hopkins University School of Medicine, Baltimore, MD) for the plasmid encoding the VSVG; T. Kitamura (The University of Tokyo) for PLAT-GP cells; C. L. Cepko (Harvard Medical School) for the pCAGEN plasmid; The One-Stop Sharing Facility Center for Future Drug Discoveries (The University of Tokyo) for FACS; R. Nagayoshi, and Y. Kakeya (The University of Tokyo) for technical assistance; and members of the Gotoh laboratory for discussion. This research was supported by AMED-CREST (JP20gm1310004 to Y.G.), by AMED-PRIME (JP20gm6110021 to Y.K.), and by MEXT/JSPS KAKENHI (JP16H06481 and JP16H06479 to Y.G.; JP19H05253, JP20H03179, JP20H05383, and JP16H06279 to Y.K.).

Author Contributions

N.K., Y.K., Y.H. and Y.G. designed the study and wrote the manuscript. N.K., Y.K., Y.M., Y.N., and Y.H. performed the experiments and analyzed the data. Y.S. performed high-throughput sequencing analysis. H.K. generated *Ring1a*$^{-/-}$;*Ring1b*$^{\text{flox/flox}}$ mice. Y.K., Y.H., and Y.G. supervised the study.

Declaration of Interests

The authors declare no competing interests.

References

Albert, M., Kalebic, N., Florio, M., Lakshmanaperumal, N., Haffner, C., Brandl, H.,

Henry, I., Huttner, W.B., 2017. Epigenome profiling and editing of neocortical

progenitor cells during development. The EMBO Journal 36, 2642–2658.

doi:10.15252/embj.201796764

Amemiya, H.M., Kundaje, A., Boyle, A.P., 2019. The ENCODE Blacklist:

Identification of Problematic Regions of the Genome. Sci Rep 9, 9354–5.

doi:10.1038/s41598-019-45839-z

Anthony, T.E., Mason, H.A., Gridley, T., Fishell, G., Heintz, N., 2005. Brain lipid-

binding protein is a direct target of Notch signaling in radial glial cells. Genes &

Development 19, 1028–1033. doi:10.1101/gad.1302105

Betizeau, M., Cortay, V., Patti, D., Pfister, S., Gautier, E., Bellemin-Ménard, A.,

Afanassieff, M., Huissoud, C., Douglas, R.J., Kennedy, H., Dehay, C., 2013.

Precursor diversity and complexity of lineage relationships in the outer

subventricular zone of the primate. Neuron 80, 442–457.

doi:10.1016/j.neuron.2013.09.032

Boyle, A.P., Guinney, J., Crawford, G.E., Furey, T.S., 2008. F-Seq: a feature density

estimator for high-throughput sequence tags. Bioinformatics 24, 2537–2538.

doi:10.1093/bioinformatics/btn480

Calés, C., Roman-Trufero, M., Pavón, L., Serrano, I., Melgar, T., Endoh, M., Pérez, C.,

Koseki, H., Vidal, M., 2008. Inactivation of the polycomb group protein Ring1B

unveils an antiproliferative role in hematopoietic cell expansion and cooperation

with tumorigenesis associated with Ink4a deletion. Molecular and Cellular Biology

28, 1018–1028. doi:10.1128/MCB.01136-07

Catez, F., Yang, H., Tracey, K.J., Reeves, R., Misteli, T., Bustin, M., 2004. Network of

dynamic interactions between histone H1 and high-mobility-group proteins in

chromatin. Molecular and Cellular Biology 24, 4321–4328.

doi:10.1128/mcb.24.10.4321-4328.2004

Chang, T.-C., Zeitels, L.R., Hwang, H.-W., Chivukula, R.R., Wentzel, E.A., Dews, M.,

Jung, J., Gao, P., Dang, C.V., Beer, M.A., Thomas-Tikhonenko, A., Mendell, J.T.,

2009. Lin-28B transactivation is necessary for Myc-mediated let-7 repression and

proliferation. Proceedings of the National Academy of Sciences 106, 3384–3389.

doi:10.1073/pnas.0808300106

Chau, K.F., Shannon, M.L., Fame, R.M., Fonseca, E., Mullan, H., Johnson, M.B.,

Sendamarai, A.K., Springel, M.W., Laurent, B., Lehtinen, M.K., 2018.

Downregulation of ribosome biogenesis during early forebrain development. Elife

7. doi:10.7554/eLife.36998

Cleynen, I., Brants, J.R., Peeters, K., Deckers, R., Debiec-Rychter, M., Sciot, R., Van

De Ven, W.J.M., Petit, M.M.R., 2007. HMGA2 regulates transcription of the Imp2

gene via an intronic regulatory element in cooperation with nuclear factor-kappaB.

Mol. Cancer Res. 5, 363–372. doi:10.1158/1541-7786.MCR-06-0331

Colombo, D.F., Burger, L., Baubec, T., Schübeler, D., 2017. Binding of high mobility

group A proteins to the mammalian genome occurs as a function of AT-content.

PLoS Genet 13, e1007102–36. doi:10.1371/journal.pgen.1007102

Conesa, A., Nueda, M.J., Ferrer, A., Talón, M., 2006. maSigPro: a method to identify

significantly differential expression profiles in time-course microarray experiments.

Bioinformatics 22, 1096–1102. doi:10.1093/bioinformatics/btl056

Copley, M.R., Babovic, S., Benz, C., Knapp, D.J.H.F., Beer, P.A., Kent, D.G., Wohrer,

S., Treloar, D.Q., Day, C., Rowe, K., Mader, H., Kuchenbauer, F., Humphries,

R.K., Eaves, C.J., 2013. The Lin28b-let-7-Hmga2 axis determines the higher self-

renewal potential of fetal haematopoietic stem cells. Nature Cell Biology.

doi:10.1038/ncb2783

Dehay, C., Kennedy, H., Kosik, K.S., 2015. The outer subventricular zone and primate-

specific cortical complexification. Neuron 85, 683–694.

doi:10.1016/j.neuron.2014.12.060

Eiraku, M., Watanabe, K., Matsuo-Takasaki, M., Kawada, M., Yonemura, S.,

Matsumura, M., Wataya, T., Nishiyama, A., Muguruma, K., Sasai, Y., 2008. Self-

organized formation of polarized cortical tissues from ESCs and its active

manipulation by extrinsic signals. Cell Stem Cell 3, 519–532.

doi:10.1016/j.stem.2008.09.002

Ellis, P., Fagan, B.M., Magness, S.T., Hutton, S., Taranova, O., Hayashi, S., McMahon,

A., Rao, M., Pevny, L., 2004. SOX2, a persistent marker for multipotential neural

stem cells derived from embryonic stem cells, the embryo or the adult. Dev.

Neurosci. 26, 148–165. doi:10.1159/000082134

Endoh, M., Endo, T.A., Endoh, T., Fujimura, Y.-I., Ohara, O., Toyoda, T., Otte, A.P.,

679 Okano, M., Brockdorff, N., Vidal, M., Koseki, H., 2008. Polycomb group proteins
680  Ring1A/B are functionally linked to the core transcriptional regulatory circuitry to
681  maintain ES cell identity. Development 135, 1513–1524. doi:10.1242/dev.014340

682 Englund, C., Fink, A., Lau, C., Pham, D., Daza, R.A.M., Bulfone, A., Kowalczyk, T.,
683  Hevner, R.F., 2005. Pax6, Tbr2, and Tbr1 are expressed sequentially by radial glia,
684  intermediate progenitor cells, and postmitotic neurons in developing neocortex.
685  Journal of Neuroscience 25, 247–251. doi:10.1523/JNEUROSCI.2899-04.2005

686 Eze, U.C., Bhaduri, A., Nowakowski, T.J., Kriegstein, A., 2020, n.d. Heterogeneity of
687  Human Neuroepithelial Cells and Early Radial Glia. biorxiv.
688  doi:10.1101/2020.03.06.981423

689 Fasano, C.A., Dimos, J.T., Ivanova, N.B., Lowry, N., Lemischka, I.R., Temple, S.,
690  2007. shRNA knockdown of Bmi-1 reveals a critical role for p21-Rb pathway in
691  NSC self-renewal during development. Cell Stem Cell 1, 87–99.
692  doi:10.1016/j.stem.2007.04.001

693 Fietz, S.A., Kelava, I., Vogt, J., Wilsch-Bräuninger, M., Stenzel, D., Fish, J.L., Corbeil,
694  D., Riehn, A., Distler, W., Nitsch, R., Huttner, W.B., 2010. OSVZ progenitors of
695  human and ferret neocortex are epithelial-like and expand by integrin signaling.
696  Nature Neuroscience 13, 690–699. doi:10.1038/nn.2553

697 Fujii, Y., Kishi, Y., Gotoh, Y., 2013. IMP2 regulates differentiation potentials of mouse
698  neocortical neural precursor cells. Genes to Cells 18, 79–89. doi:10.1111/gtc.12024

699 Gaiano, N., Kohtz, J.D., Turnbull, D.H., Fishell, G., 1999. A method for rapid gain-of-
700  function studies in the mouse embryonic nervous system. Nature Neuroscience 2,
701  812–819. doi:10.1038/12186

702 Gaiano, N., Nye, J.S., Fishell, G., 2000. Radial glial identity is promoted by Notch1
703  signaling in the murine forebrain. Neuron 26, 395–404. doi:10.1016/s0896-
704  6273(00)81172-1

705 Gao, P., Postiglione, M.P., Krieger, T.G., Hernandez, L., Wang, C., Han, Z., Streicher,
706  C., Papusheva, E., Insolera, R., Chugh, K., Kodish, O., Huang, K., Simons, B.D.,
707  Luo, L., Hippenmeyer, S., Shi, S.-H., 2014. Deterministic progenitor behavior and
708  unitary production of neurons in the neocortex. Cell 159, 775–788.
709  doi:10.1016/j.cell.2014.10.027

710 Gaspard, N., Bouschet, T., Hourez, R., Dimidschstein, J., Naeije, G., van den Ameele,
711  J., Espuny-Camacho, I., Herpoel, A., Passante, L., Schiffmann, S.N., Gaillard, A.,

Vanderhaeghen, P., 2008. An intrinsic mechanism of corticogenesis from

embryonic stem cells. Nature 455, 351–357. doi:10.1038/nature07287

GOtz, M., Huttner, W.B., 2005. The cell biology of neurogenesis. Nat Rev Mol Cell

Biol 6, 777–788. doi:10.1038/nrm1739

Hansen, D.V., Lui, J.H., Parker, P.R.L., Kriegstein, A.R., 2010. Neurogenic radial glia

in the outer subventricular zone of human neocortex. Nature 464, 554–561.

doi:10.1038/nature08845

Hartfuss, E., Galli, R., Heins, N., Götz, M., 2001. Characterization of CNS precursor

subtypes and radial glia. Developmental Biology 229, 15–30.

doi:10.1006/dbio.2000.9962

Hatakeyama, J., Bessho, Y., Katoh, K., Ookawara, S., Fujioka, M., Guillemot, F.,

Kageyama, R., 2004. Hes genes regulate size, shape and histogenesis of the nervous

system by control of the timing of neural stem cell differentiation. Development

131, 5539–5550. doi:10.1242/dev.01436

Hirabayashi, Y., Gotoh, Y., 2010. Epigenetic control of neural precursor cell fate during

development. Nat Rev Neurosci 11, 377–388. doi:10.1038/nrn2810

Hirabayashi, Y., Suzki, N., Tsuboi, M., Endo, T.A., Toyoda, T., Shinga, J., Koseki, H.,

Vidal, M., Gotoh, Y., 2009. Polycomb limits the neurogenic competence of neural

precursor cells to promote astrogenic fate transition. Neuron 63, 600–613.

doi:10.1016/j.neuron.2009.08.021

Huang, D.W., Sherman, B.T., Lempicki, R.A., 2009. Bioinformatics enrichment tools:

paths toward the comprehensive functional analysis of large gene lists. Nucleic

Acids Research 37, 1–13. doi:10.1093/nar/gkn923

Kim, D., Paggi, J.M., Park, C., Bennett, C., Salzberg, S.L., 2019. Graph-based genome

alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol 37,

907–915. doi:10.1038/s41587-019-0201-4

Kishi, Y., Fujii, Y., Hirabayashi, Y., Gotoh, Y., 2012. HMGA regulates the global

chromatin state and neurogenic potential in neocortical precursor cells. Nature

Neuroscience 15, 1127–1133. doi:10.1038/nn.3165

Kishi, Y., Gotoh, Y., 2018. Regulation of Chromatin Structure During Neural

Development. Front Neurosci 12, 874. doi:10.3389/fnins.2018.00874

Kornack, D.R., Rakic, P., 1998. Changes in cell-cycle kinetics during the development

and evolution of primate neocortex. Proc. Natl. Acad. Sci. U.S.A. 95, 1242–1246.

doi:10.1073/pnas.95.3.1242

Kriegstein, A., Alvarez-Buylla, A., 2009. The Glial Nature of Embryonic and Adult

Neural Stem Cells. Annu. Rev. Neurosci. 32, 149–184.

doi:10.1146/annurev.neuro.051508.135600

Ku, M., Koche, R.P., Rheinbay, E., Mendenhall, E.M., Endoh, M., Mikkelsen, T.S.,

Presser, A., Nusbaum, C., Xie, X., Chi, A.S., Adli, M., Kasif, S., Ptaszek, L.M.,

Cowan, C.A., Lander, E.S., Koseki, H., Bernstein, B.E., 2008. Genomewide

analysis of PRC1 and PRC2 occupancy identifies two classes of bivalent domains.

PLoS Genet 4, e1000242. doi:10.1371/journal.pgen.1000242

Lange, C., Turrero Garcia, M., Decimo, I., Bifari, F., Eelen, G., Quaegebeur, A., Boon,

R., Zhao, H., Boeckx, B., Chang, J., Wu, C., le Noble, F., Lambrechts, D.,

Dewerchin, M., Kuo, C.J., Huttner, W.B., Carmeliet, P., 2016. Relief of hypoxia by

angiogenesis promotes neural stem cell differentiation by targeting glycolysis. The

EMBO Journal 35, 924–941. doi:10.15252/embj.201592372

Langmead, B., Trapnell, C., Pop, M., Salzberg, S.L., 2009. Ultrafast and memory-

efficient alignment of short DNA sequences to the human genome. Genome

Biology 10, R25–10. doi:10.1186/gb-2009-10-3-r25

Lee, Y.S., Dutta, A., 2007. The tumor suppressor microRNA let-7 represses the

HMGA2 oncogene. Genes & Development 21, 1025–1030.

doi:10.1101/gad.1540407

Li, H., Liefke, R., Jiang, J., Kurland, J.V., Tian, W., Deng, P., Zhang, W., He, Q., Patel,

D.J., Bulyk, M.L., Shi, Y., Wang, Z., 2017. Polycomb-like proteins link the PRC2

complex to CpG islands. Nature 549, 287–291. doi:10.1038/nature23881

Li, Z., Gilbert, J.A., Zhang, Y., Zhang, M., Qiu, Q., Ramanujan, K., Shavlakadze, T.,

Eash, J.K., Scaramozza, A., Goddeeris, M.M., Kirsch, D.G., Campbell, K.P., Brack,

A.S., Glass, D.J., 2012. An HMGA2-IGF2BP2 axis regulates myoblast proliferation

and myogenesis. Developmental Cell 23, 1176–1188.

doi:10.1016/j.devcel.2012.10.019

Liao, Y., Smyth, G.K., Shi, W., 2014. featureCounts: an efficient general purpose

program for assigning sequence reads to genomic features. Bioinformatics 30, 923–

930. doi:10.1093/bioinformatics/btt656

Matsui, A., Yoshida, A.C., Kubota, M., Ogawa, M., Shimogori, T., 2011. Mouse in

utero electroporation: controlled spatiotemporal gene transfection. J Vis Exp.

doi:10.3791/3024

Mayr, C., Hemann, M.T., Bartel, D.P., 2007. Disrupting the pairing between let-7 and

Hmga2 enhances oncogenic transformation. Science 315, 1576–1579.

doi:10.1126/science.1137999

McLone, D.G., Bondareff, W., 1975. Developmental morphology of the subarachnoid

space and contiguous structures in the mouse. Am. J. Anat. 142, 273–293.

doi:10.1002/aja.1001420302

Mikkelsen, T.S., Ku, M., Jaffe, D.B., Issac, B., Lieberman, E., Giannoukos, G.,

Alvarez, P., Brockman, W., Kim, T.-K., Koche, R.P., Lee, W., Mendenhall, E.,

O'Donovan, A., Presser, A., Russ, C., Xie, X., Meissner, A., Wernig, M., Jaenisch,

R., Nusbaum, C., Lander, E.S., Bernstein, B.E., 2007. Genome-wide maps of

chromatin state in pluripotent and lineage-committed cells. Nature 448, 553–560.

doi:10.1038/nature06008

Miyata, T., Kawaguchi, A., Saito, K., Kuramochi, H., Ogawa, M., 2002. Visualization

of cell cycling by an improvement in slice culture methods. J. Neurosci. Res. 69,

861–868. doi:10.1002/jnr.10335

Mohn, F., Weber, M., Rebhan, M., Roloff, T.C., Richter, J., Stadler, M.B., Bibel, M.,

Schübeler, D., 2008. Lineage-specific polycomb targets and de novo DNA

methylation define restriction and potential of neuronal progenitors. Molecular Cell

30, 755–766. doi:10.1016/j.molcel.2008.05.007

Morimoto-Suzki, N., Hirabayashi, Y., Tyssowski, K., Shinga, J., Vidal, M., Koseki, H.,

Gotoh, Y., 2014. The polycomb component Ring1B regulates the timed termination

of subcerebral projection neuron production during mouse neocortical development.

Development 141, 4343–4353. doi:10.1242/dev.112276

Mukhtar, T., Taylor, V., 2018. Untangling Cortical Complexity During Development. J

Exp Neurosci 12, 1179069518759332. doi:10.1177/1179069518759332

Nakagawa, T., Wada, Y., Katada, S., Kishi, Y., 2020. Epigenetic regulation for

acquiring glial identity by neural stem cells during cortical development. Glia 14,

67–14. doi:10.1002/glia.23818

Nguyen, H., Kerimoglu, C., Pirouz, M., Pham, L., Kiszka, K.A., Sokpor, G., Sakib,

M.S., Rosenbusch, J., Teichmann, U., Seong, R.H., Stoykova, A., Fischer, A.,

Staiger, J.F., Tuoc, T., 2018. Epigenetic Regulation by BAF Complexes Limits

Neural Stem Cell Proliferation by Suppressing Wnt Signaling in Late Embryonic

Development. Stem Cell Reports. doi:10.1016/j.stemcr.2018.04.014

Nishino, J., Kim, I., Chada, K., Morrison, S.J., 2008. Hmga2 promotes neural stem cell

self-renewal in young but not old mice by reducing p16Ink4a and p19Arf

Expression. Cell 135, 227–239. doi:10.1016/j.cell.2008.09.017

Noctor, S.C., Flint, A.C., Weissman, T.A., Dammerman, R.S., Kriegstein, A.R., 2001.

Neurons derived from radial glial cells establish radial units in neocortex. Nature

409, 714–720. doi:10.1038/35055553

Nowakowski, T.J., Bhaduri, A., Pollen, A.A., Alvarado, B., Mostajo-Radji, M.A., Di

Lullo, E., Haeussler, M., Sandoval-Espinosa, C., Liu, S.J., Velmeshev, D.,

Ounadjela, J.R., Shuga, J., Wang, X., Lim, D.A., West, J.A., Leyrat, A.A., Kent,

W.J., Kriegstein, A.R., 2017. Spatiotemporal gene expression trajectories reveal

developmental hierarchies of the human cortex. Science 358, 1318–1323.

doi:10.1126/science.aap8809

Otani, T., Marchetto, M.C., Gage, F.H., Simons, B.D., Livesey, F.J., 2016. 2D and 3D

Stem Cell Models of Primate Cortical Development Identify Species-Specific

Differences in Progenitor Behavior Contributing to Brain Size. Cell Stem Cell 18,

467–480. doi:10.1016/j.stem.2016.03.003

Pereira, J.D., Sansom, S.N., Smith, J., Dobenecker, M.-W., Tarakhovsky, A., Livesey,

F.J., 2010. Ezh2, the histone methyltransferase of PRC2, regulates the balance

between self-renewal and differentiation in the cerebral cortex. Proceedings of the

National Academy of Sciences 107, 15957–15962. doi:10.1073/pnas.1002530107

Qian, X., Goderie, S.K., Shen, Q., Stern, J.H., Temple, S., 1998. Intrinsic programs of

patterned cell lineages in isolated vertebrate CNS ventricular zone cells.

Development 125, 3143–3152.

Quah, B.J.C., Parish, C.R., 2010. The use of carboxyfluorescein diacetate succinimidyl

ester (CFSE) to monitor lymphocyte proliferation. J Vis Exp. doi:10.3791/2259

Quinlan, A.R., Hall, I.M., 2010. BEDTools: a flexible suite of utilities for comparing

genomic features. Bioinformatics 26, 841–842. doi:10.1093/bioinformatics/btq033

Rakic, P., 2009. Evolution of the neocortex: a perspective from developmental biology.

Nat Rev Neurosci 10, 724–735. doi:10.1038/nrn2719

Rakic, P., 1995. A small step for the cell, a giant leap for mankind: a hypothesis of

neocortical expansion during evolution. Trends in Neurosciences 18, 383–388.

doi:10.1016/0166-2236(95)93934-p

Rakic, P., 1972. Mode of cell migration to the superficial layers of fetal monkey

neocortex. J. Comp. Neurol. 145, 61–83. doi:10.1002/cne.901450105

Sahara, S., O'Leary, D.D.M., 2009. Fgf10 regulates transition period of cortical stem

cell differentiation to radial glia controlling generation of neurons and basal

progenitors. Neuron 63, 48–62. doi:10.1016/j.neuron.2009.06.006

Sakai, H., Fujii, Y., Kuwayama, N., Kawaji, K., Gotoh, Y., Kishi, Y., 2019. Plag1

regulates neuronal gene expression and neuronal differentiation of neocortical

neural progenitor cells. Genes to Cells 24, 650–666. doi:10.1111/gtc.12718

Sanosaka, T., Namihira, M., Asano, H., KOHYAMA, J., Aisaki, K., Igarashi, K.,

Kanno, J., Nakashima, K., 2008. Identification of genes that restrict astrocyte

differentiation of midgestational neural precursor cells. Neuroscience 155, 780–788.

doi:10.1016/j.neuroscience.2008.06.039

Sasagawa, Y., Nikaido, I., Hayashi, T., Danno, H., Uno, K.D., Imai, T., Ueda, H.R.,

2013. Quartz-Seq: a highly reproducible and sensitive single-cell RNA sequencing

method, reveals non-genetic gene-expression heterogeneity. Genome Biology 14,

R31. doi:10.1186/gb-2013-14-4-r31

Shen, Q., Wang, Y., Dimos, J.T., Fasano, C.A., Phoenix, T.N., Lemischka, I.R.,

Ivanova, N.B., Stifani, S., Morrisey, E.E., Temple, S., 2006. The timing of cortical

neurogenesis is encoded within lineages of individual progenitor cells. Nature

Neuroscience 9, 743–751. doi:10.1038/nn1694

Shi, Y., Kirwan, P., Smith, J., Robinson, H.P.C., Livesey, F.J., 2012. Human cerebral

cortex development from pluripotent stem cells to functional excitatory synapses.

Nature Neuroscience 15, 477–86– S1. doi:10.1038/nn.3041

Siegenthaler, J.A., Ashique, A.M., Zarbalis, K., Patterson, K.P., Hecht, J.H., Kane,

M.A., Folias, A.E., Choe, Y., May, S.R., Kume, T., Napoli, J.L., Peterson, A.S.,

Pleasure, S.J., 2009. Retinoic acid from the meninges regulates cortical neuron

generation. Cell 139, 597–609. doi:10.1016/j.cell.2009.10.004

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,

Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., Mesirov, J.P., 2005. Gene

set enrichment analysis: A knowledge-based approach for interpreting genome-

wide expression profiles. Proc. Natl. Acad. Sci. U.S.A. 102, 15545–15550.

doi:10.1073/pnas.0506580102

Sun, T., Hevner, R.F., 2014. Growth and folding of the mammalian cerebral cortex:

from molecules to malformations. Nat Rev Neurosci 15, 217–232.

doi:10.1038/nrn3707

Tabata, H., Nakajima, K., 2001. Efficient in utero gene transfer system to the

developing mouse brain using electroporation: visualization of neuronal migration

in the developing cortex. Neuroscience 103, 865–872. doi:10.1016/s0306-

4522(01)00016-1

Telley, L., Govindan, S., Prados, J., Stévant, I., Nef, S., Dermitzakis, E., Dayer, A.,

Jabaudon, D., 2016. Sequential transcriptional waves direct the differentiation of

newborn neurons in the mouse neocortex. Science 351, 1443–1446.

doi:10.1126/science.aad8361

Tessari, M.A., Gostissa, M., Altamura, S., Sgarra, R., Rustighi, A., Salvagno, C.,

Caretti, G., Imbriano, C., Mantovani, R., Del Sal, G., Giancotti, V., Manfioletti, G.,

2003. Transcriptional activation of the cyclin A gene by the architectural

transcription factor HMGA2. Molecular and Cellular Biology 23, 9104–9116.

Tsuboi, M., Hirabayashi, Y., Gotoh, Y., 2019. Diverse gene regulatory mechanisms

mediated by Polycomb group proteins during neural development. Current Opinion

in Neurobiology 59, 164–173. doi:10.1016/j.conb.2019.07.003

Tsuboi, M., Kishi, Y., Yokozeki, W., Koseki, H., Hirabayashi, Y., Gotoh, Y., 2018.

Ubiquitination-Independent Repression of PRC1 Targets during Neuronal Fate

Restriction in the Developing Mouse Neocortex. Developmental Cell 47, 758–

772.e5. doi:10.1016/j.devcel.2018.11.018

Vignali, R., Marracci, S., 2020. HMGA Genes and Proteins in Development and

Evolution. IJMS 21, 654–39. doi:10.3390/ijms21020654

Winter, N., Nimzyk, R., Bösche, C., Meyer, A., Bullerdiek, J., 2011. Chromatin

immunoprecipitation to analyze DNA binding sites of HMGA2. PLoS ONE 6,

e18837. doi:10.1371/journal.pone.0018837

Yoon, K., Nery, S., Rutlin, M.L., Radtke, F., Fishell, G., Gaiano, N., 2004. Fibroblast

growth factor receptor signaling promotes radial glial identity and interacts with

Notch1 signaling in telencephalic progenitors. Journal of Neuroscience 24, 9497–

9506. doi:10.1523/JNEUROSCI.0993-04.2004

Yoon, K.-J., Ringeling, F.R., Vissers, C., Jacob, F., Pokrass, M., Jimenez-Cyrus, D., Su,

Y., Kim, N.-S., Zhu, Y., Zheng, L., Kim, S., Wang, X., Doré, L.C., Jin, P., Regot,

S., Zhuang, X., Canzar, S., He, C., Ming, G.-L., Song, H., 2017. Temporal Control

of Mammalian Cortical Neurogenesis by m(6)A Methylation. Cell 171, 877–

889.e17. doi:10.1016/j.cell.2017.09.003

Zahr, S.K., Yang, G., Kazan, H., Borrett, M.J., Yuzwa, S.A., Voronova, A., Kaplan,

D.R., Miller, F.D., 2018. A Translational Repression Complex in Developing

Mammalian Neural Stem Cells that Regulates Neuronal Specification. Neuron 97,

520–537.e6. doi:10.1016/j.neuron.2017.12.045

Zhang, X., Chen, M.H., Wu, X., Kodani, A., Fan, J., Doan, R., Ozawa, M., Ma, J.,

Yoshida, N., Reiter, J.F., Black, D.L., Kharchenko, P.V., Sharp, P.A., Walsh, C.A.,

2016. Cell-Type-Specific Alternative Splicing Governs Cell Fate in the Developing

Cerebral Cortex. Cell 166, 1147–1162.e15. doi:10.1016/j.cell.2016.07.025

 Figure Legends

 Figure 1. In utero injection of mouse embryos before neural tube closure. (**a** and **b**)

 The uterine horn of an anesthetized pregnant mouse was exposed at E7.0 or E8.0. (**c** and

 d) The uterus was pinched between an optical fiber and finger (**c**) for injection via a

 sharpened glass capillary of a solution colored with a dye (FastGreen) and containing

 either CFSE, a retrovirus, or a plasmid (**d**).

 Figure 2. In utero gene introduction by retroviral infection in embryos before

 neural tube closure. (**a**) Mouse embryos in the same uterus were injected or not with a

 retrovirus encoding GFP together with an shRNA targeted to luciferase mRNA at E8.0.

 The embryos were dissected out from the uterus at E11.0, fixed, and examined for GFP

 fluorescence with a fluorescence stereo microscope. The posterior half of the control

 embryo was excised for the purpose of identification. Scale bars, 1 mm. (**b**) An embryo

 at E7.0 was injected as in (**a**) and examined for GFP fluorescence at E11.0. Scale bar, 1

 mm. (**c**) Survival rate and labeling rate for embryos injected as in (**a**) and (**b**). (**d**) Flow

 cytometric analysis of the efficiency of gene introduction. Neocortical cells dissociated

 from E12.0 embryos that had been injected with a retrovirus for GFP at E8.0 were

 subjected to flow cytometric analysis of GFP expression. (**e**) A sagittal section of an

 E10.0 embryo injected at E8.0 as in (**a**) was subjected to immunohistofluorescence

 staining with antibodies to GFP. Nuclei were counterstained with Hoechst 33342.

 Representative infected tissues are indicated. Scale bar, 1 mm. (**f**, **g**) Higher

 magnification images of the bronchus (**f**) and spinal cord (**g**) in the section shown in (**e**).

 Scale bars, 100 μm.

 Figure 3. Hierarchical clustering of differentially expressed genes in NPCs at early

 developmental stages. (**a**) CD133[high] cells were isolated as NPCs from the

 telencephalon at E8.0 and E9.0 and from the neocortex at E10.0, E11.0, and E12.0 by

 FACS and were subjected to RNA-seq analysis. (**b**–**f**) Differentially expressed genes in

 NPCs at the various developmental stages isolated as in (**a**) were categorized into five

 clusters (clusters 1–5) on the basis of the pattern of expression changes with the use of

 hierarchical clustering analysis. The expression levels of the genes in each gene cluster

 are shown as box plots, with the boxes representing the median and upper and lower

quartiles and the whiskers indicating the range (left). Enriched GO terms and their *p*

values determined by functional annotation of each gene cluster with DAVID software

are also shown (right). RPKM, reads per kilobase of mRNA per million total reads. (**g**)

The five genes expressed at the highest levels in cluster 5 at E10.0. (**h**) NPCs isolated as

in (**a**) from embryos at E9.0, E10.0, E11.0, and E12.0 were subjected to RT-qPCR

analysis of *Hmga2* mRNA. Data were normalized by *Actb* mRNA abundance and are

means + SEM from five independent experiments. *$p < 0.05$, **$p < 0.01$ (one-way

ANOVA followed by the Bonferroni-Holm multiple-comparison test).

Figure 4. Hmga2 promotes the onset of neurogenesis in the mouse neocortex. (**a**)

Mouse embryos at E8.0 were subjected to in utero infection with retroviruses encoding

GFP and either control or Hmga2 shRNAs, and the brain was isolated from the embryos

at E10.0 or E11.0 for isolation of CD133[high], GFP[+] NPCs by FACS. (**b**) NPCs isolated

from control and Hmga2–knockdown (KD) embryos as in (**a**) were subjected to RT-

qPCR analysis of *Hmga2* mRNA. Data are expressed relative to the corresponding

control value and are means + SEM from five independent experiments. *$p < 0.05$, **p

< 0.01 versus corresponding control value (two-tailed Student's paired *t* test). (**c–g**) The

brain of embryos treated as in (**a**) was subjected to immunohistofluorescence analysis

with antibodies to GFP and to Hmga2 at E10.0 (**c**), with those to GFP and toTbr2 at

E11.0 (**d**), and with those to GFP and to Sox2 at E11.0 (**f**). Nuclei were counterstained

with Hoechst 33342. Scale bars, 20 μm (**c**) or 50 μm (**d** and **f**). Arrowheads indicate

GFP[+] cells (**c**). The percentages of Tbr2[+] cells at E10.0 or E11.0 (**e**) or of Sox2[+] cells at

E11.0 (**g**) among GFP[+] cells were also determined. Data are means + SEM from six

(Tbr2) or three (Sox2) independent experiments (total of 12 control and nine Hmga2-

knockdown embryos for Tbr2, and eight control and seven Hmga2-knockdown embryos

for Sox2). *$p < 0.05$ versus corresponding control value (two-tailed Welch's *t* test).

Figure 5. Hmga2 promotes the change in transcriptional profile of NPCs at the E-

to-N transition. (**a**) Embryos were subjected to in utero infection with retroviruses

encoding GFP and either control or Hmga2 shRNAs at E8.0, and the brain was isolated

at E10.0 or E11.0 for isolation of GFP[+], CD133[high] NPCs by FACS and subsequent RNA

(Quartz)-seq analysis. Three or four independent experiments were performed with cells

isolated at E10.0 or E11.0, respectively. (**b**) Expression levels (RPKM) of *Hmga2* and

Igf2bp2 in Hmga2–knockdown (KD) and control NPCs determined as in (**a**). Data are

means + SEM and are expressed relative to the corresponding control value. *$p < 0.05$,

***$p < 0.01$ versus corresponding control value (two-tailed Student's paired *t* test). (**c**)

NPCs isolated at E10.0 as in (**a**) were subjected to GSEA for signature genes of the

expansion phase (those in cluster 2 with an RPKM of ≥50 at E9.0) (Figure 3c). The

family-wise error rate (FWER) *p* value and false discovery rate (FDR) *q* value are

shown.

Figure 6. Hmga2 promotes the expression of PcG target genes in NPCs. (**a**) Cells

isolated from the mouse neocortex at E11.0 were subjected to ChIP-seq for Hmga2

(four independent experiments). CD133^high cells sorted by FACS were used for the

H3K27me3 analysis (three independent experiments). The genome-wide correlation

heat map based on the Jaccard index for determined peaks in the samples is shown. (**b**)

The Quartz-seq data for NPCs at E11.0 in Figure 5 were subjected to GSEA of PcG

target genes previously identified by Ring1B knockout (KO) in an NPC culture

(Hirabayashi et al., 2009). The FWER *p* value and FDR *q* value are shown. (**c**) The

Quartz-seq data for NPCs at E11.0 in Figure 5 were subjected to GSEA of PcG target

genes that we identified by RNA-seq analysis of NPCs isolated by FACS at E18 from

the neocortex of *Ring1a*^{−/−};*Ring1b*^{flox/flox};*Rosa-CreERT2*^{−/− or +/−} mouse embryos that had

been treated with tamoxifen at E15 (see Materials and Methods). The FWER *p* value

and FDR *q* value are shown. (**d**) GO terms and their *p* values determined by functional

annotation of core enrichment genes in the GSEA analyses shown in (**b**) and (**c**) with the

use of DAVID software.

1011

 Supplemental Figure Legends

 Figure S1. In utero injection of CFSE into mouse embryos before neural tube

 closure. (**a**) Embryos at E7.0 (left) or E8.0 (right) were injected with CFSE, dissected

 out from the uterus at E9.0 or E11.0, respectively, and fixed for visualization of CFSE

 fluorescence with a fluorescence stereo microscope. Scale bars, 0.5 mm (left) or 1 mm

 (right). (**b**) Survival rate and labeling rate for in utero introduction of CFSE into mouse

 embryos as in (**a**).

 Figure S2. In utero electroporation of embryos before neural tube closure. (**a**)

 Embryos at E7.0 (left) or E8.0 (right) were injected with a plasmid for mCherry and

 subjected to in utero electroporation. They were dissected out from the uterus at E9.0 or

 E10.0, respectively, fixed, and examined for mCherry fluorescence with a fluorescence

 stereo microscope. Scale bars, 0.5 mm (left) or 1 mm (right). (**b**) Survival rate and

 labeling rate for in utero electroporation of embryos as in (**a**).

Fig. 1

a

Anesthetized
pregnant mouse
(E7.0 or E8.0)

b

Exposed uterine horn

c

A uterus pinched between an
optical fiber and finger

d

Sharpened glass capillary

1-2 mm

Center of
the uterus

Uterine horn

Injected embryos

Injection of CFSE, retroviruses, or plasmids

Fig. 2

a
E8.0 injection → E11.0

Control (no injection)　　GFP-expressing virus

GFP

b
E7.0 injection → E11.0

GFP

c

	Injected embryos	Survival rate (% of injected)	GFP-positive (% of injected)
E7.0 injection → E9.0	28	27 (96%)	2 (7%)
E8.0 injection → E10.0	19	17 (89%)	14 (74%)

d
E8.0 injection → E12.0

Control (no injection)

Cell number

0%

GFP

GFP-expressing virus

Cell number

70%

GFP

e
E8.0 injection → E10.0

GFP/Hoechst

Fourth ventricle

Epidermis

Spinal cord

g

f

Optic stalk

Lateral ventricle

Endodermal tissues

f
Bronchus

g
Spinal cord

Fig. 3

a
E8.0 whole brain
E9.0 whole brain
E10.0 neocortex → FACS (CD133^high),
E11.0 neocortex   RNA-seq
E12.0 neocortex

b Cluster 1 (1175 genes)
Tissue morphogenesis
Embryo development
Anatomical structure formation involved in morphogenesis
Circulatory system development
Cardiovascular system development

c Cluster 2 (1552 genes)
Ribosome biogenesis
ncRNA metabolic process
Ribonucleoprotein complex biogenesis
ncRNA processing
rRNA processing

d Cluster 3 (1736 genes)
Nervous system development
Generation of neurons
Neurogenesis
Regulation of nervous system development
Neuron differentiation

e Cluster 4 (1597 genes)
Cell projection morphogenesis
Cell part morphogenesis
Cell projection organization
Cell morphogenesis
Regulation of cell morphogenesis

f Cluster 5 (284 genes)
Cell surface receptor signaling pathway
Chondrocyte differentiation
Connective tissue development
Negative regulation of response to stimulus
Regulation of cell communication

g

Gene symbol	Expression level (RPKM)
Atp5a1	339
Hmga2	279
Lrrn1	217
Calm3	129
Ccnd1	127

h *Hmga2/Actb* mRNA ratio (arbitrary units)
E9.0, E10.0, E11.0, E12.0

Fig. 4

Fig. 5

a

b

Control
Hmga2 KD

c

Fig. 6

a

E11.0 WT neocortex $\longrightarrow$ Hmga2 ChIP-seq

E11.0 WT neocortex $\longrightarrow$ FACS (CD133^{high}), H3K27me3 ChIP-seq

Hmga2, replicate 1
Hmga2, replicate 2
Hmga2, replicate 3
Hmga2, replicate 4
H3K27me3, replicate 1
H3K27me3, replicate 2
H3K27me3, replicate 3

1
0.5
0

b Derepressed genes by Ring1B KO in cultured NPCs

Enrichment score

0.3

Highly expressed in control — Highly expressed in Hmga2 KD

$p = 0.023$, $q = 0.069$

c Derepressed genes by Ring1A/B dKO in NPCs *in vivo*

Enrichment score

0.5

Highly expressed in control — Highly expressed in Hmga2 KD

$p = 0.0$, $q = 0.0$

d Core enrichment genes in Ring1 KO cells

p value

10^0 10^{-3} 10^{-6}

Head development

Brain development

Nervous system development

Central nervous system development

Regulation of neural precursor cell proliferation

Fig. S1

a

E7.0 injection → E9.0   E8.0 injection → E11.0

CFSE

b

	Injected embryos	Survival rate (% of injected)	CFSE-positive (% of injected)
E7.0 injection → E9.0	28	25 (89%)	18 (64%)
E8.0 injection → E10.0	30	24 (80%)	23 (77%)

Fig. S2

a

E7.0 IUE → E9.0   E8.0 IUE → E10.0

b

	Injected embryos	Survival rate (% of injected)	mCherry-positive (% of injected)
E7.0 injection → E9.0	36	26 (72%)	7 (19%)
E8.0 injection → E10.0	20	11 (55%)	8 (40%)