Superior Colliculus-Projecting GABAergic Retinal Ganglion Cells Mediate Looming-

Evoked Flight Response

Xue Luo[1,12¶], Danrui Cai[1¶], Kejiong Shen[1], Qinqin Deng[1], Xinlan Lei[1], Sen Jin[2], Wen-Bo

Zeng[3], Hua Li[4], Fuqiang Xu[5,6], Lu Huang[7,8], Chaoran Ren[8,9], Min-Hua Luo[3], Ting Xie[4, 10],

Yin Shen[1,11]*

[1]Eye Center, Renmin Hospital of Wuhan University, Wuhan University, Wuhan, Hubei,

China

[2]HUST-Suzhou Institute for Brainsmatics, JITRI Institute for Brainsmatics, Suzhou, China

[3]State Key Laboratory of Virology, CAS Center for Excellence in Brain Science and

Intelligence Technology (CEBSIT), Wuhan Institute of Virology, Chinese Academy of

Sciences, Wuhan, China

[4]Stowers Institute for Medical Research, Kansas City, MO, USA

[5]Brain Cognition and Brain Disease Institute (BCBDI), Shenzhen Institutes of Advanced

Technology, Chinese Academy of Sciences, Shenzhen-Hong Kong Institute of Brain Science-

Shenzhen Fundamental Research Institutions, Shenzhen, China

[6]Centre for Brain Science, State Key Laboratory of Magnetic Resonance and Atomic

Molecular Physics, Key Laboratory of Magnetic Resonance in Biological Systems, Wuhan

Institute of Physics and Mathematics, CAS Centre for Excellence in Brain Science and

Intelligence Technology, Chinese Academy of Sciences, Wuhan, China

[7]Department of Neurology and Stroke Center, The First Affiliated Hospital of Jinan

University, Guangzhou, China

[8]Guangdong-Hongkong-Macau Institute of CNS Regeneration, Ministry of Education CNS

Regeneration Collaborative Joint Laboratory, Jinan University, Guangzhou, China

[9]Guangzhou Regenerative Medicine and Health Guangdong Laboratory, Guangzhou, China

[10]Department of Anatomy and Cell Biology, University of Kansas School of Medicine,

Kansas City, KS, USA

[11]Frontier Science Center for Immunology and Metabolism, Medical Research Institute,

Wuhan University, Wuhan, China

[12]Division of Life Science, The Hong Kong University of Science and Technology, Clear

Water Bay, Kowloon, Hong Kong, China

[¶]These authors contributed equally to this work.

*Corresponding author

E-mail: yinshen@whu.edu.cn. (YS)

Grant support:

This work was supported by grants from the National Key R&D Program of China

(2017YFE0103400 to Y.S.), the National Nature Science Foundation of China (81470628 Y.

S. 81800872 to Q.D.).

Running title: spgRGCs mediate innate fear response

Key words: spgRGCs, innate defensive behavior, GABA, looming, superior colliculus, flight

response

Abstract

The looming stimulus-evoked flight response is an experimental paradigm for studying innate defensive behaviors. However, how the visual looming stimulus is transmitted from the retina to the brain remains poorly understood. Here, we report that superior colliculus (SC)-projecting RGCs transmit the looming signal from the retina to the brain to mediate the looming-evoked flight behavior by releasing GABA. In the mouse retina, GABAergic RGCs are capable of projecting to many brain areas, including the SC. Superior colliculus (SC)-projecting GABAergic RGCs (spgRGCs) are mono-synaptically connected to the parvalbumin-positive SC neurons known to be required for the looming-evoked flight response. Optogenetic activation of spgRGCs triggers GABA-mediated inhibition in SC neurons. The ablation or silence of spgRGCs compromises looming-evoked flight response but not image-forming functions. Therefore, this study shows that spgRGCs control the looming-evoked flight response by regulating SC neurons via GABA, providing novel insight into the regulation of innate defensive behaviors.

Introduction

The ability to detect the naturally occurring looming stimulus, which mimics

approaching objects, is a conserved innate defensive behavior important for the survival of

animal species, ranging from *Drosophila* to mammals, including humans [1-3]. The looming

stimulus-induced flight response in mice has been recently established as an effective model

for studying innate defensive behaviors in animals [4-6]. The neural circuits for mediating the

looming-evoked innate defensive behavior in the mouse brain have recently been shown to

involve different brain areas, including the superior colliculus (SC). However, how the

looming stimulus signal is transmitted from the retina to the brain remains largely elusive.

The looming-evoked innate defensive behavior has recently been shown to involve the

brain neural circuits mediated through the SC, the parabigeminal nucleus (PBGN), lateral

posterior thalamic nucleus (LPTN), the ventral tegmental area (VTA), the zona incerta (ZI)

and the amygdala in the mouse [4]. The SC is generally believed to serve as a sensory-motor

center for processing and controlling innate visual behaviors [7, 8]. Parvalbumin positive

(PV$^+$) glutamatergic excitatory neurons in the superficial layer of the SC project to different

brain areas to mediate the looming-evoked escape and freezing behavior [4]. Interestingly, the

looming stimulus simultaneously activate two distinct groups of PV$^+$ GABAergic SC neurons

projecting the PBGN and the LPTN, which both innervate the amygdala where the fear

response is initiated, to mediate escape and freezing responses, respectively [9]. Also,

glutamatergic SC neurons innervate and send excitatory signals to GABAergic neurons in the

VTA, which further project to the central nucleus of the amygdala, mediating the looming-

evoked defensive behavior [10]. Although GABAergic neurons in the ZI directly innervate

excitatory neurons in the dorsolateral and ventrolateral compartments of periaqueductal gray

(PAG) to drive escape and freezing responses, it remains unclear how these neurons are

connected to the SC [11].

The visual looming stimulus must be transmitted through retinal ganglion cells (RGCs)

since they are the only cell type connecting the eye to the brain. Interestingly, GABAergic

dorsal raphe nuclei (DRN) neurons antagonize the serotonergic DRN neurons to activate the

SC-LPTN-amygdala pathway to mediate the looming-evoked flight response [6]. It still

remains unclear how the SC/DRN-projecting RGCs transmit the looming stimulus signal.

Recently a subset of melanopsin-expressing intrinsically photosensitive RGCs (ipRGCs) in

mice that release GABA at the non-image-forming brain targets, including suprachiasmatic

nucleus (SCN) and olivary pretectal nucleus (OPN), was identified. Meanwhile this

GABAergic circuit decreases the sensitivity of the non–image-forming visual system at low

light levels [12]. Here we identified SC-projecting GABAergic RGCs (spgRGCs) transmit the

looming stimulus signal and mediate the looming-evoked flight response by releasing GABA.

Results

The mouse retina contains a population of GABAergic RGCs projecting to different

regions of the brain

Earlier studies suggest the existence of GABA-immunoreactive RGCs in turtle [13],

rabbit [14, 15] and , and mouse retinas. To furthermore confirm the existence and functions of

GABAergic RGCs in the mouse retina, the recombinant adeno-associated virus (AAV)

expressing EYFP-EYFP in a Cre-dependent manner were injected into the vitreous of adult

eyes of *Gad2-Cre* or *vGAT-Cre* mice, which are specifically expressed in GABAergic

neurons [16-20] (Fig 1A). In addition to EYFP-labeled GABAergic amacrine cells in the

inner nuclear layer (INL), EYFP-labeled GABAergic RGCs in the ganglion cell layer (GCL)

also existed based on their EYFP-positive optic nerve and laminar positions [21-23] (Fig 1B).

Those EYFP-labeled RGCs were not caused by virus leakage since there were no EYFP-

labeled retinal cells in negative controls (S1 Fig). Consistently, 13% of RGCs (n=6225) are

positive for *Gad2* mRNA expression based on the previously published scRNA results [24]

(Fig 1C). Those labeled YFP-labeled GABAergic RGCs project their axons in the optic track

(Opt) in the brain, and mainly innervate the SCN, the lateral geniculate nucleus (LGN) and

the SC, but not the dorsal raphe nucleus (DRN) (Fig 1D-H). In the LGN, the axonal arbors of

the GABAergic RGCs spread across the LGN, but appear to be more abundant in the dorsal

side (DLG) and intergeniculate leaflet (ILG) than in the ventral side (VLG) (Fig 1F). In the

SC, the axonal arbors of GABAergic RGCs are more restricted to the superficial and

intermediate layers (SL and IL) in the SC (Fig 1G). Taken together, GABAergic RGCs exist

in the adult mouse eye and are capable of projecting to the brain areas participating in image-

forming and non-image-forming functions.

Fig 1. GABAergic RGCs project to multiple areas of the mouse brain. (**A**) Schematic

diagrams explaining the anterograde labeling of GABAergic RGCs and the gene structure for

Cre-dependent expression of two copies of EYFP in the AAV. (**B**) EYFP-labeled GABAergic

neurons in the *Gad2-Cre* (left) and *vGAT-Cre* (right) retinas. EYFP-labeled RGCs are

expressing gamma-aminobutyric acid (GABA) (shown in high magnifications, 100X), and

can also extend EYFP-positive axons in the optic nerve (ONL: outer nuclear layer; INL: inner

nuclear layer; GCL: ganglion cell layer; Scale bar: 20 μm). (**C**) *Gad2* mRNA expression in

mouse RGC subpopulations based on the previously published scRNA results. (**D-H**) Brain

projections from the EYFP-labeled RGCs revealed by anterograde labeling. These

retinorecipient brain regions (shown in high magnifications; right) correspond to the green-

filled areas in the mouse brain atlas (left) (SCN: suprachiasmatic nucleus; Opt: optic tract;

SC: superior colliculus; DLG: dorsal lateral geniculate nucleus; IGL: intergeniculate leaf;

VLG: ventral lateral geniculate nucleus; DRN: dorsal raphe nucleus. Scale bar: 100 μm).

spgRGCs are capable of releasing GABA to elicit physiological responses in SC neurons

To determine if spgRGCs can functionaly release GABA to elicit post-synaptic

responses in SC neurons, channel rhodopsin 2 and EYFP fusion protein (ChR2-EYFP) were

expressed in the AAV-infected GABAergic RGCs. When spgRGCs were activated by 470 nm

laser light, post-synaptic SC responses were recorded (Figs 2A and 2B). Brief pulses of light

can induce sustained inward currents in the ChR2-EYFP-expressing RGCs, indicating that

ChR2-EYFP can be functional in GABAergic RGCs (Fig 2C). In the SC, light-mediated

ChR2 activation in the ChR2-EYFP-expressing RGC axonal fibers induces inhibitory

postsynaptic currents (IPSCs) when $V_{holding}$ was set to 0 mV, which is close to the reveasl

potential for cations [25] (Figs 2D and 2E). The addition of bicuculline (10 µM), $GABA_A$

receptor antagonist, to the recording bath solution (TTX+D-AP5+CNQX）could reversibly

suppress the light-evoked responses (n=26; 21.7±4.2 pA) (Fig 2E). To further characterize the

properties of ChR2-induced currents, in all the neurons that we recorded, we also found

ChR2-evoked excitatory (-70 mV) and inhibitory (0 mV) currents in some postsynaptic

neurons (n=14) in the presence of TTX (Figs 2F and 2G). Our findings indicate that spgRGCs

innervate and functionally inhibit SC neurons through releasing GABA.

Fig 2. ChR2-mediated activation causes GABAergic RGCs to release GABA and evoke

electrophysiological responses in SC neurons. (**A**, **B**) Schematic representation of

optogenetic stimulation of GABAergic RGCs and recordings in labelled RGCs and SC

neurons in tissue slices. (**C**) Patch-clamp recording of the AAV-ChR2-EYFP infected RGCs

in the *vGAT-Cre*/*Gad2-Cre* retina. Light stimulation of ChR2 (470 nm, 500 ms) induces

typical inward currents in the ChR2-EYFP-expressing RGCs. Scale bar: 25 µm. (**D**)

Recording of an SC neuron receiving inputs from GABAergic RGCs (scale bar: 10 µm). (**E**)

Postsynaptic IPSCs of a GABAergic RGC-projecting SC neuron in response to ChR2-

mediated activation of RGCs (470nm blue light for 5 ms) are reversibly inhibited by $GABA_A$

receptor antagonist Bicuculline (10 µM). All recordings are performed with the 0 mV holding

potential ($V_{holding}$) in the presence of D-AP5, strychnine, CNQX and TTX. Right:

quantification results on IPSCs. Light stimulation (470 nm) is marked by a blue bar. (F) The

properties of ChR2-induced currents. IPSC/（IPSC+EPSC）ratios for $ChR2^+$ neurons are

analyzed from vGAT-Cre/Gad2-Cre mice. (G) Example averaged traces of ChR2-evoked

IPSC (upper) and IPSC+EPSC (lower). IPSC: ChR2 evoked excitatory (-70 mV), but no

inhibitory (0 mV) currents in the presence of TTX; IPSC+EPSC: ChR2-evoked excitatory (-

70 mV) and inhibitory (0 mV) currents in the presence of TTX. Statistical analysis: one-way

Student's t-test; ***: $P < 0.001$.

spgRGCs are composed of ON, OFF and ON-OFF types

To further characterize the distribution, dendritic morphology and light responses of the

spgRGCs, Cre-dependent EYFP-EYFP-expressing AAV viruses were injected into the SC

region of the *vGAT-Cre* or *Gad2-Cre* brain for retrogradely labeling spgRGCs (Figs 3A and

3B). Retrogradely labeled RGCs are GABAergic since the EYFP-labeled RGCs express

GABA (n=12 mice) (Fig 3C). By comparing to the Brn3a-positive ($Brn3a^+$) RGCs (Center:

2070.25 ± 166.45 per mm^2; Midperiphery: 2105.09 ± 158.21 per mm^2; Periphery: $1550.68 \pm$

102.62 per mm^2), which are more dense in the central retina than in the peripheral retina [26],

the EYFP-labeled spgRGCs showed a higher density in the peripheral retina (220.39±34.86

per mm^2) than in the central retina (152.78±27.47 per mm^2) (Figs 3D-F) and midperipheral

retina (211.13±24.37 per mm^2). To investigate the proportion of spgRGCs in all SC

projecting RGCs, we injected AAV2/2-Ef1α-EYFP into the SC site of C57BL/6J mice (n=4).

Four weeks later the retina tissues were collected and the labelled cells were counted (Center:

466.79±95.1 per mm^2; Midperiphery: 960.71±134.04 per mm^2; Periphery: 1604.51±

302.42 per mm^2) (Data not shown). The percents of GABA-positive cells in the SC-projecting

RGCs were 32.73%, 21.98% and 13.74% in the central, midperipheral and peripheral retina.

74% of EYFP-labeled spgRGCs had the soma diameter smaller than 12.5 µm, which was

considered to be a small RGC soma [27]. Based on stratification patterns and light-induced

responses, RGCs are classified into ON, OFF, and ON-OFF subtypes. spgRGCs can also be

assigned into ON, OFF, and ON-OFF subtypes [28] (ON cells: n=5; OFF cells: n=3; ON-OFF

cells: n=4) based on stratification and induced-response pattern (Figs 3G-I). Thus, spgRGCs

exhibit preferential periphery distribution in the retina.

Fig 3. Morphological and functional characterization of spgRGCs. (**A**, **B**) Experimental

design: *AAV-EF1a::DIO-EYFP-EYFP* viruses are delivered into the SC to retrogradely trace

spgRGCs. Scale bar: 200 µm. (**C**) EYFP-labeled RGCs also express GABA. Scale bar: 20

µm. (**D**) Representative images of the distribution of EYFP (**upper**) or Brn3a (**lower**) positive

RGCs in the flat mount retina of *vGAT-Cre* mice (Scale bar: 500 µm). Highly magnified

regions (Scale bar: 20 µm) and counterstained cells (Scale bar: 10 µm) are shown in the right.

(**E**) Retinal distributions of the SC-projecting GABAergic RGCs versus Brn3a[+] RGCs (n=24;

means ± SEM). (**F**) Histogram showing the soma size distribution of the SC-projecting

GABAergic RGCs. (**G**-**I**) Light responses and dendritic morphologies of the SC-projecting

ON (**G**), OFF (**H**) and ON-OFF (**I**) GABAergic RGCs. Top: flat view; bottom: side view.

Scale bar: 20 µm.

spgRGCs mediate the looming-evoked flight response

To further determine the downstream neurons of spgRGC-projecting SC neurons, we

injected the herpes simplex virus (HSV) expressing double-floxed four copies of tandemly

repeated GFP genes (HSV-LSL-TK-G4) into the vitreous of *vGAT-Cre* mice to label trans-

synaptic neurons (Fig 4A). HSV-LSL-TK-G4 as an anterograde transsynaptic virus tool，is

capable of identifying potential postsynaptic targets and facilitates the identification of the

cell types. GFP was detected in the SC, the PBG nucleus, the DLG (the relay center in

thalamus for the visual pathway) and the pontine nucleus (Pn; involved in motor activity) in

the brain (Fig 4B). In the SC, the trans-synaptically labeled SC neurons were parvalbumin-

positive (PV[+]) (Fig 4B). These results indicate that spgRGCs are synaptically connected to

PV[+] SC neurons, which are known to form the circuitry controlling the looming-induced

flight behavior [4, 9], suggesting that spgRGCs might be involved in the looming-evoked

flight response.

To directly test their requirement for the looming-evoked flight behavior, we sought to

selectively ablate spgRGCs by expressing diphtheria toxin subunit A (DTA), which

inactivates elongation factor 2 to terminate protein synthesis and thus kill DTA-expressing

cells [25, 29]. AAV2/2-fDIO-DTA was injected vitreously and AAV2/retro-Ef1α -DIO-FLP

was injected into the SC of the *vGAT-Cre* or *Gad2-Cre* mice, to simultaneously infect the

EYFP-labeled spgRGCs for allowing the Cre-dependent expression of FLP that then drives

FLP-dependent DTA expression specifically in spgRGCs(Fig 4C). All injected animals were

preselected for fast return to shelter through two rounds of looming stimulus. Two months

after dual virus infection, EYFP-labeled spgRGCs have been reduced by approximately 90%

(before ablation/phase 1: 204±10.68 EYFP-positive spgRGCs/mm^2, n=11; after

ablation/phase 3: 25.97±2.17 EYFP-positive spgRGCs/mm^2, n=10; *P*<0.001). In the looming

stimulus-induced flight behavior test , there was a significant increase in both "the time to the

shelter" and "the total distance" for the mice depleting spgRGCs (Figs 4D-G, Supplemental

Movie 1-4). Moreover, the defective looming-evoked flight response caused by spgRGC

ablation persists 4 months after the ablation (Figs 4F and 4G). Consistently, the ablation of

spgRGCs significantly decreased the number of looming stimulation-induced c-Fos-positive

neurons in both the SC and the PBGN, further supporting that spgRGCs are required for

looming-induced neuronal activities in the SC-PBGN pathway (Figs 4H and 4I). c-Fos

expression is indicative of neuronal activation upon looming signaling-induced stimulation.

Our findings indicate that spgRGCs are required for the looming-evoked flight response by

activating the SC-PBGN pathway.

Fig 4. spgRGCs are required for the looming-evoked flight response. (**A**) Experiment

design for the trans-synaptic labeling of GABAergic RGCs. (**B**) By infecting the *Gad2-Cre* or

vGAT-Cre mouse eye with Cre-dependent HSV viruses, the SC, the PBG and the Pn in the

brain are labeled by GFP. (Left: Scale bar: 500 µm; Right: Scale bar: 10 µm). (**C**)

Experimental strategy for ablating the SC-projecting GABAergic RGCs (phase 1: Expressing

FLP in the EYFP-labeled GABAergic RGCs; phase 2: DTA expression via the FLP/FDIO

system; phase 3: DTA-induced apoptosis). The SC-projecting GABAergic RGCs can be

effectively eliminated. Scale bar: 50 µm (**D-G**). With the expanding looming stimulus at 2-20

degrees for 0.6 second, representative movement traces of the control (VEH) and DTA-

expressed (DTA+FLP) *vGAT-Cre* (top, **E**) and *Gad2-Cre* (bottom, **E**) mice. **F** and **G**:

quantification results on the movement distance and the time interval from the start to the end

point, respectively. (**H-J**) c-Fos positive cells in SC and PBG in looming stimulated control

mice (**H**) and DTA+FLP treated mice (**I**). **J**: quantification results on c-Fos positive cells in

SC (top) and PBG (bottom) in each group of mice. Scale bar: 50 µm. Mann-Whitney U test;

**: $P < 0.01$, ***: $P < 0.001$, ****: $P < 0.0001$.

spgRGCs are dispensable for image forming functions

Electroretinography (ERG) and the optomotor response were used to investigate if

spgRGCs are required for image-forming functions. There were no significant changes in the

a-wave and b-wave amplitudes of scotopic ERGs between the control and ablation groups,

indicating that spgRGCs are dispensable for overall retinal functions (Figs 5A and 5B).

Although the a wave and b wave correspond to photoreceptor and bipolar cell activity, and

therefore should not be impaired by a lack of spgRGCs. The optomotor response was used to

determine visual acuity based on head movements in response to the moving vertical grating

stimulus. There was no significant difference in the visual acuity between the control and

ablation groups (Figs 5C-G). These results indicate that spgRGCs are not required for visual

image formation.

Fig 5. spgRGCs are dispensable for innate visual functions. (**A, B**) ERG amplitudes of a-

wave (left) and b-wave (right) show no significant changes in the ablation groups. (**C-G**) A

grating stimulation-based OMR setup (**C**), there are no significant differences on the

optomotor response (**D**, **F**) and the maximum spatial frequency (visual acuity; **E**, **G**) under

scotopic and photopic conditions in the control or DTA+FLP induced ablation groups.

Inhibition of spgRGCs suppresses the looming-evoked flight response

To determine whether inhibitions of SC-projecting GABAergic RGCs are required for

looming-evoked response, using the dual-virus intersectional strategy we chemogenetically

inhibited spgRGCs by injecting rAAV-DIO-FLP which encoding Cre-dependent FLP

bilaterally into the SC and at the same time rAAV-FDIO-hM4D-EGFP (n=5) which encoding

FLP- dependent hM4D-EGFP, or rAAV-FDIO-EYFP (n=3) as the control (VEH), into the

vitreous chamber bilaterally of vGAT-cre mouse , which upon FLP excision would result in

expression of hM4D in spgRGCs (Fig 6A). All injected animals were preselected. Post hoc

histological analysis revealed that the hM4D-EGFP expressed in the whole retina (Fig 6B).

Behaviorally, we found that CNO administration significantly increased the total duration in

hM4D mice but not in the vehicles, furthermore the looming-evoked response of partial

animals vanished due to they did not return to shelter during entire stimulation 14.6 s. Taken

together, these results demonstrate that inhibition of spgRGCs suppresses looming-evoked

flight response.

Aiming at removing the effect of other neurotransmitters except GABA of spgRGCs, AAV

with short hairpin (shRNA) was adopted to knock down vGAT expression by injecting

rAAV-U6-Loxp-CMV-EGFP-SV40-polyA-Loxp-shRNA (vGAT shRNA group) into the

vitreous of the vGAT-Cre mice (S2A Fig). In order to assess the effectiveness and tissue-

specificity of the vGAT-Cre combining with LoxP-shRNA system, western blot analysis

indicated that the system was in position to efficiently silence vGAT in retina. Retina

transfected with vGAT shRNA in two months manifested an inferior expression of vGAT

protein to the Control (scramble shRNA group) (S2B Fig). Afterwards, the looming behavior

produced by knockdown vGAT in the GABAergic RGCs was compared, vGAT shRNA

group conveyed an increased escape latencies and duration during the trials in contrast with

the Control group (S2C-D Figs, Supplemental Movie 5-6). In addition, there was no

significant diversity between the Scramble and the vGAT shRNA group in the optomotor

response (S2E Fig). Nevertheless, specificly knockdown vGAT in spgRGCs remains to be

confirmed.

Fig 6. Chemogenetic inhibition of spgRGCs suppresses looming-evoked response. (A)

Schematics of the dual-virus intersectional approach to target spgRGCs. (**B**) Representative

images showing hM4D expression as revealed by immunostaining of GFP in the flat mount

retina. 20X, Scale bar: 30 µm. (**C**) Typical movement traces of the saline and CNO groups.

(**D**) Quantification results on the movement distance and the time interval from the start to the

end point of the vehicle and hM4D groups . Mann-Whitney U test; ***: $P < 0.001$.

Discussion

The ability to detect rapidly approaching objects, which mimic predators, is a highly

conserved innate defensive behavior in *Drosophila*, naïve rodents [30] and primates [31].

Although the looming-evoked neural circuits in the mouse brain have been nicely studied, it

remains poorly understood how the visual looming stimulus is transmitted from the retina to

the brain. Here, we show that mouse spgRGCs directly innervate PV$^+$ SC neurons to mediate

the looming-evoked flight behavior through the SC-PBGN pathway. In the adult mouse

retina, GABAergic RGCs project their axons to multiple brain areas, including SC, SCN,

OPN and LGN. Moreover, approximately 13% RGCs are GABAergic and belong to ON,

OFF and ON-OFF RGC subtypes. In addition, the SC-projecting GABAergic RGCs can

release the inhibitory GABA to their mono-synapsed SC neurons when activated. Finally,

DTA-mediated ablation of those spgRGCs causes the loss of the looming-evoked flight

response. Therefore, we identify a novel GABAergic RGC population in the mouse retina that

projects and releases GABA to the SC, and have further showed that spgRGCs function as a

part of the neural circuitry mediating the looming-evoked defensive behavior.

In the mammalian eye, RGCs are the only cell type sending long axons to connect the

retina to the brain. Earlier studies suggest the existence of GABA-like immunoreactive RGCs

in turtle [13] , bufo [32] and rabbit retinas [14]. Since GABAergic neurons are believed to

function within short distances to regulate local neural networks [33-35], lately a subset of

ipRGCs were showed to release GABA and affected non–image-forming behaviors including

pupillary light reflex and circadian photoentrainment [12]. In this study, we have provided

three pieces of experimental evidence to demonstrate the existence of GABAergic RGCs in

the adult mouse retina. First, *Gad2-Cre* and *vGAT-Cre*-driven Cre-dependent expression of

EYFP reliably label RGCs that project axons to the brain. *Gad2* and *vGAT* (also known as

Slc32a1) encode a GABA synthesizing enzyme and a vesicular GABA transporter,

respectively, which are both known to be expressed in GABAergic neurons [17, 19, 36].

Consistently, our EYFP-labeled RGCs also GABA positive. Second, we have found that 13%

of RGCs express *Gad2* by re-analyzing the recently published RGC SC-seq dataset [37].

Interestingly, most of the 40 RGC subgroups identified by SC-seq, including melanopsin-

positive intrinsically photosensitive RGCs (ipRGCs), also express *Gad2* mRNA. Third,

optogenetic activation drives EFYP-labeled RGCs to release GABA and elicit the IPSC

responses in the projected SC neurons, indicating that those EYFP-labeled RGCs are bona

fide GABAergic neurons.   In addition, EYFP-labeled GABAergic RGCs project their axons

to brain regions as SC, LGN, SCN and OPN. The SC is required for both non-image-forming

and image-forming functions, whereas the LGN is required for only visual image-forming

functions [38, 39]. The SCN and OPN brain areas are required for non-image-forming

functions, circadian photoentrainment [40-42], and the pupillary light reflex (PLR) [12, 43],

respectively. The LGN is required for visual image-forming functions, whereas the SC is

required for both non-image-forming and image-forming functions [38, 39]. Taken together,

multiple types of GABAergic RGCs exist in the adult mouse retina and likely have distinct

biological functions.

This study functionally demonstrates that spgRGCs participate in the looming-evoked

innate flight response via the SC-PBGN pathway. Based on stratification within the IPL and

light-mediated responses, spgRGCs are composed of ON, OFF and ON-OFF types. Those

spgRGCs primarily target the superficial and intermediate layers of the SC, which are known

to be required for the looming-evoked defensive behavior [8, 44]. Indeed, DTA-mediated

ablation of spgRGCs diminishes the looming-evoked flight response, but does not affect

image-forming and other non-image-forming functions. Dual SC/DRN-projecting RGCs have

been previously identified for the looming-evoked flight response[45]. Since newly identified

spgRGCs do not project to the DRN, SC/DRN-projecting RGCs and spgRGCs represent two

distinct RGC populations for mediating the looming-evoked flight response. Our trans-

synaptic tracing results indicate that the GABAergic RGC-projecting SC neurons are also PV$^+$

and are capable of innervating PBGN, PN and DLG in the brain, which should be the same as

the previously identified PV$^+$ glutamatergic SC neurons [4]. Those PV$^+$ SC neurons activate

the PBGN-amygdala circuitry to control the looming-evoked flight response [4, 9]. Results of

c-Fos expression also support that spgRGCs activate PV$^+$ SC and PGBN neurons to mediate

the looming-evoked flight response. Therefore, this study shows that spgRGCs are mono-

synaptically connected to the SC circuitry for mediating the looming-evoked flight behavior

through GABA.

Materials and Methods

Animals

The mouse strains used: C57BL/6, *vGAT-Cre* [*B6J.129S6(FVB)-Slc32a1$^{tm2(cre)Lowl}$/ MwarJ*;

Jackson Laboratory] and *Gad2-Cre* (*B6J.Cg-Gad2$^{tm2(cre)Zjh}$/MwarJ*; Jackson Laboratory). All

mice were group-housed and maintained on a 12h light/12h dark cycle with food and water

provided *ad libitum*. 6-8-week old adult males were randomly divided into experimental and

control groups. All the experiments were done in accordance with the guidelines of Wuhan

University Laboratory Animal Care Committee.

Stereotaxic injection of AAV and HSV

For cell-type-specific retrograde tracing, animals were first anesthetized by intraperitoneally

injecting ketamine and xylazine (60 and 16 mg/kg body weight, respectively). Following the

making of a small craniotomy hole by a dental drill (OmniDrill35, WPI), a micropipette

controlled by a Quintessential Stereotaxic Injector system (Stoelting, Wood Dale, IL, USA,

Micro4; WPI, Sarasota, USA) was used to inject 700 nl *AAV2/retro-EF1a-DIO-EYFP-EYFP-*

WPRE-pA (6.48×10^{12} vg/ml, BrainVTA, Wuhan) virus into the SC of control, *Gad2-Cre* and

vGAT-Cre mice. Coordinates targeting the SC were as follows: 3.9 mm back from the bregma

(anteroposterior), 0.45 mm lateral to the midline (mediolateral), and 1.7 mm deep

(dorsoventral) below the pial surface. The pipette was held in place for 10 min, and

then withdrawn slowly [46].

For anterograde RGCs tracing, a small incision was made in the conjunctiva to expose the

sclera, and 1.5 μl of *AAV2/2-EF1a-DIO-EYFP-EYFP-WPRE-pA* (2.12×10^{12} vg/ml,

BrainVTA, Wuhan) suspension was injected into the center of the vitreous cavity through the

ora serrata with a Hamilton syringe (Hamilton Company, Reno, NV). For anterograde trans-

synaptic tracing, 1.5 μl of *HSV-LSL-TK-G4* (1×10^{10} vg/ml) suspension was injected into the

vitreous chamber, and brain slice were obtained at 6 days [47]. For ablating SC-projecting

GABAergic RGCs, two viruses were performed sequentially. 700 nl of *AAV2/retro-EF1a-*

DIO-FLP-WPRE-pA (7.33×10^{12} vg/ml, BrainVTA, Wuhan) was first injected into the SC of

vGAT-Cre/Gad2-Cre mice, and 1.5 μl of *AAV2/2-EF1α-FDIO-DTA-WPRE-pA* (6.2×10^{12}

vg/ml, BrainVTA, Wuhan) or PBS (as a control) was then injected into the vitreous chamber

bilaterally 3 days later. GABAergic RGC degeneration were detected after one month [46,

48]. Viral-mediated transfection of DREADD receptors. A chemogenetic approach was used

to specifically inhibit spgRGCs. 700 nl of *AAV2/retro-EF1a-DIO-FLP-WPRE-pA* (7.33×10^{12}

vg/ml, BrainVTA, Wuhan) was first injected into the SC, and 2 μl of *AAV2/retro- nEF1α-*

FDIO-hM4D(Gi)-EGFP-WPRE (5.31 × 10^{12} vg/ml, BrainVTA, Wuhan) or *AAV2/retro-*

nEF1a-FDIO-EYFP-WPRE-pA (5.35 × 10^{12} vg/ml, BrainVTA, Wuhan; as a control) was then

injected into the vitreous chamber 3 days later [49]. All the experiments with viruses were

performed in bio-safety level 2 (BSL-2) laboratory and animal facilities.

To remove the putative GABAergic regulation on looming-evoked flight response from

spgRGCs, we used vGAT-selective hort hairpin (shRNA) constructs to knock down vGAT

expression in the retina by injecting vectors into the vitreous of the *vGAT-Cre* mice. The

modified U6 promoter harbors a LoxP-CMV-EGFP-LoxP inside. When the vGAT positive cell

was coinfected with LoxP-shRNA vectors, Cre-mediated recombination results in subsequently

cuts the LoxP-CMV-eGFP-LoxP inside the U6 promoter and activates the U6 promoter. The

activated U6 promoter then drives the down-stream shRNA expression and silences the target

gene. rAAV-U6-Loxp-CMV-EGFP-SV40-polyA-Loxp-shRNA (shvgat) and rAAV-U6-Loxp-

CMV-EGFP-Loxp-shRNA (scramble/control) were injected into the vitreous chamber. AAVs

carrying an shRNA outside two loxP sites (AAV-U6-LoxP-CMV-GFP-LoxP-shRNA, 6.3 × 10^{12}

vg/ml) (BrainVTA, Wuhan) were used for Cre-dependent silencing of vGAT in the *vGAT-Cre*

mice. The shRNA oligonucleotides for targeting vGAT mRNA (shvgat; 5′-TGCTGTTG

ACAGTGAGCGCGGTGTGCTCGTGGTGAATAAGTAGTGAAGCCACAGATGTACTTAT

TCACCACGAGCACACCATGCCTACTGCCTCGGA-3′) and control shRNA (Control; 5′

-TGCTGTTGACAGTGAGCGGCCGCGATTAGGCTGTTATAATAGTGAAGCCACAGAT

GTATTATAACAGCCTAATCGCGGCTGCCTACTGCCTCGGA-3′) were created following

the information described in the literature [50-52]. Control shRNA sequences were used to

construct a non-targeting control virus. Mice were given injections of either vGAT shRNA or

non-targeting scramble shRNA (control shRNA).

Histological procedures

For retina whole mounts, eyes were enucleated after cervical dislocation, and eye cups were

fixed in 4% paraformaldehyde (PFA; pH 7.4) at room temperature for 40 min [53]. Retina

was isolated from the eye cup and flattened. For retina cryosections, eyecups were fixed in

4% paraformaldehyde for 40 min, washed with PBS three times for 5 minutes, incubated

sequentially for 1 h in 10% and 20% sucrose, overnight in 30% sucrose. Then the eyes were

embedded in O.C.T compound (Sakura Finetek, Japan) and frozen at -80°C. Retinal sections

were cut at 14 μm thickness on a freezing microtome (Leica, CM1950, Germany) and

mounted on gelatin-coated slides [54]. For brain cryosections, after transcranial perfusion

with 0.9% saline followed by 4% paraformaldehyde in 0.1 M of PBS, the brain was removed

and post-fixed with 4% paraformaldehyde for 1 day at 4 °C, then transferred into a 30%

sucrose solution for 3 days until sectioning with a cryostat. A series of 50 μm sections were

collected for staining.

Immunostaining

The procedures of immunohistochemistry refer to the previous work [55] (the retina and brain

slices were first incubated for 2 hours in a blocking solution containing 5% bovine serum

albumin (BSA), and 0.2% Triton X-100 in PBS (pH 7.4). And then incubated in primary

antibodies at 4°C overnight. After washing with PBS for 3x15 min, the samples were

incubated in the appropriate secondary antibodies for 2 hours at room temperature. The

samples were washed again in PBS for 3x15 min and mounted in DAPI (4-,6-diamidino-2-

phenylindole; Life Technologies, D1306) for 5 min. The antibodies used in this work were

list as follows: chicken polyclonal antibody against GFP for immunolabeling EYFP$^+$/GFP$^+$

neurons (1:2000, Abcam, ab13970); mouse polyclonal antibody against Brn3a (1:500,

Millipore, MAB1585); goat polyclonal antibody against ChAT (1:200, Millipore, AB144P);

rabbit polyclonal antibody against GABA (1:200, Sigma, A2052), rabbit polyclonal antibody

against c-Fos (1:500; PC38T, Calbiochem). Secondary antibodies were anti-isotypic Alexa

Fluor conjugates (1:500, Invitrogen), incubation for 2 h at room temperature, and after washes

in PBS, the retinas were subsequently mounted in antifade mounting medium.

Confocal microscopy and three-dimensional reconstruction

Images were collected with a confocal laser-scanning microscope (Leica, SP8) and

Brightfield, Fluorescence & FISH Digital Pathology Scanner (Leica, Aperio VERSA 8). For

three-dimensional reconstruction of injected or virus labelled RGCs, the Z-axis interval was

set at 0.4 μm and areas of interest were scanned with a 100X oil immersion objective. A

montage of optical section stacks was created, projected to a 0° X-Y plane and a 90° X-Z

plane to obtain a 3-D reconstruction. Each stack of optical sections covered a retinal area of

325.75 x 325.75 μm^2 (1,024x1,024 pixels). Details of three-dimensional reconstruction and

confocal calibration procedures were described elsewhere [56]. Contrast and brightness were

adjusted. Total soma and dendritic field size of each filled cell were analyzed.

Electrophysiological recording on tissue slices

Brain slices containing the SC were prepared recording to the procedures described

previously [45]. In brief, mice were anesthetized with isoflurane. After decapitation, brains

were immediately immersed in ice-cold oxygenated (95% O_2 and 5% CO_2) cutting solution

which containing (in mM): Choline chloride 97, Ascorbic acid 11, KCl 2.5, $NaHCO_3$ 25,

NaH_2PO_4 1.26, Sodium Pyruvate 3, $CaCl_2$ 0.5, $MgCl_2$ 7.2, glucose 25. Brain slices (350 μm)

was cut using a vibratome (VT1200S, Leica, Germany). The slices were incubated in 34ºC

artificial cerebrospinal fluid (ACSF: NaCl 118, KCl 2.5, $CaCl_2$ 2, NaH_2PO_4 1, $MgCl_2$ 2,

$NaHCO_3$ 26 and glucose 22 in mM) and bubbled with 95% O_2 and 5% CO_2 for 30 minutes

before transfer to the recoding chamber. After an hour incubation at room temperature (25ºC),

slice was transferred to the recording chamber, visualized with an X40 water immersion lens,

DIC optics (BX51WI, Olympus, Japan) and a CCD camera (IR1000, DAGE-MTI, USA). The

data were digitized and recorded on-line using Clampex 10.6 (Axon Instruments, Molecular

Devices). Signals were amplified by MultiClamp 700B (Molecular

Devices), filtered at 1 kHz and sampled at 10 kHz. Electrodes were fabricated from borosilicate

glass (Sutter instrument) using a pipette puller (Sutter, P1000); with resistance 5-7 MΩ. The

internal solution for the recording glass pipette was (in mM): Cs-methanesulfonate 115, CsCl

20, HEPES 10, $MgCl_2$ 2, ATP-Mg 4, GTP-Na 0.4, Na-phosphocreatine 10, EGTA 0.6 (pH 7.2

adjusted by CsOH, mOsm 290). After establishing the whole-cell configuration, the holding

potential was maintained at 0 mV and GABA receptor-mediated IPSCs were pharmacologically

isolated using an antagonist cocktail (D-AP5 50 μM; strychnine 10 μM, CNQX 10 μM and

TTX 1 μM) [57]. ChR2-induced synaptic current was recorded while holding at -70 mV. The

external solution contained NaCl 125, KCl 2.5, $CaCl_2$ 2, $MgSO_4$ 1, $NaHCO_3$ 26, NaH_2PO_4 1.25

and Glucose 20. All drugs were obtained from Sigma-Aldrich or Tocris Bioscience.

For light stimulation, an optic fiber was positioned right above the slice (Mightex, BioLED

Light). Synchronized light-evoked response was triggered by Clampex 10 to deliver light

pulses (470 nm, 5 ms, 0.05 Hz). For retinal slice recording, the isolated retinas were cut into

15-μm-thick slices in Ringer's using a manual cutter (ST-20, Narishige, Japan). The slices

were transferred into a recording chamber with the cut side up and held in place with vaseline.

Whole-cell recordings were obtained from selected neurons. ChR2 positive GABAergic

neurons was confirmed by enhanced yellow fluorescent protein (eYFP) labeling.

CNO administration

The DREADDs program involves injecting an engineered G protein-coupled muscarinic

receptor modified to respond only to the synthetic compound clozapine N-oxide (CNO)

which activates the DREADD producing the spgRGCs inhibition [58]. CNO (Absin,

abs812140) was dissolved in DMSO then diluted to a final concentration of 5 mg/ml CNO in

0.5% DMSO in saline solution. Vehicle injections were 0.5% DMSO in saline solution [59].

To confirm the specificity of DREADD-CNO interactions， DREADD-free mice with

EYFP-CNO group was included. Animals received either equal volume vehicle or CNO (5

mg/kg, i.p.) 30 min before the behavior assays [49].

Looming stimulation test

The looming stimulation test was performed in an open-top acrylic box, as described by

Huang L, et al [6] . The arena had dim lighting from the screen of the monitor. The floor and

three walls were covered with a frosted coating to prevent reflections of the stimulus. A

camera from the side was used to track the movements. An LED monitor was embedded in

the ceiling to present the looming stimulus. The stimulus program was displayed on a LED

monitor (HPZ24i, HP, USA). The looming stimulus, which consisted of an expanding black

disc, appeared at a diameter of a visual angle from 2° to 20° in 0.6 s, for 15 times in 14.6 s.

The stimulation was initiated manually when the mouse was in the center of the arena [5, 6,

48]. Looming tests were conducted in Control mice (n=22 for *Gad2-Cre* mice; 28 for *vGAT-*

Cre mice) and DTA+FLP mice (n=43 for *Gad2-Cre* mice; 35 for *vGAT-Cre* mice).

Optomotor response

Optomotor responses of animals in Control (n=20 for *Gad2-Cre* mice; n=21 for *vGAT-Cre*

mice) and DTA+FLP (n=17 for *GAD-Cre* mice; n=18 for *vGAT-Cre* mice) groups were

measured as described [60]. Briefly, mice were placed on a platform (9 cm diameter, 17.5 cm

above the bottom of the mirror) surrounded by four LCD screens (Lenovo, L1900 PA) that

displayed rotating black-and-white stripes at various spatial frequencies. The platform was set

in the middle of the arena for mouse to stand and move freely. Stimuli were presented on four

computer screens surrounding the animal covering the whole field of view with a texture

projected onto the surface of a virtual sphere. Mice were videotaped with a camera mounted

on top of the platform for subsequent scoring of head tracking movements. Animals were

dark-adapted for 3-4 hours, and then presented with vertical black and white stripes of a

defined spatial frequency. These stripes were rotated alternately clockwise and anticlockwise

for 30 s in each direction and paused for 10 s. The animals were tested with spatial

frequencies increasing from 0.1 to 0.6 cycles per degree (cyc/deg): 0.1, 0.15, 0.2, 0.25, 0.3,

0.35, 0.4, 0.45, 0.5 and 0.6 cyc/deg. Procedures for measuring optomotor responses under

photopic condition were similar to the scotopic condition except that animals were subjected

to 400 Lux light stimulus for 10 mins to allow them to adapt to the light.

Electroretinography

Retinal functions of control (n=16 for *Gad2-Cre* and *vGAT-Cre*) and DTA+FLP (n=24 for

Gad2-Cre and *vGAT-Cre*) mice were measured with a commercial ERG system (RetiMINER,

China). After overnight dark adaptation [61], the mice were anesthetized by intraperitoneally

injecting ketamine and xylazine (60 mg/kg and 16 mg/kg bodyweight, respectively), and then

lightly secured to a stage with fastener strips across the back to ensure a stable, reproducible

position for ERG recordings. All procedures were performed under dim red light. Customized

gold-loop wire electrodes were placed on the center of each eye to measure the electrical

response of the eye to flash stimulation. Reference electrodes were placed subcutaneously

between the ears and a ground electrode was placed in the base of the tail. The ERG consisted

of a 9-step series of full-field flash stimuli dark-adapted (-3 to 1 lg cd·s/m^2) conditions. After

the ERG recording, erythromycin was used on the ocular surface, and the animals were

allowed to recover on a heating pad [62].

Western blotting

Western blotting was performed according to previous work on retinal extracts [54]. They were

homogenized with RIPA lysis buffer (Thermo Scientific, USA) supplemented with a PMSF

(Servicebio, Wuhan). Protein concentration in the supernatant was measured using BCA Protein

Assay kit (Servicebio, Wuhan). Proteins (40 μg) in retinal samples were resolved by 10%

sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) and transferred onto

and electro-blotted to nitro-cellulose membranes (Millipore, USA). The membranes were

subsequently blocked with 5% nonfat milk for one hour and incubated overnight at 4 °C with

primary antibodies, including rabbit anti-SLC32A1 (1:1000; Absin, abs136723), rabbit anti-

GAPDH (1:6000; Antgene, ANT012). The membranes were washed for 30 min in TBS-Tween

20 and incubated with HRP-conjugated goat anti-rabbit IgG (1:5000, Servicebio) for 2 h at RT.

After the second antibody, the membranes were washed again and immunoreactive bands were

subsequently detected using ECL reagents and recorded by X-ray films.

Data quantification and statistical Analysis

All experiments were performed with anonymized samples in which the experimenter was

unaware of the experimental condition of mice. Statistical analysis was performed using SPSS

software (IBM, Armonk, NY, USA). Unpaired *t-test* or Mann-Whitney U test were used.

Error bars represent the mean ± standard error of mean (SEM). $p < 0.05$ is considered as

significant difference between two samples [54, 63].

Author Contributions

Conception and design of the experiments: Y. Shen and X. Luo. Performed the experiments:

X. Luo, Q. Deng, D. Cai and K. Shen. Analysis and interpretation of data: Y. Shen, T. Xie, X.

Luo, D. Cai, K. Shen and H. Li. Wrote the paper: Y. Shen, X. Luo and D. Cai. All authors

approved the final version for publication.

Conflict of interest

554 The authors of this work do not have any conflicts of interest.

References

1. Gibson WT, Gonzalez CR, Fernandez C, Ramasamy L, Tabachnik T, Du RR, et al. Behavioral
responses to a repetitive visual threat stimulus express a persistent state of defensive arousal in
Drosophila. Curr Biol. 2015;25(11):1401-15.doi:10.1016/j.cub.2015.03.058
2. Schiff W, Caviness JA, Gibson JJ. Persistent fear responses in rhesus monkeys to the optical
stimulus of "looming". Science. 1962;136(3520):982-3.doi:10.1126/science.136.3520.982
3. Ball W, Tronick E. Infant responses to impending collision: optical and real. Science.
1971;171(3973):818-20.doi:10.1126/science.171.3973.818
4. Shang C, Liu Z, Chen Z, Shi Y, Wang Q, Liu S, et al. BRAIN CIRCUITS. A parvalbumin-positive
excitatory visual pathway to trigger fear responses in mice. Science. 2015;348(6242):1472-
7.doi:10.1126/science.aaa8694
5. Yilmaz M, Meister M. Rapid innate defensive responses of mice to looming visual stimuli. Curr
Biol. 2013;23(20):2011-5.doi:10.1016/j.cub.2013.08.015
6. Huang. L, T. Y, M. T, Y. X, Y. H, Q. T, et al. A retinoraphe projection regulates serotonergic activity
and looming-evoked defensive behaviour. Nat Commun. 2017;8:14908.doi:10.1038/ncomms14908
7. Zhao X, Liu M, Cang J. Visual cortex modulates the magnitude but not the selectivity of looming-
evoked responses in the superior colliculus of awake mice. Neuron. 2014;84(1):202-
13.doi:10.1016/j.neuron.2014.08.037
8. Ito S, Feldheim DA. The Mouse Superior Colliculus: An Emerging Model for Studying Circuit
Formation and Function. Front Neural Circuits. 2018;12:10.doi:10.3389/fncir.2018.00010
9. Shang C, Liu A, Li D, Xie Z, Chen Z, Huang M, et al. A subcortical excitatory circuit for sensory-
triggered predatory hunting in mice. Nat Neurosci. 2019;22(6):909-20.doi:10.1038/s41593-019-0405-
4

10.   Zhou Z, Liu X, Chen S, Zhang Z, Liu Y, Montardy Q, et al. A VTA GABAergic Neural Circuit Mediates

Visually Evoked Innate Defensive Responses. Neuron. 2019.doi:10.1016/j.neuron.2019.05.027

11.   Chou XL, Wang X, Zhang ZG, Shen L, Zingg B, Huang J, et al. Inhibitory gain modulation of defense

behaviors by zona incerta. Nat Commun. 2018;9(1):1151.doi:10.1038/s41467-018-03581-6

12.   Sonoda T, Li JY, Hayes NW, Chan JC, Okabe Y, Belin S, et al. A noncanonical inhibitory circuit

dampens behavioral sensitivity to light. Science. 2020;368(6490):527-31.doi:10.1126/science.aay3152

13.   Hurd LB, Eldred WD. Localization of GABA- and GAD-like immunoreactivity in the turtle retina.

Vis Neurosci. 1989;3(1):9-20.doi:10.1017/s0952523800012463

14.   Yu BC, Watt CB, Lam DM, Fry KR. GABAergic ganglion cells in the rabbit retina. Brain Res.

1988;439(1-2):376-82.doi:10.1016/0006-8993(88)91498-9

15.   Hirano AA, Brandstatter JH, Brecha NC. Cellular distribution and subcellular localization of

molecular components of vesicular transmitter release in horizontal cells of rabbit retina. J Comp

Neurol. 2005;488(1):70-81.doi:10.1002/cne.20577

16.   Fang YY, Yamaguchi T, Song SC, Tritsch NX, Lin D. A Hypothalamic Midbrain Pathway Essential for

Driving Maternal Behaviors. Neuron. 2018;98(1):192-207 e10.doi:10.1016/j.neuron.2018.02.019

17.   Garcia-Junco-Clemente P, Ikrar T, Tring E, Xu X, Ringach DL, Trachtenberg JT. An inhibitory pull-

push circuit in frontal cortex. Nat Neurosci. 2017;20(3):389-92.doi:10.1038/nn.4483

18.   Taniguchi H, He M, Wu P, Kim S, Paik R, Sugino K, et al. A resource of Cre driver lines for genetic

targeting of GABAergic neurons in cerebral cortex. Neuron. 2011;71(6):995-

1013.doi:10.1016/j.neuron.2011.07.026

19.   Chan KY, Jang MJ, Yoo BB, Greenbaum A, Ravi N, Wu WL, et al. Engineered AAVs for efficient

noninvasive gene delivery to the central and peripheral nervous systems. Nat Neurosci.

2017;20(8):1172-9.doi:10.1038/nn.4593

20.   Milstein AD, Bloss EB, Apostolides PF, Vaidya SP, Dilly GA, Zemelman BV, et al. Inhibitory Gating

of Input Comparison in the CA1 Microcircuit. Neuron. 2015;87(6):1274-

89.doi:10.1016/j.neuron.2015.08.025

21.   Zhang C, Kolodkin AL, Wong RO, James RE. Establishing Wiring Specificity in Visual System

Circuits: From the Retina to the Brain. Annu Rev Neurosci. 2017;40:395-424.doi:10.1146/annurev-

neuro-072116-031607

22.   Baden T, Berens P, Franke K, Roman Roson M, Bethge M, Euler T. The functional diversity of

retinal ganglion cells in the mouse. Nature. 2016;529(7586):345-50.doi:10.1038/nature16468

23.   Pang JJ, Wu SM. Morphology and immunoreactivity of retrogradely double-labeled ganglion cells

in the mouse retina. Invest Ophthalmol Vis Sci. 2011;52(7):4886-96.doi:10.1167/iovs.10-5921

24.   Rheaume BA, Jereen A, Bolisetty M, Sajid MS, Yang Y, Renna K, et al. Single cell transcriptome

profiling of retinal ganglion cells identifies cellular subtypes. Nat Commun.

2018;9(1):2759.doi:10.1038/s41467-018-05134-3

25.   Mitamura T, Higashiyama S, Taniguchi N, Klagsbrun M, Mekada E. Diphtheria toxin binds to the

epidermal growth factor (EGF)-like domain of human heparin-binding EGF-like growth

factor/diphtheria toxin receptor and inhibits specifically its mitogenic activity. J Biol Chem.

1995;270(3):1015-9.doi:10.1074/jbc.270.3.1015

26.   O'Sullivan ML, Punal VM, Kerstein PC, Brzezinski JAt, Glaser T, Wright KM, et al. Astrocytes follow
ganglion cell axons to establish an angiogenic template during retinal development. Glia.
2017;65(10):1697-716.doi:10.1002/glia.23189
27.   Sun W, Li N, He S. Large-scale morphological survey of mouse retinal ganglion cells. J Comp
Neurol. 2002;451(2):115-26.doi:10.1002/cne.10323
28.   Sernagor E, Eglen SJ, Wong ROL. Development of retinal ganglion cell structure and function.
Prog Retin Eye Res. 2001;20(2):139-74.doi:10.1016/s1350-9462(00)00024-0
29.   Collier RJ. Understanding the mode of action of diphtheria toxin: a perspective on progress
during the 20th century. Toxicon. 2001;39(11):1793-803.doi:10.1016/s0041-0101(01)00165-9
30.   Blanchard RJ, Mast M, Blanchard DC. Stimulus control of defensive reactions in the albino rat. J
Comp Physiol Psychol. 1975;88(1):81-8.doi:10.1037/h0076213
31.   Wiener SG, Levine S. Behavioral and physiological responses of mother and infant squirrel
monkeys to fearful stimuli. Dev Psychobiol. 1992;25(2):127-36.doi:10.1002/dev.420250205
32.   Gabriel R, Straznicky C, Wye-Dvorak J. GABA-like immunoreactive neurons in the retina of Bufo
marinus: evidence for the presence of GABA-containing ganglion cells. Brain Res. 1992;571(1):175-
9.doi:10.1016/0006-8993(92)90528-h
33.   Kepecs A, Fishell G. Interneuron cell types are fit to function. Nature. 2014;505(7483):318-
26.doi:10.1038/nature12983
34.   Harris KD, Shepherd GM. The neocortical circuit: themes and variations. Nat Neurosci.
2015;18(2):170-81.doi:10.1038/nn.3917
35.   Huang ZJ. Toward a genetic dissection of cortical circuits in the mouse. Neuron. 2014;83(6):1284-
302.doi:10.1016/j.neuron.2014.08.041
36.   Latremoliere A, Cheng L, DeLisle M, Wu C, Chew S, Hutchinson EB, et al. Neuronal-Specific TUBB3
Is Not Required for Normal Neuronal Function but Is Essential for Timely Axon Regeneration. Cell Rep.
2018;24(7):1865-79 e9.doi:10.1016/j.celrep.2018.07.029
37.   Kakizaki T, Oriuchi N, Yanagawa Y. GAD65/GAD67 double knockout mice exhibit intermediate
severity in both cleft palate and omphalocele compared with GAD67 knockout and VGAT knockout
mice. Neuroscience. 2015;288:86-93.doi:10.1016/j.neuroscience.2014.12.030
38.   Sylvester R, Haynes JD, Rees G. Saccades differentially modulate human LGN and V1 responses in
the presence and absence of visual stimulation. Curr Biol. 2005;15(1):37-
41.doi:10.1016/j.cub.2004.12.061
39.   Marcelja S. Initial processing of visual information within the retina and the LGN. Biol Cybern.
1979;32(4):217-26.doi:10.1007/bf00337645
40.   Hattar S, Kumar M, Park A, Tong P, Tung J, Yau KW, et al. Central projections of melanopsin-
expressing retinal ganglion cells in the mouse. J Comp Neurol. 2006;497(3):326-
49.doi:10.1002/cne.20970
41.   Brown TM, Gias C, Hatori M, Keding SR, Semo M, Coffey PJ, et al. Melanopsin contributions to
irradiance coding in the thalamo-cortical visual system. PLoS Biol.
2010;8(12):e1000558.doi:10.1371/journal.pbio.1000558
42.   Ecker JL, Dumitrescu ON, Wong KY, Alam NM, Chen SK, LeGates T, et al. Melanopsin-expressing
retinal ganglion-cell photoreceptors: cellular diversity and role in pattern vision. Neuron.
2010;67(1):49-60.doi:10.1016/j.neuron.2010.05.023

43.   Bonmati-Carrion MA, Hild K, Isherwood C, Sweeney SJ, Revell VL, Skene DJ, et al. Relationship

between Human Pupillary Light Reflex and Circadian System Status. PLoS One.

2016;11(9):e0162476.doi:10.1371/journal.pone.0162476

44.   Liang F, Xiong XR, Zingg B, Ji XY, Zhang LI, Tao HW. Sensory Cortical Control of a Visually Induced

Arrest Behavior via Corticotectal Projections. Neuron. 2015;86(3):755-

67.doi:10.1016/j.neuron.2015.03.048

45.   Huang. L, Xi Y, Peng Y, Yang Y, Huang X, Fu Y, et al. A Visual Circuit Related to Habenula Underlies

the Antidepressive Effects of Light Therapy. Neuron. 2019;102(1):128-

42.e8.doi:10.1016/j.neuron.2019.01.037

46.   Gauvain G, Murphy GJ. Projection-specific characteristics of retinal input to the brain. J Neurosci.

2015;35(16):6575-83.doi:10.1523/JNEUROSCI.4298-14.2015

47.   Lo L, Anderson DJ. A Cre-dependent, anterograde transsynaptic viral tracer for mapping output

pathways of genetically marked neurons. Neuron. 2011;72(6):938-

50.doi:10.1016/j.neuron.2011.12.002

48.   Hu Y, Chen Z, Huang L, Xi Y, Li B, Wang H, et al. A translational study on looming-evoked

defensive response and the underlying subcortical pathway in autism. Sci Rep.

2017;7(1):14755.doi:10.1038/s41598-017-15349-x

49.   Zha X, Wang L, Jiao Z, Yang R, Xu C, Xu X. VMHvl-Projecting Vglut1+ Neurons in the Posterior

Amygdala Gate Territorial Aggression. Cell Rep. 2020;31(3):107517.doi:10.1016/j.celrep.2020.03.081

50.   Stegmeier F, Hu G, Rickles RJ, Hannon GJ, Elledge SJ. A lentiviral microRNA-based system for

single-copy polymerase II-regulated RNA interference in mammalian cells. Proc Natl Acad Sci USA.

2005;102(37):13212-7.doi:10.1073/pnas.0506306102

51.   Gangadharan G, Shin J, Kim S-W, Kim A, Paydar A, Kim D-S, et al. Medial septal GABAergic

projection neurons promote object exploration behavior and type 2 theta rhythm. Proc Natl Acad Sci

USA. 2016;113(23):6550-5.doi:10.1073/pnas.1605019113

52.   Peng Y-F, Shi Y-H, Ding Z-B, Zhou J, Qiu S-J, Hui B, et al. α-Fetoprotein promoter-driven Cre/LoxP-

switched RNA interference for hepatocellular carcinoma tissue-specific target therapy. PLoS One.

2013;8(2):e53072.doi:10.1371/journal.pone.0053072

53.   Jain V, Srivastava I, Palchaudhuri S, Goel M, Sinha-Mahapatra SK, Dhingra NK. Classical

Photoreceptors Are Primarily Responsible for the Pupillary Light Reflex in Mouse. PLoS One.

2016;11(6):e0157226.doi:10.1371/journal.pone.0157226

54.   Chen Y, Luo X, Liu S, Shen Y. Neuroprotective effect of cannabinoid receptor 1 antagonist in the

MNU-induced retinal degeneration model. Exp Eye Res. 2018;167:145-

51.doi:10.1016/j.exer.2017.11.001

55.   Kircher N, Crippa SV, Martin C, Kawasaki A, Kostic C. Maturation of the Pupil Light Reflex Occurs

Until Adulthood in Mice. Front Neurol. 2019;10:56.doi:10.3389/fneur.2019.00056

56.   Ren C, Luan L, Wui-Man Lau B, Huang X, Yang J, Zhou Y, et al. Direct retino-raphe projection

alters serotonergic tone and affective behavior. Neuropsychopharmacology. 2013;38(7):1163-

75.doi:10.1038/npp.2013.35

57.   Yang K, Ma H. Blockade of GABA(B) receptors facilitates evoked neurotransmitter release at

spinal dorsal horn synapse. Neuroscience. 2011;193:411-20.doi:10.1016/j.neuroscience.2011.07.033

58. Urban D, Roth B. DREADDs (designer receptors exclusively activated by designer drugs):

chemogenetic tools with therapeutic utility. Annu Rev Pharmacol Toxicol. 2015;55:399-

417.doi:10.1146/annurev-pharmtox-010814-124803

59. MacLaren D, Browne R, Shaw J, Krishnan Radhakrishnan S, Khare P, España R, et al. Clozapine N-

Oxide Administration Produces Behavioral Effects in Long-Evans Rats: Implications for Designing

DREADD Experiments. eNeuro. 2016;3(5).doi:10.1523/eneuro.0219-16.2016

60. Abdeljalil J, Hamid M, Abdel-Mouttalib O, Stephane R, Raymond R, Johan A, et al. The optomotor

response: a robust first-line visual screening method for mice. Vision Res. 2005;45(11):1439-

46.doi:10.1016/j.visres.2004.12.015

61. Mees LM, Coulter MM, Chrenek MA, Motz CT, Landis EG, Boatright JH, et al. Low-Intensity

Exercise in Mice Is Sufficient to Protect Retinal Function During Light-Induced Retinal Degeneration.

Invest Ophthalmol Vis Sci. 2019;60(5):1328-35.doi:10.1167/iovs.18-25883

62. Su X, Tan QS, Parikh BH, Tan A, Mehta MN, Sia Wey Y, et al. Characterization of Fatty Acid

Binding Protein 7 (FABP7) in the Murine Retina. Invest Ophthalmol Vis Sci. 2016;57(7):3397-

408.doi:10.1167/iovs.15-18542

63. Dinculescu A, Stupay RM, Deng WT, Dyka FM, Min SH, Boye SL, et al. AAV-Mediated Clarin-1

Expression in the Mouse Retina: Implications for USH3A Gene Therapy. PLoS One.

2016;11(2):e0148874.doi:10.1371/journal.pone.0148874

Supplemental Information

S1 Fig. No virus-mediated labeling in control mice following *rAAV2-DIO-EYFP-

EYFP infection.** (**A**) Experimental strategy for viral delivery. (**B**) No EYFP-positive

cells are detected in the flat-mounted retina 42 days after virus infection, confirming

the Cre-depended EYFP expression in this study. Scale bar: 100 μm.

S2 Fig. Looming-evoked flight response requires functional release of GABA from

RGCs. (**A**) The hairpin (sh) oligonucleotide (sholigo), targeting the vGAT transcript, was

used to knock down vGAT expression. (**B**) Western blot analysis show that it can efficiently

knockdown vGAT. Representative immunoreactive bands of vGAT protein levels, the bottom

panel show the mean optical density (OD) of the band corresponding to vGAT normalized to

GAPDH in the two groups. (**C-D**) Typical movement traces of the Control shRNA and vGAT

shRNA groups. (**D**) Before or 30 days after injection. Quantification results on the movement

distance and the time interval from the start to the end point, respectively. (**E-F**) Functional

release of GABA from spgRGCs are dispensable for the optomotor response. Before or 30

days after injection. there are no significant differences on the optomotor response under

scotopic and photopic conditions in the Control shRNA or vGAT shRNA group.Mann-

Whitney U test; ***: $P<0.001$, ****: $P<0.0001$.

Supplemental Movie 1

Example of looming-evoked defensive behavior in *Gad2-Cre* mice.

Supplemental Movie 2

Example of looming-evoked defensive behavior in *vGAT-Cre* mice.

Supplemental Movie 3

Effects of spgRGCs deletion on looming-evoked flight response in *Gad2-Cre* mice.

Supplemental Movie 4

Effects of spgRGCs deletion on looming-evoked flight response in *vGAT-Cre* mice.

Supplemental Movie 5

Example of looming-evoked defensive behavior in vGAT shRNA group.

Supplemental Movie 6

Example of looming-evoked defensive behavior in control shRNA group.

A

Day 0 — rAAV2-retro-DIO-EYFP-EYFP

Day 28 — Collect tissue

SC

B

EYFP   SC

Zo
SuG
Op
InG
InWh

Bregma -4.04 mm

C

vGAT-Cre

GFP   GABA   Merge

GAD2-Cre

GFP   GABA   Merge

D

EYFP/Brn3a   *vGAT-Cre*

Center   Midperiphery   Periphery

EYFP

Brn3a

S
T   N
I

E

Cell Density

■ Brn3a
■ EYFP

Cells/ mm²

C   M   P

F

Soma size

Percentage (%)

4 8 12 16 20 24 28 32
(μm)

G   ON GABAergic RGC

n=5

20 mV

1 s

OFF
ON

H   OFF GABAergic RGC

n=3

20 mV

1 s

OFF
ON

I   ON-OFF GABAergic RGC

n=4

20 mV

1 s

OFF
ON

A *vGAT-Cre*

Day 0 — HSV-LSL-TK-G4

Day 6 — Collect brain slice

B GFP/DAPI

SC

PBG →

SC PV/GFP/DPAI

PBG GFP/DPAI

Retina
GABAergic
RGCs

SC

PBG

C AAV2/retro-DIO-FLP-WPRE-pA Virus A

Virus B

SC

AAV2/retro-FDIO-DTA-WPRE-pA

Phase-1

SC

Retro-AAV-
DIO-EYFP-EYFP

GABAergic RGCs

Phase-2

Virus A

Virus B

Phase-3

Dead cell Dying cell

D Looming test

0.6 s

20 deg

S

E VEH DTA+FLP

GAD2-Cre

Start
End

vGAT-Cre

Start
End

F

Reactive Distance (cm)

*** ** ***

*** ** ***

Time (d)

G VEH DTA+FLP

From start to end (s)

*** **** *** ****

n=19
n=43

**** *** ****

n=30
n=36

Time (d)

H VEH

SC PBG

SL

IL

DL

DAPI/c-fos

I DTA+FLP

SC PBG

SL

IL

DL

DAPI/c-fos

J Number of c-Fos+ cells

SC

n=6

***

VEH DTA+FLP

PBG

n=6

***

VEH DTA+FLP

A

Virus A
AAV2/retro-DIO-FLP-WPRE-pA

Virus B

SC

AAV2/retro-FDIO-hM4D(Gi)-EGFP
or AAV2/retro-FDIO-EYFP

B

retina

EGFP

C

Saline   CNO

hM4D

● Start
● End

D

Distance (cm)

N.S.   ***

VEH   hM4D

Total Time (s)

N.S.   ***

VEH   hM4D

● Saline
● CNO

A

Day 0 — **rAAV2-DIO-EYFP-EYFP**

Day 42 — **Collect retina**

B WT   EYFP