Using singleton densities to detect recent selection in *Bos taurus*

Matthew Hartfield[1,2], Nina Aagaard Poulsen[3], Bernt Guldbrandtsen[4,5,6], Thomas Bataillon[1]

1) Bioinformatics Research Centre, Aarhus University, DK–8000 Aarhus, Denmark.

2) Institute of Evolutionary Biology, University of Edinburgh, Edinburgh EH9 3FL, UK.

3) Department of Food Science, Aarhus University, Agro Food Park 48, 8200 Aarhus N, Denmark.

4) Center for Quantitative Genetics and Genomics, Department of Molecular Biology and Genetics, Aarhus University, DK–8830 Tjele, Denmark.

5) Rheinische Friedrich-Wilhelms-Universität Bonn, Institut für Tierwissenschaften, Tierzucht, Endenicher Allee 15, D-53115 Bonn, Germany.

6) Department of Veterinary Sciences, Copenhagen University, DK–1870 Frederiksberg C, Denmark.

Corresponding author: Matthew Hartfield (m.hartfield@ed.ac.uk).

Key words: selection, genomics, *Bos taurus*, milk protein, milk fat, stature.

Abstract: Many quantitative traits are subject to polygenic selection, where several genomic regions undergo small, simultaneous changes in allele frequency that collectively alter a phenotype. The widespread availability of genome data, along with novel statistical techniques, has made it easier to detect these changes. We apply one such method, the 'Singleton Density Score', to the Holstein breed of *Bos taurus* to detect recent selection (arising up to around 740 years ago). We identify several genes as candidates for targets of recent selection, including some relating to cell regulation, catabolic processes, neural-cell adhesion and immunity. We do not find strong evidence that three traits that are important to humans – milk protein content, milk fat content, and stature – have been subject to directional selection. Simulations demonstrate that since *B. taurus* recently experienced a population bottleneck, singletons are depleted so the power of SDS methods are reduced. These results inform on which genes underlie recent genetic change in *B. taurus*, while providing information on how polygenic selection can be best investigated in future studies.

Impact statement: Many traits of ecological or economic importance (including height,

disease propensity, climatic adaptation) are 'polygenic'. That is, they are affected by a

large number of genetic variants, with each one only making a small contribution to a

trait, but collectively influence variation. As selection acts on all of these variants

simultaneously, it only changes the frequency of each one by a small amount, making

it hard to detect such selection from genome data. This situation has changed in recent

years, with the proliferation of whole–genome data from many individuals, along with

the development of methods to detect the subtle effects of polygenic selection. Here,

we use data from 102 genomes from domesticated cattle (*Bos taurus*) that has

experienced intense artificial selection since domestication, and test whether we can

detect signatures of recent selection (arising up to 740 years ago). Domesticated

species are appealing for this kind of study, as they are subject to extensive genome

sequencing studies, and genetic variants can be related to traits under selection. We

carried out our analysis in two parts. We first performed a genome–wide scan to find

individual genetic regions that show signatures of recent selection. We identify some

relating to cell regulation, catabolic processes, neural-cell adhesion and immunity. In

the second part, we then analysed genetic regions associated with three key traits:

milk protein content, milk fat content, and stature. We tested whether these regions

collectively showed a signature of selection, but did not find a significant result in either

case. Simulations suggest that the domestication history of cattle affected the power

of these methods. We end with a discussion on how to best detect polygenic selection

in future studies.

Introduction

Determining which genomic regions have been subject to selection is a major research goal in evolutionary genetics. Traditional methods have focused on detecting strong selection affecting individual genes (Nielsen, 2005; Vitti *et al.*, 2013; Stephan, 2019). An alternative process is 'polygenic selection', where many loci contribute to genetic variation in a trait, so selection acting on it is expected to generate small and simultaneous allele frequency changes at multiple loci (Pritchard & Di Rienzo, 2010; Pritchard *et al.*, 2010). Many polygenic models have been formulated to account for both the response to phenotypic selection, and the maintenance of genetic variance in quantitative traits [reviewed by Sella & Barton (2019); Barghi *et al.* (2020)]. Among them is Fisher's infinitesimal model, which is important for its historical role in uniting population and quantitative genetics, and its recent renaissance in the context of genome–wide association studies (Fisher, 1918; Barton & Keightley, 2002; Barton *et al.*, 2017; Charlesworth & Edwards, 2018; Visscher & Goddard, 2019). However, whereas it has been possible to identify which genetic regions contribute to trait variation, it has historically been hard to infer which alleles have been involved in the polygenic selection response. Extensive theoretical studies of how alleles at multiple loci act when a population adapts to a new optimum generally find that 'large–effect' alleles, which strongly affect a trait, are the first to spread and fix while 'small–effect' alleles take much longer to reach high frequencies (de Vladar & Barton, 2014; Wollstein & Stephan, 2014; Jain & Stephan, 2015, 2017a, 2017b; Stetter *et al.*, 2018; Thornton, 2019; Hayward & Sella, 2019). Furthermore, if epistasis exists between variants, many selected alleles do not reach fixation as they eventually become deleterious (de Vladar & Barton, 2014; Jain & Stephan, 2017b).

The spread of large–effect alleles may also be impeded if a faster adaptive response

can be otherwise realised through changes at many small–effect alleles (Lande,

1983; Chevin & Hospital, 2008; Pavlidis *et al.*, 2012; Chevin, 2019). Alternatively, if

the optimum shift is sufficiently big, then large-effect mutations that first go to fixation

can subsequently be replaced by small–effect variants over longer timescales (on the

order of the population size; Hayward and Sella (2019)). Overall, only a small

proportion of loci affected by polygenic selection are expected to fix sufficiently

quickly to leave selection signatures in genomic data (Pavlidis *et al.*, 2012; Thornton,

2019).

Due to this difficulty, earlier methods for detecting polygenic selection focused

on cases where selection favours distinct phenotypes in different populations, so trait

differentiation amongst populations will be greater than expected under neutral drift.

Tests for this form of selection relied on comparing Q_{st} and F_{st} statistics, which

respectively measured mean genetic differentiation at the trait itself and a set of

neutral loci (Whitlock, 2008; Le Corre & Kremer, 2012; Savolainen *et al.*, 2013). Yet

these methods do not determine which genomic regions are subject to selection. This

situation has now changed with the increased number of genome–wide association

study (GWAS) data that link genotypes and phenotypes, as exemplified by the

development of large cohort studies [e.g., the UK Biobank; Bycroft *et al.* (2018)]. The

release of these data spurred a series of studies and new methods designed

specifically to detect polygenic selection. These methods usually involve determining,

which SNPs affecting a phenotype show correlated changes in frequency (Berg &

Coop, 2014; Racimo *et al.*, 2018; Sanjak *et al.*, 2018; Josephs *et al.*, 2019; Berg *et*

al., 2019a, 2019b; Uricchio *et al.*, 2019; Edge & Coop, 2019; Kreiner *et al.*, 2020;

Wieters *et al.*, 2021; Gramlich *et al.*, 2021); which sets of alleles are associated with

certain environmental or climatic variations (Coop *et al.*, 2010; Turchin *et al.*, 2012;

Robinson *et al.*, 2015; Yeaman *et al.*, 2016; Exposito-Alonso *et al.*, 2018; Zan &

Carlborg, 2018; Exposito-Alonso *et al.*, 2019; MacLachlan *et al.*, 2021; Ehrlich *et al.*,

2021; Fuhrmann *et al.*, 2021; Rowan *et al.*, 2021); or determining which SNPs or

genetic regions explain a large fraction of phenotypic variance and trait heritability

(Zhou *et al.*, 2013; Yang *et al.*, 2015; Gazal *et al.*, 2017; Zeng *et al.*, 2018; Schoech

et al., 2019; Exposito-Alonso *et al.*, 2020; Duntsch *et al.*, 2020; Zeng *et al.*, 2021).

Some of these approaches use overlapping methods.

Detecting recent polygenic selection is much harder, as long periods of time

(number of generations on the order of the population size; Hayward and Sella, 2019;

Thornton, 2019) may be needed to cause detectable frequency changes in alleles

with small effect sizes. Over shorter timescales, these frequency changes are

expected to be more modest and harder to detect (Stephan, 2016; Jain & Stephan,

2017a). A recent breakthrough in detecting these subtle changes was the

development of the 'Singleton Density Score' (SDS), a statistic tailored to detect

recent and coordinated allele frequency changes over many SNPs (Field *et al.*,

2016). Recent selection at a locus favouring one variant will lead to a reduction in the

number of singletons (i.e., variants that are only observed once) around it. The SDS

detects regions that exhibit a reduction in the density of singletons, to determine

candidate regions that have been subject to recent selection. Using this approach,

Field et al. (2016) found correlations between SDS scores at SNPs and their

associated GWAS effect sizes for several polygenic traits in the modern UK human

population, including increased height, infant head circumference and fasting insulin.

Their findings suggested that these traits have been subject to recent selection

during the most recent 75 or so generations (about 2,000 years). However, these

(and other) results that detect selection for increased height may instead reflect

previously unaccounted–for population structure (Novembre and Barton, 2018;

Barton et al., 2019; Sohail et al., 2019; Berg et al., 2019; Uricchio et al., 2019; Edge

and Coop, 2019).

The SDS method is ideally suited to organisms where large amount of whole-

genome data are available, along with QTL or GWAS information that link genotypes

to phenotypes, Domesticated species are attractive systems for studying recent

selection, as selected phenotypes are often already known and these species are

subject to large–scale sequencing studies. Investigating the genetic architecture

underlying rapid selection in these species is also important to determine how they

respond to agricultural practices, and uncover selection targets that can be used to

improve breeding programs (Georges *et al.*, 2018). Domestic cattle *Bos taurus* has

been subject to intensive genomics analyses to improve artificial selection for traits

that are important for human use, including milk protein yield, milk fat content, and

stature (Hayes *et al.*, 2009; Meuwissen *et al.*, 2013; Wray *et al.*, 2019). These traits

are influenced in part by an individual's genome, with significant heritability estimates

being recorded, some as high as 80% (Soyeurt *et al.*, 2007; Haile-Mariam *et al.*,

2013; Buitenhuis *et al.*, 2016). Previous selection scans on *B. taurus* reported

individual regions that were likely to be subject to recent selection, some of which

were close to genetic regions for stature and milk protein content (Lemay *et al.*, 2009;

MacEachern *et al.*, 2009; Qanbari *et al.*, 2010; Boitard & Rocha, 2013; Qanbari *et al.*,

2014; Zhao *et al.*, 2015; Boitard *et al.*, 2016a; Bouwman *et al.*, 2018). However,

stature and milk protein content are polygenic traits, with several genetic regions and

QTLs associated with each (Lemay *et al.*, 2009; Boitard *et al.*, 2016a; Bouwman *et*

al., 2018; van den Berg *et al.*, 2020). While recent methods have been developed to

detect polygenic environmental adaptation (Rowan *et al.,* 2021), there has yet to be a

formal test of whether these intrinsic traits show evidence of polygenic selection.

Here, we applied the SDS method to whole–autosome sequencing data from

102 *B. taurus* Holstein individuals. We first determined genetic regions that have

been subject to recent directional selection, and subsequently tested whether

evidence exists for recent selection acting on a set of QTLs underlying either milk

protein content, milk fat content, or stature in this breed.

Results

Methods outline

We filtered the data to retain only bi–allelic SNPs that had a sensible level of

coverage and did not lie in putatively over–assembled regions (i.e., duplicated

sections that caused many reads to assemble at a specific genetic location). Over–

assembled regions appear as highly heterozygous with elevated coverage, and can

exhibit false signatures of recent selection. We also obtained a set of singletons and

filtered them to retain high-quality variants where both alleles were equally well

covered to remove potentially erroneous calls. We polarised test SNPs using

outgroup sequences and applied the SDS test of Field et al. (2016) to detect recent

selection, with increased SDS values reflecting selection favouring derived SNPs

over ancestral variants. We standardised SDS scores with those of a similar

frequency, so they are normally distributed [similar normalisation was also carried out

by Field et al. (2016)]. These values are denoted sSDS for 'standardised SDS'.

Further details are available in the *Methods* in the Supplementary Text.

Estimating timescale of selection

We first determined the timescale over which we expect to detect selection in

B. taurus using the SDS method. SDS measures the changes in singleton numbers

around putatively selected SNPs, relative to background numbers in the absence of

selection. As singletons arise on the tips of the underlying gene trees, the average tip

length in the genealogy of sequenced samples determines the timescale over which

the SDS detects a signal (Field et al., 2016). As more haploid genomes are included

in the study, the time to first coalescence between two samples decreases, reducing

the tip lengths and therefore shortening the timescale over which SDS detects

selection (Field *et al.,* 2016). We hence simulate tip-ages over a range of sample

sizes to investigate how this timescale changes accordingly.

To calculate the mean tip age, we simulated gene genealogies under two

scenarios. We first simulated the Holstein population demography inferred by Boitard

et al. (2016b), which suggested that this population experienced a sudden decline in

effective population size (N_e) since domestication, but with a present–day N_e (~793)

that is much larger than that inferred from pedigree data [~49; Sørensen et al.

(2005)] or from temporal variation in SNP frequencies (~48; Jiménez–Mena et al.

2016). Hence, we also simulated genealogies under a second model that used the

Boitard et al. (2016b) demographic model, but with the present–day N_e set to 49.

These scenarios will be referred to as the 'High N_0' and 'Low N_0' models,

respectively.

Figure 1 shows simulation results. Depending on the assumed present–day

N_e, the tip length in our sample of 204 alleles (i.e., assuming two per diploid

individual) goes back either 65 or 148 generations. Assuming 5 years per generation

(Boitard *et al.*, 2016b), this timescale corresponds to between 325 and 740 years

ago. Since *B. taurus* domestication started around 10,000 years ago (Zeder, 2008)

the sample size used in this study will only capture selection acting in the very recent

past that is more relevant for breed formation, rather than selection during *B. taurus*

domestication. Sample sizes and tip-ages are linearly related on a log-log scale,

meaning that an increase in sample size will greatly decrease the timescale over

which SDS detects selection. For example, with 500 haplotypes then SDS will detect

selection acting no more than 50 generations ago, depending on the underlying

demographic model.

We will focus on detecting selection signatures assuming the high N_0 model.

Results using the low N_0 model to calibrate scores were broadly similar. They are

outlined in the Supplementary Text; we will highlight when differences arise.

Genome–wide sSDS

Figure 2 plots sSDS values (at SNPs with minor allele frequency greater than

5%) across all autosomes, excluding chromosome 25 (due to an insufficient number

of singletons needed to obtain SDS scores after filtering). Many SNPs have elevated

sSDS scores (158 SNPs at *FDR* < 0.05; 306 for the low N_0 model). Several regions

contain SNPs with significantly high sSDS values (Bonferroni–corrected nominal *P* <

0.05; actual *P* < ~2.7 x 10^{-8}). To further investigate potential selection targets, we

looked for genes that either overlapped significant SNPs or lay 10kb up– or

downstream of them. Linkage disequilibrium (LD), as measured by r^2, decays to

around 0.2 over 50kb in Danish Holstein breeds (Buitenhuis *et al.,* 2016), so genes

within 10kb should be in LD with regions harbouring high sSDS scores. Table 1 lists

these genes, with more targets present under the low N_0 model. Most of these genes

are of unknown function (as listed on UniProt); the list also includes an snRNA.

PPM1L is involved with cellular regulation and the activation of stress–activated

protein kinases. *TDO2* is involved in tryptophan–related catabolic processes, while

NTM is implicated in neural cell adhesion. SNPs with significantly elevated scores

are also found on chromosome 23 near the MHC region, which may reflect over–

dominant selection. All Bonferroni–significant SNPs were removed from subsequent

tests of recent polygenic selection to prevent directional selection from skewing the

underlying sSDS distributions. Figure S1 shows results for the low N_0 model.

Testing for polygenic selection acting on milk protein and stature

If polygenic selection were acting on specific traits, we expect a positive

correlation between the effect size of variant underpinning it, and selection acting on

it as measured by sSDS. We collated sSDS scores of SNPs that lie close to QTLs

reported for either milk fat percentage, milk protein percentage (van den Berg *et al.,*

2020), or those that lie close to stature QTLs (Bouwman *et al.,* 2018). The latter were

inferred from a meta–analysis of GWAS studies conducted in seven Holstein

populations, but not every QTL had an effect size reported in each population. We

hence investigated two overlapping consensus QTL sets, where an effect size was

either reported in at least 6 of 7 populations (yielding 42 QTLs with sSDS scores

associated with them), or where effect sizes were reported in at least 5 of 7

populations (58 QTLs had sSDS scores). We re-polarised SDS scores so that a

positive score reflected a trait-increasing effect; we denote these values 'tSDS'

following Field *et al.* (2016). We then determine if there was a positive correlation

between the absolute $\log_{10}$-value of the QTL *P*-value (a proxy for the effect size) and

tSDS.

Figure 3 shows the relationship between QTL *P*-values and tSDS for SNPs

that lie close to QTLs. Although positive trends are observed, they all exhibit non-

significant correlations (milk fat percentage Spearman ρ = 0.0990, *P* = 0.603; milk

protein percentage Spearman ρ = 0.0354, *P* = 0.758; stature from 6 breeds

Spearman ρ = -0.0739, *P* = 0.642; stature from 5 breeds Spearman ρ = -0.00966, *P*

= 0.943). Relationships remain non-significant after removing an outlier point for the

milk traits whose QTL has an extremely low *P*-value (Figure S2), and also under the

low N_0 model (Figure S3; see figure legends for regression *P*-values).

sSDS (and tSDS) can become correlated along the genome if focal SNPs are

in LD with one another, which was not accounted for in the preceding analyses. To

determine whether LD could have affected these correlations, we randomly

subsampled sSDS scores from SNPs that shared the same chromosome and bin of

derived-allele frequency as the SNPs used in the above analyses, and re-polarised

them to transform them into tSDS values. We then determined the Spearman's ρ

associated with these permuted values to determine whether that for the true data

was significantly elevated (see Methods for details). In all cases, the observed value

was not significantly higher than for permuted values (see Figure S4 for histograms

and exact *P*–values, which all exceed 0.05). We therefore conclude that these QTL

datasets do not harbour SNPs with significantly different tSDS scores compared to

the rest of the genome.

Discussion

Summary of results

We analysed an extensive *B. taurus* genomic dataset to identify signatures of

recent selection in the Holstein breed, and to determine whether the data contained a

signal of polygenic selection acting on milk proteins and QTLs underlying phenotypic

variation in stature. Given the sample size and the demographic history of Holsteins,

the SDS method can detect very recent selection events arising no more than

approximately 740 years ago (Figure 1). A whole–genome scan for sSDS scores

identified several targets of recent directional selection that overlap or lie close to

protein–coding genes (Figure 2; Table 1). The genes whose functions are known are

involved in protein regulation, catabolic processes, and neural-cell adhesion.

Significant values were also observed near the MHC region. We subsequently

investigated whether either milk protein genes or SNPs near stature QTLs

collectively showed evidence of polygenic selection. We did so by testing whether

there is a relationship between the QTL effect size, as measured by its *P*-value, and

tSDS values to SNPs near them. However, no relationship was observed, even after

performing a permutation test (Figures 3, S2-S4). Hence, while sSDS could reveal

specific instances of recent selection, tests based on collective scores of variants

associated with known selected traits yielded no signal of polygenic selection.

Potential reasons for a lack of polygenic selection signal

 Impact of Holstein demographic history

While the SDS method detected individual candidate genes for very recent

selection, we were unable to find strong evidence for polygenic selection acting on

three traits that were subject to artificial selection since domestication. This result is *a*

priori surprising, given that these traits have been subject to recent intense artificial

selection. Recent studies generally find non-zero heritability estimates for them,

indicating that there should be the potential for genetic variants underpinning them to

change in response to artificial selection (Soyeurt *et al.*, 2007; Haile-Mariam *et al.*,

2013; Buitenhuis *et al.*, 2016). In addition, the ratio of the mutation and recombination

rates in cattle is just over three (Boitard *et al.*, 2016b; Harland *et al.*, 2018), indicating

that several informative SNPs exist per haplotypes that should improve the power of

the SDS method [in contrast, this ratio is approximately equal to one in humans

(Field *et al.*, 2016)].

One potential reason for this lack of signal is due to the population history of

Bos taurus. The effective population size of many *B. taurus* breeds appears to have

undergone a decline since domestication (Sørensen et al., 2005; Boitard et al.,

2016b), which likely reflects successive bottlenecks due to domestication, breed

formation and intense recent selection. Population size reductions are known to

reduce the number of low-frequency variants and increases the prevalence of

intermediate-frequency variants (Harpending *et al.*, 1998), which can affect the power

of the SDS method. To understand if the history of *B. taurus* affects the detection of

recent selection in Holstein cattle using SDS, we ran coalescent simulations to

determine its ability to detect ongoing selection, given realistic Holstein population

history and genetic parameters (see Methods for details). We simulated a partial

sweep occurring in the middle of a 10Mb region, either assuming a mutation rate in

line with what has been inferred for Holstein, or one 10-fold higher to replicate

diversity expected in a genetic region with an elevated mutation rate.

For the standard mutation rate, no SDS scores were produced for any

simulations. After inspecting the simulation results, we see that there is a large skew

in the distribution of singleton numbers per individual with a large number of

individuals (over 20 on average) that do not carry singletons at the end of

simulations, preventing the calculation of a local SDS score (Figure 4). This fraction

remained the same irrespective of whether the simulated region was neutral or

subject to selection; the main effect of a sweep was to reduce the mean number of

singletons per individual, which is the signal measured by SDS (Field *et al.*, 2016).

This reduction in overall singleton numbers is consistent with the known effects of

population size contraction on reducing tip lengths (Harpending *et al.*, 1998).

With a 10-fold higher mutation rate, there were fewer cases where no

individual harboured singletons (Figure 4). Accordingly, SDS scores could be

calculated for 65 and 66 out of 100 simulations for the neutral and selective cases

respectively. In these cases, sSDS values were significantly higher in the selected

case than for the neutral case (Figure 5; two-sided Wilcox Test $P = 1.1 \times 10^{-5}$).

However, note that sSDS values is less than one for the selected case, which does

not exceed the FDR threshold in our study (for the high N_0 case, the smallest sSDS

value with FDR < 0.05 is 4.46).

Although singleton numbers differ between the two cases, a reduction in

power could also be caused by a more general reduction in diversity due to the small

recent effective population sizes of cattle. To investigate this effect, we estimated the

fixed *Ne* that would yield the same number of segregating sites in simulations using

the standard mutation rate, based on Watterson's estimator (Watterson, 1975;

Hudson, 1990; see Methods for details). In both cases where selection is present or

absent, *Ne* estimates lie at around 25,000, which is that inferred at approximately

halfway between the onset of domestication and the present day. (Boitard *et al.*,

2016b; Figure S5). Given that estimates are similar irrespective of whether a sweep

was present or not, the reduced population size caused by domestication could have

also affected power due to limiting genetic variation and thus the potential to detect

subtle sweep signatures associated with polygenic selection.

Overall, these simulations are consistent with population size reductions in *B.*

taurus both reducing the overall genetic diversity and the number of singletons, which

limits its ability to detect partial sweeps. SDS is more likely to detect signals in

regions of elevated mutation rate, suggesting there will likely be an ascertainment

bias in where signals are detected in the genome. The reduction in singletons also

reduces the power to investigate SDS values in telomeric regions. SDS values are

calculated using the distance up- and downstream from a SNP to the nearest

singleton, and are undefined if a certain number of samples do not harbour

singletons in either direction (Field *et al.*, 2016). SDS values are hence less likely to

be defined in telomeric regions, as it is generally less feasible to observe singletons

up until the end of the chromosome. This problem is exacerbated if there are few

singletons overall.

Other potential reasons for a lack of signal

Another potential reason for a lack of signal is that the selection response on

these traits may have been driven by large-effect variants that have already fixed in

the population, with a smaller contribution from small-effect mutations. Theoretical

models have shown that more major–effect QTLs are likely to fix if the population lies

further from a fitness optimum (Lande, 1983; Jain & Stephan, 2017b; Thornton,

2019). Domesticated species, which experience strong and sustained directional

artificial selection, especially in recent generations, could thereby fix more adaptive

mutation via sweep–like processes compared to populations evolving in more stable

environments (Lande, 1983; Jain & Stephan, 2017a). Furthermore, once a population

has adapted to a new environment (the domestication phenotype in this case), then

any remaining major–effect mutations are likely to be superseded by variants with

weaker effects, which are harder to detect (Hayward & Sella, 2019). The response to

polygenic selection will be further weakened in smaller populations (John & Stephan,

2020), which could be a factor given the reduced effective population sizes of *B.*

taurus (Sørensen *et al.*, 2005; Boitard *et al.*, 2016b). There is some evidence of this

explanation; selective sweeps signatures are associated with stature QTLs

(Bouwman *et al.*, 2018), and the study of van den Berg et al. (2020) was more likely

to identify milk QTLs that had a moderate to high minor allele frequency, suggesting

reduced power to detect low-frequency variants that are potential contributors to

polygenic selection. Conversely, the stature meta-analysis by Bouwman et al. (2018)

found significant SNPs that explained up to 13.8% of the variance in stature, which is

similar to that explained by significant SNPs for human height (16%), which is a

classic trait for polygenic selection studies. Hence, there may be sufficient polygenic

SNPs present to test for polygenic selection, but the power will still be reduced due to

the demographic history of Holstein cattle.

Potential solutions to increase power include increasing sample sizes; using

alternative methods; or analysing different kinds of genome data to detect polygenic

selection. Applying SDS to a larger sample size would increase the power to detect

selection acting in the recent past [Figure 1; see also Field et al. (2016)], but overall

power will still be limited by the tip-length of neutral genealogies. Recent

developments in methodology involve directly inferring trees from genome data, and

using these to identify subtle sweep signatures associated with trait variants (Edge &

Coop, 2019; Speidel *et al.*, 2019; Stern *et al.*, 2021). These methods have greater

power to detect weakly-selected mutations that may be segregating for longer than

the tip-length of the population.

Another approach would be to look beyond sequence data and focus on gene

networks [reviewed by Fagny & Austerlitz (2021)]. The recently–proposed 'omnigenic'

model (Boyle *et al.*, 2017; Liu *et al.*, 2019) posits that variation in quantitative traits is

principally affected by a plethora of 'peripheral' genes that indirectly affect them,

rather than a limited set of 'core' genes that directly modify a trait. These numerous

peripheral genes may exert their influence via regulatory effects (e.g., gene

expression changes), but are also expected to be highly pleiotropic. Fully testing the

omnigenic model will require larger datasets and novel experimental designs (Wray

et al., 2018). A recent example is from an experiment with *Drosophila melanogaster*,

where gene knockouts that do not pass a GWAS significance threshold for pupal

length still significantly affect it (Zhang *et al.,* 2021). There is also nascent evidence

that gene regulation may underlie directional polygenic selection. Boitard et al.

(2016a) found that some adaptive signatures of *B. taurus* are located in intergenic

regions; regulatory changes were also proposed to guide polygenic selection in

Arabidopsis (He *et al.*, 2016). Analyses of gene–sets associated with infection

responses or immunity also found evidence for polygenic selection in humans and

primates (Daub *et al.*, 2013, 2017; Svardal *et al.*, 2017). Immunity gene–sets might

be exceptional cases, as they are more likely to contain genes subject to very strong

selection (Castellano *et al.*, 2019). Further investigations using regulatory information

and a broader range of gene–sets could be a promising approach to determine the

impact of polygenic selection.

Materials and Methods

Full methods are available in the Supplementary Text.

Acknowledgements. We would like to thank Simon Boitard for sharing his

results on *B. taurus* demographic inference; Simon Boitard, Jon Slate and two

anonymous reviewers for providing feedback on the manuscript. MH is supported by

a NERC Independent Research Fellowship (NE/R015686/1). NAP, BG and TB are

funded by a synergistic research grant from the Faculty of Science and Technology,

Aarhus University, Denmark. MH and TB also acknowledge financial support from the

European Research Council under the European Union's Seventh Framework

Program (FP7/20072013, ERC Grant 311341). The authors declare no conflicts of

interest.

Author contributions. All authors contributed to the study design. NAP and

BG provided data. MH performed the analyses and wrote the manuscript, with

feedback from NAP, BG and TB.

Data archiving. Raw SDS scores and polarisation information has been

deposited on Dryad (https://doi.org/10.5061/dryad.547d7wm8q). Data analysis and

simulation scripts are available on GitHub

(https://github.com/MattHartfield/CattleSDS).

References

Barghi, N., Hermisson, J. & Schlötterer, C. (2020) Polygenic adaptation: a unifying framework to understand positive selection. *Nat. Rev. Genet.*, **21**, 769–781.

Barton, N.H., Etheridge, A.M. & Véber, A. (2017) The infinitesimal model: Definition, derivation, and implications. *Theor. Popul. Biol.*, **118**, 50–73.

Barton, N.H., Hermisson, J. & Nordborg, M. (2019) Why structure matters. *eLife*, **8**, e45380.

Barton, N.H. & Keightley, P.D. (2002) Understanding quantitative genetic variation. *Nat. Rev. Genet.*, **3**, 11–21.

Berg, I. van den, Xiang, R., Jenko, J., Pausch, H., Boussaha, M., Schrooten, C., *et al.* (2020) Meta-analysis for milk fat and protein percentage using imputed sequence variant genotypes in 94,321 cattle from eight cattle breeds. *Genet. Sel. Evol.*, **52**, 37.

Berg, J.J. & Coop, G. (2014) A Population Genetic Signal of Polygenic Adaptation. *PLoS Genet.*, **10**, e1004412.

Berg, J.J., Harpak, A., Sinnott-Armstrong, N., Joergensen, A.M., Mostafavi, H., Field, Y., *et al.* (2019a) Reduced signal for polygenic adaptation of height in UK Biobank. *eLife*, **8**, e39725.

Berg, J.J., Zhang, X. & Coop, G. (2019b) Polygenic Adaptation has Impacted Multiple Anthropometric Traits. *bioRxiv*, 167551.

Boitard, S., Boussaha, M., Capitan, A., Rocha, D. & Servin, B. (2016a) Uncovering Adaptation from Sequence Data: Lessons from Genome Resequencing of Four Cattle Breeds. *Genetics*, **203**, 433–450.

Boitard, S. & Rocha, D. (2013) Detection of signatures of selective sweeps in the Blonde d'Aquitaine cattle breed. *Anim. Genet.*, **44**, 579–583.

Boitard, S., Rodríguez, W., Jay, F., Mona, S. & Austerlitz, F. (2016b) Inferring

Population Size History from Large Samples of Genome-Wide Molecular Data - An

Approximate Bayesian Computation Approach. *PLoS Genet.*, **12**, e1005877.

Bouwman, A.C., Daetwyler, H.D., Chamberlain, A.J., Ponce, C.H., Sargolzaei, M.,

Schenkel, F.S., *et al.* (2018) Meta-analysis of genome-wide association studies for

cattle stature identifies common genes that regulate body size in mammals. *Nat.*

Genet., **50**, 362–367.

Boyle, E.A., Li, Y.I. & Pritchard, J.K. (2017) An Expanded View of Complex Traits:

From Polygenic to Omnigenic. *Cell*, **169**, 1177–1186.

Buitenhuis, B., Poulsen, N.A., Gebreyesus, G. & Larsen, L.B. (2016) Estimation of

genetic parameters and detection of chromosomal regions affecting the major milk

proteins and their post translational modifications in Danish Holstein and Danish

Jersey cattle. *BMC Genet.*, **17**, 114.

Bycroft, C., Freeman, C., Petkova, D., Band, G., Elliott, L.T., Sharp, K., *et al.* (2018)

The UK Biobank resource with deep phenotyping and genomic data. *Nature*, **562**,

203–209.

Castellano, D., Uricchio, L.H., Munch, K. & Enard, D. (2019) Viruses rule over

adaptation in conserved human proteins. *bioRxiv*, 555060.

Charlesworth, B. & Edwards, A.W.F. (2018) A century of variance. *Significance*, **15**,

20–25.

Chevin, L.-M. (2019) Selective Sweep at a QTL in a Randomly Fluctuating

Environment. *Genetics*, **213**, 987–1005.

Chevin, L.-M. & Hospital, F. (2008) Selective Sweep at a Quantitative Trait Locus in

the Presence of Background Genetic Variation. *Genetics*, **180**, 1645–1660.

Coop, G., Witonsky, D., Di Rienzo, A. & Pritchard, J.K. (2010) Using Environmental

Correlations to Identify Loci Underlying Local Adaptation. *Genetics*, **185**, 1411–1423.

Daub, J.T., Hofer, T., Cutivet, E., Dupanloup, I., Quintana-Murci, L., Robinson-

Rechavi, M., *et al.* (2013) Evidence for Polygenic Adaptation to Pathogens in the

Human Genome. *Mol. Biol. Evol.*, **30**, 1544–1558.

Daub, J.T., Moretti, S., Davydov, I.I., Excoffier, L. & Robinson-Rechavi, M. (2017)

Detection of Pathways Affected by Positive Selection in Primate Lineages Ancestral

to Humans. *Mol. Biol. Evol.*, **34**, 1391–1402.

Duntsch, L., Tomotani, B.M., Villemereuil, P. de, Brekke, P., Lee, K.D., Ewen, J.G., *et*

al. (2020) Polygenic basis for adaptive morphological variation in a threatened

Aotearoa | New Zealand bird, the hihi (*Notiomystis cincta*). *Proc. R. Soc. B.*, **287**,

20200948.

Edge, M.D. & Coop, G. (2019) Reconstructing the History of Polygenic Scores Using

Coalescent Trees. *Genetics*, **211**, 235–262.

Ehrlich, M.A., Wagner, D.N., Oleksiak, M.F. & Crawford, D.L. (2021) Polygenic

Selection within a Single Generation Leads to Subtle Divergence among Ecological

Niches. *Genome Biol. Evol.*, **13**, evaa257.

Exposito-Alonso, M., 500 Genomes Field Experiment Team, Burbano, H.A.,

Bossdorf, O., Nielsen, R. & Weigel, D. (2019) Natural selection on the *Arabidopsis*

thaliana genome in present and future climates. *Nature*, **573**, 126–129.

Exposito-Alonso, M., Vasseur, F., Ding, W., Wang, G., Burbano, H.A. & Weigel, D.

(2018) Genomic basis and evolutionary potential for extreme drought adaptation in

Arabidopsis thaliana. *Nat. Ecol. Evol.*, **2**, 352–358.

Exposito-Alonso, M., Wilton, P. & Nielsen, R. (2020) Non-additive polygenic models

improve predictions of fitness traits in three eukaryote model species. *bioRxiv*,

2020.07.14.194407.

Fagny, M. & Austerlitz, F. (2021) Polygenic Adaptation: Integrating Population

Genetics and Gene Regulatory Networks. *Trends Genet.*, **37**, 631–638.

Field, Y., Boyle, E.A., Telis, N., Gao, Z., Gaulton, K.J., Golan, D., *et al.* (2016)

Detection of human adaptation during the past 2000 years. *Science*, **354**, 760–764.

Fisher, R.A. (1918) The correlation between relatives on the supposition of

Mendelian inheritance. *Trans. R. Soc. Edinb.*, **52**, 399–433.

Fuhrmann, N., Prakash, C. & Kaiser, T.S. (2021) Polygenic adaptation from standing

genetic variation allows rapid ecotype formation. *bioRxiv*, 2021.04.16.440113.

Gazal, S., Finucane, H.K., Furlotte, N.A., Loh, P.-R., Palamara, P.F., Liu, X., *et al.*

(2017) Linkage disequilibrium–dependent architecture of human complex traits

shows action of negative selection. *Nat. Genet.*, **49**, 1421–1427.

Georges, M., Charlier, C. & Hayes, B. (2018) Harnessing genomic information for

livestock improvement. *Nat. Rev. Genet.*

Gramlich, S., Liu, X., Favre, A., Alex Buerkle, C. & Karrenberg, S. (2021) A polygenic

architecture with conditionally neutral effects underlies ecological differentiation in

Silene. *bioRxiv*, 2021.07.06.451304.

Haile-Mariam, M., Nieuwhof, G.J., Beard, K.T., Konstatinov, K.V. & Hayes, B.J.

(2013) Comparison of heritabilities of dairy traits in Australian Holstein-Friesian cattle

from genomic and pedigree data and implications for genomic evaluations:

Implication of genomic heritability for genomic evaluation. *J. Anim. Breed. Genet.*,

130, 20–31.

Harland, C., Coppieters, W., Mullaart, E., Charlier, C. & Georges, M. (2018) *Rate of*

de novo mutation in dairy cattle and potential impact of reproductive technologies.

Proc. World Congr. Genet. Appl. to Livest. Prod.

Harpending, H.C., Batzer, M.A., Gurven, M., Jorde, L.B., Rogers, A.R. & Sherry, S.T.

(1998) Genetic traces of ancient demography. *Proc. Natl. Acad. Sci. USA*, **95**, 1961–

1967.

Hayes, B.J., Bowman, P.J., Chamberlain, A.J. & Goddard, M.E. (2009) Invited review:

Genomic selection in dairy cattle: Progress and challenges. *J. Dairy Sci.*, **92**, 433–

443.

Hayward, L.K. & Sella, G. (2019) Polygenic adaptation after a sudden change in

environment. *bioRxiv*, 792952.

He, F., Arce, A.L., Schmitz, G., Koornneef, M., Novikova, P., Beyer, A., *et al.* (2016)

The Footprint of Polygenic Adaptation on Stress-Responsive Cis-Regulatory

Divergence in the *Arabidopsis* Genus. *Mol. Biol. Evol.*, **33**, 2088–2101.

Hudson, R.R. (1990) Gene Genealogies and the Coalescent Process. In *Oxford*

Surveys in Evolutionary Biology (ed. by Futuyma, D.J. & Antonovics, J.). Oxford Univ.

Press, Oxford, pp. 1–42.

Jain, K. & Stephan, W. (2015) Response of Polygenic Traits Under Stabilizing

Selection and Mutation When Loci Have Unequal Effects. *G3*, **5**, 1065–1074.

Jain, K. & Stephan, W. (2017a) Modes of Rapid Polygenic Adaptation. *Mol. Biol.*

Evol., **34**, 3169–3175.

Jain, K. & Stephan, W. (2017b) Rapid Adaptation of a Polygenic Trait After a Sudden

Environmental Shift. *Genetics*, **206**, 389–406.

Jiménez-Mena, B., Tataru, P., Brøndum, R.F., Sahana, G., Guldbrandtsen, B. &

Bataillon, T. (2016) One size fits all? Direct evidence for the heterogeneity of genetic

drift throughout the genome. *Biol. Lett.*, **12**.

John, S. & Stephan, W. (2020) Important role of genetic drift in rapid polygenic

adaptation. *Ecol. Evol.*, **10**, 1278–1287.

Josephs, E.B., Berg, J.J., Ross-Ibarra, J. & Coop, G. (2019) Detecting Adaptive

Differentiation in Structured Populations with Genomic Data and Common Gardens.

Genetics, **211**, 989–1004.

Kreiner, J.M., Tranel, P.J., Weigel, D., Stinchcombe, J.R. & Wright, S.I. (2020) The

genetic architecture and genomic context of glyphosate resistance in *Amaranthus*

tuberculatus. *bioRxiv*, 2020.08.19.257972.

Lande, R. (1983) The response to selection on major and minor mutations affecting a

metrical trait. *Heredity*, **50**, 47–65.

Le Corre, V. & Kremer, A. (2012) The genetic differentiation at quantitative trait loci

under local adaptation. *Mol. Ecol.*, **21**, 1548–1566.

Lemay, D.G., Lynn, D.J., Martin, W.F., Neville, M.C., Casey, T.M., Rincon, G., *et al.*

(2009) The bovine lactation genome: insights into the evolution of mammalian milk.

Genome Biol., **10**, R43.

Liu, X., Li, Y.I. & Pritchard, J.K. (2019) Trans Effects on Gene Expression Can Drive

Omnigenic Inheritance. *Cell*, **177**, 1022-1034.e6.

MacEachern, S., Hayes, B., McEwan, J. & Goddard, M. (2009) An examination of

positive selection and changing effective population size in Angus and Holstein cattle

populations (*Bos taurus*) using a high density SNP genotyping platform and the

contribution of ancient polymorphism to genomic diversity in Domestic cattle. *BMC*

Genom., **10**, 181.

MacLachlan, I.R., McDonald, T.K., Lind, B.M., Rieseberg, L.H., Yeaman, S. & Aitken,

S.N. (2021) Genome-wide shifts in climate-related variation underpin responses to

selective breeding in a widespread conifer. *Proc. Natl. Acad. Sci. USA*, **118**,

e2016900118.

Meuwissen, T., Hayes, B. & Goddard, M. (2013) Accelerating Improvement of

Livestock with Genomic Selection. *Annu. Rev. Anim. Biosci.*, **1**, 221–237.

Nielsen, R. (2005) Molecular Signals of Natural Selection. *Annu. Rev. Genet.*, **39**,

197–218.

Novembre, J. & Barton, N.H. (2018) Tread Lightly Interpreting Polygenic Tests of

Selection. *Genetics*, **208**, 1351–1355.

Pavlidis, P., Metzler, D. & Stephan, W. (2012) Selective Sweeps in Multilocus Models

of Quantitative Traits. *Genetics*, **192**, 225–239.

Pritchard, J.K. & Di Rienzo, A. (2010) Adaptation - not by sweeps alone. *Nat. Rev.*

Genet., **11**, 665–667.

Pritchard, J.K., Pickrell, J.K. & Coop, G. (2010) The Genetics of Human Adaptation:

Hard Sweeps, Soft Sweeps, and Polygenic Adaptation. *Curr. Biol.*, **20**, R208–R215.

Qanbari, S., Pausch, H., Jansen, S., Somel, M., Strom, T.M., Fries, R., *et al.* (2014)

Classic Selective Sweeps Revealed by Massive Sequencing in Cattle. *PLoS Genet.*,

10, e1004148.

Qanbari, S., Pimentel, E.C.G., Tetens, J., Thaller, G., Lichtner, P., Sharifi, A.R., *et al.*

(2010) A genome-wide scan for signatures of recent selection in Holstein cattle.

Anim. Genet.

Racimo, F., Berg, J.J. & Pickrell, J.K. (2018) Detecting Polygenic Adaptation in

Admixture Graphs. *Genetics*, **208**, 1565–1584.

Robinson, M.R., Hemani, G., Medina-Gomez, C., Mezzavilla, M., Esko, T.,

Shakhbazov, K., *et al.* (2015) Population genetic differentiation of height and body

mass index across Europe. *Nat. Genet.*, **47**, 1357.

Rowan, T.N., Durbin, H.J., Seabury, C.M., Schnabel, R.D. & Decker, J.E. (2021)

Powerful detection of polygenic selection and evidence of environmental adaptation

in US beef cattle. *PLoS Genet.*, **17**, e1009652.

Sanjak, J.S., Sidorenko, J., Robinson, M.R., Thornton, K.R. & Visscher, P.M. (2018)

Evidence of directional and stabilizing selection in contemporary humans. *Proc. Natl.*

Acad. Sci. USA, **115**, 151–156.

Savolainen, O., Lascoux, M. & Merila, J. (2013) Ecological genomics of local

adaptation. *Nat. Rev. Genet.*, **14**, 807–820.

Schoech, A.P., Jordan, D.M., Loh, P.-R., Gazal, S., O'Connor, L.J., Balick, D.J., *et al.*

(2019) Quantification of frequency-dependent genetic architectures in 25 UK Biobank

traits reveals action of negative selection. *Nat. Commun.*, **10**, 790.

Sella, G. & Barton, N.H. (2019) Thinking About the Evolution of Complex Traits in the

Era of Genome-Wide Association Studies. *Annu. Rev. Genom. Hum. Genet.*, **20**,

461–493.

Sohail, M., Maier, R.M., Ganna, A., Bloemendal, A., Martin, A.R., Turchin, M.C., *et al.*

(2019) Polygenic adaptation on height is overestimated due to uncorrected

stratification in genome-wide association studies. *eLife*, **8**, e39702.

Sørensen, A.C., Sørensen, M.K. & Berg, P. (2005) Inbreeding in Danish Dairy Cattle

Breeds. *J. Dairy Sci.*, **88**, 1865–1872.

Soyeurt, H., Gillon, A., Vanderick, S., Mayeres, P., Bertozzi, C. & Gengler, N. (2007)

Estimation of Heritability and Genetic Correlations for the Major Fatty Acids in Bovine

Milk. *J. Dairy Sci.*, **90**, 4435–4442.

Speidel, L., Forest, M., Shi, S. & Myers, S.R. (2019) A method for genome-wide

genealogy estimation for thousands of samples. *Nat Genet.*, **51**, 1321–1329.

Stephan, W. (2016) Signatures of positive selection: from selective sweeps at

individual loci to subtle allele frequency changes in polygenic adaptation. *Mol. Ecol.*,

25, 79–88.

Stephan, W. (2019) Selective Sweeps. *Genetics*, **211**, 5–13.

Stern, A.J., Speidel, L., Zaitlen, N.A. & Nielsen, R. (2021) Disentangling selection on

genetically correlated polygenic traits via whole-genome genealogies. *Am. J. Hum.*

Genet., **108**, 219–239.

Stetter, M.G., Thornton, K. & Ross-Ibarra, J. (2018) Genetic architecture and

selective sweeps after polygenic adaptation to distant trait optima. *PLoS Genet.*, **14**,

e1007794.

Svardal, H., Jasinska, A.J., Apetrei, C., Coppola, G., Huang, Y., Schmitt, C.A., *et al.*

(2017) Ancient hybridization and strong adaptation to viruses across African vervet

monkey populations. *Nat. Genet.*, **49**, 1705–1713.

Thornton, K.R. (2019) Polygenic Adaptation to an Environmental Shift: Temporal

Dynamics of Variation Under Gaussian Stabilizing Selection and Additive Effects on a

Single Trait. *Genetics*, **213**, 1513–1530.

Turchin, M.C., Chiang, C.W., Palmer, C.D., Sankararaman, S., Reich, D., Genetic

Investigation of ANthropometric Traits (GIANT) Consortium, *et al.* (2012) Evidence of

widespread selection on standing variation in Europe at height-associated SNPs. *Nat*

Genet., **44**, 1015–1019.

Uricchio, L.H., Kitano, H.C., Gusev, A. & Zaitlen, N.A. (2019) An evolutionary

compass for detecting signals of polygenic selection and mutational bias. *Evol. Lett.*,

3, 69–79.

Visscher, P.M. & Goddard, M.E. (2019) From R.A. Fisher's 1918 Paper to GWAS a

Century Later. *Genetics*, **211**, 1125–1130.

Vitti, J.J., Grossman, S.R. & Sabeti, P.C. (2013) Detecting Natural Selection in

Genomic Data. *Annu. Rev. Genet.*, **47**, 97–120.

Vladar, H.P. de & Barton, N. (2014) Stability and Response of Polygenic Traits to

Stabilizing Selection and Mutation. *Genetics*, **197**, 749–767.

Watterson, G.A. (1975) On the number of segregating sites in genetical models

without recombination. *Theor. Popul. Biol.*, **7**, 256–276.

Whitlock, M.C. (2008) Evolutionary inference from F_{ST}. *Mol. Ecol.*, **17**, 1885–1896.

Wieters, B., Steige, K.A., He, F., Koch, E.M., Ramos-Onsins, S.E., Gu, H., *et al.*

(2021) Polygenic adaptation of rosette growth in Arabidopsis thaliana. *PLoS Genet.*,

17, e1008748.

Wollstein, A. & Stephan, W. (2014) Adaptive Fixation in Two-Locus Models of

Stabilizing Selection and Genetic Drift. *Genetics*, **198**, 685–697.

Wray, N.R., Kemper, K.E., Hayes, B.J., Goddard, M.E. & Visscher, P.M. (2019)

Complex Trait Prediction from Genome Data: Contrasting EBV in Livestock to PRS in

Humans. *Genetics*, **211**, 1131–1141.

Wray, N.R., Wijmenga, C., Sullivan, P.F., Yang, J. & Visscher, P.M. (2018) Common

Disease Is More Complex Than Implied by the Core Gene Omnigenic Model. *Cell*,

173, 1573–1580.

Yang, J., Bakshi, A., Zhu, Z., Hemani, G., Vinkhuyzen, A.A.E., Lee, S.H., *et al.* (2015)

Genetic variance estimation with imputed variants finds negligible missing heritability

for human height and body mass index. *Nat. Genet.*, **47**, 1114–1120.

Yeaman, S., Hodgins, K.A., Lotterhos, K.E., Suren, H., Nadeau, S., Degner, J.C., *et*

al. (2016) Convergent local adaptation to climate in distantly related conifers.

Science, **353**, 1431.

Zan, Y. & Carlborg, Ö. (2018) A Polygenic Genetic Architecture of Flowering Time in

the Worldwide Arabidopsis thaliana Population. *Mol. Biol. Evol.*, **36**, 141–154.

Zeder, M.A. (2008) Domestication and early agriculture in the Mediterranean Basin:

Origins, diffusion, and impact. *Proc. Natl. Acad. Sci. USA*, **105**, 11597–11604.

Zeng, J., Vlaming, R. de, Wu, Y., Robinson, M.R., Lloyd-Jones, L.R., Yengo, L., *et al.*

(2018) Signatures of negative selection in the genetic architecture of human complex

traits. *Nat. Genet.*, **50**, 746–753.

Zeng, J., Xue, A., Jiang, L., Lloyd-Jones, L.R., Wu, Y., Wang, H., *et al.* (2021)

Widespread signatures of natural selection across human complex traits and

functional genomic categories. *Nat. Commun.*, **12**, 1164.

Zhang, W., Reeves, G.R. & Tautz, D. (2021) Testing Implications of the Omnigenic

Model for the Genetic Analysis of Loci Identified through Genome-wide Association.

Curr. Biol., **31**, 1092-1098.e6.

Zhao, F., McParland, S., Kearney, F., Du, L. & Berry, D.P. (2015) Detection of

selection signatures in dairy and beef cattle using high-density genomic information.

Genet. Sel. Evol., **47**, 49.

Zhou, X., Carbonetto, P. & Stephens, M. (2013) Polygenic Modeling with Bayesian

Sparse Linear Mixed Models. *PLoS Genet.*, **9**, e1003264.

Figure 1: Simulated mean tip age for *B. taurus*, as a function of the number of haploid

samples. Simulations assumed either (a) demography as inferred by Boitard et al.

(2016b) (the 'High N_0' model), or (b) the same but with a smaller present–day N_e of

49 (the 'Low N_0' model). Points are the mean values; bars show 95% confidence

intervals. The solid line is the best linear fit to the log of both values; dotted lines

show the predicted tip age for 204 alleles.

715
Figure 2: *P*–values of sSDS scores across *B. taurus* autosomes, as plotted on a

negative Log_{10} scale, as a function of the chromosome. Alternating black and grey

points show (non–significant) values from different chromosomes. Blue points are

SNPs with *FDR* < 0.05, with the cut-off denoted by a horizontal dotted line. Red

points are SNPs with Bonferroni–corrected *P*–value < 0.05 (actual *P*–value < ~2.7 x

10^{-8}), with the cut-off denoted by a horizontal dashed line. Figure S1 shows results

722   for the Low N_0 model.

724 Figure 3: Relationship between tSDS scores near milk or stature QTLs, as noted in

725 the subheadings, and the absolute log *P*-value of QTLs. Lines show a linear model

726 regression fit. Figure S3 shows results assuming a low N_0 model.

729   Figure 4: Mean distribution of singleton numbers per individual for each simulation,

730   wither assuming a standard mutation rate (left) or a 10-fold higher mutation rate

731   (right). Bars represent 95% confidence intervals.

Figure 5: Distribution of simulated sSDS scores assuming a high mutation rate, for

the neutral and selected cases. Numbers above each box plot denotes how many

simulations produced SDS scores and were included in the plot.

Chromosome	Gene Name	Start Position	End Position	Gene Biotype	High, Low N_0
1	PPM1L	106405113	106727070	Protein Coding	High, Low
5	TMCC3	24306913	24595494	Protein Coding	High, Low
5	CEP83	24070404	24345243	Protein Coding	High, Low
17	U6	43381106	43381209	snRNA	Low
17	CTSO	43364999	43381605	Protein Coding	Low
17	TDO2	43386894	43403747	Protein Coding	High, Low
23	OR12D2H	29291787	29292713	Protein Coding	High, Low
23	OR12D2E	29305933	29309785	Protein Coding	High, Low
24	GAREM1	24694637	24927333	Protein Coding	High, Low
29	NTM	34576918	34994005	Protein Coding	High, Low

741   Table 1: Genes that overlap or lie close to Bonferroni–significant sSDS regions. The

742   'High, Low N_0' column specifies which genes are close to significant SNPs for each

743   N_0 model.