Using singleton densities to detect recent selection in *Bos taurus*

Matthew Hartfield[1,2], Nina Aagaard Poulsen[3], Bernt Guldbrandtsen[4,5,6], Thomas Bataillon[1]

1) Bioinformatics Research Centre, Aarhus University, DK–8000 Aarhus, Denmark.

2) Institute of Evolutionary Biology, University of Edinburgh, Edinburgh EH9 3FL, UK.

3) Department of Food Science, Aarhus University, Agro Food Park 48, 8200 Aarhus N, Denmark.

4) Center for Quantitative Genetics and Genomics, Department of Molecular Biology and Genetics, Aarhus University, DK–8830 Tjele, Denmark.

5) Rheinische Friedrich-Wilhelms-Universität Bonn, Institut für Tierwissenschaften, Tierzucht, Endenicher Allee 15, D-53115 Bonn, Germany.

6) Department of Veterinary Sciences, Copenhagen University, DK–1870 Frederiksberg C, Denmark.

Corresponding author: Matthew Hartfield (m.hartfield@ed.ac.uk).

Key words: selection, genomics, *Bos taurus*, milk protein, milk fat, stature.

Abstract: Many quantitative traits are subject to polygenic selection, where several genomic regions undergo small, simultaneous changes in allele frequency that collectively alter a phenotype. The widespread availability of genome data, along with novel statistical techniques, has made it easier to detect these changes. We apply one such method, the 'Singleton Density Score', to the Holstein breed of *Bos taurus* to detect recent selection (arising up to around 740 years ago). We identify several genes as candidates for targets of recent selection, including some relating to cell regulation, catabolic processes, neural-cell adhesion and immunity. We do not find strong evidence that three traits that are important to humans – milk protein content, milk fat content, and stature – have been subject to directional selection. Simulations demonstrate that since *B. taurus* recently experienced a population bottleneck, singletons are depleted so the power of SDS methods are reduced. These results inform on which genes underlie recent genetic change in *B. taurus*, while providing information on how polygenic selection can be best investigated in future studies.

Impact statement: Many traits of ecological or economic importance (including height,

disease propensity, climatic adaptation) are 'polygenic'. That is, they are affected by a

large number of genetic variants, with each one only making a small contribution to a

trait, but collectively influence variation. As selection acts on all of these variants

simultaneously, it only changes the frequency of each one by a small amount, making

it hard to detect such selection from genome data. This situation has changed in recent

years ,with the proliferation of whole–genome data from many individuals, along with

the development of methods to detect the subtle effects of polygenic selection. Here,

we use data from 102 genomes from domesticated cattle (*Bos taurus*) that has

experienced intense artificial selection since domestication, and test if we can detect

signatures of recent selection (arising up to 740 years ago). Domesticated species are

appealing for this kind of study, as they are subject to extensive genome sequencing

studies, and genetic variants can be related to traits under selection. We carried out

our analysis in two parts. We first performed a genome–wide scan to find individual

genetic regions that show signatures of recent selection. We identify some relating to

cell regulation, catabolic processes, neural-cell adhesion and immunity. In the second

part, we then analysed genetic regions associated with three key traits: milk protein

content, milk fat content, and stature. We tested if these regions collectively showed a

signature of selection, but did not find a significant result in either case. Simulations

suggest that the domestication history of cattle affected the power of these methods.

We end with a discussion on how to best detect polygenic selection in future studies.

Introduction

Determining which genomic regions have been subject to selection is a major research goal in evolutionary genetics. Traditional methods have focused on detecting strong selection affecting individual genes (Nielsen, 2005; Vitti *et al.*, 2013; Stephan, 2019). An alternative process is 'polygenic selection', where many loci contribute to genetic variation in a trait, so selection acting on it is expected to generate small and simultaneous allele frequency changes at multiple loci (Pritchard & Di Rienzo, 2010; Pritchard *et al.*, 2010). Many polygenic models have been formulated to account for both the response to phenotypic selection, and the maintenance of genetic variance in quantitative traits [reviewed by (Sella & Barton, 2019; Barghi *et al.*, 2020)]. Among them is Fisher's infinitesimal model, which is important for its historical role in uniting population and quantitative genetics, and its recent renaissance in the context of genome–wide association studies (Fisher, 1918; Barton & Keightley, 2002; Barton *et al.*, 2017; Charlesworth & Edwards, 2018; Visscher & Goddard, 2019). However, whereas it has been possible to identify which genetic regions contribute to trait variation, it has historically been hard to infer which alleles have been involved in the polygenic selection response. Extensive theoretical studies of how alleles at multiple loci act when a population adapts to a new optimum generally find that 'large–effect' alleles, which strongly affect a trait, are the first to spread and fix while 'small–effect' alleles take much longer to reach high frequencies (de Vladar & Barton, 2014; Wollstein & Stephan, 2014; Jain & Stephan, 2015, 2017a, 2017b; Stetter *et al.*, 2018; Thornton, 2019; Hayward & Sella, 2019). Furthermore, if epistasis exists between variants, many selected alleles do not reach fixation as they eventually become deleterious (de Vladar & Barton, 2014; Jain & Stephan, 2017b).

The spread of large–effect alleles may also be impeded if a faster adaptive response

can be otherwise realised through changes at many small–effect alleles (Lande,

1983; Chevin & Hospital, 2008; Pavlidis *et al.*, 2012; Chevin, 2019). Alternatively, if

the optimum shift is sufficiently big, then large-effect mutations that first go to fixation

can subsequently be replaced by small–effect variants over longer timescales (on the

order of the population size; Hayward and Sella (2019)). Overall, only a small

proportion of loci affected by polygenic selection are expected to fix sufficiently

quickly to leave selection signatures in genomic data (Pavlidis *et al.*, 2012; Thornton,

2019).

Due to this difficulty, earlier methods for detecting polygenic selection focused

on cases where selection favours distinct phenotypes in different populations, so trait

differentiation amongst populations will be greater than expected under neutral drift.

Tests for this form of selection relied on comparing Q_{st} and F_{st} statistics, which

respectively measured mean genetic differentiation at the trait itself and a set of

neutral loci (Whitlock, 2008; Le Corre & Kremer, 2012; Savolainen *et al.*, 2013). Yet

these methods do not determine which genomic regions are subject to selection. This

situation has now changed with the increased number of genome–wide association

study (GWAS) data that link genotypes and phenotypes, as exemplified by the

development of large cohort studies [e.g., the UK Biobank; Bycroft et al. (2018)]. The

release of these data spurred a series of studies and new methods designed

specifically to detect polygenic selection. These methods usually involve determining

which SNPs affecting a phenotype show correlated changes in frequency (Berg &

Coop, 2014; Racimo *et al.*, 2018; Sanjak *et al.*, 2018; Josephs *et al.*, 2019; Berg *et*

al., 2019a, 2019b; Uricchio *et al.*, 2019; Edge & Coop, 2019; Kreiner *et al.*, 2020;

Wieters *et al.*, 2021; Gramlich *et al.*, 2021); which sets of alleles are associated with

certain environmental or climatic variations (Coop *et al.*, 2010; Turchin *et al.*, 2012;

Robinson *et al.*, 2015; Yeaman *et al.,* 2016; Exposito-Alonso *et al.*, 2018; Zan &

Carlborg, 2018; Exposito-Alonso *et al.*, 2019; MacLachlan *et al.*, 2021; Ehrlich *et al.*,

2021; Fuhrmann *et al.*, 2021; Rowan *et al.*, 2021); or determining which SNPs or

genetic regions explain a large fraction of phenotypic variance and trait heritability

(Zhou *et al.*, 2013; Yang *et al.*, 2015; Gazal *et al.*, 2017; Zeng *et al.*, 2018; Schoech

et al., 2019; Exposito-Alonso *et al.*, 2020; Duntsch *et al.*, 2020; Zeng *et al.*, 2021).

Some of these approaches use overlapping methods.

Detecting recent polygenic selection is much harder, as long periods of time

(number of generations on the order of the population size; Hayward and Sella, 2019;

Thornton, 2019) may be needed to cause detectable frequency changes in alleles

with small effect sizes. Over shorter timescales, these frequency changes are

expected to be more modest and harder to detect (Stephan, 2016; Jain & Stephan,

2017a). A recent breakthrough in detecting these subtle changes was the

development of the 'Singleton Density Score' (SDS), a statistic tailored to detect

recent and coordinated allele frequency changes over many SNPs (Field *et al.*,

2016). Recent selection at a locus favouring one variant will lead to a reduction in the

number of singletons (i.e., variants that are only observed once) around it. The SDS

detects regions that exhibit a reduction in the density of singletons, to determine

candidate regions that have been subject to recent selection. Using this approach,

Field et al. (2016) found correlations between SDS scores at SNPs and their

associated GWAS effect sizes for several polygenic traits in the modern UK human

population, including increased height, infant head circumference and fasting insulin.

Their findings suggested that these traits have been subject to recent selection

during the most recent 75 or so generations (about 2,000 years). However, these

(and other) results that detect selection for increased height may instead reflect

previously unaccounted–for population structure (Novembre and Barton, 2018;

Barton et al., 2019; Sohail et al., 2019; Berg et al., 2019; Uricchio et al., 2019; Edge

and Coop, 2019).

The SDS method is ideally suited to organisms where large amount of whole-

genome data are available, along with QTL or GWAS information that link genotypes

to phenotypes, Domesticated species are attractive systems for studying recent

selection, as selected phenotypes are often already known and these species are

subject to large–scale sequencing studies. Investigating the genetic architecture

underlying rapid selection in these species is also important to determine how they

respond to agricultural practices, and uncover selection targets that can be used to

improve breeding programs (Georges *et al.*, 2018). Domestic cattle *Bos taurus* has

been subject to intensive genomics analyses to improve artificial selection for traits

that are important for human use, including milk protein yield, milk fat content, and

stature (Hayes *et al.*, 2009; Meuwissen *et al.*, 2013; Wray *et al.*, 2019). These traits

are influenced in part by an individual's genome, with significant heritability estimates

being recorded, some as high as 80% (Soyeurt *et al.*, 2007; Haile-Mariam *et al.*,

2013; Buitenhuis *et al.*, 2016). Previous selection scans on *B. taurus* reported

individual regions that were likely to be subject to recent selection, some of which

were close to genetic regions for stature and milk protein content (Lemay *et al.*, 2009;

MacEachern *et al.*, 2009; Qanbari *et al.*, 2010; Boitard & Rocha, 2013; Qanbari *et al.*,

2014; Zhao *et al.*, 2015; Boitard *et al.*, 2016a; Bouwman *et al.*, 2018). Recently,

Rowan *et al.* (2021) introduced a method to detect polygenic environmental

adaptation. However, stature and milk protein content are polygenic traits, with

several genetic regions and QTLs associated with each (Lemay *et al.*, 2009; Boitard

et al., 2016a; Bouwman *et al.*, 2018; van den Berg *et al.*, 2020). There has also yet to

be a formal test of whether these traits show evidence of polygenic selection.

Here, we applied the SDS method to whole–autosome sequencing data from

102 *B. taurus* Holstein individuals. We first determined genetic regions that have

been subject to recent directional selection, and subsequently tested whether

evidence exists for recent selection acting on a set of QTLs underlying either milk

protein content, milk fat content, or stature in this breed.

Results

Methods outline

We filtered the data to retain only bi–allelic SNPs that had a sensible level of

coverage and did not lie in putatively over–assembled regions (i.e., duplicated

sections that caused many reads to assemble at a specific genetic location). Over–

assembled regions appear as highly heterozygous with elevated coverage, and can

exhibit false signatures of recent selection. We also obtained a set of singletons and

filtered them to retain high-quality variants where both alleles were equally well

covered to remove potentially erroneous calls. We polarised test SNPs using

outgroup sequences and applied the SDS test of Field et al. (2016) to detect recent

selection, with increased SDS values reflecting selection favouring derived SNPs

over ancestral variants. We standardised SDS scores with those of a similar

frequency, so they are normally distributed [similar normalisation was also carried out

by Field et al. (2016)]. These values are denoted sSDS for 'standardised SDS'.

Further details are available in the *Methods* section.

Estimating timescale of selection

We first determined the timescale over which we expect to detect selection in

B. taurus using the SDS method. SDS measures the changes in singleton numbers

around putatively selected SNPs, relative to background numbers in the absence of

selection. As singletons arise on the tips of the underlying gene trees, the average tip

length in the genealogy of sequenced samples determines the timescale over which

the SDS detects a signal (Field et al., 2016). As more haploid genomes are included

in the study, the time to first coalescence between two samples decreases, reducing

the tip lengths and therefore shortening the timescale over which SDS detects

selection (Field *et al.,* 2016). We hence simulate tip-ages over a range of sample

sizes to investigate how this timescale changes accordingly.

To calculate the mean tip age, we simulated gene genealogies under two

scenarios. We first simulated the Holstein population demography inferred by Boitard

et al. (2016b), which suggested that this population experienced a sudden decline in

effective population size (N_e) since domestication, but with a present–day N_e (~793)

that is much larger than that inferred from pedigree data [~49; Sørensen et al.

(2005)] or from temporal variation in SNP frequencies (~48; Jiménez–Mena et al.

2016). Hence, we also simulated genealogies under a second model that used the

Boitard et al. (2016b) demographic model, but with the present–day N_e set to 49.

These scenarios will be referred to as the 'High N_0' and 'Low N_0' models,

respectively.

Figure 1 shows simulation results. Depending on the assumed present–day

N_e, the tip length in our sample of 204 alleles (i.e., assuming two per diploid

individual) goes back either 65 or 148 generations. Assuming 5 years per generation

(Boitard *et al.*, 2016b), this timescale corresponds to between 325 and 740 years

ago. Since *B. taurus* domestication started around 10,000 years ago (Zeder, 2008)

the sample size used in this study will only capture selection acting in the very recent

past that is more relevant for breed formation, rather than selection during *B. taurus*

domestication. Sample sizes and tip-ages are linearly related on a log-log scale,

meaning that an increase in sample size will greatly decrease the timescale over

which SDS detects selection. For example, with 500 haplotypes then SDS will detect

selection acting no more than 50 generations ago, depending on the underlying

demographic model.

We will focus on detecting selection signatures assuming the high N_0 model.

Results using the low N_0 model to calibrate scores were broadly similar. They are

outlined in the Supplementary Text; we will highlight when differences arise.

Genome–wide sSDS

Figure 2 plots sSDS values (at SNPs with minor allele frequency greater than

5%) across all autosomes, excluding chromosome 25 (due to an insufficient number

of singletons needed to obtain SDS scores after filtering). Many SNPs have elevated

sSDS scores (158 SNPs at *FDR* < 0.05; 306 for the low N_0 model). Several regions

contain SNPs with significantly high sSDS values (Bonferroni–corrected nominal *P* <

0.05; actual *P* < ~2.7 x 10^{-8}). To further investigate potential selection targets, we

looked for genes that either overlapped significant SNPs or lay 10kb up– or

downstream of them. Linkage disequilibrium (LD), as measured by r^2, decays to

around 0.2 over 50kb in Danish Holstein breeds (Buitenhuis *et al.*, 2016), so genes

within 10kb should be in LD with regions harbouring high sSDS scores. Table 1 lists

these genes, with more targets present under the low N_0 model. Most of these genes

are of unknown function (as listed on UniProt); the list also includes an snRNA.

PPM1L is involved with cellular regulation and the activation of stress–activated

protein kinases. *TDO2* is involved in tryptophan–related catabolic processes, while

NTM is implicated in neural cell adhesion. SNPs with significantly elevated scores

are also found on chromosome 23 near the MHC region, which may reflect over–

dominant selection. All Bonferroni–significant SNPs were removed from subsequent

tests of recent polygenic selection to prevent directional selection from skewing the

underlying sSDS distributions. Figure S1 shows results for the low N_0 model.

Testing for polygenic selection acting on milk protein and stature

If polygenic selection were acting on specific traits, we expect a positive

correlation between the effect size of variant underpinning it, and selection acting on

it as measured by sSDS. We collated sSDS scores of SNPs that lie close to QTLs

reported for either milk fat percentage, milk protein percentage (van den Berg *et al.*,

2020), or those that lie close to stature QTLs (Bouwman *et al.*, 2018). The latter were

inferred from a meta–analysis of GWAS studies conducted in seven Holstein

populations, but not every QTL had an effect size reported in each population. We

hence investigated two overlapping consensus QTL sets, where an effect size was

either reported in at least 6 of 7 populations (yielding 42 QTLs with sSDS scores

associated with them), or where effect sizes were reported in at least 5 of 7

populations (58 QTLs had sSDS scores). We re-polarised SDS scores so that a

positive score reflected a trait-increasing effect; we denote these values 'tSDS'

following Field *et al.* (2016). We then determine if there was a positive correlation

between the absolute $\log_{10}$-value of the QTL *P*-value (a proxy for the effect size) and

tSDS.

Figure 3 shows the relationship between QTL *P*-values and tSDS for SNPs

that lie close to QTLs. Although positive trends are observed, they all exhibit non-

significant regression slopes (milk fat percentage $P = 0.596$; milk protein percentage

$P = 0.873$; stature from 6 breeds $P = 0.558$; stature from 5 breeds $P = 0.344$).

Relationships remain non-significant after removing an outlier point for the milk traits

whose QTL has an extremely low *P*-value (Figure S2), and also under the low N_0

model (Figure S3; see figure legends for regression *P*-values).

sSDS (and tSDS) can become correlated along the genome if focal SNPs are

in LD with one another, which was not accounted for in the preceding analyses. To

determine whether LD could have impacted upon these correlations, we randomly

subsampled sSDS scores from SNPs that shared the same chromosome and bin of

derived-allele frequency as the SNPs used in the above analyses, and re-polarised

them to transform them into tSDS values. We then determined the *F* statistic

associated with the linear model fit for these permuted values to determine whether

that for the true data was significantly elevated (see Methods for details). In all cases,

the observed value was not significantly higher than for permuted values (see Figure

S4 for histograms and exact *P*–values, which all exceed 0.05). We therefore

conclude that these QTL datasets do not harbour SNPs with significantly different

tSDS scores compared to the rest of the genome.

Discussion

Summary of results

We analysed an extensive *B. taurus* genomic dataset to identify signatures of

recent selection in the Holstein breed, and to determine whether the data contained a

signal of polygenic selection acting on milk proteins and QTLs underlying phenotypic

variation in stature. Given the sample size and the demographic history of Holsteins,

the SDS method can detect very recent selection events arising no more than

approximately 740 years ago (Figure 1). A whole–genome scan for sSDS scores

identified several targets of recent directional selection that overlap or lie close to

protein–coding genes (Figure 2; Table 1). The genes whose functions are known are

involved in protein regulation, catabolic processes, and neural-cell adhesion.

Significant values were also observed near the MHC region. We subsequently

investigated whether either milk protein genes or SNPs near stature QTLs

collectively showed evidence of polygenic selection. We did so by testing whether

there is a relationship between the QTL effect size, as measured by its *P*-value, and

tSDS values to SNPs near them. However, no relationship was observed, even after

performing a permutation test (Figures 3, S2-S4). Hence, while sSDS could reveal

specific instances of recent selection, tests based on collective scores of variants

associated with known selected traits yielded no signal of polygenic selection.

Potential reasons for a lack of polygenic selection signal

Impact of Holstein demographic history

While the SDS method detected individual candidate genes for very recent

selection, we were unable to find strong evidence for polygenic selection acting on

three traits that were subject to artificial selection since domestication. This result is *a*

priori surprising, given that these traits have been subject to recent intense artificial

selection. Recent studies generally find non-zero heritability estimates for them,

indicating that there should be the potential for genetic variants underpinning them to

change in response to artificial selection (Soyeurt *et al.*, 2007; Haile-Mariam *et al.*,

2013; Buitenhuis *et al.*, 2016). In addition, the ratio of the mutation and recombination

rates in cattle is just over three (Boitard *et al.*, 2016b; Harland *et al.*, 2018), indicating

that several informative SNPs exist per haplotypes that should improve the power of

the SDS method [in contrast, this ratio is approximately equal to one in humans

(Field *et al.*, 2016)].

One potential reason for this lack of signal is due to the population history of

Bos taurus. The effective population size of many *B. taurus* breeds appears to have

undergone a decline since domestication (Sørensen et al., 2005; Boitard et al.,

2016b), which likely reflects successive bottlenecks due to domestication, breed

formation and intense recent selection. Population size reductions are known to

reduce the number of low-frequency variants and increases the prevalence of

intermediate-frequency variants (Harpending *et al.*, 1998), which can affect the power

of the SDS method. To understand if the history of *B. taurus* affects the detection of

recent selection in Holstein cattle using SDS, we ran coalescent simulations to

determine its ability to detect ongoing selection, given realistic Holstein population

history and genetic parameters (see Methods for details). We simulated a partial

sweep occurring in the middle of a 10Mb region, either assuming a mutation rate in

line with what has been inferred for Holstein, or one 10-fold higher to replicate

diversity expected in a genetic region with an elevated mutation rate.

For the standard mutation rate, no SDS scores were produced for any

simulations. After inspecting the simulation results, we see that there is a large skew

in the distribution of singleton numbers per individual with a large number of

individuals (over 20 on average) that do not carry singletons at the end of

simulations, preventing the calculation of a local SDS score (Figure 4). This fraction

remained the same irrespective of whether the simulated region was neutral or

subject to selection; the main effect of a sweep was to reduce the mean number of

singletons per individual, which is the signal measured by SDS (Field *et al.*, 2016).

This reduction in overall singleton numbers is consistent with the known effects of

population size contraction on reducing tip lengths (Harpending *et al.*, 1998).

With a 10-fold higher mutation rate, there were fewer cases where no

individual harboured singletons (Figure 4). Accordingly, SDS scores could be

calculated for 65 and 66 out of 100 simulations for the neutral and selective cases

respectively. In these cases, sSDS values were significantly higher in the selected

case than for the neutral case (Figure 5; two-sided Wilcox Test $P = 1.1 \times 10^{-5}$).

However, note that sSDS values is less than one for the selected case, which does

not exceed the FDR threshold in our study (for the high N_0 case, the smallest sSDS

value with FDR < 0.05 is 4.46).

Overall, these simulations are consistent with population size reductions in *B.*

taurus reducing the number of singletons, which limits its ability to detect partial

sweeps. SDS is more likely to detect signals in regions of elevated mutation rate,

suggesting there is likely an ascertainment bias in where signals were detected;

variants contributing to polygenic selection are more likely to be detected by SDS in

regions of elevated mutation. The reduction in singletons does not just affect the

ability of SDS to detect sweeps overall, it also reduces the power to investigate SDS

values in telomeric regions. SDS values are calculated using the distance up- and

downstream from a SNP to the nearest singleton, and are undefined if a certain

number of samples do not harbour singletons in either direction (Field *et al.*, 2016).

SDS values are hence less likely to be defined in telomeric regions, as it is generally

less feasible to observe singletons up until the end of the chromosome. This problem

is exacerbated if there are few singletons overall.

Other potential reasons for a lack of signal

Another potential reason for a lack of signal is that the selection response on

these traits may have been driven by large-effect variants that have already fixed in

the population, with a smaller contribution from small-effect mutations. Theoretical

models have shown that more major–effect QTLs are likely to fix if the population lies

further from a fitness optimum (Lande, 1983; Jain & Stephan, 2017b; Thornton,

2019). Domesticated species, which experience strong and sustained directional

artificial selection, especially in recent generations, could thereby fix more adaptive

mutation via sweep–like processes compared to populations evolving in more stable

environments (Lande, 1983; Jain & Stephan, 2017a). Furthermore, once a population

has adapted to a new environment (the domestication phenotype in this case), then

any remaining major–effect mutations are likely to be superseded by variants with

weaker effects, which are harder to detect (Hayward & Sella, 2019). The response to

polygenic selection will be further weakened in smaller populations (John & Stephan,

2020), which could be a factor given the reduced effective population sizes of *B.*

taurus (Sørensen *et al.*, 2005; Boitard *et al.*, 2016b). There is some evidence of this

explanation; selective sweeps signatures are associated with stature QTLs

(Bouwman *et al.*, 2018), and the study of van den Berg et al. (2020) was more likely

to identify milk QTLs that had a moderate to high minor allele frequency, suggesting

reduced power to detect low-frequency variants that are potential contributors to

polygenic selection. Conversely, the stature meta-analysis by Bouwman et al. (2018)

found significant SNPs that explained up to 13.8% of the variance in stature, which is

similar to that explained by significant SNPs for human height (16%), which is a

classic trait for polygenic selection studies. Hence, there may be sufficient polygenic

SNPs present to test for polygenic selection, but the power will still be reduced due to

the demographic history of Holstein cattle.

Potential solutions to increase power include increasing sample sizes; using

alternative methods; or analysing different kinds of genome data to detect polygenic

selection. Applying SDS to a larger sample size would increase the power to detect

selection acting in the recent past [Figure 1; see also Field et al. (2016)], but overall

power will still be limited by the tip-length of neutral genealogies. Recent

developments in methodology involve directly inferring trees from genome data, and

using these to identify subtle sweep signatures associated with trait variants (Edge &

Coop, 2019; Speidel *et al.*, 2019; Stern *et al.*, 2021). These methods have greater

power to detect weakly-selected mutations that may be segregating for longer than

the tip-length of the population.

Another approach would be to look beyond sequence data and focus on gene

networks [reviewed by Fagny & Austerlitz (2021)]. The recently–proposed 'omnigenic'

model (Boyle *et al.*, 2017; Liu *et al.*, 2019) posits that variation in quantitative traits is

principally affected by a plethora of 'peripheral' genes that indirectly affect them,

rather than a limited set of 'core' genes that directly modify a trait. These numerous

peripheral genes may exert their influence via regulatory effects (e.g., gene

expression changes), but are also expected to be highly pleiotropic. Fully testing the

omnigenic model will require larger datasets and novel experimental designs (Wray

et al., 2018). A recent example is from an experiment with *Drosophila melanogaster*,

where gene knockouts that do not pass a GWAS significance threshold for pupal

length still significantly affect it (Zhang *et al.*, 2021). There is also nascent evidence

that gene regulation may underlie directional polygenic selection. Boitard et al.

(2016a) found that some adaptive signatures of *B. taurus* are located in intergenic

regions; regulatory changes were also proposed to guide polygenic selection in

Arabidopsis (He *et al.*, 2016). Analyses of gene–sets associated with infection

responses or immunity also found evidence for polygenic selection in humans and

primates (Daub *et al.*, 2013, 2017; Svardal *et al.*, 2017). Immunity gene–sets might

be exceptional cases, as they are more likely to contain genes subject to very strong

selection (Castellano *et al.*, 2019). Further investigations using regulatory information

and a broader range of gene–sets could be a promising approach to determine the

impact of polygenic selection.

Materials and Methods

Simulating Holstein tip ages

Neutral genealogies were simulated using *msprime* (Kelleher *et al.*, 2016) to

determine the mean tip length, and hence the background distribution of SDS in the

absence of selection. We either simulated the Holstein population demography

inferred by Boitard et al. (2016b), rounding estimated population sizes to the nearest

integer, or with the present–day N_e equal to 49 (Sørensen et al., 2005). We refer to

these outputs as the 'High N_0' and 'Low N_0' models. 1,000 simulations were

performed for each number of samples, ranging from 10 to 1,050. The mean tip

length was calculated over all 1,000 simulations; 95% confidence intervals were

calculated from 1,000 bootstraps. We fitted a linear model to the log of mean tip

length against the log number of samples, and used it to predict the average tip age

for 204 alleles, which is the number of diploid haplotypes used in the study. *B. taurus*

are somewhat inbred (Sørensen *et al.*, 2005), which increases within–individual

relatedness, and could reduce the number of unique alleles (see Nordborg and

Donnelly (1997) for an example with self–fertilisation). Estimates of the inbreeding

coefficient *F* (Wright, 1951), which measure the reduction in heterozygosity, range

from –0.15 to 0.35, with a mean of 0.059 (Figure S5; methods outlined below). Given

this low mean value, we assumed two unique alleles per individual.

Genome Data Extraction

Whole genome sequencing for 102 Holstein bulls and cows were done by

Illumina and BGI short-read paired-end sequencing in various laboratories with read

lengths of 100 base pairs and up. No raw reads were shorter than 90 basepairs.

Bulls were selected for sequencing had high genetic representation in the present–

day Holstein population. Sequencing of close relatives was avoided. Individuals were

born between approximately 1980 and 2010. DNA was extracted from tissue, blood,

or semen samples. Read data were processed according to the 1,000 Bull Genomes

Project (Daetwyler *et al.*, 2014). Briefly, data were trimmed and quality filtered using

Trimmomatic version 0.38 (Bolger *et al.*, 2014). Reads were aligned to the ARS-

UCD-1.2 bovine genome assembly (Rosen *et al.*, 2018)

(https://sites.ualberta.ca/~stothard/1000_bull_genomes/ARS-

UCD1.2_Btau5.0.1Y.fa.gz) with the *B. taurus* Y chromosome assembly from BTau-

5.0.1 added. Alignment was performed with bwa version 0.17 (Li & Durbin, 2009).

Samtools (Li *et al.*, 2009) was used for sorting and marking of PCR duplicates. Base

qualities were recalibrated using Genome Analysis Toolkit (GATK; McKenna et al.

[2010]) version 3.8 using a set of known variable sites (Schnabel and Chamberlain,

unpubl). GVCF files were formed using GATK's Haplotype Caller. Genotypes were

called using GATK's GenotypeGVCFs.

Initial filtering

Figure S6 outlines a schematic of the data filtering. We first used *VCFtools*

(Danecek *et al.*, 2011) to obtain a baseline list of biallelic SNPs. We removed indels

and sites where the genotype was unknown in any individual. For each autosome,

we obtained the mean depth for each remaining site using *VCFtools*' '--site-mean-

depth' option. Figure S7 shows the depth distribution for these sites after initial

filtering. We fitted a Poisson distribution to these data that had the same mean (9.76)

as observed in the dataset. We determined the expected coverage range based on

458 the 99.5% quantile range of the fitted distributions, which equalled 2 to 20. We

459 subsequently removed sites that had mean coverage outside this range. We hence

460 retained 6,873,371 of 20,010,175 initial variants (all entries in each autosome VCF

461 file, including indels); we denote this dataset as the 'filtered' dataset.

463 *Finding putatively over–assembled regions*

464   Scaffolds of different genetic segments that each carry highly identical

465 repeated regions can be 'over–assembled', where very similar chromosome regions

466 were anchored to a single location (Chaisson *et al.*, 2015). These over–assembled

467 regions (hereafter OARs) manifest themselves in aligned data as having high levels

468 of heterozygosity, and elevated apparent coverage. If not corrected for, they can be

469 misclassified as selected sites (e.g., subject to partial sweeps or balancing selection).

470 We used a sliding window method to identify putative OARs in the dataset. For each

471 chromosome, in each window we calculated (i) the number of sites where the

472 reference allele had a frequency between 49% – 51%, (ii) the mean heterozygosity

473 for each SNP (defined as the number of heterozygotes among the 102 individuals),

474 and (iii) the mean summed allele depth. We used overlapping windows, each of size

475 500 SNPs with a step size of 10 SNPs. We first analysed all chromosomes to

476 determine the distribution of values produced per window. We then re–ran the

477 analyses, classifying windows as OARs if values for all three statistics belonged to

478 the top 99.5% of their respective distributions, merging overlapping windows. We

479 subtracted 1 from the start position of each region so that the leftmost boundary

480 would also be excluded (if using '--bed-exclude' in *VCFtools*). Figure S8 shows an

481 example where a region at the beginning of chromosome 1 was identified as an

482 OAR. Overall, 5 OARs comprising 5,880 SNPs were identified (Table S1), which

483 were subsequently masked from the rest of the pipeline.

485 *Calculating inbreeding coefficients*

486  Inbreeding coefficients were estimated using the '--het' option of *VCFtools*,

487 which reports *F*-statistics for each chromosome per individual. Individual *F*-values

488 (Figure S5) were calculated by taking the mean over all chromosomes, weighted by

489 the chromosome size.

491 *Obtaining SDS analysis inputs*

492  ***Test SNPs:*** Focal SNPs were those with an alternate allele frequency

493 between 5% and 95%, and where each genotype was observed at least once

494 amongst all samples. We then polarised values to determine the ancestral and

495 derived states. Genome data, mostly from Wu *et al.* (2018), was downloaded from

496 the NCBI SRA (https://www.ncbi.nlm.nih.gov/sra) for Bison (*Bison bison*,

497 SRR6448737, SRR6448738, SRR6448739, SRR6448740), Wisent (*Bos bonasus*,

498 SRR6448670, SRR6448682, SRR6448683, SRR6448684), Gaur (SRR6448732,

499 SRR6448733, SRR6448734, SRR6448735), Gayal (*Bos frontalis*, SRR6448578,

500 SRR6448580, SRR6448581), Banteng (*Bos javanicus*, SRR6448720 and

501 SRR6448721), and as well as data for Yak (*Bos grunniens*, SRR5641601). Ancestral

502 SNPs were determined by counting which allele was present in a majority of these

503 related species; note that some SNPs for some species carried neither the reference

504 nor the alternate alleles. We only retained SNPs where the called ancestral allele

505 matched either the reference or alternate allele in our dataset. 3,405,023 SNPs were

retained for testing.

Singletons: Raw singleton data was extracted from the filtered Holstein

dataset using *VCFtools*' '--singleton' option. This option identified both true singletons

and private doubletons (i.e., where an allele is unique to an individual but present as

a homozygote). Only true singletons were retained for analyses. To test whether a

singleton had the same coverage as the non–singleton allele, we extracted the

sequence depth for both alleles and retained sites satisfying the following criteria.

First, the total allele depth was between 2 and 20 inclusive. Second, either (a) if the

summed depth over both alleles exceeded 5, then the binomial probability of the

observed allele depth exceeded 0.1; or (b) a stricter manual cut–off was applied if the

total allele depth equalled 5 or less. Table S2 outlines the cut–off values used;

554,402 of 765,822 singletons were subsequently retained.

Other parameters: The SDS method requires a 'singleton observability'

probability, to predict how likely it is that a singleton will be detected by genome

sequencing. Following Field et al. (2016) we used the mean depth per individual,

obtained using the '--depth' option in *VCFtools.* It was also necessary to state the

genetic boundaries between which analyses were carried out; we used a starting

point of 1 and end points equal to the reported size of each autosome in *B. taurus*, as

obtained from the ARS–UCD 1.2 genome assembly

(https://www.ncbi.nlm.nih.gov/genome/?term=txid9913[orgn])*.*

Raw SDS values were calculated by fitting a gamma distribution to observed

singleton distances, and comparing it to the expected distribution for the neutral

case. We used the scripts provided by Field et al. (2016)

(https://github.com/yairf/SDS) to generate the expected shape values for the gamma

distribution for both the high and low N_0 models. Finally, we used a value of 10^{-7} to

initiate the search for a maximum value in likelihood space.

Calculating SDS scores and their significance for individual SNPs

Out of 3,405,023 input SNPs, we retained and assigned scores to 1,877,770

of these. SDS scores were not assigned to a SNP if more than 5% of individuals did

not harbour any singletons upstream or downstream of the SNP. This cut–off tended

to exclude SNPs in telomeric regions. Furthermore, SDS scores were not calculated

for chromosome 25 as an insufficient number of singletons were present across all

individuals after data filtering. Raw SDS scores were standardized using 18 bins,

based on alternate allele frequencies at the scored SNP (i.e., from 5% to 10%, from

10% to 15%, etc.). Standardised SDS scores (denoted sSDS) were obtained by

subtracting the bin mean score from individual measures, and dividing by the bin

standard deviation.

Statistical analyses were carried out in R (R Core Team, 2021). *P*–values for

each SNP were determined by calculating the probability of the observed sSDS

values from a standardised normal distribution. Significance was determined using a

Bonferroni–corrected cut–off of $0.05/(1{,}877{,}770) \approx 2.7 \times 10^{-8}$. The false–discovery

rate (FDR) of each SNP was calculated using the 'qvalue' package (Storey *et al.*,

2021); we highlighted SNPs with an FDR of less than 0.05.

Data sources

Version 104 of the GTF gene annotation file for the ARS–UCD 1.2 assembly

was downloaded from Ensembl (available from

https://www.ensembl.org/Bos_taurus/Info/Index). Bedtools v2.29.0 (Quinlan & Hall,

2010) was used to obtain genetic annotations 10kb up– and downstream of each

Bonferroni–significant SNP; overlapping windows were merged.

Milk fat percentage and milk protein percentage QTLs were obtained from van

den Berg et al. (2020). We retained only those QTLs for which effect sizes were

reported in both Holstein breeds that were studied (French and German Holstein),

and the effect was in the same direction in each. sSDS scores were then obtained for

the nearest SNP to each QTL. Some QTLs lie close to the telomeres, where sSDS

statistics were not available. These QTLs were not considered further; for each of the

remaining QTLs, we identified the SNP nearest to it and assigned the sSDS value at

that site to the QTL. Since QTL effect directions were given for the alternate allele (I.

v. d. Berg, personal communication), we re-polarised sSDS scores for SNPs where

the ancestral allele did not match the reference allele by switching its sign. We

additionally re-polarised scores so that a positive value indicates selection acting on

the trait-increasing allele (Field et al., 2016). 30 of 138 milk fat QTLs had scores

assigned to them, while 78 of 176 milk protein QTLs had scores assigned to them.

Stature QTLs were obtained from Bouwman et al. (2018), which identified 164

QTLs in several B. taurus breeds, including 7 Holstein populations from different

countries. We initially extracted 114 QTLs for which an effect was inferred from at

least 5 of 7 Holstein populations. Positions were given relative to the UMD 3.1

assembly; we subsequently extracted sequence 100bp up– and downstream of the

position and remapped to ARS–UCD 1.0.25. 106 QTLs were re–aligned without

gaps; of the remaining 8, 4 were located close to rearrangements and discarded,

while the remaining 4 were kept. After removing those on chromosome 25, 107 QTLs

were retained for downstream analysis. We analysed this full QTL set and a subset

where effect sizes were reported in 6 of 7 Holstein breeds (containing 78 QTLs). We

obtained the sSDS value of SNPs near QTLs and re-polarised them so that positive

scores indicated selection on the trait-increasing allele. Overall, sSDS values were

assigned to 58 QTLs with effect sizes in at least 5 of 7 Holstein populations, and 42

QTLs with effect sizes in at least 6 of 7 Holstein populations.

Statistical analyses of polygenic selection

To determine whether evidence exists for polygenic selection acting on QTLs

underpinning these traits, we applied the 'lm' function in R to fit a linear model

between sSDS values of the nearest SNP to the QTL, and the absolute $\log_{10}$ *p*-value

of it as proxy for effect size. To implement the permutation test, we subsampled

SNPs from the genome that were from the same chromosome and derived-allele

frequency bin as the SNPs used in the polygenic selection tests analyses; we also

changed the sign of the sSDS value of the sampled SNP if this change was also

made to the score associated with the corresponding true SNP. We then performed a

linear model fit between these permuted sSDS values and the QTL *p*-values, and

obtained the *F*-statistic for it. 1,000 subsampled datasets were used to generate the

neutral distribution.

Power simulations given Holstein demography

To estimate the power of the SDS methods when applied to the Holstein

breed, we implemented a simulation pipeline as used in previous studies of polygenic

adaptation (Field *et al.*, 2016; Speidel *et al.*, 2019). We used the *simuPOP* program

(Peng & Kimmel, 2005) to simulate the trajectory of either a neutral variant, or a

partial selective sweep with homozygote selection coefficient of 0.2 (and a

heterozygote value of half that value). The target mutation appeared *de novo* and we

sampled individuals when it reached a frequency of 70%. We assumed the Holstein

population demography as inferred by Boitard et al. (2016). Simulated trajectories

were then piped into the coalescent simulation program *mbs* (Teshima & Innan,

2009) to generate polymorphism data for 204 haplotypes. We assumed a per-site

recombination rate of 3.66×10^{-9} (Boitard *et al.*, 2016b), and a mutation rate of either

1.21×10^{-8} as inferred from parent-offspring trios of dairy cattle (Harland *et al.*, 2018),

or one ten-fold higher. A region of 10Mb long was simulated, with the target mutation

appearing in the centre of it. Diploid genotypes were formed by randomly pairing

haplotypes. SDS scores were calculated using the same pipeline and parameter

values as for the data analyses. We also recorded the number of singletons per

individual, and their distribution per simulation. 100 replicate simulations were run for

each case. The confidence intervals for singleton distributions were calculated using

1,000 bootstraps. When plotting the distribution of SDS scores for the high-mutation

case (Figure 5), they were first normalised by subtracting the mean of the neutral

case from each value, then dividing by the standard deviation of neutral scores.

Acknowledgements. We would like to thank Simon Boitard for sharing his

results on *B. taurus* demographic inference; Simon Boitard, Jon Slate and two

anonymous reviewers for providing feedback on the manuscript. MH is supported by

a NERC Independent Research Fellowship (NE/R015686/1). NAP, BG and TB are

funded by a synergistic research grant from the Faculty of Science and Technology,

Aarhus University, Denmark. MH and TB also acknowledge financial support from the

European Research Council under the European Union's Seventh Framework

Program (FP7/20072013, ERC Grant 311341). The authors declare no conflicts of

interest.

Author contributions. All authors contributed to the study design. NAP and

BG provided data. MH performed the analyses and wrote the manuscript, with

feedback from NAP, BG and TB.

Data archiving. Raw SDS scores and polarisation information has been

deposited on Dryad (under curation, provisional DOI

https://doi.org/10.5061/dryad.547d7wm8q). Data analysis and simulation scripts are

available on GitHub (https://github.com/MattHartfield/CattleSDS).

References

Barghi, N., Hermisson, J. & Schlötterer, C. (2020) Polygenic adaptation: a unifying framework to understand positive selection. *Nat. Rev. Genet.*, **21**, 769–781.

Barton, N.H., Etheridge, A.M. & Véber, A. (2017) The infinitesimal model: Definition, derivation, and implications. *Theor. Popul. Biol.*, **118**, 50–73.

Barton, N.H., Hermisson, J. & Nordborg, M. (2019) Why structure matters. *eLife*, **8**, e45380.

Barton, N.H. & Keightley, P.D. (2002) Understanding quantitative genetic variation. *Nat. Rev. Genet.*, **3**, 11–21.

Berg, I. van den, Xiang, R., Jenko, J., Pausch, H., Boussaha, M., Schrooten, C., *et al.* (2020) Meta-analysis for milk fat and protein percentage using imputed sequence variant genotypes in 94,321 cattle from eight cattle breeds. *Genet. Sel. Evol.*, **52**, 37.

Berg, J.J. & Coop, G. (2014) A Population Genetic Signal of Polygenic Adaptation. *PLoS Genet.*, **10**, e1004412.

Berg, J.J., Harpak, A., Sinnott-Armstrong, N., Joergensen, A.M., Mostafavi, H., Field, Y., *et al.* (2019a) Reduced signal for polygenic adaptation of height in UK Biobank. *eLife*, **8**, e39725.

Berg, J.J., Zhang, X. & Coop, G. (2019b) Polygenic Adaptation has Impacted Multiple Anthropometric Traits. *bioRxiv*, 167551.

Boitard, S., Boussaha, M., Capitan, A., Rocha, D. & Servin, B. (2016a) Uncovering Adaptation from Sequence Data: Lessons from Genome Resequencing of Four Cattle Breeds. *Genetics*, **203**, 433–450.

Boitard, S. & Rocha, D. (2013) Detection of signatures of selective sweeps in the Blonde d'Aquitaine cattle breed. *Anim. Genet.*, **44**, 579–583.

Boitard, S., Rodríguez, W., Jay, F., Mona, S. & Austerlitz, F. (2016b) Inferring

Population Size History from Large Samples of Genome-Wide Molecular Data - An

Approximate Bayesian Computation Approach. *PLoS Genet.*, **12**, e1005877.

Bolger, A.M., Lohse, M. & Usadel, B. (2014) Trimmomatic: a flexible trimmer for

Illumina sequence data. *Bioinformatics*, **30**, 2114–2120.

Bouwman, A.C., Daetwyler, H.D., Chamberlain, A.J., Ponce, C.H., Sargolzaei, M.,

Schenkel, F.S., *et al.* (2018) Meta-analysis of genome-wide association studies for

cattle stature identifies common genes that regulate body size in mammals. *Nat.*

Genet., **50**, 362–367.

Boyle, E.A., Li, Y.I. & Pritchard, J.K. (2017) An Expanded View of Complex Traits:

From Polygenic to Omnigenic. *Cell*, **169**, 1177–1186.

Buitenhuis, B., Poulsen, N.A., Gebreyesus, G. & Larsen, L.B. (2016) Estimation of

genetic parameters and detection of chromosomal regions affecting the major milk

proteins and their post translational modifications in Danish Holstein and Danish

Jersey cattle. *BMC Genet.*, **17**, 114.

Bycroft, C., Freeman, C., Petkova, D., Band, G., Elliott, L.T., Sharp, K., *et al.* (2018)

The UK Biobank resource with deep phenotyping and genomic data. *Nature*, **562**,

203–209.

Castellano, D., Uricchio, L.H., Munch, K. & Enard, D. (2019) Viruses rule over

adaptation in conserved human proteins. *bioRxiv*, 555060.

Chaisson, M.J.P., Wilson, R.K. & Eichler, E.E. (2015) Genetic variation and the *de*

novo assembly of human genomes. *Nat. Rev. Genet.*, **16**, 627–640.

Charlesworth, B. & Edwards, A.W.F. (2018) A century of variance. *Significance*, **15**,

20–25.

Chevin, L.-M. (2019) Selective Sweep at a QTL in a Randomly Fluctuating

Environment. *Genetics*, **213**, 987–1005.

Chevin, L.-M. & Hospital, F. (2008) Selective Sweep at a Quantitative Trait Locus in

the Presence of Background Genetic Variation. *Genetics*, **180**, 1645–1660.

Coop, G., Witonsky, D., Di Rienzo, A. & Pritchard, J.K. (2010) Using Environmental

Correlations to Identify Loci Underlying Local Adaptation. *Genetics*, **185**, 1411–1423.

Daetwyler, H.D., Capitan, A., Pausch, H., Stothard, P., Binsbergen, R. van, Brøndum,

R.F., *et al.* (2014) Whole-genome sequencing of 234 bulls facilitates mapping of

monogenic and complex traits in cattle. *Nat. Genet.*, **46**, 858–865.

Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., *et al.*

(2011) The variant call format and VCFtools. *Bioinformatics*, **27**, 2156–2158.

Daub, J.T., Hofer, T., Cutivet, E., Dupanloup, I., Quintana-Murci, L., Robinson-

Rechavi, M., *et al.* (2013) Evidence for Polygenic Adaptation to Pathogens in the

Human Genome. *Mol. Biol. Evol.*, **30**, 1544–1558.

Daub, J.T., Moretti, S., Davydov, I.I., Excoffier, L. & Robinson-Rechavi, M. (2017)

Detection of Pathways Affected by Positive Selection in Primate Lineages Ancestral

to Humans. *Mol. Biol. Evol.*, **34**, 1391–1402.

Duntsch, L., Tomotani, B.M., Villemereuil, P. de, Brekke, P., Lee, K.D., Ewen, J.G., *et*

al. (2020) Polygenic basis for adaptive morphological variation in a threatened

Aotearoa | New Zealand bird, the hihi (*Notiomystis cincta*). *Proc. R. Soc. B.*, **287**,

20200948.

Edge, M.D. & Coop, G. (2019) Reconstructing the History of Polygenic Scores Using

Coalescent Trees. *Genetics*, **211**, 235–262.

Ehrlich, M.A., Wagner, D.N., Oleksiak, M.F. & Crawford, D.L. (2021) Polygenic

Selection within a Single Generation Leads to Subtle Divergence among Ecological

Niches. *Genome Biol. Evol.*, **13**, evaa257.

Exposito-Alonso, M., 500 Genomes Field Experiment Team, Burbano, H.A.,

Bossdorf, O., Nielsen, R. & Weigel, D. (2019) Natural selection on the *Arabidopsis*

thaliana genome in present and future climates. *Nature*, **573**, 126–129.

Exposito-Alonso, M., Vasseur, F., Ding, W., Wang, G., Burbano, H.A. & Weigel, D.

(2018) Genomic basis and evolutionary potential for extreme drought adaptation in

Arabidopsis thaliana. Nat. Ecol. Evol., **2**, 352–358.

Exposito-Alonso, M., Wilton, P. & Nielsen, R. (2020) Non-additive polygenic models

improve predictions of fitness traits in three eukaryote model species. *bioRxiv*,

2020.07.14.194407.

Fagny, M. & Austerlitz, F. (2021) Polygenic Adaptation: Integrating Population

Genetics and Gene Regulatory Networks. *Trends Genet.*, **37**, 631–638.

Field, Y., Boyle, E.A., Telis, N., Gao, Z., Gaulton, K.J., Golan, D., *et al.* (2016)

Detection of human adaptation during the past 2000 years. *Science*, **354**, 760–764.

Fisher, R.A. (1918) The correlation between relatives on the supposition of

Mendelian inheritance. *Trans. R. Soc. Edinb.*, **52**, 399–433.

Fuhrmann, N., Prakash, C. & Kaiser, T.S. (2021) Polygenic adaptation from standing

genetic variation allows rapid ecotype formation. *bioRxiv*, 2021.04.16.440113.

Gazal, S., Finucane, H.K., Furlotte, N.A., Loh, P.-R., Palamara, P.F., Liu, X., *et al.*

(2017) Linkage disequilibrium–dependent architecture of human complex traits

shows action of negative selection. *Nat. Genet.*, **49**, 1421–1427.

Georges, M., Charlier, C. & Hayes, B. (2018) Harnessing genomic information for

livestock improvement. *Nat. Rev. Genet.*

Gramlich, S., Liu, X., Favre, A., Alex Buerkle, C. & Karrenberg, S. (2021) A polygenic

architecture with conditionally neutral effects underlies ecological differentiation in

Silene. bioRxiv, 2021.07.06.451304.

Haile-Mariam, M., Nieuwhof, G.J., Beard, K.T., Konstatinov, K.V. & Hayes, B.J.

(2013) Comparison of heritabilities of dairy traits in Australian Holstein-Friesian cattle

from genomic and pedigree data and implications for genomic evaluations:

Implication of genomic heritability for genomic evaluation. *J. Anim. Breed. Genet.*,

130, 20–31.

Harland, C., Coppieters, W., Mullaart, E., Charlier, C. & Georges, M. (2018) *Rate of*

de novo mutation in dairy cattle and potential impact of reproductive technologies.

Proc. World Congr. Genet. Appl. to Livest. Prod.

Harpending, H.C., Batzer, M.A., Gurven, M., Jorde, L.B., Rogers, A.R. & Sherry, S.T.

(1998) Genetic traces of ancient demography. *Proc. Natl. Acad. Sci. USA*, **95**, 1961–

1967.

Hayes, B.J., Bowman, P.J., Chamberlain, A.J. & Goddard, M.E. (2009) Invited review:

Genomic selection in dairy cattle: Progress and challenges. *J. Dairy Sci.*, **92**, 433–

443.

Hayward, L.K. & Sella, G. (2019) Polygenic adaptation after a sudden change in

environment. *bioRxiv*, 792952.

He, F., Arce, A.L., Schmitz, G., Koornneef, M., Novikova, P., Beyer, A., *et al.* (2016)

The Footprint of Polygenic Adaptation on Stress-Responsive Cis-Regulatory

Divergence in the *Arabidopsis* Genus. *Mol. Biol. Evol.*, **33**, 2088–2101.

Jain, K. & Stephan, W. (2015) Response of Polygenic Traits Under Stabilizing

Selection and Mutation When Loci Have Unequal Effects. *G3*, **5**, 1065–1074.

Jain, K. & Stephan, W. (2017a) Modes of Rapid Polygenic Adaptation. *Mol. Biol.*

Evol., **34**, 3169–3175.

Jain, K. & Stephan, W. (2017b) Rapid Adaptation of a Polygenic Trait After a Sudden

Environmental Shift. *Genetics*, **206**, 389–406.

Jiménez-Mena, B., Tataru, P., Brøndum, R.F., Sahana, G., Guldbrandtsen, B. &

Bataillon, T. (2016) One size fits all? Direct evidence for the heterogeneity of genetic

drift throughout the genome. *Biol. Lett.*, **12**.

John, S. & Stephan, W. (2020) Important role of genetic drift in rapid polygenic

adaptation. *Ecol. Evol.*, **10**, 1278–1287.

Josephs, E.B., Berg, J.J., Ross-Ibarra, J. & Coop, G. (2019) Detecting Adaptive

Differentiation in Structured Populations with Genomic Data and Common Gardens.

Genetics, **211**, 989–1004.

Kelleher, J., Etheridge, A.M. & McVean, G. (2016) Efficient Coalescent Simulation

and Genealogical Analysis for Large Sample Sizes. *PLoS Comput. Biol.*, **12**,

e1004842.

Kreiner, J.M., Tranel, P.J., Weigel, D., Stinchcombe, J.R. & Wright, S.I. (2020) The

genetic architecture and genomic context of glyphosate resistance in *Amaranthus*

tuberculatus. *bioRxiv*, 2020.08.19.257972.

Lande, R. (1983) The response to selection on major and minor mutations affecting a

metrical trait. *Heredity*, **50**, 47–65.

Le Corre, V. & Kremer, A. (2012) The genetic differentiation at quantitative trait loci

under local adaptation. *Mol. Ecol.*, **21**, 1548–1566.

Lemay, D.G., Lynn, D.J., Martin, W.F., Neville, M.C., Casey, T.M., Rincon, G., *et al.*

(2009) The bovine lactation genome: insights into the evolution of mammalian milk.

Genome Biol., **10**, R43.

Li, H. & Durbin, R. (2009) Fast and accurate short read alignment with Burrows–

Wheeler transform. *Bioinformatics*, **25**, 1754–1760.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., *et al.* (2009) The

Sequence Alignment/Map format and SAMtools. *Bioinformatics*, **25**, 2078–2079.

Liu, X., Li, Y.I. & Pritchard, J.K. (2019) Trans Effects on Gene Expression Can Drive

Omnigenic Inheritance. *Cell*, **177**, 1022-1034.e6.

MacEachern, S., Hayes, B., McEwan, J. & Goddard, M. (2009) An examination of

positive selection and changing effective population size in Angus and Holstein cattle

populations (*Bos taurus*) using a high density SNP genotyping platform and the

contribution of ancient polymorphism to genomic diversity in Domestic cattle. *BMC*

Genom., **10**, 181.

MacLachlan, I.R., McDonald, T.K., Lind, B.M., Rieseberg, L.H., Yeaman, S. & Aitken,

S.N. (2021) Genome-wide shifts in climate-related variation underpin responses to

selective breeding in a widespread conifer. *Proc. Natl. Acad. Sci. USA*, **118**,

e2016900118.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., *et*

al. (2010) The Genome Analysis Toolkit: A MapReduce framework for analyzing next-

generation DNA sequencing data. *Genome Res.*, **20**, 1297–1303.

Meuwissen, T., Hayes, B. & Goddard, M. (2013) Accelerating Improvement of

Livestock with Genomic Selection. *Annu. Rev. Anim. Biosci.*, **1**, 221–237.

Nielsen, R. (2005) Molecular Signals of Natural Selection. *Annu. Rev. Genet.*, **39**,

197–218.

Nordborg, M. & Donnelly, P. (1997) The Coalescent Process With Selfing. *Genetics*,

146, 1185–1195.

Novembre, J. & Barton, N.H. (2018) Tread Lightly Interpreting Polygenic Tests of

Selection. *Genetics*, **208**, 1351–1355.

Pavlidis, P., Metzler, D. & Stephan, W. (2012) Selective Sweeps in Multilocus Models

of Quantitative Traits. *Genetics*, **192**, 225–239.

Peng, B. & Kimmel, M. (2005) simuPOP: a forward-time population genetics

simulation environment. *Bioinformatics*, **21**, 3686–3687.

Pritchard, J.K. & Di Rienzo, A. (2010) Adaptation - not by sweeps alone. *Nat. Rev.*

Genet., **11**, 665–667.

Pritchard, J.K., Pickrell, J.K. & Coop, G. (2010) The Genetics of Human Adaptation:

Hard Sweeps, Soft Sweeps, and Polygenic Adaptation. *Curr. Biol.*, **20**, R208–R215.

Qanbari, S., Pausch, H., Jansen, S., Somel, M., Strom, T.M., Fries, R., *et al.* (2014)

Classic Selective Sweeps Revealed by Massive Sequencing in Cattle. *PLoS Genet.*,

10, e1004148.

Qanbari, S., Pimentel, E.C.G., Tetens, J., Thaller, G., Lichtner, P., Sharifi, A.R., *et al.*

(2010) A genome-wide scan for signatures of recent selection in Holstein cattle.

Anim. Genet.

Quinlan, A.R. & Hall, I.M. (2010) BEDTools: a flexible suite of utilities for comparing

genomic features. *Bioinformatics*, **26**, 841–842.

R Core Team. (2021) *R: A Language and Environment for Statistical Computing*. R

Foundation for Statistical Computing, Vienna, Austria.

Racimo, F., Berg, J.J. & Pickrell, J.K. (2018) Detecting Polygenic Adaptation in

Admixture Graphs. *Genetics*, **208**, 1565–1584.

Robinson, M.R., Hemani, G., Medina-Gomez, C., Mezzavilla, M., Esko, T.,

Shakhbazov, K., *et al.* (2015) Population genetic differentiation of height and body

mass index across Europe. *Nat. Genet.*, **47**, 1357.

Rosen, B., Bickhart, D.M., Schnabel, R.D., Koren, S., Elsik, C., Zimin, A.V., *et al.*

(2018) Modernizing the Bovine Reference Genome Assembly. *Proc. World Congr.*

Genet. Appl. Livest. Prod., 11–16.

Rowan, T.N., Durbin, H.J., Seabury, C.M., Schnabel, R.D. & Decker, J.E. (2021)

Powerful detection of polygenic selection and evidence of environmental adaptation

in US beef cattle. *PLoS Genet.*, **17**, e1009652.

Sanjak, J.S., Sidorenko, J., Robinson, M.R., Thornton, K.R. & Visscher, P.M. (2018)

Evidence of directional and stabilizing selection in contemporary humans. *Proc. Natl.*

Acad. Sci. USA, **115**, 151–156.

Savolainen, O., Lascoux, M. & Merila, J. (2013) Ecological genomics of local

adaptation. *Nat. Rev. Genet.*, **14**, 807–820.

Schoech, A.P., Jordan, D.M., Loh, P.-R., Gazal, S., O'Connor, L.J., Balick, D.J., *et al.*

(2019) Quantification of frequency-dependent genetic architectures in 25 UK Biobank

traits reveals action of negative selection. *Nat. Commun.*, **10**, 790.

Sella, G. & Barton, N.H. (2019) Thinking About the Evolution of Complex Traits in the

Era of Genome-Wide Association Studies. *Annu. Rev. Genom. Hum. Genet.*, **20**,

461–493.

Sohail, M., Maier, R.M., Ganna, A., Bloemendal, A., Martin, A.R., Turchin, M.C., *et al.*

(2019) Polygenic adaptation on height is overestimated due to uncorrected

stratification in genome-wide association studies. *eLife*, **8**, e39702.

Sørensen, A.C., Sørensen, M.K. & Berg, P. (2005) Inbreeding in Danish Dairy Cattle

Breeds. *J. Dairy Sci.*, **88**, 1865–1872.

Soyeurt, H., Gillon, A., Vanderick, S., Mayeres, P., Bertozzi, C. & Gengler, N. (2007)

Estimation of Heritability and Genetic Correlations for the Major Fatty Acids in Bovine

Milk. *J. Dairy Sci.*, **90**, 4435–4442.

Speidel, L., Forest, M., Shi, S. & Myers, S.R. (2019) A method for genome-wide

genealogy estimation for thousands of samples. *Nat Genet.*, **51**, 1321–1329.

Stephan, W. (2016) Signatures of positive selection: from selective sweeps at

individual loci to subtle allele frequency changes in polygenic adaptation. *Mol. Ecol.*,

25, 79–88.

Stephan, W. (2019) Selective Sweeps. *Genetics*, **211**, 5–13.

Stern, A.J., Speidel, L., Zaitlen, N.A. & Nielsen, R. (2021) Disentangling selection on

genetically correlated polygenic traits via whole-genome genealogies. *Am. J. Hum.*

Genet., **108**, 219–239.

Stetter, M.G., Thornton, K. & Ross-Ibarra, J. (2018) Genetic architecture and

selective sweeps after polygenic adaptation to distant trait optima. *PLoS Genet.*, **14**,

e1007794.

Storey, J.D., Bass, A.J., Dabney, A. & Robinson, D. (2021) qvalue: Q-value

estimation for false discovery rate control [WWW Document]. URL

http://github.com/StoreyLab/qvalue [accessed on 2021].

Svardal, H., Jasinska, A.J., Apetrei, C., Coppola, G., Huang, Y., Schmitt, C.A., *et al.*

(2017) Ancient hybridization and strong adaptation to viruses across African vervet

monkey populations. *Nat. Genet.*, **49**, 1705–1713.

Teshima, K.M. & Innan, H. (2009) mbs: modifying Hudson's ms software to generate

samples of DNA sequences with a biallelic site under selection. *BMC Bioinform.*, **10**,

166.

Thornton, K.R. (2019) Polygenic Adaptation to an Environmental Shift: Temporal

Dynamics of Variation Under Gaussian Stabilizing Selection and Additive Effects on a

Single Trait. *Genetics*, **213**, 1513–1530.

Turchin, M.C., Chiang, C.W., Palmer, C.D., Sankararaman, S., Reich, D., Genetic

Investigation of ANthropometric Traits (GIANT) Consortium, *et al.* (2012) Evidence of

widespread selection on standing variation in Europe at height-associated SNPs. *Nat*

Genet., **44**, 1015–1019.

Uricchio, L.H., Kitano, H.C., Gusev, A. & Zaitlen, N.A. (2019) An evolutionary

compass for detecting signals of polygenic selection and mutational bias. *Evol. Lett.*,

3, 69–79.

Visscher, P.M. & Goddard, M.E. (2019) From R.A. Fisher's 1918 Paper to GWAS a

Century Later. *Genetics*, **211**, 1125–1130.

Vitti, J.J., Grossman, S.R. & Sabeti, P.C. (2013) Detecting Natural Selection in

Genomic Data. *Annu. Rev. Genet.*, **47**, 97–120.

Vladar, H.P. de & Barton, N. (2014) Stability and Response of Polygenic Traits to

Stabilizing Selection and Mutation. *Genetics*, **197**, 749–767.

Whitlock, M.C. (2008) Evolutionary inference from F_{ST}. *Mol. Ecol.*, **17**, 1885–1896.

Wieters, B., Steige, K.A., He, F., Koch, E.M., Ramos-Onsins, S.E., Gu, H., *et al.*

(2021) Polygenic adaptation of rosette growth in Arabidopsis thaliana. *PLoS Genet.*,

17, e1008748.

Wollstein, A. & Stephan, W. (2014) Adaptive Fixation in Two-Locus Models of

Stabilizing Selection and Genetic Drift. *Genetics*, **198**, 685–697.

Wray, N.R., Kemper, K.E., Hayes, B.J., Goddard, M.E. & Visscher, P.M. (2019)

Complex Trait Prediction from Genome Data: Contrasting EBV in Livestock to PRS in

Humans. *Genetics*, **211**, 1131–1141.

Wray, N.R., Wijmenga, C., Sullivan, P.F., Yang, J. & Visscher, P.M. (2018) Common

Disease Is More Complex Than Implied by the Core Gene Omnigenic Model. *Cell*,

173, 1573–1580.

Wright, S. (1951) The genetical structure of populations. *Ann. Eugen.*, **15**, 323–354.

Wu, D.-D., Ding, X.-D., Wang, S., Wójcik, J.M., Zhang, Y., Tokarska, M., *et al.* (2018)

Pervasive introgression facilitated domestication and adaptation in the *Bos* species

complex. *Nat. Ecol. Evol.*, **2**, 1139–1145.

Yang, J., Bakshi, A., Zhu, Z., Hemani, G., Vinkhuyzen, A.A.E., Lee, S.H., *et al.* (2015)

Genetic variance estimation with imputed variants finds negligible missing heritability

for human height and body mass index. *Nat. Genet.*, **47**, 1114–1120.

Yeaman, S., Hodgins, K.A., Lotterhos, K.E., Suren, H., Nadeau, S., Degner, J.C., *et*

al. (2016) Convergent local adaptation to climate in distantly related conifers.

Science, **353**, 1431.

Zan, Y. & Carlborg, Ö. (2018) A Polygenic Genetic Architecture of Flowering Time in

the Worldwide Arabidopsis thaliana Population. *Mol. Biol. Evol.*, **36**, 141–154.

Zeder, M.A. (2008) Domestication and early agriculture in the Mediterranean Basin:

Origins, diffusion, and impact. *Proc. Natl. Acad. Sci. USA*, **105**, 11597–11604.

Zeng, J., Vlaming, R. de, Wu, Y., Robinson, M.R., Lloyd-Jones, L.R., Yengo, L., *et al.*

(2018) Signatures of negative selection in the genetic architecture of human complex

traits. *Nat. Genet.*, **50**, 746–753.

Zeng, J., Xue, A., Jiang, L., Lloyd-Jones, L.R., Wu, Y., Wang, H., *et al.* (2021)

Widespread signatures of natural selection across human complex traits and

functional genomic categories. *Nat. Commun.*, **12**, 1164.

Zhang, W., Reeves, G.R. & Tautz, D. (2021) Testing Implications of the Omnigenic

Model for the Genetic Analysis of Loci Identified through Genome-wide Association.

Curr. Biol., **31**, 1092-1098.e6.

Zhao, F., McParland, S., Kearney, F., Du, L. & Berry, D.P. (2015) Detection of

selection signatures in dairy and beef cattle using high-density genomic information.

Genet. Sel. Evol., **47**, 49.

Zhou, X., Carbonetto, P. & Stephens, M. (2013) Polygenic Modeling with Bayesian

Sparse Linear Mixed Models. *PLoS Genet.*, **9**, e1003264.

Figure 1: Simulated mean tip age for *B. taurus*, as a function of the number of haploid

samples. Simulations assumed either (a) demography as inferred by Boitard et al.

(2016b) (the 'High N_0' model), or (b) the same but with a smaller present–day N_e of

49 (the 'Low N_0' model). Points are the mean values; bars show 95% confidence

intervals. The solid line is the best linear fit to the log of both values; dotted lines

942   show the predicted tip age for 204 alleles.

sSDS Results for *Bos taurus* Autosomes

943
944

Figure 2: *P*–values of sSDS scores across *B. taurus* autosomes, as plotted on a

negative Log$_{10}$ scale, as a function of the chromosome. Alternating black and grey

points show (non–significant) values from different chromosomes. Blue points are

SNPs with *FDR* < 0.05, with the cut-off denoted by a horizontal dotted line. Red

points are SNPs with Bonferroni–corrected *P*–value < 0.05 (actual *P*–value < ~2.7 x

10^{-8}), with the cut-off denoted by a horizontal dashed line. Figure S1 shows results

951   for the Low *N*$_0$ model.

Figure 3: Relationship between tSDS scores near milk or stature QTLs, as noted in

the subheadings, and the absolute log *P*-value of QTLs. Lines show a linear model

regression fit. Figure S3 shows results assuming a low N_0 model.

Figure 4: Mean distribution of singleton numbers per individual for each simulation,

wither assuming a standard mutation rate (left) or a 10-fold higher mutation rate

(right). Bars represent 95% confidence intervals.

Figure 5: Distribution of simulated sSDS scores assuming a high mutation rate, for the neutral and selected cases. Numbers above each box plot denotes how many simulations produced SDS scores and were included in the plot.

Chromosome	Gene Name	Start Position	End Position	Gene Biotype	High, Low N_0
1	PPM1L	106405113	106727070	Protein Coding	High, Low
5	TMCC3	24306913	24595494	Protein Coding	High, Low
5	CEP83	24070404	24345243	Protein Coding	High, Low
17	U6	43381106	43381209	snRNA	Low
17	CTSO	43364999	43381605	Protein Coding	Low
17	TDO2	43386894	43403747	Protein Coding	High, Low
23	OR12D2H	29291787	29292713	Protein Coding	High, Low
23	OR12D2E	29305933	29309785	Protein Coding	High, Low
24	GAREM1	24694637	24927333	Protein Coding	High, Low
29	NTM	34576918	34994005	Protein Coding	High, Low

970 Table 1: Genes that overlap or lie close to Bonferroni–significant sSDS regions. The

971 'High, Low N_0' column specifies which genes are close to significant SNPs for each

972 N_0 model.