Inhibitor of apoptosis proteins 1 and 2 are required for the homeostasis of murine

RORγt[+] γδ T cells

John Rizk[1], Rasmus Agerholm[1], Darshana Kadekar[1,2], Monica Torrellas Viñals [1] and Vasileios

Bekiaris*[1]

[1]Department of Health Technology, Technical University of Denmark, Kemitorvet, Bldg 202,

2800 Kgs Lyngby, Denmark.

[2]current address: Department of Bioscience, Aarhus University. Ny Munkegade 116, 8000

Aarhus C, Denmark.

*Corresponding author: Vasileios Bekiaris vasbek@dtu.dk

Abstract

RORγt[+] γδ T cells, known as γδT17, are an innate-like subset of T cells that produce

interleukin (IL)-17A and initiate type 3 immune responses during infections or autoimmune

pathologies. Herein we show that the cellular inhibitor of apoptosis proteins cIAP1 and 2 are

required for the peripheral homeostasis of γδT17 but not for their thymic development. γδT17

cells that were deficient in both cIAP1 and 2 were profoundly reduced in the peripheral lymph

nodes and skin. Likewise, both RORγt[+] innate lymphoid cells (ILC3) and RORγt[+] Tbet[+] γδ T

cells were reduced in the lamina propria of adult mice. Further, cIAP1 and 2 were required for

the expression of the transcription factors RORγt and cMAF in γδT17 cells during neonatal and

adult life. Single deficiency of either cIAP1 or 2 did not affect the homeostasis or transcription

factor profile of γδT17 cells. Inhibition of the kinase RIPK1 partially reversed the deficiency of

γδT17 cells but not the downregulation of RORγt in cIAP1 and cIAP2 double deficient animals.

Moreover, bone marrow reconstitutions and transfer of neonatal γδ T cells to RAG1[-/-] hosts

showed that both intrinsic and extrinsic factors contribute to the loss of γδT17 cells in cIAP1/2

double deficient mice, while only extrinsic signals were responsible for the decrease of ILC3

cells. Deficiency of γδT17 cells in cIAP1 and cIAP2 double deficient animals, or the presence of

functionally defective γδT17 cells did not confer protection against IMQ-induced psoriasis.

Collectively, our data reveal a previously undescribed role for cIAP1 and cIAP2 in the

homeostasis of γδT17 and ILC3 cells.

Introduction

The interleukin (IL)-17 producing γδ T cells (γδT17) are an innate-like subset of T cells that is

enriched at the mucosal barriers and peripheral lymph nodes (Chien *et al*, 2014). Like their IFNγ

producing counterparts, they arise from thymic progenitors and egress the thymus during

perinatal life (Prinz *et al*, 2013). Mouse γδT17 leave the thymus as pre-programmed effectors;

thus, they only require cytokine stimulation by IL-7, IL-23 or IL-1β to produce copious amounts

of IL-17A, IL-17F and IL-22. Hence, γδT17 are the main producers of IL-17A in the early stages

of infections by *Mycobacterium tuberculosis*, *Candida albicans* or *Listeria monocytogenes* and

are critical for initiating an immune response against such infections (Papotto *et al*, 2017).

Conversely, γδT17 have also been implicated in augmenting autoimmune conditions as psoriasis

and multiple sclerosis. Thus, IL-17A production by γδT17 cells is required for the progression of

disease in EAE, which is a model of multiple sclerosis. Similarly, dermal γδT17 play a critical in

the pathogenesis of psoriasis-like symptoms in imiquimod (IMQ)-induced psoriasis.(Sutton *et al*,

2009; van der Fits *et al*, 2009).

During perinatal thymic development, γδT17 cells undergo transcriptional and epigenetic

programming that imprints their effector phenotypes(Schmolka *et al*, 2013). Thus, thymic

development of γδT17 is dependent on the consecutive expression of the transcription factors

Sox13, cMAF followed by the lineage defining expression of RORγt. Conversely, the IFNγ

producing γδT cells are characterized by the transcription factors Tbet and Tcf1 (Sagar *et al*,

2018; Zuberbuehler *et al*, 2019). Further, canonical and non-canonical NF-κB pathway signaling

in the thymic microenvironment are essential for the development of γδT17. For instance, the

non-canonical NF-κB inducing kinase (NIK) signaling in the thymic epithelia is essential for the

development of γδT17 (Mair *et al*, 2015). Moreover, γδT17 development requires lymphotoxin β

receptor (LTβR) signaling, through the non-canonical NF-κB transcription factor RelB.

Alternatively, the canonical NF-κB transcription factor RelA is required for the optimal

expression of LTβR ligands such as LIGHT and lymphotoxin(LT)α in thymocytes, which are

required for γδT17 development (Powolny-Budnicka *et al*, 2011; Silva-Santos *et al*, 2005).

However, the role of NF-κB signaling in the homeostasis of adult γδT17 is not well described.

Cellular inhibitor of apoptosis (cIAP)1 and 2 and conserved E3 ubiquitin ligases that control

the balance between canonical and non-canonical NF-κB activation downstream of TNF receptor

superfamily(Gyrd-Hansen & Meier, 2010). In unstimulated cells, cIAP1 and 2 associate in a

heterocomplex with TNF receptor associated factor (TRAF)2, TRAF3 and NIK whereby they

induce lysine(K)-48 ubiquitination of NIK, resulting in its proteasomal degradation (Zarnegar *et*

al, 2008; Varfolomeev *et al*, 2007). However, ligation of TNFRII, LTβR, CD40 or OX40

recruits the heterocomplex to the receptor, which results in dissociation of NIK followed by

activation of the non-canonical NF-κB pathway(Vallabhapurapu *et al*, 2008). Conversely, cIAP1

and 2 positively regulate the activation of canonical NF-κB downstream of TNFRI by K-63

stabilizing ubiquitination of receptor interacting kinase (RIPK)1(Bertrand *et al*, 2008).

Furthermore, cIAP1 and 2 inhibit apoptosis and necroptosis through degradative K-48

ubiquitination of RIPK1 in the ripoptosome complex (Annibaldi & Meier, 2018). cIAP1 and 2

have been frequently studied in the context of conventional αβ T cells as well as in cancers,

however their role in γδT cells is obscure.

Herein we investigated the functions of cIAP1 and cIAP2 in the development and homeostasis

of γδT17 and ILC3 cells. We show that mice that lacked cIAP2 and conditionally lacked cIAP1

in all RORγt expressing cells had significantly reduced frequencies and abundance of γδT17

cells in peripheral lymph nodes, skin of adult mice and lacked RORgt$^+$ Tbet$^+$ γδ T cells and ILC3

cells in the gut lamina propria. Consistently, production of IL-17 from γδT cells was severely

diminished in cIAP1 and cIAP2 double deficient mice. Single deficiency of either cIAP1 or

cIAP2 minimally impacted γδT17 development or homeostasis. Intriguingly, we found that

thymic development of γδT17 was not affected in cIAP1 and cIAP2 double deficient mice.

However, cIAP1 and 2 deficient γδT17 in peripheral lymph nodes downregulated the

transcription factors cMAF and RORγt from postnatal day 21 and were progressively reduced in

numbers culminating in marked reduction of γδT17 by week 12. Further, using chow that

contained RIPK1 inhibitor or control chow, we show that the decline in frequency and

abundance of γδT17 was partially dependent on RIPK1 while the downregulation of RORγt was

independent of RIPK1. Also, we found that ectopic expression of the non-canonical NF-κB

transcription factor RelB in γδT17 cells downregulates the expression of cMAF. Using bone

marrow reconstitution and transfer of neonatal γδ T cells to RAG1 deficient hosts, we show that

in mice that lack cIAP2 and conditionally lack cIAP1, both cell extrinsic and intrinsic factors

contribute to the loss of γδT17 cells, while the loss of ILC3 cells were due to cell extrinsic

signals only. Finally, we show that the cIAP1 and cIAP2 -induced downregulation of cMAF and

RORγt renders γδT17 cells functionally defective and inhibits their response to IMQ-induced

skin inflammation but does not affect the develpment of IMQ-induced skin inflammation.

Results

cIAP1 and cIAP2 are redundantly required for the homeostasis of γδT17 cells

In order to understand the functions of cIAP1 and cIAP2 in regulating γδT17 cells, we crossed RORγt-Cre (Ebertl *et al*, 2004) to cIAP1 floxed (cIAP1$^{f/f}$)or cIAP1$^{f/f}$ cIAP2 deficient mice (cIAP1$^{f/f}$ cIAP2$^{-/-}$). Thus, 4 strains (Fig 1A) were generated as follows: RORγt^{cre-} cIAP1$^{f/f}$ (hereafter referred to as littermate controls), RORγt^{cre+} cIAP1$^{f/f}$ (ΔcIAP1), RORγt^{cre-} cIAP1$^{f/f}$ cIAP2$^{-/-}$ (ΔcIAP2) and RORγt^{cre+} cIAP1$^{f/f}$ cIAP2$^{-/-}$ (ΔcIAP1/2). All 4 strains were viable and showed no apparent defects in the development of peripheral lymph nodes (inguinal, brachial and axillary; collectively referred to as pLNs) or peyer's patches (data not shown); this indicated that there was no developmental effect on lymphoid tissue inducer (Lti) cells, which express RORγt and are required for formation of lymph nodes and preyer's patches perinatally (Sawa *et al*, 2010).

We then analyzed γδ T cell populations in the pLNs, skin and small or large intestinal lamina propria of adult mice. In pLNs, we identified γδT17 cells as being TCRγδ$^+$ CD27$^-$ CD44^{hi} CCR6$^+$ cells. We found that ΔcIAP1/2 mice had significantly reduced proportion and numbers of γδT17 compared to either littermate controls, ΔcIAP1 or ΔcIAP2 mice (Fig 1B). The reduction in γδT17 in ΔcIAP1/2 mice affected both Vγ4$^+$ and Vγ4$^-$ γδT17 cells (Vγ chain nomenclature following the Heilig & Tonegawa, 1986) (Fig S1C). We further noticed an increase in TCRγδ$^+$ CD27$^+$ population in ΔcIAP1/2 compared to other strains (Fig S1A). Consistently, there was a significant reduction in IL-17A and IL-22 production and a concomitant increase in IFNγ production by γδ T cells in the peripheral lymph nodes, reflecting the skewed proportions of the γδ T cell populations (Fig S1B). In the skin, we found that ΔcIAP1/2 mice had reduced numbers of CD3$^+$ Vγ5$^-$ TCRγδ$^+$ cells (γδT17), without a concomitant increase in the abundance of TCRβ$^+$

cells (Fig 1C) Further, both Vγ4$^+$ or Vγ4$^-$ skin γδT17 cells were reduced in ΔcIAP1/2 mice,

compared to other strains (Fig S1D).

We have previously characterized a population of RORγt$^+$ Tbet$^+$ γδ T cells in the small

intestinal (siLP) and colonic lamina propria (cLP) (Kadekar *et al*, 2019). Thus, we investigated

the impact of cIAP1 and cIAP2 deficiency on the different RORγt$^+$ γδ T cells in gut lamina

propria. In both small intestinal and colonic lamina propria we found a significant reduction of

RORγt$^+$ Tbet$^+$ γδ T cells and a concomitant increase in Tbet$^+$ γδ T cells in cIAP1 and 2 double

deficient mice (Fig 2A-B).

RORγt$^+$ innate lymphoid cells (ILC3) cells are essential for the homeostasis of intestinal

mucosal tissue and for protection against pathogens. ILC3 cells are generally divided into Tbet$^-$

and Tbet$^+$ subsets (Klose *et al*, 2013; Melo-Gonzalez & Hepworth, 2017). Akin to intestinal γδ T

cells, we found that ΔcIAP1/2 had reduced the frequency and numbers of ILC3 cells in siLP.

However, both Tbet$^+$ and Tbet$^-$ ILC3 cells were equally reduced in ΔcIAP1/2 (Fig 2C).

Collectively, this data show that both cIAP1 and 2 are required for the homeostasis of γδT17 in

the pLNs, skin of adult mice while single deficiency of either has no effect. It also shows that in

the gut lamina propria, cIAP1 and 2 are critical for RORγt$^+$ Tbet$^+$ γδ T and ILC3 cells.

cIAP1 and 2 maintain the expression of cMaf and RORγt in γδT17 during neonatal life

but not thymic development.

The development of γδT17 cells occurs in the thymus at approximately embryonic day 16,

concomitant with the expression of the transcription factors *Rorc* and *Maf* (Haas *et al*, 2012;

Zuberbuehler *et al*, 2019). Thus, we aimed to track the perinatal and neonatal development and

homeostasis of γδT17 in ΔcIAP1/2, ΔcIAP1, ΔcIAP2 and littermate control mice. First, we

analyzed γδT17 in the thymi of 1-day old pups and found that the thymic output of γδT17 in

ΔcIAP1/2 pups was comparable to littermate controls. Thus, there was no difference in the

numbers of γδT17 cells nor in the expression of RORγt and cMaf by γδT cells between 1-day old

ΔcIAP1/2 pups and controls (Fig 3A-B). Further, the proliferation of γδT17 from 1-day old pups,

as measured by Ki67 staining, was also comparable to littermate controls (Fig S2A).

We found that in γδT17 from pLNs, there was a downregulation of both RORγt and cMaf that

started at postnatal day 21 and persisted afterwards, which occurred only ΔcIAP1/2 but not in

littermate controls (Fig 3D). However, the absolute numbers of γδT17 cells (identified as being

CD44^{hi} CD27$^-$) in the pLNs of ΔcIAP1/2 remained comparable to those in littermate control

mice until approximately 5 weeks of age (Fig 3C). This culminated in a marked reduction of

γδT17 cells, identified by either surface markers or RORγt flow cytometry staining, in ΔcIAP1/2

mice at 12 weeks of age (Fig 3C & E). Collectively, this data suggests that cIAP1 and cIAP2 are

required for the adult homeostasis, but not perinatal development, of γδT17 cells in pLNs and

points to a role for cIAP1 and cIAP2 in regulating the expression of RORγt and cMaf in γδT17

cells.

Loss of γδT17, but not RORγt expression, in ΔcIAP1/2 mice is partially dependent on

RIPK1

cIAP1 and 2 regulate the stability and function of RIPK1 by either K63 stabilizing

ubiquitination or K48 ubiquitination that marks RIPK1 for proteasomal degradation; thus, in the

absence of cIAP1 and 2, RIPK1 accumulates in the cytoplasm and seeds the formation of the

ripoptosome complex thereby facilitating apoptosis or necroptosis (Silke & Meier, 2013).

Therefore, we asked whether the progressive reduction of γδT17 in ΔcIAP1/2 mice could be due

to increased RIPK1-dependent cell death. To test this possibility, we weaned mice from all 4

strains on either control chow or chow that contained a RIPK1 inhibitor compound (Ri). We then

analyzed γδT cell populations in pLNs and siLP and ILC3 cells in siLP at when the mice were at

least 11 weeks of age.

Based on surface phenotype, inhibition of RIPK1 in ΔcIAP1/2 partially rescued the frequency

and overall numbers of γδT17 in the pLNs, compared to littermate control mice that had received

control chow. There was no significant difference in frequency nor total numbers of γδT17 in the

pLNs between ΔcIAP1/2 and littermate controls that received Ri chow; this probably reflected

the tendency towards lower abundance of γδT17 in littermate controls on Ri chow compared to

controls receiving regular chow (Fig 4A). Surprisingly, transcription factor staining revealed that

RIPK1 inhibition did not affect the significant downregulation of RORγt in γδT17 cells that were

still present in the pLNs of ΔcIAP1/2 mice (Fig 4B-C). Further, RIPK1 inhibition had no

significant impact on the anti-apoptotic protein Bcl2 in γδT17 cells from ΔcIAP1/2 mice but we

noticed a mild decrease of Bcl2 in γδT17 from ΔcIAP2 mice (Fig. S2B). In siLP, the inhibition

of RIPK1 did not halt the reduction in RORγt$^+$ Tbet$^+$ γδ T cells in double deficient mice,

compared to control mice or mice that received control chow (Fig 4D). Taken together, our data

indicate that the decline in γδT17 frequency and abundance is partially dependent on RIPK1

while the concomitant downregulation in RORγt in γδT17 in ΔcIAP1/2 mice is RIPK1

independent.

Activation of the non-canonical NF-κB in γδT17 cells downregulates cMAF expression.

We have previously shown that the non-canonical NF-κB transcription factor RelB

downregulates the expression of the transcription factor cMAF in Th17 cells; this occurs

following the activation of the non-canonical NF-κB pathway through depletion of cIAP1 and 2

by SMAC mimetic compounds (Rizk *et al*, 2019). Thus, we hypothesized that the lack of cIAP1

and 2 expression in γδT17 activates the non-canonical NF-κB pathway and mobilizes RelB to

downregulate cMAF. To test this hypothesis, we purified γδT17 cells from thymi old 1-day old

wildtype mice and infected them with an empty retroviral vector or vector expressing RelB and

cultured the cells with IL-7. Subsequently, we FACS sorted infected γδT17 cells (TCRγδ$^+$

CD44^{hi} CD27$^-$ GFP$^+$) and extracted RNA to assess the expression of key γδT17 transcription

factors. As expected, the RelB-RV infected cells expressed more RelB and NFκB2 compared to

empty-RV infected cells (Fig 5A). Furthermore, we found that RelB induced a downregulation

of *Maf*, but did not affect the expression of *Rorc* or *Sox13* (Fig 5B). This indicates that similar to

what happens in Th17 cells, activation of the non-canonical NFκB in γδT17 cells downregulates

cMAF.

Intrinsic and extrinsic factors contribute to the loss of γδT17 in ΔcIAP1/2 mice.

cIAP1 and cIAP2 regulate the cells response to extrinsic stimuli to TNF and other TNFRSF

ligands but are also important to inhibit cell death through control over the formation of the

ripoptosome. Therefore, we set to investigate whether cell extrinsic or intrinsic signals were

responsible for the loss of γδT17 cells in the ΔcIAP1/2 mice. To this end we set up mixed bone

marrow chimeras where wildtype CD45.1/CD45.2 hosts were sublethally irradiated and

reconstituted with a mixture of 1:1 wildtype (CD45.1) and ΔcIAP1/2 (CD45.2) bone marrow

cells (Fig 6A). We found that, under these conditions, γδT17 and CD27$^+$ γδ T cells from

ΔcIAP1/2 bone marrow reconstituted the pLNs of the host as competitively as their wildtype

counterparts (Fig 6B). Further, only Tbet$^+$ γδ T cells reconstituted the colonic lamina propria and

were slightly dominated by cells that originated from ΔcIAP1/2 bone marrow (BM) (Fig 6C).

Conversely, colonic ILC3 cells were effectively reconstituted and were encompassed

approximately equal proportions of CD45.1 (WT) and CD45.2 (ΔcIAP1/2) cells (Fig 6D). This

indicates that the presence of wildtype hematopoietic and stromal components prevents the loss

of γδT17 from ΔcIAP1/2 origin.

We then asked whether the reconstitution of ΔcIAP1/2 mice with mixed BM chimeras would

be enough to stop the loss of γδT17 cells or not. To this end, we irradiated ΔcIAP1/2 mice that

were at least 12 weeks old, to ensure host mice lacked γδT17 cells, and reconstituted them with

either wildtype (CD45.1) or a mixture of 1:1 wildtype (CD45.1) and ΔcIAP1/2 (CD45.2) BM

cells (Fig 7A). In both instances we found that all γδT17 cells in the pLNs were from wildtype

(CD45.1) BM; conversely, CD27$^+$ γδ T cells were either equally split into CD45.1 and CD45.2

components in mice that had received mixed BM chimeras or only CD45.1 cells in mice that had

received wildtype cells only (Fig 8B). In the siLP, almost all γδ T cells came from wildtype

(CD45.1) BM and expressed various levels of Tbet and some RORγt (Fig 7C). Although ILC3

that were of both wildtype (CD45.1) and ΔcIAP1/2 (CD45.2) origin could be found in the siLP,

wildtype ILC3 cells were more abundant than ILC3 cells from ΔcIAP1/2 BM (Fig 7D). Thus, the

presence of wildtype hematopoietic cells in ΔcIAP1/2 mice is not sufficient to reverse the loss of

γδT17 cells lacking cIAP1 and cIAP2 but is sufficient to reverse the loss of ILC3 cells.

As γδT17 cells are generated perinatally in the thymus, we thus reasoned that γδT17 cells

generated after BM chimera reconstitutions might be different from γδT17 cells that egress the

neonatal thymus. To address this issue, we purified γδ T cells from the thymi of 1-day old Ctrl,

ΔcIAP1, ΔcIAP2 or ΔcIAP1/2 mice and transferred them to RAG1$^{-/-}$ recipient mice (Fig 8A).

After 12 weeks we analyzed the pLNs and cLP and found that γδT17 and CD27$^+$ γδ T cells could

be recovered from the pLNs of RAG1$^{-/-}$ mice (Fig 8B) but again only Tbet$^+$ γδ T cells could be

found in the cLP, regardless of the donor strain (Fig 8C).

Taken together our data show that γδT17 cells that lack cIAP1 and cIAP2 are intrinsically

defective in their response to an extrinsic stimulus that is altered in the ΔcIAP1/2 mice.

Conversely, ILC3 cells were intrinsically less dependent on the presence of cIAP1 and cIAP2 as

they could be rescued by the presence of wildtype hematopoietic, stromal compartment or both.

RORγt^{cre} cIAP1$^{f/f}$ cIAP2$^{-/-}$ mice are not resistant to IMQ-psoriasis.

The TLR7 ligand imiquimod (IMQ) induces a psoriasis-like pathology when applied to the

skin of mice; the development of disease symptoms is dependent on the presence of functional

γδT17 that produces IL-17A in response to increased IL-23 and IL-1β in the skin milieu (van der

Fits *et al*, 2009; Sandrock *et al*, 2018). As such, we aimed to test how the homeostatic reduction

in γδT17 in the skin of ΔcIAP1/2 affects the outcome of skin inflammatory conditions as

psoriasis. Thus, we applied IMQ on the ear skin of littermate controls and cIAP1 and 2 single or

double deficient mice and measured the inflammatory response in the ear skin, auricular and

cervical lymph nodes. Hematoxylin and Eosin (H&E) staining revealed comparable thickening

of the skin epidermis in all strains that received IMQ treatment (Fig S3A). Nevertheless, the

abundance of γδT17 cells in skin or lymph nodes of IMQ treated ΔcIAP1/2 remained

significantly reduced compared to treated or untreated littermate controls or cIAP1 or 2 deficient

mice (Fig S3B). Likewise, IL-17A production by lymph nodes γδ T cells was severely reduced in

both treated and untreated ΔcIAP1/2 mice compared to other strains (Fig S3C). Conversely, we

noticed an increase in IL-17$^+$ CD4$^+$ CD44$^+$ T cells in the lymph nodes of IMQ treated ΔcIAP1/2

mice compared to treated or untreated littermate controls (Fig S3D)

We then sought to assess how the ΔcIAP1/2 mice would respond to IMQ-induced

inflammation in the presence of transcriptionally altered γδT17 cells. To this end, we applied

IMQ on the ear skin of 4 weeks old mice of all 4 genotypes and again assessed the inflammatory

response. We found that the epidermal thickening in the ear skin of ΔcIAP1/2 mice was

comparable to that of WT mice (Fig 9A). Further, we found that both in the skin and the lymph

nodes, γδT17 cells from ΔcIAP1/2 mice failed to expand in response to IMQ-inflammation and

were significantly less abundant than in the IMQ-treated control mice. Conversely, CD4$^+$ T cells

in skin of IMQ-treated ΔcIAP1/2 mice were significantly more abundant than in control mice

although there was no significant difference in the lymph nodes (Fig 9B). Functionally, we found

that γδT17 cells from IMQ-treated ΔcIAP1/2 produced significantly less IL-17A in the lymph

nodes or skin, compared to IMQ-treated control mice; instead, CD4$^+$ T cells from ΔcIAP1/2

produced copious amounts of IL-17A in both organs compared to controls (Fig 9C-D). Likewise,

lymph nodes γδT17 deficient in cIAP1 and cIAP2 failed to produce any IL-22 following IMQ-

psoriasis, while the CD4$^+$ T cells in the same mice produced significantly more IL-22 than in the

IMQ-treated control mice (Fig 9C); intriguingly, IL-22 production by γδT17 in the skin of IMQ-

treated ΔcIAP1/2 was comparable to that of control mice, yet still there was significantly more

IL-22 produced by the CD4+ T cells in the skin of ΔcIAP1/2 mice following IMQ-treatment,

compared to controls (Fig 9E).

Thus, our data show that although adult ΔcIAP1/2 mice are severely deficient in γδT17 cells

they are not protected from IMQ-induced psoriatic inflammation; similarly, reduced

functionality of γδT17 cells in young ΔcIAP1/2 mice did not protect the mice from IMQ-induced

inflammation. In both instances, the IMQ-induced skin inflammation appears to have been a

function of aggravated Th17 responses. These results are consistent with earlier report showing

that CD4$^+$ T cells are responsible for the IMQ induced psoriasis following loss of γδ T cells

during adult life (Sandrock *et al*, 2018). Further, our data point out that during IMQ-induced

inflammation not only can CD4$^+$ T cells compensate for the lack of γδ T cells, but also for their

compromised functionality.

Discussion

Herein we showed that cIAP1 and cIAP2 are required for the adult homeostasis of γδT17 cells

in peripheral lymph nodes, skin and gut lamina propria. Although cIAP1 and cIAP2 were not

required for the thymic development of γδT17 cells, mice that lacked cIAP2 and conditionally

lacked cIAP1 in RORγt expressing cells suffered progressive reduction in γδT17 cells in the

lymph nodes during late neonatal and adult life. Further, lack of cIAP1 and cIAP2 correlated

with a downregulation of RORγt and cMaf in γδT17 cells that preceded the loss of γδT17 cells.

Although inhibition of RIPK1 led to a partial rescue of γδT17 cells in the lymph nodes, it failed

to halt the downregulation of RORγt in cIAP1 and cIAP2 double deficient γδT17 cells; however,

RIPK1 inhibition did not affect the reduction in RORγt$^+$ Tbet$^+$ γδT cells in the siLP of cIAP1 and

cIAP2 double deficient mice. We further show that the both cell intrinsic and extrinsic factors

contribute to the decreased abundances of γδT17 cells in adult RORγt^{cre} cIAP1$^{f/f}$ cIAP2$^{-/-}$ mice.

Finally, we show that adult RORγt^{cre} cIAP1$^{f/f}$ cIAP2$^{-/-}$ mice were not protected against IMQ-

induced skin inflammation despite the severe reduction in γδT17 in these mice and that the

presence of functionally defective γδT17 cells in younger RORγt^{cre} cIAP1$^{f/f}$ cIAP2$^{-/-}$ was not

sufficient to inhibit IMQ-induced psoriasis.

cIAP1 and cIAP2 are ubiquitously expressed proteins that are frequently upregulated in

cancers (Tamm *et al*, 2000; Wu *et al*, 2010). Downstream TNFR superfamily, they fine tune the

balance between NF-κB activation or induction of cell death. Thus, they are important targets in

the context of cancer therapy, whereby their antagonization can sensitize cancer cells to cell

death or modulate the functions of cytotoxic T cells (Gyrd-Hansen & Meier, 2010; Bai *et al*,

2014) . However, their role in γδ T cells has not been previously addressed. Consensus is that

cIAP1 and cIAP2 are functionally redundant (Mahoney *et al*, 2008); thus, mice deficient in either

proteins display no phenotypic changes. As the genetic knockout of both cIAP1 and cIAP2 is

embryonically lethal (Moulin *et al*, 2012), the RORγt driven deletion of cIAP1 combined with

full deletion of cIAP2 provides a targeted approach to probe their function in RORγt expressing

γδ T cells. The data presented herein reveals a critical, but redundant, role for cIAP1 and cIAP2

in the homeostasis of γδT17 cells.

We have previously shown that antagonization of cIAP1 and cIAP2 in Th17 cells

downregulates *Il17a* and *Maf* through the non-canonical NF-κB transcription factors RelB and

p52 (Rizk *et al*, 2019). Thus, it is plausible that a similar mechanism exists in γδT17 cells; as

such, the lack of cIAP1 and cIAP2 in γδT17 activates the non-canonical NF-κB, which

downregulates *Maf* expression. Consequently, *Rorc* is downregulated as cMaf is required to

maintain its expression in γδT17. Eventually, such cells would progressively perish as cMaf and

RORγt are required for the homeostasis of γδT17 (Barros-Martins *et al*, 2016; Zuberbuehler *et*

al, 2019). The partial rescue of γδT17 in the cIAP1/2 double deficient mice by RIPK1 could be

explained by the fact that cMaf deficiency in γδT17 results in increased apoptosis(Zuberbuehler

et al, 2019), which could be dependent upon RIPK1 .

Nonetheless, several outstanding questions remain to be resolved in such model. Specifically,

the persistence of expression of γδT17 surface markers (CCR6[+] CD44[hi]) despite the significant

downregulation of RORγt and cMaf in cIAP1 and 2 deficient γδT17 cells is intriguing.

Furthermore, such downregulation of RORγt and cMaf occurred in the periphery but not in the

thymus and started at approximately 3 weeks of age. Likewise, the intrinsic and extrinsic factors

contributing to the loss of γδT17 in the cIAP1/2 double deficient mice remain unresolved.

References

Annibaldi A & Meier P (2018) Checkpoints in TNF-Induced Cell Death: Implications in Inflammation and Cancer. *Trends Mol. Med.* **24:** 49–65

Bai L, Smith DC & Wang S (2014) Small-molecule SMAC mimetics as new cancer therapeutics. *Pharmacol. Ther.* **144:** 82–95

Barros-Martins J, Schmolka N, Fontinha D, Pires de Miranda M, Simas JP, Brok I, Ferreira C, Veldhoen M, Silva-Santos B & Serre K (2016) Effector γδ T Cell Differentiation Relies on Master but Not Auxiliary Th Cell Transcription Factors. *J. Immunol.* **196:** 3642–3652

Bertrand MJM, Milutinovic S, Dickson KM, Ho WC, Boudreault A, Durkin J, Gillard JW, Jaquith JB, Morris SJ & Barker PA (2008) cIAP1 and cIAP2 Facilitate Cancer Cell Survival by Functioning as E3 Ligases that Promote RIP1 Ubiquitination. *Mol. Cell* **30:** 689–700

Chien Y-H, Meyer C & Bonneville M (2014) γδ T Cells: First Line of Defense and Beyond. *Annu. Rev. Immunol.* **32:** 121–155

Ebertl G, Litman DR, Eberl G & Litman DR (2004) Thymic origin of intestinal αβ T cells revealed by fate mapping of RORγt+ cells. *Science (80-.).* **305:** 248–251

van der Fits L, Mourits S, Voerman JSA, Kant M, Boon L, Laman JD, Cornelissen F, Mus A-M, Florencia E, Prens EP & Lubberts E (2009) Imiquimod-Induced Psoriasis-Like Skin Inflammation in Mice Is Mediated via the IL-23/IL-17 Axis. *J. Immunol.* **182:** 5836–5845

Gyrd-Hansen M & Meier P (2010) IAPs: from caspase inhibitors to modulators of NF-kappaB, inflammation and cancer. *Nat. Rev. Cancer* **10:** 561–74

Haas JD, Ravens S, Düber S, Sandrock I, Oberdörfer L, Kashani E, Chennupati V, Föhse L, Naumann R, Weiss S, Krueger A, Förster R & Prinz I (2012) Development of Interleukin-17-Producing γδ T Cells Is Restricted to a Functional Embryonic Wave. *Immunity* **37:** 48–59

Heilig JS & Tonegawa S (1986) Diversity of murine gamma genes and expression in fetal and adult T lymphocytes. *Nature* **322:** 836–840

Kadekar D, Agerholm R, Rizk J, Neubauer H, Suske T, Maurer B, Viñals MT, Comelli E, Taibi A, Moriggl R & Bekiaris V (2019) The neonatal microenvironment programs conventional and intestinal Tbet+ γδT17 cells through the transcription factor STAT5. *bioRxiv*: 658542

Klose CSN, Kiss EA, Schwierzeck V, Ebert K, Hoyler T, D'Hargues Y, Göppert N, Croxford AL, Waisman A, Tanriver Y & Diefenbach A (2013) A T-bet gradient controls the fate and function of CCR6-RORγt + innate lymphoid cells. *Nature* **494:** 261–265

Mahoney DJ, Cheung HH, Mrad RL, Plenchette S, Simard C, Enwere E, Arora V, Mak TW, Lacasse EC, Waring J & Korneluk RG (2008) Both cIAP1 and cIAP2 regulate TNF - mediated NF- B activation. *Proc. Natl. Acad. Sci.* **105:** 11778–11783

Mair F, Joller S, Hoeppli R, Onder L, Hahn M, Ludewig B, Waisman A & Becher B (2015) The NFκB-inducing kinase is essential for the developmental programming of skin- resident and IL-17-producing Γδ Tδ cells. *Elife* **4:**

Melo-Gonzalez F & Hepworth MR (2017) Functional and phenotypic heterogeneity of group 3
innate lymphoid cells. *Immunology* **150:** 265–275

Moulin M, Anderton H, Voss AK, Thomas T, Wong WWL, Bankovacki A, Feltham R, Chau D,
Cook WD, Silke J & Vaux DL (2012) IAPs limit activation of RIP kinases by TNF receptor
1 during development. *EMBO J.* **31:** 1679–1691

Papotto PH, Ribot JC & Silva-Santos B (2017) IL-17 + γδ T cells as kick-starters of
inflammation. *Nat. Immunol.* **18:** 604–611

Powolny-Budnicka I, Riemann M, Tänzer S, Schmid RM, Hehlgans T & Weih F (2011) RelA
and RelB transcription factors in distinct thymocyte populations control lymphotoxin-
dependent interleukin-17 production in γδ T cells. *Immunity* **34:** 364–374

Prinz I, Silva-Santos B & Pennington DJ (2013) Functional development of γδ T cells. *Eur. J.*
Immunol. **43:** 1988–1994

Rizk J, Kaplinsky J, Agerholm R, Kadekar D, Ivars F, Agace WW, Wei-Lynn Wong W, Szucs
MJ, Myers SA, Carr SA, Waisman A & Bekiaris V (2019) SMAC mimetics promote NIK-
dependent inhibition of CD4+ TH17 cell differentiation. *Sci. Signal.* **12:**

Sagar, Pokrovskii M, Herman JS, Naik S, Sock E, Lausch U, Wegner M, Tanriver Y, Littman
DR & Grün D (2018) Deciphering the Regulatory Landscape of γδ T Cell Development by
Single-Cell RNA-Sequencing. *bioRxiv***: 478529

Sandrock I, Reinhardt A, Ravens S, Binz C, Wilharm A, Martins J, Oberdörfer L, Tan L,
Lienenklaus S, Zhang B, Naumann R, Zhuang Y, Krueger A, Förster R & Prinz I (2018)
Genetic models reveal origin, persistence and non-redundant functions of IL-17–producing
γδ T cells. *J. Exp. Med.* **215:** 3006–3018

Sawa S, Cherrier M, Lochner M, Satoh-Takayama N, Fehling HJ, Langa F, Di Santo JP & Eberl
G (2010) Lineage Relationship Analysis of RORγt + Innate Lymphoid Cells. *Science (80-.*
). **330:** 665–669

Schmolka N, Serre K, Grosso AR, Rei M, Pennington DJ, Gomes AQ & Silva-Santos B (2013)
Epigenetic and transcriptional signatures of stable versus plastic differentiation of
proinflammatory γδ T cell subsets. *Nat. Immunol.* **14:** 1093–1100

Silke J & Meier P (2013) Inhibitor of apoptosis(IAP)proteins-modulators of cell death and
inflammation. *Cold Spring Harb. Perspect. Biol.* **5:** a008730

Silva-Santos B, Pennington DJ & Hayday AC (2005) Lymphotoxin-mediated regulation of γδ
cell differentiation by αβ T cell progenitors. *Science (80-.).* **307:** 925–928

Sutton CE, Lalor SJ, Sweeney CM, Brereton CF, Lavelle EC & Mills KHG (2009) Interleukin-1
and IL-23 Induce Innate IL-17 Production from γδ T Cells, Amplifying Th17 Responses
and Autoimmunity. *Immunity* **31:** 331–341

Tamm I, Kornblau SM, Segall H, Krajewski S, Welsh K, Kitada S, Scudiero DA, Tudor G, Qui
YH, Monks A, Andreeff M & Reed JC (2000) Expression and prognostic significance of
IAP-family genes in human cancers and myeloid leukemias. *Clin. Cancer Res.* **6:** 1796–803

Vallabhapurapu S, Matsuzawa A, Zhang WZ, Tseng PH, Keats JJ, Wang H, Vignali DAA,
Bergsagel PL & Karin M (2008) Nonredundant and complementary functions of TRAF2
and TRAF3 in a ubiquitination cascade that activates NIK-dependent alternative NF-κB
signaling. *Nat. Immunol.* **9:** 1364–1370

Varfolomeev E, Blankenship JW, Wayson SM, Fedorova A V., Kayagaki N, Garg P, Zobel K,
Dynek JN, Elliott LO, Wallweber HJA, Flygare JA, Fairbrother WJ, Deshayes K, Dixit VM
& Vucic D (2007) IAP Antagonists Induce Autoubiquitination of c-IAPs, NF-κB
Activation, and TNFα-Dependent Apoptosis. *Cell* **131:** 669–681

Wu HH, Wu JY, Cheng YW, Chen CY, Lee MC, Goan YG & Lee H (2010) cIAP2 upregulated
by E6 oncoprotein via epidermal growth factor receptor/phosphatidylinositol 3-kinase/AKT
pathway confers resistance to cisplatin in human papillomavirus 16/18-infected lung cancer.
Clin. Cancer Res. **16:** 5200–5210

Zarnegar BJ, Wang Y, Mahoney DJ, Dempsey PW, Cheung HH, He J, Shiba T, Yang X, Yeh
WC, Mak TW, Korneluk RG & Cheng G (2008) Noncanonical NF-κB activation requires
coordinated assembly of a regulatory complex of the adaptors cIAP1, cIAP2, TRAF2 and
TRAF3 and the kinase NIK. *Nat. Immunol.* **9:** 1371–1378

Zuberbuehler MK, Parker ME, Wheaton JD, Espinosa JR, Salzler HR, Park E & Ciofani M
(2019) The transcription factor c-Maf is essential for the commitment of IL-17-producing
γδ T cells. *Nat. Immunol.* **20:** 73–85

Materials and methods

Mice

All animals were bred and maintained in-house at DTU health tech with the approval of the Danish animal experiments inspectorate. cIAP1$^{f/f}$ and cIAP1$^{f/f}$ cIAP2$^{-/-}$ mice were provided by Prof. W. Wei-Lynn Wong at the University of Zurich, Switzerland with the permission of Prof. John Silke, VIC Australia. RORγt^{CRE} mice were provided by Prof. Gerard Eberl at Pasteur Institute, Paris, France.

Cell culture media and buffers

For all preparations of single cell suspensions and cell cultures RPMI 1460 (Invitrogen) supplemented with 10% heat inactivated FBS (GIBCO), 20mM Hepes pH 7.4 (Gibco), 50 µM 2-mercaptoethanol, 2 mM L-glutamine (Gibco) and 10,000 U/ mL penicillin-streptomycin (Gibco), was used. FACS buffer was prepared by supplementing PBS with 3% heat inactivated FBS.

Lymphocyte isolation from mouse organs

Lymphocytes were isolated from peripheral lymph nodes (axial, brachial and inguinal), thymus, ear skin, small intestinal and colonic lamina propria following the previously described protocols (Kadekar *et al*, 2019). Lymphocytes were isolated from cervical and auricular lymph nodes in case of IMQ-induced psoriasis.

IMQ-induced psoriasis

Psoriasis was induced in mice by applying 7 mg of Aldara cream (containing 5% imiquimod) to the dorsal side of each ear for 7 days. Histological sections were prepared by fixing ear tissue in 10% formalin overnight and then paraffin embedded. The paraffin embedded section were stained by H&E.

Flow cytometry staining

Surface antigens, intracellular cytokines and transcription factors were stained for flow

cytometry as previously described (Rizk *et al*, 2019). The following antibodies were used herein

at 1:200 dilution unless otherwise indicated:

anti-IL-17A (TC11-18H10; BV786 and PE), anti-IFNγ (XMG1.2; PE-Cy7 and APC), anti-IL-22

(1H8PWSR; PE), anti-cMAF (symOF1; PE and Percp-Cy5.5; 5 μL/test), anti-CD4 (GK1.5;

BUV395), anti-TCRγδ (GL3; BV421), anti-CD27 (LG.3A10; PE-Cy7), anti-CCR6 (140706;

Alexa Fluor 647), anti-CD44 (1M7; V500), anti-CD19 (6D5; FITC), anti-TCRβ (H57-597; APC-

eflour780), anti-CD3e (BM10-37, PeCF594 and PE), anti-Tbet(4B10; PeCy7), anti- CD8 (53-

6.7;FITC), anti-Vγ5 (536; FITC), anti-Vγ4(UC3-10A6; Percp-eflour710), anti-GATA3(TWAJ;

Percp-eFlour710; 1:30), anti-CD45(30-F11;PE and V500) and anti-RORγt (B2D; APC).

Retroviral infection of neonatal γδT17 cells.

First, pooled thymi from one-day old wildtype mice were isolated and crushed against 70 μm

filter to prepare single cell solution. Subsequently, γδT17 cells were enriched by magnetic

depletion of CD4$^+$, CD8$^+$, TCRβ$^+$ and CD27$^+$ cells as follows: total thymocytes were re-

suspended in MACS buffer at 1e8 cells/ml containing 50 μL/ml normal rat serum and 1:200

biotin labelled anti- CD4$^+$, CD8$^+$, TCRβ$^+$ and CD27$^+$ antibodies; the cells were incubated for 10

minutes at room temperature and then incubated with 75 μL/ml EasySep RaphidSphere

streptavidin beads (#50001) for 2.5 minutes then transferred to EasySep magnet for 2.5 minutes.

The non-bound fraction was collected by decantation and centrifugated for 5 minutes at 400g at

4ºC. Approximately 2e5 of the enriched γδT17 cells were then cultured overnight with 20 ng/ml

recombinant IL-7 in round-bottom 96-well tissue culture plate. The cells were then spin-infected

460 by adding 4 μL of the desired retrovirus and 5 μg Retronectin (TAKARA) and centrifugation at

461 1800 rpm for 30 minutes at 32°C with deceleration set to 0. Subsequently, the cells were

462 incubated for 90 minutes at 32°C before transfer to 37°C incubator. Recombinant IL-7 at 20

463 ng/ml was again added to the cultures after 48 hours and the cells were FACS sorted 24 hours

464 later.

465 **Transfer of neonatal γδ T cells to RAG1$^{-/-}$ hosts.**

466 First, thymi from 1-2 days old mice, from all 4 strains, were isolated and crushed individually

467 against 70 μm filter to prepare single cell solutions. Subsequently, total γδ T cells were enriched

468 by magnetic depletion of CD4$^+$, CD8$^+$, TCRβ$^+$ cells as aforementioned. The enriched γδ T cells

469 from each donor mouse were re-suspended in PBS and then i.v. injected into the tail vein of a

470 RAG1$^{-/-}$ host. The RAG1$^{-/-}$ hosts were euthanized for organs after 12 weeks.

471 **Bone marrow chimeras**

472 The bone marrow cells for reconstitution were isolated by flushing the tibia and femur, which

473 were dissected from donor mice, with culture media. Total bone marrow cells were then

474 centrifugated at 400g for 5 minutes at 4°C. The cells were then re-suspended and passed through

475 70 μm filter. Subsequently, red blood cells were then lysed using RBC lysis buffer (Biolegend)

476 and a single cell suspension of bone marrow cells was the prepared by passing the cells through

477 40 μm filter. The prepared cells were then counted and mixed as appropriate.

478 Conversely, host mice were sub-lethally irradiated by 2 doses of 4.5 Gy that were at least 4 hours

479 apart. After 24 hours, the hosts were reconstituted with 10e6 bone marrow cells that were i.v.

480 injected into the tail vein of the host mice. The hosts were euthanized for organs after at least 12

481 weeks.

482 **Figures**

483 **Figure 1**

A

Ctrls	RORγt Cre⁻ cIAP1^f/f
ΔcIAP1	RORγt Cre⁺ cIAP1^f/f
ΔcIAP2	RORγt Cre⁻ cIAP1^f/f cIAP2⁻/⁻
ΔcIAP1/2	RORγt Cre⁺ cIAP1^f/f cIAP2⁻/⁻

B

C

Figure 1. cIAP1 and cIAP2 are required for the homeostasis of γδT17 cells in pLNs and skin.

(A) Mice strains used herein were generated by crossing RORγt^{CRE} strain with cIAP1$^{f/f}$ or cIAP1$^{f/f}$cIAP2$^{-/-}$ strains; the resulting 4 strains were as follows RORγt^{CRE} cIAP1$^{f/f}$ (Cre$^-$: ctrls, Cre$^+$: ΔcIAP1) and RORγt^{CRE} cIAP1$^{f/f}$cIAP2$^{-/-}$ (Cre$^-$: ΔcIAP2, Cre$^+$: ΔcIAP1/2). (B) Flow cytometric analysis plots (left) and quantification (right) of TCRγδ$^+$ populations in the pLNs of all 4 strains. γδT17 cells were identified as CD44^{hi} CD27$^-$ TCR γδ$^+$. (C) Flow cytometric analysis plots (left) and quantification (right) of TCRγδ$^+$ and TCRβ$^+$ in the skin of all 4 strains. In the skin, γδT17 cells were identified as CD45$^+$ Vγ5$^-$ CD3$^+$ TCRγδ$^+$. In graphs, each symbol represents a mouse and line the mean. **p < 0.01, ****p < 0.0001 using Mann-Whitney test.

Figure 2

Figure 2. cIAP1 and cIAP2 are required for the homeostasis of intestinal RORγt⁺ Tbet⁺ γδ T and ILC3 cells.

Flow cytometric analysis plots (left) and quantification (right) of TCRγδ$^+$ populations in the siLP (A) or cLP(B) of all 4 strains. γδT cell populations were identified as CD45$^+$ CD3$^+$ TCRβ$^-$ TCRγδ$^+$. (C) Flow cytometric analysis plots of (left) and quantification (right) of ILC3 cells in the siLP of all 4 strains. ILC cells were identified as CD45^{int} CD3$^-$ CD127$^+$ TCRγδ$^-$ TCRβ$^-$ cells. In graphs, each symbol represents a mouse and line the mean. *p < 0.05, **p < 0.01, ***p < 0.001,   ****p < 0.0001 using Mann-Whitney test.

Figure 3

Figure 3. cIAP1 and cIAP2 are not required for thymic development of γδT17 cells but are
required for cMaf and RORγt expression during neonatal and adult life.
(A) Numbers of γδT17 cells in the day1 (d1) thymi of all 4 strains (gated as in Fig1). (B) Flow
cytometric analysis of RORγt and cMaf expression in γδ T cells from d1 thymi of all 4 strains.
(C) Numbers of γδT17 cells in the pLNs at the indicated timepoints. (D) analysis of γδ T cell
populations from d21 pLNs of the indicated strains, while arrows show RORγt and cMaf
expression within γδT17 cells compartment. (E) frequency of RORγt and cMaf expression in
γδT17 in the pLNs of 12-weeks old from all strains. In graphs each symbol represents a mouse
and line the mean. *$p < 0.05$ using Mann-Whitney test.
521
522

 Figure 4

Figure 4. Inhibition of RIPK1 partially rescues the reduction in γδT17 in ΔcIAP1/2 mice but not the downregulation of RORγt

(A) Flow cytometric analysis plots (left) and quantification (right) of TCRγδ$^+$ populations in the pLNs of indicated strains that received regular chow (control) or chow that contained RIPK1 inhibitor (Ri). γδT17 cells were identified by surface phenotyping as CD44^{hi} CD27$^-$ TCRγδ$^+$. (B) Flow cytometric analysis plots showing TCRγδ$^+$ populations in the pLNs of littermate controls and ΔcIAP1/2 that received control chow or Ri chow; overlaid is RORγt$^+$ (red), Tbet$^+$ (blue) or RORγt$^-$ Tbet$^-$ (black) cells within the γδT cell compartment. (C) Histogram (upper) and quantification (lower) of the mean fluorescence intensity (MFI) of RORγt in CD44^{hi} CD27$^-$ TCRγδ$^+$ in the pLNs of the indicated strains that received control or Ri chow. (D) Flow cytometric analysis plots (left) and quantification (right) of TCRγδ$^+$ populations in the siLP of the indicated strains that received control or Ri chow. In graphs each symbol represents a mouse and line the mean. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001 using Mann-Whitney test.

Figure 5

Figure 5. Activation of the non-canonical NF-κB pathway downregulates the expression of
***Maf* in γδT17 cells.**
(A-B) RT-qPCR analysis of the indicated genes in γδT17 cells that were infected with empty
retrovirus (pMIG) or retrovirus expressing RelB. only GFP+ cells (retrovirus infected cells) were
FACS sorted and analyzed. In graphs each symbol represents an independent experiment.

Figure 6

Figure 6. Cell extrinsic signals mediate the loss of cIAP1 and cIAP2 double deficient γδT17
and ILC3 cells in ΔcIAP1/2 mice.
(A) schematic representation of the experimental design whereby sub-lethally irradiated wildtype
hosts (CD45.1/45.2) were reconstituted with 1:1 mixture of WT (CD45.1) and ΔcIAP1/2
(CD45.2) bone marrow. (B-C) Flow cytometry analysis of γδT cells in the pLNs and cLP of host
mice after 12 weeks of reconstitution and indicating the frequency of WT, host, or ΔcIAP1/2
γδT17 and CD27[+] γδ T cells. Graphs show frequency of WT, host, or ΔcIAP1/2 in γδ T cell
populations in pLN of host mice amalgamated from 2 independent experiments. (D) Flow
cytometry analysis of ILC3 cells in the cLP of host mice. Graph show frequency of WT, host, or
ΔcIAP1/2 of all ILC3 cells in cLP of host mice amalgamated from 2 independent experiments. In
graphs each symbol represents a mouse and line the mean.

Figure 7

Figure 7. Reconstitution of ΔcIAP1/2 mice with wildtype hematopoietic system rescues the
loss of ILC3 cells but not γδT17 cells.
(A) schematic representation of the experimental design whereby sub-lethally irradiated
ΔcIAP1/2 (CD45.2) were reconstituted with either WT (CD45.1) or 1:1 mixture of WT (CD45.1)
and ΔcIAP1/2 (CD45.2) bone marrow. Flow cytometry analysis of γδT cells in the pLNs (B) and
siLP (C) of host mice after 12 weeks of reconstitution and indicating the frequency of WT or
ΔcIAP1/2 γδT17 and CD27$^+$ γδT cells. Graphs show frequency of WT or ΔcIAP1/2 in γδ T cell
populations in pLN and siLP. (D) Flow cytometry analysis of ILC3 cells in the cLP of host mice.
Graph show frequency of WT or ΔcIAP1/2 of all ILC3 cells in siLP. In graphs each symbol
represents a mouse and line the mean.

 Figure 8

Figure 8. Neonatal γδT17 lacking cIAP1 and cIAP2 can reconstitute RAG⁻/⁻ hosts.
(A) schematic representation of the experimental design whereby γδ T cells were enriched from
the thymi from one-day old mice of all 4 strains and i.v. injected into RAG⁻/⁻ hosts. (B) Flow
cytometry analysis of γδT cells in the pLNs of host RAG⁻/⁻ mice at 12 weeks after transfer of the

γδT cells from the indicated strain. Graphs show frequency of recovered γδT17 or CD27$^+$ γδT

cells of total γδT cells. In graphs each symbol represents a donor mouse and line the median.

Data is amalgamated from 3 independent experiments. (C) Flow cytometry analysis of γδT cells

in cLP of host RAG$^{-/-}$ mice at 12 weeks after transfer of the γδT cells from the indicated strain.

Figure 9

Figure 9. cIAP1 and 2 double deficient γδT17 cells are functionally defective and do not respond to IMQ-induced psoriasis.

(A) Representative light microscopy images (right) indicating ear skin sections stained for H&E before and after IMQ-induced psoriasis in the indicated strains (left) Quantification of epidermal

thickening from H&E stained sections. (B) Numbers of γδT17 and CD4$^+$ TCRβ$^+$ cells in the pLN or skin of all strains of IMQ treated or control mice. In the pLN, γδT17 were identified as CD44^{hi} CD27$^-$ TCRγδ$^+$, while in the skin, they were identified as CD45$^+$ Vγ5$^-$ CD3$^+$ TCRγδ$^+$. Quantification of IL-17A (C, D) or IL-22 (C, E) production by γδ T and CD4 T cells in the pLN (C) or skin (E) in control or IMQ-treated mice of all strains. In graphs, each symbol represents a mouse and line the mean. Data is pool of 3 experiments. *p < 0.05, **p < 0.01using Mann-Whitney test.