The cIAP ubiquitin ligases sustain type 3 γδ T and innate lymphoid cells during aging to allow normal cutaneous and mucosal responses

John Rizk[1], Urs M. Mörbe[1], Rasmus Agerholm[1], Isabel Ulmert[1], Elisa Catafal Tardos[1], Darshana Kadekar[1], Maria V. Baglioni[1], Monica Torrellas Viñals[1] and Vasileios Bekiaris*[1]

[1]Department of Health Technology, Technical University of Denmark, Kemitorvet, Bldg 202, 2800 Kgs Lyngby, Denmark.

*Corresponding author: Vasileios Bekiaris vasbek@dtu.dk

Abstract

Environmental and molecular cues early in life are often associated with the permanent shaping of our immune system during adulthood. Although increasing, our knowledge of the signaling pathways that operate in early life and their temporal mode of action is limited. Herein, we demonstrate that the cellular inhibitor of apoptosis proteins 1 and 2 (cIAP1/2), which are E3 ubiquitin ligases and master regulators of the nuclear factor-kappa B (NF-κB) pathway, function during late neonatal and prepubescent life to sustain interleukin(IL)-17-producing gamma delta T cells (γδT17) and group 3 innate lymphoid cells (ILC3). We show that cell-intrinsic deficiency in cIAP1/2 at 3-4 weeks of life leads to downregulation of the transcription factors cMAF and RORγt, and failure to enter cytokine-induced cell cycle. This is followed by progressive loss of γδT17 cells and ILC3 while mice are aging. Mice deficient in cIAP1/2 have severely reduced γδT17 cells and ILC3, present with suboptimal γδT17 responses in the skin, lack small intestinal isolated lymphoid follicles and cannot control intestinal bacterial infection. Mechanistically, these effects appear to be dependent on overt activation of the non-canonical NF-κB pathway.

Our data identify the cIAP E3 ubiquitin ligases as critical early life molecular switches for

establishing effective type-3 immunity during aging.

Introduction

The neonatal period is the time when our immune system is imprinted with life-long

functional characteristics that maintain immunity to infection and prevent autoimmune

pathology. Microbial colonization, and developmentally regulated transcriptional

programs cooperate to shape innate and adaptive lymphocytes into distinct specialized

lineages that co-exist in equilibrium and respond ad hoc (Eberl, 2016). Failure to convey

these environmental and molecular cues during neonatal life, often results in irreversible

dysfunction later on. Hence, early dysbiosis impairs type-3 immunity and potentiates

susceptibility to type-2 driven allergy (Cahenzli *et al*, 2013) . Similarly, blockade of key

signaling pathways during neonatal life can permanently change cellular niches

(Kadekar *et al*, 2020). Therefore, elucidating the molecular signatures that operate early

in life is of great importance for understanding how immunity develops.

Mouse γδ T cells present a well-established example of an immune population that

is heavily dependent on an unperturbed neonatal period. In this regard, intestinal

intraepithelial (IE) γδ T cells develop during neonatal and prepubescent life through

butyrophylin-driven interactions with the epithelia (Di Marco Barros *et al*, 2016) . This

provides a necessary defense mechanism against infection within the IEL compartment

(Hoytema van Konijnenburg *et al*, 2017). Lamina propria (LP) interleukin(IL)-17-

producing γδ T cells establish mixed type-3 and type-1 transcriptional programs within

the first week of life through the transcription factor STAT5 (Kadekar *et al*, 2020) . Thus,

early life establishment of the γδT17 compartment is critical to protect from neonatal

and adult infections (Chen *et al*, 2020; Sheridan *et al*, 2013). In a similar manner,

impaired microbial colonization of the ocular or oral mucosa results in drastically altered

IL-17-producing γδ T (γδT17) cell numbers in the conjunctiva (St. Leger *et al*, 2017) and

cervical lymph nodes (LN) (Fleming *et al*, 2017). Again, paucity in such γδT17 cell

populations is associated with impaired anti-microbial responses in eye and oral cavity

(Conti *et al*, 2014; St. Leger *et al*, 2017), and resistance to pathogenic inflammation (Cai

et al, 2011; Sandrock *et al*, 2018; McGinley *et al*, 2020).

The innate lymphoid cell (ILC) compartment is also dependent on early life events,

while their function during the neonatal period is critical for the establishment of the

intestinal immune system (Spits *et al*, 2013). In this regard, although dysbiosis does not

affect ILC development, it results in altered transcriptional and epigenetic profiles of all

ILC subsets (Gury-BenAri *et al*, 2016). Similar to LP γδT17 cells, group 3 ILC (ILC3)

acquire expression of the transcription factor Tbet and the type-1 cytokine interferon-γ

(IFN-γ) during neonatal life, which allows them to clear intracellular bacterial infections

(Klose *et al*, 2013) . Moreover, group 2 ILC (ILC2) undergo an IL-33-dependent

maturation step in the neonatal lung, allowing their cytokine responsiveness in adult

mice (Steer *et al*, 2020). Importantly, ILC3 induce the maturation of intestinal

cryptopathces into isolated lymphoid follicles (ILFs) during the first 3-4 weeks of life

(Kiss *et al*, 2011; Kruglov *et al*, 2013). Evidently, perturbations of γδT17 and ILC3

development during the early stages of life will have a substantial impact on the quality

of immunity while aging. The molecular pathways that control the transition of these

cells from neonatal life to adolescence and adulthood are poorly understood.

The E3 ubiquitin ligases cellular inhibitor of apoptosis protein (cIAP)1 and 2

(cIAP1/2) catalyze both degradative lysine(K)-48 and stabilizing K-63 ubiquitination and

act as the main molecular switches for the activation of the canonical and non-canonical

nuclear factor-kappa B (NF-κB) pathway (Silke & Meier, 2013) . The presence of

cIAP1/2 downstream of TNF receptor 1 (TNFR1) determines whether a cell will initiate

the canonical NF-κB pathway or die by apoptosis or necroptosis in response to TNF

(Annibaldi & Meier, 2018). They achieve this by ubiquitinating receptor interacting

kinase-1 (RIPK1) (Silke & Meier, 2013) . However, cIAP1/2 are mostly recognized as

negative regulators of the non-canonical NF-κB pathway. Hence, in all cell types

cIAP1/2 associate in a heterocomplex with TNF receptor associated factor (TRAF)2,

TRAF3 and NF-κB-inducing kinase (NIK), whereby they induce K-48 ubiquitination of

NIK, resulting in its continuous proteasomal degradation (Zarnegar *et al*, 2008;

Varfolomeev *et al*, 2007; Vince *et al*, 2007). Breakdown of the TRAF2-TRAF3-cIAP1/2-

NIK complex either following ligation of TNF superfamily receptors that recruit TRAF2-

TRAF3 in their intracellular domain (e.g. TNFR2, LTβR, CD40) or by cIAP1/2 depletion,

liberates NIK, which initiates the cascade necessary for nuclear translocation of the

non-canonical NF-κB transcription factors RelB and p52 (Vallabhapurapu *et al*, 2008;

Matsuzawa *et al*, 2008)

In the present study we demonstrate a necessary role for cIAP1/2 in sustaining

γδT17 cells and ILC3 at the late neonatal and prepubescent stages of life, and thus

impacting the magnitude of inflammatory and anti-bacterial immune responses.

Deficiency in cIAP1/2 begun to have an impact only during late neonatal life by reducing

expression of the lineage defining transcription factors cMAF and RORγt, which was

followed by an apparent block in cytokine-induced proliferation. When animals entered

prepubescence and early adolescence, cIAP1/2 deficiency resulted in progressive loss

γδT17 cells. This was independent of TNFR1 induced canonical NF-κB or cell death. In

contrast, cIAP1/2-deficient prepubescent γδT17 cells displayed enhanced nuclear

translocation of RelB, which demonstrates evidence of overt activation of the non-

canonical NF-κB pathway. Intestinal ILC3 also relied on intact cIAP1/2 during the same

time period, with their numbers being drastically reduced in adulthood. Paucity in ILC3

coincided with ILF involution. Mice with targeted deletion of cIAP1/2 in γδT17 cells and

ILC3 responded sub-optimally to cutaneous inflammatory challenge and failed to control

intestinal bacterial infection.

Results

Paucity of γδT17 cells in the absence of the E3 ubiquitin ligases cIAP1 and cIAP2

Using acute, SMAC mimetic (SM) driven antagonization and in vitro techniques, we

showed before that cIAP1/2 are important for T_H17 differentiation through modulation of

the non-canonical NF-κB pathway (Rizk *et al*, 2019). In order to understand the in vivo

importance of cIAP1 and cIAP2 in RORγt-expressing immune cells, we crossed *Rorc*-

Cre (RORγt^CRE) mice (Eberl & Litman, 2004) with mice that were floxed for *Birc2*

(cIAP1^F/F) and knocked out for *Birc3* (cIAP2^-/-) (Gardam *et al*, 2011). This generated

mice with RORγt-driven deletion of cIAP1 (referred to thereafter as ΔIAP1) and

generalized deletion of cIAP2 (referred to thereafter as ΔIAP2), as well as the

corresponding Cre-negative littermate controls (wild-type; WT) (Fig 1A). ΔIAP1 and

ΔIAP1/2 mice were viable, produced offspring at expected rates, and did not develop

any observable spontaneous disease phenotypes. They contained a full set of lymph

nodes (inguinal, brachial, axillary and mesenteric) and Peyer's patches indicating

unperturbed lymphoid tissue development, while total numbers of CD4$^+$ T and B cells

were normal but γδ T were slightly elevated (Fig S1A).

We next analyzed some of the major IL-17-producing populations in lymph node

(LN) and small intestinal and colonic lamina propria (siLP; cLP). Compared to littermate

controls, ΔIAP1/2 mice produced slightly elevated levels of IL-17A within the

CD4$^+$TCRβ$^+$ compartment in the LN (pool of inguinal, brachial, axillary) but not the gut

(Fig S1B), suggesting that in these animals, steady-state production of IL-17A by CD4$^+$

T cells is not defective. Staining for IL-22 following overnight stimulation with IL-23

yielded the same answer (Fig S1C). However, there was a marked reduction in γδ-

associated IL-17A and IL-22 production in LN (Fig 1B) and IL-17A in the gut (Fig 1C).

This was accompanied by a dramatic loss in LN TCRγδ$^+$CD27$^-$CD44^{hi}CCR6$^+$, which are

the IL-17-producing γδ T cells (Ribot *et al*, 2009; Haas *et al*, 2009) (Fig 1D and Fig

S1D), and gut Tbet$^+$RORγt$^+$ (Kadekar *et al*, 2020) (Fig 1E) γδT17 cell numbers.

Although cIAP1 and cIAP2 individually did not contribute to this phenotype in the LN,

ΔIAP2 mice had significantly reduced Tbet$^+$RORγt$^+$ γδ T cell numbers in the gut (Fig

1E). We additionally found significantly reduced γδT17 cells in the lungs of ΔIAP2 mice

(Fig S1E). Similar to γδT17 cells, there were significantly reduced non-CD4 IL-17-

producing lymphocytes in the LNs of ΔIAP1/2 mice (Fig S1F).

In the skin, CD3^{lo}Vγ5$^-$TCRγδ$^+$CCR6$^+$ cells, which represent the γδT17 population

(Haas *et al*, 2009, 2012), were also reduced significantly in the absence of cIAP1 and

cIAP2 (Fig 2A). When we analyzed the two major γδT17 subpopulations (Vγ4$^+$ versus

Vγ4$^-$; Vγ nomenclature by Heilig and Tonegawa; (Heilig & Tonegawa, 1986)), we found

that in the skin cIAP1 but not cIAP2 was required for Vγ4$^-$ cells, whereas the Vγ4-

expressing population was only affected by the absence of both cIAP1 and cIAP2 (Fig

2B). In the LN, we did not observe differential regulation of either Vγ4$^+$ or Vγ4$^-$ cells (Fig

S2A). Collectively, this data suggests that cIAP1/2 are important for the development

and/or homeostatic maintenance of γδT17 cells. Our findings additionally pinpoint a

differential and non-redundant role of cIAP1 and cIAP2 in these cells that is organ and

subset specific. In this regard, whereas skin γδT17 cells depended more on cIAP1, gut

γδT17 cells depended more on cIAP2.

Cell-intrinsic requirement for cIAP1 and cIAP2 in γδT17 cells

Next, we investigated whether the defect we observed in ΔIAP1/2 mice was cell-

intrinsic. To this end we set up mixed bone marrow (BM) chimeras where WT

CD45.1$^+$CD45.2$^+$ hosts were sub-lethally irradiated and reconstituted with a mixture of

1:1 CD45.1$^+$ WT and CD45.2$^+$ ΔIAP1/2 BM cells (Fig 3A). We found that, under these

conditions, WT LN γδT17 cells outcompeted their ΔIAP1/2 counterparts (Fig 3B),

indicating the phenotype we observed in intact mice was cell intrinsic. Interestingly,

CD27$^+$ γδ T cells derived from ΔIAP1/2 BM were slightly less competitive than WT (Fig

3B). In contrast, both CD3$^-$ populations and B cells from ΔIAP1/2 BM were more

competitive than their WT counterparts (Fig 3C). This indicated that the reduced

competitiveness of γδT17 and CD27$^+$ γδ T cells was not due to defective ΔIAP1/2 BM

reconstitution. We could not reconstitute gut RORγt$^+$Tbet$^+$ γδT17 cells irrespective of

the BM source (Fig 3D), suggesting that this population requires either thymus-

originated γδ T cells or a neonatal microenvironment to develop fully. In contrast, lack of

cIAP1 and cIAP2 did not impinge on the reconstitution of gut Tbet$^+$RORγt$^-$ γδ T cells

(Fig 3E). Likewise, we could only recover WT LN γδT17 cells when we reconstituted

ΔIAP1/2 hosts with WT or a 1:1 mix of WT and ΔIAP1/2 BM (Fig S3A-B), while CD27$^+$

γδ T cells from WT or ΔIAP1/2 BM cells were equally competitive (Fig S3B). As before,

we could not reconstitute RORγt⁺Tbet⁺ γδT17 cells in the gut (Fig S3C).

As γδT17 cells develop perinatally in the thymus and undergo a rapid neonatal re-

programming within the tissues they localize at, we reasoned that if generated from BM

stem cells, they might have different developmental or homeostatic requirements for

cIAP1 and cIAP2. To address this issue, we purified γδ T cells from the thymi of 1-day

old WT or ΔIAP1/2 mice and transferred them to RAG1$^{-/-}$ recipients (Fig 3F). We found

that 12 weeks post transfer, the γδT17 cell compartment was reconstituted in the LN,

however, we recovered significantly more WT than ΔIAP1/2 cells (Fig 3G). As with the

BM chimeras, we could not reconstitute intestinal RORγt⁺Tbet⁺ γδT17 cells, suggesting

that this population requires a neonatal microenvironment (Fig S3D). Reconstitution of

CD27⁺ γδ T cells was independent of cIAP1 and cIAP2 (Fig 3G). Taken together our

data show that γδT17 cells require cIAP1 and cIAP2 intrinsically.

The impact of cIAP1 and cIAP2 on γδT17 cells is independent of TNF induced

canonical NF-κB and cell death

In addition to preventing spontaneous activation of the non-canonical NF-κB pathway,

cIAP1/2 are necessary to convey the canonical NF-κB downstream of TNFR1 whereas

in their absence, TNF-TNFR1 interactions can lead to RIPK1-mediated cell death via

apoptosis or necroptosis (Annibaldi & Meier, 2018). Mice deficient in TNFR1 had an

intact γδT17 cell population (Fig S4A), suggesting that the canonical NF-κB pathway

downstream of TNFR1 is not responsible for the phenotype of ΔIAP1/2 mice. Since TNF

is highly upregulated during the weaning reaction (Al Nabhani *et al*, 2019), we next

investigated whether TNF induced cell death played a role. To achieve this, we initially

analyzed mice that were deficient in cIAP2 and expressed a ubiquitin-associated (UBA)

domain mutant form of cIAP1 unable to K48 ubiquitylate and suppress RIPK1 (Annibaldi

et al, 2018). Thus, these mice are more sensitive to TNF induced cell death (Annibaldi

et al, 2018). In UBA-mutant mice, γδT17 cells were not affected (Fig S4B), suggesting

that these cells are not susceptible to death by homeostatic levels of TNF. In order to

test this directly in ΔIAP1/2 mice, we began injecting 1-week old neonates with

neutralizing anti-TNF antibody and until animals were 12-week old (Fig S4C). We could

not rescue the γδT17 population in either gut or LNs (Fig S4D-E), indicating that TNF

induced death is unlikely to play a major role in regulating these cells in the absence of

cIAP1/2. Therefore, TNF-TNFR1 interactions are not responsible for the ΔIAP1/2

phenotype, suggesting that overt activation of the non-canonical NF-κB pathway could

play a key role.

cIAP1 and cIAP2 are required for γδT17 cell cycle progression and expression of

cMAF and RORγt during aging

In order to assess the impact of cIAP1/2 on embryonic γδT17 cell development, we

enumerated thymic cell numbers in newborn ΔIAP1/2 mice and found them similar to

littermate controls (Fig 4A), despite efficient deletion of *Birc2* (cIAP1) (Fig S5A).

Production of IL-17A/F were unchanged at this stage (Fig S5B). This suggested that the

major impact of cIAP1/2 occurs post-embryonically. We therefore tracked LN γδT17

cells, defined phenotypically as CD27$^-$CD44^{hi}, during neonatal, post-neonatal (average

weaning time at 3 weeks) and adult life (mating age of 8 weeks). We did not find any

differences in cell numbers until week 5 of age (Fig 4B). This suggested that cIAP1 and

cIAP2 are only required to sustain γδT17 numbers following weaning. After week 5,

ΔIAP1/2 γδT17 cells failed to expand and began to decline progressively during aging

(Fig 4B). In order to confirm that the cells are missing from adult life and have not

converted to a non-γδT17 population, we crossed ΔIAP1/2 with the ROSA26-LSL-RFP

strain, so that RFP permanently marks all current and "ex" RORγt-expressing cells. We

found no evidence of γδT17 conversion to other populations (Fig 4C). This data

suggested that cIAP1/2 regulate a checkpoint in early adult life that manifests during

aging.

We next investigated what this checkpoint was. The inability of the cells to increase

in numbers during aging, raised the hypothesis that cIAP1/2 may be regulating

responsiveness to cytokines that induce proliferation. We thus isolated 4-week old LN

cells and treated them in vitro with IL-7 or a combination of IL-1β+IL-23. We found that

ΔIAP1/2 γδT17 cells were slower in entering cell cycle with most cells stuck in G0 (Fig

4D-E). This defect in proliferation was also observed following stimulation with either IL-

2 or TCR cross-linking (Fig S5C). Therefore, cIAP1/2 are important for γδT17 cell cycle

progression.

We have shown before that ablation of cIAP1/2 in T cells downregulates cMAF, a

lineage determining transcription factor for γδT17 cells (Zuberbuehler *et al*, 2019), in a

NIK- and RelB-dependent mechanism (Rizk *et al*, 2019). We thus hypothesized that

lack of cIAP1/2 may influence expression of cMAF. In newborn thymus expression of

cMAF as well as RORγt was unchanged (Fig S6A). At week 1 after birth we observed a

slight reduction in the expression of RORγt and cMAF (Fig S6B). However, at week 3 of

age, expression of RORγt and cMAF was significantly reduced (Fig 5A). Furthermore,

we observed a modest but significant reduction in IL-17A production by 3-week old

ΔIAP1/2 γδT17 cells (Fig 5B). In the intestine, RORγt[+] γδ T cells express high levels of

CD127 (IL-7Rα) and intermediate levels of CD45 (Fig 5C). Due to lack of other reliable

surface markers to identify these cells in the gut, we gated TCRγδ$^+$CD45^{int}CD127$^+$ cells

and quantified numbers as well as expression of RORγt and cMAF. Similar to the LNs,

numbers in the siLP did not change in 4-week old ΔIAP1/2 mice (Fig 5D), however there

was a significant reduction in the levels of RORγt and cMAF (Fig 5E).

Furthermore, we investigated the expression of RelB in newborn thymic γδ T cells

and found a significant upregulation of *Relb* mRNA in ΔIAP1/2 γδT17 but not in CD27$^+$

γδ T cells (Fig S6C). We additionally examined the extent of RelB nuclear translocation

in γδT17 cells from 4-week old WT, ΔIAP2 and ΔIAP1/2 mice. We found that the levels

of nuclear RelB in ΔIAP1/2 γδT17 cells were significantly higher by comparison to WT

cells (Fig 5F), arguing for a role of the non-canonical NF-κB pathway in downregulation

of RORγt and cMAF. Therefore, cIAP1/2 are required during late neonatal life in order to

maintain expression of the transcription factors RORγt and cMAF and to sustain normal

γδT17 numbers.

Inflammation partially restores γδT17 responses in the absence of cIAP1 and

cIAP2

Next, we investigated whether cytokines that activate γδT17 cells could regulate

expression of RORγt and cMAF from 4-week old mice. Culture with IL-7 did not

influence expression of either transcription factors (Fig 6A-C), however, a combination

of IL-1β and IL-23 resulted in partial restoration of RORγt but not cMAF in ΔIAP1/2

γδT17 cells (Fig 6A-C). Interestingly, IL-1β+IL-23 resulted in downregulation of cMAF in

WT cells (Fig 6A-B). We additionally observed that the ΔIAP1/2 γδT17 cells that

acquired RORγt, were the cells that entered G1 in response to IL-1β+IL-23 and to a

lesser extent in response to IL-7 (Fig S7). In the imiquimod(IMQ)-driven psoriasiform

dermatitis model, IL-23, IL-1β and IL-7 drive γδT17 cell expansion as well as production

of IL-17 and IL-22 in the LN and skin (Michel *et al*, 2012; Cai *et al*, 2011, 2014). We thus

treated WT, ΔIAP1, ΔIAP2, and ΔIAP1/2 4-week old mice with IMQ for seven days and

assessed expression of RORγt and cMAF in γδT17 cells. We found that IMQ treatment

partially restored expression of both RORγt and cMAF in ΔIAP1/2 γδT17 cells in the

LNs (Fig 6D), suggesting that inflammation can rescue the ΔIAP1/2 phenotype.

We then investigated whether rescue of RORγt and cMAF was sufficient for

ΔIAP1/2 γδT17 cells to mount an immune response. We observed that despite an

increase in Ki67 expression (Fig S8A), ΔIAP1/2 γδT17 numbers did not increase in

either the LNs or skin (Fig 6E). Evaluation of cytokine production revealed substantial

regional differences between LN and skin in ΔIAP1/2 mice. Thus, whereas in the LN,

ΔIAP1/2 γδT17 cells increased (albeit significantly less than their WT, ΔIAP1 and ΔIAP2

counterparts) their production of IL-17A following IMQ treatment, this was not the case

in the skin (Fig 6F). In contrast, IL-22 production in LNs was significantly reduced while

it was relatively normal in the skin (Fig 6G). The CD4[+] T cell response to IMQ was not

defective and slightly stronger in ΔIAP1/2 mice (Fig S8B). The extent of skin

inflammation, as measured by epidermal thickening, was not different between ΔIAP1/2

and control mice, reflecting both the partial rescue of the γδT17 as well as the slightly

exaggerated CD4[+] T cell response (Fig S8C).

The data suggest that although at a young age cIAP1/2 regulate proliferation,

transcriptional stability and cytokine production, strong inflammatory stimuli can, to a

certain extent, overcome this regulatory checkpoint and revive γδT17 cell responses.

These results additionally indicate that the extrathymic expression and biological impact

thereafter of RORγt and cMAF can be dynamic and under the control of multiple

microenvironment cues.

cIAP1 and cIAP2 are required for intestinal ILC3 during aging and for sustaining

ILF integrity

ILC3 share many functional characteristics and transcription factor requirements with

γδT17 cells, including constitutive expression of RORγt and cMAF (Zuberbuehler *et al*,

2019; Parker *et al*, 2019). We therefore investigated the impact of cIAP1 and cIAP2

deficiency in intestinal ILC3 populations. Similar to γδT17 cells, LP Tbet$^+$ and Tbet$^-$ ILC3

numbers were reduced in ΔIAP1/2 mice (Fig 7A-C and Fig S9A). As expected ILC2

numbers were not affected (Fig S9B). Next, we tested whether ΔIAP1/2 ILC3 converted

to an RORγt$^-$ population and hence performed a lineage tracing experiment using the

RORγt-RFP reporter mice described above. We found that within the ILC population,

there was a 10-fold reduction in RFP$^+$ cell numbers, suggesting that in the absence of

cIAP1 and cIAP2, there is loss of ILC3 rather than conversion to a non-ILC3 population

(Fig S9C). Similar to γδT17 cells, we found that ILC3 numbers did not expand post

weaning (Fig 7D). Next, we investigated whether the ILC3 defect in ΔIAP1/2 mice was

cell-intrinsic. To this end, using mixed BM chimeras, we found that WT ILC3

outcompeted their ΔIAP1/2 counterparts (Fig 7E-F), indicating that the phenotype we

observed in intact mice was cell-intrinsic. There was equal reconstitution capacity of

GATA3-expressing ILC2 derived from WT or ΔIAP1/2 BM (Fig S9D), demonstrating the

specificity of the defect within RORγt-expressing populations. We obtained similar

results when we reconstituted ΔIAP1/2 hosts with a 1:1 mix of WT and ΔIAP1/2 BM (Fig

7G and Fig S9E). Further, cIAP1 and 2 deficient ILC3 cells were not rescued by

treatment with anti-TNF (Fig S9F).

ILC3 are necessary for the maturation of cryptopatches to ILFs during the first

weeks of life. ΔIAP1/2 mice had severely defective ILFs (Fig 7H). ILFs in these mice

were either absent or reduced in size (Fig 7H). Despite the lack of ILFs, production of

IgA was not defective (Fig S9G). Collectively, this data shows that cIAP1 and cIAP2 are

necessary for intestinal ILC3 to expand during the post-weaning period, and to induce

formation of ILFs.

cIAP1 and cIAP2 are necessary to protect against *Citrobacter rodentium* infection

It has been demonstrated that ILC3 are important to control infection by the attaching

and effacing bacterium *Citrobacter rodentium* (Bauché *et al*, 2020; Guo *et al*, 2015,

2014), a widely used model for human enteropathogenic *E. coli* infections (Silberger *et*

al, 2017) .We therefore reasoned that ΔIAP1/2 mice may be defective in mounting a

protective response to *C. rodentium*. We infected ΔIAP1/2 mice and their respective

controls with 2×10^9 CFU of *C. rodentium* through oral gavage and followed weight loss

as a surrogate marker for disease. We found that by 11 days after infection ΔIAP1/2

mice lost approximately 20% of their body weight, while all other strains did not (Fig 8A).

At this time point and due to ethical constraints, all animals were sacrificed and we

analyzed bacterial loads and the immune response in the colon. ΔIAP1/2 mice had

significantly higher colonic bacterial load than controls (Fig 8B). This was associated

with compromised IL-22 production from the ILC3 compartment (Fig 8C-D).

Although ILC3 are important to protect from *C. rodentium* infection, a T_H17 and

T_H22 response is also required, as evidenced by susceptibility of RAG1$^{-/-}$ mice to this

pathogen (Silberger *et al*, 2017). We therefore additionally analyzed the CD4[+] T cell

response in the colon. Numbers of total CD4[+] T cells were not changed in infected

ΔIAP1/2 mice (Fig 8E). However, there was a significant reduction in RORγt[+]Tbet[−] CD4[+]

T cells (Fig 8E), which was accompanied by reduced levels of IL-17A (Fig 8F). Despite

normal numbers of RORγt[+]Tbet[+] CD4[+] T cells (Fig 8E), IL-17A[+]IFN-γ[+] cells were

significantly reduced in infected ΔIAP1/2 mice (Fig 8F). However, production of IL-22

was not defective in the absence of cIAP1 and cIAP2 (Fig 8F). Collectively, our data

suggest that cIAP1 and cIAP2 are required within the ILC3 and T_H17 compartments to

control intestinal bacterial infection.

Discussion

In the present study we demonstrate that the E3 ubiquitin ligases cIAP1 and cIAP2 are

necessary for γδT17 cells to transition through to prepubescent life by regulating

cytokine-mediated proliferation and stable expression of the lineage defining

transcription factors cMAF and RORγt. Thus, during aging, cIAP1 and cIAP2 are

required in a cell-intrinsic manner to maintain cMAF and RORγt levels and to allow cells

to enter cell cycle in response to IL-7, IL-1β and IL-23. Consequently, γδT17-driven

inflammatory responses in the skin and draining LNs of prepubescent ΔIAP1/2 mice are

blunted despite normal cell numbers, while by the time animals reach adulthood, γδT17

populations are deficient in gut, skin and LNs. Mechanistically, our data suggest that

this is independent of TNF and TNFR1 and most likely through overt activation of the

non-canonical NF-κB pathway. Similar to γδT17, ILC3 required cIAP1 and cIAP2

expression during the post-weaning period in order to expand, be maintained until adult

life, and induce formation of intestinal ILFs. The ILC3 deficit in ΔIAP1/2 mice together

with a defective T$_H$17 response, correlated with a profound inability to control intestinal

bacterial infection.

The IL-17-producing γδ T cell subset is an innate-like unconventional lymphocyte

that is important in many immunological processes ranging from anti-microbial

protection to pathogenic inflammation and cancer (Patil *et al*, 2015). γδT17 cells are

pre-programmed and functionally mature in the embryonic thymus in mouse and human

(Ribot *et al*, 2009; Haas *et al*, 2012). They are exported into peripheral and secondary

lymphoid tissues after birth, and evidence suggests that they go through a second wave

of transcriptional and functional programming during neonatal life within the tissues they

occupy (Kadekar *et al*, 2020; Wiede *et al*, 2017). The molecular cues that γδT17 cells

receive within the tissues during that period are obscure. Our data show that the E3

ligases cIAP1 and cIAP2 are required during late neonatal and early prepubescent life

in a cell-intrinsic manner for cytokine-induced proliferation, to sustain transcriptional

stability and allow optimal inflammatory responses. This work establishes cIAP1/2 as

critical molecular regulators of committed tissue-resident γδT17 cells, and underpins the

existence and importance of post-thymic temporal events necessary for these cells to

be maintained during aging.

cIAP1/2 are central for TNFR1 induced canonical NF-κB activation and cell death

(Mahoney *et al*, 2008) , and necessary to suppress overt non-canonical NF-κB signaling

(Vallabhapurapu *et al*, 2008; Zarnegar *et al,* 2008) . TNFR1 induced apoptosis and

necroptosis are fundamental biological processes regulating cell growth during

development, homeostatic turnover and even inflammatory diseases (Kalliolias &

Ivashkiv, 2016). The two NF-κB pathways on the other hand are synonymous with cell

survival, proliferation and differentiation in ubiquitous cell populations (Hayden & Ghosh,

2011). In T cells they are mostly active downstream of TNF superfamily receptors and

the TCR (Oh & Ghosh, 2013). Although the role of several TNF superfamily receptors

and ligands have been studied in γδ T cells (Powolny-Budnicka *et al*, 2011; Shibata *et*

al, 2011; Silva-Santos *et al*, 2005), the importance of the signaling components of the

NF-κB pathway had not been thoroughly investigated. Genetic and pharmacological

perturbations of the TNFR1 signaling, combined with aberrant nuclear translocation of

RelB that it is the cIAP1/2-mediated control of non-canonical NF-κB that is required for

the maintenance of γδT17 cells. This agrees with CD27, a TNF superfamily receptor

and potent activator of non-canonical NF-κB (Ramakrishnan *et al*, 2004), suppressing

the γδT17 differentiation program (Ribot *et al*, 2009). Importantly, deletion of NIK, the

kinase targeted by cIAP1/2 and responsible for activating the non-canonical NF-κB

cascade, did not affect γδT17 cell development or homeostasis (Mair *et al*, 2015). This

strongly suggests that it is the "brake" imposed by cIAP1/2 in order to avoid over

activation of non-canonical NF-κB that is critical and not its baseline activity.

There are a number of transcription factors that are important for the development

of γδT17 cells (Parker & Ciofani, 2020) . Ciofani and co-workers showed that cMAF acts

early in embryogenesis to allow robust expression of RORγt and thus promote

specification and stability of the γδT17 lineage (Zuberbuehler *et al*, 2019). How cMAF

and RORγt expression is regulated, however, in γδT17 cells is not well-defined. Herein,

we demonstrate that loss of cIAP1/2 results in the progressive downmodulation of cMAF

and RORγt after birth, providing a molecular understanding of how lineage defining

transcription factors are regulated in these cells. Although loss of cMAF and RORγt

during embryonic development resulted in rapid loss of γδT17 cells or their progenitors

in the thymus (Zuberbuehler *et al*, 2019; Shibata *et al*, 2011) , we observed that in

ΔIAP1/2 mice, cells persist for at least 2 weeks without either transcription factor. Thus,

it appears that during neonatal life the impact of cMAF and RORγt in γδT17 cells is less

pronounced. The exact molecular steps leading to cIAP1/2-dependent regulation of

cMAF and RORγt are currently unclear. Our previous work showed that following

cIAP1/2 inhibition, NIK-mediated RelB nuclear translocation suppressed expression of

cMAF in T_H17 cells (Rizk *et al*, 2019). It is plausible therefore, that accumulation of non-

canonical NF-κB signaling directly suppresses cMAF, which subsequently suppresses

RORγt. Intriguingly, cytokine stimulation and inflammation could partially restore

expression of cMAF and RORγt, indicating a certain degree of transcriptional plasticity.

In addition to γδT17 cells, cIAP1/2 were necessary for the establishment of a

normal ILC3 population during the post-weaning period in the gut and the formation of

ILFs, as well as protection from intestinal extracellular bacterial infection. During

infection, we additionally found that cIAP1/2, were critical for the generation of IL-17$^+$

and IL-17$^+$IFN-γ$^+$ CD4+ T cells, which have been associated with protection against

pathogens, or tissue damage in the context of inflammation (Omenetti *et al*, 2019). We

and others have previously reported that the cIAP-non-canonical NF-κB axis is

necessary for T_H17 differentiation and successful IL-17-driven responses (Rizk *et al*,

2019; Kawalkowska *et al*, 2019), while NIK was shown to be important for the

generation of neuropathogenic T_H17 cells (Lacher *et al*, 2018). Moreover, NIK

expression and activation of the non-canonical NF-κB pathway in dendritic cells

indirectly regulates maintenance of both T_H17 cells and ILC3 (Jie *et al*, 2018). Our

current data, extend and broaden the immunological importance of this pathway. We

would like to propose that through regulation of non-canonical NF-κB, cIAP1/2 are

master regulators of innate and adaptive type-3 immunity. Their requirement is

necessary during neonatal life to establish functional innate and innate-like type-3

immune cell populations, whereas in the adult they support differentiation of antigen-

dependent adaptive type-3 cells.

References

Annibaldi A & Meier P (2018) Checkpoints in TNF-Induced Cell Death: Implications in

Inflammation and Cancer. *Trends Mol Med* 24: 49–65

Annibaldi A, Wicky John S, Vanden Berghe T, Swatek KN, Ruan J, Liccardi G, Bianchi

K, Elliott PR, Choi SM, Van Coillie S, *et al* (2018) Ubiquitin-Mediated Regulation of

RIPK1 Kinase Activity Independent of IKK and MK2. *Mol Cell* 69: 566-580.e5

Bauché D, Joyce-Shaikh B, Fong J, Villarino A V, Ku KS, Jain R, Lee Y-C, Annamalai L,

Yearley JH & Cua DJ (2020) IL-23 and IL-2 activation of STAT5 is required for

optimal IL-22 production in ILC3s during colitis

Bouladoux N, Harrison OJ & Belkaid Y (2017) The Mouse Model of Infection with

Citrobacter rodentium. *Curr Protoc Immunol* 119: 19.15.1-19.15.25

Cahenzli J, Köller Y, Wyss M, Geuking MB & McCoy KD (2013) Intestinal Microbial

Diversity during Early-Life Colonization Shapes Long-Term IgE Levels. *Cell Host*

Microbe 14: 559–570

Cai Y, Shen X, Ding C, Qi C, Li K, Li X, Jala VR, Zhang H ge, Wang T, Zheng J, *et al*

(2011) Pivotal Role of Dermal IL-17-Producing gd T Cells in Skin Inflammation.

Immunity 35: 596–610

Cai Y, Xue F, Fleming C, Yang J, Ding C, Ma Y, Liu M, Zhang HG, Zheng J, Xiong N, *et*

al (2014) Differential developmental requirement and peripheral regulation for

dermal Vγ34 and Vγ36T17 cells in health and inflammation. *Nat Commun* 5: 1–14

Chen YS, Chen IB, Pham G, Shao TY, Bangar H, Way SS & Haslam DB (2020) IL-17-

producing γδ T cells protect against Clostridium difficile infection. *J Clin Invest* 130:

2377–2390

Conti HR, Peterson AC, Brane L, Huppler AR, Herna´ndez-Santos N, Whibley N, Garg

A V., Simpson-Abelson MR, Gibson GA, Mamo AJ, *et al* (2014) Oral-resident

natural Th17 cells and γδ T cells control opportunistic Candida albicans infections.

J Exp Med 211: 2075–2084

Eberl G (2016) Immunity by equilibrium. *Nat Rev Immunol* 16: 524–532

Eberl G & Litman DR (2004) Thymic origin of intestinal αβ T cells revealed by fate

mapping of RORγt+ cells. *Science (80-)* 305: 248–251

Fleming C, Cai Y, Sun X, Jala VR, Xue F, Morrissey S, Wei YL, Chien YH, Zhang HG,

Haribabu B, *et al* (2017) Microbiota-activated CD103+ DCs stemming from

microbiota adaptation specifically drive γδT17 proliferation and activation.

Microbiome 5: 1–15

Gardam S, Turner VM, Anderton H, Limaye S, Basten A, Koentgen F, Vaux DL, Silke J

& Brink R (2011) Deletion of cIAP1 and cIAP2 in murine B lymphocytes

constitutively activates cell survival pathways and inactivates the germinal center

response. *Blood* 117: 4041–4051

Guo X, Liang Y, Zhang Y, Lasorella A, Kee BL & Fu YX (2015) Innate Lymphoid Cells

Control Early Colonization Resistance against Intestinal Pathogens through ID2-

Dependent Regulation of the Microbiota. *Immunity* 42: 731–743

Guo X, Qiu J, Tu T, Yang X, Deng L, Anders RA, Zhou L & Fu YX (2014) Induction of

innate lymphoid cell-derived interleukin-22 by the transcription factor STAT3

mediates protection against intestinal infection. *Immunity*

Gury-BenAri M, Thaiss CA, Serafini N, Winter DR, Giladi A, Lara-Astiaso D, Levy M,

Salame TM, Weiner A, David E, *et al* (2016) The Spectrum and Regulatory

Landscape of Intestinal Innate Lymphoid Cells Are Shaped by the Microbiome. *Cell*

166: 1231-1246.e13

Haas JD, Malinarich González FH, Schmitz S, Chennupati V, Föhse L, Kremmer E,

Förster R & Prinz I (2009) CCR6 and NK1.1 distinguish between IL-17A and IFN-γ-

producing γδ effector T cells. *Eur J Immunol* 39: 3488–3497

Haas JD, Ravens S, Düber S, Sandrock I, Oberdörfer L, Kashani E, Chennupati V,

Föhse L, Naumann R, Weiss S, *et al* (2012) Development of Interleukin-17-

Producing γδ T Cells Is Restricted to a Functional Embryonic Wave. *Immunity* 37:

48–59

Hayden MS & Ghosh S (2011) NF-κB in immunobiology. *Cell Res* 21: 223–244

Heilig JS & Tonegawa S (1986) Diversity of murine gamma genes and expression in

fetal and adult T lymphocytes. *Nature* 322: 836–840

Hoytema van Konijnenburg DP, Reis BS, Pedicord VA, Farache J, Victora GD & Mucida

D (2017) Intestinal Epithelial and Intraepithelial T Cell Crosstalk Mediates a

Dynamic Response to Infection. *Cell* 171: 783-794.e13

Jie Z, Yang JY, Gu M, Wang H, Xie X, Li Y, Liu T, Zhu L, Shi J, Zhang L, *et al* (2018)

NIK signaling axis regulates dendritic cell function in intestinal immunity and

homeostasis. *Nat Immunol* 19: 1224–1235

Kadekar D, Agerholm R, Rizk J, Neubauer HA, Suske T, Maurer B, Viñals MT, Comelli

EM, Taibi A, Moriggl R, *et al* (2020) The neonatal microenvironment programs

innate γδ T cells through the transcription factor STAT5. *J Clin Invest* 130: 2496–

2508

Kalliolias GD & Ivashkiv LB (2016) TNF biology, pathogenic mechanisms and emerging

therapeutic strategies. *Nat Rev Rheumatol* 12: 49–62

Kawalkowska JZ, Ogbechi J, Venables PJ & Williams RO (2019) CIAP1/2 inhibition

synergizes with TNF inhibition in autoimmunity by down-regulating IL-17A and

inducing Tregs. *Sci Adv* 5: eaaw5422

Kiss EA, Vonarbourg C, Kopfmann S, Hobeika E, Finke D, Esser C & Diefenbach A

(2011) Natural aryl hydrocarbon receptor ligands control organogenesis of intestinal

lymphoid follicles. *Science (80-)* 334: 1561–1565

Klose CSN, Kiss EA, Schwierzeck V, Ebert K, Hoyler T, D'Hargues Y, Göppert N,

Croxford AL, Waisman A, Tanriver Y, *et al* (2013) A T-bet gradient controls the fate

and function of CCR6-RORγt + innate lymphoid cells. *Nature* 494: 261–265

Kruglov AA, Grivennikov SI, Kuprash D V., Winsauer C, Prepens S, Seleznik GM, Eberl

G, Littman DR, Heikenwalder M, Tumanov A V., *et al* (2013) Nonredundant function

of soluble ltα3 produced by innate lymphoid cells in intestinal homeostasis. *Science*

(80-) 342: 1243–1246

Lacher SM, Thurm C, Distler U, Mohebiany AN, Israel N, Kitic M, Ebering A, Tang Y,

Klein M, Wabnitz GH, *et al* (2018) NF-κB inducing kinase (NIK) is an essential post-

transcriptional regulator of T-cell activation affecting F-actin dynamics and TCR

signaling. *J Autoimmun* 94: 110–121

St. Leger AJ, Desai J V., Drummond RA, Kugadas A, Almaghrabi F, Silver P,

Raychaudhuri K, Gadjeva M, Iwakura Y, Lionakis MS, *et al* (2017) An Ocular

Commensal Protects against Corneal Infection by Driving an Interleukin-17

Response from Mucosal γδ T Cells. *Immunity* 47: 148-158.e5

Mahoney DJ, Cheung HH, Mrad RL, Plenchette S, Simard C, Enwere E, Arora V, Mak

TW, Lacasse EC, Waring J, *et al* (2008) Both cIAP1 and cIAP2 regulate TNF -
mediated NF- B activation. *Proc Natl Acad Sci* 105: 11778–11783

Mair F, Joller S, Hoeppli R, Onder L, Hahn M, Ludewig B, Waisman A & Becher B
(2015) The NFκB-inducing kinase is essential for the developmental programming
of skin- resident and IL-17-producing Γδ Tδ cells. *Elife* 4

Di Marco Barros R, Roberts NA, Dart RJ, Vantourout P, Jandke A, Nussbaumer O,
Deban L, Cipolat S, Hart R, Iannitto ML, *et al* (2016) Epithelia Use Butyrophilin-like
Molecules to Shape Organ-Specific γδ T Cell Compartments. *Cell* 167: 203-
218.e17

Matsuzawa A, Tseng PH, Vallabhapurapu S, Luo JL, Zhang W, Wang H, Vignali DAA,
Gallagher E & Karin M (2008) Essential cytoplasmic translocation of a cytokine
receptor-assembled signaling complex. *Science (80-)* 321: 663–668

McGinley AM, Sutton CE, Edwards SC, Leane CM, DeCourcey J, Teijeiro A, Hamilton
JA, Boon L, Djouder N & Mills KHG (2020) Interleukin-17A Serves a Priming Role
in Autoimmunity by Recruiting IL-1β-Producing Myeloid Cells that Promote
Pathogenic T Cells. *Immunity* 52: 342-356.e6

Michel M-L, Pang DJ, Haque SFY, Potocnik AJ, Pennington DJ & Hayday AC (2012)
Interleukin 7 (IL-7) selectively promotes mouse and human IL-17-producing cells.
Proc Natl Acad Sci 109: 17549–17554

Al Nabhani Z, Dulauroy S, Marques R, Cousu C, Al Bounny S, Déjardin F, Sparwasser
T, Bérard M, Cerf-Bensussan N & Eberl G (2019) A Weaning Reaction to
Microbiota Is Required for Resistance to Immunopathologies in the Adult Cell Press

Oh H & Ghosh S (2013) NF-κB: Roles and regulation in different CD4+ T-cell subsets.
Immunol Rev 252: 41–51

Omenetti S, Bussi C, Metidji A, Iseppon A, Lee S, Tolaini M, Li Y, Kelly G, Chakravarty

P, Shoaie S, *et al* (2019) The Intestine Harbors Functionally Distinct Homeostatic

Tissue-Resident and Inflammatory Th17 Cells. *Immunity* 51: 77-89.e6

Parker ME, Barrera A, Wheaton JD, Zuberbuehler MK, Allan DSJ, Carlyle JR, Reddy

TE & Ciofani M (2019) c-Maf regulates the plasticity of group 3 innate lymphoid

cells by restraining the type 1 program. *J Exp Med*: jem.20191030

Parker ME & Ciofani M (2020) Regulation of γδ T Cell Effector Diversification in the

Thymus. *Front Immunol* 11: 42

Patil RS, Bhat SA, Dar AA & Chiplunkar S V (2015) The Jekyll and Hyde story of IL17-

Producing γδT Cells. *Front Immunol* 6: 37

Powolny-Budnicka I, Riemann M, Tänzer S, Schmid RM, Hehlgans T & Weih F (2011)

RelA and RelB transcription factors in distinct thymocyte populations control

lymphotoxin-dependent interleukin-17 production in γδ T cells. *Immunity* 34: 364–

374

Ramakrishnan P, Wang W & Wallach D (2004) Receptor-specific signaling for both the

alternative and the canonical NF-κB activation pathways by NF-κB-inducing kinase.

Immunity 21: 477–489

Ribot JC, DeBarros A, Pang DJ, Neves JF, Peperzak V, Roberts SJ, Girardi M, Borst J,

Hayday AC, Pennington DJ, *et al* (2009) CD27 is a thymic determinant of the

balance between interferon-γ- and interleukin 17-producing γδ T cell subsets. *Nat*

Immunol 10: 427–436

Rizk J, Kaplinsky J, Agerholm R, Kadekar D, Ivars F, Agace WW, Wei-Lynn Wong W,

Szucs MJ, Myers SA, Carr SA, *et al* (2019) SMAC mimetics promote NIK-

dependent inhibition of CD4+ TH17 cell differentiation. *Sci Signal* 12

Sandrock I, Reinhardt A, Ravens S, Binz C, Wilharm A, Martins J, Oberdörfer L, Tan L,

Lienenklaus S, Zhang B, *et al* (2018) Genetic models reveal origin, persistence and

non-redundant functions of IL-17–producing γδ T cells. *J Exp Med* 215: 3006–3018

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,

Rueden C, Saalfeld S, Schmid B, *et al* (2012) Fiji: an open-source platform for

biological-image analysis. *Nat Methods* 9: 676–682

Sheridan BS, Romagnoli PA, Pham QM, Fu HH, Alonzo F, Schubert WD, Freitag NE &

Lefrançois L (2013) γδ T Cells Exhibit Multifunctional and Protective Memory in

Intestinal Tissues. *Immunity* 39: 184–195

Shibata K, Yamada H, Sato T, Dejima T, Nakamura M, Ikawa T, Hara H, Yamasaki S,

Kageyama R, Iwakura Y, *et al* (2011) Notch-Hes1 pathway is required for the

development of IL-17-producing γδ T cells. *Blood* 118: 586–593

Silberger DJ, Zindl CL & Weaver CT (2017) Citrobacter rodentium: a model

enteropathogen for understanding the interplay of innate and adaptive components

of type 3 immunity. *Mucosal Immunol* 10: 1108–1117

Silke J & Meier P (2013) Inhibitor of apoptosis(IAP)proteins-modulators of cell death

and inflammation. *Cold Spring Harb Perspect Biol* 5: a008730

Silva-Santos B, Pennington DJ & Hayday AC (2005) Lymphotoxin-mediated regulation

of γδ cell differentiation by αβ T cell progenitors. *Science (80-)* 307: 925–928

Spits H, Artis D, Colonna M, Diefenbach A, Di Santo JP, Eberl G, Koyasu S, Locksley

RM, McKenzie ANJ, Mebius RE, *et al* (2013) Innate lymphoid cells-a proposal for

uniform nomenclature. *Nat Rev Immunol* 13: 145–149

Steer CA, Mathä L, Shim H & Takei F (2020) Lung group 2 innate lymphoid cells are

trained by endogenous IL-33 in the neonatal period. *JCI Insight* 5

Vallabhapurapu S, Matsuzawa A, Zhang W, Tseng P-H, Keats JJ, Wang H, Vignali

DAA, Bergsagel PL & Karin M (2008) Nonredundant and complementary functions

of TRAF2 and TRAF3 in a ubiquitination cascade that activates NIK-dependent

alternative NF-κB signaling. *Nat Immunol* 9: 1364–1370

Varfolomeev E, Blankenship JW, Wayson SM, Fedorova A V, Kayagaki N, Garg P,

Zobel K, Dynek JN, Elliott LO, Wallweber HJA, *et al* (2007) IAP Antagonists Induce

Autoubiquitination of c-IAPs, NF-κB Activation, and TNFα-Dependent Apoptosis.

Cell 131: 669–681

Vince JE, Wong WWL, Khan N, Feltham R, Chau D, Ahmed AU, Benetatos CA,

Chunduru SK, Condon SM, McKinlay M, *et al* (2007) IAP Antagonists Target cIAP1

to Induce TNFα-Dependent Apoptosis. *Cell* 131: 682–693

Wiede F, Dudakov JA, Lu KH, Dodd GT, Butt T, Godfrey DI, Strasser A, Boyd RL &

Tiganis T (2017) PTPN2 regulates T cell lineage commitment and αß versus γδ

specification. *J Exp Med* 214: 2733–2758

Zarnegar BJ, Wang Y, Mahoney DJ, Dempsey PW, Cheung HH, He J, Shiba T, Yang X,

Yeh WC, Mak TW, *et al* (2008) Noncanonical NF-κB activation requires coordinated

assembly of a regulatory complex of the adaptors cIAP1, cIAP2, TRAF2 and

TRAF3 and the kinase NIK. *Nat Immunol* 9: 1371–1378

Zuberbuehler MK, Parker ME, Wheaton JD, Espinosa JR, Salzler HR, Park E & Ciofani

M (2019) The transcription factor c-Maf is essential for the commitment of IL-17-

producing γδ T cells. *Nat Immunol* 20: 73–85

Materials and methods

Mice

All animals were bred and maintained in-house at DTU health tech with the approval of the Danish animal experiments inspectorate. cIAP1$^{f/f}$ and cIAP1$^{f/f}$ cIAP2$^{-/-}$ mice were provided by Prof. W. Wei-Lynn Wong at the University of Zurich, Switzerland with the permission of Prof. John Silke, VIC Australia. RORγt^{CRE} mice were provided by Prof. Gerard Eberl at Pasteur Institute, Paris, France. ROSA26-floxSTOPflox-RFP mice were from the Swiss Immunological Mouse Repository (SwImMR). Lymph nodes from TNFR1$^{-/-}$ mice were provided by Prof. William Agace at Lund University, Sweden, while Lymph nodes from cIAP1^{UBA} mutant mice were provided by Prof. Pascal Meier at The Institute of Cancer research, UK.

Cell culture media and buffers

For all preparations of single cell suspensions and cell cultures RPMI 1460 (Invitrogen) supplemented with 10% heat inactivated FBS (GIBCO), 20mM Hepes pH 7.4 (Gibco), 50 µM 2-mercaptoethanol, 2 mM L-glutamine (Gibco) and 10,000 U/ mL penicillin-streptomycin (Gibco), was used. Where indicated, IMDM (Invitrogen) was used instead of RPMI 1460 and supplemented as aforementioned. FACS buffer was prepared by supplementing PBS with 3% heat inactivated FBS.

Lymphocyte isolation from mouse organs

Lymphocytes were isolated from peripheral lymph nodes (axial, brachial and inguinal), thymus, ear skin, small intestinal and colonic lamina propria following the previously described protocols (Kadekar *et al*, 2020). Lymphocytes were isolated from cervical and auricular lymph nodes in case of IMQ-induced psoriasis.

Ex-vivo culturing of lymphocytes

646 For staining of cytokines from lymphocytes that were isolated from peripheral lymph

647 nodes of untreated mice, the cells were plated at a denisty of 10×10^6 cells /ml in 1ml of

648 supplemented RPMI in 12 well plates. The cells were restimulated with 50ng/ml PMA

649 (phorbol myristate acetate; Sigma Aldrich), 750 ng/ml Ionomycin (Sigma Aldrich) and

650 BD GolgiStop (containing monensin at 1:1000 dilution, BD) and cultured for 3.5 hours at

651 37°C. For estimation of IL-22 production by CD4$^+$ and γδ$^+$ T cells from homeostatic

652 mice, the lymphocytes were first cultured onvernight with 40ng/ml rmIL-23 (R&D) the

653 restimulated with PMA, Ionomycing and BD GolgiStop as aforementioned. The cells

654 were then harvested and used for flow cytometry staining.

655 In case of lymphocytes that were isolated from peripheral lymph nodes or skin in IMQ-

656 experiments, the cells were plated at a denisty of 5×10^6 cells /ml in 1ml of supplemented

657 IMDM in 24 well plates. The cells were restimulated with 50ng/ml PMA (phorbol

658 myristate acetate; Sigma Aldrich), 750 ng/ml Ionomycin (Sigma Aldrich) and BD

659 Golgiplug (containing Brefeldin A at 1:1000 dilution, BD) and cultured for 3.5 hours at

660 37°C. The cells were then harvested and used for flow cytometry staining.

661 Alternatively, lymphocytes that were isolated from mesenteric lymph nodes or colonic

662 lamina propria in *Citrobacter rodentium* infection experiments, the cells were plated at a

663 denisty of 5×10^6 cells /ml in 1ml of supplemented IMDM in 24 well plates. The cells were

664 subsequently treated with 40ng/ml rmIL-23 (R&D) for 3 hours, followed by 50ng/ml PMA

665 (phorbol myristate acetate; Sigma Aldrich), 750 ng/ml Ionomycin (Sigma Aldrich) and

666 BD Golgiplug (containing Brefeldin A at 1:1000 dilution, BD) and cultured for an

667 additional 3.5 hours at 37°C.

668 For cell cycle assay experiments, lymphocytes that were isolated from peripheral lymph

669 nodes of mice, were plated at a denisty of 5×10^6 cells /ml in 1ml of supplemented RMPI

in 24 well plates. The cells were treated with either 20ng/ml rmIL-7 (R&D), or with 10

ng/ml rmIL-1β (Biolegend) + 20 ng/ml rmIL-23 (R&D), or 20 ng/ml rhIL-2 (Biolegend), or

anti-CD3 (2 µg/ml; clone 145-2C11) for 48 hours. The cells were subsequently

harvested for flow cytometry staining.

IMQ-induced psoriasis

Psoriasis was induced in mice by applying 7 mg of Aldara cream (containing 5%

imiquimod) to the dorsal side of each ear for 7 days. Histological sections were

prepared by fixing ear tissue in 10% formalin overnight and then paraffin embedded.

The paraffin embedded sections were cut and stained by H&E.

Flow cytometry staining

Surface antigens, intracellular cytokines and cell cycle assay were stained for flow

cytometry as previously described (Rizk *et al*, 2019). For transcription factor staining,

the cells were first stained for live/dead discrimination followed by surface antigen

staining and subsequently fixed using Foxp3 fixation/permeabilization buffer (Thermo

Fisher) for 1 hour at 4°C. The cells were washed once with then stained with the desired

antibodies in Foxp3 perm/wash buffer for 1 hour at 4°C. The cells were washed once

again and resuspended in FACS buffer and analyzed using BD LSRFortessa.

The following antibodies were used herein at 1:200 dilution unless otherwise indicated:

Fixed viability stain-700 (FVS700, BD, 1:1000), anti-IL-17A (TC11-18H10; BV786 and

PE), anti-IFNγ (XMG1.2; PE-Cy7, APC, BV711 and Percp-cy5.5), anti-IL-22

(1H8PWSR; PE), anti-cMAF (symOF1; PE, eF660 or Percp-Cy5.5; 5 µL/test), anti-CD4

(GK1.5; BUV395 and FITC), anti-TCRγδ (GL3; BV421 and APC), anti-KLRG1 (2F1 ,

BV786), anti-CD27 (LG.3A10; PE-Cy7 and BV650), anti-CCR6 (140706; Alexa Fluor

647), anti-CD44 (1M7; V500), anti-CD19 (6D5; FITC), anti-TCRβ (H57-597; APC-

eflour780), anti-CD3e (145-2C11, PeCF594 and PE), anti-Tbet(4B10; PeCy7), anti-

CD8 (53-6.7;FITC), anti-Vγ5 (536; FITC), anti-Vγ4(UC3-10A6; Percp-eflour710), anti-

GATA3(TWAJ; Percp-eFlour710; 1:30), anti-CD45(30-F11;PE and V500), anti-CD127

(SB/199; BUV737) and anti-RORγt (B2D; APC and PE).

Administration of Anti-TNF

699 For neutralization of TNF, 1 week old pups were weighed and i.p. injected with the a-

700 TNF (Adalimumab, brand name HUMIRA) at 5 mg/kg body weight once a week until

701 weaning. After weaning the mice were i.p. injected with 10 mg/kg body weight twice a

702 week until euthanasia at approximately 12 weeks of age.

Transfer of neonatal γδ T cells to RAG1$^{-/-}$ hosts.

First, thymi from 1-2 days old mice were isolated and crushed individually against 70 µm

filter to prepare single cell solutions. Subsequently, total γδ T cells were enriched by

magnetic depletion of CD4$^+$, CD8$^+$, TCRβ$^+$ cells as follows: total thymocytes were re-

suspended in MACS buffer at 1e8 cells/ml containing 50 µL/ml normal rat serum and

1:200 biotin labelled anti-CD4$^+$ (GK1.5), CD8$^+$ (53-6.7) and TCRβ$^+$ (H57-597)

antibodies; the cells were incubated for 10 minutes at room temperature and then

incubated with 75 µL/ml EasySep RaphidSphere streptavidin beads (#50001) for 2.5

minutes then transferred to EasySep magnet for 2.5 minutes. The non-bound fraction

was collected by decantation and centrifugated for 5 minutes at 400g at 4^{o}C.. The

enriched γδ T cells from each donor mouse were re-suspended in PBS and then i.v.

injected into the tail vein of a RAG1$^{-/-}$ host. The RAG1$^{-/-}$ hosts were euthanized for

collection of organs after 12 weeks.

Bone marrow chimeras

The bone marrow cells for reconstitution were isolated by flushing the tibia and femur,

which were dissected from donor mice, with culture media. Total bone marrow cells

were then centrifuged at 400g for 5 minutes at 4°C. The cells were then re-suspended

and passed through 70 µm filter. Subsequently, red blood cells were then lysed using

RBC lysis buffer (Biolegend) and a single cell suspension of bone marrow cells was the

prepared by passing the cells through 40 µm filter. The prepared cells were then

counted and mixed as appropriate.

Conversely, host mice were sub-lethally irradiated by 2 doses of 4.5 Gy that were at

least 4 hours apart. After 24 hours, the hosts were reconstituted with 10e6 bone marrow

cells that were i.v. injected into the tail vein of the host mice. The hosts were euthanized

for organs after at least 12 weeks.

Immunofluorescent imaging of intestinal tissue

To assess the presence of SILT in the intestines of WT or ΔIAP1/2 by confocal laser

microscopy, the distal ileum was taken and flushed once with HBSS (Thermo Fisher) to

remove intestinal contents. Cleaned intestines were fixed for 8h in 4% PFA (Sigma-

Aldrich) in PBS and stored in washing buffer (PBS+5%FCS+0.2% Triton X-100 (Sigma-

Aldrich)+0.01% Thimerosal (Sigma-Aldrich)) until further use. To prepare the collected

intestines for staining, tissues were embedded in 4% UltraPure™ Low Melting Point

Agarose (Thermo Fisher) in PBS, sectioned with a swinging blade microtome (Leica

VT1200S) into 50 micron sections and permeabilized overnight using the Foxp3

Transcription Factor Staining Buffer Set (Thermo Fisher). Permeabilized sections were

stained in the supplied permbuffer with an antibody against RORγt (AFKJS-9;

unconjugated), followed by a washing step in permbuffer and incubation with a

biotinylated secondary antibody against the primary anti- RORγt antibody (Biotinylated

anti-rat; Jackson ImmunoResearch). To detect RORγt[+] ILC and B cells, sections were

washed again in permbuffer and incubated in permbuffer with antibody against B220

(RA3-6B2; AF647) and streptavidin-conjugated AF555 (Thermo Fisher), as well as

DAPI (Thermo Fisher) to stain all nucleated cells. Sections were washed one more time,

mounted on glass slides with ProLong Gold (Thermo Fisher) and analyzed using an

LSM710 confocal laser microscope (Carl Zeiss). Images of ≥5 different sections per

mouse were acquired with the Zeiss Zen v2.3 software (Carl Zeiss) and analyzed using

Imaris v8 (Bitplane/Oxford Instruments) and Fiji v2.1.0/1.53c (Schindelin *et al*, 2012).

Murine *Citrobacter rodentium* infection

Starter cultures of *Citrobacter rodentium* strain DBS100 (ATCC 51459; American Type

Culture Collection) were grown overnight at 37^{o}C in Luria-Bertani (LB) medium. The

cultures were then used at 5% v/v to inoculate sterile LB medium. The cultures were

grown at at 37^{o}C to an OD600 of 0.8- 1 and the CFU count was determined from the

OD600 measurement using the following formula: CFU/ml =$(5 \times 10^8)(OD) - 3 \times 10^7$.

Subsequently, the bacteria was collected by centrifugation at 4000g for 10 minutes. The

bacterial pellet was then resuspended in LB medium to give at 2×10^9 CFU/100 µL. To

infect adult mice, the mice were orally gavaged with either 100 µL of *Citrobacter*

rodentium or LB control. The mice were weighed before oral gavage and once daily until

termination of the experiment. At day 12 post infection all mice were euthanized,

dissected to collect organs and fecal samples.

The collected fecal samples were weighed and dissolved in PBS and then serially

diluted. The serial dilutions were plated on Brilliance™ E. coli/coliform Agar (CM0956,

Thermo Fisher) and incubated overnight at 37^{o}C. *Citrobacter rodentium* colonies were

identified as being pink colonies and enumerated, while E. coli colonies were identified

as purple colonies. CFU/g stool was then calculated as previously described

(Bouladoux *et al*, 2017).

Immunofluorescent imaging of nuclear RelB in γδ T cells

Total lymphocytes were isolated from peripheral (axial, brachial and inguinal), cervical

and auricular lymph nodes of 4 weeks old mice as described above. Subsequently, total

γδ T cells were enriched by magnetic depletion of $CD4^+$, $CD8^+$, $CD19^+$ and $TCRβ^+$ cells

as aforementioned. The cells were then stained with FVS700 for discrimination of live

and dead cells and then stained for surface antigens with the following antibodies: anti-

TCRγδ (GL3; APC), anti-CD27 (LG.3A10; PE-Cy7) and anti-TCRβ (H57-597; APC-

eflour780) all at 1:200 dilution. The cells were subsequently sorted into $TCRγδ^+$ $CD27^+$

or $TCRγδ^+$ $CD27^-$ cells using BD ARIA-FUSION cell sorter. The sorted cells were

collected into cell culture medium and centrifuged at 400g for 5 minutes at 4°C. The

cells were then fixed using Foxp3 fixation/permeabilization buffer (Thermo Fisher) for 1

hour at 4°C. The cells were washed once with then stained with Foxp3 perm/wash

buffer containing anti-TCRγδ (GL3; APC, 1:50), anti-CD3e (145-2C11 or 17A2; biotin,

1:100) and anti-RelB (D-4, Santa-cruz, 1:40) for 1-hour 4°C in Foxp3 perm/wash buffer.

Again, the cells were washed once and stained with streptavidin-conjugated AF488

(Biolegend, 1:100) and anti-mouse AF555 (1:100) for 1 hour 4°C in Foxp3 perm/wash

buffer. The cells were then washed once more as previous and stained with DAPI to

highlight cell nuclei and washed once more with PBS. Washed cells were mounted on a

glass slide using ProLong Gold and imaged with an LSM710 confocal laser microscope

and were acquired and analyzed with the Zen v2.3 software and and Fiji v2.1.0/1.53c

(Schindelin *et al*, 2012).

Real-Time quantitative PCR

At the indicated timepoints, γδ T cells were sorted from the thymus or lymph nodes of

mice directly into RLT buffer mixed with β-mercaptoethanol. RNA was then extracted

from using Qiagen RNAeasy microkit following the manufacturer's instructions.

Subsequently, cDNA was prepared using BioRad Iscript cDNA synthesis kit using the

manufacturer's protocol. Gene expression was then measured by RT-qPCR reactions

using BioRad SSOFast EvaGreen supermix, which were run on CFX96 (Biorad) and

analyzed using Bio-rad CFX manager software. The following primers were used for

RT-qPCR:

Actb, Fwd-GGCTGTATTCCCCTCCATCG, Rev- CCAGTTGGTAACAATGCCATGT;

RelB, Fwd- GCTGGGAATTGACCCCTACA, Rev- CATGTCGACCTCCTGATGGTT;

Birc2, Fwd- TGCCTGTGGTGGGAAACTGA, Rev- GCTCGGGTGAACAGGAACA.

Figures

Figure 1

Figure 1. cIAP1 and cIAP2 are required for the homeostasis of γδT17 cells in the

LNs and intestinal lamina propria.

(A) Graphical representation of the different mouse strains generated by crossing

RORγt[CRE] mice to cIAP1[F/F] or cIAP1[F/F] cIAP2[-/-] mice. Representative flow cytometric

analysis (B) and frequency (B-C) of IL-17$^+$ IL-22$^+$ cells within γδ T cells in the LNs (B) or

(C) IL-17$^+$ cells within γδ T in the siLP. (D) Representative flow cytometric analysis (dot

plots) and numbers (graph) of γδT17 cells in the LNs of adult WT, ΔIAP1, ΔIAP2 and

ΔIAP1/2 mice. (E) Representative flow cytometric analysis (dot plots) and numbers

(graph) of RORγt$^+$ Tbet$^+$ γδ T cells in the siLP of WT, ΔIAP1, ΔIAP2 and ΔIAP1/2 mice.

In graphs, each symbol represents a mouse, and lines represent the mean, data is pool

of 4 experiments in (B) or 5 experiments in (C-E). *P < 0.05, **P < 0.01, ***P < 0.001,

****P < 0.0001 using Kruskal-Wallis test with Dunn's correction.

Figure 2

Figure 2. cIAP1 and cIAP2 are non-redundantly required for the homeostatic

maintenance of γδT17 cell subsets in the skin.

Representative flow cytometric analysis (dot plots) and numbers (graphs) of (A) total γδ

T cells or (B) Vγ4$^+$ and Vγ4$^-$ CCR6$^+$ γδ T cells in the skin of WT, ΔIAP1, ΔIAP2 and

ΔIAP1/2 mice. In graphs, each symbol represents a mouse, and lines represent the

mean, data is pool of 5 experiments in (A-B). *P < 0.05, **P < 0.01 using Kruskal-Wallis

test with Dunn's correction.

Figure 3

Figure 3. cIAP1 and cIAP2 are intrinsically required for the homeostasis of γδT17

cells.

(A) Graphical representation of the experimental setup for competitive bone marrow

experiments. (B) Representative flow cytometric analysis (dot plots) and frequency

(graphs) of WT (CD45.1$^+$), host (CD45.1$^+$ CD45.2$^+$) or ΔIAP1/2 (CD45.2$^+$) -derived

γδT17 and CD27$^+$ γδ T cells within γδ T cells population in the LNs of reconstituted

hosts. (C) Frequency of WT (CD45.1$^+$), host (CD45.1$^+$ CD45.2$^+$) or ΔIAP1/2 (CD45.2$^+$) -

derived CD3$^-$ and CD19$^+$ cells in the LNs of reconstituted hosts. (D) Flow cytometric

analysis and (E) frequency of RORγt$^+$ and Tbet$^+$ γδ T cells in the cLP of host mice

follwing bone marrow reconstitution. (B-E) In graphs, each symbol represents a mouse,

and lines represent the mean, data is pool of 3 experiments. *$P < 0.05$, **$P < 0.01$, ***P

< 0.01 using one-way ANOVA with Tukey's correction. (F) Graphical representation of

the experimental setup for transfer of neonatal γδ T cells to RAG1$^{-/-}$ recipients. (G) Flow

cytometric analysis (contour plots) and numbers (graphs) of CCR6$^+$ CD27$^-$ or CD27$^+$ γδ

T cells in the LNs of RAG1$^{-/-}$ hosts after transfer of neonatal γδ T cells from WT or

ΔIAP1/2 pups. In graphs, each symbol represents a mouse, and lines represent the

mean, data is pool of 3 experiments (G). *$P < 0.05$ using Mann-Whitney test.

Figure 4

Figure 4. cIAP1 and 2 double deficient γδT17 cells are lost progressively during

aging and fail to progress through cell cycle.

(A) Numbers of γδT17 cells in the thymi of 1-day old WT, ΔIAP1, ΔIAP2 and ΔIAP1/2

pups. In graph, each symbol represents a mouse, and lines represent the mean, data is

a pool of 3 experiments. (B) Numbers of γδT17 cells in the LN of WT and ΔIAP1/2 at the

indicated timepoints. Each symbol represents the mean amalgamated data from each

timepoint and the error bars represents the SEM. ****P < 0.01 using Two-way ANOVA

with Holm-Sidak correction. (C) Graphical representation of the genetic makeup of the

ΔIAP1/2 mice when crossed to the ROSA26-LSL-RFP strain, and representative flow

cytometry analysis of LN γδT17 cells from ΔIAP1/2 x ROSA26-LSL-RFP mice. (D)

Representative flow cytometric analysis and (E) frequency of cells in G0, G1, S or G2/M

cell cycle stages within γδT17 cells that were ex-vivo cultured with the indicated

cytokines for 48 hours. In graphs, each symbol represents a mouse, and bars represent

the mean, data is pool of 3 experiments. ***$P < 0.001$, ****$P < 0.0001$ using two-way

ANOVA with Tukey's correction.

Figure 5

Figure 5. cIAP1 and 2 maintain the transcription factor landscape of γδT17 cells in the lymph nodes and intestinal lamina propria.

(A) Flow cytometric analysis (dot plots) and frequency (graphs) of RORγt$^+$ cMAF$^+$ cells within γδT17 cells from the LNs of 3-week-old WT and ΔIAP1/2 mice. (B) frequency of IL-17$^+$ cells within γδT17 cells (top) or within all γδ T cells (bottom) from the LNs of 3-week-old WT and ΔIAP1/2 mice. In graphs, each symbol represents a mouse, and the line represent the mean, data is pool of 2 experiments. **P < 0.01 using Mann-Whitney test. (C) Flow cytometric analysis showing the gating strategy and expression of RORγt by CD45^{int} CD127$^+$ γδ T cells in siLP of adult WT mice. (D) Numbers of CD45^{int} CD127$^+$ γδ T cells in the siLP of 4-week-old WT, ΔIAP1, ΔIAP2 and ΔIAP1/2 mice. (E) Flow cytometric analysis (dot plots) and quantification (graphs) of RORγt or cMAF expression by CD45^{int} CD127$^+$ γδ T cells in the siLP of 4-week-old WT, ΔIAP1, ΔIAP2 and ΔIAP1/2 mice. In graphs, each symbol represents a mouse, and the line represent the mean, data is pool of 4 experiments. (F) Representative immunofluorescent microscopy analysis of γδT17 cells (CD27$^-$ TCRγδ$^+$ cells) and normalized nuclear RelB fluorescence intensity in sorted from the LNs of 4-week-old WT or ΔIAP1/2. Images are representative of four independent experiments. *P < 0.05, **P < 0.01, ****P < 0.001 using using Kruskal-Wallis test with Dunn's correction.

Figure 6

Figure 6. Inflammation partially overcomes cIAP1 and cIAP2 deficiency in γδT17

cells.

(A) Representative flow cytometric analysis and (B-C) quantification of RORγt and

cMAF expression by γδT17 cells from the LNs of 4-week-old WT or ΔIAP1/2 mice

following ex-vivo culture with the indicated cytokines for 48 hours. In graphs, each

symbol represents a mouse, and the line represent the mean, data is pool of 3

experiments. (D) Representative flow cytometric analysis (dot plot) and quantification

(graph) of RORγt and cMAF expression by γδT17 in LNs of 4-week-old control or IMQ-

treated ΔIAP1/2 mice. (E) Numbers of γδT17 cells in the LNs (right) or skin (left) of 4-

week-old control or IMQ-treated WT, ΔIAP1, ΔIAP2 or ΔIAP1/2 mice. (F-G) Frequency

of IL-17A$^+$ (F) or IL-22$^+$ (G) cells within γδT17 cells in the LNs or skin of 4-week-old

control or IMQ-treated WT, ΔIAP1, ΔIAP2 or ΔIAP1/2 mice. In graphs, each symbol

represents a mouse, and the line represent the median, data is pool of 3 experiments.

*P< 0.05, **P < 0.01, ****P < 0.0001 using Mann-Whitney test.

Figure 7

Figure 7. cIAP1 and cIAP2 are intrinsically required for the maintanance of intestinal ILC3 and for ILF integrity

Representative flow cytometric analysis of (A) total CD45$^+$ CD3$^-$ CD19$^-$ CD127$^+$ ILCs or (B) CD45$^+$ CD3$^-$ CD19$^-$ CD127$^+$ GATA3$^-$ cells in the siLP of adult WT, ΔIAP1, ΔIAP2 or ΔIAP1/2 mice. (C) Numbers of total ILC3s, CD4$^+$, Tbet$^+$ or Tbet$^-$ ILC3s in the siLP of adult WT, ΔIAP1, ΔIAP2 or ΔIAP1/2 mice. In graphs, each symbol represents a mouse, and the line represent the mean, data is pool of 5 experiments. *P< 0.05, **P < 0.01 using Kruskal-Wallis test with Dunn's correction. (D) Numbers of ILC3s in the siLP of WT and ΔIAP1/2 at the indicated timepoints. Each symbol represents the mean amalgamated data from each timepoint and the error bars represents the SEM. **P < 0.01, ****P < 0.0001 using Two-way ANOVA with Holm-Sidak correction. (E) representative flow cytometric analysis (dot plots) and (F) frequency (graph) of WT (CD45.1$^+$), host (CD45.1$^+$ CD45.2$^+$) or ΔIAP1/2 (CD45.2$^+$)- derived ILC3 in the cLP of reconstituted hosts. In the graph, each symbol represents a mouse, and lines represent the mean, data is pool of 3 experiments. ***P < 0.01 using one-way ANOVA with Tukey's correction. (G) Frequency of WT (CD45.1$^+$) or ΔIAP1/2(CD45.2$^+$)- derived ILC3 in the siLP of bone marrow reconstituted ΔIAP1/2 hosts. In graph, each symbol represents a mouse, and lines represent the mean, data is pool of 2 experiments. *P < 0.05 using Wilcoxon-rank t-test. (H) representative immunofluorescent microscopy images showing ILF structures in distal ileum sections from adult WT or ΔIAP1/2 mice. Images are representative of two independent experiments.

Figure 8

Figure 8. RORγt^Cre+ cIAP1^F/F cIAP2^-/- succumb to *Citrobacter rodentium* infections.

(A) Percentage body weight change of *C. rodentium* infected WT, ΔIAP1, ΔIAP2 and

ΔIAP1/2 mice. Each symbol represents the mean amalgamated data from each

timepoint and the error bars represents the SD. ***P < 0.001 using Two-way ANOVA

with Holm-Sidak correction. (B) Colony forming units (CFU) of *C. rodentium* in fecal

matter of infected and uninfected WT, ΔIAP1, ΔIAP2 and ΔIAP1/2 mice. (C) Numbers of

ILC3s in the cLP of infected and uninfected mice. (D) representative flow cytometric

analysis (dot blots) and frequency of IL-22$^+$ cells within ILCs in the cLP of *C. rodentium*

infected WT and ΔIAP1/2 mice. (E) Numbers of total CD4$^+$ T cells, Tbet$^+$ RORγt$^+$ CD4$^+$

T cells, and RORγt$^+$ Tbet- CD4$^+$ T cells in the cLP of infected and uninfected mice. (F)

Frequency of IL-17A$^+$, IL-17A$^+$ IFNγ$^+$ or IL-22$^+$ cells within CD4$^+$ T cells in the cLP of

infected and uninfected mice. In graphs, each symbol represents a mouse, and lines

represent the median, data is pool of 3 experiments. *P < 0.05, **P < 0.01, ***P < 0.001

using Mann-Whittney test.