Multiple domains in ARHGAP36 regulate
PKA degradation and Gli activation

Patricia R. Nano[1], Takamasa Kudo[1], Nancie A. Mooney[2],
Jun Ni[1,5], Janos Demeter[2], Peter K. Jackson[2], and James K. Chen[1,3,4,*]

[1]Department of Chemical and Systems Biology, Stanford University School of Medicine, Stanford, CA
[2]Department of Microbiology and Immunology, Stanford University School of Medicine, Stanford, CA
[3]Department of Developmental Biology, Stanford University School of Medicine, Stanford, CA
[4]Department of Chemistry, Stanford University, Stanford, CA
[5]Present address: Brain Science and Advanced Technology Institute, Wuhan University of Science and Technology, Wuhan, China

* Correspondence should be addressed to J.K.C. (jameschen@stanford.edu)

ABSTRACT

ARHGAP36 is a Rho GTPase-activating protein (GAP) family member that contributes to spinal cord development and tumorigenesis. This multidomain protein is composed of N-terminal sequences that vary between splice isoforms, the GAP-like region, and a unique C-terminal domain, and previous studies have revealed an N-terminal arginine-rich region that promotes protein kinase A catalytic subunit (PKA_{cat}) degradation and Gli transcription factor activation. Although these signaling mechanisms can account for at least some aspects of ARHGAP36-dependent physiology, the functional roles of other structural elements in ARHGAP36 are not well understood. To address this question, we have mapped the ARHGAP36 structure-activity landscape with both domain- and amino-acid-level resolution. We observe that ARHGAP36-mediated Gli activation in cultured mammalian cells can be autoinhibited by isoform-specific N-terminal sequences, which also regulate ARHGAP36 trafficking and the subcellular targeting of PKA_{cat} pools. ARHGAP36 autoinhibition is counteracted by the GAP-like and C-terminal domains, which promote protein trafficking to the plasma membrane and primary cilium, respectively. In addition, our studies provide evidence that the GAP-like domain conditionally suppresses the N-terminal arginine-rich region and regulates ARHGAP36 binding to several interactors, including the prolyl oligopeptidase-like protein PREPL and the E3 ubiquitin ligase PRAJA2. Thus, multiple domains within ARHGAP36 can modulate its activity state, subcellular localization, and protein binding, potentially providing a means for achieving tissue-specific ARHGAP36 functions.

INTRODUCTION

Gli transcription factors (Gli1-3) are essential regulators of cell proliferation and differentiation, controlling fate specification in the neural tube (*Briscoe, et al., 2000; Stamataki, et al., 2005*) and limb bud (*Hill, et al., 2009; te Welscher, et al., 2002*) and the maintenance of granule neuron precursors in the developing cerebellum (*Lewis, et al., 2004; Wallace, 1999; Wechsler-Reya, et al., 2001*). Accordingly, misregulation of Gli activity can lead to uncontrolled cell growth, resulting in basal cell carcinoma, medulloblastoma, and other human cancers (*Hui, et al., 2011*). Gli functions are primarily regulated by the Hedgehog (Hh) pathway, and in the absence of Hh ligands, GLI2 and GLI3 are bound to the scaffolding protein Suppressor of Fused (SUFU) (*Stone, et al., 1999; Wang, Chengbing, et al., 2010*), which promotes their sequential phosphorylation by protein kinase A (PKA), glycogen synthase kinase β (GSK3β), and casein kinase 1 (CK1) (*Pan, et al., 2006; Pan, et al., 2007; Tempe, et al., 2006; Wang, Baolin, et al., 2006*). Proteasomal machinery is then recruited to the phosphorylated Gli proteins, resulting in GLI2 degradation and proteolytic conversion of GLI3 into a transcriptional repressor (*Pan, et al., 2007*).

Hh ligands suppress these intracellular processes, acting via the transmembrane receptors Patched1 (PTCH1) and Smoothened (SMO). Through mechanisms that remain unclear, SMO promotes the dissociation of GLI2 and GLI3 from SUFU, uncoupling the transcription factors from proteasomal regulation and allowing the full-length proteins to become transcriptional activators (*Humke, et al., 2010; Tukachinsky, et al., 2010*). SMO activity is suppressed by PTCH1 (*Murone, et al., 1999; Rohatgi, et al., 2007; Taipale, J., et al., 2002*), which is in turn directly inhibited by Hh ligands (*Incardona, et al., 2000; Stone, et al., 1996*). Hh signaling therefore induces SMO activation and the expression of Hh target genes, including those that encode PTCH1 (*Ågren, et al., 2004*) and the constitutively active transcription factor GLI1 (*Bai, et al., 2004; Dai, et al., 1999*). The primary cilium serves a key center for these signaling events (*Dorn, et al., 2012; Haycraft, et al., 2005; Kim, et al., 2009; May, et al., 2005; Rohatgi, et al., 2007;*

Wang, Yu, et al., 2009; Wen, et al., 2010), and this cell-surface protrusion is required for both Gli

activator and repressor formation (*Huangfu, et al., 2005; Liu, Aimin, et al., 2005*).

In addition to these signal transduction mechanisms, there is growing evidence for non-

canonical Gli regulation in both normal physiology (*Dennler, et al., 2007; Flora, et al., 2009; Riobó,*

et al., 2006) and in cancer (*Beauchamp, et al., 2009; Dennler, et al., 2007; Elsawa, et al., 2011;*

Han, et al., 2015; Kasper, et al., 2006; Liu, Z., et al., 2014; Long, et al., 2014). Our laboratory

previously established ARHGAP36 as a non-canonical Gli activator that acts in a SMO-

independent manner (*Rack, et al., 2014*). We identified this Rho GTPase-activating protein (GAP)

family member in a genome-scale screen for Hh pathway agonists (*Rack, et al., 2014*), and

subsequent studies have uncovered an essential role for ARHGAP36 in the specification of lateral

motor column neurons (*Nam, et al., 2019; Rack, et al., 2014*). Endogenous *Arhgap36* transcription

in the developing mouse spinal cord coincides with Hh pathway activation, and its overexpression

leads to ectopic induction of the Hh target genes *Ptch1* and *Gli1* (*Nam, et al., 2019*). In addition,

Arhgap36 expression has been found to correlate with SMO inhibitor resistance in Hh pathway-

driven murine medulloblastomas (*Buonamici, et al., 2010; Rack, et al., 2014*), and upregulating

this Rho GAP family member in neural progenitor cells is sufficient to induce medulloblastomas

in mice (*Beckmann, et al., 2019*). ARHGAP36 may promote tumor growth through multiple

mechanisms, as elevated *ARHGAP36* expression also has been associated with Hh pathway-

independent subtypes of medulloblastoma and neuroblastoma (*Beckmann, et al., 2019; Lee, et*

al., 2019).

Despite the emerging importance of ARHGAP36 in neuronal development and cancer, the

biochemical and cellular mechanisms that regulate and transduce its activity are not well

understood. The ARHGAP36 protein consists of unique N- and C-terminal domains and a central

region that is homologous to Rho GAPs. In addition, the five annotated isoforms of human

ARHGAP36 have varying N-terminal structures due to alternative splicing (Figure 1A). Four of the

ARHGAP36 isoforms can activate Gli proteins, with the gene product harboring the longest N-

terminal domain (isoform 1) being the sole exception (*Rack, et al., 2014*). Isoform 1 is also the

only ARHGAP36 protein that does not localize to the plasma membrane, and it instead adopts a

perinuclear distribution (*Müller, et al., 2020; Rack, et al., 2014*). In addition, the shortest

ARHGAP36 protein (isoform 3) accumulates in the primary cilium, whereas other ARHGAP36

isoforms cannot be detected in this signaling center under steady-state conditions (*Rack, et al.,*

2014). More recently, it has been shown that an N-terminal arginine-rich motif conserved in all

human ARHGAP36 isoforms can bind directly to catalytic subunits of PKA (PRKACA and

PRKACB; henceforth referred to as PKA_{cat}) (*Eccles, et al., 2016*). In the context of isoform 2, this

motif mediates the degradation of PKA_{cat}, and a 77-amino-acid N-terminal fragment that includes

this arginine-rich region has been shown to be necessary and sufficient for cellular PKA_{cat}

depletion (*Eccles, et al., 2016*). These findings suggest a role for N-terminal sequences in

targeting ARHGAP36 to specific subcellular compartments and establish PKA_{cat} inhibition as a

potential basis for ARHGAP36-mediated Gli activation.

In comparison, the functions of the GAP-like and C-terminal domains in ARHGAP36 have

not yet been elucidated. Rho GAP family members typically attenuate the function of Rho

GTPases by stimulating GTP hydrolysis (*Moon, 2003*). However, the GAP-like region in

ARHGAP36 lacks the "arginine finger" motif conserved in catalytically active homologs (*Rack, et*

al., 2014; Scheffzek et al., 1998), and ARHGAP36 has no effect on the activities of Rac1, Cdc42,

and Rho A (*Müller, et al., 2020*). In addition, ARHGAP36 residues that are structurally equivalent

to those previously associated with Rho GAP-catalyzed GTP hydrolysis are not required for

ARHGAP36-mediated Gli activation (*Rack, et al., 2014*). Non-catalytic mechanisms have been

reported for several Rho GAP family members (*Amin, et al., 2016; Faucherre, et al., 2003;*

Marchesi, et al., 2014), and it is possible that the GAP-like domain in ARHGAP36 similarly

interacts with Rho GTPases or other signaling proteins in a stoichiometric manner. How the C-

terminal domain might contribute to ARHGAP36 function is even more enigmatic since it lacks

sequence homology with other proteins.

Deciphering the molecular and cellular mechanisms that regulate ARHGAP36 activity

requires a deeper understanding of the relationship between ARHGAP36 structure and function.

Here we describe our systematic mapping of the ARHGAP36 structure-activity landscape using

individual ARHGAP36 isoforms, truncated variants, and a high-throughput mutagenesis screen.

Our findings demonstrate that ARHGAP36-dependent Gli activation and cellular PKA_{cat} depletion

are separable activities and reveal isoform-specific differences in subcellular PKA_{cat} targeting.

While the ARHGAP36 N-terminal domain is necessary and sufficient for Gli activation, an N-

terminal region in isoform 2 (residues 1-105; $N2_{1-105}$) can inhibit this function and suppress protein

localization to the plasma membrane. This autoinhibitory mechanism is counteracted by the GAP-

like and C-terminal domains, which promote ARHGAP36 recruitment to the plasma membrane

and primary cilium, respectively. Finally, we have discovered several residues within the GAP-

like domain that are necessary for Gli activation by full-length ARHGAP36 isoforms. These

residues are predicted to cluster within the GAP-like domain structure, at a site distal to the Rho

GTPase-binding pocket, and they are required for ARHGAP36 recruitment to the plasma

membrane. We have also leveraged these mutants to discover factors that bind specifically to the

wild-type protein, identifying potential mediators of ARHGAP36 function. Taken together, our work

supports a model in which ARHGAP36 activity state, subcellular localization, and effector binding

are regulated by structural elements distributed throughout protein. In combination with the

differential expression of ARHGAP36 isoforms, such mechanisms could allow ARHGAP36 to

control Gli activity and other PKA_{cat}-regulated processes in a tissue-specific manner.

RESULTS

Gli activation and cellular PKA$_{cat}$ depletion are separable ARHGAP36 functions

To explore the relationship between ARHGAP36-mediated Gli activation and PKA$_{cat}$ degradation, we measured the effect of each human ARHGAP36 isoform on both cellular processes. Individual isoforms were retrovirally transduced into NIH-3T3 mouse fibroblasts, a commonly used line for studying Hh signal transduction (*Taipale, J, et al., 2000*) that also exhibits ARHGAP36 responsiveness (*Eccles, et al., 2016; Rack, et al., 2014*). The resulting levels of *Gli1* mRNA and PKA$_{cat}$ protein were assessed by qRT-PCR and western blot, respectively. Overexpression of isoforms 2, 4, or 5 was sufficient to activate Gli and deplete the cells of PKA$_{cat}$, while isoform 1 exhibited neither activity (Figure 1B). In contrast, transduction of isoform 3 induced *Gli1* expression without reducing cellular PKA$_{cat}$ levels to a discernable extent. These results indicate that total PKA$_{cat}$ depletion is not required for ARHGAP36-mediated Gli activation, raising the possibility that ARHGAP36 regulates Gli proteins by targeting a specific subcellular pool of PKA$_{cat}$ and/or through PKA-independent mechanisms.

We investigated these two models by further comparing the activities of ARHGAP36 isoform 3 with those of isoform 2. Using immunofluorescence microscopy, we observed that isoform 2 globally depleted PKA$_{cat}$ in NIH-3T3 cells, corroborating our western blot analyses (Figure 1C). In contrast, isoform 3 reduced PKA$_{cat}$ pools predominantly in the Golgi. These findings are consistent with the accumulation of isoform 3 in the primary cilium (*Rack, et al., 2014*), as the cilium base communicates directly with the Golgi through vesicular trafficking (*Pedersen, et al., 2016*). We next examined how the activities of ARHGAP36 isoforms 2 and 3 are affected by forskolin, an adenylate cyclase agonist that increases cAMP levels and PKA$_{cat}$ activity. We transduced the ARHGAP36 constructs into NIH-3T3 fibroblasts stably expressing a Gli-dependent firefly luciferase reporter (SHH-LIGHT2 cells) (*Taipale, J, et al., 2000*) and treated the cells with the PKA$_{cat}$ activator. Although the isoform 2-expressing cells exhibited almost two-fold

Figure 1. ARHGAP36 isoforms differentially induce PKA_cat depletion and Gli activation. (A) Domain architecture of the five human ARHGAP36 isoforms, with the N-terminal domain (N) shown in dark blue, arginine-rich motif (R) in light blue, GAP-like domain in green, and C-terminal domain (C) in orange. The yellow region indicates an amino acid sequence unique to isoform 4. (B) Gli1 mRNA and PKA_cat protein levels in NIH-3T3 cells retrovirally transduced with the indicated FLAG-tagged ARHGAP36 isoforms. Uninfected cells treated with or without Sonic Hedgehog (SHH) ligand were included as positive and negative controls, respectively. Data are the average fold change relative to the negative control for three biological replicates ± s.e.m. Single and double asterisks indicate $P < 0.05$ and $P < 0.01$, respectively. A representative western blot for each condition is also shown. (C) PKA_cat localization in NIH-3T3 cells transduced with FLAG-tagged ARHGAP36 isoform 2 or 3. Representative immunofluorescence micrographs are shown with staining for PKA_cat, GM130 (cis-Golgi), and DAPI (nucleus). Scale bar: 20 μm. (D) Forskolin dose-response curves for SHH-LIGHT2 cells stimulated with SHH or transduced with FLAG-tagged ARHGAP36 isoform 2 or 3. Data are the average Gli reporter activities for at least three biological replicates ± s.e.m., normalized to the maximum response in SHH-treated cells.

higher Gli reporter activity than those expressing isoform 3, forskolin inhibited Gli function in both

lines with comparable IC50s (Figure 1D). Maximal doses of the PKA_{cat} activator also suppressed

about 50% of the Gli reporter activity induced by each isoform. Together, these results

demonstrate that N-terminal sequences in ARHGAP36 can regulate its ability to target PKA_{cat} in

specific subcellular compartments and raise the possibility that ARHGAP36 activates Gli proteins

through both PKA_{cat}-dependent and -independent mechanisms.

ARHGAP36 is autoinhibited by an N-terminal region that is counteracted by the GAP-like and C-

terminal domains

We continued to investigate functional differences between ARHGAP36 isoforms 2 and 3

by determining the activities of various truncation mutants (Figure 2A). By retrovirally expressing

these constructs in NIH-3T3 cells, we observed that the N-terminal domain of ARHGAP36

isoform 2 (residues 1-194; N2) is necessary and sufficient for its effects on Gli and PKA_{cat}

(Figure 2B) corroborating previous reports (*Eccles, et al., 2016*). However, N2 was less effective

at activating *Gli1* expression than the N-terminal domain of isoform 3 (N3), even though N2 could

induce PKA_{cat} degradation. N2 was also markedly less active than full-length isoform 2. In

contrast, N3 and full-length isoform 3 could induce *Gli1* expression to similar extents (Figure 2B).

These results indicate that the N2 region absent in isoform 3 (residues 1-105; $N2_{1-105}$) represses

the Gli-activating function of the remaining N-terminal domain. Moreover, our findings suggest

that ARHGAP36 sequences in the GAP-like and/or C-terminal domains can influence $N2_{1-105}$

function.

To discern how the GAP-like and C-terminal domains contribute to ARHGAP36 function,

we examined the activities of N2-GAP and N3-GAP constructs in NIH-3T3 cells. N2-GAP was

moderately more active than N2 with respect to *Gli1* expression, but it was still much less active

than full-length isoform 2 (Figure 2B). N3-GAP activity was comparable to that of N3 and full-

Figure 2. N-terminal, GAP-like, and C-terminal domains have opposing effects on ARHGAP36 function. (A) Schematic representation of ARHGAP36 isoform 2 or 3 truncation mutants. Residue numbers are based on the amino acid sequence of isoform 2. (B) *Gli1* mRNA and PKA$_{cat}$ protein levels in NIH-3T3 cell retrovirally transduced with the indicated FLAG-tagged ARHGAP36 truncation mutants. Data are the average fold change relative to uninfected cells for three biological replicates ± s.e.m. Single and double asterisks indicate $P < 0.05$ and $P < 0.01$, respectively. A representative western blot for each condition is also shown (lanes from the same blot image have been cropped and re-ordered for clarity). (C) Subcellular distributions of the indicated FLAG-tagged ARHGAP36 constructs in NIH-3T3 cells. Representative immunofluorescence micrographs are shown with staining for FLAG, ARL13B (primary cilium) and DAPI (nucleus). Insets highlight ciliated regions in the dashed boxes. Scale bar: 20 μm. Images were processed to establish comparable maximum pixel intensities in order to highlight differences in localization.

length isoform 3. Thus, both the GAP-like and C-terminal domains can counteract the

autoinhibitory function of N2$_{1-105}$, with the C-terminal region playing a particularly important role.

In the absence of N2$_{1-105}$, ARHGAP36 does not require its GAP-like and C-terminal domains to

achieve high levels of Gli activity.

We next sought to determine how the N2$_{1-105}$ region, GAP-like domain, and C-terminal

domain regulate ARHGAP36 function. The differing subcellular distributions of full-length

isoforms 2 and 3 indicate that the N2$_{1-105}$ region influences ARHGAP36 trafficking (*Rack, et al.,*

2014), and we therefore examined the localizations of the truncation mutants. N2 accumulated in

both punctate structures and the plasma membrane, whereas N2-GAP was robustly recruited to

the latter (Figure 2C). In comparison, both N3 and N3-GAP predominantly associated with the

plasma membrane, and unlike the full-length isoform 3, neither construct accumulated in the

primary cilium.

These domain-dependent changes in protein localization indicate that N2$_{1-105}$ impedes and

the GAP-like domain facilitates ARHGAP36 translocation to the plasma membrane. In principle,

these opposing activities could involve direct interactions between the two regions in ARHGAP36

or parallel functions involving other factors. In addition, the C-terminal domain acts independently

of the N2$_{1-105}$ region to promote ciliary accumulation of ARHGAP36. The disparate roles of these

domains in ARHGAP36 trafficking correlate with their divergent effects on Gli activation, providing

further evidence that ARHGAP36 targets specific subcellular compartments to regulate Gli

proteins.

Identification of ARHGAP36 residues that are required for Gli activation

To characterize these regulatory ARHGAP36 sequences with amino-acid resolution, we

developed a high-throughput mutagenesis screen for ARHGAP36 residues that are essential for

Gli activation (Figure 3A). We used error-prone PCR to create a library of ARHGAP36 isoform 2

Figure 3. Identification of essential residues within the N-terminal and GAP-like domains. (A) Schematic representation of the high-throughput mutagenesis screen used to identify individual residues that contribute to ARHGAP36 function. (B) Histogram depicting the fold change in mutation frequency between pre- and post-selected population for each residue in ARHGAP36. (C) Homology model of the ARHGAP36 GAP-like domain structure based on the crystal structure of β2-chimaerin (PDB ID: 1xa6). Residues with > 4-fold change in mutation frequency are shown in red, and the site (Thr227) that is structurally equivalent to the arginine finger is shown in blue. (D) Activities of selected ARHGAP36 variants in SHH-EGFP cells, as assessed by flow cytometry-based measurements of Gli reporter fluorescence. The distributions (left) and relative medians (right) of EGFP fluorescence are shown for each ARHGAP36 construct. Median fluorescence data represent the average values relative to that of cells expressing wild-type ARHGAP36 for three biological replicates ± s.e.m. Single and double asterisks indicate $P < 0.05$ and $P < 0.01$, respectively.

variants C-terminally fused to mCherry, obtaining a collection of approximately 100,000 single

point mutants (27% of all library constructs and 22-fold theoretical coverage of the 4,374 possible

variants). The library was retrovirally transduced into NIH-3T3 fibroblasts expressing a Gli-

dependent green-fluorescent reporter (SHH-EGFP cells) (*Hyman, et al., 2009*) using a multiplicity

of infection (MOI) of 0.3 to maximize the number of cells with single integration events. Cells

expressing full-length ARHGAP36 proteins were then isolated by fluorescence-activated cell

sorting (FACS) according to their mCherry fluorescence. We next cultured these cells under Hh

signaling-competent conditions to allow active ARHGAP36 mutants to induce Gli-dependent

EGFP expression, after which mCherry+/EGFP– cells were isolated by FACS. To ensure that

these ARHGAP36-expressing cells still harbored a functional EGFP reporter, they were

subsequently cultured with the SMO agonist SAG (*Chen, et al., 2002*). The resulting

mCherry+/EGFP+ cells were obtained by FACS, yielding a population of cells expressing putative,

inactive ARHGAP36-mCherry mutants.

To identify inactivating point mutations, we used genomic PCR and deep sequencing to

compare the mutation frequency of each amino acid position in the pre- and post-selection

populations. This analysis revealed several residues that could be required for Gli activation,

including two N-terminal arginines (R153 and R154) that were previously shown to be required

for ARHGAP36-mediated PKA_{cat} inhibition (Figure 3B, Supplementary File 1) (*Eccles, et al.,*

2016). The majority of these putative essential residues were located in the GAP-like region, and

structure homology modeling of this domain predicts that these amino acids cluster together at a

site that is distal to the predicted Rho GTPase binding pocket (Figure 3C). We then used flow

cytometry-based assays to validate a subset of these ARHGAP36 variants, prioritizing residues

that were mutated > 4-fold more frequently in the inactive mutant pool. In these experiments, 17

individual ARHGAP36 mutants were retrovirally transduced into cells at an MOI of 0.3 to

standardize their expression levels. With the exception of S25Y, N326S, and G510S, all of these

point mutations significantly decreased ARHGAP36 activity in SHH-EGFP reporter cells

(Figure 3D).

GAP-like domain mutations disrupt ARHGAP36 recruitment to the plasma membrane

271    Guided by the results of our mutagenesis screen, we investigated how individual point

mutations might regulate specific aspects of ARHGAP36 function. Our studies focused on the

three residues with the greatest fold change in pre- and post-selection mutation frequencies:

N-terminal domain residue L86 and two sites within the GAP-like domain, C212 and L317

(Figure 4A). We first examined how mutations at these sites affect the ability of ARHGAP36

isoform 2 to induce *Gli1* expression and PKA_{cat} degradation in NIH-3T3 cells. All three variants

exhibited diminished *Gli1* expression and PKA_{cat} depletion, with the L317P mutation resulting in

complete loss of both ARHGAP36-dependent activities (Figure 4B). We next determined the

effects of these mutations on the subcellular localization of ARHGAP36 isoform 2. The L86P

variant retained the ability to localize to the plasma membrane, but both mutations in the GAP-

like domain rendered ARHGAP36 cytosolic (Figure 4C).

282    To further elaborate the contributions of individual residues to ARHGAP36 trafficking, we

also assessed the localization of other ARHGAP36 point mutants that were inactive in SHH-EGFP

cells (see Figure 3D). The R153W and R154Q variants accumulated in the plasma membrane,

and the GAP-like domain point mutants M387R and L394R remained primarily in the cytosol

(Figure 4 – figure supplement 1). Together, these results provide further evidence that the GAP-

like domain recruits ARHGAP36 to the cell membrane, counteracting the function of $N2_{1-105}$. We

then investigated whether plasma membrane recruitment by the GAP-like domain requires $N2_{1-105}$

or the C-terminal domain, both of which affect ARHGAP36 localization. As in the full-length

isoform 2, C212Y and L317P mutations impeded the ability of N2-GAP to localize to the plasma

membrane and activate Gli proteins (Figure 5A-B). Structurally equivalent mutations (C107Y and

Figure 4. Point mutations in the GAP-like domain cause ARHGAP36 mislocalization. (A) Schematic representation of ARHGAP36 isoform 2 point mutants. (B) *Gli1* mRNA and PKA$_{cat}$ protein levels in NIH-3T3 cells retrovirally transduced with the indicated FLAG-tagged ARHGAP36 constructs. Uninfected cells were used as controls. Data are the average fold change relative to cells expressing wild-type ARHGAP36 (*Gli1* expression) or to untreated cells (PKA$_{cat}$ levels) for three biological replicates ± s.e.m. Single and double asterisks indicate $P < 0.05$ and $P < 0.01$, respectively. A representative western blot for each condition is also shown (lanes from the same blot image have been cropped and re-ordered for clarity). (C) Subcellular distributions of the indicated FLAG-tagged ARHGAP36 constructs in NIH-3T3 cells. Representative immunofluorescence micrographs are shown with staining for FLAG and DAPI (nucleus). Scale bar: 20 μm.

Figure 5. Point mutations in the GAP-like domain act independently of N21-105 and the C-terminal domain. (A and C) Effects of GAP-like domain point mutations in N2-GAP- or isoform 3-mediated Gli activation, as assessed in the flow cytometry-based SHH-EGFP assay. The distributions (left) and relative medians (right) of EGFP fluorescence are shown for each ARHGAP36 construct. Median fluorescence data represent the average values relative to that of cells expressing unmutated N2-GAP or isoform 3 for three biological replicates ± s.e.m. Double asterisks indicate $P < 0.01$. (B and D) Subcellular distributions of the indicated mCherry-tagged ARHGAP36 constructs in SHH-EGFP cells. Representative immunofluorescence micrographs are shown with staining for mCherry, ARL13B (primary cilium), and DAPI (nucleus). Scale bars: 20 μm.

L212P) in ARHGAP36 isoform 3, which lacks the $N2_{1-105}$ region, also rendered the protein

cytosolic and attenuated Gli activation (Figure 5C-D). Thus, the two GAP-like domain mutations

disrupt ARHGAP36 function independently of $N2_{1-105}$ and the C-terminal domain.

GAP-like domain mutations alter the ARHGAP36 interactome

In addition to uncovering functional roles for specific ARHGAP36 structures, our collection

of inactive variants provided a means for identifying binding proteins that could participate in

ARHGAP36-mediated Gli activation. Toward this goal, we compared the interactomes of wild-

type and L317P ARHGAP36 isoform 2 by retrovirally transducing NIH-3T3 cells with vectors

encoding each ARHGAP36 construct fused to a C-terminal LAP tag (S-peptide-PreScission

protease site-EGFP) (*Ding, et al., 2016; Hsu, et al., 2019; Kanie, et al., 2017; Li, Bin, et al., 2017;*

Torres, et al., 2009; Wright, et al., 2011) (Figure 6A). To avoid total PKA_{cat} depletion by wild-type

ARHGAP36 in these studies, we also limited the cells to a 4-hour incubation in retroviral medium

and a subsequent 20-hour growth phase. The fibroblasts were then lysed, and each ARHGAP36

construct and its interacting proteins were isolated by tandem affinity purification and

proteolytically digested as previously described (*Ding, et al., 2016; Hsu, et al., 2019; Kanie, et al.,*

2017; Li, Bin, et al., 2017; Torres, et al., 2009; Wright, et al., 2011). The resulting peptides were

sequenced and quantified using tandem mass spectrometry, and spectral counts were normalized

to account for variabilities in protein size, LAP tag purification efficiency, and ARHGAP36

expression. Using this approach, we identified 566 putative interactors for wild-type and/or L317P

ARHGAP36 that were observed across three biological replicates (Figure 6B, Supplementary

File 2). PKA_{cat} subunits were the only canonical Hh pathway regulators detected in these pulldown

experiments, and they interacted with wild-type and L317P ARHGAP36 to similar extents

(Figure 6B, Figure 6 – figure supplement 1). Eleven proteins preferentially bound to wild-type

ARHGAP36 by at least 5-fold, with prolyl oligopeptidase-like protein PREPL and the E3 ubiquitin

Figure 6. PREPL and PRAJA2 interact with ARHGAP36 in a GAP-like domain-dependent manner. (A) Schematic representation of the tandem affinity purification workflow for identifying ARHGAP36-binding proteins. (B) ARHGAP36 interactors ranked in order of their relative binding to the wild-type versus L317P proteins. Fold changes in normalized spectral counts represent the average values for three biological replicates ± s.e.m., shown as grey bars.

ligase PRAJA2 exhibiting the greatest selectivity (52- and 35-fold, respectively). PREPL displays

high sequence homology to a family of serine hydrolases, though its physiological substrates and

functions remain largely unknown (*Jaeken, et al., 2006; Radhakrishnan, et al., 2013; Szeltner, et*

al., 2005). PRAJA2 has been shown to increase PKA activity by promoting the ubiquitination and

degradation of regulatory PKA (PKA_{reg}) subunits, a function that is enhanced by PKA_{cat}

phosphorylation as part of a positive-feedback mechanism (*Lignitto, et al., 2011*).

Both PREPL and PRAJA2 have been classified as putative ARHGAP36-binding partners

in large-scale interactome studies (*Huttlin, et al., 2017; Müller, et al., 2020*); however, their

functional significance relative to the other candidates in those lists has been unclear. Our

comparative interactome analyses therefore corroborate the results of these prior investigations

and suggest that ARHGAP36-dependent Gli activation involves PREPL and PRAJA2 functions.

DISCUSSION

By systematically exploring the ARHGAP36 structure-activity landscape, we have

identified key functional elements throughout this multidomain protein. Previous investigations

identified an N-terminal arginine-rich region that is necessary and sufficient for PKA_{cat} degradation

(*Eccles, et al., 2016; Rack, et al., 2014*), establishing ARHGAP36 as a novel antagonist of PKA

signaling. Isoform-specific differences have also implicated N-terminal sequences in ARHGAP36

trafficking (*Rack, et al., 2014*). Our studies provide further evidence for these mechanisms of

ARHGAP36 action, uncover a new functional module within the N-terminal domain, and

demonstrate important regulatory roles for the GAP-like and C-terminal domains (Figure 7).

Among our key findings is the discovery of an N-terminal autoinhibitory region that is

present in isoforms 1 and 2. In the context of isoform 2, this sequence ($N2_{1-105}$) represses Gli1

activation by the N-terminal domain and impedes its recruitment to the plasma membrane. $N2_{1-105}$

likely regulates these activities through distinct mechanisms, since the L86P mutation in this

Figure 7. A regulatory model for ARHGAP36 function. Schematic representation of the structural elements that regulate ARHGAP36 localization and its ability to target Gli-regulating pools of PKA$_{cat}$ in the primary cilium. N-terminal regulatory regions that vary between ARHGAP36 isoforms are depicted with hash marks.

region diminishes the ability of isoform 2 to modulate PKA_{cat}/Gli signaling without disrupting

membrane localization. ARHGAP36 isoform 3, which lacks the region corresponding to $N2_{1-105}$,

not only traffics to the plasma membrane but also accumulates in the primary cilium. The ciliary

localization of isoform 3 coincides with its ability to selectively degrade Golgi-localized PKA_{cat},

likely due to vesicular transport between the cilium base and the Golgi (*Pedersen, et al., 2016*).

Our results suggest that the Gli-regulating pool of PKA_{cat} traffics through these organelles,

corroborating previous reports that Hh signaling regulates PKA_{cat} activity in the basal body (*Barzi,*

et al., 2009; Tuson, et al., 2011; Zhang, et al., 2019). This mechanism of ARHGAP36-mediated

Gli activation parallels the actions of other Hh pathway regulators that modulate ciliary PKA

activity, such as adenylate cyclases 5 and 6 and the G-protein coupled receptor GPR161

(*Chávez, et al., 2015; Garcia-Gonzalo, et al., 2015; Moore, et al., 2016; Mukhopadhyay, et al.,*

2013; Vuolo, et al., 2015). ARHGAP36 might concurrently modulate Gli function via PKA-

independent mechanisms, since forskolin can only inhibit ARHGAP36-induced Gli activity by

50%, even when maximal compound doses and different levels of Gli activation are employed.

Although the ARHGAP36 N-terminal domain alone can target PKA_{cat} for degradation and

harbors regulatory elements, its function is further modulated by other structures in the full-length

protein. The C-terminal domain strongly counteracts the autoinhibitory activity of $N2_{1-105}$, as does

the GAP-like region to a more moderate extent. These regulatory actions likely involve protein

trafficking since the GAP-like and C-terminal domains promote ARHGAP36 recruitment to the

plasma membrane and primary cilium, respectively. However, we cannot rule out the possibility

that the two domains influence $N2_{1-105}$ function through additional mechanisms. Interestingly, the

ability of point mutations in the GAP-like domain to abrogate Gli activation by isoforms 2 and 3

contrasts the sufficiency of N2 and N3 for this process. The mutations also render these

ARHGAP36 proteins cytosolic. These differences could be explained if the unbound GAP-like

domain can repress the N-terminal region conserved between N2 and N3, which contains the

PKA$_{cat}$-targeting arginine-rich motif and a plasma membrane-targeting sequence. Recruitment of

specific cellular factors to the GAP-like domain could then modulate this interaction and possibly

mediate other ARHGAP36 activities (see Figure 7).

Which interacting proteins might regulate or transduce ARHGAP36 function remains to be

determined, but our investigations provide valuable leads. By comparing the interactomes of wild-

type ARHGAP36 and an inactive mutant, we have identified several proteins that bind to

ARHGAP36 in a GAP-like domain-dependent manner. Although previous studies have shown

that ARHGAP36 can co-immunoprecipitate with PTCH1 (*Zhang, et al., 2019*) and SUFU (*Rack,*

et al., 2014) when they are overexpressed in cells, we do not observe analogous interactions with

the endogenous Hh signaling proteins. Rho GTPases were also notably absent from the

ARHGAP36 pulldowns, suggesting that ARHGAP36 interacts transiently with these signaling

proteins or prefers other binding partners. In contrast, PKA$_{cat}$ associated with both wild-type and

L317P ARHGAP36 with comparable efficacies, indicating that the GAP-like domain mutation

abrogates PKA$_{cat}$ degradation without compromising its binding. Among the ARHGAP36

interactors discovered in our studies, PREPL and PRAJA2 emerged as the two most sensitive to

the L317P mutation in the GAP-like domain. Both proteins were candidates in previous

ARHGAP36 interactome datasets (*Huttlin, et al., 2017; Müller, et al., 2020*), but the functional

relevance of these factors and other ARHGAP36-binding proteins has yet to be determined. By

comparing the wild-type and mutant ARHGAP36 interactomes, we can provide a functional

context for these binding proteins, implicating PREPL and PRAJA2 in ARHGAP36-dependent Gli

activation.

In principle, the GAP-like domain could bind directly to PREPL or PRAJA2, or it could

allosterically regulate their interactions with other regions in the ARHGAP36 protein. PREPL is

highly homologous to the prolyl oligopeptidase (PREP) family of serine hydrolases, but its

hydrolytic targets remain uncharacterized (*Jaeken, et al., 2006; Szeltner, et al., 2005*). Putative

peptidic substrates could include factors that regulate PKA_{cat} or Gli activity. Alternatively, there is

evidence that PREP enzymes can modulate the metabolism of phosphoinositides (*Schulz, et al.,*

2002; Williams, et al., 1999), a family of lipids with reported roles in the formation of Golgi-derived

vesicles (*Heldwein, et al., 2004; Wang, Ying Jie, et al., 2003*) and the ciliary trafficking of Hh

pathway regulators (*Chávez, et al., 2015; Garcia-Gonzalo, et al., 2015*). The interaction of

ARHGAP36 with PRAJA2 is seemingly paradoxical since PRAJA2 has well-established roles in

promoting PKA_{reg} degradation and PKA_{cat} activation (*Lignitto, et al., 2013; Lignitto, et al., 2011;*

Sepe, et al., 2014). However, one possibility is that PRAJA2-mediated PKA_{reg} degradation leads

to the mislocalization of Gli-regulating pools of PKA_{cat}. We note that neither PREPL nor PRAJA2

were identified as hits in CRISPR knockout screens for Hh pathway regulators (*Breslow, et al.,*

2018; Pusapati, et al., 2018), suggesting that they participate in an ARHGAP36-specific pathway

for Gli activation. Determining how these factors contribute to ARHGAP36 action could uncover

novel aspects of Gli regulation.

Taken together, our findings provide new insights into the mechanisms that underlie non-

canonical Gli activation by ARHGAP36, and they provide a general framework for understanding

ARHGAP36 function. In particular, our studies reveal how ARHGAP36 can translate multiple

cellular inputs into distinct signaling outputs. By controlling the expression of ARHGAP36 isoforms

and/or ARHGAP36-interacting proteins, cells can direct this signaling protein to specific PKA_{cat}

populations and elicit tissue-specific responses. We anticipate that these mechanisms not only

contribute to Gli-dependent spinal cord development and medulloblastoma progression but also

other PKA_{cat}-dependent processes in normal physiology and disease. Moreover, the experimental

methods utilized for our study could be redeployed to elucidate these unique ARHGAP36

functions.

METHODS

Reagents and cell lines

Antibody sources and working dilutions are listed in Supplementary File 3. Forskolin was purchased from Calbiochem, and SAG was purchased from Tocris. SHH-LIGHT2 (*Taipale, J, et al., 2000*) and SHH-EGFP cells (*Hyman, et al., 2009*) were described previously, and NIH-3T3 and HEK-293T cells were purchased from the American Type Culture Collection.

Expression vectors

The pDONR223 vector was provided by J. Hartley and D. Esposito. The following constructs have been described previously (*Rack, et al., 2014; Wright, et al., 2011*): Gateway cloning destination vectors pBMN-3xFLAG-IRES-mCherry-DEST, pBMN-mCherry-DEST, and pG-LAP7-DEST, Gateway cloning entry vectors pDONR223-ARHGAP36 (isoforms 1 - 5), and expression vectors pcDNA3.2-ARHGAP36 (isoform 2)-V5, pBMN-3xFLAG-IRES-mCherry, and pBMN-mCherry. pCL-ECO retrovirus packaging vector was purchased from Imegenex.

Retroviral expression vectors for FLAG-tagged ARHGAP36 isoforms 1 - 5 were produced by transferring cDNAs from the appropriate pDONR223 entry vectors into pBMN-3xFLAG-IRES-mCherry-DEST in an LR Clonase II (Invitrogen, Waltham, MA)-mediated recombination reaction. Retroviral expression vectors for mCherry-tagged ARHGAP36 isoforms 1 and 2 were produced in an analogous manner with pBMN-mCherry-DEST. Restriction sites were then inserted upstream (XhoI) and downstream (SacII) of the *ARHGAP36* sequence in the initial pBMN-ARHGAP36 (isoform 1)-mCherry and pBMN-ARHGAP36 (isoform 2)-mCherry products. This was achieved by first amplifying *ARHGAP36* cDNA from the pBMN-derived vectors (Supplementary File 3; primers 1 - 3) and then inserting the resulting PCR product into BamHI-digested pBMN-mCherry using Gibson assembly (New England Biolabs, Ipswich, MA). The resulting pBMN-

ARHGAP36-mCherry vectors with XhoI and SacII restriction sites were subsequently used in all

experiments described herein.

Retroviral expression vectors for ARHGAP36 isoform 2 truncation mutants were

generated by amplifying the cDNA for each variant from pcDNA3.2-ARHGAP36 (isoform 2)-V5,

using primers containing attB adapter sequences (Supplementary File 3; primers 4-8). The PCR

products were transferred into pDONR223 in a BP Clonase II (Invitrogen)-mediated

recombination reaction, and the *ARHGAP36*-derived cDNAs in these pDONR223 entry vectors

were then transferred to pBMN-3xFLAG-IRES-mCherry-DEST using LR Clonase II. The resulting

constructs were also used as templates to amplify cDNAs for the analogous FLAG-tagged

ARHGAP36 isoform 3 truncation mutants and the downstream IRES-mCherry sequence

(Supplementary File 3; primers 9 and 10). These PCR products were then inserted into XcmI-

digested pBMN-ARHGAP36 (isoform 2)-3xFLAG-IRES-mCherry using Gibson assembly.

Individual ARHGAP36 isoform 2 point mutants, with the exception of L171P, were

generated using site-directed mutagenesis with PfuUltra II Fusion polymerase (Agilent, Santa

Clara, CA) (Supplementary File 3; primers 11-42) and either pBMN-ARHGAP36 (isoform 2)-

3xFLAG-IRES-mCherry or the pBMN-ARHGAP36 (isoform 2)-mCherry as template. ARHGAP36

isoform 2 L171P mutant constructs were generated by Gibson assembly using a XhoI- and SacII-

digested pBMN-ARHGAP36 (isoform 2)-mCherry plasmid, inserts amplified from pBMN-

ARHGAP36 (isoform 2)-mCherry (Supplementary File 3; primers 3 and 43-45), and a double-

stranded oligonucleotide encoding the L171P mutation (Integrated DNA Technologies, Coralville,

IA) (Supplementary File 3; entry 46) .

Retroviral expression vectors for mCherry-tagged wild-type, C107Y, and L212P

ARHGAP36 isoform 3 were generated by amplifying the cDNAs encoding this isoform from the

corresponding mutant pBMN-ARHGAP36 (isoform 2)-mCherry plasmids (Supplementary File 3;

primers 3 and 47) and amplifying the mCherry tag from pBMN-ARHGAP36 (isoform2)-mCherry

plasmid (Supplementary File 3; primers 10 and 48). The resulting amplicons were inserted into

XcmI-digested pBMN-ARHGAP36 (isoform 2)-3xFLAG-IRES-mCherry using Gibson assembly.

Retroviral expression vectors for mCherry-tagged wild-type, C212Y, and L317P ARHGAP36 N2-

GAP were generated by amplifying the cDNAs encoding ARHGAP36 N2-GAP from the

corresponding mutant pBMN-ARHGAP36 (isoform 2)-mCherry plasmids (Supplementary File 3;

primers 45 and 49) and inserting into XhoI- and SacII-digested pBMN-ARHGAP36 (isoform 2)-

mCherry using Gibson assembly.

To generate retroviral expression vectors for LAP-tagged ARHGAP36 constructs,

ARHGAP36 cDNA in the pDONR223-ARHGAP36 (isoform 2) entry vector was transferred to pG-

LAP7-DEST using LR Clonase II. The cDNA encoding LAP-tagged ARHGAP36 was amplified

from the resulting pG-ARHGAP36 (isoform 2)-LAP7 plasmid (Supplementary File 3; primers 10

and 50) and inserted into XcmI-digested pBMN-ARHGAP36 (isoform 2)-3xFLAG-IRES-mCherry

using Gibson assembly. LAP-tagged L317P ARHGAP36 isoform 2 retroviral expression

constructs were generated using an analogous Gibson assembly with *ARHGAP36* cDNA

amplified from pBMN-ARHGAP36 (isoform 2 L317P)-3xFLAG-IRES-mCherry (Supplementary

File 3; primers 50 and 51) and LAP tag cDNA amplified from pG-ARHGAP36 (isoform 2)-LAP7

(Supplementary File 3; primers 10 and 48).

With the exception of the constructs generated by site-directed mutagenesis described

above, all PCR products were generated with Phusion polymerase (New England Biolabs). All

plasmids were sequence-verified.

Retrovirus production

HEK-293T cells were seeded into individual wells of a 6-well plate at a density of

1.0×10^6 cells/well. The cells were cultured for 24 hours in HEK-293T growth medium (DMEM

containing 10% fetal bovine serum, 2 mM L-glutamine, 1 mM sodium pyruvate, 100 U/mL

penicillin, and 0.1 mg/mL streptomycin) and then transfected as follows. pBMN vectors containing

the appropriate ARHGAP36 construct (1.33 µg) and the pCL-ECO retrovirus packaging vector

(0.67 µg) were diluted in OMEM medium (75 µL), and the solution was added to OMEM (75 µL)

containing 6 µL Fugene HD reagent (Promega, Madison, WI). The mixture was incubated at room

temperature for 10 minutes and gently added to the growth medium on the cultured cells. After

24 hours, the medium was replaced with DMEM containing 1.8 mM L-glutamine, 4% fetal bovine

serum, 6% calf serum, 1 mM sodium pyruvate, 100 U/mL penicillin, and 0.1 mg/mL streptomycin.

Retrovirus-containing supernatant was then collected two times at 20-hour intervals, passed

through a 0.45-µm filter, and stored at –80 °C. Large-scale retrovirus production was conducted

using HEK-293T cells seeded on 10-cm plates at a density of 5.0×10^6 cells/well transfected with

6.48 µg of the ARHGAP36 construct, 4 µg of pCL-ECO, and 35 µL of Fugene HD in 750 µL

OMEM medium.

Generation of ARHGAP36-expressing NIH-3T3 cell lines

NIH-3T3 cells were seeded into individual wells of 24-well or 6-well plates at a density of

7.5×10^4 or 2.0×10^5 cells/well, respectively. The cells were cultured for 24 hours in NIH-3T3

growth medium (DMEM containing 10% calf serum, 1 mM sodium pyruvate, 100 U/mL penicillin,

and 0.1 mg/mL streptomycin) and then transduced with 4 µg/mL polybrene and retrovirus for the

appropriate ARHGAP36-3xFLAG-IRES-mCherry construct to achieve a multiplicity of infection

(MOI) < 0.5. After 24 hours, the medium was exchanged, and the cells were expanded for

fluorescence-activated cell sorting (FACS).

For FACS, the cells were washed with PBS buffer, dissociated with TrypLE (Invitrogen)

for 3-5 minutes at 37 °C, and centrifuged at 106 *g* for 5 minutes at 4 ºC. Cell pellets were then

resuspended in FACS buffer (PBS containing 1% calf serum), passed through a 70-µm cell

strainer (BD Biosciences, San Jose, CA), and added to round-bottom FACS tubes. Cell

populations with comparable mCherry fluorescence intensities were then obtained using a BD

FACSAria II (532-nm laser and 600-nm longpass filter, or 561-nm laser and 610/20-nm bandpass

filter) or BD Influx (561-nm laser and 610/20-nm bandpass filter) sorter.

Quantitative reverse transcription-PCR (qRT-PCR) analyses

NIH-3T3 cell lines expressing the indicated FLAG-tagged ARHGAP36 constructs were

seeded into 6-well plates at a density of 5.2 x 10^5 cells/well and cultured in NIH-3T3 growth

medium. An uninfected condition was also prepared as a negative control. After two days, fully

confluent cells were treated with NIH-3T3 low-serum medium (DMEM containing 0.5% calf serum,

sodium pyruvate and antibiotics) with or without 10% SHH-N-conditioned medium for 30 hours.

The media was replaced with ice-cold PBS, and the cells were collected by manual scraping.

Each resulting cell suspension was divided into two tubes (one each for qRT-PCR and Western

blot analyses) and centrifuged at 750 *g* for 7 minutes at 4 °C.

Cell pellets were prepared for qRT-PCR analyses as follows. RNA was isolated using the

Monarch Total RNA miniprep kit (New England BioLabs), and equivalent amounts of RNA were

used to synthesize cDNA using the SuperScript III First-Strand Synthesis System (Invitrogen).

qRT-PCR was performed on a Lightcycler 480 II (Roche, Penzberg, Germany) using the following

TaqMan probes: Gli1-Mm00494645_m1, Beta-2-Microglobulin-Mm00437762_m1 (Applied

Biosystems, Waltham, MA). Gene expression levels were normalized to β-2-microglobulin. For

ARHGAP36 isoform and truncation mutant analyses, the normalized gene expression levels were

compared to that of uninfected cells, and for point mutant analyses, compared to that of wild-type

ARHGAP36. The resulting gene expression levels were averaged across three biological

replicates, and *P* values were determined using either a Student's one-tailed t-test (ARHGAP36

isoform and truncation mutant analyses) or two-tailed t-test (point mutant analyses).

 Western blot analyses

Cell pellets were resuspended in 1x Laemmli sample buffer (10% glycerol, 2% SDS,

17 mM DTT, 0.01% bromophenol blue, 60 mM Tris-HCl, pH 6.8, and protease and phosphatase

inhibitors (Roche)). After incubation for 20 minutes at 4 ºC, cell lysates were boiled for 10 minutes

and sonicated in a water bath for 15 seconds. Equivalent amounts of total protein per lysate were

loaded onto Criterion XT 4-12% Bis-Tris polyacrylamide gels (Bio-Rad, Hercules, CA), transferred

onto PVDF membranes (Bio-Rad), and detected using the antibodies listed in Supplementary

File 3 with either SuperSignal West Dura or SuperSignal Femto kits (Pierce, Waltham, MA) and a

ChemiDoc XRS imaging system (Bio-Rad). Band intensities were quantified using ImageLab

software (Bio-Rad) and normalized to KPNB1 band intensities in the corresponding sample. For

each replicate, the normalized band intensity in each condition was normalized to that of

uninfected cells. The resulting relative band intensities for each condition was averaged across

three biological replicates, and *P* values were determined using a Student's two-tailed t-test.

Immunofluorescence studies

The subcellular localizations of FLAG-tagged ARHGAP36 constructs were assessed as

follows. NIH-3T3 cells were seeded onto individual wells of a 6-well plate at a density of

2.0×10^5 cells/well. Cells were cultured for 24 hours in NIH-3T3 growth medium, then transduced

with 4 µg/mL polybrene and retrovirus for the appropriate ARHGAP36-3xFLAG-IRES-mCherry

construct to achieve an MOI < 0.5. After 24 hours, cells were re-seeded at a 1:8 dilution into 24-

well plates containing poly-D-lysine-coated 12-mm glass coverslips and cultured for 1-2 days in

growth medium. Cells were fixed in PBS containing 4% paraformaldehyde for 10 minutes at room

temperature and washed 3 times with PBS. Cells were next permeabilized with PBS containing

0.5% Triton X-100 for 5 minutes, washed 2 times with PBS, and incubated in blocking buffer (PBS

containing 1% BSA and 0.1% Triton X-100) for 1 hour at room temperature. The cells were then

incubated for 1 hour at room temperature with primary antibodies diluted in blocking buffer,

washed 4 × 5 minutes with PBS containing 0.1% Triton X-100, incubated for 1 hour with the

appropriate secondary antibodies diluted in PBS containing 0.2% Triton X-100, and washed 4 ×

5 minutes with PBS. The coverslips were rinsed briefly in water and mounted onto slides using

Prolong Gold Antifade reagent with DAPI (Invitrogen).

The subcellular localizations of mCherry-tagged ARHGAP36 isoform 3 and N2-GAP

constructs were assessed as follows. SHH-EGFP cells were seeded into individual wells of a 24-

well plate at a density of 7.5 x 10^4 cells/well. The cells were cultured for 24 hours in SHH-EGFP

growth medium (NIH-3T3 growth medium containing 150 µg/mL zeocin) for 24 hours and then

transduced with 4 µg/mL polybrene and retrovirus for the appropriate ARHGAP36-mCherry

construct to achieve an MOI < 0.5. After 24 hours, the cells were passaged into a new 24-well

plate at a 1:1.5 dilution and cultured in growth medium for an additional 2 days to achieve 100%

confluency. Confluent cells were then treated for 24 hours with SHH-EGFP low-serum medium

(DMEM containing 0.5% calf serum, sodium pyruvate, zeocin, and antibiotics). Cells were next

passaged at a 1:3 dilution into 24-well plates containing poly-D-lysine-coated 12-mm glass

coverslips and cultured for 1 day in growth medium. Cells were then fixed, blocked,

immunostained, and mounted as described above. The subcellular localizations of mCherry-

tagged isoform 2 constructs were similarly assessed, with the exception that the cells were

passaged onto coverslip-containing 24-well plates 24 hours after retroviral transduction then

cultured for two days prior to being fixed.

The effects of ARHGAP36 isoforms on PKA_{cat} localization were assessed as follows. NIH-

3T3 cell lines stably expressing FLAG-tagged ARHGAP36 isoforms 2 or 3 were seeded into 24-

well plates containing poly-D-lysine-coated 12-mm glass coverslips at a density of 1.2 x 10^5

cells/well. Cells were then cultured in NIH-3T3 growth medium for 2 days, at which time the cells

were fixed in PBS containing 2% paraformaldehyde for 20 minutes at room temperature and then

treated with methanol for 5 minutes at –20 ºC. The fixed cells were incubated in blocking buffer

for 1 hour at room temperature and incubated with primary antibodies diluted in blocking buffer

overnight at 4 ºC. Subsequent PBS washes, secondary antibody incubation, and mounting were

conducted as described above.

Fluorescence images were obtained using either a Zeiss LSM 700 or 800 confocal

microscope equipped with a 63x oil-immersion objective. Maximum-intensity Z-stack projections

were created using either ZEN Black (Zeiss, Oberkochen, Germany), ZEN Blue (Zeiss), or FIJI

(*Schindelin, et al., 2012*) software, fluorescence intensities were adjusted using FIJI, and images

were cropped using Photoshop CC (Adobe, San Jose, CA).

Luciferase assays

NIH-3T3 cells were seeded into 10-cm plates at a density of 1.0×10^6 cells/plate and

cultured in NIH-3T3 growth medium. After 24 hours of growth, cells were transduced with 4 µg/mL

polybrene and retrovirus for either the appropriate ARHGAP36-3xFLAG-IRES-mCherry construct

or for a 3xFLAG-IRES-mCherry construct for another 24 hours. The cells were then re-seeded

into individual wells of a 96-well plate at a density of 3.5×10^4 cells/well. All cells were cultured for

an additional 24 hours, at which time the fully confluent cells were treated for 30 hours with varying

conditions of forskolin in NIH-3T3 low-serum medium with or without 10% SHH-N-conditioned

medium. The cells were then lysed and luciferase activities were measured using a Dual-

Luciferase Reporter Assay System (Promega) on a Veritas luminometer (Turner BioSystems,

Sunnyvale, CA). At least three technical replicates were conducted for each condition.

Mutant library generation

A library encoding ARHGAP36 isoform 2 mutants was created via error-prone PCR

(epPCR) using the GeneMorph II Random Mutagenesis Kit (Agilent). To determine the optimal

epPCR conditions for library generation, the mutation frequency was estimated for epPCRs

consisting of 15, 20, 25, or 29 cycles. All reactions were conducted according to the

manufacturer's instructions, using 1.64 µg of the pBMN-ARHGAP36 (isoform2)-mCherry plasmid

as template, 0.4 µM of each primer (Supplementary File 3; primers 45 and 52), and 4% DMSO.

The product yield for each condition was estimated by resolving 10% of the reaction on a 1%

EtBr-agarose gel and quantifying the band intensity of the resulting amplicon with ImageLab

software (Bio-Rad). The 1.6-kb amplicon was gel-extracted using the QiaQuick Gel Extraction Kit

(Qiagen, Hilden, Germany) and ligated into a XhoI- and SacII-digested pBMN-ARHGAP36

(isoform 2)-mCherry vector using Gibson assembly. To estimate the mutation frequency for each

epPCR-generated library, XL-10 Gold *E. coli (*Agilent) were chemically transformed with 1:4

diluted Gibson assembly products and plated onto ampicillin-agarose plates. Forty colonies from

each plate were sequenced using rolling circle amplification and Sanger sequencing (Sequetech,

Mountain View, CA) (Supplementary File 3; primers 53-58). High-quality *ARHGAP36* reads were

aligned to the coding sequence for wild-type ARHAGP36 isoform 2, and the number of nucleotide

mutations within the coding sequence was counted for each read. This analysis yielded the

distribution of mutated nucleotides across the library.

Through these pilot studies, we found that the 15-cycle epPCR conditions maximized the

percentage of *ARHGAP36* variants with single-nucleotide changes. We next generated a large-

scale library using the 15-cycle epPCR and Gibson assembly strategy described above. The

undiluted Gibson reaction (8 µL) was electroporated into 160 µL of MegaX DH10B T1^R

Electrocomp cells (Invitrogen). The electroporated cells were immediately transferred to 480 mL

of Superior Broth containing 75 µg/mL ampicillin, and 100 µL of the culture was plated on

ampicillin-agar plates to estimate the number of colony-forming units. The final library was found

to contain approximately 4×10^5 colony-forming units, which corresponds to an equivalent number

of library elements. The liquid culture was incubated at 30 ºC until it reached an OD_{600} of 1, after

which plasmids were isolated using the NucleoBond Xtra Midi Plus Kit (Macherey-Nagel, Düren,

Germany).

Retroviral medium harboring the ARHGAP36 mutant library was generated in the following

manner. One 10-cm plate of HEK-293T cells at 90% confluency was transfected with 6.5 µg of

the pBMN-ARHGAP36 (isoform 2)-mCherry mutant library and 4 µg of pCL-ECO using the

FuGene HD transfection reagent (Promega). The medium was replaced after 24 hours with

DMEM containing 1.8 mM L-glutamine, 4% fetal bovine serum, 6% calf serum, 1% sodium

pyruvate, 100 U/mL penicillin, and 0.1 mg/mL streptomycin. Retrovirus-containing supernatant

was then collected two times at 24-hour intervals, passed through a 0.45-µm filter, and stored at

−80 °C.

FACS-based screening

SHH-EGFP cells were seeded onto a 15-cm plate at a density of 1.0×10^6 cells/plate and

cultured in SHH-EGFP growth medium for 2 days, then transduced with the retroviral library of

ARHGAP36 mutants and 4 µg/mL polybrene to achieve an MOI < 0.5. After 24 hours, the cells

were expanded to 4 x 15-cm plates and cultured for an additional 2 days. SHH-EGFP cells treated

with 10% SHH-N-conditioned media or transduced with retrovirus encoding wild-type ARHGAP36

isoform 2 served as positive controls for flow cytometry. For negative controls, untreated SHH-

EGFP cells or cells transduced with retrovirus encoding ARHGAP36 isoform 1 were used.

To isolate mCherry+ cells by FACS, the transduced SHH-EGFP cells were washed with

PBS, dissociated with TrypLE for 3-5 minutes at 37° C, and centrifuged at 750 *g* for 7 minutes at

4 °C. The resulting cell pellets were resuspended in FACS buffer (PBS containing 1% calf serum),

passed through a 70-µm cell strainer, and added to round-bottom FACS tubes. Cell sorting was

performed on a BD FACSAria II configured with a 561-nm laser, a 595-nm longpass filter, and a

616/23-nm bandpass filter for mCherry detection. Data was collected with FACSDiva software

(BD Biosciences) and analyzed using FlowJo (FlowJo, Ashland, Oregon). 2.7×10^7 cells were

analyzed by FACS analysis.

This first sort produced a population of 3×10^6 mCherry+ cells, which were cultured for

3 days until they reached 100% confluency. The cells were then cultured in SHH-EGFP low-

serum medium to enable ARHGAP36-mediated Gli activation. After 24 hours, cells were

expanded at a 1:2 dilution, cultured for 24 hours to achieve full confluency, and serum-starved

again for another 24 hours.

To isolate cells expressing inactive forms of ARHGAP36 isoform 2 (mCherry+/EGFP–),

1.1×10^7 cells from the expanded mCherry+ population were washed and dissociated as

described above. Approximately 3×10^6 cells were separated and expanded to 4×10^7 cells to

establish a pre-selection population, which was then washed with PBS, dissociated, and pelleted

by centrifugation at 750 g for 7 minutes at 4 °C. The pellet was stored at –80 °C until used for

genomic DNA extraction. The remaining 8×10^6 mCherry+ cells were analyzed by FACS to select

for those expressing inactive ARHGAP36-mCherry mutants. Cells were sorted as described

above using the BD FACSAria II configured with a 488-nm dye laser, a 495-nm longpass filter,

and a 530/30-nm bandpass filter for EGFP detection and the laser/filter configurations described

above for mCherry detection. Approximately 4×10^5 mCherry+/EGFP– cells were obtained from

this second sort, and they were cultured for 2 days to reach full confluency and then subjected to

2 rounds of serum starvation. FACS sorting of this enriched population yielded 1.6×10^5

mCherry+/EGFP– cells, which were expanded, frozen in SHH-EGFP growth medium containing

10% DMSO, and stored in liquid nitrogen.

We then assessed if the mCherry+/EGFP– cells were still capable of Gli-dependent EGFP

expression under canonical Hh pathway activation conditions. Frozen aliquots of cells from the

third sort were thawed, expanded, and subjected to two rounds of serum starvation.

Approximately 1.0×10^7 cells were sorted with a BD FACSAria IIu configured with a 488-nm laser,

a 502-nm longpass filter, and a 525/50-nm bandpass filter for EGFP detection and a 488-nm

laser, a 595-nm longpass filter, and a 610/20-nm bandpass filter for mCherry detection. 1.0×10^6

mCherry+/EGFP– cells were collected and cultured for 7 days to achieve full confluency. The

cells were then treated with 200 nM SAG in SHH-EGFP low-serum medium for 24 hours,

expanded at a 1:2 dilution, cultured for 24 hours to enable full confluency, and treated again with

200 nM SAG for 24 hours. 2×10^7 cells were then sorted with the BD FACSAria IIu to obtain $3 \times$

10^6 mCherry+/EGFP+ cells. This selected population was expanded to 5×10^6 cells, which were

then washed with PBS, dissociated, and pelleted by centrifugation at 750 g for 7 minutes at 4 ºC.

The pellet was stored at –80 ºC until used for genomic DNA extraction.

Deep-sequencing analyses of pre- and post-selection pools

Genomic DNA was extracted from frozen pellets using the QIAamp DNA Blood Maxi Kit

(Qiagen) according to manufacturer's instructions. For each sample, *ARHGAP36* inserts were

isolated from genomic DNA by PCR using 500 ng of genomic DNA, 0.5 µM each of primer (Tables

S3, primers 59 and 60), 0.2 µM dNTP mix, and Phusion High-Fidelity DNA Polymerase in HF

buffer (New England BioLabs). A total of 182 PCRs were used to isolate *ARHGAP36* inserts from

91 µg of pre-selection genomic DNA, while 89 PCRs were used to isolate inserts from 45 µg of

post-selection genomic DNA. For each condition, the respective reactions were pooled, purified

using the QIAquick PCR Purification Kit (Qiagen), and resolved on a 0.8% EtBr–agarose gel. The

1.6-kb amplicon was then gel-extracted using the QIAquick Gel Extraction Kit. Amplicons were

quantified using a Bioanalyzer 2100 with high-sensitivity DNA kits (Agilent), sheared into 150-bp

fragments with an S220 focused-ultrasonicator (Covaris, Woburn, MA), and sequenced on a

NextSeq 500 Sequencer using High-Output v2 kits (Illumina, San Diego, CA). Raw data have

been deposited into the Dryad Digital Repository with the dataset identifier

10.5061/dryad.dz08kprv9.

FASTQ files were aligned to the wild-type *ARHGAP36* coding sequence (Bowtie2 v2.3

(*Langmead, et al., 2012*) and sorted by read name (SAMtools v1.3.1) (*Li, Heng, et al., 2009*). The

resulting mapped and sorted reads were then analyzed with an in-house Python script. Briefly,

high-quality reads were aligned to the wild-type *ARHGAP36* coding sequence. Reads with greater

than 3 high-quality mutations or with internal stop codons were discarded from further analysis.

The remaining reads were translated, identifying the ARHGAP36 amino acid mutations present

in the population. For each residue, the fold-change in its mutation frequency between the pre-

and post-selection populations was calculated.

Flow cytometry-based assays

SHH-EGFP cells were seeded into individual wells of a 24-well plate at a density of

7.5×10^4 cells/well and cultured for 24 hours in SHH-EGFP growth medium. The cells were then

transduced with retrovirus for the appropriate ARHGAP36-mCherry construct and 4 µg/mL

polybrene to achieve an MOI < 0.5. An uninfected condition was also prepared as a negative

control. After 24 hours, the cells were passaged onto a new 24-well well at a 1:1.5 dilution and

cultured in growth medium for an additional 2 days to achieve 100% confluency. Confluent cells

were then treated with SHH-EGFP low-serum medium for 24 hours. Cells were then passaged at

a 1:1.5 dilution onto a new 24-well well and cultured for 24 hours to achieve full confluency for a

second round of 24-hour serum-starvation with or without SHH-N-conditioned medium.

For flow cytometry analyses, the cells were washed with PBS, dissociated with TrypLE for

3-5 minutes at 37 °C, and centrifuged 750 *g* for 7 minutes at 4 ºC. Cell pellets were resuspended

in FACS buffer (PBS containing 1% calf serum) and analyzed on a DxP FACScan (561-nm laser

and 616/25-nm bandpass filter for mCherry detection; 488-nm laser, 560-nm shortpass filter, and

525/50-nm bandpass filter for EGFP detection) or a BD LSRII (561-nm laser, 600-nm longpass

band filter, and a 610/20-nm bandpass filter for mCherry detection; 488-nm laser, 505-nm

longpass band filter, and 525/50-nm bandpass filter for EGFP detection). Data was collected with

Cypod (Cytek, Fremont, CA) and FACSDiva software and analyzed using FlowJo. Fluorescence

data was collected for at least 2.5×10^4 cells, and three biological replicates were analyzed for

each condition.

Data analyses for all conditions except for the uninfected controls excluded mCherry–

cells, which is indicative of a lack of ARHGAP36 expression, and the median EGFP fluorescence

was calculated to measure Gli activity in each condition. For each replicate, the median EGFP

fluorescence was normalized to that of wild-type ARHGAP36-expressing cells, and a Student's

one-tailed t-test was used to identify mutations that significantly altered median EGFP

fluorescence levels ($P \leq 0.05$).

Tandem affinity purification and quantitative proteomics

NIH-3T3 cells were seeded onto 15-cm plates (6 per condition) at a density of 2×10^6

cells/plate and cultured in NIH-3T3 growth medium. After 24 hours, cells were transduced with

retrovirus for either wild-type or L317P ARHGAP36 with a C-terminal LAP tag and 4 µg/mL

polybrene to achieve an MOI > 1.5. The media was exchanged for growth medium after 4 hours,

and cells were cultured for an additional 20 hours. Cells were then washed with cold PBS,

manually scraped off each dish, and transferred into Falcon tubes. Cell suspensions expressing

the same ARHGAP36 construct were combined, and 0.5% of the resulting pool was reserved for

downstream flow cytometry analyses to confirm the MOI. The remaining cells were centrifuged at

750 *g* for 7 minutes at 4 ºC. The supernatant was aspirated, and the remaining cell pellet was

flash frozen in liquid nitrogen and stored at –80 ºC prior to LAP-tagged mediated tandem affinity

purification. Three biological replicates were conducted for each wild-type and L317P

ARHGAP36-LAP comparison.

Tandem affinity purifications and mass spectrometry analyses were conducted as

described previously (*Ding, et al., 2016; Hsu, et al., 2019; Kanie, et al., 2017; Li, Bin, et al., 2017;*

Torres, et al., 2009; Wright, et al., 2011). Pellets of ARHGAP36-LAP-expressing cells were re-

suspended in LAP resuspension buffer (300 mM KCl, 50 mM HEPES-KOH (pH 7.4), 1 mM EGTA,

1 mM MgCl$_2$, 10% glycerol, 0.5 mM DTT, and protease inhibitors (Thermo Scientific)). Cells were

lysed with the gradual addition of 10% NP-40 to a final concentration of 0.3%, followed by a 10-

minute incubation at 4 ℃. The lysate was then centrifuged at 27,000 *g* at 4 °C for 10 minutes,

and the resulting supernatant was centrifuged at 100,000 *g* for 1 hour at 4 °C. The high-speed

supernatant was next incubated with anti-GFP-antibody-coupled beads (*Ding, et al., 2016; Hsu,*

et al., 2019; Kanie, et al., 2017; Li, Bin, et al., 2017; Torres, et al., 2009; Wright, et al., 2011) for

1 hour at 4 °C to capture GFP-tagged proteins. The beads were washed five times with LAP200N

buffer (200 mM KCl, 50 mM HEPES-KOH (pH 7.4), 1 mM EGTA, 1 mM MgCl$_2$, 10% glycerol,

protease inhibitors, and 0.05% NP40) and incubated with PreScission protease in LAP200N

buffer at 4 °C for 16 hours. All subsequent steps were performed in a laminar flow hood.

PreScission protease-eluted supernatant was added to S-protein agarose beads (EMD Millipore,

Burlington, MA) and incubated rocking for 3 hours at 4 ℃. S-protein agarose beads were then

washed three times with LAP200N buffer and twice with LAP100 buffer (100 mM KCl, 50 mM

HEPES-KOH (pH 7.4), 1 mM EGTA and 10% glycerol). Beads were stored in 50mM HEPES (pH

7.5), 1 mM EGTA, 1 mM MgCl$_2$, 10% glycerol at 4 ℃ prior to on-bead digestion.

Proteins were eluted from S-protein agarose beads with an on-bead reduction, alkylation,

and tryptic digestion as follows. Samples were reduced with 10 mM DTT in ammonium

bicarbonate for an initial 5-minute incubation at 55 ℃ followed by 25 minutes at room temperature.

The proteins were then alkylated with a 30-minute incubation in 30 mM acrylamide at room

temperature, and finally eluted from the beads with an overnight digest performed at room

temperature using Trypsin/LysC (Promega) and 0.02% ProteaseMax (Promega). The digests

were acidified with 1% formic acid, de-salted with C18 Monospin reversed phase columns (GL

Sciences, Tokyo, Japan), dried on a SpeedVac, and reconstituted in 12.5 µL of 2% acetonitrile

and 0.1% formic acid. 4 µL of each sample were used for liquid-chromatography-mass

spectrometry analyses performed on an Acquity M-Class UPLC (Waters Corporation, Milford, MA)

and either an Orbitrap Q-Exactive HFX mass spectrometer (Thermo Scientific, San Jose, CA) or

an Orbitrap Fusion Tribrid mass spectrometer (Thermo Scientific). For each biological replicate,

the sample from the ARHGAP36 L317P-expressing cells was run immediately before that of the

wild-type ARHGAP36-expressing cells. Analysis of the resulting .RAW data files was conducted

using Byonic (Protein Metrics, San Carlos, CA), with the assumption of tryptic proteolysis and a

maximum allowance of two missed cleavage sites. Precursor and MS/MS fragment mass

accuracies were held within 12 ppm and 0.4 Da, respectively. A false discovery rate of 1% was

used for protein identification (*Elias, et al., 2007*). Raw data have been deposited to the

ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier

PXD019056 and 10.6019/PXD019056.

The resulting list of identified proteins was compared to an NCBI FASTA database

containing all mouse proteomic isoforms with the exception of the tandem affinity bait construct

sequence and common contaminant proteins. Post-processing of spectral counts was conducted

with an in-house R script. For each protein, spectral counts detected across all isoforms were

combined and normalized to the mean amino acid length of all isoforms. The resulting normalized

spectral count was divided by the sum of normalized spectral counts calculated for all proteins in

the pulldown sample, generating a normalized spectral abundance factor (NSAF) for each protein.

To account for variability in bait ARHGAP36 expression across different pulldown samples, the

NSAF of each protein was divided by that of the bait ARHGAP36 (relative NSAF). Proteins that

were detected in all biological replicates for a given condition were tabulated, resulting in a dataset

of 566 candidate ARHGAP36-binding proteins. For each protein, the fold-change in relative NSAF

between the wild-type and L317P mutant ARHGAP36 pulldown samples of the same replicate

was calculated. The average fold-change in relative NSAF across of all three replicates was then

calculated for each protein.

To assess the robustness of our comparative interactome analyses, we calculated a

modified Z score that compares the protein enrichment in either interactome (wild-type vs. L317P

ARHGAP36) against the experimental variability in protein abundance measurements. Protein

enrichment was represented by the $\log_2$-transformation of the average fold change in relative

NSAF between wild-type and L317P ARHGAP36 interactomes [$\log_2$ (WT NSAF:L317 NSAF)]. We

estimated the error in measuring protein abundance in a given interactome by normalizing the

relative NSAFs for each replicate to the average value across all three replicates (mean-

normalized relative NSAFs). This transformation approximates how much the variation between

replicates can contribute to an apparent fold change. To place equal weight on upward and

downward variations from the mean, we calculated the absolute value of the $\log_2$-transformed

mean-normalized relative NSAFs. These calculations were conducted for the relative NSAFs of

a protein in both the wild-type and L317P mutant ARHGAP36 interactomes, and the two resulting

values were summed to produce a final error estimate. The modified Z score of each protein was

calculated by dividing the $\log_2$ (WT NSAF:L317 NSAF) by the final error estimate and then

calculating the absolute value of the resulting quotient. Proteins with higher modified Z scores are

those that are enriched in a given interactome to a degree that is greater than the estimated

experimental error.

SUPPLEMENTAL DATA

837 Figure 4 – figure supplement 1

838 Figure 6 – figure supplement 1

839 Supplementary File 1. Mutation frequencies in pre- and post-selection populations.

840 Supplementary File 2. Wild-type and L317P ARHGAP36 isoform 2 interactomes

841 Supplementary File 3. Antibody and primer resources

ACKNOWLEDGEMENTS

This work was supported by the National Institutes of Health (R35 GM127030 to J.K.C.;

S10 RR025518-01 and S10 RR027431-01 to the Stanford Shared FACS Facility; and P30

CA124435 to Stanford University Mass Spectrometry), the Rachel Molly Markoff Foundation

(J.K.C.), and an Alex's Lemonade Stand Foundation Young Investigator Award (J.N.). Cell sorting

and flow cytometry analyses were conducted at the Stanford Shared FACS Facility. Deep

sequencing analyses were performed at the Stanford Functional Genomics Facility. Quantitative

proteomics were executed at the Vincent Coates Foundation Mass Spectrometry Laboratory,

Stanford University Mass Spectrometry.

AUTHOR CONTRIBUTIONS

P.R. N. designed and conducted experiments, analyzed the data, and wrote the paper.

T.K. developed software to analyze deep sequencing data. N.A.M. performed tandem affinity

purifications. J.N. generated expression vectors for ARHGAP36 isoforms and isoform 2 truncation

mutants. J.D. contributed to the development of software to analyze proteomic data. P.K.J.

provided resources and supervision for proteomics experiments. J.K.C. designed the

experiments, analyzed the data, and wrote the paper.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

REFERENCES

Ågren, M., Kogerman, P., Kleman, M.I., Wessling, M. & Toftgård, R. 2004. Expression of the PTCH1 tumor suppressor gene is regulated by alternative promoters and a single functional Gli-binding site. *Gene* **330:** 101-114. doi: https://doi.org/10.1016/j.gene.2004.01.010.

Amin, E. *et al.* 2016. Deciphering the molecular and functional basis of RHOGAP family proteins: A systematic approach toward selective inactivation of RHO family proteins. *Journal of Biological Chemistry* **291:** 20353-20371. doi: https://doi.org/10.1074/jbc.M116.736967.

Bai, C.B., Stephen, D. & Joyner, A.L. 2004. All mouse ventral spinal cord patterning by Hedgehog is Gli dependent and involves an activator function of Gli3. *Developmental Cell* **6:** 103-115. doi: https://doi.org/10.1016/S1534-5807(03)00394-0.

Barzi, M., Berenguer, J., Menendez, A., Alvarez-Rodriguez, R. & Pons, S. 2009. Sonic-hedgehog-mediated proliferation requires the localization of PKA to the cilium base. *Journal of Cell Science* **123:** 62-69. doi: https://doi.org/10.1242/jcs.060020, PMID: 5661997.

Beauchamp, E. *et al.* 2009. GLI1 is a direct transcriptional target of EWS-FLI1 oncoprotein. *The Journal of Biological Chemistry* **284:** 9074-9082. doi: https://doi.org/10.1074/jbc.M806233200, PMID: 19189974.

Beckmann, P.J. *et al.* 2019. Sleeping beauty insertional mutagenesis reveals important genetic drivers of central nervous system embryonal tumors. *Cancer Research* **79:** 905-917. doi: https://doi.org/10.1158/0008-5472.CAN-18-1261, PMID: 30674530.

Breslow, D.K. *et al.* 2018. A CRISPR-based screen for Hedgehog signaling provides insights into ciliary function and ciliopathies. *Nature Genetics* **50:** 460-471. doi: https://doi.org/10.1038/s41588-018-0054-7.

Briscoe, J., Pierani, A., Jessell, T.M. & Ericson, J. 2000. A homeodomain protein code specifies

progenitor cell identity and neuronal fate in the ventral neural tube. *Cell* **101:** 435-445. doi:

https://doi.org/10.1016/S0092-8674(00)80853-3, PMID: 10830170.

Buonamici, S. *et al.* 2010. Interfering with resistance to smoothened antagonists by inhibition of

the PI3K pathway in medulloblastoma. *Sci. Transl. Med.* 10.1126/scitranslmed.3001599

51-70. doi: https://doi.org/10.1126/scitranslmed.3001599, PMID: 20881279.

Chávez, M. *et al.* 2015. Modulation of ciliary phosphoinositide content regulates trafficking and

Sonic Hedgehog signaling output. *Developmental Cell* **34:** 338-350. doi:

https://doi.org/10.1016/j.devcel.2015.06.016.

Chen, J.K., Taipale, J., Young, K.E., Maiti, T. & Beachy, P.A. 2002. Small molecule modulation

of Smoothened activity. *Proceedings of the National Academy of Sciences of the United*

States of America **99:** 14071-14076. doi: https://doi.org/10.1073/pnas.182542899.

Dai, P. *et al.* 1999. Sonic hedgehog-induced activation of the Gli1 promoter is mediated by GLI3.

Journal of Biological Chemistry **274:** 8143-8152. doi:

https://doi.org/10.1074/jbc.274.12.8143.

Dennler, S. *et al.* 2007. Induction of sonic hedgehog mediators by transforming growth factor-

beta: Smad3-dependent activation of Gli2 and Gli1 expression in vitro and in vivo. *Cancer*

Research **67:** 6981-6986. doi: https://doi.org/10.1158/0008-5472.CAN-07-0491, PMID:

17638910.

Ding, S. *et al.* 2016. Comparative proteomics reveals strain-specific β-TrCP degradation via

rotavirus NSP1 hijacking a host Cullin-3-Rbx1 complex. *PLoS Pathogens* **12:** e1005929.

doi: https://doi.org/10.1371/journal.ppat.1005929.

Dorn, K.V., Hughes, C.E. & Rohatgi, R. 2012. A Smoothened-Evc2 complex transduces the

Hedgehog signal at primary cilia. *Developmental Cell* **23:** 823-835. doi:

https://doi.org/10.1016/j.devcel.2012.07.004.

Eccles, R.L. *et al.* 2016. Bimodal antagonism of PKA signalling by ARHGAP36. *Nature*

Communications **7:** 12963. doi: https://doi.org/10.1038/ncomms12963, PMID: 27713425.

Elias, J.E. & Gygi, S.P. 2007. Target-decoy search strategy for increased confidence in large-

scale protein identifications by mass spectrometry. *Nature Methods* **4:** 207-214. doi:

https://doi.org/10.1038/nmeth1019.

Elsawa, S.F. *et al.* 2011. GLI2 transcription factor mediates cytokine cross-talk in the tumor

microenvironment. *Journal of Biological Chemistry* **286:** 21524-21534. doi:

https://doi.org/10.1074/jbc.M111.234146.

Faucherre, A. *et al.* 2003. Lowe syndrome protein OCRL1 interacts with Rac GTPase in the trans-

Golgi network. *Human Molecular Genetics* **12:** 2449-2456. doi:

https://doi.org/10.1093/hmg/ddg250.

Flora, A., Klisch, T.J., Schuster, G. & Zoghbi, H.Y. 2009. Deletion of Atoh1 disrupts Sonic

Hedgehog signaling in the developing cerebellum and prevents medulloblastoma. *Science*

326: 1424-1427. doi: https://doi.org/10.1126/science.1181453, PMID: 19965762.

Garcia-Gonzalo, F.R. *et al.* 2015. Phosphoinositides regulate ciliary protein trafficking to

modulate Hedgehog signaling. *Developmental Cell* **34:** 400-409. doi:

https://doi.org/10.1016/j.devcel.2015.08.001.

Han, B. *et al.* 2015. FOXC1 activates Smoothened-independent Hedgehog signaling in basal-

like breast cancer. *Cell Reports* **13:** 1046-1058. doi:

https://doi.org/10.1016/j.celrep.2015.09.063.

Haycraft, C.J. *et al.* 2005. Gli2 and Gli3 localize to cilia and require the intraflagellar transport

protein Polaris for processing and function. *PLoS Genetics* **1:** e53. doi:

https://doi.org/10.1371/journal.pgen.0010053.

Heldwein, E.E. *et al.* 2004. Crystal structure of the clathrin adaptor protein 1 core. *Proceedings*

of the National Academy of Sciences of the United States of America **101:** 14108-14113.

doi: https://doi.org/10.1073/pnas.0406102101.

Hill, P., Götz, K. & Rüther, U. 2009. A SHH-independent regulation of Gli3 is a significant

determinant of anteroposterior patterning of the limb bud. *Developmental Biology* **328:**

506-516. doi: https://doi.org/10.1016/j.ydbio.2009.02.017.

Hsu, J. *et al.* 2019. E2F4 regulates transcriptional activation in mouse embryonic stem cells

independently of the RB family. *Nature Communications* **10:** 2939. doi:

https://doi.org/10.1038/s41467-019-10901-x.

Huangfu, D. & Anderson, K.V. 2005. Cilia and Hedgehog responsiveness in the mouse.

Proceedings of the National Academy of Sciences of the United States of America **102:**

11325-11330. doi: https://doi.org/10.1073/pnas.0505328102.

Hui, C.-c. & Angers, S. 2011. Gli proteins in development and disease. *Annual Review of Cell*

and Developmental Biology **27:** 513-537. doi: https://doi.org/10.1146/annurev-cellbio-

092910-154048, PMID: 21801010.

Humke, E.W., Dorn, K.V., Milenkovic, L., Scott, M.P. & Rohatgi, R. 2010. The output of Hedgehog

signaling is controlled by the dynamic association between Suppressor of Fused and the

Gli proteins. *Genes & Development* **24:** 670-682. doi:

https://doi.org/10.1101/gad.1902910, PMID: 20360384.

Huttlin, E.L. *et al.* 2017. Architecture of the human interactome defines protein communities and

disease networks. *Nature* **545:** 505-509. doi: https://doi.org/10.1038/nature22366, PMID:

28514442.

Hyman, J.M. *et al.* 2009. Small-molecule inhibitors reveal multiple strategies for Hedgehog

pathway blockade. *Proceedings of the National Academy of Sciences of the United States*

of America **106:** 14132-14137. doi: https://doi.org/10.1073/pnas.0907134106, PMID: 19666565.

Incardona, J.P. *et al.* 2000. Receptor-mediated endocytosis of soluble and membrane-tethered sonic hedgehog by patched-1. *Proceedings of the National Academy of Sciences of the United States of America* **97:** 12044-12049. doi: https://doi.org/10.1073/pnas.220251997.

Jaeken, J. *et al.* 2006. Deletion of PREPL, a gene encoding a putative serine oligopeptidase, in patients with hypotonia-cystinuria syndrome. *American Journal of Human Genetics* **78:** 38-51. doi: https://doi.org/10.1086/498852, PMID: 16385448.

Kanie, T. *et al.* 2017. The CEP19-RABL2 GTPase complex binds IFT-B to Iinitiate intraflagellar transport at the ciliary base. *Developmental Cell* **42:** 22-36.e12. doi: https://doi.org/10.1016/j.devcel.2017.05.016.

Kasper, M. *et al.* 2006. Selective modulation of Hedgehog/GLI target gene expression by epidermal growth factor signaling in human keratinocytes. *Molecular and Cellular Biology* **26:** 6283-6298. doi: https://doi.org/10.1128/mcb.02317-05.

Kim, J., Kato, M. & Beachy, P.A. 2009. Gli2 trafficking links Hedgehog-dependent activation of Smoothened in the primary cilium to transcriptional activation in the nucleus. *Proceedings of the National Academy of Sciences of the United States of America* **106:** 21666-21671. doi: https://doi.org/10.1073/pnas.0912180106, PMID: 19996169.

Langmead, B. & Salzberg, S.L. 2012. Fast gapped-read alignment with Bowtie 2. *Nature Methods* **9:** 357-359. doi: https://doi.org/10.1038/nmeth.1923, PMID: 22388286.

Lee, Y.s. *et al.* 2019. A computational framework for genome-wide characterization of the human disease landscape. *Cell Systems* **8:** 152-162. doi: https://doi.org/10.1016/j.cels.2018.12.010.

Lewis, P.M., Gritli-Linde, A., Smeyne, R., Kottmann, A. & McMahon, A.P. 2004. Sonic hedgehog signaling is required for expansion of granule neuron precursors and patterning of the

mouse cerebellum. *Developmental Biology* **270:** 393-410. doi:

https://doi.org/10.1016/j.ydbio.2004.03.007, PMID: 15183722.

Li, B. *et al.* 2017. Drebrin restricts rotavirus entry by inhibiting dynamin-mediated endocytosis.

Proceedings of the National Academy of Sciences of the United States of America **114:**

E3642-E3651. doi: https://doi.org/10.1073/pnas.1619266114.

Li, H. *et al.* 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25:** 2078-

2079. doi: https://doi.org/10.1093/bioinformatics/btp352, PMID: 19505943.

Lignitto, L. *et al.* 2013. Proteolysis of MOB1 by the ubiquitin ligase praja2 attenuates Hippo

signalling and supports glioblastoma growth. *Nature Communications* **4:** 1822. doi:

https://doi.org/10.1038/ncomms2791.

Lignitto, L. *et al.* 2011. Control of PKA stability and signalling by the RING ligase praja2. *Nature*

Cell Biology **13:** 412-422. doi: https://doi.org/10.1038/ncb2209.

Liu, A., Wang, B. & Niswander, L.A. 2005. Mouse intraflagellar transport proteins regulate both

the activator and repressor functions of Gli transcription factors. *Development* **132:** 3103-

3111. doi: https://doi.org/10.1242/dev.01894.

Liu, Z., Li, T., Reinhold, M.I. & Naski, M.C. 2014. MEK1-RSK2 contributes to Hedgehog signaling

by stabilizing GLI2 transcription factor and inhibiting ubiquitination. *Oncogene* **33:** 65-73.

doi: https://doi.org/10.1038/onc.2012.544, PMID: 23208494.

Long, J. *et al.* 2014. The BET bromodomain inhibitor I-BET151 acts downstream of smoothened

protein to abrogate the growth of hedgehog protein-driven cancers. *Journal of Biological*

Chemistry **289:** 35494-35502. doi: https://doi.org/10.1074/jbc.M114.595348.

Marchesi, S. *et al.* 2014. DEPDC1B coordinates de-adhesion events and cell-cycle progression

at mitosis. *Developmental Cell* **31:** 420-433. doi:

https://doi.org/10.1016/j.devcel.2014.09.009.

May, S.R. *et al.* 2005. Loss of the retrograde motor for IFT disrupts localization of Smo to cilia and prevents the expression of both activator and repressor functions of Gli. *Developmental Biology* **287:** 378-389. doi: https://doi.org/10.1016/j.ydbio.2005.08.050, PMID: 16229832.

Moon, S. 2003. Rho GTPase-activating proteins in cell regulation. *Trends in Cell Biology* **13:** 13-22. doi: https://doi.org/10.1016/S0962-8924(02)00004-1.

Moore, B.S. *et al.* 2016. Cilia have high cAMP levels that are inhibited by Sonic Hedgehog-regulated calcium dynamics. *Proceedings of the National Academy of Sciences of the United States of America* **113:** 13069-13074. doi: https://doi.org/10.1073/pnas.1602393113, PMID: 27799542.

Mukhopadhyay, S. *et al.* 2013. The ciliary G-protein-coupled receptor Gpr161 negatively regulates the Sonic hedgehog pathway via cAMP signaling. *Cell* **152:** 210-223. doi: https://doi.org/10.1016/j.cell.2012.12.026, PMID: 23332756.

Müller, P.M. *et al.* 2020. Systems analysis of RhoGEF and RhoGAP regulatory proteins reveals spatially organized RAC1 signalling from integrin adhesions. *Nature Cell Biology* **22:** 498–511. doi: https://doi.org/10.1038/s41556-020-0488-x, PMID: 32203420.

Murone, M., Rosenthal, A. & De Sauvage, F.J. 1999. Sonic hedgehog signaling by the patched-smoothened receptor complex. *Current Biology* **9:** 76-84. doi: https://doi.org/10.1016/S0960-9822(99)80018-9, PMID: 10021362.

Nam, H. *et al.* 2019. Critical roles of ARHGAP36 as a signal transduction mediator of shh pathway in lateral motor columnar specification. *eLife* **8:** e46683. doi: https://doi.org/10.7554/eLife.46683.

Pan, Y., Bai, C.B., Joyner, A.L. & Wang, B. 2006. Sonic hedgehog signaling regulates Gli2 transcriptional activity by suppressing its processing and degradation. *Molecular and*

Cellular Biology **26:** 3365-3377. doi: https://doi.org/10.1128/MCB.26.9.3365-3377.2006,

PMID: 16611981.

Pan, Y. & Wang, B. 2007. A novel protein-processing domain in Gli2 and Gli3 differentially blocks

complete protein degradation by the proteasome. *The Journal of Biological Chemistry*

282: 10846-10852. doi: https://doi.org/10.1074/jbc.M608599200, PMID: 17283082.

Pedersen, L.B., Mogensen, J.B. & Christensen, S.T. 2016. Endocytic control of cellular signaling

at the primary cilium. *Trends in Biochemical Sciences* **41:** 787-797. doi:

https://doi.org/10.1016/j.tibs.2016.06.002.

Pusapati, G.V. *et al.* 2018. CRISPR screens uncover genes that regulate target cell sensitivity to

the morphogen Sonic Hedgehog. *Developmental Cell* **44:** 113-129.e118. doi:

https://doi.org/10.1016/j.devcel.2017.12.003.

Rack, P.G. *et al.* 2014. Arhgap36-dependent activation of Gli transcription factors. *Proceedings*

of the National Academy of Sciences of the United States of America **111:** 11061-11066.

doi: https://doi.org/10.1073/pnas.1322362111, PMID: 25024229.

Radhakrishnan, K., Baltes, J., Creemers, J.W.M. & Schu, P. 2013. Trans-Golgi network

morphology and sorting is regulated by prolyl-oligopeptidase-like protein PREPL and the

AP-1 complex subunit μ1A. *Journal of Cell Science* **126:** 1155-1163. doi:

https://doi.org/10.1242/jcs.116079.

Riobó, N.A., Lu, K., Ai, X., Haines, G.M. & Emerson, C.P. 2006. Phosphoinositide 3-kinase and

Akt are essential for Sonic Hedgehog signaling. *Proceedings of the National Academy of*

Sciences of the United States of America **103:** 4505-4510. doi:

https://doi.org/10.1073/pnas.0504337103, PMID: 16537363.

Rohatgi, R., Milenkovic, L. & Scott, M.P. 2007. Patched1 regulates hedgehog signaling at the

primary cilium. *Science* **317:** 372-376. doi: https://doi.org/10.1126/science.1139740,

PMID: 17641202.

Scheffzek, K., Ahmadian, M.R. & Wittinghofer, A. 1998. GTPase-activating proteins: helping

hands to complement an active site. *Trends in Biochemical Sciences* **23:** 257-262. doi:

https://doi.org/10.1016/S0968-0004(98)01224-9.

Schindelin, J. *et al.* 2012. Fiji: An open-source platform for biological-image analysis. *Nature*

Methods **9:** 676-682. doi: https://doi.org/10.1038/nmeth.2019, PMID: 22743772.

Schulz, I. *et al.* 2002. Modulation of inositol 1,4,5-triphosphate concentration by prolyl

endopeptidase inhibition. *European Journal of Biochemistry* **269:** 5813-5820. doi:

https://doi.org/10.1046/j.1432-1033.2002.03297.x, PMID: 12444969.

Sepe, M. *et al.* 2014. Proteolytic control of neurite outgrowth inhibitor NOGO-A by the cAMP/PKA

pathway. *Proceedings of the National Academy of Sciences of the United States of*

America **111:** 15729-15734. doi: https://doi.org/10.1073/pnas.1410274111.

Stamataki, D., Ulloa, F., Tsoni, S.V., Mynett, A. & Briscoe, J. 2005. A gradient of Gli activity

mediates graded Sonic Hedgehog signaling in the neural tube. *Genes & Development* **19:**

626-641. doi: https://doi.org/10.1101/gad.325905.

Stone, D.M. *et al.* 1996. The tumour-suppressor gene patched encodes a candidate receptor for

Sonic hedgehog. *Nature* **384:** 129-134. doi: https://doi.org/10.1038/384129a0.

Stone, D.M. *et al.* 1999. Characterization of the human Suppressor of fused a negative regulator

of the zinc-finger transcription factor Gli. *Journal of Cell Science* **112:** 4437-4448. PMID:

10564661.

Szeltner, Z., Alshafee, I., Juhász, T., Parvari, R. & Polgár, L. 2005. The PREPL A protein, a new

member of the prolyl oligopeptidase family, lacking catalytic activity. *Cellular and*

Molecular Life Sciences **62:** 2376-2381. doi: https://doi.org/10.1007/s00018-005-5262-5.

Taipale, J. *et al.* 2000. Effects of oncogenic mutations in Smoothened and Patched can be

reversed by cyclopamine. *Nature* **406:** 1005-1009. doi: https://doi.org/10.1038/35023008,

PMID: 10984056.

Taipale, J., Cooper, M.K., Maiti, T. & Beachy, P.A. 2002. Patched acts catalytically to suppress

the activity of smoothened. *Nature* **418:** 892-897. doi:

https://doi.org/10.1038/nature00989, PMID: 12192414.

te Welscher, P. *et al.* 2002. Progression of vertebrate limb development through SHH-mediated

counteraction of GLI3. *Science* **298:** 827-830. doi:

https://doi.org/10.1126/science.1075620.

Tempe, D., Casas, M., Karaz, S., Blanchet-Tournier, M.-F. & Concordet, J.-P. 2006. Multisite

protein kinase A and glycogen synthase kinase 3 phosphorylation leads to Gli3

ubiquitination by SCF β-TrCP. *Molecular and Cellular Biology* **26:** 4316-4326. doi:

https://doi.org/10.1128/mcb.02183-05.

Torres, J.Z., Miller, J.J. & Jackson, P.K. 2009. High-throughput generation of tagged stable cell

lines for proteomic analysis. *Proteomics* **9:** 2888-2891. doi:

https://doi.org/10.1002/pmic.200800873.

Tukachinsky, H., Lopez, L.V. & Salic, A. 2010. A mechanism for vertebrate Hedgehog signaling:

recruitment to cilia and dissociation of SuFu-Gli protein complexes. *The Journal of Cell*

Biology **191:** 415-428. doi: https://doi.org/10.1083/jcb.201004108, PMID: 20956384.

Tuson, M. *et al.* 2011. Protein kinase A acts at the basal body of the primary cilium to prevent

Gli2 activation and ventralization of the mouse neural tube. *Development* **138:** 4921-4930.

doi: https://doi.org/10.1242/dev.070805, PMID: 22007132.

Vuolo, L., Herrera, A., Torroba, B., Menendez, A. & Pons, S. 2015. Ciliary adenylyl cyclases

control the Hedgehog pathway. *Journal of Cell Science* **128:** 2928-2937. doi:

https://doi.org/10.1242/jcs.172635.

Wallace, V.A. 1999. Purkinje-cell-derived Sonic hedgehog regulates granule neuron precursor

cell proliferation in the developing mouse cerebellum. *Current Biology* **9:** 445-448. doi:

https://doi.org/10.1016/S0960-9822(99)80195-X.

Wang, B. & Li, Y. 2006. Evidence for the direct involvement of β-TrCP in Gli3 protein processing. *Proceedings of the National Academy of Sciences of the United States of America* **103:** 33-38. doi: https://doi.org/10.1073/pnas.0509927103.

Wang, C., Pan, Y. & Wang, B. 2010. Suppressor of fused and Spop regulate the stability processing and function of Gli2 and Gli3 full-length activators but not their repressors. *Development* **137:** 2001-2009. doi: https://doi.org/10.1242/dev.052126, PMID: 20463034.

Wang, Y., Zhou, Z., Walsh, C.T. & McMahon, A.P. 2009. Selective translocation of intracellular Smoothened to the primary cilium in response to Hedgehog pathway modulation. *Proceedings of the National Academy of Sciences of the United States of America* **106:** 2623-2628. doi: https://doi.org/10.1073/pnas.0812110106.

Wang, Y.J. *et al.* 2003. Phosphatidylinositol 4 phosphate regulates targeting of clathrin adaptor AP-1 complexes to the Golgi. *Cell* **114:** 299-310. doi: https://doi.org/10.1016/S0092-8674(03)00603-2, PMID: 12914695.

Wechsler-Reya, R. & Scott, M.P. 2001. The developmental biology of brain tumors. *Annual Review of Neuroscience* **24:** 385-428. doi: https://doi.org/10.1146/annurev.neuro.24.1.385, PMID: 11283316.

Wen, X. *et al.* 2010. Kinetics of Hedgehog-dependent full-length Gli3 accumulation in primary cilia and subsequent degradation. *Molecular and Cellular Biology* **30:** 1910-1922. doi: https://doi.org/10.1128/mcb.01089-09.

Williams, R.S.B., Eames, M., Ryves, W.J., Viggars, J. & Harwood, A.J. 1999. Loss of a prolyl oligopeptidase confers resistance to lithium by elevation of inositol (1,4,5) trisphosphate. *EMBO Journal* **18:** 2734-2745. doi: https://doi.org/10.1093/emboj/18.10.2734, PMID: 10329620.

Wright, K.J. *et al.* 2011. An ARL3-UNC119-RP2 GTPase cycle targets myristoylated NPHP3 to

the primary cilium. *Genes and Development* **25:** 2347-2360. doi:

https://doi.org/10.1101/gad.173443.111, PMID: 22085962.

Zhang, B. *et al.* 2019. Patched1-ArhGAP36-PKA-Inversin axis determines the ciliary

translocation of Smoothened for Sonic Hedgehog pathway activation. *Proceedings of the*

National Academy of Sciences of the United States of America **116:** 874-879. doi:

https://doi.org/10.1073/pnas.1804042116, PMID: 30598432.

1138

mCherry DAPI

Figure 4 – figure supplement 1. Subcellular localization of ARHGAP36 isoform 2 point mutants. Representative immunofluorescence micrographs of NIH-3T3 cells retrovirally transduced with the indicated mCherry-tagged ARHGAP36 constructs. Scale bar: 20 μm.

Figure 6 – figure supplement 1. Comparative analyses of the wild-type and L317P ARHGAP36 interactomes. Scatter plot of ARHGAP36 isoform 2-binding proteins according to their normalized spectral abundance in the wild-type and L317P interactomes. Each data point represents the average value for three biological replicates, and a modified Z-score was calculated for each fold change in abundance (see Methods)