Identification of *Avramr1* from *Phytophthora infestans* using long read and cDNA pathogen-enrichment sequencing (PenSeq)

Xiao Lin[1], Tianqiao Song[1], Sebastian Fairhead[1], Kamil Witek[1], Agathe Jouet[1], Florian Jupe[1], Agnieszka I. Witek[1], Vivianne G. A. A. Vleeshouwers[2], Ingo Hein[3] and Jonathan D. G. Jones[1*]

[1]The Sainsbury Laboratory, Norwich Research Park, Norwich, NR4 7UH, UK; [2]Plant Breeding, Wageningen University and Research, Droevendaalsesteeg 1, 6708 PB. Wageningen, The Netherlands; [3]School of Life Sciences, Division of Plant Sciences, University of Dundee and The James Hutton Institute, Dundee, DD2 5DA, UK

*Corresponding author:
Jonathan D. G. Jones
Tel: +44 (0)1603 450400
Email: jonathan.jones@tsl.ac.uk

Word count: Total: 3,788 words; Summary: 185; Introduction: 537; Materials and Methods: 957; Results: 1,590; Discussion: 486; Acknowledgements: 33.

Number of Figures: 5 (colour)

Number of Table: 1

Number of supplementary Figures: 2

Number of supplementary Tables: 3

Number of supplementary Notes: 4

Running title: *Avramr1* cloning by long read and cDNA PenSeq

Keywords: *Avramr1*, late blight, oomycete, PenSeq, *Phytophthora infestans*, *Rpi-amr1*, RxLR effector

Summary

- Potato late blight, caused by the oomycete pathogen *Phytophthora infestans,* significantly hampers potato production. Recently, a new *Resistance to Phytophthora infestans* (*Rpi*) gene, *Rpi-amr1*, was cloned from a wild *Solanum* species, *Solanum americanum*. Identification of the corresponding recognized effector (*Avirulence,* or *Avr)* genes from *P. infestans* is key to elucidating their naturally occurring sequence variation, which in turn informs the potential durability of the cognate late blight resistance.

- To identify the *P. infestans* effector recognized by *Rpi-amr1*, we screened available effector libraries and used long read and cDNA pathogen-enrichment sequencing (PenSeq) on four *P. infestans* isolates to explore the untested effectors.

- By using SMRT and cDNA PenSeq, we identified 47 highly expressed effectors from *P. infestans*, including PITG_07569 which triggers a highly specific cell death response when transiently co-expressed with *Rpi-amr1* in *Nicotiana benthamiana*, suggesting that *PITG_07569* is *Avramr1*.

- Here we demonstrate that long read and cDNA PenSeq enables the identification of full-length RxLR effector families, and their expression profile. This study has revealed key insights into the evolution and polymorphism of a complex RxLR effector family that is associated with the recognition by *Rpi-amr1*.

Introduction

Potato late blight, caused by the hemi-biotrophic oomycete pathogen *Phytophthora infestans,* triggered the Irish and European famine in the late 1840s, and still causes severe losses to world potato production.

To reduce losses, breeders sought resistance genes in wild relatives of potato. Early in the 20[th] century, *Solanum demissum*, a highly resistant hexaploid (2n=72) wild potato was found to be a useful source of Resistance to *P. infestans (Rpi)* genes (Salaman, 1937). Since then, many resistance traits have been transferred to cultivated potatoes by introgression breeding (Toxopeus, 1956), and many *Rpi* genes have been cloned from wild potatoes, e.g. *R1*, *R3a*, *R8*, *Rpi-blb1* and *Rpi-vnt1* (Ballvora *et al.*, 2002; van der Vossen *et al.*, 2003; Huang *et al.*, 2005; Foster *et al.*, 2009; Pel *et al.*, 2009; Vossen *et al.*, 2016). Unlike wild potatoes, *Solanum nigrum* and *Solanum americanum* have been reported to be non-hosts for *P. infestans* (Colon *et al.*, 1993)*.* Two *Rpi* genes encoding NLR proteins, *Rpi-amr3* and *Rpi-amr1,* were cloned from *S. americanum* and confer late blight resistance in potato (Witek *et al.*, 2016; 2020).

Identification of the recognized effectors for *Rpi-amr3* and *Rpi-amr1* would open the way to investigate their virulence function and distribution in *P. infestans* populations. Moreover, it could also help to diagnose *Rpi* gene repertoires in resistant plants, and individually confirm their activity in genetically modified potatoes carrying multiple *Rpi* genes. In oomycetes, all the cloned Avr proteins contain a signal peptide and RxLR motif (Rehmany *et al.*, 2005), and the genomic sequencing of *P. infestans* revealed 563 RxLR effectors in the T30-4 reference genome (Haas *et al.*, 2009). This enabled a high-throughput effectoromics approach for functional screening of the candidate effectors in plants (Vleeshouwers *et al.*, 2008; 2011), and many *Avr* genes were identified by this approach, including *Avrblb1*, *Avrblb2* and *Avrvnt1* (Vleeshouwers *et al.*, 2008; Oh *et al.*, 2009; Pel, 2010).

However, available RxLR effector libraries do not contain recombinant clones of all *P. infestans* RxLR effectors, the effector candidates were defined on the basis of expression profile, motif analysis and distribution between *P. infestans* races (Vleeshouwers *et al.*, 2008; Oh *et al.*, 2009; Haas *et al.*, 2009). In total, ~300/563 RxLR effectors were previously cloned into expression vectors for functional screening (Rietman, 2011).

To further explore the diversity of RxLR effectors from *P. infestans*, a pathogen enrichment

sequencing (PenSeq) approach was adopted to study allelic variation of RxLR effectors and

population genomics of oomycetes. A bait library of RxLR effectors and some other pathogen-

related genes was synthesized and used for enrichment prior to sequencing (Jouet *et al.*, 2018;

Thilliez *et al.*, 2018). However, the previous PenSeq analyses used Illumina reads and genomic

DNA (gDNA), making it difficult to differentiate individual effector alleles and closely related

paralogs, or to find out which effectors are expressed.

Here, to identify the recognized effector of the newly-cloned Rpi-amr1 protein from *S.*

americanum (Witek *et al.*, 2020)*,* we screened all currently available RxLR effectors for

recognition but without success. Therefore, we used PenSeq with long read (PacBio) and

cDNA methods, and extended the list of candidate effectors that could be screened. Amongst

these additional candidate RXLR genes, we identified *Avramr1* and defined orthologs and

paralogs from four different isolates of *P. infestans*.

Materials and Methods:

Sample preparation

To collect the mycelium of *Phytophthora infestans* for DNA extraction, *P. infestans* strains

were grown on RSA solid medium for 7 days and then moved to Plich liquid media for 14 days.

Mycelia were washed and harvested, freeze-dried using a vacuum pump and stored at -80°C

until DNA or RNA extraction.

To collect the infection samples or zoospores for RNA extraction, *P. infestans* strains were

cultured for 10 days on RSA medium. Grown mycelia were covered with cold (4 °C) sterile

water and then incubated at 4-6 °C for 2-3h. The concentration of the inoculum was adjusted

to about 50,000 zoospores/mL and 10 μL drops of inoculum were placed on the detached leaves

of potato plants. Detached leaf assays (DLA) were incubated at 20 °C in high humidity for a

required time post inoculation. Leaf discs of the infection area were collected and stored at -

80°C until DNA or RNA extraction.

DNA and RNA extraction

DNA was extracted using phenol/chloroform. *P. infestans* mycelium samples or infected leaf

discs were ground into powder in liquid nitrogen. Ground material was resuspended in 500 μl

of Shorty buffer (20% 1M Tris HCl pH 9, 20% 2M LiCl, 5% 0.5M EDTA, 10% SDS 10%, 45%

dH2O) and one volume of phenol: chloroform: isoamyl alcohol (25: 24: 1) was added. The

upper aqueous phase containing DNA was mixed with one volume of 100% ice-cold

isopropanol to precipitate DNA. The pellet was washed twice using 70% ethanol, heated at

70°C for 2-5 minutes to completely remove ethanol and resuspended in sterile water.

Resuspended DNA was then heated at 65°C for 20 minutes to inactivate DNases before RNase

treatment was performed (2 μl of 10 mg/ml^{-1} RNase A, 37°C, 1h) and RNase A removed by

chlorophorm precipitation. Genomic DNA was resuspended in water and sheared into 3-5 kb

fragments using the S220 Focused-ultrasonicator (Covaris Inc., MA, USA).

RNA samples were extracted with Direct-zol^{TM} RNA MiniPrep kit (Zymo Research, Tustin,

CA, USA) according to the manufacturer's instructions.

124
125 **PacBio and Illumina PenSeq capture**
126
PacBio library was constructed with DNA samples from the mycelium of four *P. infestans*

strains, EU_13_A2 (2006_3928A), EC1_A1 (EC1_3626), EU_6_A1 (2006_3920A) and US23.

The library construction and target DNA sequence capture were performed according to (Witek

et al., 2016), with minor modifications. Qubit Fluorometer (ThermoFisher, Dubuque, IA, USA)

was used to quantify the barcoded DNA library from each isolate. Equimolar amounts of DNA

from the four individually barcoded samples were pooled to obtain 250 ng of total DNA and

then subjected to sequence capture. A 10x excess of non-adaptor-ligated *P. infestans* DNA at

about 500-1,000 bp was added for the hybridization. The final mixture of the amplicons of the

captured library was further size selected by SageELF electrophoresis system (Sage Science,

MA, USA) according to the instructions of the manufacturer.

Illumina library was constructed with RNA samples from zoospores of the four *P. infestans*

strains, from the corresponding infected leaf discs harvested at 12 hours post infection (hpi), 1,

2 and 3 days post infection (dpi), and from mycelium of EU_13_A2. An Illumina library for

each sample was constructed with KAPA mRNA HyperPrep Kit for Illumina® Platforms

(KR1352 – v5.17) following the manufacturer's instructions. mRNA was fragmented to 300-

400 bp. The barcoded libraries were mixed together at a ratio of 16: 8: 4: 1: 1 for 12 hpi, 1 dpi,

2 dpi, 3 dpi, zoospores and mycelium samples, respectively.

Both types of libraries were subjected to sequence capture using the bait library as described
previously (Jouet *et al.*, 2018, Thilliez *et al.*, 2018). Before and after sequence capture, qPCR
was performed on Bio-Rad CFX96 real-time detection system with an input of 1 ng DNA to
assess the efficiency of capture.

149
150 **Sequencing**
151
152 PacBio PenSeq libraries were sequenced at the Earlham Institute (Norwich, UK) using Sequel
153 platform. Illumina PenSeq cDNA libraries were sequenced at Novogene (Hong Kong, China)
154 using HiSeq, PE250.

155
gDNA PenSeq assembly

PacBio raw reads were processed as described in (Witek et al., 2016) to generate ROI reads
and demultiplexed using custom script (Van de Weyer et al., 2019). Demultiplexed ROI were
assembled using Geneious R8 (http://www.geneious.com/) using settings as in (Witek et al.,
2016).

Analysis of cDNA PenSeq
All RxLR effectors from the *P. infestans* reference genome T30-4 were used to generate an
artificial "RxLRome" contig, where RxLR effectors' sequences were separated by stretches of
500 "Ns". The contig also contained nine non-RxLR control genes (Jouet et al., 2018). The
cDNA PenSeq reads from all treatments were mapped to the T30-4 RxLRome, and the
expression analyses were performed and visualized using Geneious R10 (Kearse et al., 2012).

168
169 **New candidate RxLR effectors**
170

For the previously untested RxLR effectors, we first selected the effectors showing differential
expression at different stages and ranked them based on the raw transcript counts. Next, local
alignment searches (BLAST) were performed against the 563 predicted RxLR effectors (Haas
et al., 2009) to remove the previously tested effectors. This analysis revealed 47 candidate
RxLR effectors which were not included in previous functional study. The 47 RxLR effectors
were synthesized by Twist Bioscience (San Francisco, CA, USA). The signal peptides were
removed, the sequences were domesticated for Golden Gate cloning, and overhangs containing
BsaI restriction sites were added to both ends of all effector sequences.

All the effectors were cloned into vector pICSL86977 (TSL SynBio) with CaMV 35S promoter
and OCS terminator. The constructs were transformed to *Agrobacterium* strain GV3101 for
agro-infiltration.

183
184
185 **Cell death assay**
186
187 Transient expression of RxLR effectors and *Rpi-amr1* in *Nicotiana benthamiana* was
188 performed as described previously (Bos *et al.*, 2006). *Agrobacterium* was infiltrated at
189 OD600=1, and each effector was co-infiltrated with *Rpi-amr1-2273* (Witek *et al.*, 2020). The
190 cell death phenotype was observed at 4 dpi.

191
192 **Data availability**
193 Raw PacBio and cDNA PenSeq read sequences have been deposited in the Sequence Read
194 Archive (SRA) under BioProject IDs PRJNA623167 and PRJNA598824.

195
196
197
198 **Results**
199
200
Available recombinant RxLR effector libraries do not contain *Avramr1*
To identify *Avramr1*, we tested 278 available RxLR effectors (Table S1) by co-expressing
them with *Rpi-amr1-2273* in *N. benthamiana* (Rietman, 2011; Witek *et al.*, 2020). However,
no effector activated *Rpi-amr1*-dependent HR, so we concluded that *Avramr1* was absent from
the available RxLR effector libraries. Notably, *Avr8* was not originally included in the core
effector selection, because *Avr8* expression goes up earlier then 2 dpi (Jo, 2013), showing that
the criteria adopted to define core effectors did not reveal all recognized effectors.

208
To find *Avramr1*, we proposed three hypotheses: 1) *Avramr1* is an RxLR effector but it is not
present in the assembled version of *P. infestans* T30-4 reference genome 2) *Avramr1* is an
RxLR effector but it was not yet tested in previous functional studies/libraries; 3) *Avramr1* is
not a typical RxLR effector. To address hypothesis 1, we performed PacBio PenSeq to
sequence the effector alleles in the four diverse *P. infestans* isolates, EU_13_A2 (2006_3928A),
EC1_A1 (EC1_3626), EU_6_A1 (2006_3920A) and US23, all of them avirulent on potato
plants carrying *Rpi-amr1*, it indicates they all carry the recognized effector. To address

hypothesis 2, we performed cDNA PenSeq to try to identify other RxLR effectors that are
expressed during infection but not reported or defined in previous functional studies.

218
219 **PacBio PenSeq of four *P. infestans* isolates EU_13_A2, EC1_A1, EU_6_A1 and US23**
220

PacBio gDNA PenSeq was performed on four *P. infestans* isolates of genotypes EU_13_A2,
EC1_A1, EU_6_A1 and US23 (Fig. 1a). To evaluate the enrichment efficiency, qPCR was
performed with the DNA pre- and post- capture. In general, the targeted genes of different
length were well enriched at Concentration x time (Cot) value <20, while the untargeted genes
were almost undetectable, with Cot value >27 (Peterson *et al.*, 2002) (Fig. S1). Furthermore,
we found that the capture efficiency was increased by including a 10-fold molar excess of non-
adaptor-ligated fragmented *P. infestans* DNA (500-1,000 bp) in the reannealing reaction, to
reduce the extent to which sequences were recovered due to concatenation of transposon-
containing sequences adjacent to *RxLR* genes. After sequence capture, enrichment of most
effector genes was more efficient when non-adaptor-ligated *P. infestans* DNA was included
(Fig. S1).

Following the enrichment sequencing, circular consensus sequencing (CCS) reads were
assembled (Fig. 1a) and contigs with fewer than 10 reads were removed. The average length
of the contigs of coverage >10 reads was 7 kb (Table 1), and the size of the largest contig was
over 50 kb. This suggests that the PacBio PenSeq successfully captured the target effector
genes and the adjacent flanking DNA sequences. In total, 1,137, 1,054, 1,283 and 925 contigs
were obtained from EU_13_A2, EC1_A1, EU_6_A1 and US23 respectively, of which 687,
650, 741 and 571 contigs contain RxLR effectors. (Table 1, Notes S1-S4). The remaining
contigs contained non-RxLR effectors which were included in the bait library design for other
purposes (Thilliez *et al.*, 2018).

The PacBio PenSeq data allowed us to detect new RxLR effector alleles from different
haplotypes of various *P. infestans* isolates, and even in polyploid genotypes like EU_13_A2
(Li *et al.*, 2017). This dataset can also be used to extensively study allelic variation,
presence/absence (P/A) polymorphism and effector evolution. For example, *Avr1*
(PITG_16663) and a paralogous *Avr1-like* gene (PITG_06432) are located on supercontigs
1.51 and 1.8 of the reference T30-4 genome, respectively. The *R1*-breaking clonal lineage
EU_13_A2 was reported to have an 18 kb deletion comprising the *Avr1* locus (Cooke *et al.*,

2012). Also, the Illumina PenSeq data showed that the *Avr1* locus is missing in EU_13_A2,

EC1_A1 and US23 (Thilliez *et al.*, 2018). We mapped the four *Avr1* contigs from EU_13_A2

(contig 192, 261, 296 and 329) to supercontig 1.51 and 1.8, and found that all four contigs map

to the *Avr1-like* supercontig 1.8. Two contigs (contig 261 and 286) mapped to the *Avr1-like*

locus, and two other contigs (contig 192 and 329) mapped to a locus next to *Avr1-like* that was

not previously annotated (Fig. S2), though the genes in those two contigs might be pseudogenes

as the signal peptide is missing in both of them. Additionally, in EU_6_A1 and US23, two

Avr1 contigs did not map to *Avr1* or *Avr1-like* loci of T30-4. Thus, our PacBio PenSeq dataset

can provide means to detect novel RxLR effector paralogs absent from the reference genome.

As another example, our dataset carries in total 504 of the 563 predicted RxLR effectors from

the reference genome T30-4 (Haas *et al.*, 2009). To investigate P/A polymorphism of RxLR

effectors in the four sequenced isolates, we performed a basic local alignment search (BLAST)

of the 504 effectors against the PacBio contigs, with hits with < 50% coverage defined as absent

(Table S2). We found that 17, 28, 15 and 33 RxLR effectors out of the 504 are missing in

EU_13_A2, EC1_A1, EU_6_A1 and US23, respectively.

Taken together, we have generated a rich dataset that could help to define full length RxLR

effector genes, deliver robust information on alleles and paralogs, and reveal conserved or race-

specific effectors from different isolates. It is available in full in Notes S1-S4.

cDNA PenSeq enables effector expression detection in early stages of infection

To clarify whether the untested effectors might be putative *Avr* genes, we performed cDNA

PenSeq for the four *P. infestans* isolates EU_13_A2, EC1_A1, EU_6_A1 and US23, at

different time points post-infection (12 hpi, 1, 2 and 3 dpi) and in mycelium and zoospores

(Fig. 1b). To analyse and visualize the cDNA PenSeq data, we built an artificial DNA sequence

contig ("RxLRome") for the RxLR effectors. In addition, nine non-RxLR genes from the bait

library were included as controls (Jouet et al., 2018; Fig. 2). The cDNA PenSeq reads were

mapped to the RxLRome and gene expression compared over time (Fig. 1b).

Most of the RxLR effectors which were included in previous effector libraries show an up-

regulation of expression in the early stages of infection (Fig. 2). Some of the untested RxLR

effectors show a similar pattern of expression, and might also represent potential *Avr* genes,

while others are poorly expressed in some isolates. The details of the cDNA PenSeq are
available in Table S3.

Identification of *Avramr1*
288
To test if *Avramr1* is among the untested effectors, we selected 47 highly expressed effectors
(Fig. 3) present in all tested lineages that had not previously been investigated. The effectors
were synthesized, cloned into an expression vector with 35S promoter and transformed into
Agrobacterium GV3101 for agro-infiltration in *N. benthamiana* (Fig. 4a). All the effectors
were infiltrated alone or co-infiltrated with *Rpi-amr1-2273* (Witek *et al.*, 2020). Among the 47
effectors, PITG_07569 was the only effector which triggered an HR when co-expressed with
Rpi-amr1-2273 (Fig. 4b). Hence, we concluded PITG_07569 is *Avramr1*.

***Avramr1* homologs in different *P. infestans* isolates and other *Phytophthora* species**
298
Avramr1 is a canonical RxLR effector with RYLR and EER motifs and an N-terminal signal
peptide (Fig. 5b). *Avramr1* locates on supercontig 1.11 of the *P. infestans* reference genome
T30-4. *Avramr1-like* (hereafter *Avramr1L*), a truncated paralog (PITG_07566) maps adjacent
to *Avramr1* (Fig. 5a and 5b). Two known *Avr* effectors, *Avr8* (PITG_07558) and *Avrsmira1*
(PITG_07550), are physically close to the *Avramr1* locus in the T30-4 genome (Fig. 5a)
(Rietman *et al.*, 2012).

To study the sequence polymorphism of *Avramr1* homologs in *P. infestans*, we used BLAST
to search for *Avramr1* homologs in the PacBio PenSeq assemblies generated in this study. It
revealed that EU_13_A2, EC1_A1, EU_6_A1 and US23 carry six, four, three and six *Avramr1*
homologs, respectively. Next, we aligned the corresponding Avramr1 amino acid sequences
and generated a neighbourhood joining (NJ) tree for phylogenetic analysis (Fig. 6a). Two
Avramr1 homologs from *Phytophthora parasitica* and *Phytophthora cactorum* were identified
from public database, and they were used as an outgroup (Fig. 5b and 6a). Based on the
phylogenetic tree, we distinguished four Avramr1 clades, clade A (containing Avramr1 from
T30-4) and clade C (with Avramr1L from T30-4), and two more clades, B and D (Fig. 5a). For
a more detailed analysis, we selected one *Avramr1* homolog from clade B and one from D
(*Avramr1-13B1* and *Avramr1-13D1* from EU_13_A2) and aligned them with *Avramr1*
homologs from clade A and C, and with *P. parasitica* and *P. cactorum* homologs. Significant

sequence polymorphisms are observed between effectors from different clades (Fig. 5b).
Meanwhile, the *Avramr1* homologs within the same clade are almost identical (Fig. 6a).

Differential expression of *Avramr1* homologs in different *P. infestans* isolates

To investigate the expression patterns of *Avramr1* homologs defined in the PacBio PenSeq
data, we mapped the corresponding cDNA PenSeq reads to the PacBio PenSeq contigs from
EU_13_A2, EC1_A1, EU_6_A1 and US23. The transcript per million (TPM) values for each
time point are visualized in Fig. 6b. The clade A homologs *Avramr1-23A1*, *Avramr1-13A1* and
Avramr1-13A2 are highly expressed at almost all stages, and the *Avramr1* homologs from clade
B show a similar expression pattern. For clade C, some homologs, like *Avramr1-13C*,
Avramr1-6C1 or *Avramr1L* gene from T30-4, are weakly expressed at all stages. However, two
other *Avramr1L* homologs, *Avramr1-23C1* and *Avramr1-13C2,* show moderately elevated
expression in zoospores, and at 1 dpi and 3 dpi. Interestingly, the *Avramr1* homologs from
clade D, which are missing in the reference genome T30-4, show an intermediate expression
level compared to Clade A, B and Clade C, and most *Avramr1* homologs in Clade D show an
increase in expression at the zoospore stage, and at 1, 2, 3 dpi.

In summary, our PacBio PenSeq analysis created a rich dataset to reveal new *Avr* variants from
different *P. infestans* isolates, and to quantify their expression profile individually. This
facilitates the analysis of the polymorphism of pathogen effectors and their potential
differential recognition patterns with the corresponding *Rpi* genes (Witek *et al.*, 2020).

340
341

Discussion

The availability of the *P. infestans* genome sequence enabled a step-change in the rate of
investigation of this pathogen, accelerating the discovery of recognized effectors, and of new
Rpi genes (Haas et al., 2009; Vleeshouwers et al., 2008; 2011). However, some questions
remain open. For example, how different are the effector repertoires in different *P. infestans*
isolates? To what extent do they show differential expression between races? The study of
plant *NLR* gene repertoires faces similar challenges, and sequence capture, combined with
long-read sequencing technologies, has enabled the refinement of tools to cost-effectively
investigate diversity, such as RenSeq, SMRT RenSeq, RLP/KSeq and AgRenSeq (Arora et al.,

2019; Lin *et al.*, 2020; Jupe et al., 2013; Witek et al., 2016). Recently, the pan-NLRome of 65

diverse *Arabidopsis thaliana* accessions was determined by a similar strategy, revealing that

any one accession lacks many of the NLRs found in the species pan-NLRome (Van de Weyer

et al., 2019).

Pathogen-enrichment sequencing (PenSeq) was developed to facilitate cost-effective

investigation of pathogen diversity on infected plants, and polymorphism of pathogen effectors

(Jouet et al., 2018; Thilliez et al., 2018). The first PenSeq studies, however, were conducted

using Illumina short reads. This significantly limited their resolving power as many oomycete

genomes are highly heterozygous, and some effectors belong to large gene families with

multiple sequence-related paralogs that can lead to false assemblies (Gilroy et al., 2011; Oliva

et al., 2015).

In this study, we combined long read PenSeq and cDNA Penseq, enabling a detailed analysis

of the RxLR genes and their expression patterns in different *P. infestans* isolates. The cDNA

PenSeq dataset allowed us to define an additional set of 47 RxLR genes expressed during

infection that were not previously investigated. Amongst these, we identified *Avramr1*, which

encodes the cognate recognized effector for *Rpi-amr1* from *S. americanum* (Witek *et al.*, 2020).

The long read PenSeq data helped us to obtain full-length RxLR effector haplotypes with their

flanking sequences. This allowed us to distinguish individual alleles from polyploid isolates

like EU_13_A2, and also distinct effector paralogs. The sequences flanking the *RxLR* genes

enabled us to understand the possible translocation events and identify new *RxLR* loci. We

were also able to identify multiple new *Avramr1* homologs from different isolates, and

identified a new *Avramr1* Clade D which is not present in T30-4. The PenSeq dataset

constitutes a valuable community resource for investigating the allelic and expression diversity

of multiple recognized effectors.

So far, no *Rpi-amr1*-breaking *P. infestans* isolates have been found (Witek *et al.*, 2020), and

therefore we propose that *Avramr1* might be crucial for the virulence of *P. infestans*. The

identification of *Avramr1* will enable us to study its virulence function, its polymorphism in

the *P. infestans* population and its recognition by *Rpi-amr1*. Collectively, these data and

methods will contribute to understanding this fast-evolving and destructive oomycete pathogen,

and to achieving durable late blight resistance in potato.

Reference

Ballvora A, Ercolano MR, Weiß J, Meksem K, Bormann CA, Oberhagemann P, Salamini F, Gebhardt C. **2002**. The *R1* gene for potato resistance to late blight (*Phytophthora infestans*) belongs to the leucine zipper/NBS/LRR class of plant resistance genes. *The Plant Journal* **30**: 361–371.

Bos JIB, Kanneganti TD, Young C, Cakir C, Huitema E, Win J, Armstrong MR, Birch PRJ, Kamoun S. **2006**. The C-terminal half of *Phytophthora infestans* RXLR effector AVR3a is sufficient to trigger R3a-mediated hypersensitivity and suppress INF1-induced cell death in *Nicotiana benthamiana*. *The Plant Journal* **48**: 165–176.

Colon IT, Eijlander R, Budding DJ, Van Ijzendoorn MT, Pieters MMJ, Hoogendoorn J. **1993**. Resistance to potato late blight (*Phytophthora infestans* (Mont.) de Bary) in *Solanum nigrum*, *S. villosum* and their sexual hybrids with *S. tuberosum* and *S. demissum*. *Euphytica* **66**: 55–64.

Cooke DEL, Cano LM, Raffaele S, Bain RA, Cooke LR, Etherington GJ, Deahl KL, Farrer RA, Gilroy EM, Goss EM, et al. **2012**. Genome analyses of an aggressive and invasive lineage of the Irish potato famine pathogen. *PLoS Pathogens* **8**: e1002940.

Foster SJ, Park T-H, Pel M, Brigneti G, Śliwka J, Jagger L, van der Vossen E, Jones JDG. **2009**. *Rpi-vnt1.1*, a *Tm-22* homolog from *Solanum venturii*, confers resistance to potato late blight. *Molecular Plant-Microbe Interactions* **22**: 589–600.

Gilroy EM, Breen S, Whisson SC, Squires J, Hein I, Kaczmarek M, Turnbull D, Boevink PC, Lokossou A, Cano LM, et al. **2011**. Presence/absence, differential expression and sequence polymorphisms between *PiAVR2* and *PiAVR2*-like in *Phytophthora infestans* determine virulence on *R2* plants. *New Phytologist* **191**: 763–776.

Haas BJ, Kamoun S, Zody MC, Jiang RHY, Handsaker RE, Cano LM, Grabherr M, Kodira CD, Raffaele S, Torto-Alalibo T, et al. **2009**. Genome sequence and analysis of the Irish potato famine pathogen *Phytophthora infestans*. *Nature* **461**: 393–398.

Huang S, van der Vossen EAG, Kuang H, Vleeshouwers VGAA, Zhang N, Borm TJA, Van Eck HJ, Baker B, Jacobsen E, Visser RGF. **2005**. Comparative genomics enabled the isolation of the *R3a* late blight resistance gene in potato. *The Plant Journal* **42**: 251–261.

Jo KR. **2013**. Unveiling and deploying durability of late blight resistance in potato. PhD thesis, Wageningen UR.

Jouet A, Saunders DGO, McMullan M, Ben Ward, Furzer O, Jupe F, Cevik V, Hein I, Thilliez GJA, Holub E, et al. **2018**. *Albugo candida* race diversity, ploidy and host-associated microbes revealed using DNA sequence capture on diseased plants in the field. *New Phytologist* **93**: 959.

Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, et al. **2012**. Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28**: 1647–1649.

Li Y, Shen H, Zhou Q, Qian K, van der Lee T, Huang S. **2017**. Changing ploidy as a strategy:
the Irish potato famine pathogen shifts ploidy in relation to its sexuality. *Molecular Plant-*
Microbe Interactions **30**: 45–52.

Lin X, Armstrong M, Baker K, Wouters D, Visser RGF, Wolters PJ, Hein I,
Vleeshouwers VGAA. **2020**. RLP/K enrichment sequencing; a novel method to identify
receptor-like protein (*RLP*) and receptor-like kinase (*RLK*) genes. *New Phytologist*.
https://doi.org/10.1111/nph.16608

Oh S-K, Young C, Lee M, Oliva R, Bozkur TO, Cano LM, Win J, Bos JIB, Liu H-Y, van
Damme M, *et al*. 2009. In planta expression screens of *Phytophthora infestans* RXLR effectors
reveal diverse phenotypes, including activation of the *Solanum bulbocastanum* disease
resistance protein *Rpi-blb2*. *The Plant Cell* **21**: 2928–2947.

Oliva RF, Cano LM, Raffaele S, Win J, Bozkur TO, Belhaj K, Oh S-K, THINES M,
Kamoun S. **2015**. A recent expansion of the RXLR effector gene *Avrblb2* is maintained in
global populations of *Phytophthora infestans* indicating different contributions to virulence.
Molecular Plant-Microbe Interactions **28**: 901–912.

Pel M. **2010**. Mapping, isolation and characterization of genes responsible for late blight
resistance in potato. PhD thesis, Wageningen UR.

Pel MA, Foster SJ, Park T-H, Rietman H, van Arkel G, Jones JDG, Van Eck HJ,
Jacobsen E, Visser RGF, van der Vossen EAG. **2009**. Mapping and cloning of late blight
resistance genes from *Solanum venturii* using an interspecific candidate gene approach.
Molecular Plant-Microbe Interactions **22**: 601–615.

Peterson DG, Wessler SR, Genetics APTI, 2002. **2002**. Efficient capture of unique sequences
from eukaryotic genomes. *Journal of Integrative Agriculture* **18**: 547–550.

Rehmany AP, Gordon A, Rose LE, Allen RL, Armstrong MR, Whisson SC, Kamoun S,
Tyler BM, Birch PRJ, Beynon JL. **2005**. Differential recognition of highly divergent downy
mildew avirulence gene alleles by *RPP1* resistance genes from two *Arabidopsis* lines. *The*
Plant Cell **17**: 1839–1850.

Rietman H. **2011**. Putting the *Phytophthora infestans* genome sequence at work: multiple
novel avirulence and potato resistance gene candidates revealed. PhD thesis, Wageningen UR.

Rietman H, Bijsterbosch G, Cano LM, Lee H-R, Vossen JH, Jacobsen E, Visser RGF,
Kamoun S, Vleeshouwers VGAA. **2012**. Qualitative and quantitative late blight resistance in
the potato cultivar Sarpo Mira is determined by the perception of five distinct RXLR effectors.
Molecular Plant-Microbe Interactions **25**: 910–919.

Salaman R. **1937**. *Potato variety production: a new departure*. *Gardeners Chronicle* **102**: 326-
327.

Thilliez GJA, Armstrong MR, Lim TY, Baker K, Jouet A, Ben Ward, van Oosterhout C,
Jones JDG, Huitema E, Birch PRJ, *et al*. 2018. Pathogen enrichment sequencing (PenSeq)
enables population genomic studies in oomycetes. *New Phytologist* **4**: 903.

Toxopeus HJ. **1956**. Reflections on the origin of new physiologic races in *Phytophthora*
infestans and the breeding for resistance in potatoes. *Euphytica* **5**: 221–356.

**Van de Weyer A-L, Monteiro F, Furzer OJ, Nishimura MT, Cevik V, Witek K, Jones
JDG, Dangl JL, Weigel D, Bemm F**. **2019**. A species-wide inventory of *NLR* genes and alleles
in *Arabidopsis thaliana*. *Cell* **178**: 1260–1272.e14.

**van der Vossen E, Sikkema A, Hekkert BTL, Gros J, Stevens P, Muskens M, Wouters D,
Pereira A, Stiekema W, Allefs S**. **2003**. An ancient *R* gene from the wild potato species
Solanum bulbocastanum confers broad-spectrum resistance to *Phytophthora infestans* in
cultivated potato and tomato. *The Plant Journal* **36**: 867–882.

**Vleeshouwers VGAA, Raffaele S, Vossen JH, Champouret N, Oliva R, Segretin ME,
Rietman H, Cano LM, Lokossou A, Kessel G,** *et al.* **2011**. Understanding and exploiting late
blight resistance in the age of effectors. *Annual Review of Phytopathology* **49**: 507–531.

**Vleeshouwers VGAA, Rietman H, Krenek P, Champouret N, Young C, Oh S-K, Wang
M, Bouwmeester K, Ben Vosman, Visser RGF,** *et al.* **2008**. Effector genomics accelerates
discovery and functional profiling of potato disease resistance and *Phytophthora infestans*
avirulence genes. *PLoS ONE* **3**: e2875.

Vossen JH, van Arkel G, Bergervoet M, Jo K-R, Jacobsen E, Visser RGF. **2016**. The
Solanum demissum R8 late blight resistance gene is an *Sw-5* homologue that has been deployed
worldwide in late blight resistant varieties. *Theoretical and Applied Genetics* **129**: 1785–1796.

Witek K, Jupe F, Witek AI, Baker D, Clark MD, Jones JDG. **2016**. Accelerated cloning of
a potato late blight–resistance gene using RenSeq and SMRT sequencing. *Nature
Biotechnology* **34**: 656–660.

**Witek K, Lin X, Karki HS, Jupe F, Steuernagel B, Stam R, van Oosterhout C, Fairhead
S, Cocker JM, Bhanvadia S,** *et al.* **2020**. A complex non-host resistance locus in *Solanum
americanum* recognizes a conserved *Phytophthora* effector. *bioRxiv*: 1–25.

Acknowledgements:

492 This research was finance from BBSRC grants BB/P021646/1. We would like to thank

493 Sophien Kamoun and Joe Win for valuable discussion, and TSL Bioinformatics Team,

494 SynBio Team and horticultural team for their support.

504 **Table:**

Table 1. PacBio PenSeq for EU_13_A2, EC1_A1, EU_6_A1 and US23

	Total contigs	RxLR effector	Non_RxLR	Average length (bp)	Minimum length (bp)	Maximum length (bp)
EU_13_A2	1,137	687	450	7,732	2,617	30,571
EC1_A1	1,054	650	404	7,663	2,696	33,203
EU_6_A1	1,283	741	542	7,463	3,052	50,275
US23	925	571	354	7,598	2,648	29,901

 Figures:

Fig. 1 The pipelines of PacBio and cDNA PenSeq.

(a) The pipeline of PacBio gDNA PenSeq. Briefly, the gDNA isolated from various
Phytophthora infestans was enriched for RxLR effectors, sequenced by PacBio and de-
novo assembled for data mining.

(b) The pipeline of cDNA PenSeq. The cDNA were synthetized using RNA sampled from
various *P. infestans* at different stages (mycelium, zoospore, 12 hpi, 1, 2 and 3 dpi).
The libraries enriched for RxLR effectors were sequenced, reads were mapped to the
RxLRome of the reference *P. infestans* genome T30-4 and the expression levels of
samples were calculated and compared.

Black lines with dots represent the baits, the enriched fragments are depicted in blue
(EU_13_A2), yellow (EC1_A1), pink (EU_6_A1) and green (US23). The black boxes
indicate RxLR effectors. EU_13_A2, EC1_A1, EU_6_A1, US23, *P. infestans* genotypes.

545

Fig. 2 cDNA PenSeq of RxLR effectors from EU_13_A2, EC1_A1, EU_6_A1 and US23

547

The cDNA PenSeq data for the RxLR effectors from four *Phytophthora infestans* at different stages were mapped to an artificial contig (RxLRome) of 499 RxLR effectors and nine non-RxLR genes, demarcated by bright green arrows on the outer edge of the diagram. Black lines separate the previously tested RxLR effectors (grey bar), new effector candidates with differential expression (red bar), unexpressed effectors (blue bar) and non-RxLR controls (cyan). The concentric circles in blue, yellow, pink and green represent data from *P. infestans* EU_13_A2, EC1_A1, EU_6_A1 and US23, respectively. The arrows on them indicate differential expression (red, up-regulation, blue, down-regulation; no fill, no difference), where the more intense the colour, the bigger the difference. The data are plotted as follows: a, mycelium vs zoospores (for EU_13_A2 only); b, zoospores vs 12 hpi; c, 12 hpi vs 1 dpi; d, 1 dpi vs 2 dpi; e, 2 dpi vs 3 dpi. Six known *Avr* genes, *AvrSmira1*, *Avr3a*, *Avrvnt1*, *Avrblb2*, *Avrblb1* and *Avr2* are indicated by black arrows.

560

561

562

Fig. 3 Raw transcript counts for the new candidate RxLR effectors.

47 most differentially expressed RxLR effectors from the previously untested set were selected, and the raw transcript counts were visualized as a heat map across time points and treatments. Each square indicates a single data point derived from two independent biological replicates. The colours red, orange, yellow and blue represent >100, 50-100, 0-50, or 0 raw transcripts, respectively.

Fig. 4 Identification of *Avramr1*.

(a) All 47 selected effectors without signal peptides (SP) were synthesized and cloned into

an expression vector under CaMV-35S promoter for *Agrobacterium*-mediated transient

expression.

(b) Transient expression of candidate effectors on their own or with *Rpi-amr1-2273* in

Nicotiana benthamiana. Dashed circles demarcate the infiltration sites. Only

PITG_07569 triggers HR when co-expressed with *Rpi-amr1-2273*. All other effectors

that did not trigger HR are represented by PITG_06375 and PITG_06305. A known

R/Avr gene pair was used as positive control (+). This experiment was repeated more

than 10 times with the same results.

Fig. 5 Genomic localization and amino-acid alignment of *Avramr1*.

(a) The localization of *Avramr1* (*PITG_07569,* red arrow) on supercontig 1.11 of the

592   reference *Phytophthora infestans* T30-4 genome. A paralog *Avramr1L* gene

593   (*PITG_07566*, blue arrow) is located close to *Avramr1*. The supercontig contains

594   another two known *Avr* genes, *Avrsmira1* (*PITG_07550*, pink arrow) and *Avr8*

595   (*PITG_07558*, green arrow).

596   (b) The alignment of protein sequences of Avramr1 and selected homologs and paralogs

597   from *P. infestans*, *P. capsici* (Pc) and *P. palmivora* (Pp). The dark green bars on top of

598   the alignment indicate 100 % identity while olive green and red bars indicate various

599   degrees of polymorphism between the sequences. RxLR and EER motifs are

600   highlighted by red boxes.

Fig. 6 Phylogeny and expression profile of *Avramr1* homologs from EU_13_A2, EC1_A1,

EU_6_A1 and US23.

(a) Neighbor joining tree of the protein sequences of the *Avramr1* homologs. Pc-Avramr1

and Pp-Avramr1 were used as outgroups.

(b) The expression profile of *Avramr1* homologs at different stages and time points

(zoospores, 12hpi, 1, 2 and 3 dpi). Transcripts per kilobase million (TPM) for each

effector homologs were visualized as follows: black, data not available; red, 350,000-

1,000,000 TPM; orange, 200,000-350,000 TPM; yellow, 0-200,000 TPM; beige, 0

TPM.

Supporting Information:

Notes S1: PacBio PenSeq contigs of EU_13_A2.

Notes S2: PacBio PenSeq contigs of EC1_A1.

Notes S3: PacBio PenSeq contigs of EU_6_A1.

Notes S4: PacBio PenSeq contigs of US23.

Table S1: 278 RxLR effectors from previously available effector libraries.

Table S2: P/A polymorphism of RxLR effectors from EU_13_A2, EC1_A1, EU_6_A1 and US23.

Table S3: cDNA PenSeq for EU_13_A2, EC1_A1, EU_6_A1 and US23.

Fig. S1: Enrichment efficiency with/ without non-adaptor-ligated DNA.

Fig. S2: Compare EU_13_A2 *Avr1* contigs and the T30-4 reference genome.