Human Dicer helicase domain recruits PKR and dsRNA binding proteins during viral infection

Thomas C. Montavon[1,*], Morgane Baldaccini[1,*], Mathieu Lefèvre[1,*], Erika Girardi[1], Béatrice Chane-Woon-Ming[1], Mélanie Messmer[1], Philippe Hammann[2], Johana Chicher[2], Sébastien Pfeffer[1,‡]

[1] Université de Strasbourg, Architecture et Réactivité de l'ARN, Institut de Biologie Moléculaire et Cellulaire du CNRS, 2 allée Konrad Roentgen, 67084 Strasbourg France

[2] Université de Strasbourg, Institut de Biologie Moléculaire et Cellulaire du CNRS, Plateforme Protéomique Strasbourg – Esplanade, 2 allée Konrad Roentgen, 67084 Strasbourg France

*These authors contributed equally

‡To whom correspondence should be addressed: spfeffer@unistra.fr

Abstract

The antiviral innate immune response mainly involves type I interferon (IFN) in mammalian

cells. The contribution of the RNA silencing machinery remains to be established, but several

recent studies indicate that the ribonuclease DICER can generate viral siRNAs in specific

conditions. It has also been proposed that type I IFN and RNA silencing could be mutually

exclusive antiviral responses. In order to decipher the implication of DICER during infection

of human cells with the Sindbis virus, we determined its interactome by proteomics analysis.

We show that DICER specifically interacts with several double-stranded RNA binding proteins

and RNA helicases during viral infection. In particular, proteins such as DHX9, ADAR-1 and

the protein kinase RNA-activated (PKR) are enriched with DICER in virus-infected cells. We

demonstrate the importance of DICER helicase domain in its interaction with PKR and showed

that it has functional consequences for the cellular response to viral infection.

Introduction

In mammalian cells, the main antiviral defense system involves the activation of a signaling cascade relying on production of type I interferon (IFN I). This pathway depends on the recognition of extrinsic signals or pathogen associated molecular patterns (PAMPs) by dedicated host receptors. Double-stranded (ds) RNA, which can originate from viral replication or convergent transcription, is a very potent PAMP and can be sensed in the cell by various proteins among which a specific class of DExD/H-box helicases called RIG-I-like receptors (RLRs) (Ivashkiv and Donlin, 2014). RLRs comprise RIG-I, MDA5 and LGP2 and transduce viral infection signals to induce expression of IFN I cytokines that act in autocrine and paracrine fashion. These cytokines then trigger the expression of hundreds of interferon stimulated genes (ISGs) to stop the virus in its tracks (Borden et al, 2007). Among those ISGs, dsRNA-activated protein kinase R (PKR) plays an important role in antiviral defense by blocking cellular and viral translation upon direct binding to long dsRNA (Williams, 1999). PKR is a serine-threonine kinase that dimerizes and auto-phosphorylates upon activation. It then phosphorylates numerous cellular targets among which the translation initiation factor eIF2α, which results in the inhibition of cap-dependent translation (Lemaire et al., 2008). Accordingly, translation of many RNA viruses, including alphaviruses, is inhibited by PKR (Fros and Pijlman, 2016; Pfaller et al., 2011; Ryman et al., 2002). PKR is also involved in other cellular pathways including apoptosis, autophagy and cell cycle (Kim et al., 2014; Williams, 1999).

RNAi is another evolutionary conserved pathway triggered by long dsRNA sensing (Meister and Tuschl, 2004). One key component in this pathway is the type III ribonuclease DICER, which is also essential for micro (mi)RNA biogenesis (Hutvágner et al., 2001; Wienholds et al., 2003). These small regulatory RNAs are sequentially produced by the two ribonucleases DROSHA and DICER, before being loaded into an Argonaute (AGO) effector protein in order to regulate their target mRNAs (Bartel, 2018). Whatever its substrate, be it long

dsRNA or miRNA precursor, DICER relies on interacting with co-factors to be fully functional. In mammalian cells, the TAR-RNA binding protein (TRBP), a dsRNA binding protein (dsRBP), was shown to play a role in the selection of DICER substrates, its stabilization, strand selection and incorporation into AGO2 (Chendrimada et al., 2005). The interaction with TRBP is well characterized and depends on the helicase domain of DICER and the third dsRNA binding domain (dsRBD) of TRBP (Daniels et al., 2009). Another dsRBP, the protein activator of interferon-induced protein kinase R (PACT), was also described as an important cofactor of DICER. Although its function is not fully understood, PACT seems to also participate in miRNA loading and strand selection (Heyam et al., 2015; Kok et al., 2007) *via* protein-protein interaction between the DICER helicase domain and the third dsRBD of PACT (Lee et al., 2006).

It is now common knowledge that RNAi is the main antiviral defense system in several phyla such as plants, arthropods and nematodes (reviewed in (Guo et al., 2019)). However, its exact contribution in the mammalian antiviral response remains unclear (Cullen, 2006; Maillard et al., 2019; tenOever, 2016). Recent studies indicate that a functional antiviral RNAi does exist in mammals in specific cases. A functional antiviral RNAi response was first detected in undifferentiated mouse embryonic stem cells (Maillard et al., 2013) lacking the IFN response, suggesting that these two pathways could be incompatible. Indeed, in mammalian somatic cells deficient for MAVS or IFNAR, two components of the interferon response, an accumulation of DICER-dependent siRNAs derived from exogenous long dsRNA was detected (Maillard et al., 2016). In addition, the RLR LGP2 was found interacting with both DICER and TRBP, blocking respectively siRNA production and miRNA maturation (Takahashi et al., 2018a, 2018b; van der Veen et al., 2018). Moreover, AGO4 was recently shown to be involved in antiviral RNAi against Influenza A virus (IAV), Vesicular stomatitis virus (VSV) and Encephalomyocarditis virus (EMCV) (Adiliaghdam et al., 2020). Finally, viral suppressors of RNAi (VSRs) have been

shown to prevent DICER from playing an antiviral role in mammalian cells (Qiu et al., 2017,

2020). Nonetheless, several studies reported no detection of viral siRNAs in mammalian

somatic cells infected with several viruses (Girardi et al., 2013; Parameswaran et al., 2010;

Schuster et al., 2019). In somatic cells, only a helicase-truncated form of human DICER could

produce siRNAs from IAV genome (Kennedy et al., 2015), but it also turned out that these

siRNAs cannot confer an antiviral state (Tsai et al., 2018).

Based on these conflicting observations, we decided to study the involvement of DICER

during infection of human cells with the Sindbis virus (SINV). SINV is a member of the

Togaviridae family in the alphavirus genus, which is transmitted by mosquitoes to mammals

and can induce arthritogenic as well as encephalitic diseases (Griffin, 2007). It is widely used

as a laboratory alphaviruses model as it infects several cell types and replicates to high titers.

SINV has a positive stranded RNA genome of about 12 kb, which codes for two polyproteins

that give rise to non-structural and structural proteins, including the capsid. Moreover, upon

viral replication, a long dsRNA intermediate, which can be sensed by the host antiviral

machinery, accumulates. Of note, SINV dsRNA can be cleaved into siRNA in insects as well

as in human cells expressing the Drosophila DICER-2 protein (Girardi et al., 2015).

Nonetheless, although human DICER has the potential to interact with the viral RNA duplex,

we did not find evidence that SINV dsRNA could be processed into siRNAs in somatic

mammalian cells (Girardi et al., 2013, 2015). We thus hypothesized that specific proteins could

interfere with DICER during SINV infection by direct interaction and limit its accessibility

and/or activity. To address this hypothesis, we generated HEK293T cells expressing a tagged

version of human DICER that could be immunoprecipitated in mock or SINV-infected cells in

order to perform a proteomic analysis of its interactome. Among the proteins co-

immunoprecipitated with DICER and that were specifically enriched upon infection, we

identified dsRBPs such as ADAR1, DHX9, PACT and PKR. We further validated the direct

interaction between DICER and PKR upon SINV infection. We also demonstrated that the

interactions of the endogenous DICER with PKR, PACT and DHX9 could also be detected in

SINV-infected, but not mock-infected, HCT116 cells. We dissected the protein domains

necessary for this interaction and we found that DICER helicase domain plays a fundamental

role as a recruitment platform for PKR but also for other co-factors. Finally, we also show that

expression of a helicase-truncated version of DICER has a negative effect on SINV infection.

Our results indicate that DICER interactome is highly dynamic and directly link components

of RNAi and IFN pathways in modulating the cellular response to viral infection.

Results

Establishment of a HEK293T cell line expressing FLAG-HA tagged DICER

In order to be able to study the interactome of the human DICER protein during viral infection,

we transduced *Dicer* knock-out HEK293T cells (NoDice 2.20) (Bogerd et al., 2014a) with

either a lentiviral construct expressing a FLAG-HA-tagged wild type DICER protein

(FHA:DICER WT #4) or a construct without insert as a negative control (FHA:ctrl #1). After

monoclonal selection of stably transduced cells, we first characterized one clone of both

FHA:DICER WT and of the FHA:ctrl cell lines. We first confirmed that the expression of the

tagged version of DICER restored the miRNA biogenesis defect observed in the NoDice cells

(Fig. S1A). We then monitored the phenotype of these cells during SINV infection by using as

a readout of viral infection the modified version of SINV able to express GFP from a duplicated

sub-genomic promoter (SINV-GFP) (López et al, 2020). At 24 hours post-infection (hpi) and a

multiplicity of infection (MOI) of 0.02, the GFP fluorescence observed in FHA:DICER WT #4

cells and HEK293T cells was similar. However, the NoDice FHA:ctrl #1 cells displayed a

decrease in GFP signal (Fig. 1A). Western blot analysis of GFP expression confirmed the

observations by epifluorescence microscopy, *i.e*. a significantly lower accumulation of GFP in

the absence of the DICER protein (Fig. 1B). We therefore wished to confirm the effect of

DICER loss on SINV-GFP infection in another NoDice cell line, *i.e.* the NoDice clone 4.25

(Bogerd et al., 2014b), and in another clone of the NoDice 2.20 FHA:ctrl cells (NoDice

FHA:ctrl #2). We observed a similar decrease of SINV-GFP infection in NoDice 2.20 cells and

two independent NoDice FHA:ctrl clones compared to HEK293T cells as shown by GFP

microscopy (Fig. S1B), by titration of the virus (Fig. S1C) and by western blot analysis (Fig.

S1D). However, the independent NoDice 4.25 *Dicer* knock-out clone appeared mostly

unaffected compared to HEK293T cells in term of GFP accumulation and viral titer (Fig. S1B-

D). This suggests that, despite the observed slight effect on SINV-GFP in NoDice 2.20 cells

(Fig. 1), DICER antiviral effect is not reproductible in an independent clone and therefore could

not be generalized.

In order to evaluate whether different expression levels of DICER in a NoDice

background could rescue the SINV infection phenotype observed in HEK293T cells, we also

infected both the FHA:DICER WT polyclonal and an independent FHA:DICER WT clone

(FHA:DICER WT #17) with SINV-GFP (Fig. 1A and Fig. 1C-D). We confirmed that the GFP

fluorescence observed by microscopy (Fig. 1A), as well as the viral titers and the GFP protein

accumulation (Fig. 1C-D) in all tested FHA:DICER lines were comparable to the ones observed

in HEK293T cells. Moreover, there was no striking differences in AGO2 expression between

the FHA:DICER lines (Fig. 1D).

Altogether, these results indicate that the FHA-tagged DICER protein can functionally

complement the lack of DICER in terms of miRNA biogenesis (Fig. S1A) and can therefore be

used for proteomics studies. Moreover, because we could not observe significant differences in

terms of SINV infection (Fig. 1) between the different FHA:DICER clones tested, we decided

to select one line, namely FHA:DICER WT#4, for further analysis.

Analysis of DICER interactome during SINV infection by mass spectrometry

Our molecular tool being validated, we then focused on determining the interactome of FHA:DICER during SINV infection. We wanted to look at DICER interactome at an early infection time point to isolate cellular factors that could potentially modulate either DICER accessibility or its effect on viral dsRNA. As SINV replicates quickly upon cellular entry, we chose to set up the infection conditions to a duration of 6 hours at an MOI of 2.

We performed an anti-HA immunoprecipitation experiment (HA IP) coupled to label-free LC-MS/MS analysis in FHA:DICER WT #4 cells either mock-infected or infected for 6 h at an MOI of 2 with SINV-GFP. In parallel, we performed an anti-MYC immunoprecipitation as a negative control (CTL IP). The experiments were performed in technical triplicate in order to have statistically reproducible data for the differential analysis, which was performed using spectral counts. Prior to the detailed analysis of the results, we verified that there was no confounding factor in the experimentation by performing a Principal Component Analysis (PCA). This allowed us to see that the replicates were very homogenous and that the different samples were well separated based on the conditions.

To check the specificity of the HA immunoprecipitation, we first compared the proteins identified in the HA IP with the ones identified in the CTL IP in mock-infected cells. Differential expression analysis allowed us to calculate a fold change and an adjusted p-value for each protein identified and to generate a volcano plot representing the differences between HA and CTL IP samples. Applying a fold change threshold of 2 ((LogFC) > 1), an adjusted p-value threshold of 0.05 and a cutoff of at least 5 spectral counts in the most abundant condition, we identified 258 proteins differentially immunoprecipitated between the two conditions out of 1318 proteins (Fig. 2A and Supp. Table 1). Among these, 123 proteins were specifically enriched in the HA IP. The most enriched protein was DICER, followed by its known co-factors TRBP and PACT (also known as PRKRA) (Chendrimada et al., 2005; Lee et al., 2006). We

were also able to retrieve AGO2, indicating that the RISC loading complex was

immunoprecipitated and that proteins retrieved in our HA IP are specific to DICER

immunoprecipitation.

We next performed the differential expression analysis of proteins retrieved in the HA

IP in SINV-GFP compared to mock-infected cells. Among 1342 proteins, 296 were

differentially retrieved between conditions (Fig. 2B and Supp. Table 2). Of these, 184 proteins,

including viral ones, were at least 2-fold enriched in SINV-GFP-infected cells. GO-term

analysis showed a significant enrichment in RNA binding proteins including double-stranded

RNA binding proteins and RNA helicases (Fig. 2C). Among the RNA binding proteins

retrieved, the top and most specific DICER interactor is the interferon-induced, double-stranded

(ds) RNA-activated protein kinase PKR (also known as E2AK2), which is enriched more than

250 times in virus-infected cells (Fig. 2B and D). We were also able to identify the dsRNA-

specific adenosine deaminase protein ADAR-1 (also known as DSRAD), as well as PACT,

which were enriched 5.9 and 4.2 times respectively in SINV-GFP-infected cells compared to

mock-infected cells (Fig. 2B and D). Among the isolated RNA helicases, we identified the

ATP-dependent RNA helicase A protein DHX9, which is implicated in *Alu* element-derived

dsRNA regulation and in RISC loading (Aktaş et al., 2017; Robb and Rana, 2007). In order to

verify if the observed interactions were specific to SINV we performed the same experiments

with another virus of the *Togaviridae* family, the Semliki forest virus (SFV). In this analysis,

we were able to retrieve ADAR-1, DHX9, PACT and PKR, specifically enriched in SFV-

infected samples (Fig. S2 and Supp. Tables 3 and 4). These results show that these interactions

can be retrieved in *Togaviridae*-infected cells.

Taken together, our data indicate that several proteins interacting with DICER in virus-

infected cells are involved in dsRNA sensing and/or interferon-induced antiviral response.

DICER and PKR interact *in vivo* in the cytoplasm during SINV infection

To validate the LC-MS/MS analysis, we performed a co-immunoprecipitation (co-IP) followed by western blot analysis in FHA:DICER WT #4 cells infected with SINV-GFP at an MOI of 2 for 6 h. Whereas TRBP interacted equally well with FHA:DICER in mock and SINV-GFP-infected cells, ADAR-1, PKR, DHX9 and PACT were only retrieved in the HA IP in SINV-infected cells (Fig. 3A). We verified that these interactions could also be observed at a later time post-infection by performing the HA IP in FHA:DICER WT #4 cells infected with SINV-GFP for 24 h at an MOI of 0.02. This indicates that the specific interactions between DICER and ADAR-1, DHX9, PACT or PKR occur at an early stage of the SINV infection and remain stable in time in virus-infected cells (Fig. S3A).

In order to verify whether these interactions were mediated by RNA, we performed an anti-HA co-IP experiment on an RNase A/T1 treated total extract from FHA:DICER WT #4 cells infected with SINV-GFP at an MOI of 2 for 6h. We confirmed the efficiency of the RNase treatment by ethidium bromide staining visualisation of total RNA on an agarose gel (Fig. S3B). TRBP equally interacted with FHA:DICER, with or without RNase treatment, in mock and SINV-GFP-infected cells (Fig. 3B). Instead, the virus-induced interactions between DICER and PKR or PACT upon SINV-GFP infection were almost totally lost in the RNase-treated samples. Upon virus infection, PKR is phosphorylated to be activated and exert its antiviral function (Lemaire et al., 2008). Using an antibody targeting the phosphorylated form of PKR (p-PKR), we looked for p-PKR before and after RNase treatment. The virus-enriched interactions between DICER and p-PKR or DHX9 were completely lost upon RNase treatment. These results therefore indicate that RNA molecules (either single- or double-stranded) facilitate DICER interaction with DHX9, PACT and PKR and its active form, although the complex may also partially interact in an RNA-independent manner.

Because of the involvement of PKR in antiviral response (García et al., 2007) and the

fact that it shares common co-factors with DICER, namely TRBP and PACT (Haase et al.,

2005; Patel et al., 2000), we decided to focus our analysis on the DICER-PKR interaction. To

confirm the biological relevance of this interaction, we first performed a reverse co-IP to

immunoprecipitate the endogenous PKR protein in HEK293T cells infected or not with SINV-

GFP. While PACT interacted with PKR both in mock and in SINV-GFP-infected cells as

expected (Fig. 3C lower panel), DICER co-immunoprecipitated with the endogenous PKR only

in virus-infected cells as previously observed (Fig. 3C upper panel).

To further determine whether DICER and PKR could directly interact *in vivo*, we set up

a bi-molecular fluorescent complementation assay (BiFC) experiment (Lepur et al., 2016). To

this end, we fused the N- or C-terminal halves of the Venus protein ($^{N-ter}$Venus or $^{C-ter}$Venus) to

DICER and to PKR but also to TRBP and PACT. Since we showed above that an N-terminally

tagged DICER was functional, we fused the Venus fragments at the N-terminal end of DICER.

For the other three proteins, we fused the Venus fragments at the N- or C-terminus and selected

the best combination. To avoid interaction with the endogenous DICER and PKR proteins, we

conducted all BiFC experiments in NoDice/ΔPKR HEK293T cells (Kennedy et al., 2015). In

order to control the BiFC experiments, we chose to exploit the well characterized DICER-

TRBP interaction, which is known to occur via the DICER DEAD-box helicase domain

(Daniels et al., 2009). We therefore used the wild-type DICER protein as a positive control and

a truncated version of DICER protein lacking part of this helicase domain and called DICER

N1 (Kennedy et al., 2015) as a negative control (Fig. S3C). We first confirmed the expression

of the tagged proteins by western blot analysis (Fig. S3D) and then, we tested the interactions

between DICER and TRBP or PACT or PKR. We co-transfected the Venus constructs for 24 h

and then infected cells with SINV or not for 6 h at a MOI of 2. A comparable fluorescent signal

was observed both in mock- and SINV-infected cells when $^{N-ter}$Venus:DICER was co-

transfected with either PACT or TRBP fusion construct (Fig. 3D). Although we initially

expected an increase of the Venus fluorescence in SINV-infected cells, overall we observed a

similar signal for the DICER-PKR interaction both in mock- and SINV-infected cells, probably

due to the fact that both proteins are transiently overexpressed in this experiment.

As a control and to rule out any aspecific interactions between the different proteins

tested, we also monitored the DICER-N1-TRBP interaction by BiFC. As expected, no

fluorescent signal was observed in cells co-transfected with [N-ter]Venus:DICER N1 and

TRBP:Venus[C-ter] (Fig. S3E), confirming that DICER helicase domain is required for its

interaction with TRBP (Daniels et al., 2009) and validating the specificity of the BiFC

approach.

To further confirm that the absence of PKR did not influence the interactions of TRBP

or PACT with DICER, we also performed a BiFC analysis in HEK293T cells. After verifying

that in this context as well, fusion proteins were expressed as expected (Fig. S3F), we observed

that the results were similar as in NoDiceΔPKR cells (Fig. S3G).

To gain more insight into the subcellular localization of these interactions during SINV

infection, we performed the BiFC experiments, fixed the cells and observed them under a

confocal microscope. We observed a cytoplasmic fluorescent signal for DICER-TRBP and

DICER-PACT interactions (Fig. 3E upper and middle panels), which is in agreement with their

canonical localization for the maturation of most miRNAs (Bernstein et al., 2001; Hutvágner

et al., 2001). Similarly, co-transfection of DICER and PKR led to a strong Venus signal

homogeneously distributed in the cytoplasm (Fig. 3E lower panel).

Collectively, these results formally confirm that DICER interacts with several RNA

helicases and dsRNA-binding proteins in virus-infected cells, among which PKR, and that for

the latter this interaction occurs in the cytoplasm.

DICER interactome changes upon SINV infection are not cell-type specific

To further validate our previous DICER interactome results and generalize them to another biological system, we performed co-IP experiments on the endogenous DICER in a different cell type. To this end, the FLAG-HA-GFP tag was knocked into (KI) the *Dicer* locus in human colon carcinoma cells (HCT116) by CRISPR-Cas9-mediated homologous recombination (Fig. S4A-C). A guide RNA (gRNA) targeting the region corresponding to Dicer ATG and a DNA template for homologous recombination bearing the FLAG-HA-GFP sequence surrounded by the upstream and downstream arms of Dicer were used to generate the resulting cell line referred to as HCT116 KI-DICER cells. The expected insertion of the tag in one of the two Dicer allele was assessed by PCR amplification and Sanger sequencing (Fig. S4A-C). In agreement, we could detect two bands for DICER protein by western blot in the HCT116 KI-DICER cells, which confirmed that this cell line is heterozygous (Fig. 4A).

We additionally verified the expression of specific DICER-interacting proteins, such as AGO2, PKR or TRBP, in HCT116 KI-DICER cells compared to the parental HCT116 cells and to HEK293T cells (Fig. 4A). We also measured the production of mature miRNAs, such as miR-16, by northern blot analysis and confirmed that miRNA expression is maintained in HCT116 KI-DICER cells (Fig. 4B). Of note, the GFP inserted at the *Dicer* locus could not be detected by epifluorescence microscopy in the HCT116 KI-DICER cells, which probably reflects the low abundance of the DICER protein.

We then determined whether SINV-GFP infection was comparable in HCT116 cells and HEK293T cells. We infected HCT116, HCT116 KI-DICER and HEK293T cells with SINV-GFP at three different MOI (0.02, 0.1 and 1) and measured GFP fluorescence by microscopy at 24 hpi (Fig. 4C). Both HCT116 and HCT116 KI-DICER cells expressed GFP upon infection with SINV-GFP, although with a lower intensity than HEK293T cells. We also verified by western blot analysis the accumulation of GFP and the phosphorylation of both PKR and eIF2α

upon SINV-GFP infection of HCT116 KI-DICER and HEK293T cells (Fig. S4D) and chose as

optimal SINV-GFP condition of infection in HCT116 KI-DICER cells the MOI of 0.1 for 24h.

To validate the DICER interactions observed in HEK293T FHA:DICER cells, we then

performed anti-HA co-IP experiments followed by western blot analysis in HCT116 KI-DICER

cells infected or not with SINV-GFP. We successfully retrieved TRBP interacting with DICER

in both mock and infected cells, whereas DHX9, PKR (phosphorylated or not) and PACT were

only retrieved in the HA IP in infected cells (Fig. 4D). These results not only confirm that the

endogenous DICER specifically interacts with DHX9, PACT and PKR upon SINV infection,

but also that these interactions are not restricted to one specific cell type.

The helicase domain of DICER is required for its interaction with PKR

Even though DICER and PKR are likely brought together by RNA, specific protein domains

might be involved in stabilizing the complex. We therefore next determined the domain of

DICER required for its interaction with PKR. Since its helicase domain was previously shown

to be involved in the interaction with TRBP and PACT (Daniels et al., 2009; Lee et al., 2006),

we speculated that it could also be implicated in binding PKR. To test this hypothesis, we cloned

several versions of DICER proteins wholly or partly deleted of the helicase domain (Fig. 5A

DICER N1 and N3). In addition, we also cloned the helicase domain alone (Fig. 5A DICER

Hel.) and a DICER variant deleted of its C-terminal dsRNA binding domain (Fig. 5A DICER

ΔdsRBD) since this domain could also be involved in protein-protein interaction (Doyle et al.,

2013; Lambert et al., 2016). We then transfected the different versions of DICER WT and the

deletion mutant constructs in NoDice cells. In mock and SINV-GFP infected cells, whole cell

extracts were subjected to anti-HA and anti-MYC (CTL) IP. TRBP was retrieved in both

conditions with DICER WT, Hel. and ΔdsRBD (Fig. 5B and 5C). In mock cells, PACT and

PKR were only found weakly interacting with DICER WT (Fig. 5B). In SINV-infected cells,

we observed that similar to TRBP and to a lesser extent PACT, N1 and N3 mutations strongly

reduced the binding of DICER with PKR (Fig. 5C lanes 2-3 and 7-8). Importantly, we also

noted that the helicase domain alone could bind PKR, TRBP and PACT (Fig. 5C lanes 4 and

9). Moreover, the deletion of the dsRNA binding domain of DICER did not affect its interaction

with TRBP, PACT and PKR (Fig. 5C lanes 5 and 10). We also looked for p-PKR in our co-IP

(Fig. 5C panel p-PKR). We noticed that only WT DICER and its helicase domain were able to

interact with p-PKR (Fig. 5C lanes 1&6 and 4&9). The fact that DICER ΔdsRBD did not

interact with p-PKR (Fig. 5C lanes 5&10) is striking but could indicate that the phosphorylation

of PKR may induce conformation changes preventing its interaction with some domains of

DICER. These results reveal that, like for TRBP and PACT, the helicase domain of DICER is

required for DICER-PKR/p-PKR interaction during SINV infection.

In order to confirm these co-IP experiments, we next decided to perform BiFC

experiments using the same conditions as previously. In both mock and SINV-infected cells,

only the combinations of DICER WT-PKR and DICER ΔdsRBD-PKR showed a strong Venus

signal, while neither DICER N1 nor N3 constructs revealed an interaction with PKR (Fig. 5D).

In contrast, the DICER Hel. construct did not seem to interact with PKR in mock-infected cells

but appeared to do so in SINV-infected cells as a faint Venus signal could be observed. These

results therefore confirmed the co-IP observations for the DICER-PKR interaction. In addition,

we also performed a BiFC experiment using the different DICER constructs with TRBP or

PACT. Altogether, the BiFC results mostly fitted with the co-IP experiment for the DICER-

TRBP (Fig. S5A) and DICER-PACT (Fig. S5B) interactions. TRBP indeed did not seem to

interact with the DICER N1 and only slightly with the DICER N3. However, PACT interaction

was lost with DICER N1, but not with DICER N3 in mock- and SINV-infected cells (Fig. S5B

third panel). This result may be explained by the fact that DICER interacts with PACT *via* the

helicase and DUF domains, whereas only the DICER helicase domain is required for its

interaction with TRBP (Daniels et al., 2009; Lee et al., 2006). In agreement, the Venus signal observed between the DICER Hel. and PACT seemed weaker than the one we observed with TRBP (Fig. S5A and B fourth panels).

Taken together these results indicate that DICER interacts with both PKR and its phosphorylated form during SINV infection and that this interaction requires the helicase domain of DICER.

Functional importance of DICER helicase domain during SINV infection

We finally sought to study the functional role of DICER-PKR interaction during viral infection. For this purpose, we decided to use DICER helicase deletion mutants to study SINV infection. To do so, we first generated NoDice HEK293T cells stably expressing FHA-tagged DICER N1 (FHA:DICER N1) by lentiviral transduction. As for the FHA:DICER WT cell line, we first selected a clone expressing the tagged DICER N1 at a level similar level to the endogenous DICER protein in HEK293T cells (Fig. 6A). DICER N1 protein has been shown to still be able to produce miRNAs (Kennedy et al., 2015). We thus verified by northern blot analysis that DICER N1 is indeed able to process miRNAs similarly to WT DICER in HEK 293T and FHA:DICER cells, thereby validating the functionality of the tagged protein (Fig. 6B). We next infected HEK293T, FHA:DICER WT #4 and FHA:DICER N1 #6 cells with SINV-GFP and measured virus accumulation by assessing GFP expression by microscopy analysis. Interestingly, the GFP protein level was drastically reduced in FHA:DICER N1 #6 cells compared to FHA:DICER WT #4 and HEK293T cells (Fig. 6C). Encouraged by this observation, we decided to infect with SINV-GFP additional DICER deletion mutants, namely N3 and Hel. We generated stable cell lines for these various mutants by lentiviral transduction in the NoDice 2.20 background and infected those cells with SINV-GFP at an MOI of 0.02 for 24h. We verified by western blot analysis that all selected DICER mutant clones, namely N1

#6, N3 #2.13 and Hel. #2.6, expressed the tagged protein at the expected size and at levels mostly similar to the FHA:DICER WT #4 line (Fig. 6D, first two panels). We also verified the DICER mutants contribution to the endogenous miRNA biogenesis by performing a northern blot analysis on miR-16 accumulation (Fig. S6A).

We additionally verified the impact of these DICER mutants on SINV-GFP infection by measuring the GFP intensity of fluorescence by microscopy (Fig. S6B). Our results indicated that GFP accumulation is similar in HEK293T, NoDice 2.20, FHA:DICER WT, Hel. and ctrl cells. However, almost no fluorescence was detected in FHA:DICER N1 #6 and N3 #2.13 cells compared to HEK293T cells (Fig. S6B). The reduction of virus-encoded GFP accumulation and viral production were confirmed by western blot (Fig. 6D) and by plaque assay, respectively (Fig. 6E).

Altogether, these results therefore indicate that expressing a helicase truncated version of DICER, which is unable to interact with PKR, appears to confer an antiviral phenotype to SINV infection.

Discussion

The role of DICER in antiviral defense in human cells remains a topic of intense discussion (Cullen et al., 2013; Maillard et al., 2013, 2019; Cullen, 2017). In particular there have been contradictory reports regarding its capacity to produce siRNAs from viral RNAs (Bogerd et al., 2014a; Parameswaran et al., 2010; Donaszi-Ivanov et al., 2013; Backes et al., 2014). These observations could be due to the fact that several mammalian viruses potentially encode VSR proteins, thereby masking the effect of RNAi (Li et al., 2013, 2016; Maillard et al., 2013; Qiu et al., 2017, 2020). Another putative but non-exclusive explanation could be that there is a mutual regulation of type I IFN and RNAi pathways (Berkhout, 2018; Takahashi and Ui-Tei, 2020). Thus, it has already been shown that PACT can regulate MDA5 and RIG-I during virus

infection and therefore the induction of type I IFN response (Kok et al., 2011; Lui et al., 2017).

To date, it is not clear whether the activity of the DICER protein as well could be regulated by

potential interactors, or inversely whether it could itself modulate the activity of proteins

involved in the IFN pathway. To answer this question, we determined the changes in the

interactome of human DICER upon SINV and SFV infection. This analysis allowed us to reveal

that a lot of proteins associating with DICER during viral infection are dsRNA-binding proteins

and RNA helicases. A number of these proteins are known to be involved in antiviral defense

pathways, thereby indicating the possible formation of one or several complexes between

DICER and these proteins, which are very likely brought together by the accumulation of

dsRNA during virus infection.

Among these proteins, we chose to focus on the well-known ISG PKR, which is

involved in many cellular pathways such as apoptosis, cellular differentiation, development and

antiviral defense (Clemens, 1997; Gal-Ben-Ari et al., 2018; Kim et al., 2014; Lemaire et al.,

2008). PKR is one of the main actors of the Integrative Stress Response (ISR) in human cells,

and its activation or inhibition needs to be tightly regulated in order to have a properly balanced

response to stress. Our results indicate that DICER interacts via its helicase domain with PKR

in the cytoplasm during SINV infection. The helicase domain of DICER, which is also required

for its interaction with TRBP and PACT, belongs to the helicase superfamily 2, which is also

found in RLRs such as RIG-I, MDA5 or LGP2 (Ahmad and Hur, 2015; Fairman-Williams et

al., 2010). These proteins act as sensors of viral infection and through the activation of proteins

such as MAVS, mediate the induction of type I IFN pathway (Ahmad and Hur, 2015). We

hypothesize that even though the human DICER helicase has evolved mainly to act in

miRNA/siRNA pathway, it still retained the capacity to act as an RLR. However, as opposed

to RIG-I and MDA5, our data suggest that DICER would act more as an inhibitor rather than

inducer of the immune response. Therefore, we propose that this domain serves as a platform

for the recruitment of different proteins to diversify the function of DICER.

433   One such regulatory effect appears to be on the antiviral activity of PKR, as cells

expressing a truncated form of DICER unable to interact with PKR become resistant to SINV

infection. This is in agreement with previous observations that ectopic expression of the

Drosophila DICER2 protein in human cells perturbs IFN signaling pathways and antagonizes

PKR-mediated antiviral immunity (Girardi et al., 2015). The exact mechanism involved is still

unclear and will require further work, but it seems that the two proteins are likely brought

together *via* their interaction with RNA, most probably of viral origin. Indeed, we showed that

the co-IP interaction was partially RNase sensitive. However, we confirmed that the interaction

is not artificially created during the co-immunoprecipitation procedure, since we could show

that DICER and PKR interact in BiFC assay, a technique that favors the detection of direct

interactions (Lepur et al., 2016). Most of the time, the inhibition of PKR activity relies on its

inhibition to bind to dsRNA or to auto-phosphorylate. For example, the human tRNA-

dihydrouridine synthase 2 (hDus2) binds the first dsRBD of PKR and prevents its activation

(Mittelstadt et al., 2008). TRBP binds dsRNAs but also PKR directly hindering its dimerization.

In normal condition, TRBP is also associated with PACT thus preventing PKR activation by

PACT (Park et al., 1994; Nakamura et al., 2015; Daher et al., 2009; Singh et al., 2011). Since

we showed that DICER can bind the activated phospho-PKR, we hypothesize that this

interaction does not result in the inhibition of PKR autophosphorylation. In fact, in condition

of infection with a high virus dose, we showed that phospho-PKR levels are similar in cells

expressing DICER WT or helicase deletion mutants N1 and N3, but the activated PKR does

not associate with these truncated versions of DICER. Therefore, one possibility could be that

DICER interaction with PKR prevents the latter from acting upon some of its targets, which

remain to be identified, to fine-tune the antiviral response.

As of now, we cannot formally rule out that the effect of DICER on PKR is mediated

by other proteins. TRBP and PACT have been shown to regulate PKR activity, the former

normally acting as a repressor and the latter as an activator (Nakamura et al., 2015; Patel et al.,

2000; Singh et al., 2011). Interestingly, in lymphocytic Jurkat cells infected by HIV-1, PACT

can also act as a repressor of PKR (Clerzius et al., 2013). It is thus tempting to speculate that

these two proteins participate in the formation of the DICER-PKR complex. However, our

results show that this may not necessarily be the case. Indeed, in the BiFC experiment, the

DICER N3 mutant still interacted with PACT but not with PKR indicating that PACT binding

is not sufficient to confer the association with PKR.

Besides PKR, other proteins were specifically enriched upon viral infection in the

DICER IP. These are also interesting candidates to explain the putative regulatory role of

DICER. Among these proteins, DHX9 and ADAR-1 are especially intriguing. DHX9, also

known as RNA helicase A (RHA), associates with RISC, helping the RISC loading (Robb and

Rana, 2007). Moreover, DHX9 is directly involved in removing toxic dsRNAs from the cell to

prevent their processing by DICER (Aktaş et al., 2017). It has also been implicated in HIV-1

replication and knockdown of DXH9 leads to the production of less infectious HIV-1 virions

(Li et al., 1999; Fujii et al., 2001; Roy et al., 2006). Finally, DXH9 interacts with and is

phosphorylated by PKR in mouse embryonic fibroblasts. This phosphorylation precludes the

association of DHX9 with RNA, thus inhibiting its proviral effect (Sadler et al., 2009). In light

of these observations and ours, we can speculate that the inhibitory effect of DICER on PKR

activity could also be linked to DHX9 phosphorylation. ADAR-1 is one of the well-known

RNA-editing factors (Herbert et al., 1997). ADAR-1 is linked to both miRNA biogenesis (Iizasa

et al., 2010; Ota et al., 2013; Yang et al., 2006) and virus infection. Indeed, ADAR-1 has an

antiviral effect against Influenza virus, but most of the time, its depletion leads to a decrease of

the viral titer, as was reported for VSV or HIV-1 (Li et al., 2010; Samuel, 2011). It has been

shown that ADAR-1 and PKR interact directly during HIV-1 infection. This interaction triggers

the inhibition of PKR activation, and thus a reduction of eIF2α phosphorylation leading to an

increase of virus replication (Pfaller et al., 2011; Clerzius et al., 2009). Interestingly, over-

expression of ADAR-1 enhances drastically the replication of the alphaviruses Chikungunya

virus (CHIKV), and Venezuelan equine encephalitis virus (VEEV) most likely by interfering

with the IFN induction (Schoggins et al., 2011).

Finally, we cannot formally rule out that the virus resistance phenotype of the DICER

N1 and N3 cell lines is due to an increased processivity of these truncated proteins on long

dsRNA substrates (Kennedy et al., 2015), which would render DICER RNAi proficient.

However, PKR is still expressed in these cells, and is therefore expected to be predominant over

DICER for viral dsRNA sensing. It is nonetheless possible that its interaction with several

dsRNA binding proteins could also participate in limiting the efficiency of the helicase domain

of DICER, on top of its intrinsic lack of processivity (Chakravarthy et al., 2010).

Deciphering the exact role of human DICER protein during virus infection is a

challenging task and additional studies will be required to get a global picture. Nevertheless, by

assessing the interactome of this protein during SINV infection, we have unveiled a new role

for the helicase domain of DICER in regulating the cellular response to viral infection.

499 Material and methods

500 Plasmids, cloning and mutagenesis

Plasmids used for BiFC experiments were a gift from Dr. Oliver Vugrek from the Ruđer

Bošković Institute and described in (Lepur et al., 2016). The cDNAs of TRBP, PACT and PKR

were respectively amplified from (pcDNA-TRBP Addgene #15666) (Kok et al., 2007),

(pcDNA-PACT Addgene #15667) (Kok et al., 2007), (pSB819-PKR-hum Addgene #20030)

(Elde et al., 2009), and cloned into the four pBiFC vectors by Gateway recombination. DICER

N1, N3 Hel. and ΔdsRBD were generated by PCR mutagenesis from pDONR-DICER described

in (Girardi et al., 2015) and cloned into the four pBiFC and pDEST-FHA vectors by Gateway

recombination. plenti6 FHA-V5 vector was modified from plenti6-V5 gateway vector (Thermo

Fisher scientific V49610) by Gibson cloning. DICER WT, N1 from pDONOR plasmids were

cloned into plenti6 FHA-V5 by Gateway recombination. All primers used are listed in Supp.

Table 5.

Cell lines

HEK293T, HEK293T/NoDice (2.20 and 4.25), and HEK293T/NoDiceΔPKR cell lines were a

gift from Pr. Bryan Cullen and described in (Bogerd et al., 2014b; Kennedy et al., 2015).

HCT116 cell line was a gift from Dr. Christian Gaiddon.

Generation of Flag-HA-GFP-DICER knock-in cell line by CRISPR/Cas9

To generate the knock-in cell line, the sequence of Flag-HA-GFP was amplified by PCR from

the Flag-HA-GFP plasmid (Meister et al., 2005). DNA sequences corresponding to 1 Kb

upstream (left homology arm) and downstream (right homology arm) the starting codon (ATG)

of DICER gene were amplified from HCT116 cell genomic DNA using primer pairs listed in

Supp. Table 5. The three PCR products were gel-purified and cloned into a linearized pUC19

by In-fusion cloning (Clontech) to obtain the template for homologous recombination

(LarmDICER-FlagHAGFP-RarmDICER).

Design of the guide RNA targeting the region between Dicer 5'-UTR and its first coding exon

for CRISPR/Cas9 mediated knock-in was carried out using the CRISPOR Design Tool

(Concordet and Haeussler, 2018). Annealed oligonucleotide corresponding to the gRNA (Supp.

Table 5) were cloned into the vector pX459 (Addgene #48139) which also encodes *S. pyogenes*

Cas9 with 2A-Puro.

The sequence of the donor plasmid was additionally mutagenized to disrupt the PAM sequence

of the right homology arm to avoid its cleavage by the gRNA.

To obtain the knock-in (KI) cell line, 5×10^5 HCT116 cells were seeded in a 6 well plate with

Dulbecco's modified Eagle medium (DMEM, Gibco®, Life Technologies) supplemented with

10% fetal bovine serum (FBS, Clontech) in a humidified atmosphere of 5% CO_2 at 37°C and

transfected after 24 hours with the pX459-gRNADicerNterm-Cas9-2A-Puro plasmid and the

Leftarm-FlagHAGFP-RightarmDICER donor plasmids at the ratio of 1 to 1 (6 micrograms

plasmids in total) using Lipofectamine 2000 according to the manufacturer instructions. 24

hours later, puromycin (1mg/mL) was added to the cells to increase the KI efficiency and

genomic DNA was isolated from individual colonies few days later.

The presence of the Flag-HA-GFP tag in frame with hDICER coding sequence was confirmed

by sequencing PCR amplicon from KI cell gDNA. Expression of Flag-HA-GFP N-terminal

tagged Dicer protein in the KI cells was confirmed by western blot.

Cell culture and transfection

Cells were maintained in Dulbecco's modified Eagle medium (DMEM, Gibco®, Life

Technologies) supplemented with 10% fetal bovine serum (FBS, Clontech) in a humidified

atmosphere of 5% CO_2 at 37°C. Transfection was performed using Lipofectamine 2000

(Invitrogen, ThermoFisher Scientific) according to the manufacturer's instructions.

Lentivirus production and generation of stable cell lines

The lentiviral supernatant from single transfer vector was produced by transfecting HEK293T

cells (ATCC® CRL-3216™) with 20 µg of the transfer vector, 15 µg of pMDLg/p RRE and 10

μg of pRSV-Rev packaging plasmids (Addgene #12251 and Addgene #12253) and the pVSV

envelope plasmid (Addgene #8454) using Lipofectamine 2000 (Invitrogen, ThermoFisher

Scientific) reagent according to the manufacturer's protocol. Standard DMEM (Gibco®, Life

Technologies) medium supplemented with 10% Fetal bovine serum (FBS, Gibco®, Life

Technologies) and 100 U/mL of penicillin-Streptomycin (Gibco®, Life Technologies) was

used for growing HEK293T cells and for lentivirus production. One 10 cm plate of HEK293T

cells at 70-80% confluency was used for the transfection. The medium was replaced 8 hours

post-transfection. After 48 hours the medium containing viral particles was collected and

filtered through a 0.45μm PES filter. The supernatant was directly used for transfection or

stored at -80°C. A 6 well plate of HEK293T/NoDice cells at 80% confluency was transduced

using 600 μL of lentiviral supernatant either expressing FHADICER, N1 or empty vector,

supplemented with 4 ug/mL polybrene (Sigma) for 6 hours. The transduction media was then

changed with fresh DMEM for 24 hours and the resistant cell clones were selected for about 6

weeks with blasticidin (15 μg/mL) and subsequently maintained under blasticidin selection.

Viral stocks, virus infection

Viral stocks of SINV or SINV-GFP were produced as described in (Girardi et al., 2015). Cells

were infected with SINV or SINV-GFP at an MOI of 0.02, 0.1, 1 or 2 and samples were

collected at different time points as indicated in the figure legends.

Analysis of viral titer by plaque assay

Vero R cells were seeded in 96-well plates format and were infected with 10-fold serial

dilutions infection supernatants for 1 hour. Afterwards, the inoculum was removed, and cells

were cultured in 2.5% carboxymethyl cellulose for 72 hours at 37°C in a humidified atmosphere

of 5% CO_2. Plaques were counted manually under the microscope and viral titer was calculated

according to the formula: *PFU/mL = #plaques/ (Dilution*Volume of inoculum)*

Western blot analysis

Proteins were extracted from cells and homogenized in 350 μL of lysis buffer (50 mM Tris-

HCl pH 7.5, 150 mM NaCl, 5 mM EDTA, 1% Triton X-100, 0.5% SDS and Protease Inhibitor

Cocktail (complete Mini; Sigma Aldrich)). Proteins were quantified by the Bradford method

and 20 to 30 μg of total protein extract were loaded on 4-20% Mini-PROTEAN® TGX™

Precast Gels (Bio-Rad). After transfer onto nitrocellulose membrane, equal loading was verified

by Ponceau staining. For PVDF membrane, equal loading was verified by Coomassie staining

after transfer and blotting. Membranes were blocked in 5% milk and probed with the following

antibodies: anti-hDicer (1:500, F10 Santa Cruz sc-136979) and anti-hDicer (1:1000, A301-

937A Bethyl), anti-TRBP (1:500, D-5 Santa Cruz sc-514124), anti-PKR (1:2500, Abcam

ab32506), anti-PACT (1:500, Abcam, ab75749), anti-HA (1:10000, Sigma, H9658), anti-

DHX9 (1:500, Abcam ab26271), anti-p-eIF2a (1:1000, Ser-52 Santa Cruz sc-601670), anti-

hADAR-1 (1:500 Santa Cruz sc-271854) anti-p-PKR (1:1000 Abcam ab81303) anti-GFP

(1:10000, Roche 11814460001) and anti-Tubulin (1:10000, Sigma T6557). Detection was

performed using Chemiluminescent Substrate (Pierce, ThermoFisher Scientific) and visualized

on a Fusion FX imaging system (Vilber).

RNA extraction and northern blot analysis

Total RNA was extracted using Tri-Reagent Solution (Fisher Scientific; MRC, Inc) according

to the manufacturer's instructions. Northern blotting was performed on 10 μg of total RNA.

RNA was resolved on a 12% urea-acrylamide gel, transferred onto Hybond-NX membrane (GE

Healthcare). RNAs were then chemically cross-linked to the membrane during 90 min at 65°C

using 1-ethyl-3-[3-dimethylaminopropyl]carbodiimide hydrochloride (EDC) (Sigma Aldrich).

Membranes were prehybridized for 30 min in PerfectHyb™ plus (Sigma Aldrich) at 50°C.

Probes consisting of oligodeoxyribonucleotides (see Supplementary Table 2) were 5′-end

labeled using T4 polynucleotide kinase (ThermoFisher Scientific) with 25 μCi of [γ-32P]dATP.

The labeled probe was hybridized to the blot overnight at 50°C. The blot was then washed twice

at 50°C for 20 min (5× SSC/0.1% SDS), followed by an additional wash (1× SSC/0.1% SDS)

for 5 min. Northern blots were exposed to phosphorimager plates and scanned using a

Bioimager FLA-7000 (Fuji).

Immunoprecipitation

Immunoprecipitation experiments were carried out either on tagged proteins or on endogenous

proteins.

Tagged proteins: Cells were harvested, washed twice with ice-cold 1× PBS (Gibco®, Life

Technologies), and resuspended in 550 μL of lysis buffer (50 mM Tris-HCl pH 7.5, 140 mM

NaCl, 1.5 mM $MgCl_2$, 0.1% NP-40), supplemented with Complete-EDTA-free Protease

Inhibitor Cocktail (complete Mini; Sigma Aldrich). Cells were lysed by 30 min incubation on

ice and debris were removed by 15 min centrifugation at 2000 g and 4°C. An aliquot of the

cleared lysates (50 μL) was kept aside as protein Input. Samples were divided into equal parts

(250 μL each) and incubated with 15 μL of magnetic microparticles coated with monoclonal

HA or MYC antibodies (MACS purification system, Miltenyi Biotech) at 4°C for 1 hour under

rotation (10 rpm). Samples were passed through μ Columns (MACS purification system,

Miltenyi Biotech). The μ Columns were then washed 3 times with 200 μL of lysis buffer and 1

time with 100 μL of washing buffer (20 mM Tris-HCl pH 7.5). To elute the immunoprecipitated

proteins, 95°C pre-warmed 2x Western blot loading buffer (10% glycerol, 4% SDS, 62.5 mM

Tris-HCl pH 6.8, 5% (v/v) 2-β-mercaptoethanol, Bromophenol Blue) was passed through the

μ Columns. Proteins were analyzed by western blotting or by mass spectrometry.

Endogenous proteins: The day before immunoprecipitation, magnetic DynaBeads protein G

(Invitrogen, ThermoFisher Scientific) were prepared: briefly, 160 μL of magnetic DynaBeads

protein G (Invitrogen) were washed three times with lysis buffer (50 mM Tris-HCl pH 7.5, 150

mM NaCl, 5 mM EDTA, 1% Triton X-100, 0.5% SDS). The beads were then incubated with 4

μg of BSA for 1 h under agitation at room temperature. An aliquot of beads (40 μL) was kept

for preclearing step. The remaining beads were incubated either with 6 μg of PKR antibody or

control IgG overnight at 4°C under rotation.

For the IP: Cells were harvested, washed twice with ice-cold 1× PBS (Gibco®, Life

Technologies) and resuspended in 1 mL of lysis buffer (50 mM Tris-HCl pH 7.5, 150 mM

NaCl, 5 mM EDTA, 1% Triton X-100, 0.5% SDS) supplemented with Complete-EDTA-free

Protease Inhibitor Cocktail (complete Mini; Sigma Aldrich). Cells were lysed by three 15 s

sonication steps followed by 30 min incubation on ice and debris were removed by 15 min

centrifugation at 16000 g and 4°C. After centrifugation, an aliquot was kept aside as protein

Input. Samples were incubated with beads alone for preclearing step for 1 hour at 4°C on wheel.

Samples were then put on magnetic racks and the supernatant transferred on new tubes

containing either beads coupled to PKR antibody or to control IgG. Samples were incubated on

wheel for 3 hours at 4°C. After incubation, samples were put on magnetic racks and beads were

washed three times with lysis buffer. After removal of supernatant, beads were eluted with 2x

western blot loading buffer and incubated for 10 min at 95°C under agitation. Proteins were

analyzed by western blotting.

RNase treatment followed by co-IP

On tagged proteins: Cells were harvested, washed twice with ice-cold 1× PBS (Gibco®, Life

Technologies), and resuspended in 550 μL of lysis buffer (50 mM Tris-HCl pH 7.5, 140 mM

NaCl, 1.5 mM MgCl₂, 0.1% NP-40), supplemented with Complete-EDTA-free Protease

Inhibitor Cocktail (complete Mini; Sigma Aldrich). Cells were lysed by 30 min incubation on

ice and debris were removed by 15 min centrifugation at 2000 g and 4°C. Lysate was treated

or not with RNase A/T1 mix (ThermoFisher Scientific) and place at 37°C 30min. An aliquot of

the cleared lysates (25 μL) was kept aside as protein Input and another aliquot (25 μL) was kept

to assess RNase treatment efficiency. Co-IP was led as previously described.

Total RNA was extracted using Tri-Reagent Solution (Fisher Scientific; MRC, Inc) according

to the manufacturer's instructions. RNA integrity upon treatment was verified on an 1% agarose

gel containing ethidium bromide 10 mg/mL (Invitrogen, ThermoFisher Scientific) and revealed

under UV on Gel Doc™ EZ system (Bio-Rad).

BiFC assay

Experiments were carried out in two different ways. For non-fixed cells, NoDiceΔPKR or

HEK293T cells were seeded at the density of 1.2×10^5 cells per well in a 24-well plate. After

16 hours, cells were transfected with equimolar quantities of each plasmid forming BiFC

couples. After 24 hours, cells were infected with SINV at an MOI of 2 and pictures were taken

6 hours post-infection using ZOE fluorescent cell imager (Bio-Rad). Proteins were collected

with lysis buffer (50 mM Tris-HCl pH 7.5, SDS 0.05%, Triton 1%, 5 mM EDTA, 150 mM

NaCl) supplemented with Complete-EDTA-free Protease Inhibitor Cocktail (complete Mini;

Sigma Aldrich), and subjected to western blot analysis. For fixed cells, NoDice/ΔPKR cells

were seeded at the density of 8.10^4 cells per well in 8-well Millicell® EZ Slides (Merck

Millipore), transfected and infected as described previously. At 6 hours post-infection, cells

were fixed with 4% formaldehyde and 0.2% glutaraldehyde for 10 min. Cells were then washed

with 1× PBS (Gibco®, Life Technologies) and stained with 10 μg/μL DAPI (Invitrogen,

ThermoFisher Scientific) in 1× PBS solution (Invitrogen, ThermoFisher Scientific) for 5 min.

Fixed cells were mounted on a glass slide with Fluoromount-G mounting media (Southern

Biotech). Images were acquired using confocal LSM780 (Zeiss) inverted microscope with an

argon laser (514x nm) and with ×40 immersion oil objective. All pictures obtained from BiFC

experiments were treated using FigureJ software (NIH).

Mass spectrometry analysis

Protein extracts were prepared for mass spectrometry as described in a previous study (Chicois

et al., 2018). Each sample was precipitated with 0.1 M ammonium acetate in 100% methanol,

and proteins were resuspended in 50 mM ammonium bicarbonate. After a reduction-alkylation

step (dithiothreitol 5 mM – iodoacetamide 10 mM), proteins were digested overnight with

sequencing-grade porcine trypsin (1:25, w/w, Promega, Fitchburg, MA, USA). The resulting

vacuum-dried peptides were resuspended in water containing 0.1% (v/v) formic acid (solvent

A). One sixth of the peptide mixtures were analyzed by nanoLC-MS/MS an Easy-nanoLC-1000

system coupled to a Q-Exactive Plus mass spectrometer (Thermo-Fisher Scientific, USA)

operating in positive mode. Five microliters of each sample were loaded on a C-18 precolumn

(75 μm ID × 20 mm nanoViper, 3 μm Acclaim PepMap; Thermo) coupled with the analytical

C18 analytical column (75 μm ID × 25 cm nanoViper, 3 μm Acclaim PepMap; Thermo).

Peptides were eluted with a 160 min gradient of 0.1% formic acid in acetonitrile at 300 nL/min.

The Q-Exactive Plus was operated in data-dependent acquisition mode (DDA) with Xcalibur

software (Thermo-Fisher Scientific). Survey MS scans were acquired at a resolution of 70K at

200 m/z (mass range 350-1250), with a maximum injection time of 20 ms and an automatic

gain control (AGC) set to 3e6. Up to 10 of the most intense multiply charged ions (≥2) were

selected for fragmentation with a maximum injection time of 100 ms, an AGC set at 1e5 and a

resolution of 17.5K. A dynamic exclusion time of 20 s was applied during the peak selection

process.

Database search and mass-spectrometry data post-processing

Data were searched against a database containing Human and Viruses UniProtKB sequences

with a decoy strategy (GFP, Human and Sindbis Virus SwissProt sequences as well as Semliki

Forest Virus SwissProt and TrEMBL sequences (releases from January 2017, 40439

sequences)). Peptides were identified with Mascot algorithm (version 2.3, Matrix Science,

London, UK) with the following search parameters: carbamidomethylation of cysteine was set

as fixed modification; N-terminal protein acetylation, phosphorylation of serine / threonine /

tyrosine and oxidation of methionine were set as variable modifications; tryptic specificity with

up to three missed cleavages was used. The mass tolerances in MS and MS/MS were set to 10

ppm and 0.02 Da respectively, and the instrument configuration was specified as "ESI-Trap".

The resulting .dat Mascot files were then imported into Proline v1.4 package

(http://proline.profiproteomics.fr/) for post-processing. Proteins were validated with Mascot

pretty rank equal to 1, 1% FDR on both peptide spectrum matches (PSM) and protein sets

(based on score). The total number of MS/MS fragmentation spectra (Spectral count or SpC)

was used for subsequent protein quantification in the different samples.

Exploratory and differential expression analysis of LC-MS/MS data

Mass spectrometry data obtained for each sample were stored in a local MongoDB database

and subsequently analyzed through a Shiny Application built upon the R/Bioconductor

packages msmsEDA (Gregori J, Sanchez A, Villanueva J (2014). msmsEDA: Exploratory Data

Analysis of LC-MS/MS data by spectral counts. R/Bioconductor package version 1.22.0) and

msmsTests (Gregori J, Sanchez A, Villanueva J (2013). msmsTests: LC-MS/MS Differential

Expression Tests. R/Bioconductor package version 1.22.0). Exploratory data analyses of LC-

MS/MS data were thus conducted, and differential expression tests were performed using a

negative binomial regression model. The p-values were adjusted with FDR control by the

Benjamini-Hochberg method and the following criteria were used to define differentially

expressed proteins: an adjusted p-value < 0.05, a minimum of 5 SpC in the most abundant

condition, and a minimum fold change of 2 (abs(LogFC) > 1).

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange

Consortium via the PRIDE (Perez-Riverol et al., 2019) partner repository with the dataset

identifier PXD019093 and 10.6019/PXD019093.

Acknowledgments

The authors would like to thank members of the Pfeffer laboratory for discussion, Pr. Bryan

Cullen for the kind gift of the HEK293T NoDice and NoDice/ΔPKR cell lines and Dr. Oliver

Vugrek for the BiFC plasmids.

This work was funded by the European Research Council (ERC-CoG-647455 RegulRNA) and

was performed under the framework of the LABEX: ANR-10-LABX-0036_NETRNA and

ANR-17-EURE-0023, which benefits from a funding from the state managed by the French

National Research Agency as part of the Investments for the future program. This work has

also received funding from the People Programme (Marie Curie Actions) of the European

Union's Seventh Framework Program (FP7/2007-2013) under REA grant agreement n°

PCOFUND-GA-2013-609102, through the PRESTIGE program coordinated by Campus

France (to EG), and from the French Minister for Higher Education, Research and Innovation

(PhD contract to MB). The mass spectrometry instrumentation was funded by the University

of Strasbourg, IdEx "Equipement mi-lourd" 2015.

Author contributions

754   SP and ML conceived the project. TCM, ML, EG and SP designed the work. TCM, MB, ML,

755   EG and MM performed the experiments and analyzed the results. PH and JC performed the

756   mass spectrometry analysis. BCWM designed software and analyzed the mass spectrometry

757   results. SP coordinated the work and assured funding. TCM, MB and SP wrote the manuscript

758   with input from the other authors. All authors reviewed and approved the final manuscript.

References

Adiliaghdam, F., Basavappa, M., Saunders, T.L., Harjanto, D., Prior, J.T., Cronkite, D.A., Papavasiliou, N., and Jeffrey, K.L. (2020). A Requirement for Argonaute 4 in Mammalian Antiviral Defense. Cell Rep. *30*, 1690-1701.e4.

Ahmad, S., and Hur, S. (2015). Helicases in Antiviral Immunity: Dual Properties as Sensors and Effectors. Trends Biochem. Sci. *40*, 576–585.

Aktaş, T., Avşar Ilık, İ., Maticzka, D., Bhardwaj, V., Pessoa Rodrigues, C., Mittler, G., Manke, T., Backofen, R., and Akhtar, A. (2017). DHX9 suppresses RNA processing defects originating from the Alu invasion of the human genome. Nature *544*, 115–119.

Backes, S., Langlois, R.A., Schmid, S., Varble, A., Shim, J.V., Sachs, D., and tenOever, B.R. (2014). The Mammalian Response to Virus Infection Is Independent of Small RNA Silencing. Cell Rep. *8*, 114–125.

Bartel, D.P. (2018). Metazoan MicroRNAs. Cell *173*, 20–51.

Berkhout, B. (2018). RNAi-mediated antiviral immunity in mammals. Curr. Opin. Virol. *32*, 9–14.

Bernstein, E., Caudy, A.A., Hammond, S.M., and Hannon, G.J. (2001). Role for a bidentate ribonuclease in the initiation step of RNA interference. Nature *409*, 363–366.

Bogerd, H.P., Skalsky, R.L., Kennedy, E.M., Furuse, Y., Whisnant, A.W., Flores, O., Schultz, K.L.W., Putnam, N., Barrows, N.J., Sherry, B., et al. (2014a). Replication of many human viruses is refractory to inhibition by endogenous cellular microRNAs. J. Virol. *88*, 8065–8076.

Bogerd, H.P., Whisnant, A.W., Kennedy, E.M., Flores, O., and Cullen, B.R. (2014b). Derivation and characterization of Dicer- and microRNA-deficient human cells. RNA *20*, 923–937.

Borden, E.C., Sen, G.C., Uze, G., Silverman, R.H., Ransohoff, R.M., Foster, G.R., and Stark, G.R. (2007). Interferons at age 50: past, current and future impact on biomedicine. Nat. Rev. Drug Discov. *6*, 975–990.

Chakravarthy, S., Sternberg, S.H., Kellenberger, C.A., and Doudna, J.A. (2010). Substrate-Specific Kinetics of Dicer-Catalyzed RNA Processing. J. Mol. Biol. *404*, 392–402.

Chen, E.Y., Tan, C.M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G.V., Clark, N.R., and Ma'ayan, A. (2013). Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. BMC Bioinformatics *14*, 128.

Chendrimada, T.P., Gregory, R.I., Kumaraswamy, E., Norman, J., Cooch, N., Nishikura, K., and Shiekhattar, R. (2005). TRBP recruits the Dicer complex to Ago2 for microRNA processing and gene silencing. Nature *436*, 740–744.

Chicois, C., Scheer, H., Garcia, S., Zuber, H., Mutterer, J., Chicher, J., Hammann, P., Gagliardi, D., and Garcia, D. (2018). The UPF1 interactome reveals interaction networks between RNA degradation and translation repression factors in Arabidopsis. Plant J. *96*, 119–132.

Clemens, M.J. (1997). PKR-a protein kinase regulated by double-stranded RNA. Int J Biochem Cell Biol *29*, 945–949.

Clerzius, G., Gélinas, J.-F., Daher, A., Bonnet, M., Meurs, E.F., and Gatignol, A. (2009). ADAR1 interacts with PKR during human immunodeficiency virus infection of lymphocytes and contributes to viral replication. J. Virol. *83*, 10119–10128.

Clerzius, G., Shaw, E., Daher, A., Burugu, S., Gélinas, J.-F., Ear, T., Sinck, L., Routy, J.-P., Mouland, A.J., Patel, R.C., et al. (2013). The PKR activator, PACT, becomes a PKR inhibitor during HIV-1 replication. Retrovirology *10*, 96.

Concordet, J.-P., and Haeussler, M. (2018). CRISPOR: intuitive guide selection for CRISPR/Cas9 genome editing experiments and screens. Nucleic Acids Res. *46*, W242–W245.

Cullen, B.R. (2006). Is RNA interference involved in intrinsic antiviral immunity in mammals?
Nat Immunol 7, 563–567.
Cullen, B.R. (2017). RNA Interference in Mammals: The Virus Strikes Back. Immunity 46,
970–972.
Cullen, B.R., Cherry, S., and tenOever, B.R. (2013). Is RNA Interference a Physiologically
Relevant Innate Antiviral Immune Response in Mammals? Cell Host Microbe 14, 374–
378.
Daher, A., Laraki, G., Singh, M., Melendez-Peña, C.E., Bannwarth, S., Peters, A.H.F.M.,
Meurs, E.F., Braun, R.E., Patel, R.C., and Gatignol, A. (2009). TRBP control of PACT-
induced phosphorylation of protein kinase R is reversed by stress. Mol. Cell. Biol. 29, 254–
265.
Daniels, S.M., Melendez-Peña, C.E., Scarborough, R.J., Daher, A., Christensen, H.S., El Far,
M., Purcell, D.F.J., Lainé, S., and Gatignol, A. (2009). Characterization of the TRBP
domain required for dicer interaction and function in RNA interference. BMC Mol. Biol.
10, 38.
Donaszi-Ivanov, A., Mohorianu, I., Dalmay, T., and Powell, P.P. (2013). Small RNA Analysis
in Sindbis Virus Infected Human HEK293 Cells. PLoS ONE 8, e84070.
Doyle, M., Badertscher, L., Jaskiewicz, L., Güttinger, S., Jurado, S., Hugenschmidt, T., Kutay,
U., and Filipowicz, W. (2013). The double-stranded RNA binding domain of human Dicer
functions as a nuclear localization signal. RNA N. Y. N 19, 1238–1252.
Elde, N.C., Child, S.J., Geballe, A.P., and Malik, H.S. (2009). Protein kinase R reveals an
evolutionary model for defeating viral mimicry. Nature 457, 485–489.
Fairman-Williams, M.E., Guenther, U.-P., and Jankowsky, E. (2010). SF1 and SF2 helicases:
family matters. Curr. Opin. Struct. Biol. 20, 313–324.
Fros, J.J., and Pijlman, G.P. (2016). Alphavirus Infection: Host Cell Shut-Off and Inhibition of
Antiviral Responses. Viruses 8, 166.
Fujii, R., Okamoto, M., Aratani, S., Oishi, T., Ohshima, T., Taira, K., Baba, M., Fukamizu, A.,
and Nakajima, T. (2001). A Role of RNA Helicase A in cis-Acting Transactivation
Response Element-mediated Transcriptional Regulation of Human Immunodeficiency
Virus Type 1. J. Biol. Chem. 276, 5445–5451.
Gal-Ben-Ari, S., Barrera, I., Ehrlich, M., and Rosenblum, K. (2018). PKR: A Kinase to
Remember. Front. Mol. Neurosci. 11, 480.
García, M.A., Meurs, E.F., and Esteban, M. (2007). The dsRNA protein kinase PKR: Virus and
cell control. Biochimie 89, 799–811.
Girardi, E., Chane-Woon-Ming, B., Messmer, M., Kaukinen, P., and Pfeffer, S. (2013).
Identification of RNase L-dependent, 3'-end-modified, viral small RNAs in Sindbis virus-
infected mammalian cells. MBio 4, e00698-00613.
Girardi, E., Lefèvre, M., Chane-Woon-Ming, B., Paro, S., Claydon, B., Imler, J.-L., Meignin,
C., and Pfeffer, S. (2015). Cross-species comparative analysis of Dicer proteins during
Sindbis virus infection. Sci. Rep. 5, 10693.
Griffin, D. (2007). Alphaviruses, p 1023–1067. Fields Virol. 5th Ed Lippincott Williams
Wilkins Phila. PA.
Guo, Z., Li, Y., and Ding, S.-W. (2019). Small RNA-based antimicrobial immunity. Nat. Rev.
Immunol. 19, 31–44.
Haase, A.D., Jaskiewicz, L., Zhang, H., Laine, S., Sack, R., Gatignol, A., and Filipowicz, W.
(2005). TRBP, a regulator of cellular PKR and HIV-1 virus expression, interacts with Dicer
and functions in RNA silencing. EMBO Rep 6, 961–967.
Herbert, A., Alfken, J., Kim, Y.G., Mian, I.S., Nishikura, K., and Rich, A. (1997). A Z-DNA
binding domain present in the human editing enzyme, double-stranded RNA adenosine
deaminase. Proc. Natl. Acad. Sci. U. S. A. 94, 8421–8426.

Heyam, A., Lagos, D., and Plevin, M. (2015). Dissecting the roles of TRBP and PACT in
double-stranded RNA recognition and processing of noncoding RNAs. WIREs RNA *6*,
271–289.
Hutvágner, G., McLachlan, J., Pasquinelli, A.E., Balint, E., Tuschl, T., and Zamore, P.D.
(2001). A cellular function for the RNA-interference enzyme Dicer in the maturation of
the let-7 small temporal RNA. Science *293*, 834–838.
Iizasa, H., Wulff, B.-E., Alla, N.R., Maragkakis, M., Megraw, M., Hatzigeorgiou, A., Iwakiri,
D., Takada, K., Wiedmer, A., Showe, L., et al. (2010). Editing of Epstein-Barr virus-
encoded BART6 microRNAs controls their dicer targeting and consequently affects viral
latency. J. Biol. Chem. *285*, 33358–33370.
Ivashkiv, L.B., and Donlin, L.T. (2014). Regulation of type I interferon responses. Nat. Rev.
Immunol. *14*, 36–49.
Kennedy, E.M., Whisnant, A.W., Kornepati, A.V.R., Marshall, J.B., Bogerd, H.P., and Cullen,
B.R. (2015). Production of functional small interfering RNAs by an amino-terminal
deletion mutant of human Dicer. Proc. Natl. Acad. Sci. *112*, E6945–E6954.
Kim, Y., Lee, J.H., Park, J.-E., Cho, J., Yi, H., and Kim, V.N. (2014). PKR is activated by
cellular dsRNAs during mitosis and acts as a mitotic regulator. Genes Dev. *28*, 1310–1322.
Kok, K.H., Ng, M.H., Ching, Y.P., and Jin, D.Y. (2007). Human TRBP and PACT Directly
Interact with Each Other and Associate with Dicer to Facilitate the Production of Small
Interfering RNA. J Biol Chem *282*, 17649–17657.
Kok, K.-H., Lui, P.-Y., Ng, M.-H.J., Siu, K.-L., Au, S.W.N., and Jin, D.-Y. (2011). The Double-
Stranded RNA-Binding Protein PACT Functions as a Cellular Activator of RIG-I to
Facilitate Innate Antiviral Response. Cell Host Microbe *9*, 299–309.
Kuleshov, M.V., Jones, M.R., Rouillard, A.D., Fernandez, N.F., Duan, Q., Wang, Z., Koplev,
S., Jenkins, S.L., Jagodnik, K.M., Lachmann, A., et al. (2016). Enrichr: a comprehensive
gene set enrichment analysis web server 2016 update. Nucleic Acids Res. *44*, W90-97.
Lambert, M., Pépin, G., Peralta-Zaragoza, O., Matusiak, R., Ly, S., Landry, P., and Provost, P.
(2016). TWEAK Negatively Regulates Human Dicer. Non-Coding RNA *2*.
Lee, Y., Hur, I., Park, S.Y., Kim, Y.K., Suh, M.R., and Kim, V.N. (2006). The role of PACT
in the RNA silencing pathway. Embo J *25*, 522–532.
Lemaire, P.A., Anderson, E., Lary, J., and Cole, J.L. (2008). Mechanism of PKR Activation by
dsRNA. J. Mol. Biol. *381*, 351–360.
Lepur, A., Kovačević, L., Belužić, R., and Vugrek, O. (2016). Combining unique multiplex
gateway cloning and bimolecular fluorescence complementation (BiFC) for high-
throughput screening of protein–protein interactions. J. Biomol. Screen. *21*, 1100–1111.
Li, J., Tang, H., Mullen, T.M., Westberg, C., Reddy, T.R., Rose, D.W., and Wong-Staal, F.
(1999). A role for RNA helicase A in post-transcriptional regulation of HIV type 1. Proc.
Natl. Acad. Sci. U. S. A. *96*, 709–714.
Li, Y., Lu, J., Han, Y., Fan, X., and Ding, S.-W. (2013). RNA interference functions as an
antiviral immunity mechanism in mammals. Science *342*, 231–234.
Li, Y., Basavappa, M., Lu, J., Dong, S., Cronkite, D.A., Prior, J.T., Reinecker, H.-C., Hertzog,
P., Han, Y., Li, W.-X., et al. (2016). Induction and suppression of antiviral RNA
interference by influenza A virus in mammalian cells. Nat. Microbiol. *2*, 16250.
Li, Z., Wolff, K.C., and Samuel, C.E. (2010). RNA adenosine deaminase ADAR1 deficiency
leads to increased activation of protein kinase PKR and reduced vesicular stomatitis virus
growth following interferon treatment. Virology *396*, 316–322.
Lui, P.-Y., Wong, L.-Y.R., Ho, T.-H., Au, S.W.N., Chan, C.-P., Kok, K.-H., and Jin, D.-Y.
(2017). PACT Facilitates RNA-Induced Activation of MDA5 by Promoting MDA5
Oligomerization. J. Immunol. *199*, 1846–1855.

Maillard, P.V., Ciaudo, C., Marchais, A., Li, Y., Jay, F., Ding, S.W., and Voinnet, O. (2013).
Antiviral RNA interference in mammalian cells. Science *342*, 235–238.

Maillard, P.V., Veen, A.G.V. der, Deddouche-Grass, S., Rogers, N.C., Merits, A., and Sousa,
C.R. e (2016). Inactivation of the type I interferon pathway reveals long double-stranded
RNA-mediated RNA interference in mammalian cells. EMBO J. *35*, 2505–2518.

Maillard, P.V., van der Veen, A.G., Poirier, E.Z., and Reis e Sousa, C. (2019). Slicing and
dicing viruses: antiviral RNA interference in mammals. EMBO J. *38*.

Meister, G., and Tuschl, T. (2004). Mechanisms of gene silencing by double-stranded RNA.
Nature *431*, 343–349.

Meister, G., Landthaler, M., Peters, L., Chen, P.Y., Urlaub, H., Lührmann, R., and Tuschl, T.
(2005). Identification of novel argonaute-associated proteins. Curr. Biol. CB *15*, 2149–
2155.

Mittelstadt, M., Frump, A., Khuu, T., Fowlkes, V., Handy, I., Patel, C.V., and Patel, R.C.
(2008). Interaction of human tRNA-dihydrouridine synthase-2 with interferon-induced
protein kinase PKR. Nucleic Acids Res. *36*, 998–1008.

Nakamura, T., Kunz, R.C., Zhang, C., Kimura, T., Yuan, C.L., Baccaro, B., Namiki, Y., Gygi,
S.P., and Hotamisligil, G.S. (2015). A Critical Role for PKR Complexes with TRBP in
Immunometabolic Regulation and eIF2α Phosphorylation in Obesity. Cell Rep. *11*, 295–
307.

Ota, H., Sakurai, M., Gupta, R., Valente, L., Wulff, B.-E., Ariyoshi, K., Iizasa, H., Davuluri,
R.V., and Nishikura, K. (2013). ADAR1 forms a complex with Dicer to promote
microRNA processing and RNA-induced gene silencing. Cell *153*, 575–589.

Parameswaran, P., Sklan, E., Wilkins, C., Burgon, T., Samuel, M.A., Lu, R., Ansel, K.M.,
Heissmeyer, V., Einav, S., Jackson, W., et al. (2010). Six RNA viruses and forty-one hosts:
viral small RNAs and modulation of small RNA repertoires in vertebrate and invertebrate
systems. PLoS Pathog *6*, e1000764.

Park, H., Davies, M.V., Langland, J.O., Chang, H.W., Nam, Y.S., Tartaglia, J., Paoletti, E.,
Jacobs, B.L., Kaufman, R.J., and Venkatesan, S. (1994). TAR RNA-binding protein is an
inhibitor of the interferon-induced protein kinase PKR. Proc. Natl. Acad. Sci. U. S. A. *91*,
4713–4717.

Patel, C.V., Handy, I., Goldsmith, T., and Patel, R.C. (2000). PACT, a Stress-modulated
Cellular Activator of Interferon-induced Double-stranded RNA-activated Protein Kinase,
PKR. J. Biol. Chem. *275*, 37993–37998.

Perez-Riverol, Y., Csordas, A., Bai, J., Bernal-Llinares, M., Hewapathirana, S., Kundu, D.J.,
Inuganti, A., Griss, J., Mayer, G., Eisenacher, M., et al. (2019). The PRIDE database and
related tools and resources in 2019: improving support for quantification data. Nucleic
Acids Res. *47*, D442–D450.

Pfaller, C.K., Li, Z., George, C.X., and Samuel, C.E. (2011). Protein kinase PKR and RNA
adenosine deaminase ADAR1: new roles for old players as modulators of the interferon
response. Curr. Opin. Immunol. *23*, 573–582.

Qiu, Y., Xu, Y., Zhang, Y., Zhou, H., Deng, Y.-Q., Li, X.-F., Miao, M., Zhang, Q., Zhong, B.,
Hu, Y., et al. (2017). Human Virus-Derived Small RNAs Can Confer Antiviral Immunity
in Mammals. Immunity *46*, 992-1004.e5.

Qiu, Y., Xu, Y.-P., Wang, M., Miao, M., Zhou, H., Xu, J., Kong, J., Zheng, D., Li, R.-T., and
Zhang, R.-R. (2020). Flavivirus induces and antagonizes antiviral RNA interference in both
mammals and mosquitoes. Sci. Adv. *6*, eaax7989.

Robb, G.B., and Rana, T.M. (2007). RNA helicase A interacts with RISC in human cells and
functions in RISC loading. Mol. Cell *26*, 523–537.

Roy, B.B., Hu, J., Guo, X., Russell, R.S., Guo, F., Kleiman, L., and Liang, C. (2006). Association of RNA helicase a with human immunodeficiency virus type 1 particles. J. Biol. Chem. *281*, 12625–12635.

Ryman, K.D., White, L.J., Johnston, R.E., and Klimstra, W.B. (2002). Effects of PKR/RNase L-Dependent and Alternative Antiviral Pathways on Alphavirus Replication and Pathogenesis. Viral Immunol. *15*, 53–76.

Sadler, A.J., Latchoumanin, O., Hawkes, D., Mak, J., and Williams, B.R.G. (2009). An Antiviral Response Directed by PKR Phosphorylation of the RNA Helicase A. PLoS Pathog. *5*, e1000311.

Samuel, C.E. (2011). Adenosine deaminases acting on RNA (ADARs) are both antiviral and proviral. Virology *411*, 180–193.

Schoggins, J.W., Wilson, S.J., Panis, M., Murphy, M.Y., Jones, C.T., Bieniasz, P., and Rice, C.M. (2011). A diverse range of gene products are effectors of the type I interferon antiviral response. Nature *472*, 481–485.

Schuster, S., Overheul, G.J., Bauer, L., van Kuppeveld, F.J.M., and van Rij, R.P. (2019). No evidence for viral small RNA production and antiviral function of Argonaute 2 in human cells. Sci. Rep. *9*, 1–11.

Singh, M., Castillo, D., Patel, C.V., and Patel, R.C. (2011). Stress-induced phosphorylation of PACT reduces its interaction with TRBP and leads to PKR activation. Biochemistry *50*, 4550–4560.

Takahashi, T., and Ui-Tei, K. (2020). Mutual Regulation of RNA Silencing and the IFN Response as an Antiviral Defense System in Mammalian Cells. Int. J. Mol. Sci. *21*, 1348.

Takahashi, T., Nakano, Y., Onomoto, K., Murakami, F., Komori, C., Suzuki, Y., Yoneyama, M., and Ui-Tei, K. (2018a). LGP2 virus sensor regulates gene expression network mediated by TRBP-bound microRNAs. Nucleic Acids Res. *46*, 9134–9147.

Takahashi, T., Nakano, Y., Onomoto, K., Yoneyama, M., and Ui-Tei, K. (2018b). Virus Sensor RIG-I Represses RNA Interference by Interacting with TRBP through LGP2 in Mammalian Cells. Genes *9*, 511.

tenOever, B.R. (2016). The Evolution of Antiviral Defense Systems. Cell Host Microbe *19*, 142–149.

Tsai, K., Courtney, D.G., Kennedy, E.M., and Cullen, B.R. (2018). Influenza A virus-derived siRNAs increase in the absence of NS1 yet fail to inhibit virus replication. RNA N. Y. N.

van der Veen, A.G., Maillard, P.V., Schmidt, J.M., Lee, S.A., Deddouche-Grass, S., Borg, A., Kjær, S., Snijders, A.P., and Reis e Sousa, C. (2018). The RIG-I-like receptor LGP2 inhibits Dicer-dependent processing of long double-stranded RNA and blocks RNA interference in mammalian cells. EMBO J. *37*, e97479.

Wienholds, E., Koudijs, M.J., Van Eeden, F.J., Cuppen, E., and Plasterk, R.H. (2003). The microRNA-producing enzyme Dicer1 is essential for zebrafish development. Nat Genet 217–218.

Williams, B.R. (1999). PKR; a sentinel kinase for cellular stress. Oncogene *18*, 6112–6120.

Yang, W., Chendrimada, T.P., Wang, Q., Higuchi, M., Seeburg, P.H., Shiekhattar, R., and Nishikura, K. (2006). Modulation of microRNA processing and expression through RNA editing by ADAR deaminases. Nat Struct Mol Biol *13*, 13–21.

Figure legends

Figure 1. Analysis of SINV infection in HEK293T cells and characterization of FHA:DICER WT cell lines.

A. GFP fluorescent microscopy pictures of HEK293T, NoDice FHA:ctrl #1 and FHA:DICER cell lines infected (polyclonal and two clones, #4 and #17) with SINV-GFP at an MOI of 0.02 for 24 h. The left panel corresponds to GFP signal from infected cells and the right panel to a merge picture of GFP signal and brightfield. Pictures were taken with a 5x magnification. hpi: hours post-infection. **B.** Western blot analysis of DICER (DICER and HA) and GFP expression in SINV-GFP-infected HEK293T, NoDice FHA:ctrl #1 and FHA:DICER cell lines shown in **A**. Gamma-Tubulin was used as loading control. **C.** Mean (+/- SEM) of SINV-GFP viral titers in the same cell lines as in **A** infected at an MOI of 0.02 for 24 h (n = 3) from plaque assay quantification. ns: non-significant, ordinary one-way ANOVA test with Bonferroni correction. **D.** Western blot analysis of DICER (DICER and HA) and AGO2 expression in HEK293T, NoDice FHA:ctrl #1 and FHA:DICER cell lines. Gamma-Tubulin was used as loading control.

Figure 2. LC-MS/MS analysis of DICER interactome during SINV infection.

A. Volcano plot for differentially expressed proteins (DEPs) between HA IP and CTL IP in FHA:DICER mock-infected cells. Each protein is marked as a dot; proteins that are significantly up-regulated in HA IP are shown in red, up-regulated proteins in CTL IP are shown in blue, and non-significant proteins are in black. The horizontal line denotes a p-value of 0.05 and the vertical lines the Log2 fold change cutoff (-1 and 1). DICER and its cofactors (TRBP, PACT, AGO2) are highlighted in yellow. **B.** Volcano plot for DEPs between SINV-GFP (MOI of 2, 6 hpi) and mock fractions of HA IP in FHA:DICER cells. Same colour code and thresholds as in A have been applied. Proteins that are discussed in the text are highlighted in yellow and SINV proteins in purple. **C.** Gene Ontology (GO) term enrichment of proteins

up-regulated in SINV-GFP fraction of HA IP using Enrichr software (Chen et al., 2013;

Kuleshov et al., 2016). The graph displays the GO term hierarchy within the "biological

process" branch sorted by p-value ranking computed from the Fisher exact test. The length of

each bar represents the significance of that specific term. In addition, the brighter the colour is,

the more significant that term is. Viral proteins have been excluded for this analysis. **D.**

Summary of the differential expression analysis of SINV-GFP vs mock fractions from HA IP

in FHA:DICER cells. The analysis has been performed using a generalized linear model of a

negative-binomial distribution and p-values were corrected for multiple testing using the

Benjamini-Hochberg method.

1035

Figure 3. Confirmation of LC-MS/MS analysis by co-IP and BiFC.

A. Western blot analysis of HA co-IP in mock or SINV-GFP-infected (MOI of 2, 6 hpi)

FHA:DICER WT #4 cells. Proteins associated to FHA:DICER were revealed by using

antibodies targeting endogenous ADAR-1, PKR, TRBP, DHX9 or PACT proteins. In parallel,

an HA antibody was used to verify the IP efficiency and GFP antibody was used to verify the

infection. Ponceau was used as loading control. **B.** Western blot analysis of HA co-IP in mock

or SINV-GFP-infected (MOI of 2, 6 hpi) FHA:DICER WT #4 cells. The lysate was treated or

not with RNase A/T1. Proteins associated to FHA:DICER were revealed by using antibodies

targeting endogenous DHX9, p-PKR, PKR, TRBP, or PACT proteins. In parallel, an HA

antibody was used to verify the IP efficiency and GFP antibody was used to verify the infection.

Ponceau was used as loading control. The arrow points to the expected size of the FHA:DICER

protein and the asterisks correspond to aspecific bands. **C.** Western blot analysis to validate the

interaction of PKR with DICER (upper panel) and PACT (lower panel) in mock or SINV-GFP-

infected HEK293T cells. Immunoprecipitated proteins obtained from PKR pulldowns were

compared to IgG pulldowns to verify the specificity of the assay. **D.** Interactions between

DICER and TRBP, PACT or PKR were visualized by BiFC. Plasmids expressing [N-ter]Venus:DICER and TRBP:, PACT: or PKR:Venus[C-ter] were co-transfected in NoDice/ΔPKR

1052

cells for 24 h and cells were either infected with SINV at an MOI of 2 for 6 h or not. The

different combinations are indicated on the left side. Reconstitution of Venus (BiFC) signal was

observed under epifluorescence microscope. For each condition, the left panel corresponds to

Venus signal and the right panel to the corresponding brightfield pictures. Scale bar: 100 μm.

hpi: hours post-infection. **E.** BiFC experiment on fixed NoDice/ΔPKR cells treated as in D.

After fixation, cells were stained with DAPI and observed under confocal microscope. Only a

merge picture of BiFC and DAPI signals of SINV-infected cells is shown here. A higher

magnification of picture showing cytoplasmic localization of the interaction represented by a

red square is shown in the bottom left corner. Scale bar: 20 μm and 10 μm.

1062

Figure 4. Confirmation of DICER interactome upon SINV infection in HCT116 KI-

DICER cells.

A. Western blot analysis of DICER, AGO2, PKR and TRBP expression in HEK293T, HCT116

and HCT116 KI-DICER cell lines. Gamma-Tubulin and ponceau were used as loading control.

B. GFP fluorescent microscopy pictures of HEK293T, HCT116 and HCT116 KI-DICER cell

lines infected with SINV-GFP at an MOI of 0.02, 0.1 and 1 for 24 h. The left panel corresponds

to GFP signal from infected cells and the right panel to a merge picture of GFP signal and

brightfield. Pictures were taken with a 5x magnification. **C**. miR-16 expression analyzed by

northern blot in in the same cell lines as in B. Expression of snRNA U6 was used as loading

control. **D.** Western blot analysis of HA co-IP in mock or SINV-GFP-infected (MOI of 0.1, 24

hpi) HCT116 KI-DICER cells. Proteins associated to FHA-GFP:DICER were revealed by using

antibodies targeting endogenous DHX9, p-PKR, PKR, PACT or TRBP proteins. In parallel, an

HA antibody was used to verify the IP efficiency and GFP antibody was used to verify the

infection. Ponceau was used as loading control.

1077

Figure 5. Identification of DICER domains involved in DICER-PKR interaction.

A. Schematic representation of Human DICER proteins used in this study. The different

conserved domains are shown in colored boxes. DUF283: Domain of Unknown Function; PAZ:

PIWI ARGONAUTE ZWILLE domain; dsRBD: dsRNA-binding domain. hDICER WT is the

full-length protein. hDICER N1 is deleted of the first N-terminal 495 amino acids. hDICER N3

is wholly deleted of the helicase domain. hDICER Hel. is the whole DICER's helicase domain.

hDICER ΔdsRBD is deleted of the C-terminal dsRBD. **B.** Western blot analysis of HA co-IP

in mock NoDice 2.20 cells transfected with different versions of FHA:DICER proteins.

Efficiency of immunoprecipitation was assessed using anti-HA and anti-DICER antibodies and

co-IPs of TRBP, PKR and PACT were examined using appropriate antibodies. Expression of

GFP in INPUT fraction was visualized as control of SINV-GFP infection. Ponceau staining of

membranes is used as loading control. **C.** Western blot analysis of HA co-IP in NoDice 2.20

cells transfected with different versions of FHA:DICER proteins and infected with SINV-GFP

(MOI of 2, 6 hpi). Efficiency of immunoprecipitation was assessed using an anti- Flag antibody

and co-IPs of PKR, TRBP, p-PKR and PACT were examined using appropriate antibodies.

Expression of GFP in INPUT fraction was visualized as control of SINV-GFP infection.

Ponceau staining of membranes is used as loading control. **D.** Plasmids expressing the different

versions of DICER proteins fused to the N-terminal part of Venus and PKR:Venus^{C-ter} plasmid

were co-transfected in NoDice/ΔPKR cells. Cells were treated as in Fig. 3D. The different

combinations are noted on the left side. The fluorescent signal was observed using an

epifluorescence microscope. For each condition, the left panel corresponds to Venus signal and

the right panel to the corresponding brightfield pictures. Scale bar: 100 μm. hpi: hours post-

infection.

1101

1102 Figure 6. Analysis of the importance of Dicer helicase domain on SINV-GFP infection in

1103 FHA:DICER mutant stable cell lines.

A. Expression of DICER (DICER, HA), TRBP and AGO2 was analysed by western blot in

HEK293T, NoDice FHA:ctrl #1, FHA:DICER WT #4 and FHA:DICER N1 #6 cell lines.

Gamma-Tubulin was used as loading control. **B.** Northern blot analysis of miR-16 expression

in the same samples as in A. Expression of snRNA U6 was used as loading control. **C.**

Representative GFP fluorescent microscopy images of HEK293T, FHA:DICER WT #4 and

FHA:DICER N1 #6 cell lines infected with SINV-GFP at an MOI of 0.02 for 24 h. The left

panel corresponds to GFP signal and the right panel to a merge picture of GFP signal and

brightfield. Pictures were taken with a 5x magnification. hpi: hours post-infection. **D.** Western

blot analysis of DICER (DICER and HA), AGO2, PKR, and GFP expression in SINV-GFP-

infected cells in the same condition as in C. Gamma-Tubulin was used as loading control. The

asterisk correspond to aspecific bands **E.** Mean (+/- SEM) of SINV-GFP viral titers fold change

over HEK293T cells in HEK293T, NoDice 2.20, FHA:DICER WT #4 and FHA:DICER

mutants cell lines infected at an MOI of 0.02 for 24 h (n = 3) from plaque assay quantification.

* p < 0.05, ns: non-significant, ordinary one-way ANOVA test with Bonferroni correction.

1118

 Supplementary material

 Supplementary figure legends

 Figure S1. Analysis of SINV-GFP infection in FHA:DICER cell lines at different MOI

 and time points.

 A. miR-16 expression analyzed by northern blot in HEK293T, NoDice FHA:ctrl #1 and

 FHA:DICER WT #4 cell lines**.** Expression of snRNA U6 was used as loading control. **B.**

 Representative GFP pictures of HEK293T, NoDice 2.20, NoDice 4.25, NoDice FHA:ctrl #1

 and NoDice FHA:ctrl #2 cells infected with SINV-GFP at an MOI of 0.02 for 24h. The left

 panel corresponds to GFP signal and the right panel to a merge of GFP signal and the

 corresponding brightfield. Pictures were taken with a 5x magnification. hpi: hours post-

 infection. **C.** Mean (+/- SEM) of SINV-GFP viral titers in cells infected at an MOI of 0.02 for

 24 h (n=3) from plaque assay quantification. * $p < 0.05$, ns: non-significant, ordinary one-way

 ANOVA test with Bonferroni correction. **D**. Western blot analysis of DICER, AGO2 and GFP

 expression in SINV-GFP-infected cells shown in B. Gamma-Tubulin was used as loading

 control.

 Figure S2. LC-MS/MS analysis of DICER interactome during SFV infection.

 A. Volcano plot for differentially expressed proteins (DEPs) between HA IP and CTL IP in

 FHA:DICER mock-infected cells. Each protein is marked as a dot; proteins that are

 significantly up-regulated in HA IP are shown in red, up-regulated proteins in CTL IP are

 shown in blue, and non-significant proteins are in black. The horizontal line denotes a p-value

 of 0.05 and the vertical lines the Log2 fold change cutoff (-1 and 1). DICER and its cofactors

 (TRBP, PACT, AGO2) are highlighted in yellow. **B.** Left panel: Volcano plot for DEPs

 between SFV (MOI of 2, 6 hpi) and mock fractions of HA IP in FHA:DICER cells. Same colour

1144 code and thresholds as in A were applied. Proteins that are discussed in the text are highlighted

1145 in yellow and SFV proteins in purple. **C.** Summary of the differential expression analysis of

1146 SFV vs mock fractions from HA IP in FHA:DICER cells. The analysis has been performed

1147 using a generalized linear model of a negative-binomial distribution and p-values were

1148 corrected for multiple testing using the Benjamini-Hochberg method.

1149

Figure S3. Confirmation of LC-MS/MS analysis by co-IP and BiFC controls.

A. FHA:DICER WT #4 cells were infected with SINV-GFP at an MOI of 0.02 for 24 h and a

HA co-IP was performed. Eluted proteins were resolved by western blot and IP efficiency was

assessed using an HA antibody. In parallel, co-IPed proteins were visualized using appropriate

antibodies. GFP antibody was used to verify the infection and Ponceau staining serves as

loading control. **B.** 1% Agarose gel analysis of RNA extracted from INPUT of the co-IP in Fig.

3B. Ribosomal RNA integrity was compared to a control HEK293T cell line. RNAs were

revealed using ethidium bromide under UV. **C.** Schematic representation of Human DICER

proteins used for BiFC positive and negative controls. The different conserved domains are

shown in colored boxes. DUF283: Domain of Unknown Function; PAZ: PIWI ARGONAUTE

ZWILLE domain; dsRBD: dsRNA-binding domain. hDICER WT is the full-length protein.

hDICER N1 is deleted of the first N-terminal 495 amino acids. **D.** Expression of BiFC plasmids

was assessed by western blot. DICER proteins (WT and N1) and PKR were visualized using

antibodies targeting endogenous proteins, whereas TRBP and PACT were detected using GFP

antibody. Antibody targeting the SINV coat protein (CP) was used as infection control. Ponceau

staining was used as loading control. **E.** Positive and negative BiFC controls on fixed

NoDice/ΔPKR cells. After co-transfection, cells were infected with SINV at an MOI of 2 for 6

h and fixed. After fixation, cells were stained with DAPI and observed under confocal

microscope. Merge pictures of BiFC and DAPI signals of SINV-infected cells are shown. A

higher magnification of images showing the interaction represented by a red square is shown

in the bottom left corner. Scale bar: 20 μm and 10 μm. **F.** Expression of BiFC plasmids was

assessed by western blot. DICER, PKR, TRBP and PACT were detected using GFP antibody.

Antibody targeting the SINV coat protein (CP) was used as infection control. Gamma-Tubulin

was used as loading control. The asterisk corresponds to an aspecific band. **G.** Interactions

between DICER and TRBP, PACT or PKR were visualized by BiFC. Plasmids expressing $^{N\text{-}}$

^{ter}Venus:DICER and TRBP:, PACT: or PKR:Venus$^{C\text{-}ter}$ were co-transfected in HEK293T cells

for 24 h and cells were either infected with SINV at an MOI of 2 for 6 h or not. The different

combinations are indicated on the left side. Reconstitution of Venus (BiFC) signal was observed

under epifluorescence microscope. For each condition, the left panel corresponds to Venus

signal and the right panel to the corresponding brightfield pictures. Scale bar: 100 μm.

1180

Figure S4. Confirmation of DICER interactome upon SINV infection in HCT116 KI-

DICER cells.

A. Schematic representation of DICER WT and Flag-HA(FHA)-GFP kocked-in (KI) alleles.

FHA sequence is in purple, GFP in green, DICER 5'UTR in orange and DICER coding region

in yellow. The gRNA used to generate the KI was designed to target the first coding exon of

DICER gene. **B.** PCR on genomic DNA extracted from WT and KI cells. **C.** An oligo outside

the homologous recombination region and an oligo within the GFP tag were used to verify the

presence of a 1040bp amplicon in HCT116 KI-DICER clone. Sequencing results corresponding

to this region are shown. **D.** Western blot analysis of DICER, p-PKR, PKR and p-eIF2α

expression in mock or SINVGFP-infected HEK293T and HCT116 KI-DICER cell lines at an

MOI of 2 for 6h or 16h and 0.02 for 24h. GFP antibody was used to verify the infection. Ponceau

and gamma-Tubulin were used as loading control.

1193

Figure S5. Interaction analysis between the different versions of DICER and TRBP or PACT using BiFC assay.

NoDice/ΔPKR cells were co-transfected for 24 h with plasmids expressing the different versions of DICER proteins fused to the N-terminal part of Venus and either TRBP:Venus^{C-ter} (**A**) or PACT:Venus^{C-ter} (**B**). Cells were then infected with SINV at an MOI of 2 for 6 h and Venus signal was observed under epifluorescence microscope. The left panel corresponds to Venus signal and the right panel to the corresponding brightfield picture. Pictures were taken with a 5x magnification. hpi: hours post-infection. Scale bar: 100 μm.

Figure S6. Analysis of the importance of Dicer helicase domain on SINV-GFP infection in FHA:DICER mutant stable cell lines.

A. Northern blot analysis of miR-16 expression in HEK293T, NoDice 2.20, NoDice FHA:ctrl #2, FHA:DICER WT polyclonal, FHA:DICER N1 #6, FHA:DICER Hel. #2.6, and FHA:DICER N3 #2.13. Expression of snRNA U6 was used as loading control. **B.** Representative GFP fluorescent microscopy images of HEK293T, NoDice 2.20, FHA:DICER mutants cell lines infected with SINV-GFP at an MOI of 0.02 for 24 h. The left panel corresponds to GFP signal and the right panel to a merge picture of GFP signal and brightfield. Pictures were taken with a 5x magnification. hpi: hours post-infection. **C**. Mean (+/- SEM) of SINV-GFP viral titers over FHA:DICER WT #4 cells in FHA:DICER N1 #6, FHA:DICER N3 #2.13, NoDice FHA:ctrl #2 and NoDice 2.20 cell lines infected at an MOI of 0.02 for 24 h (n = 3) from plaque assay quantification. *** $p < 0.001$, ns: non-significant, ordinary one-way ANOVA test with Bonferroni correction.

Supplementary tables

1218   **Supp. Table 1:** Top 100 proteins that are differentially immunoprecipitated in mock-infected

1219   FHA:DICER cells by the HA and Myc (CTL) antibodies. Related to Figure 2.

1220

1221   **Supp. Table 2:** Top 100 proteins that are differentially immunoprecipitated with the HA

1222   antibody in SINV-infected vs mock-infected FHA:DICER cells. Related to Figure 2.

1223

1224   **Supp. Table 3:** Top 100 proteins that are differentially immunoprecipitated in mock-infected

1225   FHA:DICER cells by the HA and Myc (CTL) antibodies, in the SFV infection experiment.

1226   Related to Figure S2.

1227

1228   **Supp. Table 4:** Top 100 proteins that are differentially immunoprecipitated with the HA

1229   antibody in SFV-infected vs mock-infected FHA:DICER cells. Related to Figure S2.

1230

1231   **Supp. Table 5:** List of primers used in this study.

Figure 1

A

| | GFP | GFP + Brightfield |

HEK293T

NoDice 2.20

FHA:ctrl #1

FHA:DICER WT #4

FHA:DICER polyclonal

FHA:DICER WT #17

SINV-GFP MOI 0.02 24 hpi

B

Mock   SINV-GFP MOI 0.02 24 hpi

HEK293T | FHA:ctrl #1 | FHA:DICER WT #4 | HEK293T | FHA:ctrl #1 | FHA:DICER WT #4

250 kDa — DICER
250 kDa — HA
35 kDa / 25 kDa — GFP
55 kDa — Tubulin

C

SINV-GFP titer MOI 0.02 24 hpi

Log10 viral titer PFU/mL

10^8
10^7
10^6
10^5

ns ns ns ns ns

HEK293T | FHA:ctrl #1 | FHA:DICER WT #4 | FHA:DICER WT polyclonal | FHA:DICER WT #17

NoDice 2.20

D

HEK293T | FHA:DICER polyclonal | FHA:DICER WT #4 | FHA:DICER WT #17

SINV-GFP   - +   - +   - +   - +

250 kDa — DICER
250 kDa — HA
35 kDa / 25 kDa — GFP
55 kDa — Tubulin
100 kDa — AGO2
55 kDa — Tubulin

Figure 2

A

FHA:DICER IP vs CTL IP

B

SINV-GFP vs Mock in FHA:DICER IP

C

RNA binding (GO:0003723)

ATP-dependent helicase activity (GO:0008026)

snoRNA binding (GO:0030515)

RNA helicase activity (GO:0003724)

ATP-dependent RNA helicase activity (GO:0004004)

RNA-dependent ATPase activity (GO:0008186)

double-stranded RNA binding (GO:0003725)

actin-dependent ATPase activity (GO:0030898)

ATP-dependent DNA helicase activity (GO:0004003)

RNA stem-loop binding (GO:0035613)

D

Gene name	Mean spectral count		LogFC	Adjusted p-value
	SINV-GFP IP	MOCK IP		
E2AK2 (PKR)	123.3	0.3	7.979	7.316e-102
DSRAD (ADAR-1)	181.3	28.3	2.563	1.421e-73
DHX9	247.3	50	2.194	4.012e-83
PRKRA (PACT)	63.3	13.7	2.093	1.158e-20

Figure 3

Figure 4

A

B

C

D

Figure 5

A

B

NoDice 2.20 mock

C

NoDice 2.20 infected with SINV-GFP MOI 2 6 hpi

D

Figure 6

Figure S1

A

HEK293T
NoDice FHA:ctrl #1
FHA:DICER WT #4

pre-miR-16
miR-16
snRNA U6

B

GFP   GFP + Brightfield

HEK293T

NoDice 2.20

NoDice 4.25

FHA:ctrl #1

FHA:ctrl #2

NoDice 2.20

SINV-GFP MOI 0.02 24 hpi

C

SINV-GFP titer MOI 0.02 24 hpi

Log10 viral titer PFU/mL

10^8
10^7
10^6
10^5

ns   ns   *   *

HEK293T   NoDice 2.20   NoDice 4.25   FHA:ctrl #1   FHA:ctrl #2

NoDice 2.20

D

HEK293T
NoDice 2.20
NoDice 4.25
FHA:ctrl #1
FHA:ctrl #2

- + - + - + - + - +   SINV-GFP

250 kDa   DICER
100 kDa   AGO2
35 kDa   GFP
25 kDa
55 kDa   Tubulin

Figure S2

Gene name	Mean spectral count		LogFC	Adjusted p-value
	SFV IP	Mock IP		
E2AK2 (PKR)	221	0.5	8.344	2.611e-119
PRKRA (PACT)	131.5	4.5	4.697	8.415e-57
DSRAD (ADAR-1)	268	22	3.466	9.118e-45
DHX9	265	31	2.955	1.245e-80

Figure S3

A

FHA:DICER WT #4

	INPUT		IP				
			HA	CTL	HA	CTL	SINV-GFP
	-	+	-	-	+	+	
250 kDa							HA
70 kDa							PKR
35 kDa							PACT
							Ponceau
130 kDa							ADAR-1
130 kDa							DHX9
55 kDa							TRBP
25 kDa							GFP
							Ponceau

B

FHA:DICER WT #4

HEK293T	-RNase A/T1		+RNase A/T1		
	-	+	-	+	SINV-GFP

RNA INPUT

C

hDicer WT — HEL 1 | HEL 2i | HEL 2 | DUF 283 | Platform | PAZ | Connector | RNase IIIa | RNase IIIb | dsRBD
DExD/H-box helicase
1 ... 1922

hDicer N1 — HEL 2 | DUF 283 | Platform | PAZ | Connector | RNase IIIa | RNase IIIb | dsRBD
496 ... 1922

D

N-terVenus:DICER

	TRBP:VenusC-ter		PKR:VenusC-ter		PACT:VenusC-ter		NT		N-terVenus:DICER N1 + TRBP:VenusC-ter		NT		SINV
	-	+	-	+	-	+	-	+	-	+	-	+	
250 kDa													DICER
100 kDa													PKR
55 kDa													GFP
35 kDa / 25 kDa													SINV CP
													Ponceau

E

BiFC+DAPI   BiFC+DAPI

N-terVenus:DICER + TRBP:VenusC-ter   N-terVenus:DICER N1 + TRBP:VenusC-ter

F

N-terVenus:DICER

	TRBP:VenusC-ter		PKR:VenusC-ter		PACT:VenusC-ter		SINV
	-	+	-	+	-	+	
250 kDa							GFP
100 kDa							GFP
55 kDa							
35 kDa / 25 kDa							SINV CP
55 kDa							Tubulin

G

	MOCK		SINV MOI 2 6 hpi	
	BiFC	Brightfield	BiFC	Brightfield
N-terVenus:DICER + TRBP:VenusC-ter				
N-terVenus:DICER + PACT:VenusC-ter				
N-terVenus:DICER + PKR:VenusC-ter				

Figure S4

A

B

C

D

Figure S5

Figure S6

A

HEK293T

FHA:DICER

NoDice 2.20
FHA:ctrl #2
WT polyclonal
N1 #6
Hel. #2.6
N3 #2.13

pre-miR-16

miR-16

snRNA U6

B

GFP   GFP + Brightfield

HEK293T

NoDice 2.20

FHA:DICER WT #4

FHA:DICER N1 #6

FHA:DICER N3 #2.13

FHA:DICER Hel. #2.6

FHA:ctrl #2

NoDice 2.20

SINV-GFP MOI 0.02 24 hpi

C

SINV-GFP titer MOI 0.02 24 hpi

Viral Titer PFU/mL

ns   ns

***   ***

FHA:DICER WT #4
FHA:DICER N1 #6
FHA:DICER N3 #2.13
FHA:ctrl #2
NoDice 2.20

NoDice 2.20