Human DICER helicase domain recruits PKR and modulates its antiviral activity

Thomas C. Montavon[1,*], Morgane Baldaccini[1,*], Mathieu Lefèvre[1,*], Erika Girardi[1], Béatrice Chane-Woon-Ming[1], Mélanie Messmer[1], Philippe Hammann[2], Johana Chicher[2], Sébastien Pfeffer[1,‡]

[1] Université de Strasbourg, Architecture et Réactivité de l'ARN, Institut de Biologie Moléculaire et Cellulaire du CNRS, 2 allée Konrad Roentgen, 67084 Strasbourg France

[2] Université de Strasbourg, Institut de Biologie Moléculaire et Cellulaire du CNRS, Plateforme Protéomique Strasbourg – Esplanade, 2 allée Konrad Roentgen, 67084 Strasbourg France

*These authors contributed equally

‡To whom correspondence should be addressed: spfeffer@unistra.fr

Short title: Regulation of PKR by DICER in viral infection

Abstract

The antiviral innate immune response mainly involves type I interferon (IFN) in mammalian

cells. The contribution of the RNA silencing machinery remains to be established, but several

recent studies indicate that the ribonuclease DICER can generate viral siRNAs in specific

conditions. It has also been proposed that type I IFN and RNA silencing could be mutually

exclusive antiviral responses. In order to decipher the implication of DICER during infection

of human cells with alphaviruses such as the Sindbis virus and Semliki forest virus, we

determined its interactome by proteomics analysis. We show that DICER specifically interacts

with several double-stranded RNA binding proteins and RNA helicases during viral infection.

In particular, proteins such as DHX9, ADAR-1 and the protein kinase RNA-activated (PKR)

are enriched with DICER in virus-infected cells. We demonstrate that the helicase domain of

DICER is essential for this interaction and that its deletion confers antiviral properties to

DICER in an RNAi-independent, PKR-dependent, manner.

Author summary

While RNAi has been recognized as an efficient antiviral defense system in organisms such as

plants and insects, its physiological importance in mammals remains to be determined. DICER

is an enzyme involved in cleaving long double-stranded RNAs and is essential for RNAi

induction. Using mass spectrometry analysis, we determined its interactome in human cells and

showed that RNA binding proteins such as PKR are specifically enriched upon infection with

the Sindbis virus or the Semliki forest virus. We determined that the N terminal helicase domain

of the DICER protein acts as a platform to recruit these factors during infection and that its

deletion confers an antiviral activity to DICER.

Introduction

In mammalian cells, the main antiviral defense system involves the activation of a signaling cascade relying on production of type I interferon (IFN I). This pathway depends on the recognition of extrinsic signals or pathogen associated molecular patterns (PAMPs) by dedicated host receptors. Double-stranded (ds) RNA, which can originate from viral replication or convergent transcription, is a very potent PAMP and can be sensed in the cell by various proteins among which a specific class of DExD/H-box helicases called RIG-I-like receptors (RLRs) [1]. RLRs comprise RIG-I, MDA5 and LGP2 and transduce viral infection signals to induce expression of IFN I cytokines that act in autocrine and paracrine fashion. These cytokines then trigger the expression of hundreds of interferon-stimulated genes (ISGs) to stop the virus in its tracks [2]. Among those ISGs, dsRNA-activated protein kinase R (PKR) plays an important role in antiviral defense by blocking cellular and viral translation upon direct binding to long dsRNA [3]. PKR is a serine-threonine kinase that dimerizes and auto-phosphorylates upon activation. It then phosphorylates numerous cellular targets among which the translation initiation factor eIF2α, which results in the inhibition of cap-dependent translation [4]. Accordingly, translation of many RNA viruses, including alphaviruses, is inhibited by PKR [5–7]. PKR is also involved in other cellular pathways including apoptosis, autophagy and cell cycle [3,8].

RNAi is another evolutionary conserved pathway triggered by long dsRNA sensing [9]. One key component in this pathway is the type III ribonuclease DICER, which is also essential for micro (mi)RNA biogenesis [10,11]. These small regulatory RNAs are sequentially produced by the two ribonucleases DROSHA and DICER, before being loaded into an Argonaute (AGO) effector protein in order to regulate their target mRNAs [12]. Whatever its substrate, be it long dsRNA or miRNA precursor, DICER relies on interacting with co-factors to be fully functional. In mammalian cells, the TAR-RNA binding protein (TRBP), a dsRNA binding protein (dsRBP), was shown to play a role in the selection of DICER substrates, its stabilization, strand selection and incorporation into AGO2 [13]. The interaction with TRBP is well characterized and depends on the helicase domain of DICER and the third dsRNA binding domain (dsRBD) of TRBP [14]. Another dsRBP, the protein activator of interferon-induced protein kinase R (PACT), was also described as an important cofactor of DICER. Although its function is not fully understood, PACT seems to also participate in miRNA loading and strand selection [15,16] *via* protein-protein interaction between the DICER helicase domain and the third dsRBD of PACT [17].

It is now common knowledge that RNAi is the main antiviral defense system in several
phyla such as plants, arthropods and nematodes (reviewed in [18]). However, its exact
contribution in the mammalian antiviral response remains unclear [19–21]. Recent studies
indicate that a functional antiviral RNAi does exist in mammals in specific cases. A functional
antiviral RNAi response was first detected in undifferentiated mouse embryonic stem cells [22]
lacking the IFN response, suggesting that these two pathways could be incompatible. Indeed,
in mammalian somatic cells deficient for MAVS or IFNAR, two components of the interferon
response, an accumulation of DICER-dependent siRNAs derived from exogenous long dsRNA
was detected [23]. In addition, the RLR LGP2 was found interacting with both DICER and
TRBP, blocking respectively siRNA production and miRNA maturation [24–26]. Moreover,
AGO4 was recently shown to be involved in antiviral RNAi against Influenza A virus (IAV),
Vesicular stomatitis virus (VSV) and Encephalomyocarditis virus (EMCV) [27]. Finally, viral
suppressors of RNAi (VSRs) have been shown to prevent DICER from playing an antiviral role
in mammalian cells [28,29]. Nonetheless, several studies reported no detection of viral siRNAs
in mammalian somatic cells infected with several viruses [30–32]. In somatic cells, only a
helicase-truncated form of human DICER could produce siRNAs from IAV genome [33], but
it also turned out that these siRNAs cannot confer an antiviral state [34].

Based on these conflicting observations, we decided to study the involvement of DICER
during infection of human cells with the Sindbis virus (SINV). SINV is a member of the
Togaviridae family in the alphavirus genus, which is transmitted by mosquitoes to mammals
and can induce arthritogenic as well as encephalitic diseases [35]. It is widely used as a
laboratory alphaviruses model as it infects several cell types and replicates to high titers. SINV
has a positive stranded RNA genome of about 12 kb, which codes for two polyproteins that
give rise to non-structural and structural proteins, including the capsid. Moreover, upon viral
replication, a long dsRNA intermediate, which can be sensed by the host antiviral machinery,
accumulates. Of note, SINV dsRNA can be cleaved into siRNA in insects as well as in human
cells expressing the Drosophila DICER-2 protein [36]. Nonetheless, although human DICER
has the potential to interact with the viral RNA duplex, we did not find evidence that SINV
dsRNA could be processed into siRNAs in somatic mammalian cells [30,36]. We thus
hypothesized that specific proteins could interfere with DICER during SINV infection by direct
interaction and limit its accessibility and/or activity. To address this hypothesis, we generated
HEK293T cells expressing a tagged version of human DICER that could be
immunoprecipitated in mock or SINV-infected cells in order to perform a proteomic analysis
of its interactome. Among the proteins co-immunoprecipitated with DICER and that were

specifically enriched upon infection, we identified dsRBPs such as ADAR1, DHX9, PACT and PKR. We further validated the direct interaction between DICER and PKR upon SINV infection. We also demonstrated that the interactions of the endogenous DICER with PKR, PACT and DHX9 could also be detected in SINV-infected, but not mock-infected, HCT116 cells. We dissected the protein domains necessary for this interaction and we found that DICER helicase domain plays a fundamental role as a recruitment platform for PKR but also for other co-factors. Finally, we also show that expression of a helicase-truncated version of DICER has a negative effect on SINV infection. Importantly, this antiviral phenotype is independent of RNAi, but requires the presence of PKR. Our results indicate that DICER interactome is highly dynamic and directly link components of RNAi and IFN pathways in modulating the cellular response to viral infection.

Results

Establishment of a HEK293T cell line expressing FLAG-HA tagged DICER

In order to be able to study the interactome of the human DICER protein during viral infection, we transduced *Dicer* knock-out HEK293T cells (NoDice 2.20) [37] with either a lentiviral construct expressing a FLAG-HA-tagged wild type DICER protein (FHA:DICER WT #4) or a construct without insert as a negative control (FHA:ctrl #1). After monoclonal selection of stably transduced cells, we first characterized one clone of both FHA:DICER WT and of the FHA:ctrl cell lines. We first confirmed that the expression of the tagged version of DICER restored the miRNA biogenesis defect observed in the NoDice cells (Fig. S1A). We then monitored the phenotype of these cells during SINV infection by using as a readout of viral infection the modified version of SINV able to express GFP from a duplicated sub-genomic promoter (SINV-GFP) [38]. At 24 hours post-infection (hpi) and a multiplicity of infection (MOI) of 0.02, the GFP fluorescence observed in FHA:DICER WT #4 cells and HEK293T cells was similar. However, the NoDice FHA:ctrl #1 cells displayed a decrease in GFP signal (Fig. 1A). Western blot analysis of GFP expression confirmed the observations by epifluorescence microscopy, *i.e.* a significantly lower accumulation of GFP in the absence of the DICER protein (Fig. 1B). We therefore wished to confirm the effect of DICER loss on SINV-GFP infection in another NoDice cell line, *i.e.* the NoDice clone 4.25 [39], and in another clone of the NoDice 2.20 FHA:ctrl cells (NoDice FHA:ctrl #2). We observed a similar decrease of SINV-GFP infection in NoDice 2.20 cells and two independent NoDice FHA:ctrl clones

compared to HEK293T cells as shown by GFP microscopy (Fig. S1B), by titration of the virus (Fig. S1C) and by western blot analysis (Fig. S1D). However, the independent NoDice 4.25 *Dicer* knock-out clone appeared mostly unaffected compared to HEK293T cells in term of GFP accumulation and viral titer (Fig. S1B-D). This suggests that, despite the observed slight effect on SINV-GFP in NoDice 2.20 cells (Fig. 1), DICER proviral effect is not reproductible in an independent clone and therefore could not be generalized.

In order to evaluate whether different expression levels of DICER in a NoDice background could rescue the SINV infection phenotype observed in HEK293T cells, we also infected both the FHA:DICER WT polyclonal and an independent FHA:DICER WT clone (FHA:DICER WT #17) with SINV-GFP (Fig. 1A and Fig. 1C-D). We confirmed that the GFP fluorescence observed by microscopy (Fig. 1A), as well as the viral titers and the GFP protein accumulation (Fig. 1C-D) in all tested FHA:DICER lines were comparable to the ones observed in HEK293T cells. Moreover, there was no striking difference in AGO2 expression between the FHA:DICER lines (Fig. 1D).

Altogether, these results indicate that the FHA-tagged DICER protein can functionally complement the lack of DICER in terms of miRNA biogenesis (Fig. S1A) and can therefore be used for proteomics studies. Moreover, because we could not observe significant differences in terms of SINV infection (Fig. 1) between the different FHA:DICER clones tested, we decided to select one line, namely FHA:DICER WT#4, for further analysis.

Analysis of DICER interactome during SINV infection by mass spectrometry

Our molecular tool being validated, we then focused on determining the interactome of FHA:DICER during SINV infection. We wanted to look at DICER interactome at an early infection time point to isolate cellular factors that could potentially modulate either DICER accessibility or its effect on viral dsRNA. As SINV replicates quickly upon cellular entry, we chose to set up the infection conditions to a duration of 6 hours at an MOI of 2.

We performed an anti-HA immunoprecipitation experiment (HA IP) coupled to label-free LC-MS/MS analysis in FHA:DICER WT #4 cells either mock-infected or infected for 6 h at an MOI of 2 with SINV-GFP. In parallel, we performed an anti-MYC immunoprecipitation as a negative control (CTL IP). The experiments were performed in technical triplicate in order to have statistically reproducible data for the differential analysis, which was performed using spectral counts. Prior to the detailed analysis of the results, we verified that there was no confounding factor in the experimentation by performing a Principal Component Analysis

(PCA). This allowed us to see that the replicates were very homogenous and that the different

samples were well separated based on the conditions.

To check the specificity of the HA immunoprecipitation, we first compared the proteins

identified in the HA IP with the ones identified in the CTL IP in mock-infected cells.

Differential expression analysis allowed us to calculate a fold change and an adjusted p-value

for each protein identified and to generate a volcano plot representing the differences between

HA and CTL IP samples. Applying a fold change threshold of 2 ((LogFC) > 1), an adjusted p-

value threshold of 0.05 and a cutoff of at least 5 spectral counts in the most abundant condition,

we identified 258 proteins differentially immunoprecipitated between the two conditions out of

1318 proteins (Fig. 2A and Supp. Table 1). Among these, 123 proteins were specifically

enriched in the HA IP. The most enriched protein was DICER, followed by its known co-factors

TRBP and PACT (also known as PRKRA) [13,17]. We were also able to retrieve AGO2,

indicating that the RISC loading complex was immunoprecipitated and that proteins retrieved

in our HA IP are specific to DICER immunoprecipitation.

We next performed the differential expression analysis of proteins retrieved in the HA

IP in SINV-GFP compared to mock-infected cells. Among 1342 proteins, 296 were

differentially retrieved between conditions (Fig. 2B and Supp. Table 2). Of these, 184 proteins,

including viral ones, were at least 2-fold enriched in SINV-GFP-infected cells. GO-term

analysis showed a significant enrichment in RNA binding proteins including double-stranded

RNA binding proteins and RNA helicases (Fig. 2C). We then generated a functional protein

association network using STRING on the top 100 proteins enriched in SINV-infected

compared to mock-infected cells (Fig. 2D). The resulting STRING network confirmed that a

limited number of these proteins are known to be interacting with DICER, but that they are all

engaged in other complexes (*e.g.* DHX9, DDX18) that could partly explain the presence of

some candidates in the mass spectrometry data. In addition, a large number of these proteins

are involved in RNA metabolic processes (Fig. 2D, in red), or in their regulation (in blue), while

a whole cluster is composed of dsRNA binding proteins (in green). Among the RNA binding

proteins retrieved, the top and most specific DICER interactor is the interferon-induced, double-

stranded (ds) RNA-activated protein kinase PKR (also known as E2AK2), which is enriched

more than 250 times in virus-infected cells (Fig. 2B and E). We were also able to identify the

dsRNA-specific adenosine deaminase protein ADAR-1 (also known as DSRAD), as well as

PACT, which were enriched 5.9 and 4.2 times respectively in SINV-GFP-infected cells

compared to mock-infected cells (Fig. 2B and E). Among the isolated RNA helicases, we

identified the ATP-dependent RNA helicase A protein DHX9, which is implicated in *Alu*

element-derived dsRNA regulation and in RISC loading [40,41]. In order to verify if the observed interactions were specific to SINV we performed the same experiments with another virus of the *Togaviridae* family, the Semliki forest virus (SFV). In this analysis, we were able to retrieve ADAR-1, DHX9, PACT and PKR, specifically enriched in SFV-infected samples (Fig. S2 and Supp. Tables 3 and 4). These results show that these interactions can be retrieved in *Togaviridae*-infected cells.

Taken together, our data indicate that several proteins interacting with DICER in virus-infected cells are involved in dsRNA sensing and/or interferon-induced antiviral response.

DICER and PKR interact *in vivo* in the cytoplasm during SINV infection

To validate the LC-MS/MS analysis, we performed a co-immunoprecipitation (co-IP) followed by western blot analysis in FHA:DICER WT #4 cells infected with SINV-GFP at an MOI of 2 for 6 h. Whereas TRBP interacted equally well with FHA:DICER in mock and SINV-GFP-infected cells, ADAR-1, PKR, DHX9 and PACT were only retrieved in the HA IP in SINV-infected cells (Fig. 3A). We verified that these interactions could also be observed at a later time post-infection by performing the HA IP in FHA:DICER WT #4 cells infected with SINV-GFP for 24 h at an MOI of 0.02. This indicates that the specific interactions between DICER and ADAR-1, DHX9, PACT or PKR occur at an early stage of the SINV infection and remain stable in time in virus-infected cells (Fig. S3A).

In order to verify whether these interactions were mediated by RNA, we performed an anti-HA co-IP experiment on an RNase A/T1 treated total extract from FHA:DICER WT #4 cells infected with SINV-GFP at an MOI of 2 for 6h. Since the RNase treatment was performed at relatively low salt concentration (140 mM NaCl), RNase A should cleave dsRNA [42,43] and we should therefore assess both ss and dsRNA-dependency in these conditions. We confirmed the efficiency of the RNase treatment by ethidium bromide staining visualisation of total RNA on an agarose gel (Fig. S3B). TRBP equally interacted with FHA:DICER, with or without RNase treatment, in mock and SINV-GFP-infected cells (Fig. 3B). Instead, the virus-induced interactions between DICER and PKR or PACT upon SINV-GFP infection were almost totally lost in the RNase-treated samples. Upon virus infection, PKR is phosphorylated to be activated and exert its antiviral function [4]. Using an antibody targeting the phosphorylated form of PKR (p-PKR), we looked for p-PKR before and after RNase treatment. The virus-enriched interactions between DICER and p-PKR or DHX9 were completely lost upon RNase treatment. These results therefore indicate that RNA molecules (either single- or

double-stranded) facilitate DICER interaction with DHX9, PACT and PKR and its active form,

although the complex may also partially interact in an RNA-independent manner.

Because of the involvement of PKR in antiviral response [44] and the fact that it shares

common co-factors with DICER, namely TRBP and PACT [45,46], we decided to focus our

analysis on the DICER-PKR interaction. To confirm the biological relevance of this interaction,

we first performed a reverse co-IP to immunoprecipitate the endogenous PKR protein in

HEK293T cells infected or not with SINV-GFP. While PACT interacted with PKR both in

mock and in SINV-GFP-infected cells as expected (Fig. 3C lower panel), DICER co-

immunoprecipitated with the endogenous PKR only in virus-infected cells as previously

observed (Fig. 3C upper panel).

To further determine whether DICER and PKR could directly interact *in vivo*, we set up

a bi-molecular fluorescent complementation assay (BiFC) experiment [47]. To this end, we

fused the N- or C-terminal halves of the Venus protein ([N-ter]Venus or [C-ter]Venus) to DICER and

to PKR but also to TRBP and PACT. Since we showed above that an N-terminally tagged

DICER was functional, we fused the Venus fragments at the N-terminal end of DICER. For the

other three proteins, we fused the Venus fragments at the N- or C-terminus and selected the

best combination. To avoid interaction with the endogenous DICER and PKR proteins, we

conducted all BiFC experiments in NoDice/ΔPKR HEK293T cells [33]. In order to control the

BiFC experiments, we chose to exploit the well characterized DICER-TRBP interaction, which

is known to occur via the DICER DEAD-box helicase domain [14]. We therefore used the wild-

type DICER protein as a positive control and a truncated version of DICER protein lacking part

of this helicase domain and called DICER N1 [33] as a negative control (Fig. S3C). We first

confirmed the expression of the tagged proteins by western blot analysis (Fig. S3D) and then,

we tested the interactions between DICER and TRBP or PACT or PKR. We co-transfected the

Venus constructs for 24 h and then infected cells with SINV or not for 6 h at a MOI of 2. A

comparable fluorescent signal was observed both in mock- and SINV-infected cells when [N-]

[ter]Venus:DICER was co-transfected with either PACT or TRBP fusion construct (Fig. 3D).

Although we initially expected an increase of the Venus fluorescence in SINV-infected cells,

overall we observed a similar signal for the DICER-PKR interaction both in mock- and SINV-

infected cells, probably due to the fact that both proteins are transiently overexpressed in this

experiment. The same holds true for the DICER-PACT interaction that can also be seen both

in mock- and SINV-infected cells.

As a control and to rule out any aspecific interactions between the different proteins

tested, we also monitored the DICER-N1-TRBP interaction by BiFC. As expected, no

fluorescent signal was observed in cells co-transfected with [N-ter]Venus:DICER N1 and

TRBP:Venus[C-ter] (Fig. S3E), confirming that DICER helicase domain is required for its

interaction with TRBP [14] and validating the specificity of the BiFC approach.

To further confirm that the absence of PKR did not influence the interactions of TRBP

or PACT with DICER, we also performed a BiFC analysis in HEK293T cells. After verifying

that in this context as well, fusion proteins were expressed as expected (Fig. S3F), we observed

that the results were similar as in NoDiceΔPKR cells (Fig. S3G).

To gain more insight into the subcellular localization of these interactions during SINV

infection, we performed the BiFC experiments, fixed the cells and observed them under a

confocal microscope. We observed a cytoplasmic fluorescent signal for DICER-TRBP and

DICER-PACT interactions (Fig. 3E upper and middle panels), which is in agreement with their

canonical localization for the maturation of most miRNAs [10,48]. Similarly, co-transfection

of DICER and PKR led to a strong Venus signal homogeneously distributed in the cytoplasm

(Fig. 3E lower panel).

Collectively, these results formally confirm that DICER interacts with several RNA

helicases and dsRNA-binding proteins in virus-infected cells, among which PKR, and that for

the latter this interaction occurs in the cytoplasm.

DICER interactome changes upon SINV infection are not cell-type specific

To further validate our DICER interactome results and generalize them to another biological

system, we performed co-IP experiments on the endogenous DICER in a different cell type. To

this end, the FLAG-HA-GFP tag was knocked into (KI) the *Dicer* locus in human colon

carcinoma cells (HCT116) by CRISPR-Cas9-mediated homologous recombination (Fig. S4A-

C). A guide RNA (gRNA) targeting the region corresponding to Dicer ATG and a DNA

template for homologous recombination bearing the FLAG-HA-GFP sequence surrounded by

the upstream and downstream arms of Dicer were used to generate the resulting cell line

referred to as HCT116 KI-DICER cells. The expected insertion of the tag in one of the two

Dicer allele was assessed by PCR amplification and Sanger sequencing (Fig. S4A-C). In

agreement, we could detect two bands for DICER protein by western blot in the HCT116 KI-

DICER cells, which confirmed that this cell line is heterozygous (Fig. 4A).

We additionally verified the expression of specific DICER-interacting proteins, such as

AGO2, PKR or TRBP, in HCT116 KI-DICER cells compared to the parental HCT116 cells

and to HEK293T cells (Fig. 4A). We also measured the production of mature miRNAs, such
as miR-16, by northern blot analysis and confirmed that miRNA expression is maintained in
HCT116 KI-DICER cells (Fig. 4B). Of note, the GFP inserted at the *Dicer* locus could not be
detected by epifluorescence microscopy in the HCT116 KI-DICER cells, which probably
reflects the low abundance of the DICER protein.

We then determined whether SINV-GFP infection was comparable in HCT116 cells and
HEK293T cells. We infected HCT116, HCT116 KI-DICER and HEK293T cells with SINV-
GFP at three different MOI (0.02, 0.1 and 1) and measured GFP fluorescence by microscopy at
24 hpi (Fig. 4C). Both HCT116 and HCT116 KI-DICER cells expressed GFP upon infection
with SINV-GFP, although with a lower intensity than HEK293T cells. We also verified by
western blot analysis the accumulation of GFP and the phosphorylation of both PKR and eIF2α
upon SINV-GFP infection of HCT116 KI-DICER and HEK293T cells (Fig. S4D) and chose as
optimal SINV-GFP condition of infection in HCT116 KI-DICER cells the MOI of 0.1 for 24h.

To validate the DICER interactions observed in HEK293T FHA:DICER cells, we then
performed anti-HA co-IP experiments followed by western blot analysis in HCT116 KI-DICER
cells infected or not with SINV-GFP. We successfully retrieved TRBP interacting with DICER
in both mock and infected cells, whereas DHX9, PKR (phosphorylated or not) and PACT were
only retrieved in the HA IP in infected cells (Fig. 4D). These results not only confirm that the
endogenous DICER specifically interacts with DHX9, PACT and PKR upon SINV infection,
but also that these interactions are not restricted to one specific cell type.

The helicase domain of DICER is required for its interaction with PKR
Even though DICER and PKR are likely brought together by RNA, specific protein domains
might be involved in stabilizing the complex. Therefore, we next determined the domain of
DICER required for its interaction with PKR. Since its helicase domain was previously shown
to be involved in the interaction with TRBP and PACT [14,17], we speculated that it could also
be implicated in binding PKR. To test this hypothesis, we cloned several versions of DICER
proteins wholly or partly deleted of the helicase domain (Fig. 5A DICER N1 and N3). In
addition, we also cloned the helicase domain alone (Fig. 5A DICER Hel.) and a DICER variant
deleted of its C-terminal dsRNA binding domain (Fig. 5A DICER ΔdsRBD) since this domain
could also be involved in protein-protein interaction [49,50]. We then transfected the different
versions of DICER WT and the deletion mutant constructs in NoDice cells. In mock and SINV-
GFP infected cells, whole cell extracts were subjected to anti-HA and anti-MYC (CTL) IP.

TRBP was retrieved in both conditions with DICER WT, Hel. and ΔdsRBD (Fig. 5B and 5C). In mock cells, PACT and PKR were only found weakly interacting with DICER WT (Fig. 5B). In SINV-infected cells, we observed that similar to TRBP and to a lesser extent PACT, N1 and N3 mutations strongly reduced the binding of DICER with PKR (Fig. 5C lanes 2-3 and 7-8). Importantly, we also noted that the helicase domain alone could bind PKR, TRBP and PACT (Fig. 5C lanes 4 and 9). Moreover, the deletion of the dsRNA binding domain of DICER did not affect its interaction with TRBP, PACT and PKR (Fig. 5C lanes 5 and 10). We also looked for p-PKR in our co-IP (Fig. 5C panel p-PKR). We noticed that only WT DICER and its helicase domain were able to interact with p-PKR (Fig. 5C lanes 1&6 and 4&9). The fact that DICER ΔdsRBD did not interact with p-PKR (Fig. 5C lanes 5&10) is striking but could indicate that the phosphorylation of PKR may induce conformation changes preventing its interaction with some domains of DICER. These results reveal that, like for TRBP and PACT, the helicase domain of DICER is required for DICER-PKR/p-PKR interaction during SINV infection.

In order to confirm these co-IP experiments, we next decided to perform BiFC experiments using the same conditions as previously. In both mock and SINV-infected cells, only the combinations of DICER WT-PKR and DICER ΔdsRBD-PKR showed a strong Venus signal, while neither DICER N1 nor N3 constructs revealed an interaction with PKR (Fig. 5D). In contrast, the DICER Hel. construct did not seem to interact with PKR in mock-infected cells but appeared to do so in SINV-infected cells as a faint Venus signal could be observed. These results therefore confirmed the co-IP observations for the DICER-PKR interaction. In addition, we also performed a BiFC experiment using the different DICER constructs with TRBP or PACT. Altogether, the BiFC results mostly fitted with the co-IP experiment for the DICER-TRBP (Fig. S5A) and DICER-PACT (Fig. S5B) interactions. TRBP indeed did not seem to interact with the DICER N1 and only slightly with the DICER N3. However, PACT interaction was lost with DICER N1, but not with DICER N3 in mock- and SINV-infected cells (Fig. S5B third panel). This result may be explained by the fact that DICER interacts with PACT *via* the helicase and DUF domains, whereas only the DICER helicase domain is required for its interaction with TRBP [14,17]. In agreement, the Venus signal observed between the DICER Hel. and PACT seemed weaker than the one we observed with TRBP (Fig. S5A and B fourth panels).

Taken together these results indicate that DICER interacts with both PKR and its phosphorylated form during SINV infection, and that this interaction requires the helicase domain of DICER.

Functional importance of DICER helicase domain during SINV infection

We then sought to study the functional role of DICER-PKR interaction during viral infection. For this purpose, we decided to use DICER helicase deletion mutants to study SINV infection. To do so, we first generated NoDice HEK293T cells stably expressing FHA-tagged DICER N1 (FHA:DICER N1) by lentiviral transduction. As for the FHA:DICER WT cell line, we first selected a clone expressing the tagged DICER N1 at a level similar level to the endogenous DICER protein in HEK293T cells (Fig. 6A). DICER N1 protein has been shown to still be able to produce miRNAs [33]. We thus verified by northern blot analysis that DICER N1 is indeed able to process miRNAs similarly to WT DICER in HEK 293T and FHA:DICER cells, thereby validating the functionality of the tagged protein (Fig. 6B). We next infected HEK293T, FHA:DICER WT #4 and FHA:DICER N1 #6 cells with SINV-GFP and measured virus accumulation by assessing GFP expression by microscopy analysis. Interestingly, the GFP protein level was drastically reduced in FHA:DICER N1 #6 cells compared to FHA:DICER WT #4 and HEK293T cells (Fig. 6C). Encouraged by this observation, we decided to infect with SINV-GFP additional DICER deletion mutants, namely N3 and Hel. We generated stable cell lines for these various mutants by lentiviral transduction in the NoDice 2.20 background and infected those cells with SINV-GFP at an MOI of 0.02 for 24h. We verified by western blot analysis that all selected DICER mutant clones, namely N1 #6, N3 #2.13 and Hel. #2.6, expressed the tagged protein at the expected size and at levels mostly similar to the FHA:DICER WT #4 line (Fig. 6D, first two panels). We also verified the DICER mutants contribution to the endogenous miRNA biogenesis by performing a northern blot analysis on miR-16 accumulation (Fig. S6A).

We additionally verified the impact of these DICER mutants on SINV-GFP infection by measuring the GFP intensity of fluorescence by microscopy (Fig. S6B). Our results indicated that GFP accumulation is similar in HEK293T, NoDice 2.20, FHA:DICER WT, Hel. and ctrl cells. However, almost no fluorescence was detected in FHA:DICER N1 #6 and N3 #2.13 cells compared to HEK293T cells (Fig. S6B). The reduction of virus-encoded GFP accumulation and viral production were confirmed by western blot (Fig. 6D) and by plaque assay, respectively (Fig. 6E).

Altogether, these results therefore indicate that expressing a helicase truncated version of DICER, which is unable to interact with PKR, appears to confer an antiviral phenotype to SINV infection.

The antiviral phenotype of the helicase-truncated DICER mutants is independent of AGO2

We finally carried out a functional analysis of the helicase-domain-truncated DICER N1 and N3 mutants to investigate the mechanism of the antiviral phenotype. First, to investigate a potential implication of the RNAi pathway, we performed a knock-down of the AGO2 protein prior to the infection of NoDice cells expressing either WT, N1 or N3 FHA:DICER. AGO2 is the main effector protein in RNA silencing pathways [51] and has been previously shown to be a crucial antiviral RNAi factor against Influenza A virus in mouse embryonic fibroblasts (MEFs) [52]. We transfected either control siRNAs, or siRNAs targeting AGO2 for 48h in NoDice cells stably expressing either an empty vector (FHA:ctrl #2) or WT, N1 or N3 FHA:DICER constructs. Cells were then infected with SINV-GFP at an MOI of 0.02 for 24h, and virus accumulation was first assessed by looking at GFP expression by microscopy analysis (Fig. 7A). In all cell lines, no major difference in GFP fluorescence could be observed when comparing cells transfected with the control siRNA or AGO2-specific siRNAs. We verified the knock-down efficiency by western blot analysis and confirmed the microscopy observation by measuring GFP protein accumulation (Fig. 7B). Finally, we measured virus accumulation by plaque assay, and we observed that the antiviral phenotype was clearly visible in FHA:DICER N1 #6 and FHA:DICER N3 #2.13 cell lines but was not complemented upon AGO2 knock-down (Fig. 7C).

Altogether, these results indicate that the antiviral phenotype against SINV observed in cells expressing helicase-truncated mutant DICER proteins does not depend on the presence of AGO2, thereby ruling out an involvement of RNAi.

The antiviral phenotype due to the DICER helicase-domain deletion requires PKR

In order to determine the functional role of the PKR-DICER interaction in the antiviral response to SINV, we generated NoDiceΔPKR cells stably expressing either the full length FHA:DICER WT or the helicase deletion mutants FHA:DICER N1 or N3, or the empty vector as a control (FHA:ctrl) by lentiviral transduction. After monoclonal selection of each cell line, we infected them with SINV-GFP at an MOI of 0.02 and assessed virus accumulation by looking at GFP fluorescence by microscopy analysis (Fig. 8A). As expected, an increase in GFP fluorescence was observed in NoDiceΔPKR FHA:ctrl cells compared to HEK293T cells at 24 hpi. In contrast we could not observe any difference in GFP fluorescence between NoDiceΔPKR FHA:ctr cells and those expressing FHA:DICER WT, FHA:DICER N1 or N3 proteins. To verify whether any significant difference in terms of virus accumulation could be observed in NoDiceΔPKR cells

expressing WT or helicase truncated DICER proteins, we measured GFP protein levels by

western blot analysis (Fig. 8B) and virus production by plaque assay (Fig. 8C). As opposed to

the observations done in cells NoDice cells expressing PKR (Fig. 6), both GFP accumulation

and viral titers remained unchanged between NoDiceΔPKR FHA:ctrl cells and those expressing

FHA:DICER WT, N1 or N3 constructs. Taken together, these results demonstrate that the

antiviral phenotype of helicase-truncated DICER mutants depends on the presence of PKR.

Therefore, our data suggest that the helicase domain of DICER sequesters PKR and when this

interaction is lost, the antiviral effect of PKR is exacerbated, thereby explaining the phenotype

observed in cells expressing helicase-truncated DICER mutants.

Discussion

The role of DICER in antiviral defense in human cells remains a topic of intense discussion

[21,22,53,54]. In particular there have been contradictory reports regarding its capacity to

produce siRNAs from viral RNAs [31,37,55,56]. These observations could be due to the fact

that several mammalian viruses potentially encode VSR proteins, thereby masking the effect of

RNAi [22,28,29,52,57]. Another putative but non-exclusive explanation could be that there is

a mutual regulation of type I IFN and RNAi pathways [58,59]. Thus, it has already been shown

that PACT can regulate MDA5 and RIG-I during virus infection and therefore the induction of

type I IFN response [60,61]. To date, it is not clear whether the activity of the DICER protein

as well could be regulated by potential interactors, or inversely whether it could itself modulate

the activity of proteins involved in the IFN pathway. To answer this question, we determined

the changes in the interactome of human DICER upon SINV and SFV infection. This analysis

allowed us to reveal that a lot of proteins associating with DICER during viral infection are

dsRNA-binding proteins and RNA helicases. A number of these proteins are known to be

involved in antiviral defense pathways, thereby indicating the possible formation of one or

several complexes between DICER and these proteins, which are very likely brought together

by the accumulation of dsRNA during virus infection.

Among these proteins, we chose to focus on the well-known ISG PKR, which is

involved in many cellular pathways such as apoptosis, cellular differentiation, development and

antiviral defense [4,8,62,63]. PKR is one of the main actors of the Integrative Stress Response

(ISR) in human cells, and its activation or inhibition needs to be tightly regulated in order to

have a properly balanced response to stress. Our results indicate that DICER interacts via its

helicase domain with PKR in the cytoplasm during SINV infection. The helicase domain of

DICER, which is also required for its interaction with TRBP and PACT, belongs to the helicase

superfamily 2, which is also found in RLRs such as RIG-I, MDA5 or LGP2 [64,65]. These

proteins act as sensors of viral infection and through the activation of proteins such as MAVS,

mediate the induction of type I IFN pathway [65]. We hypothesize that even though the human

DICER helicase has evolved mainly to act in miRNA/siRNA pathway, it still retained the

capacity to act as an RLR. However, as opposed to RIG-I and MDA5, our data suggest that

DICER would act more as an inhibitor rather than inducer of the immune response. Therefore,

we propose that this domain serves as a platform for the recruitment of different proteins to

diversify the function of DICER.

One such regulatory effect appears to be on the antiviral activity of PKR, as cells

expressing a truncated form of DICER unable to interact with PKR become resistant to SINV

infection. This is in agreement with previous observations that ectopic expression of the

Drosophila DICER2 protein in human cells perturbs IFN signaling pathways and antagonizes

PKR-mediated antiviral immunity [36]. Although the precise molecular mechanism involved

will require further work to be fully deciphered, it seems that the two proteins are likely brought

together *via* their interaction with RNA, most probably of viral origin. Indeed, we showed that

the co-IP interaction was partially RNase sensitive. However, we confirmed that the interaction

is not artificially created during the co-immunoprecipitation procedure, since we could show

that DICER and PKR interact in BiFC assay, a technique that favors the detection of direct

interactions [47]. Most of the time, the inhibition of PKR activity relies on its inhibition to bind

to dsRNA or to auto-phosphorylate. For example, the human tRNA-dihydrouridine synthase 2

(hDus2) binds the first dsRBD of PKR and prevents its activation [66]. TRBP binds dsRNAs

but also PKR directly hindering its dimerization. In normal condition, TRBP is also associated

with PACT thus preventing PKR activation by PACT [67–70]. Since we showed that DICER

can bind the activated phospho-PKR, we hypothesize that this interaction does not result in the

inhibition of PKR autophosphorylation. In fact, in condition of infection with a high virus dose,

we showed that phospho-PKR levels are similar in cells expressing DICER WT or helicase

deletion mutants N1 and N3, but the activated PKR does not associate with these truncated

versions of DICER. Therefore, one possibility could be that DICER interaction with PKR

prevents the latter from acting upon some of its targets, which remain to be identified, to fine-

tune the antiviral response.

As of now, we cannot formally rule out that the effect of DICER on PKR is mediated

by other proteins. TRBP and PACT have been shown to regulate PKR activity, the former

normally acting as a repressor and the latter as an activator [46,68,70]. Interestingly, in

lymphocytic Jurkat cells infected by HIV-1, PACT can also act as a repressor of PKR [71]. It

is thus tempting to speculate that these two proteins participate in the formation of the DICER-

PKR complex. However, our results show that this may not necessarily be the case. Indeed, in

the BiFC experiment, the DICER N3 mutant still interacted with PACT but not with PKR

indicating that PACT binding is not sufficient to confer the association with PKR.

Besides PKR, other proteins were specifically enriched upon viral infection in the

DICER IP. These are also interesting candidates to explain the putative regulatory role of

DICER. Among these proteins, DHX9 and ADAR-1 are especially intriguing. DHX9, also

known as RNA helicase A (RHA), associates with RISC, helping the RISC loading [41].

Moreover, DHX9 is directly involved in removing toxic dsRNAs from the cell to prevent their

processing by DICER [40]. It has also been implicated in HIV-1 replication and knockdown of

DXH9 leads to the production of less infectious HIV-1 virions [72–74]. Finally, DXH9 interacts

with and is phosphorylated by PKR in MEFs. This phosphorylation precludes the association

of DHX9 with RNA, thus inhibiting its proviral effect [75]. In light of these observations and

ours, we can speculate that the inhibitory effect of DICER on PKR activity could also be linked

to DHX9 phosphorylation. ADAR-1 is one of the well-known RNA-editing factors [76].

ADAR-1 is linked to both miRNA biogenesis [77–79] and virus infection. Indeed, ADAR-1

has an antiviral effect against Influenza virus, but most of the time, its depletion leads to a

decrease of the viral titer, as was reported for VSV or HIV-1 [80,81]. It has been shown that

ADAR-1 and PKR interact directly during HIV-1 infection. This interaction triggers the

inhibition of PKR activation, and thus a reduction of eIF2α phosphorylation leading to an

increase of virus replication [5,82]. Interestingly, over-expression of ADAR-1 enhances

drastically the replication of the alphaviruses Chikungunya virus (CHIKV), and Venezuelan

equine encephalitis virus (VEEV) most likely by interfering with the IFN induction [83].

One hypothesis to explain the virus resistance phenotype of the DICER N1 and N3 cell

lines could be an increased processivity of these truncated proteins on long dsRNA substrates

[33], which would render DICER RNAi proficient. However, our results are not in favor of this

hypothesis, since we show that knocking-down AGO2 does not allow to make cells expressing

DICER N1 or N3 more sensitive to SINV infection. AGO2 being the only slicer-proficient

Argonaute protein expressed at physiological levels in HEK293T cells, we can confidently

conclude that the observed phenotype is RNAi-independent.

Finally, we demonstrated that the phenotype of helicase-truncated DICER isoforms

depends on PKR expression, because it was completely lost in PKR knockout cell lines. We

therefore propose that, at least during infection with SINV, DICER prevents PKR to be fully active by interacting with and potentially sequestrating it. Deciphering the exact molecular mechanism at play will require additional studies in order to get the full picture. Nevertheless, by assessing the interactome of DICER during SINV infection, we have unveiled a new, PKR-dependent, role for the helicase domain of DICER in regulating the cellular response to viral infection.

Material and methods

Plasmids, cloning and mutagenesis

Plasmids used for BiFC experiments were a gift from Dr. Oliver Vugrek from the Ruđer Bošković Institute and described in [47]. The cDNAs of TRBP, PACT and PKR were respectively amplified from (pcDNA-TRBP Addgene #15666) [16], (pcDNA-PACT Addgene #15667) [16], (pSB819-PKR-hum Addgene #20030) [84], and cloned into the four pBiFC vectors by Gateway recombination. DICER N1, N3 Hel. and ΔdsRBD were generated by PCR mutagenesis from pDONR-DICER described in [36] and cloned into the four pBiFC and pDEST-FHA vectors by Gateway recombination. plenti6 FHA-V5 vector was modified from plenti6-V5 gateway vector (Thermo Fisher scientific V49610) by Gibson cloning. DICER WT, N1, N3 and Hel. from pDONR plasmids were cloned into plenti6 FHA-V5 by Gateway recombination. All primers used are listed in Supp. Table 5.

Cell lines

HEK293T, HEK293T/NoDice (2.20 and 4.25), and HEK293T/NoDiceΔPKR cell lines were a gift from Pr. Bryan Cullen and described in [33,39]. HCT116 cell line was a gift from Dr. Christian Gaiddon.

Generation of Flag-HA-GFP-DICER knock-in cell line by CRISPR/Cas9

To generate the knock-in cell line, the sequence of Flag-HA-GFP was amplified by PCR from the Flag-HA-GFP plasmid [85]. DNA sequences corresponding to 1 Kb upstream (left homology arm) and downstream (right homology arm) the starting codon (ATG) of DICER gene were amplified from HCT116 cell genomic DNA using primer pairs listed in Supp. Table 5. The three PCR products were gel-purified and cloned into a linearized pUC19 by In-fusion cloning (Clontech) to obtain the template for homologous recombination (LarmDICER-FlagHAGFP-RarmDICER).

Design of the guide RNA targeting the region between Dicer 5'-UTR and its first coding exon

for CRISPR/Cas9 mediated knock-in was carried out using the CRISPOR Design Tool [86].

Annealed oligonucleotide corresponding to the gRNA (Supp. Table 5) were cloned into the

vector pX459 (Addgene #48139) which also encodes *S. pyogenes* Cas9 with 2A-Puro.

The sequence of the donor plasmid was additionally mutagenized to disrupt the PAM sequence

of the right homology arm to avoid its cleavage by the gRNA.

To obtain the knock-in (KI) cell line, 5×10^5 HCT116 cells were seeded in a 6 well plate with

Dulbecco's modified Eagle medium (DMEM, Gibco®, Life Technologies) supplemented with

10% fetal bovine serum (FBS, Clontech) in a humidified atmosphere of 5% CO_2 at 37°C and

transfected after 24 hours with the pX459-gRNADicerNterm-Cas9-2A-Puro plasmid and the

Leftarm-FlagHAGFP-RightarmDICER donor plasmids at the ratio of 1 to 1 (6 micrograms

plasmids in total) using Lipofectamine 2000 according to the manufacturer instructions. 24

hours later, puromycin (1mg/mL) was added to the cells to increase the KI efficiency and

genomic DNA was isolated from individual colonies few days later.

The presence of the Flag-HA-GFP tag in frame with hDICER coding sequence was confirmed

by sequencing PCR amplicon from KI cell gDNA. Expression of Flag-HA-GFP N-terminal

tagged Dicer protein in the KI cells was confirmed by western blot.

Cell culture and transfection

Cells were maintained in Dulbecco's modified Eagle medium (DMEM, Gibco®, Life

Technologies) supplemented with 10% fetal bovine serum (FBS, Clontech) in a humidified

atmosphere of 5% CO_2 at 37°C. Transfection was performed using Lipofectamine 2000

(Invitrogen, ThermoFisher Scientific) according to the manufacturer's instructions.

Lentivirus production and generation of stable cell lines

The lentiviral supernatant from single transfer vector was produced by transfecting HEK293T

cells (ATCC® CRL-3216™) with 20 μg of the transfer vector, 15 μg of pMDLg/p RRE and 10

μg of pRSV-Rev packaging plasmids (Addgene #12251 and Addgene #12253) and the pVSV

envelope plasmid (Addgene #8454) using Lipofectamine 2000 (Invitrogen, ThermoFisher

Scientific) reagent according to the manufacturer's protocol. Standard DMEM (Gibco®, Life

Technologies) medium supplemented with 10% Fetal bovine serum (FBS, Gibco®, Life

Technologies) and 100 U/mL of penicillin-Streptomycin (Gibco®, Life Technologies) was

used for growing HEK293T cells and for lentivirus production. One 10 cm plate of HEK293T

cells at 70-80% confluency was used for the transfection. The medium was replaced 8 hours

post-transfection. After 48 hours the medium containing viral particles was collected and

filtered through a 0.45μm PES filter. The supernatant was directly used for transfection or

stored at -80°C. A 6 well plate of HEK293T/NoDice or HEK293T/NoDiceΔPKR cells at 80%

confluency was transduced using 600 μL of lentiviral supernatant either expressing

FHA:DICER, N1, N3, Hel. or empty vector, supplemented with 4 ug/mL polybrene (Sigma)

for 6 hours. The transduction media was then changed with fresh DMEM for 24 hours and the

resistant cell clones were selected for about 6 weeks with blasticidin (15 μg/mL for NoDice or

10 μg/mL for NoDiceΔPKR) and subsequently maintained under blasticidin selection.

Viral stocks, virus infection

Viral stocks of SINV or SINV-GFP were produced as described in [36]. Cells were infected

with SINV or SINV-GFP at an MOI of 0.02, 0.1, 1 or 2 and samples were collected at different

time points as indicated in the figure legends.

Analysis of viral titer by plaque assay

Vero R cells were seeded in 96-well plates format and were infected with 10-fold serial

dilutions infection supernatants for 1 hour. Afterwards, the inoculum was removed, and cells

were cultured in 2.5% carboxymethyl cellulose for 72 hours at 37°C in a humidified atmosphere

of 5% CO_2. Plaques were counted manually under the microscope and viral titer was calculated

according to the formula: *PFU/mL = #plaques/ (Dilution*Volume of inoculum)*

Western blot analysis

Proteins were extracted from cells and homogenized in 350 μL of lysis buffer (50 mM Tris-

HCl pH 7.5, 150 mM NaCl, 5 mM EDTA, 1% Triton X-100, 0.5% SDS and Protease Inhibitor

Cocktail (complete Mini; Sigma Aldrich)). Proteins were quantified by the Bradford method

and 20 to 30 μg of total protein extract were loaded on 4-20% Mini-PROTEAN® TGX™

Precast Gels (Bio-Rad). After transfer onto nitrocellulose membrane, equal loading was verified

by Ponceau staining. For PVDF membrane, equal loading was verified by Coomassie staining

after transfer and blotting. Membranes were blocked in 5% milk and probed with the following

antibodies: anti-hDicer (1:500, F10 Santa Cruz sc-136979) and anti-hDicer (1:1000, A301-

937A Bethyl), anti-TRBP (1:500, D-5 Santa Cruz sc-514124), anti-PKR (1:2500, Abcam

ab32506), anti-PACT (1:500, Abcam, ab75749), anti-HA (1:10000, Sigma, H9658), anti-

DHX9 (1:500, Abcam ab26271), anti-p-eIF2a (1:1000, Ser-52 Santa Cruz sc-601670), anti-

hADAR-1 (1:500 Santa Cruz sc-271854) anti-p-PKR (1:1000 Abcam ab81303) anti-GFP (1:10000, Roche 11814460001) and anti-Tubulin (1:10000, Sigma T6557). Detection was performed using Chemiluminescent Substrate (Pierce, ThermoFisher Scientific) and visualized on a Fusion FX imaging system (Vilber).

RNA extraction and northern blot analysis

Total RNA was extracted using Tri-Reagent Solution (Fisher Scientific; MRC, Inc) according to the manufacturer's instructions. Northern blotting was performed on 10 μg of total RNA. RNA was resolved on a 12% urea-acrylamide gel, transferred onto Hybond-NX membrane (GE Healthcare). RNAs were then chemically cross-linked to the membrane during 90 min at 65°C using 1-ethyl-3-[3-dimethylaminopropyl]carbodiimide hydrochloride (EDC) (Sigma Aldrich). Membranes were prehybridized for 30 min in PerfectHyb™ plus (Sigma Aldrich) at 50°C. Probes consisting of oligodeoxyribonucleotides (see Supplementary Table 2) were 5′-end labeled using T4 polynucleotide kinase (ThermoFisher Scientific) with 25 μCi of [γ-32P]dATP. The labeled probe was hybridized to the blot overnight at 50°C. The blot was then washed twice at 50°C for 20 min (5× SSC/0.1% SDS), followed by an additional wash (1× SSC/0.1% SDS) for 5 min. Northern blots were exposed to phosphorimager plates and scanned using a Bioimager FLA-7000 (Fuji).

Immunoprecipitation

Immunoprecipitation experiments were carried out either on tagged proteins or on endogenous proteins.

Tagged proteins: Cells were harvested, washed twice with ice-cold 1× PBS (Gibco®, Life Technologies), and resuspended in 550 μL of lysis buffer (50 mM Tris-HCl pH 7.5, 140 mM NaCl, 1.5 mM MgCl$_2$, 0.1% NP-40), supplemented with Complete-EDTA-free Protease Inhibitor Cocktail (complete Mini; Sigma Aldrich). Cells were lysed by 30 min incubation on ice and debris were removed by 15 min centrifugation at 2000 g and 4°C. An aliquot of the cleared lysates (50 μL) was kept aside as protein Input. Samples were divided into equal parts (250 μL each) and incubated with 15 μL of magnetic microparticles coated with monoclonal HA or MYC antibodies (MACS purification system, Miltenyi Biotech) at 4°C for 1 hour under rotation (10 rpm). Samples were passed through μ Columns (MACS purification system, Miltenyi Biotech). The μ Columns were then washed 3 times with 200 μL of lysis buffer and 1 time with 100 μL of washing buffer (20 mM Tris-HCl pH 7.5). To elute the immunoprecipitated

proteins, 95°C pre-warmed 2x Western blot loading buffer (10% glycerol, 4% SDS, 62.5 mM Tris-HCl pH 6.8, 5% (v/v) 2-β-mercaptoethanol, Bromophenol Blue) was passed through the μ Columns. Proteins were analyzed by western blotting or by mass spectrometry.

Endogenous proteins: mock or SINV-GFP-infected HEK293T cells (MOI of 2) were lysed 6 hours post-infection using immunoprecipitation buffer (50mM tris-HCl [pH 7.5], 150mM NaCl, 5mM EDTA, 0.05% SDS, 1% triton) supplemented with Complete-EDTA-free Protease Inhibitor Cocktail (complete Mini; Sigma Aldrich). Lysates were treated for 20 min at 37°C with 1μL of DNase I (ThermoFisher Scientific) using its buffer (10mM $MgCl_2$, 5mM $CaCl_2$ and 1μL of ribolock). Lysates were cleared by centrifugation at 16000 g, 10 min at 4°C. Supernatants were precleared 1h at room temperature with magnetic beads blocked with BSA (ThermoFisher Scientific) to avoid aspecific binding. Lysates were incubated overnight on wheel at 4°C with immunoprecipitation buffer containing magnetic Protein A DynaBeads (Invitrogen, ThermoFisher Scientific) conjugated with human PKR antibody (Abcam) or negative control rabbit IgG (Cell signaling, Ozyme). Beads were washed 3 times with immunoprecipitation buffer, 3 times with wash buffer (50mM tris-HCl [pH 7.5], 200mM NaCl, 5mM EDTA, 0.05% SDS, 1% triton, supplemented with Complete-EDTA-free Protease Inhibitor Cocktail (complete Mini; Sigma Aldrich)) and twice with cold PBS 1X (Gibco®, Life Technologies). Beads were eluted with 2x western blot loading buffer and incubated for 10 min at 95°C under agitation. Proteins were analyzed by western blotting.

RNase treatment followed by co-IP

On tagged proteins: Cells were harvested, washed twice with ice-cold 1× PBS (Gibco®, Life Technologies), and resuspended in 550 μL of lysis buffer (50 mM Tris-HCl pH 7.5, 140 mM NaCl, 1.5 mM $MgCl_2$, 0.1% NP-40), supplemented with Complete-EDTA-free Protease Inhibitor Cocktail (complete Mini; Sigma Aldrich). Cells were lysed by 30 min incubation on ice and debris were removed by 15 min centrifugation at 2000 g and 4°C. Lysate was treated or not with RNase A/T1 mix (ThermoFisher Scientific) and place at 37°C 30min. An aliquot of the cleared lysates (25 μL) was kept aside as protein Input and another aliquot (25 μL) was kept to assess RNase treatment efficiency. Co-IP was led as previously described.

Total RNA was extracted using Tri-Reagent Solution (Fisher Scientific; MRC, Inc) according to the manufacturer's instructions. RNA integrity upon treatment was verified on an 1% agarose gel containing ethidium bromide 10 mg/mL (Invitrogen, ThermoFisher Scientific) and revealed under UV on Gel Doc™ EZ system (Bio-Rad).

siRNA transfection

20nM of human AGO2 or non-targeting control siRNA (Horizon discovery) were transfected in 130 000 NoDice FHA:ctrl #2, NoDice FHA:DICER WT#4, N1 #6 or N3 #2.13 cells using Lipofectamine 2000 transfection reagent (Invitrogen, ThermoFisher Scientific) according to the manufacturer's instructions. After 24 hours, the cells were again transfected with 20nM of the same siRNA and incubated overnight. Cells were infected or not with SINV-GFP at an MOI of 0.02 for 24h. Proteins and supernatants were collected and analyzed by western blotting and plaque assay, respectively.

BiFC assay

Experiments were carried out in two different ways. For non-fixed cells, NoDiceΔPKR or HEK293T cells were seeded at the density of 1.2×10^5 cells per well in a 24-well plate. After 16 hours, cells were transfected with equimolar quantities of each plasmid forming BiFC couples. After 24 hours, cells were infected with SINV at an MOI of 2 and pictures were taken 6 hours post-infection using ZOE fluorescent cell imager (Bio-Rad). Proteins were collected with lysis buffer (50 mM Tris-HCl pH 7.5, SDS 0.05%, Triton 1%, 5 mM EDTA, 150 mM NaCl) supplemented with Complete-EDTA-free Protease Inhibitor Cocktail (complete Mini; Sigma Aldrich), and subjected to western blot analysis. For fixed cells, NoDice/ΔPKR cells were seeded at the density of 8.10^4 cells per well in 8-well Millicell® EZ Slides (Merck Millipore), transfected and infected as described previously. At 6 hours post-infection, cells were fixed with 4% formaldehyde and 0.2% glutaraldehyde for 10 min. Cells were then washed with $1\times$ PBS (Gibco®, Life Technologies) and stained with 10 μg/μL DAPI (Invitrogen, ThermoFisher Scientific) in $1\times$ PBS solution (Invitrogen, ThermoFisher Scientific) for 5 min. Fixed cells were mounted on a glass slide with Fluoromount-G mounting media (Southern Biotech). Images were acquired using confocal LSM780 (Zeiss) inverted microscope with an argon laser (514x nm) and with $\times40$ immersion oil objective. All pictures obtained from BiFC experiments were treated using FigureJ software (NIH).

Mass spectrometry analysis

Protein extracts were prepared for mass spectrometry as described in a previous study [87]. Each sample was precipitated with 0.1 M ammonium acetate in 100% methanol, and proteins were resuspended in 50 mM ammonium bicarbonate. After a reduction-alkylation step (dithiothreitol 5 mM – iodoacetamide 10 mM), proteins were digested overnight with

sequencing-grade porcine trypsin (1:25, w/w, Promega, Fitchburg, MA, USA). The resulting

vacuum-dried peptides were resuspended in water containing 0.1% (v/v) formic acid (solvent

A). One sixth of the peptide mixtures were analyzed by nanoLC-MS/MS an Easy-nanoLC-1000

system coupled to a Q-Exactive Plus mass spectrometer (Thermo-Fisher Scientific, USA)

operating in positive mode. Five microliters of each sample were loaded on a C-18 precolumn

(75 μm ID × 20 mm nanoViper, 3 μm Acclaim PepMap; Thermo) coupled with the analytical

C18 analytical column (75 μm ID × 25 cm nanoViper, 3 μm Acclaim PepMap; Thermo).

Peptides were eluted with a 160 min gradient of 0.1% formic acid in acetonitrile at 300 nL/min.

The Q-Exactive Plus was operated in data-dependent acquisition mode (DDA) with Xcalibur

software (Thermo-Fisher Scientific). Survey MS scans were acquired at a resolution of 70K at

200 m/z (mass range 350-1250), with a maximum injection time of 20 ms and an automatic

gain control (AGC) set to 3e6. Up to 10 of the most intense multiply charged ions (≥2) were

selected for fragmentation with a maximum injection time of 100 ms, an AGC set at 1e5 and a

resolution of 17.5K. A dynamic exclusion time of 20 s was applied during the peak selection

process.

Database search and mass-spectrometry data post-processing

Data were searched against a database containing Human and Viruses UniProtKB sequences

with a decoy strategy (GFP, Human and Sindbis Virus SwissProt sequences as well as Semliki

Forest Virus SwissProt and TrEMBL sequences (releases from January 2017, 40439

sequences)). Peptides were identified with Mascot algorithm (version 2.3, Matrix Science,

London, UK) with the following search parameters: carbamidomethylation of cysteine was set

as fixed modification; N-terminal protein acetylation, phosphorylation of serine / threonine /

tyrosine and oxidation of methionine were set as variable modifications; tryptic specificity with

up to three missed cleavages was used. The mass tolerances in MS and MS/MS were set to 10

ppm and 0.02 Da respectively, and the instrument configuration was specified as "ESI-Trap".

The resulting .dat Mascot files were then imported into Proline v1.4 package

(http://proline.profiproteomics.fr) for post-processing. Proteins were validated with Mascot

pretty rank equal to 1, 1% FDR on both peptide spectrum matches (PSM) and protein sets

(based on score). The total number of MS/MS fragmentation spectra (Spectral count or SpC)

was used for subsequent protein quantification in the different samples.

Exploratory and differential expression analysis of LC-MS/MS data

Mass spectrometry data obtained for each sample were stored in a local MongoDB database and subsequently analyzed through a Shiny Application built upon the R/Bioconductor packages msmsEDA (Gregori J, Sanchez A, Villanueva J (2014). msmsEDA: Exploratory Data Analysis of LC-MS/MS data by spectral counts. R/Bioconductor package version 1.22.0) and msmsTests (Gregori J, Sanchez A, Villanueva J (2013). msmsTests: LC-MS/MS Differential Expression Tests. R/Bioconductor package version 1.22.0). Exploratory data analyses of LC-MS/MS data were thus conducted, and differential expression tests were performed using a negative binomial regression model. The p-values were adjusted with FDR control by the Benjamini-Hochberg method and the following criteria were used to define differentially expressed proteins: an adjusted p-value < 0.05, a minimum of 5 SpC in the most abundant condition, and a minimum fold change of 2 (abs(LogFC) > 1). GO term analysis was performed using the EnrichR web-based tool (http://amp.pharm.mssm.edu/Enrichr). The direct interaction network for proteins enriched in SINV-infected cells was generated using the STRING database (https://string-db.org).

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE [88] partner repository with the dataset identifier PXD019093 and 10.6019/PXD019093.

Acknowledgments

The authors would like to thank members of the Pfeffer laboratory for discussion, Pr. Bryan Cullen for the kind gift of the HEK293T NoDice and NoDice/ΔPKR cell lines and Dr. Oliver Vugrek for the BiFC plasmids.

This work was funded by the European Research Council (ERC-CoG-647455 RegulRNA) and was performed under the framework of the LABEX: ANR-10-LABX-0036_NETRNA and ANR-17-EURE-0023, which benefits from a funding from the state managed by the French National Research Agency as part of the Investments for the future program. This work has also received funding from the People Programme (Marie Curie Actions) of the European Union's Seventh Framework Program (FP7/2007-2013) under REA grant agreement n° PCOFUND-GA-2013-609102, through the PRESTIGE program coordinated by Campus France (to EG), and from the French Minister for Higher Education, Research and Innovation

806 (PhD contract to MB). The mass spectrometry instrumentation was funded by the University
807 of Strasbourg, IdEx "Equipement mi-lourd" 2015.
808

Author contributions

SP and ML conceived the project. TCM, ML, EG and SP designed the work. TCM, MB, ML,
EG and MM performed the experiments and analyzed the results. PH and JC performed the
mass spectrometry analysis. BCWM designed software and analyzed the mass spectrometry
results. SP coordinated the work and assured funding. TCM, MB and SP wrote the manuscript
with input from the other authors. All authors reviewed and approved the final manuscript.
815

References

1. Ivashkiv LB, Donlin LT. Regulation of type I interferon responses. Nat Rev Immunol.
2014;14: 36–49. doi:10.1038/nri3581
2. Borden EC, Sen GC, Uze G, Silverman RH, Ransohoff RM, Foster GR, et al. Interferons
at age 50: past, current and future impact on biomedicine. Nat Rev Drug Discov. 2007;6:
975–990. doi:10.1038/nrd2422
3. Williams BR. PKR; a sentinel kinase for cellular stress. Oncogene. 1999;18: 6112–6120.
doi:10.1038/sj.onc.1203127
4. Lemaire PA, Anderson E, Lary J, Cole JL. Mechanism of PKR Activation by dsRNA. J
Mol Biol. 2008;381: 351–360. doi:10.1016/j.jmb.2008.05.056
5. Pfaller CK, Li Z, George CX, Samuel CE. Protein kinase PKR and RNA adenosine
deaminase ADAR1: new roles for old players as modulators of the interferon response.
Curr Opin Immunol. 2011;23: 573–582. doi:10.1016/j.coi.2011.08.009
6. Fros JJ, Pijlman GP. Alphavirus Infection: Host Cell Shut-Off and Inhibition of Antiviral
Responses. Viruses. 2016;8: 166. doi:10.3390/v8060166
7. Ryman KD, White LJ, Johnston RE, Klimstra WB. Effects of PKR/RNase L-Dependent
and Alternative Antiviral Pathways on Alphavirus Replication and Pathogenesis. Viral
Immunol. 2002;15: 53–76. doi:10.1089/088282402317340233
8. Kim Y, Lee JH, Park J-E, Cho J, Yi H, Kim VN. PKR is activated by cellular dsRNAs
during mitosis and acts as a mitotic regulator. Genes Dev. 2014;28: 1310–1322.
doi:10.1101/gad.242644.114
9. Meister G, Tuschl T. Mechanisms of gene silencing by double-stranded RNA. Nature.
2004;431: 343–9. doi:10.1038/nature02873
10. Hutvágner G, McLachlan J, Pasquinelli AE, Balint E, Tuschl T, Zamore PD. A cellular
function for the RNA-interference enzyme Dicer in the maturation of the let-7 small
temporal RNA. Science. 2001;293: 834–8. doi:10.1126/science.1062961
11. Wienholds E, Koudijs MJ, Van Eeden FJ, Cuppen E, Plasterk RH. The microRNA-
producing enzyme Dicer1 is essential for zebrafish development. Nat Genet. 2003; 217–
218. doi:10.1038/ng1251
12. Bartel DP. Metazoan MicroRNAs. Cell. 2018;173: 20–51. doi:10.1016/j.cell.2018.03.006
13. Chendrimada TP, Gregory RI, Kumaraswamy E, Norman J, Cooch N, Nishikura K, et al.
TRBP recruits the Dicer complex to Ago2 for microRNA processing and gene silencing.
Nature. 2005;436: 740–4. doi:10.1038/nature03868
14. Daniels SM, Melendez-Peña CE, Scarborough RJ, Daher A, Christensen HS, El Far M, et
al. Characterization of the TRBP domain required for dicer interaction and function in
RNA interference. BMC Mol Biol. 2009;10: 38. doi:10.1186/1471-2199-10-38
15. Heyam A, Lagos D, Plevin M. Dissecting the roles of TRBP and PACT in double-stranded
RNA recognition and processing of noncoding RNAs. WIREs RNA. 2015;6: 271–289.
doi:10.1002/wrna.1272
16. Kok KH, Ng MH, Ching YP, Jin DY. Human TRBP and PACT Directly Interact with Each
Other and Associate with Dicer to Facilitate the Production of Small Interfering RNA. J
Biol Chem. 2007;282: 17649–57. doi:10.1074/jbc.M611768200
17. Lee Y, Hur I, Park SY, Kim YK, Suh MR, Kim VN. The role of PACT in the RNA
silencing pathway. Embo J. 2006;25: 522–32. doi:10.1038/sj.emboj.7600942
18. Guo Z, Li Y, Ding S-W. Small RNA-based antimicrobial immunity. Nat Rev Immunol.
2019;19: 31–44. doi:10.1038/s41577-018-0071-x
19. tenOever BR. The Evolution of Antiviral Defense Systems. Cell Host Microbe. 2016;19:
142–149. doi:10.1016/j.chom.2016.01.006
20. Cullen BR. Is RNA interference involved in intrinsic antiviral immunity in mammals? Nat

Immunol. 2006;7: 563–7. doi:10.1038/ni1352

21. Maillard PV, van der Veen AG, Poirier EZ, Reis e Sousa C. Slicing and dicing viruses:
antiviral RNA interference in mammals. EMBO J. 2019;38.
doi:10.15252/embj.2018100941

22. Maillard PV, Ciaudo C, Marchais A, Li Y, Jay F, Ding SW, et al. Antiviral RNA
interference in mammalian cells. Science. 2013;342: 235–238.
doi:10.1126/science.1241930

23. Maillard PV, Veen AGV der, Deddouche-Grass S, Rogers NC, Merits A, Sousa CR e.
Inactivation of the type I interferon pathway reveals long double-stranded RNA-mediated
RNA interference in mammalian cells. EMBO J. 2016;35: 2505–2518.
doi:10.15252/embj.201695086

24. Takahashi T, Nakano Y, Onomoto K, Murakami F, Komori C, Suzuki Y, et al. LGP2 virus
sensor regulates gene expression network mediated by TRBP-bound microRNAs. Nucleic
Acids Res. 2018;46: 9134–9147. doi:10.1093/nar/gky575

25. Takahashi T, Nakano Y, Onomoto K, Yoneyama M, Ui-Tei K. Virus Sensor RIG-I
Represses RNA Interference by Interacting with TRBP through LGP2 in Mammalian
Cells. Genes. 2018;9: 511. doi:10.3390/genes9100511

26. van der Veen AG, Maillard PV, Schmidt JM, Lee SA, Deddouche-Grass S, Borg A, et al.
The RIG-I-like receptor LGP2 inhibits Dicer-dependent processing of long double-
stranded RNA and blocks RNA interference in mammalian cells. EMBO J. 2018;37:
e97479. doi:10.15252/embj.201797479

27. Adiliaghdam F, Basavappa M, Saunders TL, Harjanto D, Prior JT, Cronkite DA, et al. A
Requirement for Argonaute 4 in Mammalian Antiviral Defense. Cell Rep. 2020;30: 1690-
1701.e4. doi:10.1016/j.celrep.2020.01.021

28. Qiu Y, Xu Y, Zhang Y, Zhou H, Deng Y-Q, Li X-F, et al. Human Virus-Derived Small
RNAs Can Confer Antiviral Immunity in Mammals. Immunity. 2017;46: 992-1004.e5.
doi:10.1016/j.immuni.2017.05.006

29. Qiu Y, Xu Y-P, Wang M, Miao M, Zhou H, Xu J, et al. Flavivirus induces and antagonizes
antiviral RNA interference in both mammals and mosquitoes. Sci Adv. 2020;6: eaax7989.
doi:10.1126/sciadv.aax7989

30. Girardi E, Chane-Woon-Ming B, Messmer M, Kaukinen P, Pfeffer S. Identification of
RNase L-dependent, 3'-end-modified, viral small RNAs in Sindbis virus-infected
mammalian cells. mBio. 2013;4: e00698-00613. doi:10.1128/mBio.00698-13

31. Parameswaran P, Sklan E, Wilkins C, Burgon T, Samuel MA, Lu R, et al. Six RNA viruses
and forty-one hosts: viral small RNAs and modulation of small RNA repertoires in
vertebrate and invertebrate systems. PLoS Pathog. 2010;6: e1000764.
doi:10.1371/journal.ppat.1000764

32. Schuster S, Overheul GJ, Bauer L, van Kuppeveld FJM, van Rij RP. No evidence for viral
small RNA production and antiviral function of Argonaute 2 in human cells. Sci Rep.
2019;9: 1–11. doi:10.1038/s41598-019-50287-w

33. Kennedy EM, Whisnant AW, Kornepati AVR, Marshall JB, Bogerd HP, Cullen BR.
Production of functional small interfering RNAs by an amino-terminal deletion mutant of
human Dicer. Proc Natl Acad Sci. 2015;112: E6945–E6954.
doi:10.1073/pnas.1513421112

34. Tsai K, Courtney DG, Kennedy EM, Cullen BR. Influenza A virus-derived siRNAs
increase in the absence of NS1 yet fail to inhibit virus replication. RNA N Y N. 2018.
doi:10.1261/rna.066332.118

35. Griffin D. Alphaviruses, p 1023–1067. Fields Virol 5th Ed Lippincott Williams Wilkins
Phila PA. 2007.

36. Girardi E, Lefèvre M, Chane-Woon-Ming B, Paro S, Claydon B, Imler J-L, et al. Cross-

species comparative analysis of Dicer proteins during Sindbis virus infection. Sci Rep.
2015;5: 10693. doi:10.1038/srep10693

37. Bogerd HP, Skalsky RL, Kennedy EM, Furuse Y, Whisnant AW, Flores O, et al.
Replication of many human viruses is refractory to inhibition by endogenous cellular
microRNAs. J Virol. 2014;88: 8065–8076. doi:10.1128/JVI.00985-14

38. López P, Girardi E, Mounce BC, Weiss A, Chane-Woon-Ming B, Messmer M, et al. High-
Throughput Fluorescence-Based Screen Identifies the Neuronal MicroRNA miR-124 as a
Positive Regulator of Alphavirus Infection. J Virol. 2020;94. doi:10.1128/JVI.02145-19

39. Bogerd HP, Whisnant AW, Kennedy EM, Flores O, Cullen BR. Derivation and
characterization of Dicer- and microRNA-deficient human cells. RNA. 2014;20: 923–937.
doi:10.1261/rna.044545.114

40. Aktaş T, Avşar Ilık İ, Maticzka D, Bhardwaj V, Pessoa Rodrigues C, Mittler G, et al. DHX9
suppresses RNA processing defects originating from the Alu invasion of the human
genome. Nature. 2017;544: 115–119. doi:10.1038/nature21715

41. Robb GB, Rana TM. RNA helicase A interacts with RISC in human cells and functions in
RISC loading. Mol Cell. 2007;26: 523–537. doi:10.1016/j.molcel.2007.04.016

42. Sorrentino S, Naddeo M, Russo A, D'Alessio G. Degradation of double-stranded RNA by
human pancreatic ribonuclease: crucial role of noncatalytic basic amino acid residues.
Biochemistry. 2003;42: 10182–10190. doi:10.1021/bi030040q

43. Edy VG, Szekely M, Loviny T, Dreyer C. Action of nucleases on double-stranded RNA.
Eur J Biochem. 1976;61: 563–572. doi:10.1111/j.1432-1033.1976.tb10051.x

44. García MA, Meurs EF, Esteban M. The dsRNA protein kinase PKR: Virus and cell control.
Biochimie. 2007;89: 799–811. doi:10.1016/j.biochi.2007.03.001

45. Haase AD, Jaskiewicz L, Zhang H, Laine S, Sack R, Gatignol A, et al. TRBP, a regulator
of cellular PKR and HIV-1 virus expression, interacts with Dicer and functions in RNA
silencing. EMBO Rep. 2005;6: 961–7. doi:10.1038/sj.embor.7400509

46. Patel CV, Handy I, Goldsmith T, Patel RC. PACT, a Stress-modulated Cellular Activator
of Interferon-induced Double-stranded RNA-activated Protein Kinase, PKR. J Biol Chem.
2000;275: 37993–37998. doi:10.1074/jbc.M004762200

47. Lepur A, Kovačević L, Belužić R, Vugrek O. Combining unique multiplex gateway
cloning and bimolecular fluorescence complementation (BiFC) for high-throughput
screening of protein–protein interactions. J Biomol Screen. 2016;21: 1100–1111.
doi:10.1177/1087057116659438

48. Bernstein E, Caudy AA, Hammond SM, Hannon GJ. Role for a bidentate ribonuclease in
the initiation step of RNA interference. Nature. 2001;409: 363–6. doi:10.1038/35053110

49. Doyle M, Badertscher L, Jaskiewicz L, Güttinger S, Jurado S, Hugenschmidt T, et al. The
double-stranded RNA binding domain of human Dicer functions as a nuclear localization
signal. RNA N Y N. 2013;19: 1238–1252. doi:10.1261/rna.039255.113

50. Lambert M, Pépin G, Peralta-Zaragoza O, Matusiak R, Ly S, Landry P, et al. TWEAK
Negatively Regulates Human Dicer. Non-Coding RNA. 2016;2.
doi:10.3390/ncrna2040012

51. Meister G, Landthaler M, Patkaniowska A, Dorsett Y, Teng G, Tuschl T. Human
Argonaute2 mediates RNA cleavage targeted by miRNAs and siRNAs. Mol Cell. 2004;15:
185–197. doi:10.1016/j.molcel.2004.07.007

52. Li Y, Basavappa M, Lu J, Dong S, Cronkite DA, Prior JT, et al. Induction and suppression
of antiviral RNA interference by influenza A virus in mammalian cells. Nat Microbiol.
2016;2: 16250. doi:10.1038/nmicrobiol.2016.250

53. Cullen BR, Cherry S, tenOever BR. Is RNA Interference a Physiologically Relevant Innate
Antiviral Immune Response in Mammals? Cell Host Microbe. 2013;14: 374–378.
doi:10.1016/j.chom.2013.09.011

54. Cullen BR. RNA Interference in Mammals: The Virus Strikes Back. Immunity. 2017;46:
970–972. doi:10.1016/j.immuni.2017.05.004

55. Donaszi-Ivanov A, Mohorianu I, Dalmay T, Powell PP. Small RNA Analysis in Sindbis
Virus Infected Human HEK293 Cells. Adelman ZN, editor. PLoS ONE. 2013;8: e84070.
doi:10.1371/journal.pone.0084070

56. Backes S, Langlois RA, Schmid S, Varble A, Shim JV, Sachs D, et al. The Mammalian
Response to Virus Infection Is Independent of Small RNA Silencing. Cell Rep. 2014;8:
114–125. doi:10.1016/j.celrep.2014.05.038

57. Li Y, Lu J, Han Y, Fan X, Ding S-W. RNA interference functions as an antiviral immunity
mechanism in mammals. Science. 2013;342: 231–234. doi:10.1126/science.1241911

58. Berkhout B. RNAi-mediated antiviral immunity in mammals. Curr Opin Virol. 2018;32:
9–14. doi:10.1016/j.coviro.2018.07.008

59. Takahashi T, Ui-Tei K. Mutual Regulation of RNA Silencing and the IFN Response as an
Antiviral Defense System in Mammalian Cells. Int J Mol Sci. 2020;21: 1348.
doi:10.3390/ijms21041348

60. Lui P-Y, Wong L-YR, Ho T-H, Au SWN, Chan C-P, Kok K-H, et al. PACT Facilitates
RNA-Induced Activation of MDA5 by Promoting MDA5 Oligomerization. J Immunol.
2017;199: 1846–1855. doi:10.4049/jimmunol.1601493

61. Kok K-H, Lui P-Y, Ng M-HJ, Siu K-L, Au SWN, Jin D-Y. The Double-Stranded RNA-
Binding Protein PACT Functions as a Cellular Activator of RIG-I to Facilitate Innate
Antiviral   Response.   Cell   Host   Microbe.   2011;9:   299–309.
doi:10.1016/j.chom.2011.03.007

62. Clemens MJ. PKR-a protein kinase regulated by double-stranded RNA. Int J Biochem Cell
Biol. 1997;29: 945–9. doi:10.1016/s1357-2725(96)00169-0

63. Gal-Ben-Ari S, Barrera I, Ehrlich M, Rosenblum K. PKR: A Kinase to Remember. Front
Mol Neurosci. 2018;11: 480. doi:10.3389/fnmol.2018.00480

64. Fairman-Williams ME, Guenther U-P, Jankowsky E. SF1 and SF2 helicases: family
matters. Curr Opin Struct Biol. 2010;20: 313–324. doi:10.1016/j.sbi.2010.03.011

65. Ahmad S, Hur S. Helicases in Antiviral Immunity: Dual Properties as Sensors and
Effectors. Trends Biochem Sci. 2015;40: 576–585. doi:10.1016/j.tibs.2015.08.001

66. Mittelstadt M, Frump A, Khuu T, Fowlkes V, Handy I, Patel CV, et al. Interaction of
human tRNA-dihydrouridine synthase-2 with interferon-induced protein kinase PKR.
Nucleic Acids Res. 2008;36: 998–1008. doi:10.1093/nar/gkm1129

67. Park H, Davies MV, Langland JO, Chang HW, Nam YS, Tartaglia J, et al. TAR RNA-
binding protein is an inhibitor of the interferon-induced protein kinase PKR. Proc Natl
Acad Sci U S A. 1994;91: 4713–4717. doi:10.1073/pnas.91.11.4713

68. Nakamura T, Kunz RC, Zhang C, Kimura T, Yuan CL, Baccaro B, et al. A Critical Role
for PKR Complexes with TRBP in Immunometabolic Regulation and eIF2α
Phosphorylation   in   Obesity.   Cell   Rep.   2015;11:   295–307.
doi:10.1016/j.celrep.2015.03.021

69. Daher A, Laraki G, Singh M, Melendez-Peña CE, Bannwarth S, Peters AHFM, et al. TRBP
control of PACT-induced phosphorylation of protein kinase R is reversed by stress. Mol
Cell Biol. 2009;29: 254–265. doi:10.1128/MCB.01030-08

70. Singh M, Castillo D, Patel CV, Patel RC. Stress-induced phosphorylation of PACT reduces
its interaction with TRBP and leads to PKR activation. Biochemistry. 2011;50: 4550–4560.
doi:10.1021/bi200104h

71. Clerzius G, Shaw E, Daher A, Burugu S, Gélinas J-F, Ear T, et al. The PKR activator,
PACT, becomes a PKR inhibitor during HIV-1 replication. Retrovirology. 2013;10: 96.
doi:10.1186/1742-4690-10-96

72. Li J, Tang H, Mullen TM, Westberg C, Reddy TR, Rose DW, et al. A role for RNA helicase

A in post-transcriptional regulation of HIV type 1. Proc Natl Acad Sci U S A. 1999;96:
709–714. doi:10.1073/pnas.96.2.709

73. Fujii R, Okamoto M, Aratani S, Oishi T, Ohshima T, Taira K, et al. A Role of RNA
Helicase A in cis-Acting Transactivation Response Element-mediated Transcriptional
Regulation of Human Immunodeficiency Virus Type 1. J Biol Chem. 2001;276: 5445–
5451. doi:10.1074/jbc.M006892200

74. Roy BB, Hu J, Guo X, Russell RS, Guo F, Kleiman L, et al. Association of RNA helicase
a with human immunodeficiency virus type 1 particles. J Biol Chem. 2006;281: 12625–
12635. doi:10.1074/jbc.M510596200

75. Sadler AJ, Latchoumanin O, Hawkes D, Mak J, Williams BRG. An Antiviral Response
Directed by PKR Phosphorylation of the RNA Helicase A. Gale M, editor. PLoS Pathog.
2009;5: e1000311. doi:10.1371/journal.ppat.1000311

76. Herbert A, Alfken J, Kim YG, Mian IS, Nishikura K, Rich A. A Z-DNA binding domain
present in the human editing enzyme, double-stranded RNA adenosine deaminase. Proc
Natl Acad Sci U S A. 1997;94: 8421–8426. doi:10.1073/pnas.94.16.8421

77. Yang W, Chendrimada TP, Wang Q, Higuchi M, Seeburg PH, Shiekhattar R, et al.
Modulation of microRNA processing and expression through RNA editing by ADAR
deaminases. Nat Struct Mol Biol. 2006;13: 13–21. doi:10.1038/nsmb1041

78. Iizasa H, Wulff B-E, Alla NR, Maragkakis M, Megraw M, Hatzigeorgiou A, et al. Editing
of Epstein-Barr virus-encoded BART6 microRNAs controls their dicer targeting and
consequently affects viral latency. J Biol Chem. 2010;285: 33358–33370.
doi:10.1074/jbc.M110.138362

79. Ota H, Sakurai M, Gupta R, Valente L, Wulff B-E, Ariyoshi K, et al. ADAR1 forms a
complex with Dicer to promote microRNA processing and RNA-induced gene silencing.
Cell. 2013;153: 575–589. doi:10.1016/j.cell.2013.03.024

80. Samuel CE. Adenosine deaminases acting on RNA (ADARs) are both antiviral and
proviral. Virology. 2011;411: 180–193. doi:10.1016/j.virol.2010.12.004

81. Li Z, Wolff KC, Samuel CE. RNA adenosine deaminase ADAR1 deficiency leads to
increased activation of protein kinase PKR and reduced vesicular stomatitis virus growth
following interferon treatment. Virology. 2010;396: 316–322.
doi:10.1016/j.virol.2009.10.026

82. Clerzius G, Gélinas J-F, Daher A, Bonnet M, Meurs EF, Gatignol A. ADAR1 interacts
with PKR during human immunodeficiency virus infection of lymphocytes and contributes
to viral replication. J Virol. 2009;83: 10119–10128. doi:10.1128/JVI.02457-08

83. Schoggins JW, Wilson SJ, Panis M, Murphy MY, Jones CT, Bieniasz P, et al. A diverse
range of gene products are effectors of the type I interferon antiviral response. Nature.
2011;472: 481–485. doi:10.1038/nature09907

84. Elde NC, Child SJ, Geballe AP, Malik HS. Protein kinase R reveals an evolutionary model
for defeating viral mimicry. Nature. 2009;457: 485–489. doi:10.1038/nature07529

85. Meister G, Landthaler M, Peters L, Chen PY, Urlaub H, Lührmann R, et al. Identification
of novel argonaute-associated proteins. Curr Biol CB. 2005;15: 2149–2155.
doi:10.1016/j.cub.2005.10.048

86. Concordet J-P, Haeussler M. CRISPOR: intuitive guide selection for CRISPR/Cas9
genome editing experiments and screens. Nucleic Acids Res. 2018;46: W242–W245.
doi:10.1093/nar/gky354

87. Chicois C, Scheer H, Garcia S, Zuber H, Mutterer J, Chicher J, et al. The UPF1 interactome
reveals interaction networks between RNA degradation and translation repression factors
in Arabidopsis. Plant J. 2018;96: 119–132. doi:10.1111/tpj.14022

88. Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S, Kundu DJ, et al.
The PRIDE database and related tools and resources in 2019: improving support for

quantification data. Nucleic Acids Res. 2019;47: D442–D450. doi:10.1093/nar/gky1106

89. Chen EY, Tan CM, Kou Y, Duan Q, Wang Z, Meirelles GV, et al. Enrichr: interactive and
collaborative HTML5 gene list enrichment analysis tool. BMC Bioinformatics. 2013;14:
128. doi:10.1186/1471-2105-14-128

90. Kuleshov MV, Jones MR, Rouillard AD, Fernandez NF, Duan Q, Wang Z, et al. Enrichr:
a comprehensive gene set enrichment analysis web server 2016 update. Nucleic Acids Res.
2016;44: W90-97. doi:10.1093/nar/gkw377

1072

Figure legends

Figure 1. Analysis of SINV infection in HEK293T cells and characterization of FHA:DICER WT cell lines

A. GFP fluorescent microscopy pictures of HEK293T, NoDice FHA:ctrl #1 and FHA:DICER cell lines infected (polyclonal and two clones, #4 and #17) with SINV-GFP at an MOI of 0.02 for 24 h. The left panel corresponds to GFP signal from infected cells and the right panel to a merge picture of GFP signal and brightfield. Pictures were taken with a 5x magnification. hpi: hours post-infection. **B.** Western blot analysis of DICER (DICER and HA) and GFP expression in SINV-GFP-infected HEK293T, NoDice FHA:ctrl #1 and FHA:DICER cell lines shown in **A**. Gamma-Tubulin was used as loading control. **C.** Mean (+/- SEM) of SINV-GFP viral titers in the same cell lines as in **A** infected at an MOI of 0.02 for 24 h (n = 3) from plaque assay quantification. ns: non-significant, ordinary one-way ANOVA test with Bonferroni correction. **D.** Western blot analysis of DICER (DICER and HA) and AGO2 expression in HEK293T, NoDice FHA:ctrl #1 and FHA:DICER cell lines. Gamma-Tubulin was used as loading control.

Figure 2. LC-MS/MS analysis of DICER interactome during SINV infection

A. Volcano plot for differentially expressed proteins (DEPs) between HA IP and CTL IP in FHA:DICER mock-infected cells. Each protein is marked as a dot; proteins that are significantly up-regulated in HA IP are shown in red, up-regulated proteins in CTL IP are shown in blue, and non-significant proteins are in black. The horizontal line denotes a p-value of 0.05 and the vertical lines the Log2 fold change cutoff (-1 and 1). DICER and its cofactors (TRBP, PACT, AGO2) are highlighted in yellow. **B.** Volcano plot for DEPs between SINV-GFP (MOI of 2, 6 hpi) and mock fractions of HA IP in FHA:DICER cells. Same colour code and thresholds as in A have been applied. Proteins that are discussed in the text are highlighted in yellow and SINV proteins in purple. **C.** Gene Ontology (GO) term enrichment of proteins up-regulated in SINV-GFP fraction of HA IP using Enrichr software [89,90]. The graph displays the GO term hierarchy within the "biological process" branch sorted by p-value ranking computed from the Fisher exact test. The length of each bar represents the significance of that specific term. In addition, the brighter the colour is, the more significant that term is. Viral proteins have been excluded for this analysis. **D.** STRING interaction network of the top 100 proteins enriched in SINV-infected vs. mock-infected cells. Proteins involved in RNA metabolic processes or the regulation thereof are indicated in red and blue respectively, proteins with a known dsRNA binding function are indicated in green. DICER is indicated by a red

circle. **E.** Summary of the differential expression analysis of SINV-GFP vs mock fractions from 1106
HA IP in FHA:DICER cells. The analysis has been performed using a generalized linear model 1107
of a negative-binomial distribution and p-values were corrected for multiple testing using the 1108
Benjamini-Hochberg method. 1109

1110

Figure 3. Confirmation of LC-MS/MS analysis by co-IP and BiFC 1111

A. Western blot analysis of HA co-IP in mock or SINV-GFP-infected (MOI of 2, 6 hpi) 1112
FHA:DICER WT #4 cells. Proteins associated to FHA:DICER were revealed by using 1113
antibodies targeting endogenous ADAR-1, PKR, TRBP, DHX9 or PACT proteins. In parallel, 1114
an HA antibody was used to verify the IP efficiency and GFP antibody was used to verify the 1115
infection. Ponceau was used as loading control. **B.** Western blot analysis of HA co-IP in mock 1116
or SINV-GFP-infected (MOI of 2, 6 hpi) FHA:DICER WT #4 cells. The lysate was treated or 1117
not with RNase A/T1. Proteins associated to FHA:DICER were revealed by using antibodies 1118
targeting endogenous DHX9, p-PKR, PKR, TRBP, or PACT proteins. In parallel, an HA 1119
antibody was used to verify the IP efficiency and GFP antibody was used to verify the infection. 1120
Ponceau was used as loading control. The arrow points to the expected size of the FHA:DICER 1121
protein and the asterisks correspond to aspecific bands. **C.** Western blot analysis to validate the 1122
interaction of PKR with DICER (upper panel) and PACT (lower panel) in mock or SINV-GFP- 1123
infected HEK293T cells (MOI of 2, 6 hpi). Immunoprecipitated proteins obtained from PKR 1124
pulldowns were compared to rabbit IgG pulldowns to verify the specificity of the assay. **D.** 1125
Interactions between DICER and TRBP, PACT or PKR were visualized by BiFC. Plasmids 1126
expressing [N-ter]Venus:DICER and TRBP:, PACT: or PKR:Venus[C-ter] were co-transfected in 1127
NoDice/ΔPKR cells for 24 h and cells were either infected with SINV at an MOI of 2 for 6 h 1128
or not. The different combinations are indicated on the left side. Reconstitution of Venus (BiFC) 1129
signal was observed under epifluorescence microscope. For each condition, the left panel 1130
corresponds to Venus signal and the right panel to the corresponding brightfield pictures. Scale 1131
bar: 100 μm. hpi: hours post-infection. **E.** BiFC experiment on fixed NoDice/ΔPKR cells 1132
treated as in D. After fixation, cells were stained with DAPI and observed under confocal 1133
microscope. Only a merge picture of BiFC and DAPI signals of SINV-infected cells is shown 1134
here. A higher magnification of picture showing cytoplasmic localization of the interaction 1135
represented by a red square is shown in the bottom left corner. Scale bar: 20 μm and 10 μm. 1136

1137

Figure 4. Confirmation of DICER interactome upon SINV infection in HCT116 KI-DICER cells

A. Western blot analysis of DICER, AGO2, PKR and TRBP expression in HEK293T, HCT116 and HCT116 KI-DICER cell lines. Gamma-Tubulin and ponceau were used as loading control. **B**. GFP fluorescent microscopy pictures of HEK293T, HCT116 and HCT116 KI-DICER cell lines infected with SINV-GFP at an MOI of 0.02, 0.1 and 1 for 24 h. The left panel corresponds to GFP signal from infected cells and the right panel to a merge picture of GFP signal and brightfield. Pictures were taken with a 5x magnification. **C**. miR-16 expression analyzed by northern blot in in the same cell lines as in B. Expression of snRNA U6 was used as loading control. **D.** Western blot analysis of HA co-IP in mock or SINV-GFP-infected (MOI of 0.1, 24 hpi) HCT116 KI-DICER cells. Proteins associated to FHA-GFP:DICER were revealed by using antibodies targeting endogenous DHX9, p-PKR, PKR, PACT or TRBP proteins. The TRBP immunoblot was performed by loading the same samples on a separate membrane. In parallel, an HA antibody was used to verify the IP efficiency and GFP antibody was used to verify the infection. Ponceau was used as a loading control.

Figure 5. Identification of DICER domains involved in DICER-PKR interaction

A. Schematic representation of Human DICER proteins used in this study. The different conserved domains are shown in colored boxes. DUF283: Domain of Unknown Function; PAZ: PIWI ARGONAUTE ZWILLE domain; dsRBD: dsRNA-binding domain. hDICER WT is the full-length protein. hDICER N1 is deleted of the first N-terminal 495 amino acids. hDICER N3 is wholly deleted of the helicase domain. hDICER Hel. is the whole DICER's helicase domain. hDICER ΔdsRBD is deleted of the C-terminal dsRBD. **B.** Western blot analysis of HA co-IP in mock NoDice 2.20 cells transfected with different versions of FHA:DICER proteins. Efficiency of immunoprecipitation was assessed using anti-HA and anti-DICER antibodies and co-IPs of TRBP, PKR and PACT were examined using appropriate antibodies. Expression of GFP in INPUT fraction was visualized as control of SINV-GFP infection. Ponceau staining of membranes is used as loading control. **C.** Western blot analysis of HA co-IP in NoDice 2.20 cells transfected with different versions of FHA:DICER proteins and infected with SINV-GFP (MOI of 2, 6 hpi). Efficiency of immunoprecipitation was assessed using an anti- Flag antibody and co-IPs of PKR, TRBP, p-PKR and PACT were examined using appropriate antibodies. Expression of GFP in INPUT fraction was visualized as control of SINV-GFP infection. Ponceau staining of membranes is used as loading control. The DICER Hel. band is indicated by a red asterisk. **D.** Plasmids expressing the different versions of DICER proteins fused to the

 N-terminal part of Venus and PKR:Venus$^{C\text{-ter}}$ plasmid were co-transfected in NoDice/ΔPKR

1173   cells. Cells were treated as in Fig. 3D. The different combinations are noted on the left side.

1174   The fluorescent signal was observed using an epifluorescence microscope. For each condition,

1175   the left panel corresponds to Venus signal and the right panel to the corresponding brightfield

1176   pictures. Scale bar: 100 μm. hpi: hours post-infection.

1177

Figure 6. Analysis of the importance of Dicer helicase domain on SINV-GFP infection in

FHA:DICER mutant stable cell lines

A. Expression of DICER (DICER, HA), TRBP and AGO2 was analysed by western blot in

HEK293T, NoDice FHA:ctrl #1, FHA:DICER WT #4 and FHA:DICER N1 #6 cell lines.

Gamma-Tubulin was used as loading control. **B.** Northern blot analysis of miR-16 expression

in the same samples as in A. Expression of snRNA U6 was used as loading control. **C.**

Representative GFP fluorescent microscopy images of HEK293T, FHA:DICER WT #4 and

FHA:DICER N1 #6 cell lines infected with SINV-GFP at an MOI of 0.02 for 24 h. The left

panel corresponds to GFP signal and the right panel to a merge picture of GFP signal and

brightfield. Pictures were taken with a 5x magnification. hpi: hours post-infection. **D.** Western

blot analysis of DICER (DICER and HA), AGO2, PKR, and GFP expression in SINV-GFP-

infected cells in the same condition as in C. Gamma-Tubulin was used as loading control. The

asterisk correspond to aspecific bands **E.** Mean (+/- SEM) of SINV-GFP viral titers fold change

over HEK293T cells in HEK293T, NoDice 2.20, FHA:DICER WT #4 and FHA:DICER

mutants cell lines infected at an MOI of 0.02 for 24 h (n = 3) from plaque assay quantification.

* $p < 0.05$, ns: non-significant, ordinary one-way ANOVA test with Bonferroni correction.

1194

Figure 7. The antiviral effect of helicase-deleted DICER mutants is independent from

AGO2

A. GFP fluorescent microscopy pictures of NoDice FHA:ctrl #2, NoDice FHA:DICER WT #4

and NoDice FHA:DICER mutant cell lines treated with two doses of siAGO2 at 20nM for 48

hours before a 24-hour-SINV-GFP infection at an MOI of 0.02. The left panel corresponds to

GFP signal from infected cells and the right panel to a merge picture of GFP signal and

brightfield. Pictures were taken with 5x magnification. hpi: hours post-infection. **B.** Western

blot analysis of DICER, AGO2 and GFP expression in SINV GFP-infected NoDice FHA:ctrl

#2, NoDice FHA:DICER WT #4 and NoDice FHA:DICER mutant cell lines shown in **A.** Cells

were treated with two doses of siAGO2 at 20nM for 48 hours before a 24-hour-SINV-GFP

infection at an MOI of 0.02. Gamma-Tubulin was used as loading control. **C.** Mean (+/-SEM)
of SINV-GFP viral titers in the same cell lines as in **A.** infected at an MOI of 0.02 for 24h (n =
3) from plaque assay quantification. ns: non-significant, two-tailed unpaired parametric t-test.
1208

Figure 8. The antiviral effect of helicase-deleted DICER mutants requires PKR
A. GFP fluorescent microscopy pictures of HEK293T, NoDiceΔPKR:ctrl and FHA:DICER
mutant cell lines infected with SINV-GFP at an MOI of 0.02 for 24h. The left panel corresponds
to GFP signal from infected cells and the right panel to a merge picture of GFP signal and
brightfield. Pictures were taken with 5x magnification. hpi: hours post-infection. **B.** Western
blot analysis of DICER (DICER and HA), AGO2, PKR and GFP expression in SINV-GFP-
infected HEK293T, NoDiceΔPKR FHA:ctrl and FHA:DICER mutant cell lines shown in **A.**
Gamma-Tubulin was used as loading control. **C.** Mean (+/-SEM) of SINV-GFP viral titers in
the same cell lines as in **A.** infected at an MOI of 0.02 for 24h (n = 3) from plaque assay
quantification. ns: non-significant, ordinary one-way ANOVA test with Bonferroni correction.
1219

 Supplementary material

 Supplementary figure legends

Figure S1. Analysis of SINV-GFP infection in FHA:DICER cell lines at different MOI

and time points.

A. miR-16 expression analyzed by northern blot in HEK293T, NoDice FHA:ctrl #1 and

FHA:DICER WT #4 cell lines**.** Expression of snRNA U6 was used as loading control. **B.**

Representative GFP pictures of HEK293T, NoDice 2.20, NoDice 4.25, NoDice FHA:ctrl #1

and NoDice FHA:ctrl #2 cells infected with SINV-GFP at an MOI of 0.02 for 24h. The left

panel corresponds to GFP signal and the right panel to a merge of GFP signal and the

corresponding brightfield. Pictures were taken with a 5x magnification. hpi: hours post-

infection. **C.** Mean (+/- SEM) of SINV-GFP viral titers in cells infected at an MOI of 0.02 for

24 h (n=3) from plaque assay quantification. * $p < 0.05$, ns: non-significant, ordinary one-way

ANOVA test with Bonferroni correction. **D**. Western blot analysis of DICER, AGO2 and GFP

expression in SINV-GFP-infected cells shown in B. Gamma-Tubulin was used as loading

control.

1235

Figure S2. LC-MS/MS analysis of DICER interactome during SFV infection.

A. Volcano plot for differentially expressed proteins (DEPs) between HA IP and CTL IP in

FHA:DICER mock-infected cells. Each protein is marked as a dot; proteins that are

significantly up-regulated in HA IP are shown in red, up-regulated proteins in CTL IP are

shown in blue, and non-significant proteins are in black. The horizontal line denotes a p-value

of 0.05 and the vertical lines the Log2 fold change cutoff (-1 and 1). DICER and its cofactors

(TRBP, PACT, AGO2) are highlighted in yellow. **B.** Left panel: Volcano plot for DEPs

between SFV (MOI of 2, 6 hpi) and mock fractions of HA IP in FHA:DICER cells. Same colour

code and thresholds as in A were applied. Proteins that are discussed in the text are highlighted

in yellow and SFV proteins in purple. **C.** Summary of the differential expression analysis of

SFV vs mock fractions from HA IP in FHA:DICER cells. The analysis has been performed

using a generalized linear model of a negative-binomial distribution and p-values were

corrected for multiple testing using the Benjamini-Hochberg method.

1249

1250 **Figure S3. Confirmation of LC-MS/MS analysis by co-IP and BiFC controls.**

1251 **A.** FHA:DICER WT #4 cells were infected with SINV-GFP at an MOI of 0.02 for 24 h and a

1252 HA co-IP was performed. Eluted proteins were resolved by western blot and IP efficiency was

assessed using an HA antibody. In parallel, co-IPed proteins were visualized using appropriate antibodies. GFP antibody was used to verify the infection and Ponceau staining serves as loading control. **B.** 1% Agarose gel analysis of RNA extracted from INPUT of the co-IP in Fig. 3B. Ribosomal RNA integrity was compared to a control HEK293T cell line. RNAs were revealed using ethidium bromide under UV. **C.** Schematic representation of Human DICER proteins used for BiFC positive and negative controls. The different conserved domains are shown in colored boxes. DUF283: Domain of Unknown Function; PAZ: PIWI ARGONAUTE ZWILLE domain; dsRBD: dsRNA-binding domain. hDICER WT is the full-length protein. hDICER N1 is deleted of the first N-terminal 495 amino acids. **D.** Expression of BiFC plasmids was assessed by western blot. DICER proteins (WT and N1) and PKR were visualized using antibodies targeting endogenous proteins, whereas TRBP and PACT were detected using GFP antibody. Antibody targeting the SINV coat protein (CP) was used as infection control. Ponceau staining was used as loading control. **E.** Positive and negative BiFC controls on fixed NoDice/ΔPKR cells. After co-transfection, cells were infected with SINV at an MOI of 2 for 6 h and fixed. After fixation, cells were stained with DAPI and observed under confocal microscope. Merge pictures of BiFC and DAPI signals of SINV-infected cells are shown. A higher magnification of images showing the interaction represented by a red square is shown in the bottom left corner. Scale bar: 20 μm and 10 μm. **F.** Expression of BiFC plasmids was assessed by western blot. DICER, PKR, TRBP and PACT were detected using GFP antibody. Antibody targeting the SINV coat protein (CP) was used as infection control. Gamma-Tubulin was used as loading control. The asterisk corresponds to an aspecific band. **G.** Interactions between DICER and TRBP, PACT or PKR were visualized by BiFC. Plasmids expressing [N-ter]Venus:DICER and TRBP:, PACT: or PKR:Venus[C-ter] were co-transfected in HEK293T cells for 24 h and cells were either infected with SINV at an MOI of 2 for 6 h or not. The different combinations are indicated on the left side. Reconstitution of Venus (BiFC) signal was observed under epifluorescence microscope. For each condition, the left panel corresponds to Venus signal and the right panel to the corresponding brightfield pictures. Scale bar: 100 μm.

Figure S4. Confirmation of DICER interactome upon SINV infection in HCT116 KI-DICER cells.

A. Schematic representation of DICER WT and Flag-HA(FHA)-GFP kocked-in (KI) alleles. FHA sequence is in purple, GFP in green, DICER 5'UTR in orange and DICER coding region in yellow. The gRNA used to generate the KI was designed to target the first coding exon of

DICER gene. **B.** PCR on genomic DNA extracted from WT and KI cells. **C.** An oligo outside the homologous recombination region and an oligo within the GFP tag were used to verify the presence of a 1040bp amplicon in HCT116 KI-DICER clone. Sequencing results corresponding to this region are shown. **D.** Western blot analysis of DICER, p-PKR, PKR and p-eIF2α expression in mock or SINVGFP-infected HEK293T and HCT116 KI-DICER cell lines at an MOI of 2 for 6h or 16h and 0.02 for 24h. GFP antibody was used to verify the infection. Ponceau and gamma-Tubulin were used as loading control.

Figure S5. Interaction analysis between the different versions of DICER and TRBP or PACT using BiFC assay.

NoDice/ΔPKR cells were co-transfected for 24 h with plasmids expressing the different versions of DICER proteins fused to the N-terminal part of Venus and either TRBP:Venus^{C-ter} (**A**) or PACT:Venus^{C-ter} (**B**). Cells were then infected with SINV at an MOI of 2 for 6 h and Venus signal was observed under epifluorescence microscope. The left panel corresponds to Venus signal and the right panel to the corresponding brightfield picture. Pictures were taken with a 5x magnification. hpi: hours post-infection. Scale bar: 100 μm.

Figure S6. Analysis of the importance of Dicer helicase domain on SINV-GFP infection in FHA:DICER mutant stable cell lines.

A. Northern blot analysis of miR-16 expression in HEK293T, NoDice 2.20, NoDice FHA:ctrl #2, FHA:DICER WT polyclonal, FHA:DICER N1 #6, FHA:DICER Hel. #2.6, and FHA:DICER N3 #2.13. Expression of snRNA U6 was used as loading control. **B.** Representative GFP fluorescent microscopy images of HEK293T, NoDice 2.20, FHA:DICER mutants cell lines infected with SINV-GFP at an MOI of 0.02 for 24 h. The left panel corresponds to GFP signal and the right panel to a merge picture of GFP signal and brightfield. Pictures were taken with a 5x magnification. hpi: hours post-infection. **C.** Mean (+/- SEM) of SINV-GFP viral titers over FHA:DICER WT #4 cells in FHA:DICER N1 #6, FHA:DICER N3 #2.13, NoDice FHA:ctrl #2 and NoDice 2.20 cell lines infected at an MOI of 0.02 for 24 h (n = 3) from plaque assay quantification. *** $p < 0.001$, ns: non-significant, ordinary one-way ANOVA test with Bonferroni correction.

1319 **Supplementary tables**

1320 **Supp. Table 1:** Top 100 proteins that are differentially immunoprecipitated in mock-infected
1321 FHA:DICER cells by the HA and Myc (CTL) antibodies. Related to Figure 2.

1322

1323 **Supp. Table 2:** Top 100 proteins that are differentially immunoprecipitated with the HA
1324 antibody in SINV-infected vs mock-infected FHA:DICER cells. Related to Figure 2.

1325

1326 **Supp. Table 3:** Top 100 proteins that are differentially immunoprecipitated in mock-infected
1327 FHA:DICER cells by the HA and Myc (CTL) antibodies, in the SFV infection experiment.
1328 Related to Figure S2.

1329

1330 **Supp. Table 4:** Top 100 proteins that are differentially immunoprecipitated with the HA
1331 antibody in SFV-infected vs mock-infected FHA:DICER cells. Related to Figure S2.

1332

1333 **Supp. Table 5:** List of primers used in this study.

1334

1335 **Supp. Table 6:** data and statistical tests details used in plaque assays shown in Figures 1, 6, 7,
1336 8, S1 and S6

Figure 1

A

GFP | GFP + Brightfield

HEK293T

NoDice 2.20

FHA:ctrl #1

FHA:DICER WT #4

FHA:DICER polyclonal

FHA:DICER WT #17

SINV-GFP MOI 0.02 24 hpi

B

Mock | SINV-GFP MOI 0.02 24 hpi

HEK293T | FHA:ctr l#1 | FHA:DICER WT #4 | HEK293T | FHA:ctrl #1 | FHA:DICER WT #4

250 kDa — DICER
250 kDa — HA
35 kDa / 25 kDa — GFP
55 kDa — Tubulin

C

SINV-GFP titer MOI 0.02 24 hpi

Log10 viral titer PFU/mL

10^8
10^7
10^6
10^5

HEK293T | FHA:ctrl #1 | FHA:DICER WT #4 | FHA:DICER WT polyclonal | FHA:DICER WT #17

ns ns ns ns ns

NoDice 2.20

D

HEK293T | FHA:DICER polyclonal | FHA:DICER WT #4 | FHA:DICER WT #17

SINV-GFP - + - + - + - +

250 kDa — DICER
250 kDa — HA
35 kDa / 25 kDa — GFP
55 kDa — Tubulin
100 kDa — AGO2
55 kDa — Tubulin

Figure 2

A FHA:DICER IP vs CTL IP

B SINV-GFP vs Mock in FHA:DICER IP

C

RNA binding (GO:0003723)
ATP-dependent helicase activity (GO:0008026)
snoRNA binding (GO:0030515)
RNA helicase activity (GO:0003724)
ATP-dependent RNA helicase activity (GO:0004004)
RNA-dependent ATPase activity (GO:0008186)
double-stranded RNA binding (GO:0003725)
actin-dependent ATPase activity (GO:0030898)
ATP-dependent DNA helicase activity (GO:0004003)
RNA stem-loop binding (GO:0035613)

D

E

Gene name	Mean spectral count		LogFC	Adjusted p-value
	SINV-GFP IP	MOCK IP		
E2AK2 (PKR)	123.3	0.3	7.979	7.316e-102
DSRAD (ADAR-1)	181.3	28.3	2.563	1.421e-73
DHX9	247.3	50	2.194	4.012e-83
PRKRA (PACT)	63.3	13.7	2.093	1.158e-20

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

A

SINV-GFP MOI 0.02 24hpi

B

C

SINV-GFP titer MOI 0.02 24 hpi

Figure 8

A

	GFP	GFP + Brightfield

NoDiceΔPKR

HEK 293T

FHA:ctrl

FHA:DICER WT #2

FHA:DICER N1 #6

FHA:DICER N3 #8

SINV-GFP MOI 0.02 24hpi

B

NoDiceΔPKR

FHA:DICER

	HEK 293T	FHA: ctrl	WT #2	N1 #6	N3 #8	
SINV GFP	− +	− +	− +	− +	− +	
250 kDa						DICER
250 kDa						HA
100 kDa						AGO2
70 kDa						PKR
55 kDa						Tubulin
35 kDa / 25 kDa						GFP
						Ponceau

C

SINV-GFP titer MOI 0.02 24 hpi

Log10 viral titer PFU/mL

ns ns ns ns

HEK293T, FHA:ctrl, FHA:DICER WT #2, FHA:DICER N1 #6, FHA:DICER N3 #8

NoDiceΔPKR