Human Dicer helicase domain acts as an interaction platform to recruit PKR and dsRNA binding proteins during viral infection

Thomas C. Montavon[1,*], Mathieu Lefèvre[1,*], Morgane Baldaccini[1,*], Erika Girardi[1], Béatrice Chane-Woon-Ming[1], Mélanie Messmer[1], Philippe Hammann[2], Johana Chicher[2], Sébastien Pfeffer[1,‡]

[1] Université de Strasbourg, Architecture et Réactivité de l'ARN, Institut de Biologie Moléculaire et Cellulaire du CNRS, 2 allée Konrad Roentgen, 67084 Strasbourg France

[2] Université de Strasbourg, Institut de Biologie Moléculaire et Cellulaire du CNRS, Plateforme Protéomique Strasbourg – Esplanade, 2 allée Konrad Roentgen, 67084 Strasbourg France

*These authors contributed equally

‡To whom correspondence should be addressed: spfeffer@unistra.fr

Abstract

The innate immune response against viruses mainly involves type I interferon (IFN) in

mammalian cells. The exact contribution of the RNA silencing machinery remains to be

established, but several recent studies indicate that the type III ribonuclease DICER can

generate viral siRNAs in specific conditions. In addition, it has been proposed that type I IFN

and RNA silencing could be mutually exclusive antiviral responses. In order to decipher the

implication of DICER during infection of human cells with the Sindbis virus, we determined

its interactome by immunoprecipitation and mass spectrometry analysis. Our results show that

human DICER specifically interacts with several double-stranded RNA binding proteins and

RNA helicases during viral infection. In particular, proteins such as DHX9, ADAR-1 and the

protein kinase RNA-activated (PKR) are enriched with DICER in virus-infected cells. We

demonstrated the importance of DICER helicase domain in its interaction with PKR and

showed that it has functional consequences for the cellular response to viral infection.

Introduction

In mammalian cells, the main antiviral defense system involves the activation of a signaling cascade relying on production of type I interferon (IFN I). This pathway depends on the recognition of extrinsic signals or pathogen associated molecular patterns (PAMPs) by dedicated host receptors. Double-stranded (ds) RNA, which can originate from viral replication or convergent transcription, is a very potent PAMP and can be sensed in the cell by various proteins among which a specific class of DExD/H-box helicases called RIG-I-like receptors (RLRs) (Ivashkiv and Donlin, 2014). RLRs comprise RIG-I, MDA5 and LGP2 and transduce viral infection signals to induce expression of IFN I cytokines that act in autocrine and paracrine fashion. These cytokines then trigger the expression of hundreds of interferon stimulated genes (ISGs) to stop the virus in its tracks (Borden et al, 2007). Among those ISGs, dsRNA-activated protein kinase R (PKR) plays an important role in antiviral defense by blocking cellular and viral translation upon direct binding to long dsRNA (Williams, 1999). PKR is a serine-threonine kinase that dimerizes and auto-phosphorylates upon activation. It then phosphorylates numerous cellular targets among which the translation initiation factor eIF2α, which results in the inhibition of cap-dependent translation (Lemaire et al., 2008). Accordingly, translation of many RNA viruses, including alphaviruses, is inhibited by PKR (Fros and Pijlman, 2016; Pfaller et al., 2011; Ryman et al., 2002). PKR is also involved in other cellular pathways including apoptosis, autophagy and cell cycle (Kim et al., 2014; Williams, 1999).

RNAi is another evolutionary conserved pathway triggered by long dsRNA sensing (Meister and Tuschl, 2004). One key component in this pathway is the type III ribonuclease DICER, which is also essential for micro (mi)RNA biogenesis (Hutvágner et al., 2001; Wienholds et al., 2003). These small regulatory RNAs are sequentially produced by the two ribonucleases DROSHA and DICER, before being loaded into an Argonaute (AGO) effector protein in order to regulate their target mRNAs (Bartel, 2018). Whatever its substrate, be it long

dsRNA or miRNA precursor, DICER relies on interacting with co-factors to be fully functional. In mammalian cells, the TAR-RNA binding protein (TRBP), a dsRNA binding protein (dsRBP), was shown to play a role in the selection of DICER substrates, its stabilization, strand selection and incorporation into AGO2 (Chendrimada et al., 2005). The interaction with TRBP is well characterized and depends on the helicase domain of DICER and the third dsRNA binding domain (dsRBD) of TRBP (Daniels et al., 2009). Another dsRBP, the protein activator of interferon-induced protein kinase R (PACT), was also described as an important cofactor of DICER. Although its function is not fully understood, PACT seems to also participate in miRNA loading and strand selection (Heyam et al., 2015; Kok et al., 2007). via protein-protein interaction between the DICER helicase domain and the third dsRBD of PACT (Lee et al., 2006).

It is now common knowledge that RNAi is the main antiviral defense system in several phyla such as plants, arthropods, nematodes (reviewed in (Guo et al., 2019)). However, its exact contribution in the mammalian antiviral response remains unclear (Cullen, 2006; Maillard et al., 2019; tenOever, 2016). Recent studies indicate that a functional antiviral RNAi does exist in mammals in specific cases. A functional antiviral RNAi response was first detected in undifferentiated mouse embryonic stem cells (Maillard et al., 2013) lacking the IFN response, suggesting that these two pathways could be incompatible. Indeed, in mammalian somatic cells deficient for MAVS or IFNAR, two components of the interferon response, an accumulation of DICER-dependent siRNAs derived from exogenous long dsRNA was detected (Maillard et al., 2016). In addition, the RLR LGP2 was found interacting with both DICER and TRBP, blocking respectively siRNA production and miRNA maturation (Takahashi et al., 2018a, 2018b; van der Veen et al., 2018). Moreover, AGO4 was recently showed to be involved in antiviral RNAi against Influenza A virus (IAV), Vesicular stomatitis virus (VSV) and Encephalomyocarditis virus (EMCV) (Adiliaghdam et al., 2020). Finally, viral suppressors of RNAi (VSRs) have been

shown to prevent DICER from playing an antiviral role in mammalian cells (Qiu et al., 2020,

2017). Nonetheless, several studies reported no detection of viral siRNAs in mammalian

somatic cells infected with several viruses (Girardi et al., 2013; Parameswaran et al., 2010;

Schuster et al., 2019). In somatic cells, only a helicase-truncated form of human DICER could

produce siRNAs from IAV genome(Kennedy et al., 2015), but it also turned out that these

siRNAs cannot confer an antiviral state (Tsai et al., 2018).

Based on these observations, we decided to study the involvement of DICER during

infection of human cells with the Sindbis virus (SINV). SINV is a member of the *Togaviridae*

family in the alphavirus genus, which is transmitted by mosquitoes to mammals and can induce

arthritogenic as well as encephalitic diseases (Griffin, 2007). It is widely used as a laboratory

alphaviruses model as it infects several cell types and replicates to high titers. SINV has a

positive stranded RNA genome of about 12 kb, which codes for two polyproteins that give rise

to non-structural and structural proteins, including the capsid. Moreover, upon viral replication,

a long dsRNA intermediate, which can be sensed by the host antiviral machinery, accumulates.

Of note, SINV dsRNA can be cleaved into siRNA in insects as well as in human cells expressing

the Drosophila DICER-2 protein (Girardi et al., 2015). Nonetheless, although human DICER

has the potential to interact with the viral RNA duplex, we did not find evidence that SINV

dsRNA could be processed into siRNAs in somatic mammalian cells (Girardi et al., 2015,

2013). We thus hypothesized that specific proteins could interfere with DICER during SINV

infection by direct interaction and limit its accessibility and/or activity. To address this

hypothesis, we generated HEK 293T cells expressing a tagged version of human DICER that

could be immunoprecipitated in mock or SINV-infected cells in order to perform a proteomic

analysis of its interactome. Among the proteins co-immunoprecipitated with DICER and that

were specifically enriched upon infection, we identified dsRBPs such as ADAR1, DHX9 PACT

and PKR. We further validated the direct interaction between DICER and PKR upon SINV

infection. We dissected the protein domains necessary for this interaction and we found that

DICER helicase domain plays a fundamental role as a recruitment platform for PKR but also

for other co-factors. Our results indicate that DICER interactome is highly dynamic and directly

link components of RNAi and IFN pathways in modulating the cellular response to viral

infection.

Results

Establishment of a HEK 293T cell line expressing FLAG-HA tagged DICER

In order to be able to study the interactome of the human DICER protein during viral infection,

we transduced *Dicer* knock-out HEK 293T cells (Hal P. Bogerd et al., 2014) with either a

lentiviral construct expressing a FLAG-HA-tagged wild type DICER protein (FHA:DICER) or

a construct without insert as a negative control (NoDice). After monoclonal selection of stably

transduced cells, we picked a clone that expressed FHA:DICER at levels relatively

physiological compared to HEK 293T wild type cells (Fig 1A). Of note, the expression of

AGO2, which dropped significantly in NoDice cells, was restored to levels similar to WT cells

in the FHA:DICER cell line. We then analyzed the ability of the tagged protein to produce

mature miRNAs by northern blot analysis (Fig 1B). The results show that expression of

FHA:DICER complements the defect in miRNA production observed in the NoDice cell line.

We next characterized the FHA:DICER cell line during SINV infection. We used a

modified version of SINV in which the subgenomic promoter has been duplicated to introduce

the GFP coding sequence (SINV-GFP) as a readout of viral infection (López et al., 2020). At

24 hours post-infection (hpi) and a multiplicity of infection (MOI) of 0.02, the fluorescent GFP

signal in FHA:DICER cells was similar to the one observed in HEK 293T cells, while NoDice

cells displayed a strong decrease in GFP signal (Fig 1C). Western blot analysis of GFP

accumulation (Fig 1D) as well as the analysis of viral particles production by plaque assay (Fig

1E) confirmed the epifluorescence microscope observations, *i.e.* a significantly lower

accumulation of viral protein and particles in the absence of the DICER protein. In order to

check if this phenotype was also visible at a higher MOI and earlier during infection, we

infected cells with SINV-GFP at an MOI of 2 for 6 or 16 h. Again, we observed that GFP

expression and viral particle production at 16 hpi were similar between HEK 293T and

FHA:DICER cells, while they were strongly impaired in NoDice cells (Fig S1A). At 6 hpi, we

made similar observations for the expression of the virus-encoded GFP protein (Fig S1B upper

and middle panels), but did not see any difference in viral titers between cell lines (Fig S1B

lower panel). The latter observation is most likely due to the fact that production of mature viral

particles was not yet completed at this early time point.

Altogether, these results indicate that the FHA-tagged DICER protein is functional and

can be used for proteomics studies. Moreover, these results also show that in our conditions,

the loss of DICER protein has an antiviral effect, which is in opposition with previous

observations where the replication of various viruses seems to be largely unaffected in DICER

knock-out somatic cells (Hal P. Bogerd et al., 2014; Parameswaran et al., 2010; Tsai et al.,

2018).

Analysis of DICER interactome during SINV infection by mass spectrometry

Our molecular tool being validated, we then focused on determining the interactome of

FHA:DICER during SINV infection. We performed an anti-HA immunoprecipitation

experiment (HA IP) coupled to label-free LC-MS/MS analysis in FHA:DICER cells either

mock-infected or infected for 6 h at an MOI of 2 with SINV-GFP. In parallel, we performed an

anti-MYC immunoprecipitation as a negative control (CTL IP). The experiments were

performed in technical triplicate in order to have statistically reproducible data for the

differential analysis, which was performed using spectral counts. Prior to the detailed analysis

of the results, we verified that there was no confounding factor in the experimentation by

performing a Principal Component Analysis (PCA). This allowed us to see that the replicates

were very homogenous and that the different samples were well separated based on the

conditions.

To check the specificity of the HA immunoprecipitation, we first compared the proteins

identified in the HA IP with the ones identified in the CTL IP in mock-infected cells.

Differential expression analysis allowed us to calculate a fold change and an adjusted p-value

for each protein identified and to generate a volcano plot representing the differences between

HA and CTL IP samples. Applying a fold change threshold of 2 (abs(LogFC) > 1), an adjusted

p-value threshold of 0.05 and a cutoff of at least 5 spectral counts in the most abundant

condition, we identified 258 proteins differentially immunoprecipitated between the two

conditions out of 1318 proteins (Fig 2A and Supp. Table 1). Among these, 123 proteins were

specifically enriched in the HA IP. The most enriched protein was DICER, followed by its

known co-factors TRBP and PACT (also known as PRKRA) (Chendrimada et al., 2005; Lee et

al., 2006). We were also able to retrieve AGO2, indicating that the RISC loading complex has

been immunoprecipitated and that proteins retrieved in our HA IP are specific to DICER

immunoprecipitation.

We next performed the differential expression analysis of proteins retrieved in the HA

IP in SINV-GFP compared to mock-infected cells. Among 1342 proteins, 296 were

differentially retrieved between conditions (Fig 2B and Supp. Table 2). Of these, 184 proteins,

including viral ones, were at least 2-fold enriched in SINV-GFP-infected cells. GO-term

analysis showed a significant enrichment in RNA binding proteins including double-stranded

RNA binding proteins and RNA helicases (Fig 2C). Among the RNA binding proteins

retrieved, the top and most specific DICER interactor is the interferon-induced, double-stranded

(ds) RNA-activated protein kinase PKR (also known as E2AK2), which is enriched more than

250 times in virus-infected cells (Fig 2B and D). We were also able to identify the dsRNA-

specific adenosine deaminase protein ADAR-1 (also known as DSRAD), as well as PACT,

which were enriched 5.9 and 4.2 times respectively in SINV-GFP-infected cells compared to

mock-infected cells (Fig 2B and D). Among the isolated RNA helicases, we identified the ATP-

dependent RNA helicase A protein DHX9, which is implicated in *Alu* element-derived dsRNA

regulation and in RISC loading (Aktaş et al., 2017; Robb and Rana, 2007). In order to verify if

the observed interactions were specific to SINV we performed the same experiments with

another virus of the *Togaviridae* family, the Semliki forest virus (SFV). In this analysis, we

were able to retrieve ADAR-1, DHX9, PACT and PKR, specifically enriched in SFV-infected

samples (Fig S2 and Supp. Tables 3 and 4). These results show that these interactions can be

retrieved in *Togaviridae*-infected cells.

Taken together, these results indicate that proteins interacting with DICER in virus-

infected cells are involved in dsRNA sensing and/or interferon-induced antiviral response.

Because SINV-GFP allows us to have a visual read-out of the infection, we decided to use this

virus to continue the analysis.

DICER and PKR interact *in vivo* in the cytoplasm during SINV infection

To validate the LC-MS/MS analysis, we performed a co-immunoprecipitation (co-IP) followed

by western blot analysis in FHA:DICER cells infected with SINV-GFP at an MOI of 2 for 6 h.

Whereas TRBP interacted equally well with FHA:DICER in mock and SINV-GFP-infected

cells, ADAR-1, PKR, DHX9 and PACT were only retrieved in the HA IP in SINV-infected

cells (Fig 3A). We verified that these interactions could also be observed at a later time in

infection by performing the HA IP in FHA:DICER cells infected with SINV-GFP for 24 h at

an MOI of 0.02. This allowed us to confirm the specific interactions between DICER and

ADAR-1, DHX9, PACT or PKR only in virus-infected cells (Fig S3A), indicating that these

interactions occur at an early stage of the infection and remain stable in time.

Because of the involvement of PKR in antiviral response (García et al., 2007) and the

fact that it shares common co-factors with DICER, namely TRBP and PACT (Haase et al.,

2005; Patel et al., 2000), we decided to focus our analysis on the DICER-PKR interaction. To

confirm the validity of this interaction, we first performed a reverse co-IP using magnetic beads

coupled to anti-PKR antibody in HEK 293T cells infected with SINV-GFP. While PACT

interacted as expected with PKR both in mock and in SINV-GFP-infected cells (Fig 3B upper

panel), it co-immunoprecipitated with DICER only in virus-infected cells as previously

observed (Fig 3B lower panel). In order to determine whether DICER and PKR could interact

directly *in vivo*, we set up a bi-molecular fluorescent complementation assay (BiFC) experiment

(Lepur et al., 2016). To this end, we fused the N- or C-terminal halves of the Venus protein ($^{N\text{-}}$

^{ter}Venus or $^{C\text{-}ter}$Venus) to DICER and to PKR but also to TRBP and PACT. Since we showed

above that an N-terminally tagged DICER was functional, we fused Venus fragments at the N-

terminal end of DICER. For the other three proteins, we fused Venus fragments at the N- or C-

terminus and selected the best combination. To avoid interaction with endogenous DICER and

PKR proteins, we conducted all BiFC experiments in NoDice/ΔPKR HEK 293T cells (Kennedy

et al., 2015). In order to control the BiFC experiment, we chose to exploit the well characterized

DICER-TRBP interaction, which is known to occur via the DICER DEAD-Box helicase

domain. We therefore used as a positive control the wild-type DICER protein, and as a negative

control a truncated version of DICER protein lacking part of this helicase domain and called

DICER N1 (Kennedy et al., 2015) (Fig S3B). After confirmation of the expression of the

proteins by western blot analysis (Fig S3C), we tested the interactions between DICER and

TRBP, PACT or PKR. We co-transfected the Venus constructs for 24 h and then infected cells

or not with SINV for 6 h at an MOI of 2. A fluorescent signal was observed both in mock- and

SINV-infected cells when [N-ter]Venus:DICER was co-transfected with any of the three proteins

(Fig 3C). Whereas the signal was similar in mock- and virus-infected cells for the DICER-

TRBP and DICER-PACT interactions, we observed an increase of the Venus signal for the

DICER-PKR interaction in SINV-infected cells compared to mock. These results confirm the

previous observations showing an enhanced interaction between DICER and PKR during virus

infection. To gain more insight into the subcellular localization of these interactions during

SINV infection, we performed the BiFC experiments, fixed the cells and observed them under

a confocal microscope. We observed a cytoplasmic signal for DICER-TRBP and DICER-

PACT interactions (Fig 3D upper and middle panels), which is in agreement with their

canonical localization for the maturation of most miRNAs (Bernstein et al., 2001; Hutvágner

et al., 2001). Similarly, co-transfection of DICER and PKR led to a strong Venus signal

homogeneously distributed in the cytoplasm (Fig 3D lower panel). As a control and to rule out

any aspecific interactions between the different proteins tested, we also monitored the DICER-

N1-TRBP interaction by BiFC. As expected, no fluorescent signal was observed in cells co-

transfected with [N-ter]Venus:DICER N1 and TRBP:Venus[C-ter] (Fig S3D), confirming that DICER

helicase domain is required for its interaction with TRBP (Daniels et al., 2009).

Collectively, these results formally confirm that DICER interacts with several RNA

helicases and dsRNA-binding proteins in virus-infected cells, among which PKR, and that for

the latter this interaction occurs in the cytoplasm.

The helicase domain of DICER is required for its interaction with PKR

We next determined the domain of DICER required for its interaction with PKR. Since its

helicase domain was previously shown to be involved in the interaction with TRBP and PACT

(Daniels et al., 2009; Lee et al., 2006), we speculated that it could also be implicated in binding

PKR. To test this hypothesis, we cloned several versions of DICER proteins wholly or partly

deleted of the helicase domain (Fig 4A DICER N1 and N3). In addition, we also cloned the

helicase domain alone (Fig 4A DICER Hel.) and a DICER variant deleted of its C-terminal

dsRNA binding domain (Fig 4A DICER ΔdsRBD) since this domain could also be involved in

protein-protein interaction (Doyle et al., 2013; Lambert et al., 2016). We then transfected the

different versions of DICER WT and the deletion mutant constructs in NoDice cells. After

SINV-GFP infection, whole cell extracts were subjected to HA IP. As for TRBP and to a lesser

extent for PACT, we observed that N1 and N3 mutations strongly reduced the binding of

DICER with PKR (Fig 4B lanes 2-3 and 7-8). Importantly, we noted that the helicase domain

alone could bind PKR, TRBP and PACT (Fig 4B lanes 4 and 9). Moreover, the deletion of the

dsRNA binding domain of DICER did not affect its interaction with TRBP, PACT and PKR

(Fig 4B lanes 5 and 10). Upon virus infection, PKR is phosphorylated to be activated and exert

its antiviral function (Lemaire et al., 2008). Using an antibody targeting the phosphorylated

form of PKR (p-PKR), we looked for p-PKR in our co-IP (Fig 4B panel p-PKR). We noticed

that only WT DICER and its helicase domain were able to interact with p-PKR (Fig 4B lanes

1-6 and 4-9). The fact that DICER ΔdsRBD did not interact with p-PKR (Fig 4B lanes 5-10) is

striking but could indicate that the phosphorylation of PKR may induce conformation changes

preventing its interaction with some domains of DICER. These results reveal that like for TRBP

and PACT, the helicase domain of DICER is required for DICER-PKR/p-PKR interaction

during SINV infection.

In order to confirm the co-IP experiment, we next decided to perform BiFC experiments

using the same conditions as previously. In both mock and SINV-infected cells, only the

combinations of DICER WT-PKR and DICER ΔdsRBD-PKR showed a strong Venus signal,

while neither DICER N1 nor N3 constructs revealed an interaction with PKR (Fig 4C). In

contrast, the DICER Hel. construct did not seem to interact with PKR in mock-infected cells

but appeared to do so in SINV-infected cells as a faint Venus signal could be observed. These

results therefore confirmed the co-IP observations for the DICER-PKR interaction. In addition,

we also performed a BiFC experiment using the different DICER constructs with TRBP or

PACT. Altogether, the BiFC results mostly fitted with the co-IP experiment for the DICER-

TRBP (Fig S4A) and DICER-PACT (Fig S4B) interactions. TRBP indeed did not seem to

interact with the DICER N1 and only slightly with the DICER N3. However, PACT interaction

was lost with DICER N1, but not with DICER N3 in mock- and SINV-infected cells (Fig S4B

third panel). This result may be explained by the fact that DICER interacts with PACT via the

helicase and DUF domains, whereas only the DICER helicase domain is required for its

interaction with TRBP (Daniels et al., 2009; Lee et al., 2006). In agreement, the Venus signal

observed between the DICER Hel. and PACT seemed weaker than the one we observed with

TRBP (Fig S4A and B fourth panels).

Taken together these results indicate that DICER interacts with both PKR and its

phosphorylated form during SINV infection and that this interaction requires the helicase

domain of DICER.

Functional importance of DICER helicase domain during SINV infection

We finally sought to study the functional role of DICER-PKR interaction during viral infection.

For this purpose, we decided to use DICER N1 to study SINV infection. To do so, we generated

by lentiviral transduction NoDice HEK 293T cells stably expressing FHA-tagged DICER N1

(FHA:DICER N1). As for the FHA:DICER cell line, we first selected a clone expressing the

tagged DICER N1 at a level similar level to the endogenous one in HEK 293T cells (Fig 5A).

DICER N1 protein has been shown to still be able to produce miRNAs (Kennedy et al., 2015).

We thus verified by northern blot analysis the capacity of the FHA:DICER N1 cell line to

produce mature miRNAs. The results indicate that indeed DICER N1 is still able to process

miRNAs similarly to WT DICER in HEK 293T and FHA:DICER cells, thereby validating the

functionality of the tagged protein (Fig 5B). We next infected HEK 293T, NoDice,

FHA:DICER and FHA:DICER N1 cells with SINV-GFP and measured virus accumulation by

assessing GFP protein level and virion production (Fig 5C and D and E). As in NoDice cells,

the GFP protein level was drastically reduced in FHA:DICER N1 cells compared to

FHA:DICER and HEK 293T cells (Fig 5C and D). In addition, we observed a statistically

significant reduction in viral titer in both NoDice and FHA:DICER N1 cells compared to HEK

293T or FHA:DICER cells (Fig 5E). This indicates that FHA:DICER N1, lacking the Hel 1 and

Hel 2i domains, is not able to restore the defects observed in NoDice cells.

Moreover, these results show that the antiviral effect observed in NoDice cells is

probably not due to a default in miRNA production but more likely to a loss of interaction

between DICER and PKR.

Discussion

The role of DICER in antiviral defense in human cells remains a topic of intense discussion

(Cullen et al., 2013; Maillard et al., 2013, 2019; Cullen, 2017). In particular there have been

contradictory reports regarding its capacity to produce siRNAs from viral RNAs (Hal P. Bogerd

et al., 2014; Parameswaran et al., 2010; Donaszi-Ivanov et al., 2013; Backes et al., 2014). These

observations could be due to the fact that several mammalian viruses potentially encode VSR

proteins, thereby masking the effect of RNAi (Li et al., 2016, 2013; Maillard et al., 2013; Qiu

et al., 2020, 2017). Another putative but non-exclusive explanation could be that there is a

mutual regulation of type I IFN and RNAi pathways (Berkhout, 2018; Takahashi and Ui-Tei,

2020). Thus, it has already been shown that PACT can regulate MDA5 and RIG-I during virus

infection and therefore the induction of type I IFN response (Kok et al., 2011; Lui et al., 2017).

To date, it is not clear whether the activity of the DICER protein as well could be regulated by

potential interactors, or inversely whether it could itself modulate the activity of proteins

involved in the IFN pathway. To answer this question, we determined the changes in the interactome of human DICER upon SINV and SFV infection. This analysis allowed us to reveal that a lot of proteins associating with DICER during viral infection are dsRNA-binding proteins and RNA helicases. A number of these proteins are known to be involved in antiviral defense pathways, thereby indicating the possible formation of one or several complexes between DICER and these proteins, which are very likely brought together by the accumulation of dsRNA during virus infection.

Among these proteins, we chose to focus on the well-known ISG PKR, which is involved in many cellular pathways such as apoptosis, cellular differentiation, development and antiviral defense (Clemens, 1997; Gal-Ben-Ari et al., 2018; Kim et al., 2014; Lemaire et al., 2008). PKR is one of the main actors of the Integrative Stress Response (ISR) in human cells, and its activation or inhibition needs to be tightly regulated in order to have a properly balanced response to stress. Our results indicate that DICER interacts via its helicase domain with PKR in the cytoplasm during SINV infection. The helicase domain of DICER, which is also required for its interaction with TRBP and PACT, belongs to the helicase superfamily 2, which is also found in RLRs such as RIG-I, MDA5 or LGP2 (Ahmad and Hur, 2015; Fairman-Williams et al., 2010). These proteins act as sensors of viral infection and through the activation of proteins such as MAVS, mediate the induction of type I IFN pathway (Ahmad and Hur, 2015). We hypothesize that even though the human DICER helicase has evolved mainly to act in miRNA/siRNA pathway, it still retained the capacity to act as an RLR. However, as opposed to RIG-I and MDA5, our data suggest that DICER would act more as an inhibitor rather than inducer of the immune response. Therefore, we propose that this domain serves as a platform for the recruitment of different proteins to diversify the function of DICER.

One such regulatory effect appears to be on the antiviral activity of PKR, as cells lacking DICER or expressing a truncated form of DICER unable to interact with PKR are less

permissive to SINV infection. This is in agreement with previous observations that ectopic

expression of the *Drosophila* DICER2 protein in human cells perturbs IFN signaling pathways

and antagonizes PKR-mediated antiviral immunity (Girardi et al., 2015). The exact mechanism

involved is still unclear and will require further work, but it seems that it is not solely due to a

competition of the two proteins for dsRNA binding. Indeed, DICER and PKR interact in BiFC

assay, a technique that favors the detection of direct interactions (Lepur et al., 2016). Moreover,

the DICER helicase domain alone can interact with PKR suggesting that this interaction is

independent of the dsRNA substrates. Most of the time, the inhibition of PKR activity relies on

its inhibition to bind to dsRNA or to auto-phosphorylate. For example, the human tRNA-

dihydrouridine synthase 2 (hDus2) binds the first dsRBD of PKR and prevents its activation

(Mittelstadt et al., 2008). TRBP binds dsRNAs but also PKR directly hindering its dimerization.

In normal condition, TRBP is also associated with PACT thus preventing PKR activation by

PACT (Park et al., 1994; Nakamura et al., 2015; Daher et al., 2009; Singh et al., 2011). Since

we showed that DICER can bind the activated phospho-PKR, we hypothesize that this

interaction does not result in the inhibition of PKR autophosphorylation. Therefore, one

possibility could be that DICER interaction with PKR prevents the latter from acting upon some

of its targets, which remain to be identified, to fine-tune the antiviral response.

As of now, we cannot formally rule out that the effect of DICER on PKR is mediated

by other proteins. TRBP and PACT have been shown to regulate PKR activity, the former

normally acting as a repressor and the latter as an activator (Nakamura et al., 2015; Patel et al.,

2000; Singh et al., 2011). Interestingly, in lymphocytic Jurkat cells infected by HIV-1, PACT

can also act as a repressor of PKR (Clerzius et al., 2013). It is thus tempting to speculate that

these two proteins participate in the formation of the DICER-PKR complex. However, our

results show that this may not necessarily be the case. Indeed, in the BiFC experiment, the

DICER N3 mutant still interacted with PACT but not with PKR indicating that PACT binding

is not sufficient to confer the association with PKR.

Besides PKR, other proteins were specifically enriched upon viral infection in the

DICER IP. These are also interesting candidates to explain the putative regulatory role of

DICER. Among these proteins, DHX9 and ADAR-1 are especially intriguing. DHX9, also

known as RNA helicase A (RHA), associates with RISC, helping the RISC loading (Robb and

Rana, 2007). Moreover, DHX9 is directly involved in removing toxic dsRNAs from the cell to

prevent their processing by DICER (Aktaş et al., 2017). It has also been implicated in HIV-1

replication and knockdown of DXH9 leads to the production of less infectious HIV-1 virions

(Li et al., 1999; Fujii et al., 2001; Roy et al., 2006). Finally, DXH9 interacts with and is

phosphorylated by PKR in mouse embryonic fibroblasts. This phosphorylation precludes the

association of DHX9 with RNA, thus inhibiting its proviral effect (Sadler et al., 2009). In light

of these observations and ours, we can speculate that the inhibitory effect of DICER on PKR

activity could also be linked to DHX9 phosphorylation. ADAR-1 is one of the well-known

RNA-editing factors (Herbert et al.,1997). ADAR-1 is linked to both miRNA biogenesis (Iizasa

et al., 2010; Ota et al., 2013; Yang et al., 2006, p. 1) and virus infection. Indeed, ADAR-1 has

an antiviral effect against Influenza virus, but most of the time, its depletion leads to a decrease

of the viral titer, as was reported for VSV or HIV-1 (Li et al., 2010; Samuel, 2011). It has been

shown that ADAR-1 and PKR interact directly during HIV-1 infection. This interaction triggers

the inhibition of PKR activation, and thus a reduction of eIF2α phosphorylation leading to an

increase of virus replication (Pfaller et al., 2011; Clerzius et al., 2009). Interestingly, over-

expression of ADAR-1 enhances drastically the replication of the alphaviruses Chikungunya

virus (CHIKV), and Venezuelan equine encephalitis virus (VEEV) most likely by interfering

with the IFN induction (Schoggins et al., 2011).

Deciphering the exact role of human DICER protein during virus infection is a

challenging task and additional studies will be required to get a global picture. Nonetheless, by

assessing the interactome of this protein during SINV infection, we have unveiled a new role

for the helicase domain of DICER in regulating the cellular response to viral infection.

Material and methods

Plasmids, cloning and mutagenesis

Plasmids used for BiFC experiments were a gift from Dr. Oliver Vugrek from the Ruđer

Bošković Institute and described in . The cDNAs of TRBP, PACT and PKR were respectively

amplified from (pcDNA-TRBP Addgene #15666) (Kok et al., 2007), (pcDNA-PACT Addgene

#15667) (Kok et al., 2007), (pSB819-PKR-hum Addgene #20030) (Elde et al., 2009), and

cloned into the four pBiFC vectors by Gateway recombination. DICER N1, N3 Hel. and

ΔdsRBD were generated by PCR mutagenesis from pDONR-DICER described in (Girardi et

al., 2015) and cloned into the four pBiFC and pDEST-FHA vectors by Gateway recombination.

plenti6 FHA-V5 vector was modified from plenti6-V5 gateway vector (Thermo Fisher

scientific V49610) by Gibson cloning. DICER WT, N1 from pDONOR plasmids were cloned

into plenti6 FHA-V5 by Gateway recombination. All primers used are listed in Supp. Table 5.

Cell lines

HEK 293T, HEK 293T/NoDice, and HEK 293T/NoDiceΔPKR cell lines were a gift from Pr.

Bryan Cullen and described in (H. P. Bogerd et al., 2014; Kennedy et al., 2015).

Cell culture and transfection

Cells were maintained in Dulbecco's modified Eagle medium (DMEM, Gibco) supplemented

with 10% fetal bovine serum (FBS, Clontech) in a humidified atmosphere of 5% CO_2 at 37°C.

Transfection was performed using Lipofectamine 2000 (Thermofisher) according to the

manufacturer's instructions.

Lentivirus production and generation of stable cell lines

The lentiviral supernatant from single transfer vector was produced by transfecting HEK293T

cells (ATCC® CRL-3216™) with 20 μg the transfer vector, 15 μg of pMDLg/p RRE and 10

μg of pRSV-Rev packaging plasmids (Addgene # 12251 and Addgene #12253) and the pVSV

envelope plasmid (Addgene #8454) using Lipofectamine 2000 (Thermofisher) reagent

according to manufacturer protocol. Standard DMEM (GIBCO) medium supplemented with

10% Fetal bovine serum (FBS, Gibco) and 100 U/mL of penicillin-Streptomycin (Gibco) was

used for growing HEK293T cells and for lentivirus production. One 10 cm plate of HEK293T

cells at 70-80% confluency was used for the transfection. The medium was replaced 8 hours

post-transfection. After 48 hours the medium containing viral particles was collected and

filtered through a 0.45μm PES filter. The supernatant was directly used for transfection or

stored at -80°C. A 6 well plate of HEK 293T/NoDice cells at 80% confluency was transduced

using 600 μL of lentiviral supernatant either expressing FHADICER, N1 or empty vector,

supplemented with 4 ug/mL polybrene (Sigma) for 6 hours. The transduction media was then

changed with fresh DMEM for 24 hours and the resistant cells clones were selected for about

6 weeks with blasticidin (15 μg/mL) and subsequently maintained under blasticidin selection.

Viral stocks, virus infection

Viral stocks of SINV or SINV-GFP were produced as described in (Girardi et al., 2015). Cells

were infected with SINV or SINV-GFP at an MOI of 0.02 or 2 and samples were collected at

different time points as indicated in the figure legends.

Analysis of viral titer by plaque assay

Vero R cells were seeded in 96-well plates format and were infected with 10-fold serial dilutions infection supernatants for 1 h. Afterwards, the inoculum was removed, and cells were cultured in 2.5% carboxymethyl cellulose for 72 h at 37°C in a humidified atmosphere of 5% CO_2. Plaques were counted manually under the microscope and viral titer was calculated according to the formula: *PFU/mL = #plaques/ (Dilution*Volume of inoculum)*

Western blot analysis

Proteins were extracted from cells and homogenized in 350 μL of lysis buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 5 mM EDTA, 1% Triton X-100, 0.5% SDS and Protease Inhibitor Cocktail (complete Mini; Sigma Aldrich)). Proteins were quantified by the Bradford method and 20 to 30 μg of total protein extract were loaded on 4-20% Mini-PROTEAN® TGX™ Precast Gels (Bio-Rad). After transfer onto nitrocellulose membrane, equal loading was verified by Ponceau staining. For PVDF membrane, equal loading was verified by Coomassie staining after transfer and blotting. Membranes were blocked in 5% milk and probed with the following antibodies: anti-hDicer (1:500, F10 Santa Cruz sc-136979), anti-TRBP (1:500, D-5 Santa Cruz sc-514124), anti-PKR (1:2500, Abcam ab32506), anti-PACT (1:500, Abcam, ab75749), anti-HA (1:10000, Sigma, H9658), anti-DHX9 (1:500, Abcam ab26271), anti-p-eIF2a (1:1000, Ser-52 Santa Cruz sc-601670), anti-hADAR-1 (1:500 Santa Cruz sc-271854) anti-p-PKR (1:1000 Abcam ab81303) anti-GFP (1:10000, Roche 11814460001) and anti-Tubulin (1:10000, Sigma T6557). Detection was performed using Chemiluminescent Substrate (Pierce, Thermofisher) and visualized on a Fusion FX imaging system (Vilber).

RNA extraction and northern blot analysis

Total RNA was extracted using Tri-Reagent Solution (Fisher Scientific; MRC, Inc) according

to the manufacturer's instructions. Northern blotting was performed on 10 μg of total RNA.

RNA was resolved on a 12% urea-acrylamide gel, transferred onto Hybond-NX membrane (GE

Healthcare). RNAs were then chemically cross-linked to the membrane during 90 min at 65°C

using 1-ethyl-3-[3-dimethylaminopropyl]carbodiimide hydrochloride (EDC) (Sigma Aldrich).

Membranes were prehybridized for 30 min in PerfectHyb™ plus (Sigma Aldrich) at 50°C.

Probes consisting of oligodeoxyribonucleotides (see Supplementary Table 2) were 5′-end

labeled using T4 polynucleotide kinase (Thermofisher) with 25 μCi of [γ-32P]dATP. The

labeled probe was hybridized to the blot overnight at 50°C. The blot was then washed twice at

50°C for 20 min (5× SSC/0.1% SDS), followed by an additional wash (1× SSC/0.1% SDS) for

5 min. Northern blots were exposed to phosphorimager plates and scanned using a Bioimager

FLA-7000 (Fuji).

Immunoprecipitation

Immunoprecipitation experiments were carried out either on tagged proteins or on endogenous

proteins.

Tagged proteins: Cells were harvested, washed twice with ice-cold 1× PBS (Gibco®, Life

Technologies), and resuspended in 550 μL of lysis buffer (50 mM Tris-HCl pH 7.5, 140 mM

NaCl, 1.5 mM MgCl$_2$, 0.1% NP-40), supplemented with Complete-EDTA-free Protease

Inhibitor Cocktail (complete Mini; Sigma Aldrich). Cells were lysed by 30 min incubation on

ice and debris were removed by 15 min centrifugation at 2000 g and 4°C. An aliquot of the

cleared lysates (50 μL) was kept aside as protein Input. Samples were divided into equal parts

(250 μL each) and incubated with 15 μL of magnetic microparticles coated with monoclonal

HA or MYC antibodies (MACS purification system, Miltenyi Biotech) at 4°C for 1 h under

rotation (10 rpm). Samples were passed through μ Columns (MACS purification system,

Miltenyi Biotech). The μ Columns were then washed 3 times with 200 µL of lysis buffer and 1

time with 100 µL of washing buffer (20 mM Tris-HCl pH 7.5). To elute the immunoprecipitated

proteins, 95°C pre-warmed 2x Western blot loading buffer (10% glycerol, 4% SDS, 62.5 mM

Tris-HCl pH 6.8, 5% (v/v) 2-β-mercaptoethanol, Bromophenol Blue) was passed through the

μ Columns. Proteins were analyzed by western blotting or by mass spectrometry.

Endogenous proteins: The day before immunoprecipitation, magnetic DynaBeads protein G

(Invitrogen) were prepared: briefly, 160 µL of magnetic DynaBeads protein G (Invitrogen)

were washed three times with lysis buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 5 mM

EDTA, 1% Triton X-100, 0.5% SDS). The beads were then incubated with 4 µg of BSA for 1

h under agitation at room temperature. An aliquot of beads (40 µL) was kept for preclearing

step. The remaining beads were incubated either with 6 µg of PKR antibody or control IgG

overnight at 4°C under rotation.

For the IP: Cells were harvested, washed twice with ice-cold 1× PBS (Gibco®, Life

Technologies) and resuspended in 1 mL of lysis buffer (50 mM Tris-HCl pH 7.5, 150 mM

NaCl, 5 mM EDTA, 1% Triton X-100, 0.5% SDS) supplemented with Complete-EDTA-free

Protease Inhibitor Cocktail (complete Mini; Sigma Aldrich). Cells were lysed by three 15 s

sonication steps followed by 30 min incubation on ice and debris were removed by 15 min

centrifugation at 16000 g and 4°C. After centrifugation, an aliquot was kept aside as protein

Input. Samples were incubated with beads alone for preclearing step for 1 h at 4°C on wheel.

Samples were then put on magnetic racks and the supernatant transferred on new tubes

containing either beads coupled to PKR antibody or to control IgG. Samples were incubated on

wheel for 3 h at 4°C. After incubation, samples were put on magnetic racks and beads were

washed three times with lysis buffer. After removal of supernatant, beads were eluted with 2x

western blot loading buffer and incubated for 10 min at 95°C under agitation. Proteins were

analyzed by western blotting.

BiFC assay

Experiments were carried out in two different ways. For non-fixed cells, NoDiceΔPKR cells

were seeded at the density of 1.2×10^5 cells per well in a 24-well plate. After 16 hours, cells

were transfected with equimolar quantities of each plasmid forming BiFC couples. After 24

hours, cells were infected with SINV at an MOI of 2 and pictures were taken 6 hours post-

infection using ZOE fluorescent cell imager (Bio-Rad). Proteins were collected with lysis buffer

(50 mM Tris-HCl pH 7.5, SDS 0.05%, Triton 1%, 5 mM EDTA, 150 mM NaCl) supplemented

with Complete-EDTA-free Protease Inhibitor Cocktail (complete Mini; Sigma Aldrich), and

subjected to western blot analysis. For fixed cells, NoDice/ΔPKR cells were seeded at the

density of 8.10^4 cells per well in 8-well Millicell® EZ Slides (Merck Millipore), transfected

and infected as described previously. At 6 hours post-infection, cells were fixed with 4%

formaldehyde and 0.2% glutaraldehyde for 10 min. Cells were then washed with $1 \times$ PBS

(Gibco®, Life Technologies) and stained with $10 \, \mu g/\mu L$ DAPI in $1 \times$ PBS solution (Invitrogen)

for 5 min. Fixed cells were mounted on a glass slide with Fluoromount-G mounting media

(Southern Biotech). Images were acquired using confocal LSM780 (Zeiss) inverted microscope

with an argon laser (514x nm) and with $\times$40 immersion oil objective. All pictures obtained

from BiFC experiments were treated using FigureJ software (NIH).

Mass spectrometry analysis

Protein extracts were prepared for mass spectrometry as described in a previous study (Chicois

et al., 2018). Each sample was precipitated with 0.1 M ammonium acetate in 100% methanol,

and proteins were resuspended in 50 mM ammonium bicarbonate. After a reduction-alkylation

step (dithiothreitol 5 mM – iodoacetamide 10 mM), proteins were digested overnight with

sequencing-grade porcine trypsin (1:25, w/w, Promega, Fitchburg, MA, USA). The resulting

vacuum-dried peptides were resuspended in water containing 0.1% (v/v) formic acid (solvent

A). One sixth of the peptide mixtures were analyzed by nanoLC-MS/MS an Easy-nanoLC-1000

system coupled to a Q-Exactive Plus mass spectrometer (Thermo-Fisher Scientific, USA)

operating in positive mode. Five microliters of each sample were loaded on a C-18 precolumn

(75 μm ID × 20 mm nanoViper, 3 μm Acclaim PepMap; Thermo) coupled with the analytical

C18 analytical column (75 μm ID × 25 cm nanoViper, 3 μm Acclaim PepMap; Thermo).

Peptides were eluted with a 160 min gradient of 0.1% formic acid in acetonitrile at 300 nL/min.

The Q-Exactive Plus was operated in data-dependent acquisition mode (DDA) with Xcalibur

software (Thermo-Fisher Scientific). Survey MS scans were acquired at a resolution of 70K at

200 m/z (mass range 350-1250), with a maximum injection time of 20 ms and an automatic

gain control (AGC) set to 3e6. Up to 10 of the most intense multiply charged ions (≥2) were

selected for fragmentation with a maximum injection time of 100 ms, an AGC set at 1e5 and a

resolution of 17.5K. A dynamic exclusion time of 20 s was applied during the peak selection

process.

Database search and mass-spectrometry data post-processing

Data were searched against a database containing Human and Viruses UniProtKB sequences

with a decoy strategy (GFP, Human and Sindbis Virus SwissProt sequences as well as Semliki

Forest Virus SwissProt and TrEMBL sequences (releases from January 2017, 40439

sequences)). Peptides were identified with Mascot algorithm (version 2.3, Matrix Science,

London, UK) with the following search parameters: carbamidomethylation of cysteine was set

as fixed modification; N-terminal protein acetylation, phosphorylation of serine / threonine /

tyrosine and oxidation of methionine were set as variable modifications; tryptic specificity with

up to three missed cleavages was used. The mass tolerances in MS and MS/MS were set to 10

ppm and 0.02 Da respectively, and the instrument configuration was specified as "ESI-Trap".

The   resulting   .dat   Mascot   files   were   then   imported   into   Proline   v1.4   package

(http://proline.profiproteomics.fr/) for post-processing. Proteins were validated with Mascot

pretty rank equal to 1, 1% FDR on both peptide spectrum matches (PSM) and protein sets

(based on score). The total number of MS/MS fragmentation spectra (Spectral count or SpC)

was used for subsequent protein quantification in the different samples.

Exploratory and differential expression analysis of LC-MS/MS data

Mass spectrometry data obtained for each sample were stored in a local MongoDB database

and subsequently analyzed through a Shiny Application built upon the R/Bioconductor

packages msmsEDA (Gregori J, Sanchez A, Villanueva J (2014). msmsEDA: Exploratory Data

Analysis of LC-MS/MS data by spectral counts. R/Bioconductor package version 1.22.0) and

msmsTests (Gregori J, Sanchez A, Villanueva J (2013). msmsTests: LC-MS/MS Differential

Expression Tests. R/Bioconductor package version 1.22.0). Exploratory data analyses of LC-

MS/MS data were thus conducted, and differential expression tests were performed using a

negative binomial regression model. The p-values were adjusted with FDR control by the

Benjamini-Hochberg method and the following criteria were used to define differentially

expressed proteins: an adjusted p-value < 0.05, a minimum of 5 SpC in the most abundant

condition, and a minimum fold change of 2 (abs(LogFC) > 1).

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange

Consortium via the PRIDE (Perez-Riverol et al., 2019) partner repository with the dataset

identifier PXD019093 and 10.6019/PXD019093.

Acknowledgments

The authors would like to thank members of the Pfeffer laboratory for discussion, Pr. Bryan Cullen for the kind gift of the HEK293T NoDice and NoDice/ΔPKR cell lines and Dr. Oliver Vugrek for the BiFC plasmids.

This work was funded by the European Research Council (ERC-CoG-647455 RegulRNA) and was performed under the framework of the LABEX: ANR-10-LABX-0036_NETRNA, which benefits from a funding from the state managed by the French National Research Agency as part of the Investments for the future program. This work has also received funding from the People Programme (Marie Curie Actions) of the European Union's Seventh Framework Program (FP7/2007-2013) under REA grant agreement n° PCOFUND-GA-2013-609102, through the PRESTIGE program coordinated by Campus France (to EG), and from the French Minister for Higher Education, Research and Innovation (PhD contract to MB). The mass spectrometry instrumentation was funded by the University of Strasbourg, IdEx "Equipement mi-lourd" 2015.

Authors contributions

SP and ML conceived the project. TCM, ML, EG and SP designed the work. TCM, ML, MB, EG and MM performed the experiments and analyzed the results. PH and JC performed the mass spectrometry analysis. BCWM designed softwares and analyzed the mass spectrometry results. SP coordinated the work and assured funding. TCM, MB and SP wrote the manuscript with input from the other authors. All authors reviewed and approved the final manuscript.

References

Adiliaghdam F, Basavappa M, Saunders TL, Harjanto D, Prior JT, Cronkite DA, Papavasiliou N, Jeffrey KL. 2020. A Requirement for Argonaute 4 in Mammalian Antiviral Defense. *Cell Rep* **30**:1690-1701.e4. doi:10.1016/j.celrep.2020.01.021

Ahmad S, Hur S. 2015. Helicases in Antiviral Immunity: Dual Properties as Sensors and Effectors. *Trends Biochem Sci* **40**:576–585. doi:10.1016/j.tibs.2015.08.001

Aktaş T, Avşar Ilık İ, Maticzka D, Bhardwaj V, Pessoa Rodrigues C, Mittler G, Manke T, Backofen R, Akhtar A. 2017. DHX9 suppresses RNA processing defects originating from the Alu invasion of the human genome. *Nature* **544**:115–119. doi:10.1038/nature21715

Backes S, Langlois RA, Schmid S, Varble A, Shim JV, Sachs D, tenOever BR. 2014. The Mammalian Response to Virus Infection Is Independent of Small RNA Silencing. *Cell Rep* **8**:114–125. doi:10.1016/j.celrep.2014.05.038

Bartel DP. 2018. Metazoan MicroRNAs. *Cell* **173**:20–51. doi:10.1016/j.cell.2018.03.006

Berkhout B. 2018. RNAi-mediated antiviral immunity in mammals. *Curr Opin Virol*, Engineering for viral resistance • Viruses and cancer **32**:9–14. doi:10.1016/j.coviro.2018.07.008

Bernstein E, Caudy AA, Hammond SM, Hannon GJ. 2001. Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature* **409**:363–6.

Bogerd Hal P., Skalsky RL, Kennedy EM, Furuse Y, Whisnant AW, Flores O, Schultz KLW, Putnam N, Barrows NJ, Sherry B, Scholle F, Garcia-Blanco MA, Griffin DE, Cullen BR. 2014. Replication of many human viruses is refractory to inhibition by endogenous cellular microRNAs. *J Virol* **88**:8065–8076. doi:10.1128/JVI.00985-14

Bogerd H. P., Whisnant AW, Kennedy EM, Flores O, Cullen BR. 2014. Derivation and characterization of Dicer- and microRNA-deficient human cells. *RNA* **20**:923–937. doi:10.1261/rna.044545.114

Borden EC, Sen GC, Uze G, Silverman RH, Ransohoff RM, Foster GR, Stark GR. 2007. Interferons at age 50: past, current and future impact on biomedicine. *Nat Rev Drug Discov* **6**:975–990. doi:10.1038/nrd2422

Chen EY, Tan CM, Kou Y, Duan Q, Wang Z, Meirelles GV, Clark NR, Ma'ayan A. 2013. Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics* **14**:128. doi:10.1186/1471-2105-14-128

Chendrimada TP, Gregory RI, Kumaraswamy E, Norman J, Cooch N, Nishikura K, Shiekhattar
R. 2005. TRBP recruits the Dicer complex to Ago2 for microRNA processing and gene
silencing. *Nature* **436**:740–4.

Chicois C, Scheer H, Garcia S, Zuber H, Mutterer J, Chicher J, Hammann P, Gagliardi D,
Garcia D. 2018. The UPF1 interactome reveals interaction networks between RNA
degradation and translation repression factors in Arabidopsis. *Plant J* **96**:119–132.
doi:10.1111/tpj.14022

Clemens MJ. 1997. PKR-a protein kinase regulated by double-stranded RNA. *Int J Biochem*
Cell Biol **29**:945–9.

Clerzius G, Gélinas J-F, Daher A, Bonnet M, Meurs EF, Gatignol A. 2009. ADAR1 interacts
with PKR during human immunodeficiency virus infection of lymphocytes and
contributes to viral replication. *J Virol* **83**:10119–10128. doi:10.1128/JVI.02457-08

Clerzius G, Shaw E, Daher A, Burugu S, Gélinas J-F, Ear T, Sinck L, Routy J-P, Mouland AJ,
Patel RC, Gatignol A. 2013. The PKR activator, PACT, becomes a PKR inhibitor during
HIV-1 replication. *Retrovirology* **10**:96. doi:10.1186/1742-4690-10-96

Cullen BR. 2017. RNA Interference in Mammals: The Virus Strikes Back. *Immunity* **46**:970–
972. doi:10.1016/j.immuni.2017.05.004

Cullen BR. 2006. Is RNA interference involved in intrinsic antiviral immunity in mammals?
Nat Immunol **7**:563–7.

Cullen BR, Cherry S, tenOever BR. 2013. Is RNA Interference a Physiologically Relevant
Innate Antiviral Immune Response in Mammals? *Cell Host Microbe* **14**:374–378.
doi:10.1016/j.chom.2013.09.011

Daher A, Laraki G, Singh M, Melendez-Peña CE, Bannwarth S, Peters AHFM, Meurs EF,
Braun RE, Patel RC, Gatignol A. 2009. TRBP control of PACT-induced
phosphorylation of protein kinase R is reversed by stress. *Mol Cell Biol* **29**:254–265.
doi:10.1128/MCB.01030-08

Daniels SM, Melendez-Peña CE, Scarborough RJ, Daher A, Christensen HS, El Far M, Purcell
DFJ, Lainé S, Gatignol A. 2009. Characterization of the TRBP domain required for
dicer interaction and function in RNA interference. *BMC Mol Biol* **10**:38.
doi:10.1186/1471-2199-10-38

Donaszi-Ivanov A, Mohorianu I, Dalmay T, Powell PP. 2013. Small RNA Analysis in Sindbis
Virus Infected Human HEK293 Cells. *PLoS ONE* **8**:e84070.
doi:10.1371/journal.pone.0084070

Doyle M, Badertscher L, Jaskiewicz L, Güttinger S, Jurado S, Hugenschmidt T, Kutay U,
Filipowicz W. 2013. The double-stranded RNA binding domain of human Dicer
functions as a nuclear localization signal. *RNA N Y N* **19**:1238–1252.
doi:10.1261/rna.039255.113

Elde NC, Child SJ, Geballe AP, Malik HS. 2009. Protein kinase R reveals an evolutionary
model for defeating viral mimicry. *Nature* **457**:485–489. doi:10.1038/nature07529

Fairman-Williams ME, Guenther U-P, Jankowsky E. 2010. SF1 and SF2 helicases: family
matters. *Curr Opin Struct Biol* **20**:313–324. doi:10.1016/j.sbi.2010.03.011

Fros JJ, Pijlman GP. 2016. Alphavirus Infection: Host Cell Shut-Off and Inhibition of Antiviral
Responses. *Viruses* **8**:166. doi:10.3390/v8060166

Fujii R, Okamoto M, Aratani S, Oishi T, Ohshima T, Taira K, Baba M, Fukamizu A, Nakajima
T. 2001. A Role of RNA Helicase A in cis-Acting Transactivation Response Element-
mediated Transcriptional Regulation of Human Immunodeficiency Virus Type 1. *J Biol*
Chem **276**:5445–5451. doi:10.1074/jbc.M006892200

Gal-Ben-Ari S, Barrera I, Ehrlich M, Rosenblum K. 2018. PKR: A Kinase to Remember. *Front*
Mol Neurosci **11**:480. doi:10.3389/fnmol.2018.00480

García MA, Meurs EF, Esteban M. 2007. The dsRNA protein kinase PKR: Virus and cell
control. *Biochimie* **89**:799–811. doi:10.1016/j.biochi.2007.03.001

Girardi E, Chane-Woon-Ming B, Messmer M, Kaukinen P, Pfeffer S. 2013. Identification of
RNase L-dependent, 3'-end-modified, viral small RNAs in Sindbis virus-infected
mammalian cells. *mBio* **4**:e00698-00613. doi:10.1128/mBio.00698-13

Girardi E, Lefèvre M, Chane-Woon-Ming B, Paro S, Claydon B, Imler J-L, Meignin C, Pfeffer
S. 2015. Cross-species comparative analysis of Dicer proteins during Sindbis virus
infection. *Sci Rep* **5**:10693. doi:10.1038/srep10693

Griffin D. 2007. Alphaviruses, p 1023–1067. *Fields Virol 5th Ed Lippincott Williams Wilkins*
Phila PA.

Guo Z, Li Y, Ding S-W. 2019. Small RNA-based antimicrobial immunity. *Nat Rev Immunol*
19:31–44. doi:10.1038/s41577-018-0071-x

Haase AD, Jaskiewicz L, Zhang H, Laine S, Sack R, Gatignol A, Filipowicz W. 2005. TRBP,
a regulator of cellular PKR and HIV-1 virus expression, interacts with Dicer and
functions in RNA silencing. *EMBO Rep* **6**:961–7.

Herbert A, Alfken J, Kim YG, Mian IS, Nishikura K, Rich A. 1997. A Z-DNA binding domain
present in the human editing enzyme, double-stranded RNA adenosine deaminase. *Proc*
Natl Acad Sci U S A **94**:8421–8426. doi:10.1073/pnas.94.16.8421

Heyam A, Lagos D, Plevin M. 2015. Dissecting the roles of TRBP and PACT in double-
stranded RNA recognition and processing of noncoding RNAs. *WIREs RNA* **6**:271–289.
doi:10.1002/wrna.1272

Hutvágner G, McLachlan J, Pasquinelli AE, Balint E, Tuschl T, Zamore PD. 2001. A cellular
function for the RNA-interference enzyme Dicer in the maturation of the let-7 small
temporal RNA. *Science* **293**:834–8.

Iizasa H, Wulff B-E, Alla NR, Maragkakis M, Megraw M, Hatzigeorgiou A, Iwakiri D, Takada
K, Wiedmer A, Showe L, Lieberman P, Nishikura K. 2010. Editing of Epstein-Barr
virus-encoded BART6 microRNAs controls their dicer targeting and consequently
affects viral latency. *J Biol Chem* **285**:33358–33370. doi:10.1074/jbc.M110.138362

Ivashkiv LB, Donlin LT. 2014. Regulation of type I interferon responses. *Nat Rev Immunol*
14:36–49. doi:10.1038/nri3581

Kennedy EM, Whisnant AW, Kornepati AVR, Marshall JB, Bogerd HP, Cullen BR. 2015.
Production of functional small interfering RNAs by an amino-terminal deletion mutant
of human Dicer. *Proc Natl Acad Sci* **112**:E6945–E6954. doi:10.1073/pnas.1513421112

Kim Y, Lee JH, Park J-E, Cho J, Yi H, Kim VN. 2014. PKR is activated by cellular dsRNAs
during mitosis and acts as a mitotic regulator. *Genes Dev* **28**:1310–1322.
doi:10.1101/gad.242644.114

Kok K-H, Lui P-Y, Ng M-HJ, Siu K-L, Au SWN, Jin D-Y. 2011. The Double-Stranded RNA-
Binding Protein PACT Functions as a Cellular Activator of RIG-I to Facilitate Innate
Antiviral Response. *Cell Host Microbe* **9**:299–309. doi:10.1016/j.chom.2011.03.007

Kok KH, Ng MH, Ching YP, Jin DY. 2007. Human TRBP and PACT Directly Interact with
Each Other and Associate with Dicer to Facilitate the Production of Small Interfering
RNA. *J Biol Chem* **282**:17649–57.

Kuleshov MV, Jones MR, Rouillard AD, Fernandez NF, Duan Q, Wang Z, Koplev S, Jenkins
SL, Jagodnik KM, Lachmann A, McDermott MG, Monteiro CD, Gundersen GW,
Ma'ayan A. 2016. Enrichr: a comprehensive gene set enrichment analysis web server
2016 update. *Nucleic Acids Res* **44**:W90-97. doi:10.1093/nar/gkw377

Lambert M, Pépin G, Peralta-Zaragoza O, Matusiak R, Ly S, Landry P, Provost P. 2016.
TWEAK Negatively Regulates Human Dicer. *Non-Coding RNA* **2**.
doi:10.3390/ncrna2040012

Lee Y, Hur I, Park SY, Kim YK, Suh MR, Kim VN. 2006. The role of PACT in the RNA
silencing pathway. *Embo J* **25**:522–32.

Lemaire PA, Anderson E, Lary J, Cole JL. 2008. Mechanism of PKR Activation by dsRNA. *J*
Mol Biol **381**:351–360. doi:10.1016/j.jmb.2008.05.056

Lepur A, Kovačević L, Belužić R, Vugrek O. 2016. Combining unique multiplex gateway
cloning and bimolecular fluorescence complementation (BiFC) for high-throughput
screening of protein–protein interactions. *J Biomol Screen* **21**:1100–1111.

Li J, Tang H, Mullen TM, Westberg C, Reddy TR, Rose DW, Wong-Staal F. 1999. A role for
RNA helicase A in post-transcriptional regulation of HIV type 1. *Proc Natl Acad Sci U*
S A **96**:709–714. doi:10.1073/pnas.96.2.709

Li Y, Basavappa M, Lu J, Dong S, Cronkite DA, Prior JT, Reinecker H-C, Hertzog P, Han Y,
Li W-X, Cheloufi S, Karginov FV, Ding S-W, Jeffrey KL. 2016. Induction and
suppression of antiviral RNA interference by influenza A virus in mammalian cells. *Nat*
Microbiol **2**:16250. doi:10.1038/nmicrobiol.2016.250

Li Y, Lu J, Han Y, Fan X, Ding S-W. 2013. RNA interference functions as an antiviral
immunity mechanism in mammals. *Science* **342**:231–234.
doi:10.1126/science.1241911

Li Z, Wolff KC, Samuel CE. 2010. RNA adenosine deaminase ADAR1 deficiency leads to
increased activation of protein kinase PKR and reduced vesicular stomatitis virus
growth following interferon treatment. *Virology* **396**:316–322.
doi:10.1016/j.virol.2009.10.026

López P, Girardi E, Mounce BC, Weiss A, Chane-Woon-Ming B, Messmer M, Kaukinen P,
Kopp A, Bortolamiol-Becet D, Fendri A, Vignuzzi M, Brino L, Pfeffer S. 2020. High-
Throughput Fluorescence-Based Screen Identifies the Neuronal MicroRNA miR-124 as
a Positive Regulator of Alphavirus Infection. *J Virol* **94**. doi:10.1128/JVI.02145-19

Lui P-Y, Wong L-YR, Ho T-H, Au SWN, Chan C-P, Kok K-H, Jin D-Y. 2017. PACT
Facilitates RNA-Induced Activation of MDA5 by Promoting MDA5 Oligomerization.
J Immunol **199**:1846–1855. doi:10.4049/jimmunol.1601493

Maillard PV, Ciaudo C, Marchais A, Li Y, Jay F, Ding SW, Voinnet O. 2013. Antiviral RNA
interference in mammalian cells. *Science* **342**:235–238. doi:10.1126/science.1241930

Maillard PV, van der Veen AG, Poirier EZ, Reis e Sousa C. 2019. Slicing and dicing viruses:
antiviral RNA interference in mammals. *EMBO J* **38**. doi:10.15252/embj.2018100941

Maillard PV, Veen AGV der, Deddouche-Grass S, Rogers NC, Merits A, Sousa CR e. 2016.
Inactivation of the type I interferon pathway reveals long double-stranded RNA-

mediated RNA interference in mammalian cells. *EMBO J* **35**:2505–2518.
doi:10.15252/embj.201695086

Meister G, Tuschl T. 2004. Mechanisms of gene silencing by double-stranded RNA. *Nature*
431:343–9.

Mittelstadt M, Frump A, Khuu T, Fowlkes V, Handy I, Patel CV, Patel RC. 2008. Interaction
of human tRNA-dihydrouridine synthase-2 with interferon-induced protein kinase
PKR. *Nucleic Acids Res* **36**:998–1008. doi:10.1093/nar/gkm1129

Nakamura T, Kunz RC, Zhang C, Kimura T, Yuan CL, Baccaro B, Namiki Y, Gygi SP,
Hotamisligil GS. 2015. A Critical Role for PKR Complexes with TRBP in
Immunometabolic Regulation and eIF2α Phosphorylation in Obesity. *Cell Rep* **11**:295–
307. doi:10.1016/j.celrep.2015.03.021

Ota H, Sakurai M, Gupta R, Valente L, Wulff B-E, Ariyoshi K, Iizasa H, Davuluri RV,
Nishikura K. 2013. ADAR1 forms a complex with Dicer to promote microRNA
processing   and   RNA-induced   gene   silencing.   *Cell*   **153**:575–589.
doi:10.1016/j.cell.2013.03.024

Parameswaran P, Sklan E, Wilkins C, Burgon T, Samuel MA, Lu R, Ansel KM, Heissmeyer
V, Einav S, Jackson W, Doukas T, Paranjape S, Polacek C, dos Santos FB, Jalili R,
Babrzadeh F, Gharizadeh B, Grimm D, Kay M, Koike S, Sarnow P, Ronaghi M, Ding
SW, Harris E, Chow M, Diamond MS, Kirkegaard K, Glenn JS, Fire AZ. 2010. Six
RNA viruses and forty-one hosts: viral small RNAs and modulation of small RNA
repertoires in vertebrate and invertebrate systems. *PLoS Pathog* **6**:e1000764.
doi:10.1371/journal.ppat.1000764

Park H, Davies MV, Langland JO, Chang HW, Nam YS, Tartaglia J, Paoletti E, Jacobs BL,
Kaufman RJ, Venkatesan S. 1994. TAR RNA-binding protein is an inhibitor of the
interferon-induced protein kinase PKR. *Proc Natl Acad Sci U S A* **91**:4713–4717.
doi:10.1073/pnas.91.11.4713

Patel CV, Handy I, Goldsmith T, Patel RC. 2000. PACT, a Stress-modulated Cellular Activator
of Interferon-induced Double-stranded RNA-activated Protein Kinase, PKR. *J Biol*
Chem **275**:37993–37998. doi:10.1074/jbc.M004762200

Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S, Kundu DJ, Inuganti
A, Griss J, Mayer G, Eisenacher M, Pérez E, Uszkoreit J, Pfeuffer J, Sachsenberg T,
Yilmaz S, Tiwary S, Cox J, Audain E, Walzer M, Jarnuczak AF, Ternent T, Brazma A,
Vizcaíno JA. 2019. The PRIDE database and related tools and resources in 2019:

improving support for quantification data. *Nucleic Acids Res* **47**:D442–D450.

doi:10.1093/nar/gky1106

Pfaller CK, Li Z, George CX, Samuel CE. 2011. Protein kinase PKR and RNA adenosine

deaminase ADAR1: new roles for old players as modulators of the interferon response.

Curr Opin Immunol, Special section: Cytokines/Immunogenetics and transplantation

23:573–582. doi:10.1016/j.coi.2011.08.009

Qiu Y, Xu Y, Zhang Y, Zhou H, Deng Y-Q, Li X-F, Miao M, Zhang Q, Zhong B, Hu Y, Zhang

F-C, Wu L, Qin C-F, Zhou X. 2017. Human Virus-Derived Small RNAs Can Confer

Antiviral Immunity in Mammals. *Immunity* **46**:992-1004.e5.

doi:10.1016/j.immuni.2017.05.006

Qiu Y, Xu Y-P, Wang M, Miao M, Zhou H, Xu J, Kong J, Zheng D, Li R-T, Zhang R-R. 2020.

Flavivirus induces and antagonizes antiviral RNA interference in both mammals and

mosquitoes. *Sci Adv* **6**:eaax7989.

Robb GB, Rana TM. 2007. RNA helicase A interacts with RISC in human cells and functions

in RISC loading. *Mol Cell* **26**:523–537. doi:10.1016/j.molcel.2007.04.016

Roy BB, Hu J, Guo X, Russell RS, Guo F, Kleiman L, Liang C. 2006. Association of RNA

helicase a with human immunodeficiency virus type 1 particles. *J Biol Chem*

281:12625–12635. doi:10.1074/jbc.M510596200

Ryman KD, White LJ, Johnston RE, Klimstra WB. 2002. Effects of PKR/RNase L-Dependent

and Alternative Antiviral Pathways on Alphavirus Replication and Pathogenesis. *Viral*

Immunol **15**:53–76. doi:10.1089/088282402317340233

Sadler AJ, Latchoumanin O, Hawkes D, Mak J, Williams BRG. 2009. An Antiviral Response

Directed by PKR Phosphorylation of the RNA Helicase A. *PLoS Pathog* **5**:e1000311.

doi:10.1371/journal.ppat.1000311

Samuel CE. 2011. Adenosine deaminases acting on RNA (ADARs) are both antiviral and

proviral. *Virology* **411**:180–193. doi:10.1016/j.virol.2010.12.004

Schoggins JW, Wilson SJ, Panis M, Murphy MY, Jones CT, Bieniasz P, Rice CM. 2011. A

diverse range of gene products are effectors of the type I interferon antiviral response.

Nature **472**:481–485. doi:10.1038/nature09907

Schuster S, Overheul GJ, Bauer L, van Kuppeveld FJM, van Rij RP. 2019. No evidence for

viral small RNA production and antiviral function of Argonaute 2 in human cells. *Sci*

Rep **9**:1–11. doi:10.1038/s41598-019-50287-w

Singh M, Castillo D, Patel CV, Patel RC. 2011. Stress-induced phosphorylation of PACT
reduces its interaction with TRBP and leads to PKR activation. *Biochemistry* **50**:4550–
4560. doi:10.1021/bi200104h

Takahashi T, Nakano Y, Onomoto K, Murakami F, Komori C, Suzuki Y, Yoneyama M, Ui-Tei
K. 2018a. LGP2 virus sensor regulates gene expression network mediated by TRBP-
bound microRNAs. *Nucleic Acids Res* **46**:9134–9147. doi:10.1093/nar/gky575

Takahashi T, Nakano Y, Onomoto K, Yoneyama M, Ui-Tei K. 2018b. Virus Sensor RIG-I
Represses RNA Interference by Interacting with TRBP through LGP2 in Mammalian
Cells. *Genes* **9**:511. doi:10.3390/genes9100511

Takahashi T, Ui-Tei K. 2020. Mutual Regulation of RNA Silencing and the IFN Response as
an Antiviral Defense System in Mammalian Cells. *Int J Mol Sci* **21**:1348.
doi:10.3390/ijms21041348

tenOever BR. 2016. The Evolution of Antiviral Defense Systems. *Cell Host Microbe* **19**:142–
149. doi:10.1016/j.chom.2016.01.006

Tsai K, Courtney DG, Kennedy EM, Cullen BR. 2018. Influenza A virus-derived siRNAs
increase in the absence of NS1 yet fail to inhibit virus replication. *RNA N Y N*.
doi:10.1261/rna.066332.118

van der Veen AG, Maillard PV, Schmidt JM, Lee SA, Deddouche-Grass S, Borg A, Kjær S,
Snijders AP, Reis e Sousa C. 2018. The RIG-I-like receptor LGP2 inhibits Dicer-
dependent processing of long double-stranded RNA and blocks RNA interference in
mammalian cells. *EMBO J* **37**:e97479. doi:10.15252/embj.201797479

Wienholds E, Koudijs MJ, Van Eeden FJ, Cuppen E, Plasterk RH. 2003. The microRNA-
producing enzyme Dicer1 is essential for zebrafish development. *Nat Genet* 217–218.

Williams BR. 1999. PKR; a sentinel kinase for cellular stress. *Oncogene* **18**:6112–6120.
doi:10.1038/sj.onc.1203127

Yang W, Chendrimada TP, Wang Q, Higuchi M, Seeburg PH, Shiekhattar R, Nishikura K.
2006. Modulation of microRNA processing and expression through RNA editing by
ADAR deaminases. *Nat Struct Mol Biol* **13**:13–21.

884
885

 Figure legends

Figure 1: Molecular characterization of FHA:DICER cell line.

A, Western blot analysis of DICER (DICER and HA), TRBP and AGO2 expression in HEK

293T, NoDice and FHA:DICER cell lines. Alpha-Tubulin was used as loading control. **B,** miR-

16 expression analyzed by northern blot in the same samples as in A. Expression of snRNA U6

was used as loading control. **C,** GFP fluorescent microscopy pictures of HEK 293T, NoDice

and FHA:DICER cell lines infected with SINV-GFP at an MOI of 0.02 for 24 h. The left panel

corresponds to GFP signal from infected cells and the right panel to a merge picture of GFP

signal and brightfield. Pictures were taken with a 5x magnification. hpi: hours post-infection.

D, Representative western blot of DICER (DICER and HA) and GFP expression in SINV-GFP-

infected cells in the same condition as in C. Alpha-Tubulin was used as loading control. **E,**

Mean (+/- SEM) of SINV-GFP viral titers in HEK 293T, NoDice and FHA:DICER cells

infected at an MOI of 0.02 for 24 h (n = 3) from plaque assay quantification. $* p < 0.05$, $** p <$

0.005, ordinary one-way ANOVA test.

Figure 2: LC-MS/MS analysis of DICER interactome during SINV infection.

A, Volcano plot for differentially expressed proteins (DEPs) between HA IP and CTL IP in

FHA:DICER mock-infected cells. Each protein is marked as a dot; proteins that are

significantly up-regulated in HA IP are shown in red, up-regulated proteins in CTL IP are

shown in blue, and non-significant proteins are in black. The horizontal line denotes the

adjusted p-value threshold (0.05) and the vertical lines the Log2 fold change cutoff (-1 and 1).

DICER and its cofactors (TRBP, PACT, AGO2) are highlighted in yellow. **B,** Left panel:

Volcano plot for DEPs between SINV-GFP (MOI of 2, 6 hpi) and mock fractions of HA IP in

FHA:DICER cells. Same colour code and thresholds as in A have been applied. Proteins that

are discussed in the text are highlighted in yellow and SINV proteins in purple. **C,** Gene

Ontology (GO) term enrichment of proteins up-regulated in SINV-GFP fraction of HA IP using

Enrichr software (Chen et al., 2013; Kuleshov et al., 2016). The graph displays the GO term

hierarchy within the "biological process" branch sorted by p-value ranking computed from the

Fisher exact test. The length of each bar represents the significance of that specific term. In

addition, the brighter the colour is, the more significant that term is. Viral proteins have been

excluded for this analysis. **D,** Summary of the differential expression analysis of SINV-GFP vs

mock fractions from HA IP in FHA:DICER cells. The analysis has been performed using a

generalized linear model of a negative-binomial distribution and p-values were corrected for

multiple testing using the Benjamini-Hochberg method.

Figure 3: Confirmation of LC-MS/MS analysis by co-IP and BiFC.

A, Western blot analysis of HA co-IP in mock or SINV-GFP-infected (MOI of 2, 6 hpi)

FHA:DICER cells. Proteins associated to FHA:DICER were revealed by using antibodies

targeting endogenous ADAR-1, PKR, TRBP, DHX9 or PACT proteins. In parallel, an HA

antibody was used to verify the IP efficiency and GFP antibody was used to verify the infection.

Ponceau was used as loading control. **B,** Western blot analysis to validate the interaction of

PKR with PACT (upper panel) and DICER (lower panel) in mock or SINV-GFP-infected HEK

293T cells. Immunoprecipitated proteins obtained from PKR pulldowns were compared to IgG

pulldowns to verify the specificity of the assay. **C,** Interactions between DICER and TRBP,

PKR or PACT was visualized by BiFC. Plasmids expressing $^{N\text{-ter}}$Venus:DICER and TRBP:,

PACT: or PKR:Venus$^{C\text{-ter}}$ were co-transfected in NoDice/ΔPKR cells for 24 h and cells were

either infected with SINV at an MOI of 2 for 6 h or not. The different combinations are indicated

on the left side. Reconstitution of Venus (BiFC) signal was observed under epifluorescence

microscope. For each condition, the left panel corresponds to Venus signal and the right panel

to the corresponding brightfield pictures. Scale bar: 100 μm. **D,** BiFC experiment on fixed

NoDice/ΔPKR cells treated as in C. After fixation, cells were stained with DAPI and observed

under confocal microscope. Only a merge picture of BiFC and DAPI signals of SINV-infected

cells is shown here. A higher magnification of picture showing cytoplasmic localization of the

interaction represented by a red square is shown in the bottom left corner. Scale bar: 20 μm and

10 μm.

Figure 4: Identification of DICER domains involved in DICER-PKR interaction.

A, Schematic representation of Human DICER proteins used in this study. The different

conserved domains are shown in colored boxes. DUF283: Domain of Unknown Function; PAZ:

PIWI ARGONAUTE ZWILLE domain; dsRBD: dsRNA-binding domain. hDICER WT is the

full-length protein. hDICER N1 is deleted of the first N-terminal 495 amino acids. hDICER N3

is wholly deleted of the helicase domain. hDICER Hel. is the whole DICER's helicase domain.

hDICER ΔdsRBD is deleted of the C-terminal dsRBD. **B,** Western blot analysis of HA co-IP

in NoDice cells transfected with different versions of FHA:DICER proteins and infected with

SINV-GFP (MOI of 2, 6 hpi). Efficiency of immunoprecipitation was assessed using an anti-

Flag antibody and co-IPs of PKR, TRBP, PACT and p-PKR were examined using appropriate

antibodies. Expression of GFP in INPUT fraction was visualized as control of SINV-GFP

infection. Ponceau staining of membranes is used as loading control. **C,** Plasmids expressing

the different versions of DICER proteins fused to the N-terminal part of Venus and

PKR:Venus[C-ter] plasmid were co-transfected in NoDice/ΔPKR cells. Cells were treated as in Fig

3C. The different combinations are noted on the left side. The fluorescent signal was observed

using an epifluorescence microscope. For each condition, the left panel corresponds to Venus

signal and the right panel to the corresponding brightfield pictures. Scale bar: 100 μm.

Figure 5: Analysis of SINV-GFP infection in FHA:DICER N1 cell line.

A, Expression of DICER (DICER, HA), TRBP and AGO2 was analysed by western blot in

HEK 293T, NoDice, FHA:DICER and FHA:DICER N1 cell lines. Alpha-Tubulin was used as

loading control. **B,** Northern blot analysis of miR-16 expression in the same samples as in A.

Expression of snRNA U6 was used as loading control. **C,** Representative GFP fluorescent

microscopy images of HEK 293T, NoDice, FHA:DICER and FHA:DICER N1 cell lines

infected with SINV-GFP at an MOI of 0.02 for 24 h. The left panel corresponds to GFP signal

and the right panel to a merge picture of GFP signal and brightfield. Pictures were taken with a

5x magnification. hpi: hours post-infection. **D,** Western blot analysis of DICER (DICER and

HA) and GFP expression in SINV-GFP-infected cells in the same condition as in C. alpha-

Tubulin was used as loading control. **E**, Mean (+/- SEM) of SINV-GFP viral titers fold change

over HEK 293T cells in HEK 293T, NoDice, FHA:DICER and FHA:DICER N1 cell lines

infected at an MOI of 0.02 for 24 h (n = 3) from plaque assay quantification. $* p < 0.05$, $** p <$

0.01, ordinary one-way ANOVA test.

Supplementary material

Supplementary figures legends

Figure S1: Analysis of SINV-GFP infection in FHA:DICER cell line at different MOI and time points.

HEK 293T, NoDice and FHA:DICER cells were infected with SINV-GFP at an MOI of 2 for 16 h (**A**) or 6 h (**B**). Upper panel: Representative GFP pictures of infected cells. The left panel corresponds to GFP signal and the right panel to a merge of GFP signal and the corresponding brightfield. Pictures were taken with a 5x magnification. hpi: hours post-infection. Middle panel: Western blot analysis of DICER (DICER and HA) and GFP expression in SINV-GFP-infected cells. Alpha-Tubulin was used as loading control. Bottom panel: Mean (+/- SEM) of SINV-GFP viral titers in HEK 293T, NoDice and FHA:DICER cells infected at an MOI of 2 for 16 h (left panel n = 4) or for 6 h (right panel n = 3) from plaque assay quantification. * $p < 0.05$, *** $p < 0.001$, ns non-significant. ordinary one-way ANOVA test.

Figure S2: LC-MS/MS analysis of DICER interactome during SFV infection.

A, Volcano plot for differentially expressed proteins (DEPs) between HA IP and CTL IP in FHA:DICER mock-infected cells. Each protein is marked as a dot; proteins that are significantly up-regulated in HA IP are shown in red, up-regulated proteins in CTL IP are shown in blue, and non-significant proteins are in black. The horizontal line denotes the adjusted p-value threshold (0.05) and the vertical lines the Log2 fold change cutoff (-1 and 1). DICER and its cofactors (TRBP, PACT, AGO2) are highlighted in yellow. **B,** Left panel: Volcano plot for DEPs between SFV (MOI of 2, 6 hpi) and mock fractions of HA IP in FHA:DICER cells. Same colour code and thresholds as in A were applied. Proteins that are discussed in the text are highlighted in yellow and SFV proteins in purple. **C,** Summary of the

1000   differential expression analysis of SFV vs mock fractions from HA IP in FHA:DICER cells.

1001   The analysis has been performed using a generalized linear model of a negative-binomial

1002   distribution and p-values were corrected for multiple testing using the Benjamini-Hochberg

1003   method.

1004

Figure S3: Confirmation of LC-MS/MS analysis by co-IP and BiFC controls.

A, FHA:DICER cells were infected with SINV-GFP at an MOI of 0.02 for 24 h and a HA co-

IP was performed. Eluted proteins were resolved by western blot and IP efficiency was assessed

using an HA antibody. In parallel, co-IPed proteins were visualized using appropriate

antibodies. GFP antibody was used to verify the infection and Ponceau staining serves as

loading control. **B,** Schematic representation of Human DICER proteins used for BiFC positive

and negative controls. The different conserved domains are shown in colored boxes. DUF283:

Domain of Unknown Function; PAZ: PIWI ARGONAUTE ZWILLE domain; dsRBD:

dsRNA-binding domain. hDICER WT is the full-length protein. hDICER N1 is deleted of the

first N-terminal 495 amino acids. **C,** In parallel to BiFC assay, expression of BiFC plasmids

was assessed by western blot. DICER proteins (WT and N1) and PKR were visualized using

antibodies targeting endogenous proteins, whereas TRBP and PACT were detected using GFP

antibody. Antibody targeting the SINV coat protein (CP) was used as infection control. Ponceau

staining was used as loading control. **D,** Positive and negative BiFC controls on fixed

NoDice/ΔPKR cells. After co-transfection, cells were infected with SINV at an MOI of 2 for 6

h and fixed. After fixation, cells were stained with DAPI and observed under confocal

microscope. Merge pictures of BiFC and DAPI signals of SINV-infected cells are shown. A

higher magnification of images showing the interaction represented by a red square is shown

in the bottom left corner. Scale bar: 20 μm and 10 μm.

1024

Figure S4: Interaction analysis between the different versions of DICER and TRBP or

PACT using BiFC assay.

NoDice/ΔPKR cells were co-transfected for 24 h with plasmids expressing the different

versions of DICER proteins fused to the N-terminal part of Venus and either TRBP:Venus$^{C\text{-ter}}$

(**A**) or PACT:Venus$^{C\text{-ter}}$ (**B**). Cells were then infected with SINV at an MOI of 2 for 6 h and

Venus signal was observed under epifluorescence microscope. The left panel corresponds to

Venus signal and the right panel to the corresponding brightfield picture. Pictures were taken

with a 5x magnification. hpi: hours post-infection. Scale bar: 100 μm.

1033

1034 **Supplementary tables**

1035 **Supp. Table 1:** Top 100 proteins that are differentially immunoprecipitated in Mock infected

1036 FHA:DICER cells by the HA and Myc (CTL) antibodies. Related to Figure 2.

1037

1038 **Supp. Table 2:** Top 100 proteins that are differentially immunoprecipitated with the HA

1039 antibody in SINV-infected vs. Mock-infected FHA:DICER cells. Related to Figure 2.

1040

1041 **Supp. Table 3:** Top 100 proteins that are differentially immunoprecipitated in Mock infected

1042 FHA:DICER cells by the HA and Myc (CTL) antibodies, in the SFV infection experiment.

1043 Related to Figure S2.

1044

1045 **Supp. Table 4:** Top 100 proteins that are differentially immunoprecipitated with the HA

1046 antibody in SFV-infected vs. Mock-infected FHA:DICER cells. Related to Figure S2.

1047

1048 **Supp. Table 5:** List of primers used in this study.

Figure 1

Figure 2

A

B

C

RNA binding (GO:0003723)

ATP-dependent helicase activity (GO:0008026)

snoRNA binding (GO:0030515)

RNA helicase activity (GO:0003724)

ATP-dependent RNA helicase activity (GO:0004004)

RNA-dependent ATPase activity (GO:0008186)

double-stranded RNA binding (GO:0003725)

actin-dependent ATPase activity (GO:0030898)

ATP-dependent DNA helicase activity (GO:0004003)

RNA stem-loop binding (GO:0035613)

D

Gene name	Mean spectral count		LogFC	Adjusted p-value
	SINV-GFP IP	MOCK IP		
E2AK2 (PKR)	123.3	0.3	7.979	7.316e-102
DSRAD (ADAR-1)	181.3	28.3	2.563	1.421e-73
DHX9	247.3	50	2.194	4.012e-83
PRKRA (PACT)	63.3	13.7	2.093	1.158e-20

Figure 3

Figure 4

A

B

C

Figure 5

A

B

C

SINV-GFP MOI 0.02 24 hpi

D

E

SINV-GFP titer MOI 0.02 24 hpi

Figure S1

Figure S2

A

B

C

Gene name	Mean spectral count		LogFC	Adjusted p-value
	SFV IP	Mock IP		
E2AK2 (PKR)	221	0.5	8.344	2.611e-119
PRKRA (PACT)	131.5	4.5	4.697	8.415e-57
DSRAD (ADAR-1)	268	22	3.466	9.118e-45
DHX9	265	31	2.955	1.245e-80

Figure S3

A

B

C

D

BiFC+DAPI   BiFC+DAPI

^N-ter^Venus:DICER WT
+
TRBP:Venus^C-ter^

^N-ter^Venus:DICER N1
+
TRBP:Venus^C-ter^

Figure S4