**Using a centrifuge-based method to identify novel root traits that affect**

**root-substrate adhesion in *Arabidopsis thaliana***

Bethany M Eldridge[1,*], Emily R Larson[1,*], Laura Weldon[1,3], Kevin M Smyth[1], Annabelle N

Sellin[1], Isaac V Chenchiah[2], Tanniemola B Liverpool[2] and Claire S Grierson[1,#]

[1]School of Biological Sciences, University of Bristol, Bristol Life Sciences Building, 24

Tyndall Avenue, Bristol, BS8 1TQ, UK

[2]School of Mathematics, University of Bristol, Fry Building, Woodland Road, Bristol BS8

1UG, UK

[3]Wildfowl and Wetlands Trust (WWT), Slimbridge, Gloucester, GL2 7BT, UK

[*]Authors contributed equally to this work.

#Corresponding author

Author for correspondence:

*Claire Grierson*

*Tel:* +44 117 39 41185

*Email:* lacsg@bristol.ac.uk

**ORCID:**

BME: ORCID: 0000-0002-6598-3701

ERL: ORCID: 0000-0002-5498-8152

LW: ORCID: 0000-0002-0761-4414

KMS: ORCID: 0000-0003-0105-8557

IVC: ORCID: 0000-0002-8618-620X

TBL: ORCID: 0000-0003-4376-5604

CSG: ORCID: 0000-0002-4000-0975

**Summary** (162)

- The physical presence of roots and the compounds they release affect the cohesion between roots and their environment. However, the plant traits that are most important for these interactions are unknown and most methods that quantify the contributions of these traits are time-intensive and require specialist equipment and complex substrates.

- We used an inexpensive, high-throughput phenotyping assay to test its sensitivity and versatility for discovering root traits and novel genes important for root-substrate adhesion. This assay measures the centrifugal force required to detach *Arabidopsis thaliana* seedlings from an agar medium.

- Using this method, we identified root hair shapes, vesicle trafficking pathways and root exudate composition that are important for root-substrate adhesion. We used the assay to conduct a genetic screen and discovered a previously uncharacterised ABC transporter. This demonstrates how this assay can be used to identify novel genes that affect root-substrate interactions.

- This assay is a powerful tool for identifying and quantifying genetic contributions to cohesion between roots and their environment.

## Keywords

Arabidopsis, Centrifuge assay, Forward genetics, Root exudates, Root hairs, Root-soil adhesion, Survival analysis, Vesicle trafficking

## Introduction (593)

Plants secrete compounds that help them adapt to their environment, sense and interact with other organisms, and improve water and nutrient uptake (Bailey et al., 2002; Datta et al., 2011; Haling et al., 2013). These compounds, termed exudates, can vary in composition based on plant species, developmental stage, or environment (Naveed et al., 2017). Recent studies have shown that some compounds have bioadhesive properties that stick plants and soils together and aggregate soil particles, modifying the microenvironment around plant roots (Galloway et al., 2018).

Understanding how plant biology and physiology contribute to plant-soil interactions can be confounded by multiple factors: other organisms living in the soil also secrete organic metabolites that can contribute to soil cohesion, soil composition might affect plant exudate

composition and vice versa. Also, plant root morphology including root architecture and root

hairs can alter root-soil interactions (Bailey et al., 2002; Burylo et al.; De Baets et al., 2007;

Ghestem et al., 2014; Stokes et al., 2009). This complexity has made it difficult to

characterise the relevant plant physiology.

Increasing centrifugal force has been used to measure the capacity of different

tropical arboreal ant species to adhere to a smooth Perspex surface when spun at increasing

speed in a centrifuge (Federle et al., 2000). We developed a high-throughput assay that

identifies and quantifies the contribution of root hair cells to root-substrate cohesion in a less

variable environment than the natural rhizosphere (De Baets et al., 2020). The assay uses a

centrifuge to apply force to *Arabidopsis thaliana* seedlings grown on the surface of a sterile

medium and measures the force required to detach seedlings from that surface. The

adherence of candidate lines to the gel at increasing centrifugal force is then compared to a

wild type control line. Applying a Cox proportional hazards regression (Prentice and

Kalbfleisch, 2003) to the results from centrifuging seedlings allows the adherence of

candidate Arabidopsis lines relative to wild-type plants at increasing centrifugal force to be

quantified.

While the centrifuge assay was initially developed as one component of a report on

how the absence or presence of root hairs affects root-substrate adhesion (De Baets et al.,

2020), we wanted to optimize the capability of the assay to evaluate other traits and provide a

highly quantifiable and affordable way to screen large populations of seedlings for the

identification of new genetic factors important for plant-substrate interactions. To assess this

extended use of the assay, we asked (1) if the assay could detect effects of root hair shape on

adhesion; (2) if trafficking functions that affect adhesion could be identified in mutants with

altered vesicle trafficking but no known root hair morphological defects; (3) if effects of

exudate composition could be detected in mutants with altered exudate composition but no

known root hair morphological defects; and (4) if the assay could isolate lines from a mutant

population in a forward genetic screen to identify novel genes that affect root-substrate

adhesion.

Here, we show that this centrifugation method can be used to investigate how root

hair morphology contributes to substrate adhesion and screen for biological components that

do not present a visible phenotype. Using the genetic and molecular tools available in

Arabidopsis and working in a sterile environment allows hypotheses about plant adhesive

factors to be tested. While plant-soil cohesion is a complex and dynamic interaction, our

assay provides a way to probe root cellular functions that might be masked by confounding

variables in a soil-based study system. Results from this assay provide a platform for more
comprehensive studies of plant morphology and physiology that promote root-substrate
cohesion.

**Materials and Methods** (1380)

**Plant material and growth conditions**

Candidate lines of the Columbia-0 (Col-0) ecotype of *Arabidopsis thaliana* single-
gene mutants were obtained from the Nottingham Arabidopsis Stock Centre (NASC;
Nottingham, UK) or were stocks maintained in the lab (Table 1). Wild-type plants for all
experiments were also the Col-0 ecotype. Homozygous lines of *csld3, cow1-3, lrx1-4, rol1-2,*
*gdi1-2, chc1-2, chc2-3, syp121, syp122, jin1-9, pft1-3* and *pdr2* were used in all experiments.
Homozygosity of the *abcg43* mutant alleles was confirmed by genomic PCR analysis.

For all experiments, seeds were surface sterilised for 15 min in a solution containing
20% bleach and sterile water, followed by five washes in sterile water. All sterilised seeds
were stratified in the dark at 4°C for 24-48 h in Eppendorf microtubes containing sterile
water. All plants were reared in a growth room at 21-22°C with a continuous photoperiod
(light intensity = 120-145 µmol m$^{-2}$ s$^{-1}$) at 60% humidity.

Sterilized seeds were sown onto 90-mm Petri plates (Thermo Scientific RC2260).
containing 30 ml sterile gel medium and sealed with Parafilm (Bemis, NA). The gel consisted
of half strength Murashige and Skoog basal medium (Sigma M5519) with 1% [w/v] sucrose
pH adjusted to 5.7, solidified with 1% [w/v] agar (Sigma A1296). Seed-sown plates were
placed upright at ~80° from the horizontal (lid side down) to encourage vertical growth along
the gel surface. Plants for genomic DNA extractions were grown in compost (three parts
Levington F3 compost and one-part J Arthur Bowers horticultural silver sand).

**Centrifuge assay set-up**

To measure the adhesion of Arabidopsis seedling roots using a centrifuge, seedling
roots must not penetrate the gel or each other, as both events could affect root-gel adhesion
by interfering with seedling detachment during centrifugation. Therefore, only ten seeds were
sown 1 cm apart on a 90 mm Petri plate in two parallel rows**.** The plates were then orientated
vertically at approximately 80° from horizontal and grown for 5-6 days in a growth chamber
(Fig. S1a). Individual plates and seedlings were visually evaluated to ensure that plants were
at the same developmental stage and of similar size. Any seedlings that were touching each
other, had grown into the gel, or not uniform in size were omitted from the assay. The

seedlings were then numbered so that plate positional effects could be included as a potential

covariate in the final analysis (Fig. S1b). Plates were inverted and placed into a swing-out

bucket centrifuge (Beckman Allegra X-30R Centrifuge) that held four plates at a time and

subjected to increments of increasing centrifugal force at 720, 1018, 1247, 1440 and 1611

RPM for one minute each (Fig. S1b). The proportion of seedlings that detached from the gel

surface was recorded between each centrifugal speed; a detachment event was recorded when

a seedling had partially or fully peeled away from the gel (Fig. S1b). If the gel shattered

during the assay, seedlings that had not detached from the gel were not recorded and

seedlings that remained adhered to the gel after the maximum centrifugal speed were

censored in our analyses.

**Calculating seedling detachment force**

To calculate the centrifugal force acting on a seedling, the aerial tissue mass for each

seedling was measured using an analytical balance (Microbalance ME5; Sartorius; Fig. S1c).

The aerial tissue weight of each seedling was measured because a plant with heavier aerial

tissue experiences more force than a lighter one subjected to the same centrifugal speed.

Centrifugal force, Fc (mN) acting on a seedling was calculated as previously reported in De

Baets et al. (2020). Aerial tissue weight (kg), the angular velocity ($\omega$) and the distance

between the seedling and the axis of rotation on the centrifuge (radius = 0.07m) were used to

give the following equation:

Fc= *mass* x *radius* x $\omega^2$                (Equation 1)

**Identification of *abcg43-1* mutant T-DNA insert**

Genomic DNA was extracted from a pool of vegetative tissue taken from ten three-

week old plants using an adapted protocol (Healey et al., 2014) for Illumina Next-Generation

two x paired-end sequencing conducted at the Bristol Genomics Facility. The Illumina

TruSeq Nano LT gDNA kit (Illumina Inc) was used to generate a genomic DNA sequencing

library following the manufacturer's protocol. The final library was diluted to a loading

concentration of 1.4 pM for cluster generation and 2 x 150 bp paired-end sequencing on the

Illumina NextSeq500 system (Illumina Inc) alongside a 5% PhiX spike-in control library.

Read summary statistics were generated using the RTA 2.4.6 Primary Analysis Software

(Illumina Inc). The read summaries were analysed in the Sequencing Analysis Viewer

(Illumina Inc). Filtered paired reads were subject to paired-end alignment using the Bowtie2

2.3.4.2 aligner (Langmead and Salzberg, 2012). A bespoke reference genome was produced

that combined the TAIR10 Arabidopsis genome (Lamesch et al., 2012) and the pROK2

vector sequence (Baulcombe et al., 1986). Alignments were viewed in the Integrative

Genomics Viewer IGV 2.3 (Robinson et al., 2011).

For Sanger sequencing, DNA was extracted from a pool of vegetative tissue taken

from two-week-old plants using a modified Edwards prep (Edwards et al., 1991). High

fidelity PCR was conducted on the *abcg43-1* line using the Q5 High-Fidelity 2X Master Mix

(NEB). PCR products were purified and extracted from a 1% agarose gel using the QIAquick

Gel Extraction Kit (QIAGEN), following the manufacturer's instructions. Purified PCR

product plus the *abcg43-1* forward, reverse or border genotyping primer (Table S1) was used

for Sanger sequencing (using the Mix2Seq overnight sequencing kit, Eurofins Genomics).

Chromatograms and FASTA files were obtained from Eurofin Genomics and following

manual low-quality end trimming, the final sequences were aligned to the pROK2 vector

sequence and *ABCG43* gene sequence using the MUSCLE alignment tool (Edgar, 2004).

**Genotyping**

184          To genotype the *abcg43* mutant alleles, genomic DNA was extracted from the

185    vegetative tissue of two-week-old plants using a modified Edwards prep (Edwards et al.,

186    1991). T-DNA border and gene-specific primers were used in PCR analyses to confirm the

187    genotype (Table S1).

**Morphological root trait analyses**

Five-day-old Arabidopsis seedlings were imaged with a Leica MZ FLIII microscope

(Leica) with dark-field lighting. Root hair length and density were measured using

microscope images of five-day-old Arabidopsis seedlings and Fiji version 1.0 (Schindelin et

al., 2012), using the Bio-Formats Importer plugin to load images into Fiji. At least two

experimental repeats were conducted; in each experiment we imaged eight to ten individual

plants from each line and measured at least 30 root hairs per plant.

**Statistical analyses**

All statistical analyses were conducted using RStudio, version 1.1453 (R Core Team,

2014) and all graphs were generated using the R package ggplot2 (Wickham, 2016).

A Cox proportional hazards (PH) regression model was used to statistically test for

differences between the detachment of experimental seedlings relative to the wild-type

control (Col-0) at increasing force. Cox PH regression models were conducted using the
coxph function within the survival package in R and in all cases, the assumption of
proportionality was satisfied. Comparisons of different candidate lines relative to the wild
type line were analysed *a priori*; therefore, a series of contrasts were set up using the R
function contr.treatment rather than using post-hoc testing methods. Seedling position and
individual plate number were incorporated as covariates to account for any heterogeneity;
when these covariates had no significant effect, they were removed from the model. We
censored seedlings that remained attached to the gel after the maximum centrifugal speed
because we did not determine the speed these seedlings would have detached from the gel.
For each Cox PH regression model run, we report the P value of the Wald Statistic ($z$-score)
and the hazard ratio with the upper and lower bound confidence intervals. An alpha level of
0.01 was used. The root-gel detachment of at least 70 plants from each line was assessed
within a single experiment.

Univariate analyses were conducted using the lm function in R to test for differences
in the mean root hair density or mean root hair length (of eight to ten individual plants from
each line) between candidate lines relative to wild type. In all cases, the assumptions of
normality and homoscedasticity were satisfied. Because comparisons of different mutant
lines relative to the wild type line were *a priori*, a series of contrasts were conducted using
the contr.treatment function in R. To prevent type I errors from multiple testing, we applied
the Bonferroni method and adjusted the alpha level to 0.025 (0.05/2).

**Results** (1795)

**Using the Cox survival regression model to quantify root-substrate adhesion**

To compare the root-gel adhesion properties of candidate lines relative to wild type
we conducted a semi-parametric Cox proportional hazards (PH) regression. We chose this
over other types of survival analyses to statistically evaluate the differences in detachment of
Arabidopsis seedlings at increasing force because Cox PH regression models are multivariate
and can test whether covariates affect the occurrence of an event (i.e. gel detachment) as well
as manage censored data (Bradburn et al., 2003; Cox, 1972; Devarajan and Ebrahimi, 2009).
Unlike univariate tests commonly used in survival analysis such as the Logrank test, Cox PH
regression models also provide an estimate of the difference in effect size between two
groups by calculating a hazard ratio, which is an exponential coefficient that gives the
instantaneous risk of an event (Cox and Oakes, 1984; Spruance et al., 2004). This is

advantageous because we used hazard ratios as a strength measure of root-gel adhesion by

comparing the seedling detachment risk of candidate lines relative to wild type.

We report the P-value calculated from the Wald statistic ($z$-score) and the hazard ratio

with upper and lower bound confidence intervals. Wild type was used as the baseline group

for all models, with a hazard ratio of one. Therefore, a candidate line with a higher risk of

detachment from the gel than wild type will have a hazard ratio above one, while lines with a

lower risk of detachment than wild type will have a hazard ratio below one. Because roots are

developmentally plastic and sensitive to environmental heterogeneity (Bliss et al., 2002; Gao

et al., 2004; Šmilauerová, 2001), we assessed the root-gel detachment of at least 70 seedlings

from each line in a single experiment.

**Root hair morphological effects on root-substrate adhesion**

We screened mutant lines with known root hair morphology defects to ask if the

centrifuge assay could determine if root hair shape affects root-substrate adhesion. The

mostly bald mutant *csld3*, which encodes the CESA-like 3D protein (Favery et al., 2001;

Yang et al., 2020) was included as a root hairless example. For lines that produce short or

malformed root hairs, we included the *gdi1-2* Rho GTPase GDP dissociation inhibitor

mutant, *rol1-2* rhamnose biosynthesis 1 mutant allele, *can of worms 1* (*cow1-3*)

phosphatidylinositol transferase protein mutant, and *lrx1-4* leucine rich extensin 1 mutant

(Baumberger et al., 2001; Böhme et al., 2004; Carol et al., 2005; Diet et al., 2006; Grierson et

al., 1997). The *rol1-2* and *gdi1-2* mutants have short root hairs and the root hairs on *gdi1-2*

plants can also branch (Parker et al., 2000; Ringli et al., 2008). *cow1-3* plants have short,

wide root hairs that can branch, whilst *lrx1-4* plants can develop root hairs that abort, swell,

branch, or collapse in a growth condition-dependent manner (Baumberger et al., 2001;

Grierson et al., 1997; Parker et al., 2000). Under our growth conditions, we confirmed the

reported phenotypes for each mutant line (Fig. 1a).

Consistent with previous results (De Baets et al., 2020), bald (*csld3*) seedlings

detached at lower centrifugal forces relative to wild-type seedlings, with 4.6 times the risk of

detachment compared to wild-type plants ($z$ = 8.644, P<0.001, HR = 4.636, 95% CI = 3.274 –

6.564; Fig. 2b). Mutants with deformed root hair phenotypes including *gdi1-2, rol1-2, cow1-*

*3* and *lrx1-4* had risks of detachment that were 2.5, 4, 3.8 and 3.9 times that of wild-type

plants, respectively (*gdi1-2* – $z$ = 5.199, P<0.001, HR = 2.459, 95% CI = 1.752 – 3.452; *rol1-*

*2* – $z$ = 7.927, P<0.001, HR = 4.036, 95% CI = 2.858 – 5.698; *cow1-3* – $z$ = 7.350, P<0.001,

HR = 3.790, 95% CI = 2.657 – 5.407; *lrx1-4* – $z$ = 8.049, P<0.001, HR = 3.926, 95% CI =

2.814 – 5.478; Fig. 2b-d). These results indicate that the centrifuge assay can quantify effects

of root hair shape on root-substrate cohesion. Many of the root hair mutants tested are

associated with changes in root hair cell wall composition, suggesting that alterations in

secreted compounds could affect root-substrate interactions in addition to any physical or

surface area-related factors that are specific to root hairs.

**Vesicle trafficking mutations affect root-substrate adhesion**

We hypothesized that Arabidopsis mutants with defective trafficking pathways

might alter cell wall and apoplast characteristics that contribute to root-substrate interactions

and adhesion. We chose the secretory Soluble NSF (N-ethylmaleimide sensitive fusion

protein) Attachment proteins (SNAP) REceptor (SNARE) mutants *syp121* and *syp122-1*, and

the endocytic mutants, *chc1-2* and *chc2-3* as candidates for our assay. SYP121 is the primary

secretory SNARE found on the plasma membrane (Assaad et al., 2004; Geelen et al., 2002)

and has characterized aboveground phenotypes, including small rosettes and stomatal

mobility defects (Eisenach et al., 2012; Larson et al., 2017); however, the mutant does not

have a reported root hair phenotype. Although the related SNARE SYP122 is thought to

share partial functionality with SYP121, the *syp122-1* mutant does not share these

phenotypes with *syp121* and a recent proteomic analysis reported differences in their vesicle

cargoes, which could contribute to their functional independence (Waghmare et al., 2018).

The *chc1-2* and *chc2-3* mutants are defective in the heavy chain subunits of the clathrin coat

complex, which is required for vesicle traffic at the plasma membrane (Kitakura et al., 2011).

Endo- and exocytic rates are impaired in both mutants (Larson et al., 2017), but no root hair

phenotypes have been reported. For the centrifuge assay, it was important to select mutants

that do not have root hair phenotypes given the effect root hair morphology has on root-gel

adhesion. Therefore, we evaluated the root hairs of these trafficking mutants and found that

the root hairs of these lines did not significantly differ from wild-type plants (Fig. 2a).

Overall, the risk of detachment for *syp121, syp122-1, chc1-2* and *chc2-3* plants was

3.6, 3.3, 4.5 and 3.9 times that of wild type, respectively (*syp121* – $z$ = 7.702, P<0.001, HR =

3.593, 95% CI = 2.595 – 4975; *syp122-1* – $z$ = 7.199, P<0.001, HR = 3.270, 95% CI = 2.369

– 4.515; *chc1-2* – $z$ = 8.238, P<0.001, HR = 4.482, 95% CI = 3.137 – 6.405; *chc2-3* – $z$ =

7.653, P<0.001, HR = 3.941, 95% CI = 2.774 – 5.599; Fig. 2b, c). Given that all the mutants

detached from the gel at lower centrifugal forces than wild-type seedlings, these results

indicate that the corresponding genes contribute to root-substrate adhesion and that the

centrifuge assay can be used to probe for intracellular molecular machinery that affect root-
substrate interactions.

**Exudate composition changes root detachment rates**

Over 20% of plant assimilated carbon is released as root exudates (Huang et al., 2016)
that modify the external environment in response to abiotic and biotic stimuli and contribute
to soil adhesion (Akhtar et al., 2018; Galloway et al., 2018). Root epidermal and hair cells
secrete compounds that contribute to root exudate profiles that can be plant species specific
(Naveed et al., 2017). Because mutants in both root hair development and vesicle trafficking
pathways showed root-substrate adhesion phenotypes (Figs. 1 and 2), we asked if the
centrifuge assay could identify effects of exudate composition on root-substrate adhesion. We
selected Arabidopsis mutants reported to have altered exudate composition compared to wild
type, including *pft1-3*, *jin1-9*, and *pdr2* (Badri et al., 2009; Berger et al., 1996; Carvalhais et
al., 2015; Kidd et al., 2009). *PFT1* (*MED25*) encodes the MEDIATOR25 subunit of the
Mediator nuclear protein, and *JIN1* (*MYC2*) encodes a basic helix-loop-helix leucine zipper
transcription factor; both proteins are involved in the jasmonate signalling pathway (Kidd et
al., 2009; Lorenzo et al., 2004). *pft1-3* and *jin1-9* mutants are reported to have altered root
exudate composition, including lower amounts of the amino acids asparagine, ornithine and
tryptophan than wild-type plants (Carvalhais et al., 2015). *PDR2* (*ABCG30*) encodes a
pleiotropic drug resistance (PDR) full-length ABC transporter that is involved in ABA
transport and the exudation of secondary metabolites (Badri et al., 2008; Kang et al., 2015).
We did not observe root hair growth phenotypes in these mutant lines compared to wild type
(Fig. 3a).

We found that compared to wild-type seedlings, *pdr2* seedlings resisted detachment,
with 0.38 times the risk of detaching from the gel ($z$ = -5.765, P<0.001, HR = 0.379, 95% CI
= 0.273 – 0.528; Fig. 3b). Conversely, *pft1-3* seedlings had an increased risk of detachment
1.7 times that of wild-type plants ($z$ = 3.315, P<0.001, HR = 1.698, 95% CI = 1.242 – 2.323),
and there was no difference in gel adhesion between *jin1-9* and wild-type plants ($z$ = 0.737,
P>0.05, HR = 1.124, 95% CI = 0.824 – 1.537; Fig. 3b). These results show that exudate
composition can alter root-substrate interactions, although this may not be the case for all
exudate compositional changes, or identification of the effects of the exudate composition of
the *jin1-9* mutant lay outside the sensitivity range of the assay. Since these mutant lines did
not have root hair phenotypes, our results suggest that changes in exudate composition alter

root adhesive properties without affecting plant root hair morphology, further indicating the

power of the centrifuge assay as a potential mutant screening method.

**Centrifuge assay as a method for forward genetic screening**

Having demonstrated that our centrifugation technique can be used to identify

different types of traits that affect root-substrate adhesion, we conducted a forward genetic

screen to identify novel genes involved in root-substrate adhesion. To conduct this screen, the

root-gel adhesion properties of individual plants from a pooled SALK T-DNA insertion

mutant collection were analysed (Alonso et al., 2003). Individual plants with significantly

increased or decreased root-gel adhesion relative to wild-type plants were recovered from the

screen and self-fertilized to obtain progeny and identify the mutations. We identified a line

with significantly enhanced root-gel adhesion and conducted genomic Next-Generation

sequencing followed by Sanger sequencing of T-DNA flanking PCR products to confirm a T-

DNA insertion in the ABC transporter gene, *ABCG43,* whose function is unknown. We

named this T-DNA insertion line *abcg43-*1 and identified additional mutant alleles, *abcg43-2*

(SALK_201207c) and *abcg43-3* (SALKseq_30713) (Fig. 4a).

We analysed the root-gel adhesion of homozygous insertional mutant alleles for

*ABCG43* and found *abcg43-1, abcg43-*2 *and abcg43-*3 plants resisted detachment from the

gel, with detachment risks 0.25, 0.35 and 0.2 times that of wild-type plants, respectively

(*abcg43-1 – z = -*7.823, P<0.001, HR = 0.245, 95% CI = 0.172 – 0.348; *abcg43-2 – z = -*

6.049, P<0.001, HR = 0.355, 95% CI = 0.253 – 0.496; *abcg43-3 – z =* -7.179, P<0.001, HR =

0.203 – 0.403; Fig. 4b, c). Collectively, these results suggest that disruption of *ABCG43*

expression and protein function affects root-gel adhesion. These results also illustrate the use

of the centrifuge assay as a powerful screening method for identifying novel genes involved

in root-substrate adhesion, as well as a tool for evaluating the effects of known gene function

on root-substrate interactions. Experiments are now being conducted to identify mutant

trait(s) in these lines and understand how *ABCG43* affects root-substrate adhesion.

**Discussion** (837)

To date, most published methods that quantify the contribution of particular traits to

plant-soil interactions are time-intensive and expensive (Bailey et al., 2002; De Baets and

Poesen, 2010; De Baets et al., 2020; Lin and Ying, 1994; Toukura et al., 2006). In contrast,

our assay can produce data within a week, is available to any laboratory with access to a
bench top centrifuge and does not require speciality consumables. This method is a high-
throughput and quantitative way to test the effects of root morphology and cell function on
root-substrate adhesion and to screen for new biological and molecular factors that can alter
plant-substrate interactions. Using a defined model system like Arabidopsis and sterile
medium conditions permits plant-specific characteristics to be identified that can then be
probed in other substrate- or soil-based conditions.

Survival analyses are not often utilised by plant biologists because measurements are
static or endpoints, like root length, flowering time, or fresh/dry weight and therefore, do not
require a dynamic analysis of individual datapoints over a continuous spectrum of time or
force. However, survival analysis is used across other scientific fields to study the timing of
an event, such as the time for an animal to learn a behaviour, for an individual to recover or
die from a disease, or an electrical failure to occur (Allison, 2010; Lin and Ying, 1994;
Miller, 1998). Here, we use survival analysis to study root-substrate interactions, where
seedling detachment is the event that occurs over increasing forces.

**Using the centrifuge assay to investigate the role of root hairs in root-substrate**
**interactions**

Results presented in this paper show that the centrifuge assay can quantify effects of
root hair morphology on plant-substrate adhesion. By using well-characterized Arabidopsis
root hair mutants, we show this assay can evaluate the adhesive properties of mutants with a
wide variety of root hair phenotypes. Previous work has shown that the presence of root hairs
can significantly enhance root-substrate adhesion compared to the absence of root hairs (Choi
and Cho, 2019; De Baets et al., 2020; Haling et al., 2013). Indeed, our centrifuge assay has
previously been used to quantify the forces required to detach seedlings with and without root
hairs (De Baets et al., 2020); however, the sensitivity of the assay for comparing how root
hair morphology and function affects root-substrate adhesion was not evaluated. We now
show that this assay can distinguish the adherence strength of seedlings with altered root hair
shapes and sizes (Fig. 1) and quantify the contribution these morphological traits make to
plant-substrate interactions. Some mutants with defective root hairs have mutations in cell
wall biosynthesis and modification proteins, so it is possible that altered root-substrate
interfaces compromise adhesion in these mutants, not just root hair shape. Using our method
to rapidly phenotype mutants with root hair growth defects could direct additional
experiments to characterize biochemical and morphological properties of genes of interest.

Indeed, current evidence for the importance of root hairs for the deposition of complex
polysaccharides with soil-binding properties (Galloway et al., 2020) supports the need for
assays such as this one to expedite our ability to identify genes and mechanisms that could
improve soil-remediation, top soil maintenance, and erosion tolerance of at-risk land.

**Identifying new genes involved in root-substrate adhesion**

Mutations in cellular functions that do not alter root hair growth but do affect
substrate adhesion can also be investigated using this assay. Vesicle trafficking pathways are
required for cell wall deposition and maintenance (De Caroli et al., 2011; Larson et al., 2014;
Rodriguez-Furlán et al., 2016); therefore, we hypothesized that mutants in trafficking
pathways, particularly those at the plasma membrane, could affect root-substrate adhesion.
Similarly, the composition and deposition of plant exudates help plants modify and optimize
their growth conditions (Carvalhais et al., 2015; Naveed et al., 2017), suggesting that
modification to plant exudate composition could affect root-substrate interactions. Consistent
with these hypotheses, mutants with vesicle trafficking defects and altered exudate
composition have root-substrate adhesion properties that differ from wild type, indicating that
this method can identify aspects of plant biology important for root-substrate interactions that
do not affect root or root hair morphology (Figs. 2 and 3). Our results further indicate that
while root hairs provide significant adhesive and cohesive support both on solid gel and soil
media (De Baets et al., 2020), there are additional cellular factors contributing to plant-
substrate interactions that are not directly observable without the use of expensive antibodies,
development of time-intensive transgenic lines, and microscopy equipment. Our centrifuge
assay provides an affordable and quick alternative to these current methodologies.

We used this assay as a screen to identify new genes affecting plant-substrate
interactions. As an example, we report the identification in a forward genetics screen of a
mutant population in which *abcg43* mutant seedlings were more adhesive to gel than wild
type (Fig. 4). The function of ABCG43 in Arabidopsis is uncharacterised and therefore, this
assay is a promising tool for identifying novel candidate genes that have no previously
known association with root morphological defects, adhesion or exudate composition for
future study.

**Conclusion (115)**

This assay has applications in plant cell biology and genetics to understand how gene
activity, cell function and root structure affect plant-soil adhesion and identify the molecular

pathways involved. Potential applications include measuring variables not tested in this
report, such as the effects of environmental conditions on plant-substrate adhesion and how
other plant species or ecotypes respond to detachment forces. Given the recent discovery that
root hair presence is important for Arabidopsis cohesion between roots and gel, and compost
or soil (De Baets *et al.,* 2020), this assay could contribute to plant and soil sciences more
broadly through the identification of plant traits that increase root-substrate cohesion and
potentially stabilise slopes or reduce soil erosion.
442

**Supporting Information**

443
444 **Fig. S1** Centrifuge assay workflow for determining the root-gel adhesion properties of
445 *Arabidopsis thaliana* seedlings
446 **Table S1** Genotyping primers used in this study for genotyping *Arabidopsis thaliana* mutants
447 **Table S2** Centrifuge assay troubleshooting
448

**Acknowledgements (88)**

449
This work was supported by a Leverhulme Trust project grant RPG-2013-260 to CSG and
TBL and BME was supported by a BBSRC SWBio PhD studentship (grant BB/M009122/1).
We are thankful to Don Grierson FRS and Enrico Coen FRS who independently suggested
the centrifuge assay concept, Timothy Quine for providing insightful guidance during method
development, and Thomas Denbigh for providing essential technical advice. We are grateful
to Jill Harrison, Victoria Spencer, Zoe Nemec Venza, Sophie Carpenter and Nicholas H Fair
for their helpful comments during revision of the manuscript.
457

**Author Contributions (105)**

458
CSG designed the initial project, conducted proof-of-concept pilot experiments, obtained the
funding, and recruited and advise the team. TBL and IC developed the theoretical
parametrisation of the centrifugal assay and bounds on the errors from other mechanical
forces. BME and ERL planned and conducted all experiments apart from the initial genetic
screen, which was carried out by LW and KMS. BME and ERL analysed all of the data
presented herein. AS and BME optimised the centrifuge assay data collection and data
analysis methods. BME and ERL wrote the initial manuscript. BME, ERL, and CSG edited
the manuscript. All authors reviewed and approved the final manuscript.
467

## References

Akhtar, J., Galloway, A.F., Nikolopoulos, G., Field, K.J., and Knox, P. (2018). A quantitative method for the high throughput screening for the soil adhesion properties of plant and microbial polysaccharides and exudates. Plant Soil *428*, 57–65.

Allison, P.D. (2010). Survival Analysis Using SAS: A Practical Guide, Second Edition (SAS Institute).

Alonso, J.M., Stepanova, A.N., Leisse, T.J., Kim, C.J., Chen, H., Shinn, P., Stevenson, D.K., Zimmerman, J., Barajas, P., Cheuk, R., et al. (2003). Genome-wide insertional mutagenesis of Arabidopsis thaliana. Science *301*, 653–657.

Assaad, F.F., Qiu, J.-L., Youngs, H., Ehrhardt, D., Zimmerli, L., Kalde, M., Wanner, G., Peck, S.C., Edwards, H., Ramonell, K., et al. (2004). The PEN1 Syntaxin Defines a Novel Cellular Compartment upon Fungal Attack and Is Required for the Timely Assembly of Papillae. Mol. Biol. Cell *15*, 5118–5129.

Badri, D.V., Loyola-Vargas, V.M., Broeckling, C.D., De-la-Peña, C., Jasinski, M., Santelia, D., Martinoia, E., Sumner, L.W., Banta, L.M., Stermitz, F., et al. (2008). Altered Profile of Secondary Metabolites in the Root Exudates of Arabidopsis ATP-Binding Cassette Transporter Mutants. Plant Physiology *146*, 762–771.

Badri, D.V., Quintana, N., El Kassis, E.G., Kim, H.K., Choi, Y.H., Sugiyama, A., Verpoorte, R., Martinoia, E., Manter, D.K., and Vivanco, J.M. (2009). An ABC transporter mutation alters root exudation of phytochemicals that provoke an overhaul of natural soil microbiota. Plant Physiol. *151*, 2006–2017.

Bailey, P.H.J., Currey, J.D., and Fitter, A.H. (2002). The role of root system architecture and root hairs in promoting anchorage against uprooting forces in Allium cepa and root mutants of Arabidopsis thaliana. J. Exp. Bot. *53*, 333–340.

Baulcombe, D.C., Saunders, G.R., Bevan, M.W., Mayo, M.A., and Harrison, B.D. (1986). Expression of biologically active viral satellite RNA from the nuclear genome of transformed plants. Nature *321*, 446–449.

Baumberger, N., Ringli, C., and Keller, B. (2001). The chimeric leucine-rich repeat/extensin cell wall protein LRX1 is required for root hair morphogenesis in Arabidopsis thaliana. Genes Dev. *15*, 1128–1139.

Berger, S., Bell, E., and Mullet, J.E. (1996). Two Methyl Jasmonate-Insensitive Mutants Show Altered Expression of AtVsp in Response to Methyl Jasmonate and Wounding. Plant Physiology *111*, 525–531.

Bliss, K.M., Jones, R.H., Mitchell, R.J., and Mou, P.P. (2002). Are competitive interactions influenced by spatial nutrient heterogeneity and root foraging behavior? New Phytologist *154*, 409–417.

Böhme, K., Li, Y., Charlot, F., Grierson, C., Marrocco, K., Okada, K., Laloue, M., and Nogué, F. (2004). The Arabidopsis COW1 gene encodes a phosphatidylinositol transfer protein essential for root hair tip growth. Plant J. *40*, 686–698.

Bradburn, M.J., Clark, T.G., Love, S.B., and Altman, D.G. (2003). Survival analysis part II:
multivariate data analysis--an introduction to concepts and methods. Br. J. Cancer *89*, 431–
436.

Burylo, M., Rey, F., Mathys, N., and Dutoit, T. Plant root traits affecting the resistance of
soils to concentrated flow erosion. Earth Surface Processes and Landforms *37*, 1463–1470.

Carol, R.J., Takeda, S., Linstead, P., Durrant, M.C., Kakesova, H., Derbyshire, P., Drea, S.,
Zarsky, V., and Dolan, L. (2005). A RhoGDP dissociation inhibitor spatially regulates
growth in root hair cells. Nature *438*, 1013–1016.

Carvalhais, L.C., Dennis, P.G., Badri, D.V., Kidd, B.N., Vivanco, J.M., and Schenk, P.M.
(2015). Linking Jasmonic Acid Signaling, Root Exudates, and Rhizosphere Microbiomes.
Mol. Plant Microbe Interact. *28*, 1049–1058.

Choi, H.-S., and Cho, H.-T. (2019). Root hairs enhance Arabidopsis seedling survival upon
soil disruption. Scientific Reports *9*, 11181.

Cox, D.R. (1972). Regression Models and Life-Tables. Journal of the Royal Statistical
Society: Series B (Methodological) *34*, 187–202.

Cox, D.R., and Oakes, D. (1984). Analysis of Survival Data (Boca Raton, Florida: Chapman
& Hall/CRC Press).

Datta, S., kim, C.M., Pernas, M., Pires, N.D., Proust, H., Tam, T., Vijayakumar, P., and
Dolan, L. (2011). Root hairs: development, growth and evolution at the plant-soil interface.
Plant and Soil *346*, 1–14.

De Baets, S., and Poesen, J. (2010). Empirical models for predicting the erosion-reducing
effects of plant roots during concentrated flow erosion. Geomorphology *118*, 425–432.

De Baets, S., Poesen, J., Knapen, A., and Galindo, P. (2007). Impact of root architecture on
the erosion-reducing potential of roots during concentrated flow. Earth Surface Processes and
Landforms *32*, 1323–1345.

De Baets, S., Denbigh, T.D., Smyth, K.M., Eldridge, B.M., Weldon, L.M., Higgins, B.W.,
Matyjaszkiewicz, A.W., Meersmans, J., Larson, E.R., Chenchiah, I.V., et al. (2020). Micro-
scale interactions between Arabidopsis root hairs and soil particles influence soil erosion.
Nature Comms Biol *3*, 1–11.

De Caroli, M., Lenucci, M.S., Di Sansebastiano, G.-P., Dalessandro, G., De Lorenzo, G., and
Piro, G. (2011). Dynamic protein trafficking to the cell wall. Plant Signal Behav *6*, 1012–
1015.

Devarajan, K., and Ebrahimi, N. (2009). Testing for Covariate Effect in the Cox Proportional
Hazards Regression Model. Commun Stat Theory Methods *38*, 2333–2347.

Diet, A., Link, B., Seifert, G.J., Schellenberg, B., Wagner, U., Pauly, M., Reiter, W.-D., and
Ringli, C. (2006). The Arabidopsis root hair cell wall formation mutant lrx1 is suppressed by
mutations in the RHM1 gene encoding a UDP-L-rhamnose synthase. Plant Cell *18*, 1630–
1641.

Edwards, K., Johnstone, C., and Thompson, C. (1991). A simple and rapid method for the
preparation of plant genomic DNA for PCR analysis. Nucleic Acids Res *19*, 1349.

Eisenach, C., Chen, Z.-H., Grefen, C., and Blatt, M.R. (2012). The trafficking protein
SYP121 of Arabidopsis connects programmed stomatal closure and K$^+$ channel activity with
vegetative growth. Plant J. *69*, 241–251.

Favery, B., Ryan, E., Foreman, J., Linstead, P., Boudonck, K., Steer, M., Shaw, P., and
Dolan, L. (2001). KOJAK encodes a cellulose synthase-like protein required for root hair cell
morphogenesis in Arabidopsis. Genes Dev. *15*, 79–89.

Federle, W., Rohrseitz, K., and Holldobler, B. (2000). Attachment forces of ants measured
with a centrifuge: better 'wax-runners' have a poorer attachment to a smooth surface. Journal
of Experimental Biology *203*, 505–512.

Galloway, A.F., Pedersen, M.J., Merry, B., Marcus, S.E., Blacker, J., Benning, L.G., Field,
K.J., and Knox, J.P. (2018). Xyloglucan is released by plants and promotes soil particle
aggregation. New Phytol. *217*, 1128–1136.

Galloway, A.F., Akhtar, J., Marcus, S.E., Fletcher, N., Field, K., and Knox, P. (2020). Cereal
root exudates contain highly structurally complex polysaccharides with soil-binding
properties. The Plant Journal.

Gao, D., Knight, M.R., Trewavas, A.J., Sattelmacher, B., and Plieth, C. (2004). Self-reporting
Arabidopsis expressing pH and [Ca2+] indicators unveil ion dynamics in the cytoplasm and
in the apoplast under abiotic stress. Plant Physiol. *134*, 898–908.

Geelen, D., Leyman, B., Batoko, H., Di Sansebastiano, G.-P., Moore, I., Blatt, M.R., and Di
Sansabastiano, G.-P. (2002). The abscisic acid-related SNARE homolog NtSyr1 contributes
to secretion and growth: evidence from competition with its cytosolic domain. Plant Cell *14*,
387–406.

Ghestem, M., Cao, K., Ma, W., Rowe, N., Leclerc, R., Gadenne, C., and Stokes, A. (2014). A
framework for identifying plant species to be used as "ecological engineers" for fixing soil on
unstable slopes. PLoS ONE *9*, e95876.

Grierson, C.S., Roberts, K., Feldmann, K.A., and Dolan, L. (1997). The COW1 locus of
arabidopsis acts after RHD2, and in parallel with RHD3 and TIP1, to determine the shape,
rate of elongation, and number of root hairs produced from each site of hair formation. Plant
Physiol *115*, 981–990.

Haling, R.E., Brown, L.K., Bengough, A.G., Young, I.M., Hallett, P.D., White, P.J., and
George, T.S. (2013). Root hairs improve root penetration, root-soil contact, and phosphorus
acquisition in soils of different strength. J. Exp. Bot. *64*, 3711–3721.

Healey, A., Furtado, A., Cooper, T., and Henry, R.J. (2014). Protocol: a simple method for
extracting next-generation sequencing quality genomic DNA from recalcitrant plant species.
Plant Methods *10*, 21.

Huang, Y., Wang, Y., Tan, L., Sun, L., Petrosino, J., Cui, M.-Z., Hao, F., and Zhang, M.
(2016). Nanospherical arabinogalactan proteins are a key component of the high-strength
adhesive secreted by English ivy. Proc. Natl. Acad. Sci. U.S.A. *113*, E3193-3202.

Kang, J., Yim, S., Choi, H., Kim, A., Lee, K.P., Lopez-Molina, L., Martinoia, E., and Lee, Y.
(2015). Abscisic acid transporters cooperate to control seed germination. Nat Commun *6*,
8113.

Kidd, B.N., Edgar, C.I., Kumar, K.K., Aitken, E.A., Schenk, P.M., Manners, J.M., and
Kazan, K. (2009). The mediator complex subunit PFT1 is a key regulator of jasmonate-
dependent defense in Arabidopsis. Plant Cell *21*, 2237–2252.

Kitakura, S., Vanneste, S., Robert, S., Löfke, C., Teichmann, T., Tanaka, H., and Friml, J.
(2011). Clathrin mediates endocytosis and polar distribution of PIN auxin transporters in
Arabidopsis. Plant Cell *23*, 1920–1931.

Lamesch, P., Berardini, T.Z., Li, D., Swarbreck, D., Wilks, C., Sasidharan, R., Muller, R.,
Dreher, K., Alexander, D.L., Garcia-Hernandez, M., et al. (2012). The Arabidopsis
Information Resource (TAIR): improved gene annotation and new tools. Nucleic Acids Res.
*40*, D1202-1210.

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat
Methods *9*, 357–359.

Larson, E.R., Domozych, D.S., and Tierney, M.L. (2014). SNARE VTI13 plays a unique role
in endosomal trafficking pathways associated with the vacuole and is essential for cell wall
organization and root hair growth in arabidopsis. Ann. Bot. *114*, 1147–1159.

Larson, E.R., Van Zelm, E., Roux, C., Marion-Poll, A., and Blatt, M.R. (2017). Clathrin
Heavy Chain Subunits Coordinate Endo- and Exocytic Traffic and Affect Stomatal
Movement. Plant Physiol. *175*, 708–720.

Lin, D.Y., and Ying, Z. (1994). Semiparametric Analysis of the Additive Risk Model.
Biometrika *81*, 61–71.

Lorenzo, O., Chico, J.M., Sánchez-Serrano, J.J., and Solano, R. (2004). JASMONATE-
INSENSITIVE1 encodes a MYC transcription factor essential to discriminate between
different jasmonate-regulated defense responses in Arabidopsis. Plant Cell *16*, 1938–1950.

Miller, R.G. (1998). Survival Analysis (New York: Wiley & Sons).

Naveed, M., Brown, L.K., Raffan, A.C., George, T.S., Bengough, A.G., Roose, T., Sinclair,
I., Koebernick, N., Cooper, L., Hackett, C.A., et al. (2017). Plant exudates may stabilize or
weaken soil depending on species, origin and time. European Journal of Soil Science *68*,
806–816.

Parker, J.S., Cavell, A.C., Dolan, L., Roberts, K., and Grierson, C.S. (2000). Genetic
interactions during root hair morphogenesis in Arabidopsis. Plant Cell *12*, 1961–1974.

Prentice, R.L., and Kalbfleisch, J.D. (2003). Mixed discrete and continuous Cox regression
model. Lifetime Data Anal *9*, 195–210.

R Core Team (2014). R: A language and environment for statistical computing (Vienna,
Austria: R Foundation for Statistical Computing).

Ringli, C., Bigler, L., Kuhn, B.M., Leiber, R.-M., Diet, A., Santelia, D., Frey, B., Pollmann,
S., and Klein, M. (2008). The Modified Flavonol Glycosylation Profile in the Arabidopsis
rol1 Mutants Results in Alterations in Plant Growth and Cell Shape Formation. The Plant
Cell *20*, 1470–1481.

Robinson, D.G., Scheuring, D., Naramoto, S., and Friml, J. (2011). ARF1 localizes to the
golgi and the trans-golgi network. Plant Cell *23*, 846–849; author reply 849-850.

Rodriguez-Furlán, C., Salinas-Grenet, H., Sandoval, O., Recabarren, C., Arraño-Salinas, P.,
Soto-Alvear, S., Orellana, A., and Blanco-Herrera, F. (2016). The Root Hair Specific
SYP123 Regulates the Localization of Cell Wall Components and Contributes to
Rizhobacterial Priming of Induced Systemic Resistance. Front Plant Sci *7*, 1081.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. Nat. Methods *9*, 676–682.

Šmilauerová, M. (2001). Plant root response to heterogeneity of soil resources: Effects of
nutrient patches, AM symbiosis, and species composition. Folia Geobot *36*, 337–351.

Spruance, S.L., Reid, J.E., Grace, M., and Samore, M. (2004). Hazard ratio in clinical trials.
Antimicrob. Agents Chemother. *48*, 2787–2792.

Stokes, A., Atger, C., Bengough, A.G., Fourcard, T., and Sidle, R.C. (2009). Desirable plant
root traits for protecting natural and engineered slopes against landslides. Plant and Soil *324*,
1–30.

Toukura, Y., Devee, E., and Hongo, A. (2006). Uprooting and shearing resistances in the
seedlings of four weedy species. Weed Biology and Management *6*, 35–43.

Waghmare, S., Lileikyte, E., Karnik, R., Goodman, J.K., Blatt, M.R., and Jones, A.M.E.
(2018). SNAREs SYP121 and SYP122 Mediate the Secretion of Distinct Cargo Subsets.
Plant Physiology *178*, 1679–1688.

Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis (Springer).

Yang, J., Bak, G., Burgin, T., Barnes, W.J., Mayes, H.B., Peña, M.J., Urbanowicz, B.R., and
Nielsen, E. (2020). Biochemical and Genetic Analysis Identify CSLD3 as a beta-1,4-Glucan
Synthase That Functions during Plant Cell Wall Synthesis. Plant Cell *32*, 1749–1767.

**Tables**

**Table 1** *Arabidopsis thaliana* mutant lines used in this study.

| Mutant type | Mutant | AGI code[a] | Mutant line | Referenced in |
|---|---|---|---|---|
| Root hair shape | *cow1-3* | At4g34580 | SALK_0021245 | Böhme *et al.* (2004) |
| | *csld3-1* | At3g03050 | CS899 | Favery *et al.* (2001) |
| | *lrx1-4* | At1g12040 | SALK_057038 | Kim *et al.* (2006) |
| | *rol1-2* | At1g78570 | CS16373 | Diet *et al.* (2006) |
| | *gdi1-2* | At3g07880 | SALK_035400 | Kang *et al.* (2017) |
| Vesicle trafficking | *chc1-2* | At3g11130 | SALK_103252 | Kitakura *et al.* (2011) |
| | *chc2-3* | At3g08530 | SALK_151638 | Bourdais *et al.* (2016) |
| | *syp121* | At3g11820 | - | Assaad *et al.* (2004) |
| | *syp122-1* | At3g52400 | SALK_008617 | Assaad *et al.* (2004) |
| Root exudate composition | *jin1-9* | At1g32640 | SALK_017005 | Anderson *et al.* (2004) |
| | *pdr2* | At4g15230 | SAIL_811_F08 | Badri *et al.* (2008) |
| | *pft1-3* | At1g25540 | SALK_059316 | Kidd *et al.* (2009) |
| Unknown | *abcg43-1* | At4g15236 | N75206[b] | This study |
| | *abcg43-2* | At4g15236 | SALK_201207 | This study |
| | *abcg43-3* | At4g15236 | SALKseq_30713 | This study |

**Figure legends**

**Fig. 1** Physical root hair properties contribute to root-substrate adhesion in *Arabidopsis*
*thaliana.*

(a) Root hair phenotypes of 5-d-old wild-type (Col-0), *lrx 1-4, csld3-1, cow1-3, rol1-2,* and
*gdi1-2* seedlings grown on gel medium and statistical comparisons of mean root hair density
(number per mm length of root) and mean root hair length (mm) for each mutant line relative
to wild type. White asterisks on the root hair images indicate the characteristic root hair
bulging phenotype in the *lrx1-4* mutant and the short, branching root hair phenotype in *gdi1-*
*2*. Scale bar = 0.25 mm. In the table, the mean ± standard error is given for root hair density
and root hair length of wild type and each mutant as well as a mean comparison to wild type,
which is listed in bold. "No difference" is stated when there was no statistically significant
difference between wild type and a mutant line. The statistical output of each univariate
linear model is given. Alpha = 0.025, statistical significance codes: '***' = < 0.001, '**' = <
0.01. The results shown are from a representative experiment for at least two independent
experiments. Survival curves displaying the proportion of seedlings that adhered to the gel at
increasing centrifugal force for (b) 92 wild type (Col-0 – black); 81 *lrx1-4* (light green); 70
*csld3-1* (green); (c) 82 wild type (Col-0 - black); 83 *cow1-3* (green); and (d) 77 wild type
(Col-0- black); 85 *rol1-2* (light green); and 86 *gdi1-2* (dark green). Circled red crosses on the
survival curves represent seedlings that remained adhered to the gel after the maximum
centrifugal speed (1611 RPM). The results shown are from a representative experiment for at
least two independent experiments showing a statistically significant difference in adhesion
between mutant lines relative to wild type (Cox PH regression; alpha = 0.01). A single
experiment included ≥ 70 biological replicates for each candidate line.

**Fig. 2** Vesicle trafficking mechanisms that contribute to root-substrate adhesion in
*Arabidopsis thaliana.*

(a) Root hair phenotypes of 5-d-old wild-type (Col-0), *syp121, syp122-1, chc1-2* and *chc2-3*
seedlings grown on a gel medium and statistical comparisons of mean root hair density
(number per mm length of root) and mean root hair length (mm) for each mutant line relative
to wild type. Scale bar = 0.25 mm. In the table, the mean ± standard error is given for the root
hair density and root hair length of wild type and each mutant as well as a mean comparison
to wild type, which is listed in bold. "No difference" is stated when there was no statistically
significant difference between wild type and a mutant line. The statistical output of each
univariate linear model is given. Alpha = 0.025. The results shown are from a representative
experiment for at least two independent experiments. Survival curves displaying the

proportion of seedlings that adhered to the gel at increasing centrifugal force for (b) 87 wild

type (Col-0 – black); 91 *syp121* (turquoise); 83 *syp122-1* (blue/grey); and (c) 83 wild type

(Col-0 – black); 70 *chc1-2* (dark blue)*; and* 72 *chc2-3* (medium blue)*.* Red crosses circled on

the survival curves represent seedlings that remained adhered to the gel after the maximum

centrifugal speed (1611 RPM). The results shown are from a representative experiment for at

least two independent experiments showing a statistically significant difference in adhesion

between all mutant lines relative to wild type (Cox PH regression; alpha = 0.01). A single

experiment included $\geq$ 70 biological replicates for each candidate line.

**Fig. 3** Exudate composition changes root-substrate adhesion properties in *Arabidopsis*

*thaliana.*

(a) Root hair phenotypes of 5-d-old wild-type (Col-0), *jin1-9, pdr2* and *pft1-3* seedlings

grown on a gel medium and statistical comparisons of mean root hair density (number per

mm length of root) and mean root hair length (mm) for each mutant line relative to wild type.

Scale bar = 0.25 mm. In the table, the mean ± standard error is given for the root hair density

and root hair length of wild type and each mutant as well as a mean comparison to wild type,

which is listed in bold. "No difference" is stated when there was no statistically significant

difference between wild type and a mutant line. The statistical output of each univariate

linear model is given. Alpha = 0.025. The results shown are from a representative experiment

for at least two independent experiments. (b) Survival curves displaying the proportion of

seedlings that adhered to the gel at increasing centrifugal force for 80 wild type (Col-0 –

black), 81 *jin1-9* (light pink), 80 *pdr2* (light orange), and 81 *pft1-3* (dark orange)*.* The results

shown are from a representative experiment for at least two independent experiments

showing a statistically significant difference in adhesion between mutant lines relative to wild

type (Cox PH regression; alpha = 0.01), except for *jin1-9*. A single experiment included $\geq$ 70

biological replicates for each candidate line.

**Fig. 4** Using the centrifuge assay in a forward genetic screen to identify *Arabidopsis thaliana*

root-substrate adhesion mutants.

(a) T-DNA insert locations in *ABCG43* for each *abcg43* mutant allele, with insertions

indicated by orange arrowheads. The *ABCG43* gene is located on chromosome four at

position Chr4: 8,969,677-8,702,727 and contains 23 exons (dark green) and 22 introns (white

gaps). *abcg43-1* has a T-DNA insert in the exon10/intron10 boundary, *abcg43-2*

(SALK_201207) has a T-DNA insert in exon 2 and *abcg43-3* (SALKseq_30713) has a T-

DNA insert in exon 3. (b) Genomic PCR confirming the homozygosity of each *abcg43*

mutant for the respective T-DNA inserts. Lanes **1** and **2** were loaded with the gene-specific

and T-DNA-border PCR products from *abcg43* mutant lines. Lanes **3** and **4** were loaded with

the gene-specific and T-DNA-border PCR products from a wild type (Col-0) genomic DNA

template. Lanes **5** and **6** were loaded with the water controls for the gene-specific and T-

DNA-border PCR reactions, respectively. '**L**' indicates the 100-bp ladder. The expected

product sizes for the gene-specific PCRs were ~350 nt, ~1025 nt and ~1090 nt; the T-DNA-

border PCR product sizes were ~400 nt, ~550 nt and ~750 nt for *abcg43-1, abcg43-2* and

*abcg43-3* alleles, respectively. (c) Survival curves displaying the proportion of seedlings that

adhered to the gel at increasing centrifugal force for 79 wild type (Col-0 – black), 70 *abcg43-*

*1* (dark pink), 85 *abcg43-2* (purple), and 73 *abcg43-3* (light purple). Red crosses circled on

the survival curves represent seedlings that remained adhered to the gel after the maximum

centrifugal speed (1611 RPM). The results are from a representative experiment for at least

two independent experiments showing a statistically significant difference in adhesion

between the mutant lines relative to wild type (Cox PH regression; alpha = 0.01). A single

experiment included $\geq$ 70 biological replicates for each candidate line.

(a)

| Line | Root hair density (number per mm length of root) | | Root hair length (mm) | |
|---|---|---|---|---|
| Col-0 | 30.556 ± 0.784 | - | 0.518 ± 0.009 | - |
| lrx1-4 | 31.889 ± 0.935 **No difference** | t = 1.093, P > 0.05 d.f. = 16, n = 9 | 0.388 ± 0.008 **1.3 times shorter\*\*** | t = -2.819, P < 0.001 d.f. = 16, n = 9 |
| csld3-1 | 8.778 ± 1.188 **3.5 times less\*\*\*** | t = -15.30, P < 0.001 d.f. = 16, n = 9 | 0.045 ± 0.002 **10.1 times shorter\*\*\*** | t = -11.790, P < 0.001 d.f. = 16, n = 9 |
| cow1-3 | 29.200 ± 0.772 **No difference** | t = -1.23, P > 0.05 d.f. = 17, n = 10 | 0.049 ± 0.001 **10 times shorter\*\*\*** | t = -12.37, P < 0.001 d.f. = 17, n = 10 |
| rol1-2 | 30.700 ± 1.550 **No difference** | t = 0.08, P > 0.05 d.f. = 17, n = 9 | 0.195 ± 0.003 **2.7 times shorter\*\*\*** | t = -11.790, P < 0.001 d.f. = 16, n = 9 |
| gdi1-2 | 26.200 ± 1.041 **0.17 times less\*\*** | t = -3.282, P < 0.01 d.f. = 16, n = 9 | 0.135 ± 0.006 **3.8 times shorter\*\*\*** | t = -9.04, P < 0.001 d.f. = 16, n = 9 |

(b)

(c)

(d)

(a)

Col-0    syp121    syp122-1    chc1-2    chc2-3

| Line | Root hair density (number per mm length of root) | | Root hair length (mm) | |
|---|---|---|---|---|
| Col-0 | 32.000 ± 0.632 | | 0.420 ± 0.005 | - |
| syp121 | 31.200 ± 0.952 **No difference** | t = -0.700, P > 0.05 d.f. = 18, n = 10 | 0.461 ± 0.004 **No difference** | t = 1.974, P > 0.05 d.f. = 18, n = 10 |
| syp122-1 | 32.200 ± 0.442 **No difference** | t = 0.259, P > 0.05 d.f. = 18, n = 10 | 0.452 ± 0.005 **No difference** | t = 1.379, P > 0.05 d.f. = 18, n = 10 |
| chc1-2 | 30.889 ± 0.716 **No difference** | t = -1.168, P > 0.05 d.f. = 17, n =9 | 0.441 ± 0.006 **No difference** | t = 0.684, P > 0.05 d.f. = 17, n = 9 |
| chc2-3 | 31.889 ± 0.790 **No difference** | t = -0.111, P > 0.05 d.f. = 17, n =9 | 0.463 ± 0.006 **No difference** | t = 0.182, P > 0.05 d.f. = 18, n = 10 |

(b)

(c)

**(a)**

Col-0      *jin1-9*      *pdr2*      *pft1-3*

| Line | Root hair density (number per mm length of root) | | Root hair length (mm) | |
|---|---|---|---|---|
| Col-0 | 30.111 ± 1.612 | - | 0.416 ± 0.005 | - |
| *jin1-9* | 29.444 ± 1.271 **No difference** | t = -0.324, P > 0.05 d.f. = 16, *n* = 9 | 0.429 ± 0.006 **No difference** | t = 0.555, P > 0.05 d.f. = 16, *n* = 9 |
| *pdr2* | 31.125 ± 0.854 **No difference** | t = 0.533, P > 0.05 d.f. = 15, *n* =8 | 0.428 ± 0.006 **No difference** | t = 0.521, P > 0.05 d.f. = 15, *n* = 8 |
| *pft1-3* | 29.000 ± 1.179 **No difference** | t = -0.555, P > 0.05 d.f. = 16, *n* = 9 | 0.424 ± 0.006 **No difference** | t = 0.295, P > 0.05 d.f. = 16, *n* = 9 |

**(b)**

Legend:
- Col-0
- *jin1-9*
- *pdr2*
- *pft1-3*

Percentage of seedlings adhered to the gel (%) vs Centrifugal force (mN)

**(a)**

acbcg43-2
acbcg43-3

abcg43-1

*ABCG43*
(At4g15236)

Chr4: 8,969,677                                                    Chr4: 8,702,727

**(b)**

kb    L    1    2    3    4    5    6

1.0
0.8

0.5
0.3

*abcg43-1*

1.2
1.0
0.8

0.5

*abcg43-2*

1.2
1.0

0.8

0.5

*abcg43-3*

**(c)**

Percentage of seedlings adhered to the gel (%)

Centrifugal force (mN)

Col-0
abcg43-1
abcg43-2
abcg43-3