A centrifuge-based method for identifying novel genetic traits that affect root-substrate adhesion in *Arabidopsis thaliana*

Bethany M Eldridge[1,*], Emily R Larson[1,*], Laura Weldon[1,a], Kevin M Smyth[1], Annabelle N Sellin[1], Isaac V Chenchiah[2], Tanniemola B Liverpool[2] and Claire S Grierson[1,#]

[1]School of Biological Sciences, University of Bristol, Bristol Life Sciences Building, 24 Tyndall Avenue, Bristol, BS8 1TQ, UK

[2]School of Mathematics, University of Bristol, Fry Building, Woodland Road, Bristol BS8 1UG, UK

[a] Current address: Wildfowl and Wetlands Trust (WWT), Slimbridge, Gloucester, GL2 7BT, UK

[*]Authors contributed equally to this work.

[#]Corresponding author: Claire Grierson

lacsg@bristol.ac.uk

ORCID:
BME: ORCID: 0000-0002-6598-3701
ERL: ORCID: 0000-0002-5498-8152
LW: ORCID: 0000-0002-0761-4414
KMS: ORCID: 0000-0003-0105-8557
IVC: ORCID: 0000-0002-8618-620X
TBL: ORCID: 0000-0003-4376-5604
CSG: ORCID: 0000-0002-4000-0975

Keywords: Arabidopsis, Centrifuge assay, Forward genetics, Root exudates, Root hairs, Root-substrate adhesion, Survival analysis, Vesicle trafficking and Secretion

Running title: Centrifuge-based assay measures root adhesion

ABSTRACT

The physical presence of roots and the compounds they release affect the cohesion between roots and their environment. However, the plant traits that are important for these interactions are unknown and most methods that quantify the contributions of these traits are time-intensive and require specialist equipment and complex substrates. Our lab developed an inexpensive, high-throughput phenotyping assay that quantifies root-substrate adhesion in *Arabidopsis thaliana*. We now report that this method has high sensitivity and versatility for identifying different types of traits affecting root-substrate adhesion including root hair morphology, vesicle trafficking pathways and root exudate composition. We describe a practical protocol for conducting this assay and introduce its use in a forward genetic screen to identify novel genes affecting root-substrate interactions. This assay is a powerful tool for identifying and quantifying genetic contributions to cohesion between roots and their environment.

INTRODUCTION

Plants secrete compounds that help them adapt to their environment, sense and interact with other organisms, and improve water and nutrient uptake (reviewed in Bais et al., 2006; Sasse et al., 2018; Vives-Peris et al., 2020). These compounds (exudates) can vary in composition based on plant species, developmental stage, and environment (Naveed et al., 2017). Recent studies have shown that some compounds have bioadhesive properties that stick plants and soils together and aggregate soil particles, modifying the microenvironment around plant roots (Galloway et al., 2018). However, understanding how plant biology and physiology contribute to plant-soil interactions can be confounded by multiple factors like other organisms living in the soil that also secrete organic metabolites that can contribute to soil cohesion. Soil composition might affect plant exudate composition and vice versa. Also, plant root morphology including root architecture and the presence of root hairs can alter root-soil interactions (Bailey et al., 2002; Burylo et al., 2012; De Baets et al., 2007; Ghestem et al., 2014; Stokes et al., 2009). This complexity has made it difficult to characterise the relevant plant physiology.

We developed a high-throughput assay that identifies and quantifies the contribution of root hair cells to root-substrate cohesion in a less variable environment than the natural rhizosphere (De Baets et al., 2020). The assay uses a centrifuge to apply force to *Arabidopsis thaliana* seedlings grown on the surface of a sterile medium and measures the force required to detach seedlings from that surface, using a Cox proportional hazards regression (Prentice and Kalbfleisch, 2003) to quantify the adherence of candidate Arabidopsis lines relative to

wild-type plants at increasing centrifugal force. Previous use of this assay assessed whether
presence or absence of root hairs affected adhesion to the agar (De Baets *et al.,* 2020) but did
not address subtle aspects of root hair biology. We now wanted to test whether this assay is
sensitive enough to identify other types of traits involved in root-substrate adhesion. To
validate this extended use of the assay, we asked if it could detect the adhesive effects in (1)
mutants with altered root hair shape; (2) vesicle trafficking mutants with no known root hair
morphological defects; (3) exudate compositional mutants with no known root hair
phenotypes; and (4) a forward genetic screen of a mutant population to identify novel genes
that affect root-substrate adhesion.

Here, we show that this method has the sensitivity to identify how differences in root
hair morphology and biological functions that do not present a visible root hair phenotype
contribute to substrate cohesion. Using the genetic and molecular tools available in Arabidopsis
and working in a sterile environment allows hypotheses about plant specific factors of root-
substrate interactions to be tested. While plant-soil cohesion is a complex and dynamic
interaction, our assay provides a way to efficiently probe root cellular functions that might be
masked by confounding variables in a soil-based study system. Results from this assay provide
a platform for more comprehensive studies of plant morphology and physiology that affect
root-substrate cohesion.

MATERIALS AND METHODS
Materials and Equipment
1. 20% [v/v] bleach solution
2. Sterile distilled water
3. Seed for candidate Arabidopsis lines
4. Half strength Murashige and Skoog basal medium
5. 1% [w/v] sucrose
6. 1% [w/v] agar
7. Laminar air hood
8. 90-mm triple vented Petri Plates
9. Parafilm
10. Growth room
11. Swing-out bucket centrifuge
12. Laboratory safety razor blade
13. Analytical balance

Plant Material and Growth Conditions

Candidate and wild-type lines of the Columbia-0 (Col-0) ecotype of *Arabidopsis thaliana* single-gene mutants were obtained from the Nottingham Arabidopsis Stock Centre (NASC; Nottingham, UK) or were stocks maintained in the lab (**Table 1**). Homozygous lines were used in all experiments.

For all experiments, seeds were surface sterilised for 15 min in 20% [v/v] bleach and sterile water solution, followed by five washes in sterile water. All sterilised seeds were stratified in the dark at 4°C for 24-48 h in Eppendorf microtubes containing sterile water. All plants were reared in a growth room at 21-22°C with a continuous photoperiod (light intensity = 120-145 µmol m^{-2} s^{-1}) at 60% humidity.

Half strength Murashige and Skoog basal medium was supplemented with 1% [w/v] sucrose pH adjusted to 5.7, solidified with 1% [w/v] agar. A 30 ml volume of sterilized media (at ~45 – 50°C) was pipetted into 90-mm triple vented Petri plates and left to set for ~30 min in a laminar air hood. Sterilized seeds were sown onto Petri plates containing the sterile medium and sealed with Parafilm. Plates were then placed in stacks of five at ~80° from the horizontal (lid side up) to encourage vertical growth along the gel surface. Plates were incubated in the growth chamber for 5-6 days before centrifugation. Plants for genomic DNA extractions were grown in compost (three parts Levington F3 compost and one-part J Arthur Bowers horticultural silver sand).

Centrifuge Assay Set-Up

To measure the adhesion of Arabidopsis seedling roots using a centrifuge, seedling roots must not penetrate the gel or touch each other, as both events could interfere with seedling detachment during centrifugation. Therefore, ten seeds were sown 1-cm apart on each plate in two parallel rows (**Figure 1Ai**). The plates were then grown for 5-6 days (**Figure 1Aii**) and seedlings that were touching each other, had grown into the gel, or not uniform in size were omitted from the assay. The seedlings were then numbered so that plate positional effects could be included as a potential covariate in the final analysis (**Figure 1Bi**). Plates were inverted and placed into a swing-out bucket centrifuge (Beckman Allegra X-30R Centrifuge) that held four plates at a time and subjected to increments of increasing centrifugal force at 720, 1018, 1247, 1440 and 1611 RPM for one minute each (**Figure 1Bii**). The proportion of seedlings that detached from the gel surface was recorded between each centrifugal speed; a detachment event was recorded when a seedling had partially or fully peeled away from the gel (**Figure**

1Biii). If the gel shattered during the assay, seedlings that had not detached from the gel were
not recorded and seedlings that remained adhered to the gel after the maximum centrifugal
speed were censored in the analysis. During method optimisation we encountered some
common methodological challenges for which we provide troubleshooting options in **Table**
S1.

Calculating Seedling Detachment Force

To calculate the centrifugal force acting on a seedling, the aerial tissue mass for each
seedling was separated from the root by making an incision at the hypocotyl-root junction using
a laboratory safety razor blade and immediately measured using an analytical balance (**Figure**
1Ci, ii). The aerial tissue weight of each seedling was measured because a plant with heavier
aerial tissue experiences more force than a lighter one subjected to the same centrifugal speed.
Centrifugal force, Fc (mN) acting on a seedling was calculated as previously reported in De
Baets et al. (2020). Aerial tissue weight (kg), the angular velocity (ω) and the distance between
the seedling and the axis of rotation on the centrifuge (radius = 0.07m) were used to give the
following equation:

Fc= *mass* x *radius* x ω^2 (Equation 1)

Identification of *abcg43-1* Mutant T-DNA Insert

Genomic DNA was extracted from a pool of vegetative tissue taken from ten three-
week old plants using an adapted protocol (Healey et al., 2014) for Illumina Next-Generation
two x paired-end sequencing conducted at the Bristol Genomics Facility (University of Bristol,
UK). The Illumina TruSeq Nano LT gDNA kit (Illumina Inc) was used to generate a genomic
DNA sequencing library following the manufacturer's protocol. The final library was diluted
to a loading concentration of 1.4 pM for cluster generation and 2 x 150 bp paired-end
sequencing on the Illumina NextSeq500 system (Illumina Inc) alongside a 5% PhiX spike-in
control library. Read summary statistics were generated using the RTA 2.4.6 Primary Analysis
Software (Illumina Inc). The read summaries were analysed in the Sequencing Analysis
Viewer (Illumina Inc). Filtered paired reads were subject to paired-end alignment using the
Bowtie2 2.3.4.2 aligner (Langmead and Salzberg, 2012). A bespoke reference genome was
produced that combined the TAIR10 Arabidopsis genome (Lamesch et al., 2012) and the
pROK2 vector sequence (Baulcombe et al., 1986). Alignments were viewed in the Integrative
Genomics Viewer IGV 2.3 (Robinson et al., 2011).

For Sanger sequencing, DNA was extracted from a pool of vegetative tissue taken from
two-week-old plants using a modified Edwards prep (Edwards et al., 1991). High fidelity PCR
was conducted on the *abcg43-1* line using the Q5 High-Fidelity 2X Master Mix (NEB). PCR
products were purified and extracted from a 1% agarose gel using the QIAquick Gel Extraction
Kit (QIAGEN), following the manufacturer's instructions. Purified PCR product plus the
abcg43-1 forward, reverse or border genotyping primer (**Table S2**) was used for Sanger
sequencing (using the Mix2Seq overnight sequencing kit, Eurofins Genomics).
Chromatograms and FASTA files were obtained from Eurofin Genomics and following manual
low-quality end trimming, the final sequences were aligned to the pROK2 vector sequence and
ABCG43 gene sequence using the MUSCLE alignment tool (Madeira et al., 2019).
181

Genotyping

183 To genotype the *abcg43* mutant alleles, genomic DNA was extracted from the vegetative tissue
184 of two-week-old plants using a modified Edwards prep (Edwards et al., 1991). T-DNA border
185 and gene-specific primers were used in PCR analyses to confirm the genotype (**Table S2**).
186

Morphological Root Trait Analyses

Five-day-old Arabidopsis seedlings were imaged with a Leica MZ FLIII microscope
(Leica) with dark-field lighting. Root hair length and density were measured using microscope
images of five-day-old Arabidopsis seedlings and Fiji version 1.0 (Schindelin et al., 2012) and
the Bio-Formats Importer plugin. At least two experimental repeats were conducted; in each
experiment, we imaged eight to ten individual plants from each line and measured at least 30
root hairs per plant.
194

Statistical Analyses

196   All statistical analyses were conducted using RStudio, version 1.1453 (R Core Team,
197 2014) and all graphs were generated using the R package ggplot2 (Wickham, 2016).
198

Centrifuge assay

Survival analysis was used to study when seedlings detach from the sterile gel medium (i.e.
seedling detachment is the event) at increasing force (i.e. force is modelled rather than time).
We used the Cox proportional hazards (PH) regression function, coxph, within the survival
package in R to statistically test for differences between the detachment of experimental
seedlings relative to the wild-type control (Col-0), using centrifugal force as the predictor

variable (**Figure 1Ciii**). We incorporated seedling position and individual plate number as
covariates within our Cox PH regression models to account for any environmental
heterogeneity such as minor inconsistencies in the water content or gelling of the medium.
When these covariates had no significant effect, they were removed from the model. Seedlings
that remained adhered to the gel after the maximum centrifugal speed were censored in our
Cox PH regression model. Comparisons of different candidate lines relative to the wild type
line were analysed *a priori*; therefore, a series of contrasts were set up using the R function
contr.treatment rather than using post-hoc testing methods.

For each Cox PH regression model run, we report the P value of the Wald Statistic (z-score)
and the hazard ratio with the upper and lower bound confidence intervals. Hazard ratios are
exponentiated coefficients that give the instantaneous risk of root-gel detachment (Cox &
Oakes, 1984; Bradburn *et al.*, 2003; Spruance *et al.*, 2004). For each Cox PH regression model
conducted, we used the hazard ratio as a measure of effect size to assess differences between
the root-gel adhesion of candidate lines and wild type (Cox & Oakes, 1984). Wild type was
used as the baseline group for all models, with a hazard ratio of one. A candidate line with
higher risk of detachment from gel than wild type will have a hazard ratio above one.
Conversely, lines with a lower risk of detachment than wild type will have a hazard ratio below
one.

For all Cox PH regression models, the assumption of proportionality was satisfied and an alpha
level of 0.01 was used. The root-gel detachment of at least 70 plants from each line was
assessed within a single experiment.

Root hair analyses
Two sample t-tests were conducted using the t.test function in R to test for differences in the
mean root hair density or mean root hair length (of eight to ten individual plants from each
line) between candidate lines relative to wild type (Col-0). In all cases, the assumptions of
normality and homoscedasticity were satisfied. Because comparisons of different mutant lines
relative to the wild type line were *a priori*, a series of contrasts were conducted using the
contr.treatment function in R. To prevent type I errors from multiple testing, we applied the
Bonferroni method and adjusted the alpha level to 0.025 (0.05/2).

RESULTS

Root Hair Morphological Effects on Root-Substrate Adhesion

We screened mutant lines with known root hair morphology defects to ask if the centrifuge assay could determine if root hair shape affects root-substrate adhesion. The mostly bald mutant *csld3*, which encodes the CESA-like 3D protein (Favery et al., 2001; Wang et al., 2001; Yang et al., 2020) was included as a root hairless example. For lines that produce short or malformed root hairs, we included the *gdi1-2* Rho GTPase GDP dissociation inhibitor mutant, *rol1-2* rhamnose biosynthesis 1 mutant allele, *can of worms 1* (*cow1-3*) phosphatidylinositol transferase protein mutant, and *lrx1-4* leucine rich extensin 1 mutant (Baumberger et al., 2001; Böhme et al., 2004; Carol et al., 2005; Diet et al., 2006; Grierson et al., 1997). The *rol1-2* and *gdi1-2* mutants have short root hairs and the root hairs on *gdi1-2* plants can also branch (Diet et al., 2006; Kang et al., 2017; Parker et al., 2000; Ringli et al., 2008); *cow1-3* plants have short, wide root hairs that can branch; and *lrx1-4* plants can develop root hairs that abort, swell, branch, or collapse in a growth condition-dependent manner (Baumberger et al., 2001; Böhme et al., 2004; Grierson et al., 1997; Parker et al., 2000). Assay results for ectopic and long root hairs were reported previously (De Baets et al., 2020). Under our growth conditions, we confirmed the reported phenotypes for each mutant line (**Figure 2A**). Wild type was used as the baseline group for all centrifuge experiments, with a hazard ratio of one. Therefore, a candidate line with a higher risk of detachment from the gel than wild type will have a hazard ratio above one, while lines with a lower risk of detachment than wild type will have a hazard ratio below one.

Consistent with previous results (De Baets et al., 2020), bald (*csld3*) seedlings detached at lower centrifugal forces relative to wild-type seedlings, with 4.6 times the risk of detachment than wild-type plants (P < 0.001 – **Table 2**; **Figure 2B**). Mutants with deformed root hair phenotypes including *gdi1-2, rol1-2, cow1-3* and *lrx1-4* had risks of detachment that were 2.5, 4, 3.8 and 3.9 times that of wild-type plants, respectively (for all lines P < 0.001 - **Table 2**; **Figure 2B-D**). These results indicated that the centrifuge assay can quantify the effects of root hair shape on root-substrate cohesion.

Vesicle Trafficking Mutations Affect Root-Substrate Adhesion

We hypothesized that Arabidopsis mutants with defective vesicle trafficking pathways might alter cell wall and apoplast characteristics that contribute to root-substrate interactions and adhesion. We chose the secretory Soluble NSF (N-ethylmaleimide sensitive fusion protein) Attachment proteins (SNAP) REceptor (SNARE) mutants *syp121* and *syp122-1*, and the

endocytic mutants, *chc1-2* and *chc2-3* as candidates for our assay. SYP121 is the primary

secretory SNARE found on the plasma membrane (Assaad et al., 2004; Geelen et al., 2002)

and has characterized aboveground phenotypes, including small rosettes and stomatal mobility

defects (Eisenach et al., 2012; Larson et al., 2017); however, the mutant does not have a

reported root hair phenotype. Although the related SNARE SYP122 is thought to share partial

functionality with SYP121, the *syp122-1* mutant does not share these phenotypes with *syp121*

and a recent proteomic analysis reported differences in their vesicle cargoes, which could

contribute to their functional independence (Waghmare et al., 2018). The *chc1-2* and *chc2-3*

mutants are defective in the heavy chain subunits of the clathrin coat complex, which is

required for vesicle traffic at the plasma membrane (Kitakura et al., 2011; Mbengue et al.,

2016). Endo- and exocytic rates are impaired in both mutants (Larson et al., 2017), but no root

hair phenotypes have been reported. For the centrifuge assay, it was important to select mutants

that do not have root hair phenotypes given the effect root hair morphology has on root-gel

adhesion. Therefore, we evaluated the root hairs of these vesicle trafficking mutants and found

that the root hairs of these lines did not significantly differ from wild-type plants (**Figure 3A**).

Overall, the risk of detachment for *syp121, syp122-1, chc1-2* and *chc2-3* plants was 3.6, 3.3,

4.5 and 3.9 times that of wild type, respectively (for all lines $P < 0.001$ – Table 2; **Figure**

3B,C). These results indicated that the centrifuge assay can be used to probe for intracellular

molecular machinery that affect root-substrate interactions.

Exudate Composition Changes Root Detachment Rates

Over 20% of plant assimilated carbon is released as root exudates (Huang et al., 2016).

These are complex mixtures of diverse molecular forms that can modify the external

environment in response to abiotic and biotic stimuli. Some exudate components contribute to

soil adhesion (Akhtar et al., 2018; Galloway et al., 2018). Root epidermal and hair cells secrete

compounds that contribute to root exudate profiles that can be plant species specific (Naveed

et al., 2017). Because mutants in both root hair development and vesicle trafficking pathways

showed root-substrate adhesion phenotypes (**Figure 2 and 3**), we asked if the centrifuge assay

could identify effects of exudate composition on root-substrate adhesion. We selected

Arabidopsis mutants reported to have altered exudate composition compared to wild type,

including *pft1-3*, *jin1-9*, and *pdr2* (Badri et al., 2009; Berger et al., 1996; Carvalhais et al.,

2015; Kidd et al., 2009). *PFT1* (*MED25*) encodes the MEDIATOR25 subunit of the Mediator

nuclear protein, and *JIN1* (*MYC2*) encodes a basic helix-loop-helix leucine zipper transcription

factor; both proteins are involved in the jasmonate signalling pathway (Kidd et al., 2009;

Lorenzo et al., 2004). *pft1-3* and *jin1-9* mutants are reported to have altered root exudate
composition, including lower amounts of the amino acids asparagine, ornithine and tryptophan
than wild-type plants (Carvalhais et al., 2015). *PDR2* (*ABCG30*) encodes a pleiotropic drug
resistance (PDR) full-length ABC transporter that is involved in ABA transport and the
exudation of secondary metabolites (Badri et al., 2008; Kang et al., 2015). We did not observe
root hair growth phenotypes in these mutant lines compared to wild type (**Figure 4A**).

Compared to wild-type seedlings, *pdr2* seedlings resisted detachment, with 0.38 times
the risk of detaching from the gel (P<0.001 – **Table 2**; **Figure 4B**). Conversely, *pft1-3* seedlings
had an increased risk of detachment 1.7 times that of wild-type plants (P < 0.001) and there
was no difference in gel adhesion between *jin1-9* and wild-type plants (P>0.05) (**Table 2;**
Figure 4B). These results show that exudate composition can alter root-substrate interactions,
although this may not be the case for all exudate compositional changes, which may lay outside
the sensitivity range of the assay.

Centrifuge Assay as a Method for Forward Genetic Screening

Having demonstrated that our centrifugation technique can identify different types of
traits that affect root-substrate adhesion, we used it in a forward genetic screen to find novel
genes involved in root-substrate adhesion. To conduct this screen, the root-gel adhesion
properties of individual plants from a pooled SALK T-DNA insertion mutant collection were
analysed (Alonso et al., 2003). Individual plants that detached at the lowest centrifugal speed
or remained attached to the gel after being spun at the highest centrifugal speed were recovered
from the screen and self-fertilized to obtain progeny and identify the putative mutations. The
centrifuge assay was used to rescreen the progeny (using a sample size of ≥ 70 *n*) to confirm
whether candidate lines had significantly increased or decreased root-gel adhesion relative to
wild-type plants. From this screen, we identified a line with significantly enhanced root-gel
adhesion and conducted genomic Next-Generation sequencing followed by Sanger sequencing
of T-DNA flanking PCR products to confirm a T-DNA insertion in the ABC transporter gene,
ABCG43, whose function is unknown. We named this T-DNA insertion line *abcg43-1* and
identified two additional mutant alleles, *abcg43-2* (SALK_201207c) and *abcg43-3*
(SALKseq_30713) (**Figure 5A**).

We analysed the root-gel adhesion of the homozygous insertional *ABCG43* mutant
alleles and found *abcg43-1*, *abcg43-2* and *abcg43-3* had detachment risks 0.25, 0.36 and 0.35
times that of wild-type plants, respectively (for all lines P < 0.001- **Table 2**; **Figure 5B, C**).
These results illustrate the use of the centrifuge assay as a powerful screening method for

identifying novel genes involved in root-substrate adhesion, as well as a tool for evaluating the effects of known gene function on root-substrate interactions.

DISCUSSION

To date, most published methods that quantify the contribution of particular traits to plant-soil interactions are time-intensive and expensive (Bailey et al., 2002; De Baets and Poesen, 2010; De Baets et al., 2020; Toukura et al., 2006). In contrast, our assay can produce data within a week, is available to any laboratory with access to a bench top centrifuge and does not require speciality consumables. This method is a high-throughput and quantitative way to test the effects of root morphology and cell function on root-substrate adhesion and to screen for new biological and molecular factors that can alter plant-substrate interactions. Using a defined model system like Arabidopsis and sterile medium conditions permits the identification of plant-specific characteristics that can then be probed in other substrate- or soil-based conditions.

Using the Centrifuge Assay to Investigate the Role of Root Hairs in Root-Substrate Interactions

Results presented in this paper show that the centrifuge assay can quantify effects of root hair morphology on plant-substrate adhesion. Previous work has shown that the presence of root hairs can significantly enhance root-substrate adhesion compared to the absence of root hairs (Choi and Cho, 2019; De Baets et al., 2020; Haling et al., 2013). We now show that our centrifuge assay can distinguish the adherence strength of seedlings with altered root hair shapes and sizes (**Figure 2**) and can quantify the contribution these morphological traits make to plant-substrate interactions. Some mutants with defective root hairs have mutations in cell wall biosynthesis and modification proteins, so it is possible that altered root-substrate interface compromises adhesion in these mutants, not just root hair shape. Using our method to rapidly phenotype mutants with root hair growth defects could direct additional experiments to characterize biochemical and morphological properties of genes of interest. Indeed, current evidence for the importance of root hairs for the deposition of complex polysaccharides with soil-binding properties (Galloway et al., 2020) supports the need for assays such as ours to expedite the identification of genes and mechanisms that improve soil-remediation, top soil maintenance, and erosion tolerance of at-risk land.

Identifying New Genes Involved in Root-Substrate Adhesion

Mutations in cellular functions that do not alter root hair growth but do affect substrate

adhesion can also be investigated using this assay. Vesicle trafficking pathways are required

for cell wall deposition and maintenance (De Caroli et al., 2011; Larson et al., 2014; Rodriguez-

Furlán et al., 2016); therefore, we hypothesized that mutants in vesicle trafficking pathways,

particularly those at the plasma membrane, could affect root-substrate adhesion. Similarly, the

composition and deposition of plant exudates help plants modify and optimize their growth

conditions (Carvalhais et al., 2015; Naveed et al., 2017), suggesting that modification to plant

exudate composition could affect root-substrate interactions. Consistent with these hypotheses,

mutants with vesicle trafficking defects and altered exudate composition had root-substrate

adhesion properties that differed from wild type, indicating that this method can identify

aspects of plant cell biology important for root-substrate interactions that do not affect root or

root hair morphology (**Figures 3 and 4**). Our results further indicate that while root hairs

provide significant adhesive and cohesive support both on solid gel and soil media (De Baets

et al., 2020), there are additional cellular factors contributing to plant-substrate interactions that

are not directly observable without the use of expensive reagents, development of time-

intensive transgenic lines, and microscopy equipment. Our centrifuge assay provides an

affordable and quick alternative or complement to these current methodologies.

The compounds and interactions that contribute to root-substrate adhesion are complex,

and the direct and indirect factors that participate in these interactions still remain to be

characterised in detail. Many root cell wall or secreted compounds might directly affect

adhesive characteristics of the root surface, such as arabinogalactan proteins involved in the

adhesion of English ivy (Huang et al., 2016). Other complex polysaccharides have also been

shown to enhance soil binding in both Arabidopsis and cereal root systems (Galloway et al.,

2018, 2020). Indeed, mutants in genes associated with the production of these compounds

would be potentially excellent candidates for the application of this assay, which would

quantify the contribution these compounds make to root-substrate adhesion.

We cannot expect all exudate components to present adhesive phenotypes because not

all will affect root-substrate adhesion, and some will affect adhesion in a way that is not

detected by this assay. For example, the *jin1-9* mutant tested in our exudate cohort was not

significantly more or less adhesive compared to wild-type roots (**Figure 4**) despite its altered

exudate composition (Anderson et al., 2004). Some exudate components will play no role in

adhesion, while others may require additional co-factors present in soil or non-sterile media to

participate in root-substrate interactions. Although these are examples of the limitations of the

assay, this assay does allow for the identification of direct contributions that plants make to
root-substrate cohesion and its robust statistical power and high-throughput aspects will
significantly improve genetic screening timelines and enhance large-scale experimental design.

Forward genetic screening relies on distinctive phenotypes that distinguish differences
within large, genetically diverse populations. High throughput, creative techniques for
identifying phenotypes within morphologically similar populations, such as thermal imaging
(Plessis et al., 2011) and confocal imaging platforms (Drakakaki et al., 2011; Robert et al.,
2008) require specific equipment that not all laboratories have at their disposal. In contrast, the
centrifuge assay can be used as a screen to identify new genes affecting plant-substrate
interactions without specialist equipment, making it an accessible option for research groups
that may not otherwise be able to efficiently conduct genetic screens. Moreover, the centrifuge
assay does not require morphological or conditional differences for candidate selection. As an
example, we report the identification in a forward genetics screen of a mutant population in
which *abcg43* mutant seedlings were identified and shown to be more adhesive to gel than wild
type (**Figure 5**). The function of ABCG43 in Arabidopsis is uncharacterised, indicating that
this assay can identify novel candidate genes that have no previously known association with
root morphological defects, adhesion or exudate composition for future study.

CONCLUSION

This assay has applications in plant cell biology and genetics to investigate how diverse
gene activities, cell functions and root structure affect plant-soil adhesion, and identify the
molecular pathways involved. Potential applications not tested in this report include the
measurement of the effects environmental conditions have on plant-substrate adhesion and how
other plant species or ecotypes respond to detachment forces. Given the recent discovery that
root hair presence is important for Arabidopsis cohesion between roots and gel, and compost
or soil (De Baets *et al.,* 2020), this assay can broadly contribute to plant and soil sciences
through the identification of plant traits that increase root-substrate cohesion and potentially
stabilise slopes or reduce soil erosion.

SUPPLEMENTARY INFORMATION
Table S1 Centrifuge assay troubleshooting
Table S2 Primers used in this study for genotyping *Arabidopsis thaliana* mutants

ACKNOWLEDGEMENTS

This work was supported by a Leverhulme Trust project grant RPG-2013-260 to CSG and TBL. BME was supported by a BBSRC SWBio PhD studentship (grant BB/M009122/1). We are thankful to Don Grierson FRS and Enrico Coen FRS who independently suggested the centrifuge assay concept, Timothy Quine for providing insightful guidance during method development, and Thomas Denbigh for providing technical advice. We are grateful to Jill Harrison, Victoria Spencer, Zoe Nemec Venza, Sophie Carpenter and Nicholas H Fair for their helpful comments during revision of the manuscript. This manuscript has been released as a pre-print at [http://doi.org/10.1101/2020.05.14.095448] (Eldridge et al., 2020).

AUTHOR CONTRIBUTIONS

CSG designed the initial project, conducted proof-of-concept pilot experiments, obtained funding, and recruited and advised the team. TBL and IC developed the theoretical parameters for the centrifugal assay and the bounds on the errors from other mechanical forces. BME and ERL planned and conducted all experiments apart from the initial genetic screen, which was carried out by LW and KMS. BME and ERL analysed all of the data presented herein. AS and BME optimised the centrifuge assay data collection and data analysis methods. BME and ERL wrote the initial manuscript. BME, ERL, and CSG edited the manuscript. All authors reviewed and approved the final manuscript.

REFERENCES

Akhtar, J., Galloway, A.F., Nikolopoulos, G., Field, K.J., and Knox, P. (2018). A quantitative method for the high throughput screening for the soil adhesion properties of plant and microbial polysaccharides and exudates. Plant Soil *428*, 57–65.

Alonso, J.M., Stepanova, A.N., Leisse, T.J., Kim, C.J., Chen, H., Shinn, P., Stevenson, D.K., Zimmerman, J., Barajas, P., Cheuk, R., et al. (2003). Genome-wide insertional mutagenesis of Arabidopsis thaliana. Science *301*, 653–657.

Anderson, J.P., Badruzsaufari, E., Schenk, P.M., Manners, J.M., Desmond, O.J., Ehlert, C., Maclean, D.J., Ebert, P.R., and Kazan, K. (2004). Antagonistic Interaction between Abscisic Acid and Jasmonate-Ethylene Signaling Pathways Modulates Defense Gene Expression and Disease Resistance in Arabidopsis. The Plant Cell *16*, 3460–3479.

Assaad, F.F., Qiu, J.-L., Youngs, H., Ehrhardt, D., Zimmerli, L., Kalde, M., Wanner, G., Peck, S.C., Edwards, H., Ramonell, K., et al. (2004). The PEN1 Syntaxin Defines a Novel

Cellular Compartment upon Fungal Attack and Is Required for the Timely Assembly of
Papillae. Mol. Biol. Cell *15*, 5118–5129.

Badri, D.V., Loyola-Vargas, V.M., Broeckling, C.D., De-la-Peña, C., Jasinski, M., Santelia,
D., Martinoia, E., Sumner, L.W., Banta, L.M., Stermitz, F., et al. (2008). Altered Profile of
Secondary Metabolites in the Root Exudates of Arabidopsis ATP-Binding Cassette
Transporter Mutants. Plant Physiology *146*, 762–771.

Badri, D.V., Quintana, N., El Kassis, E.G., Kim, H.K., Choi, Y.H., Sugiyama, A., Verpoorte,
R., Martinoia, E., Manter, D.K., and Vivanco, J.M. (2009). An ABC transporter mutation
alters root exudation of phytochemicals that provoke an overhaul of natural soil microbiota.
Plant Physiol. *151*, 2006–2017.

Bailey, P.H.J., Currey, J.D., and Fitter, A.H. (2002). The role of root system architecture and
root hairs in promoting anchorage against uprooting forces in Allium cepa and root mutants
of Arabidopsis thaliana. J. Exp. Bot. *53*, 333–340.

Bais, H.P., Weir, T.L., Perry, L.G., Gilroy, S., and Vivanco, J.M. (2006). The role of root
exudates in rhizosphere interactions with plants and other organisms. Annu Rev Plant Biol
57, 233–266.

Baulcombe, D.C., Saunders, G.R., Bevan, M.W., Mayo, M.A., and Harrison, B.D. (1986).
Expression of biologically active viral satellite RNA from the nuclear genome of transformed
plants. Nature *321*, 446–449.

Baumberger, N., Ringli, C., and Keller, B. (2001). The chimeric leucine-rich repeat/extensin
cell wall protein LRX1 is required for root hair morphogenesis in Arabidopsis thaliana.
Genes Dev. *15*, 1128–1139.

Berger, S., Bell, E., and Mullet, J.E. (1996). Two Methyl Jasmonate-Insensitive Mutants
Show Altered Expression of AtVsp in Response to Methyl Jasmonate and Wounding. Plant
Physiology *111*, 525–531.

Böhme, K., Li, Y., Charlot, F., Grierson, C., Marrocco, K., Okada, K., Laloue, M., and
Nogué, F. (2004). The Arabidopsis COW1 gene encodes a phosphatidylinositol transfer
protein essential for root hair tip growth. Plant J. *40*, 686–698.

Burylo, M., Rey, F., Mathys, N., and Dutoit, T. (2012). Plant root traits affecting the
resistance of soils to concentrated flow erosion. Earth Surface Processes and Landforms *37*,
1463–1470.

Carol, R.J., Takeda, S., Linstead, P., Durrant, M.C., Kakesova, H., Derbyshire, P., Drea, S.,
Zarsky, V., and Dolan, L. (2005). A RhoGDP dissociation inhibitor spatially regulates
growth in root hair cells. Nature *438*, 1013–1016.

Carvalhais, L.C., Dennis, P.G., Badri, D.V., Kidd, B.N., Vivanco, J.M., and Schenk, P.M.
(2015). Linking Jasmonic Acid Signaling, Root Exudates, and Rhizosphere Microbiomes.
Mol. Plant Microbe Interact. *28*, 1049–1058.

Choi, H.-S., and Cho, H.-T. (2019). Root hairs enhance Arabidopsis seedling survival upon
soil disruption. Scientific Reports *9*, 11181.

Collins, N.C., Thordal-Christensen, H., Lipka, V., Bau, S., Kombrink, E., Qiu, J.-L.,
Hückelhoven, R., Stein, M., Freialdenhoven, A., Somerville, S.C., et al. (2003). SNARE-
protein-mediated disease resistance at the plant cell wall. Nature *425*, 973–977.

De Baets, S., and Poesen, J. (2010). Empirical models for predicting the erosion-reducing
effects of plant roots during concentrated flow erosion. Geomorphology *118*, 425–432.

De Baets, S., Poesen, J., Knapen, A., and Galindo, P. (2007). Impact of root architecture on
the erosion-reducing potential of roots during concentrated flow. Earth Surface Processes and
Landforms *32*, 1323–1345.

De Baets, S., Denbigh, T.D., Smyth, K.M., Eldridge, B.M., Weldon, L.M., Higgins, B.W.,
Matyjaszkiewicz, A.W., Meersmans, J., Larson, E.R., Chenchiah, I.V., et al. (2020). Micro-
scale interactions between Arabidopsis root hairs and soil particles influence soil erosion.
Nature Comms Biol *3*, 1–11.

De Caroli, M., Lenucci, M.S., Di Sansebastiano, G.-P., Dalessandro, G., De Lorenzo, G., and
Piro, G. (2011). Dynamic protein trafficking to the cell wall. Plant Signal Behav *6*, 1012–
1015.

Diet, A., Link, B., Seifert, G.J., Schellenberg, B., Wagner, U., Pauly, M., Reiter, W.-D., and
Ringli, C. (2006). The Arabidopsis root hair cell wall formation mutant lrx1 is suppressed by
mutations in the RHM1 gene encoding a UDP-L-rhamnose synthase. Plant Cell *18*, 1630–
1641.

Drakakaki, G., Robert, S., Szatmari, A.-M., Brown, M.Q., Nagawa, S., Van Damme, D.,
Leonard, M., Yang, Z., Girke, T., Schmid, S.L., et al. (2011). Clusters of bioactive
compounds target dynamic endomembrane networks in vivo. Proceedings of the National
Academy of Sciences *108*, 17850–17855.

Edwards, K., Johnstone, C., and Thompson, C. (1991). A simple and rapid method for the
preparation of plant genomic DNA for PCR analysis. Nucleic Acids Res *19*, 1349.

Eisenach, C., Chen, Z.-H., Grefen, C., and Blatt, M.R. (2012). The trafficking protein
SYP121 of Arabidopsis connects programmed stomatal closure and K^+ channel activity with
vegetative growth. Plant J. *69*, 241–251.

Eldridge, B.M., Larson, E.R., Weldon, L., Smyth, K.M., Sellin, A.N., Chenchiah, I.V.,
Liverpool, T.B., and Grierson, C.S. (2020). A centrifuge-based method for identifying novel
genetic traits that affect root-substrate adhesion in Arabidopsis thaliana. BioRxiv
2020.05.14.095448.

Favery, B., Ryan, E., Foreman, J., Linstead, P., Boudonck, K., Steer, M., Shaw, P., and
Dolan, L. (2001). KOJAK encodes a cellulose synthase-like protein required for root hair cell
morphogenesis in Arabidopsis. Genes Dev. *15*, 79–89.

Galloway, A.F., Pedersen, M.J., Merry, B., Marcus, S.E., Blacker, J., Benning, L.G., Field,
K.J., and Knox, J.P. (2018). Xyloglucan is released by plants and promotes soil particle
aggregation. New Phytol. *217*, 1128–1136.

Galloway, A.F., Akhtar, J., Marcus, S.E., Fletcher, N., Field, K., and Knox, P. (2020). Cereal
root exudates contain highly structurally complex polysaccharides with soil-binding
properties. The Plant Journal *103*, 1666–1678.

Geelen, D., Leyman, B., Batoko, H., Di Sansebastiano, G.-P., Moore, I., Blatt, M.R., and Di
Sansabastiano, G.-P. (2002). The abscisic acid-related SNARE homolog NtSyr1 contributes
to secretion and growth: evidence from competition with its cytosolic domain. Plant Cell *14*,
387–406.

Ghestem, M., Cao, K., Ma, W., Rowe, N., Leclerc, R., Gadenne, C., and Stokes, A. (2014). A
framework for identifying plant species to be used as "ecological engineers" for fixing soil on
unstable slopes. PLoS ONE *9*, e95876.

Grierson, C.S., Roberts, K., Feldmann, K.A., and Dolan, L. (1997). The COW1 locus of
arabidopsis acts after RHD2, and in parallel with RHD3 and TIP1, to determine the shape,
rate of elongation, and number of root hairs produced from each site of hair formation. Plant
Physiol *115*, 981–990.

Haling, R.E., Brown, L.K., Bengough, A.G., Young, I.M., Hallett, P.D., White, P.J., and
George, T.S. (2013). Root hairs improve root penetration, root-soil contact, and phosphorus
acquisition in soils of different strength. J. Exp. Bot. *64*, 3711–3721.

Healey, A., Furtado, A., Cooper, T., and Henry, R.J. (2014). Protocol: a simple method for
extracting next-generation sequencing quality genomic DNA from recalcitrant plant species.
Plant Methods *10*, 21.

Huang, Y., Wang, Y., Tan, L., Sun, L., Petrosino, J., Cui, M.-Z., Hao, F., and Zhang, M.
(2016). Nanospherical arabinogalactan proteins are a key component of the high-strength
adhesive secreted by English ivy. Proc. Natl. Acad. Sci. U.S.A. *113*, E3193-3202.

Kang, E., Zheng, M., Zhang, Y., Yuan, M., Yalovsky, S., Zhu, L., and Fu, Y. (2017). The
Microtubule-Associated Protein MAP18 Affects ROP2 GTPase Activity during Root Hair
Growth1[OPEN]. Plant Physiol *174*, 202–222.

Kang, J., Yim, S., Choi, H., Kim, A., Lee, K.P., Lopez-Molina, L., Martinoia, E., and Lee, Y.
(2015). Abscisic acid transporters cooperate to control seed germination. Nat Commun *6*,
8113.

Kidd, B.N., Edgar, C.I., Kumar, K.K., Aitken, E.A., Schenk, P.M., Manners, J.M., and
Kazan, K. (2009). The mediator complex subunit PFT1 is a key regulator of jasmonate-
dependent defense in Arabidopsis. Plant Cell *21*, 2237–2252.

Kitakura, S., Vanneste, S., Robert, S., Löfke, C., Teichmann, T., Tanaka, H., and Friml, J.
(2011). Clathrin mediates endocytosis and polar distribution of PIN auxin transporters in
Arabidopsis. Plant Cell *23*, 1920–1931.

Lamesch, P., Berardini, T.Z., Li, D., Swarbreck, D., Wilks, C., Sasidharan, R., Muller, R.,
Dreher, K., Alexander, D.L., Garcia-Hernandez, M., et al. (2012). The Arabidopsis
Information Resource (TAIR): improved gene annotation and new tools. Nucleic Acids Res.
40, D1202-1210.

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat
Methods *9*, 357–359.

Larson, E.R., Domozych, D.S., and Tierney, M.L. (2014). SNARE VTI13 plays a unique role
in endosomal trafficking pathways associated with the vacuole and is essential for cell wall
organization and root hair growth in arabidopsis. Ann. Bot. *114*, 1147–1159.

Larson, E.R., Van Zelm, E., Roux, C., Marion-Poll, A., and Blatt, M.R. (2017). Clathrin
Heavy Chain Subunits Coordinate Endo- and Exocytic Traffic and Affect Stomatal
Movement. Plant Physiol. *175*, 708–720.

Lorenzo, O., Chico, J.M., Sánchez-Serrano, J.J., and Solano, R. (2004). JASMONATE-
INSENSITIVE1 encodes a MYC transcription factor essential to discriminate between
different jasmonate-regulated defense responses in Arabidopsis. Plant Cell *16*, 1938–1950.

Madeira, F., Park, Y.M., Lee, J., Buso, N., Gur, T., Madhusoodanan, N., Basutkar, P., Tivey,
A.R.N., Potter, S.C., Finn, R.D., et al. (2019). The EMBL-EBI search and sequence analysis
tools APIs in 2019. Nucleic Acids Res *47*, W636–W641.

Mbengue, M., Bourdais, G., Gervasi, F., Beck, M., Zhou, J., Spallek, T., Bartels, S., Boller,
T., Ueda, T., Kuhn, H., et al. (2016). Clathrin-dependent endocytosis is required for
immunity mediated by pattern recognition receptor kinases. Proc Natl Acad Sci U S A *113*,
11034–11039.

Naveed, M., Brown, L.K., Raffan, A.C., George, T.S., Bengough, A.G., Roose, T., Sinclair,
I., Koebernick, N., Cooper, L., Hackett, C.A., et al. (2017). Plant exudates may stabilize or
weaken soil depending on species, origin and time. European Journal of Soil Science *68*,
806–816.

Parker, J.S., Cavell, A.C., Dolan, L., Roberts, K., and Grierson, C.S. (2000). Genetic
interactions during root hair morphogenesis in Arabidopsis. Plant Cell *12*, 1961–1974.

Plessis, A., Cournol, R., Effroy, D., Silva Pérez, V., Botran, L., Kraepiel, Y., Frey, A., Sotta,
B., Cornic, G., Leung, J., et al. (2011). New ABA-hypersensitive Arabidopsis mutants are
affected in loci mediating responses to water deficit and Dickeya dadantii infection. PLoS
ONE *6*, e20243.

Prentice, R.L., and Kalbfleisch, J.D. (2003). Mixed discrete and continuous Cox regression
model. Lifetime Data Anal *9*, 195–210.

R Core Team (2014). R: A language and environment for statistical computing (Vienna,
Austria: R Foundation for Statistical Computing).

Ringli, C., Bigler, L., Kuhn, B.M., Leiber, R.-M., Diet, A., Santelia, D., Frey, B., Pollmann,
S., and Klein, M. (2008). The Modified Flavonol Glycosylation Profile in the Arabidopsis
rol1 Mutants Results in Alterations in Plant Growth and Cell Shape Formation. The Plant
Cell *20*, 1470–1481.

Robert, S., Chary, S.N., Drakakaki, G., Li, S., Yang, Z., Raikhel, N.V., and Hicks, G.R.
(2008). Endosidin1 defines a compartment involved in endocytosis of the brassinosteroid
receptor BRI1 and the auxin transporters PIN2 and AUX1. PNAS *105*, 8464–8469.

Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and
Mesirov, J.P. (2011). Integrative genomics viewer. Nat. Biotechnol. *29*, 24–26.

Rodriguez-Furlán, C., Salinas-Grenet, H., Sandoval, O., Recabarren, C., Arraño-Salinas, P.,
Soto-Alvear, S., Orellana, A., and Blanco-Herrera, F. (2016). The Root Hair Specific
SYP123 Regulates the Localization of Cell Wall Components and Contributes to
Rizhobacterial Priming of Induced Systemic Resistance. Front Plant Sci *7*, 1081.

Sasse, J., Martinoia, E., and Northen, T. (2018). Feed Your Friends: Do Plant Exudates Shape
the Root Microbiome? Trends Plant Sci. *23*, 25–41.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. Nat. Methods *9*, 676–682.

Stokes, A., Atger, C., Bengough, A.G., Fourcard, T., and Sidle, R.C. (2009). Desirable plant
root traits for protecting natural and engineered slopes against landslides. Plant and Soil *324*,
1–30.

Toukura, Y., Devee, E., and Hongo, A. (2006). Uprooting and shearing resistances in the
seedlings of four weedy species. Weed Biology and Management *6*, 35–43.

Vives-Peris, V., de Ollas, C., Gómez-Cadenas, A., and Pérez-Clemente, R.M. (2020). Root
exudates: from plant to rhizosphere and beyond. Plant Cell Rep. *39*, 3–17.

Waghmare, S., Lileikyte, E., Karnik, R., Goodman, J.K., Blatt, M.R., and Jones, A.M.E.
(2018). SNAREs SYP121 and SYP122 Mediate the Secretion of Distinct Cargo Subsets.
Plant Physiology *178*, 1679–1688.

Wang, X., Cnops, G., Vanderhaeghen, R., De Block, S., Van Montagu, M., and Van
Lijsebettens, M. (2001). AtCSLD3, A Cellulose Synthase-Like Gene Important for Root Hair
Growth in Arabidopsis. Plant Physiol *126*, 575–586.

Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis (Springer).

Yang, J., Bak, G., Burgin, T., Barnes, W.J., Mayes, H.B., Peña, M.J., Urbanowicz, B.R., and
Nielsen, E. (2020). Biochemical and Genetic Analysis Identify CSLD3 as a beta-1,4-Glucan
Synthase That Functions during Plant Cell Wall Synthesis. Plant Cell *32*, 1749–1767.

TABLES

Table 1 *Arabidopsis thaliana* mutant lines used in this study.

Mutant type	Mutant	AGI code[a]	Mutant line	Referenced in
Root hair shape	*cow1-3*	At4g34580	SALK_002124	Böhme *et al.* (2004)
	csld3-1	At3g03050	CS899	Wang *et al.* (2001)
	lrx1-4	At1g12040	SALK_057038	Baumberger *et al.* (2001)
	rol1-2	At1g78570	CS16373	Diet *et al.* (2006)
	gdi1-2	At3g07880	SALK_035400	Kang *et al.* (2017)
Vesicle trafficking	*chc1-2*	At3g11130	SALK_103252	Kitakura *et al.* (2011)
	chc2-3	At3g08530	SALK_151638	Mbengue *et al.* (2016)
	syp121	At3g11820	-	Collins *et al.* (2003)
	syp122-1	At3g52400	SALK_008617	Assaad *et al.* (2004)
Root exudate composition	*jin1-9*	At1g32640	SALK_017005	Anderson *et al.* (2004)
	pdr2	At4g15230	SAIL_811_F08	Badri *et al.* (2008)
	pft1-3	At1g25540	SALK_059316	Kidd *et al.* (2009)
Unknown	*abcg43-1*	At4g15236	N75206[b]	This study
	abcg43-2	At4g15236	SALK_201207	This study
	abcg43-3	At4g15236	SALKseq_30713	This study

Table 2 Output of the Cox PH regression models comparing the root-gel detachment of root hair shape, vesicle trafficking and root exudate composition mutants relative to wild type.

Mutant type	Mutant	Wald test (z-score) and corresponding P value		Hazard ratio (95% CI)
Root hair shape	cow1-3	$z = 7.350$ P < 0.001	***	3.790 (2.657, 5.407)
	csld3-1	$z = 8.644$ P < 0.001	***	4.636 (3.274, 6.564)
	gdi1-2	$z = 5.199$ P < 0.001	***	2.459 (1.752, 3.452)
	lrx1-4	$z = 8.049$ P < 0.001	***	3.926 (2.814, 5.478)
	rol1-2	$z = 7.927$ P < 0.001	***	4.036 (2.858, 5.698)
Vesicle trafficking	chc1-2	$z = 8.238$ P < 0.001	***	4.482 (3.137, 6.405)
	chc2-3	$z = 7.653$ P < 0.001	***	3.941 (2.774, 5.599)
	syp121	$z = 7.702$ P < 0.001	***	3.593 (2.595, 4.975)
	syp122-1	$z = 7.199$ P < 0.001	***	3.270 (2.369, 4.515)
Root exudate composition	jin1-9	$z = 0.737$ P > 0.05	ns	1.124 (0.824, 1.537)
	pdr2	$z = -5.765$ P < 0.001	***	0.379 (0.273, 0.528)
	pft1-3	$z = 3.315$ P < 0.001	***	1.698 (1.242, 2.323)
Unknown	abcg43-1	$z = -7.823$ P < 0.001	***	0.245 (0.172, 0.348)
	abcg43-2	$z = -6.049$ P < 0.001	***	0.355 (0.253, 0.496)
	abcg43-3	$z = -7.179$ P < 0.001	***	3.496 (0.203, 0.403)

*** refers to a statistically significant difference ≤ 0.001 and **ns** no statistically significant difference ≥ 0.05 relative to wild type, respectively.

FIGURE LEGENDS

FIGURE 1. Centrifuge assay workflow.

(A) i. Ten seeds (highlighted in pink) are sown onto the surface of sterile, solid gel growth medium in a single 90 mm Petri plate in two horizontal rows. ii. The plates are stacked in

groups of five and orientated vertically at approximately 80° to encourage the roots to grow

down the surface of the gel medium. Plates are grown in a growth chamber with constant light

(120-145 µmol m^{-2} s^{-1}) conditions at 22°C and 60% relative humidity.

(B) After 5-6 days, i. seedlings are visually analysed and numbered. ii. Petri plates are placed

into a swing-out-bucket centrifuge in an inverted orientation with their roots pointing inward

(indicated by the purple arrows). iii. After seedlings have been subjected to each one-minute

pulse of increasing centrifugal speed, seedling detachment is recorded.

(C) i. The aerial tissue mass of each seedling is determined using ii. an analytical scale to iii.

determine the root-gel adhesion properties of candidate lines are compared to wild-type to

assess whether they have increased (e.g. line y) or decreased (e.g. line x) adhesion to the sterile

gel.

FIGURE 2 Physical root hair properties contribute to root-substrate adhesion in *Arabidopsis*

thaliana.

(A) Root hair phenotypes of 5-d-old wild-type (Col-0), *lrx 1-4, csld3-1, cow1-3, rol1-2,* and

gdi1-2 seedlings grown on gel medium and statistical comparisons of mean root hair density

(number per mm length of root) and mean root hair length (mm) for each mutant line relative

to wild type. White asterisks on the root hair images indicate the characteristic root hair bulging

phenotype in the *lrx1-4* mutant and the short, branching root hair phenotype in *gdi1*-2. Scale

bar = 0.5 mm. In the table, the mean ± standard error is given for root hair density and root hair

length of wild type and each mutant as well as a mean comparison to wild type, which is listed

in bold. "No difference" is stated when there was no statistically significant difference between

wild type and a mutant line. The statistical output of each univariate linear model is given.

Alpha = 0.025, statistical significance codes: '***' = < 0.001, '**' = < 0.01. The results shown

are from a representative experiment for at least two independent experiments. Survival curves

displaying the proportion of seedlings that adhered to the gel at increasing centrifugal force for

(B) 92 wild type (Col-0 – black); 81 *lrx1-4* (light green); 70 *csld3-1* (green); **(C)** 82 wild type

(Col-0 - black); 83 *cow1-3* (green); and **(D)** 77 wild type (Col-0- black); 85 *rol1-2* (light green);

and 86 *gdi1-2* (dark green). Circled red crosses on the survival curves represent seedlings that

remained adhered to the gel after the maximum centrifugal speed (1611 RPM). Each graph

shows a representative experiment for at least two independent experiments showing a

statistically significant difference in adhesion between mutant lines relative to wild type (Cox

PH regression; alpha = 0.01). A single experiment included ≥ 70 biological replicates for each

candidate line.

FIGURE 3 Vesicle trafficking mechanisms that contribute to root-substrate adhesion in
Arabidopsis thaliana.
(A) Root hair phenotypes of 5-d-old wild-type (Col-0), *syp121, syp122-1, chc1-2* and *chc2-3*
seedlings grown on a gel medium and statistical comparisons of mean root hair density (number
per mm length of root) and mean root hair length (mm) for each mutant line relative to wild
type. Scale bar = 0.5 mm. In the table, the mean ± standard error is given for the root hair
density and root hair length of wild type and each mutant as well as a mean comparison to wild
type, which is listed in bold. "No difference" is stated when there was no statistically significant
difference between wild type and a mutant line. The statistical output of each univariate linear
model is given. Alpha = 0.025. The results shown are from a representative experiment for at
least two independent experiments. Survival curves displaying the proportion of seedlings that
adhered to the gel at increasing centrifugal force for **(B)** 87 wild type (Col-0 – black); 91 *syp121*
(turquoise); 83 *syp122-1* (blue/grey); and **(C)** 83 wild type (Col-0 – black); 70 *chc1-2* (dark
blue)*;* and 72 *chc2-3* (medium blue)*.* Red crosses circled on the survival curves represent
seedlings that remained adhered to the gel after the maximum centrifugal speed (1611 RPM).
Each graph shows a representative experiment for at least two independent experiments
showing a statistically significant difference in adhesion between all mutant lines relative to
wild type (Cox PH regression; alpha = 0.01). A single experiment included ≥ 70 biological
replicates for each candidate line.

FIGURE 4 Exudate composition changes root-substrate adhesion properties in *Arabidopsis*
thaliana.
(A) Root hair phenotypes of 5-d-old wild-type (Col-0), *jin1-9, pdr2* and *pft1-3* seedlings grown
on a gel medium and statistical comparisons of mean root hair density (number per mm length
of root) and mean root hair length (mm) for each mutant line relative to wild type. Scale bar =
0.5 mm. In the table, the mean ± standard error is given for the root hair density and root hair
length of wild type and each mutant as well as a mean comparison to wild type, which is listed
in bold. "No difference" is stated when there was no statistically significant difference between
wild type and a mutant line. The statistical output of each univariate linear model is given.
Alpha = 0.025. The results shown are from a representative experiment for at least two
independent experiments. **(B)** Survival curves displaying the proportion of seedlings that
adhered to the gel at increasing centrifugal force for 80 wild type (Col-0 – black), 81 *jin1-9*
(light pink), 80 *pdr2* (light orange), and 81 *pft1-3* (dark orange)*.* The results shown are from a

representative experiment for at least two independent experiments showing a statistically

significant difference in adhesion between mutant lines relative to wild type (Cox PH

regression; alpha = 0.01), except for *jin1-9*. A single experiment included $\geq$ 70 biological

replicates for each candidate line.

FIGURE 5 Using the centrifuge assay in a forward genetic screen to identify *Arabidopsis*

thaliana root-substrate adhesion mutants.

(A) T-DNA insert locations in *ABCG43* for each *abcg43* mutant allele, with insertions

indicated by orange arrowheads. The *ABCG43* gene is located on chromosome four at position

Chr4: 8,969,677-8,702,727 and contains 23 exons (dark green) and 22 introns (white gaps).

abcg43-1 has a T-DNA insert in the exon10/intron10 boundary, *abcg43-2* (SALK_201207)

has a T-DNA insert in exon 2 and *abcg43-3* (SALKseq_30713) has a T-DNA insert in exon 3.

(B) Genomic PCR confirming the homozygosity of each *abcg43* mutant for the respective T-

DNA inserts. Lanes **1** and **2** were loaded with the gene-specific and T-DNA-border PCR

products from *abcg43* mutant lines. Lanes **3** and **4** were loaded with the gene-specific and T-

DNA-border PCR products from a wild type (Col-0) genomic DNA template. Lanes **5** and **6**

were loaded with the water controls for the gene-specific and T-DNA-border PCR reactions,

respectively. '**L**' indicates the 100-bp ladder. The expected product sizes for the gene-specific

PCRs were ~350 nt, ~1025 nt and ~1090 nt; the T-DNA-border PCR product sizes were ~400

nt, ~550 nt and ~750 nt for *abcg43-1, abcg43-2* and *abcg43-3* alleles, respectively. **(C)**

Survival curves displaying the proportion of seedlings that adhered to the gel at increasing

centrifugal force for 79 wild type (Col-0 – black), 70 *abcg43-1* (dark pink), 85 *abcg43-2*

(purple), and 73 *abcg43-3* (light purple). Red crosses circled on the survival curves represent

seedlings that remained adhered to the gel after the maximum centrifugal speed (1611 RPM).

The results are from a representative experiment for at least two independent experiments

showing a statistically significant difference in adhesion between the mutant lines relative to

wild type (Cox PH regression; alpha = 0.01). A single experiment included $\geq$ 70 biological

replicates for each candidate line.

A

Col-0   lrx1-4   csld3-1   cow1-3   rol1-2   gdi1-2

Line	Root hair density (number per mm length of root)		Root hair length (mm)	
Col-0	30.556 ± 0.784	-	0.518 ± 0.009	-
lrx1-4	31.889 ± 0.935 **No difference**	t = 1.093, P > 0.05 d.f. = 16, n = 9	0.388 ± 0.008 **1.3 times shorter****	t = -2.819, P < 0.001 d.f. = 16, n = 9
csld3-1	8.778 ± 1.188 **3.5 times less****	t = -15.30, P < 0.001 d.f. = 16, n = 9	0.045 ± 0.002 **10.1 times shorter*****	t = -11.790, P < 0.001 d.f. = 16, n = 9
cow1-3	29.200 ± 0.772 **No difference**	t = -1.23, P > 0.05 d.f. = 17, n = 10	0.049 ± 0.001 **10 times shorter*****	t = -12.37, P < 0.001 d.f. = 17, n = 10
rol1-2	30.700 ± 1.550 **No difference**	t = 0.08, P > 0.05 d.f. = 17, n = 9	0.195 ± 0.003 **2.7 times shorter*****	t = -8.417, P < 0.001 d.f. = 17, n = 10
gdi1-2	26.200 ± 1.041 **0.17 times less****	t = -3.282, P < 0.01 d.f. = 17, n = 9	0.135 ± 0.006 **3.8 times shorter*****	t = -9.04, P < 0.001 d.f. = 16, n = 9

B

Percentage of seedlings adhered to the gel (%) vs Centrifugal force (mN)

— Col-0
— lrx1-4
— csld3-1

C

Percentage of seedlings adhered to the gel (%) vs Centrifugal force (mN)

— Col-0
— cow1-3

D

Percentage of seedlings adhered to the gel (%) vs Centrifugal force (mN)

— Col-0
— rol1-2
— gdi1-2

Line	Root hair density (number per mm length of root)		Root hair length (mm)	
Col-0	32.000 ± 0.632	-	0.420 ± 0.005	-
syp121	31.200 ± 0.952 **No difference**	t = -0.700, P > 0.05 d.f. = 18, *n* = 10	0.461 ± 0.004 **No difference**	t = 1.974, P > 0.05 d.f. = 18, *n* = 10
syp122-1	32.200 ± 0.442 **No difference**	t = 0.259, P > 0.05 d.f. = 18, *n* = 10	0.452 ± 0.005 **No difference**	t = 1.379, P > 0.05 d.f. = 18, *n* = 10
chc1-2	30.889 ± 0.716 **No difference**	t = -1.168, P > 0.05 d.f. = 17, *n* = 9	0.441 ± 0.006 **No difference**	t = 0.684, P > 0.05 d.f. = 17, *n* = 9
chc2-3	31.889 ± 0.790 **No difference**	t = -0.111, P > 0.05 d.f. = 17, *n* = 9	0.463 ± 0.006 **No difference**	t = 0.182, P > 0.05 d.f. = 18, *n* = 10

A

Line	Root hair density (number per mm length of root)		Root hair length (mm)	
Col-0	30.111 ± 1.612	-	0.416 ± 0.005	-
jin1-9	29.444 ± 1.271 **No difference**	t = -0.324, P > 0.05 d.f. = 16, *n* = 9	0.429 ± 0.006 **No difference**	t = 0.555, P > 0.05 d.f. = 16, *n* = 9
pdr2	31.125 ± 0.854 **No difference**	t = 0.533, P > 0.05 d.f. = 15, *n* = 8	0.428 ± 0.006 **No difference**	t = 0.521, P > 0.05 d.f. = 15, *n* = 8
pft1-3	29.000 ± 1.179 **No difference**	t = -0.555, P > 0.05 d.f. = 16, *n* = 9	0.424 ± 0.006 **No difference**	t = 0.295, P > 0.05 d.f. = 16, *n* = 9

B