A centrifuge-based method for identifying novel genetic traits that affect root-substrate adhesion in *Arabidopsis thaliana*

Bethany M Eldridge[1,*], Emily R Larson[1,*], Laura Weldon[1,a], Kevin M Smyth[1], Annabelle N Sellin[1], Isaac V Chenchiah[2], Tanniemola B Liverpool[2] and Claire S Grierson[1,#]

[1]School of Biological Sciences, University of Bristol, Bristol Life Sciences Building, 24 Tyndall Avenue, Bristol, BS8 1TQ, UK

[2]School of Mathematics, University of Bristol, Fry Building, Woodland Road, Bristol BS8 1UG, UK

[a] Current address: Wildfowl and Wetlands Trust (WWT), Slimbridge, Gloucester, GL2 7BT, UK

[*]Authors contributed equally to this work.

[#]Corresponding author: Claire Grierson

lacsg@bristol.ac.uk

ORCID:

BME: ORCID: 0000-0002-6598-3701

ERL: ORCID: 0000-0002-5498-8152

LW: ORCID: 0000-0002-0761-4414

KMS: ORCID: 0000-0003-0105-8557

IVC: ORCID: 0000-0002-8618-620X

TBL: ORCID: 0000-0003-4376-5604

CSG: ORCID: 0000-0002-4000-0975

Keywords: Arabidopsis, Centrifuge assay, Forward genetics, Root exudates, Root hairs, Root-soil adhesion, Survival analysis, Vesicle trafficking

ABSTRACT

The physical presence of roots and the compounds they release affect the cohesion between roots and their environment. However, the plant traits that are important for these interactions are unknown and most methods that quantify the contributions of these traits are time-intensive and require specialist equipment and complex substrates. Our lab developed an inexpensive, high-throughput phenotyping assay that quantifies root-substrate adhesion in *Arabidopsis thaliana*. We now report that this method has high sensitivity and versatility for identifying different types of traits affecting root-substrate adhesion including root hair morphology, vesicle trafficking pathways and root exudate composition. We describe a practical protocol for conducting this assay and introduce its use in a forward genetic screen to identify novel genes affecting root-substrate interactions. This assay is a powerful tool for identifying and quantifying genetic contributions to cohesion between roots and their environment.

INTRODUCTION

Plants secrete compounds that help them adapt to their environment, sense and interact with other organisms, and improve water and nutrient uptake (Bailey et al., 2002; Datta et al., 2011; Haling et al., 2013). These compounds (exudates) can vary in composition based on plant species, developmental stage, and environment (Naveed et al., 2017). Recent studies have shown that some compounds have bioadhesive properties that stick plants and soils together and aggregate soil particles, modifying the microenvironment around plant roots (Galloway et al., 2018). However, understanding how plant biology and physiology contribute to plant-soil interactions can be confounded by multiple factors: other organisms living in the soil also secrete organic metabolites that can contribute to soil cohesion, soil composition might affect plant exudate composition and vice versa. Also, plant root morphology including root architecture and the presence of root hairs can alter root-soil interactions (Bailey et al., 2002; Burylo et al., 2012; De Baets et al., 2007; Ghestem et al., 2014; Stokes et al., 2009). This complexity has made it difficult to characterise the relevant plant physiology.

We developed a high-throughput assay that identifies and quantifies the contribution of root hair cells to root-substrate cohesion in a less variable environment than the natural rhizosphere (De Baets et al., 2020). The assay uses a centrifuge to apply force to *Arabidopsis thaliana* seedlings grown on the surface of a sterile medium and measures the force required

to detach seedlings from that surface and applies a Cox proportional hazards regression

(Prentice and Kalbfleisch, 2003) to quantify the adherence of candidate Arabidopsis lines

relative to wild-type plants at increasing centrifugal force. Because previous use of this assay

only assessed whether presence or absence of root hairs affected adhesion to the agar (De

Baets *et al.,* 2020) and did not address subtle aspects of root hair biology, we were interested

in whether this assay is sensitive enough to identify other types of traits involved in root-

substrate adhesion. To test this extended use of the assay, we asked if it could detect the

adhesive effects in (1) mutants with altered root hair shape; (2) vesicle trafficking mutants

with no known root hair morphological defects; (3) exudate compositional mutants with no

known root hair phenotypes; and (4) a forward genetic screen of a mutant population to

identify novel genes that affect root-substrate adhesion.

Here, we describe how this method has the sensitivity to identify how differences in

root hair morphology and biological functions that do not present a visible root hair

phenotype contribute to substrate cohesion. Using the genetic and molecular tools available

in Arabidopsis and working in a sterile environment allows hypotheses about plant specific

factors of root-substrate interactions to be tested. While plant-soil cohesion is a complex and

dynamic interaction, our assay provides a way to efficiently probe root cellular functions that

might be masked by confounding variables in a soil-based study system. Results from this

assay provide a platform for more comprehensive studies of plant morphology and

physiology that affect root-substrate cohesion.

MATERIALS AND METHODS

Plant Material and Growth Conditions

Candidate and wild-type lines of the Columbia-0 (Col-0) ecotype of *Arabidopsis*

thaliana single-gene mutants were obtained from the Nottingham Arabidopsis Stock Centre

(NASC; Nottingham, UK) or were stocks maintained in the lab (**Table 1**). Homozygous lines

were used in all experiments.

For all experiments, seeds were surface sterilised for 15 min in 20% (v/v) bleach and

sterile water solution, followed by five washes in sterile water. All sterilised seeds were

stratified in the dark at 4°C for 24-48 h in Eppendorf microtubes containing sterile water. All

plants were reared in a growth room at 21-22°C with a continuous photoperiod (light

intensity = 120-145 μmol m^{-2} s^{-1}) at 60% humidity.

Half strength Murashige and Skoog basal medium (Sigma M5519) was supplemented

with 1% [w/v] sucrose pH adjusted to 5.7, solidified with 1% [w/v] agar (Sigma A1296).

Sterilized seeds were sown onto 90-mm Petri plates (Thermo Scientific RC2260) containing

30 ml sterile medium and sealed with Parafilm (Bemis, NA). Plates were then placed in

stacks of five at ~80° from the horizontal (lid side up) to encourage vertical growth along the

gel surface. That plates were incubated in the growth chamber for 5-6 days before

centrifugation Plants for genomic DNA extractions were grown in compost (three parts

Levington F3 compost and one-part J Arthur Bowers horticultural silver sand).

Centrifuge Assay Set-Up

To measure the adhesion of Arabidopsis seedling roots using a centrifuge, seedling

roots must not penetrate the gel or touch each other, as both events could interfere with

seedling detachment during centrifugation. Therefore, ten seeds were sown 1-cm apart on

each plate in two parallel rows (**Figure 1Ai**). The plates were then grown for 5-6 days

(**Figure 1Aii**) and seedlings that were touching each other, had grown into the gel, or not

uniform in size were omitted from the assay. The seedlings were then numbered so that plate

positional effects could be included as a potential covariate in the final analysis (**Figure 1Bi**).

Plates were inverted and placed into a swing-out bucket centrifuge (Beckman Allegra X-30R

Centrifuge) that held four plates at a time and subjected to increments of increasing

centrifugal force at 720, 1018, 1247, 1440 and 1611 RPM for one minute each (**Figure 1Bii**).

The proportion of seedlings that detached from the gel surface was recorded between each

centrifugal speed; a detachment event was recorded when a seedling had partially or fully

peeled away from the gel (**Figure 1Biii**). If the gel shattered during the assay, seedlings that

had not detached from the gel were not recorded and seedlings that remained adhered to the

gel after the maximum centrifugal speed were censored in the analysis. During method

optimisation we encountered some common methodological challenges for which we provide

troubleshooting options in **Table S1**.

Calculating Seedling Detachment Force

To calculate the centrifugal force acting on a seedling, the aerial tissue mass for each

seedling was measured using an analytical balance (Microbalance ME5; Sartorius; **Figure**

1Ci, ii). The aerial tissue weight of each seedling was measured because a plant with heavier

aerial tissue experiences more force than a lighter one subjected to the same centrifugal

speed. Centrifugal force, Fc (mN) acting on a seedling was calculated as previously reported

in De Baets et al. (2020). Aerial tissue weight (kg), the angular velocity (ω) and the distance

between the seedling and the axis of rotation on the centrifuge (radius = 0.07m) were used to

give the following equation:

Fc= *mass* x *radius* x ω^2 (Equation 1)

Identification of *abcg43-1* Mutant T-DNA Insert

Genomic DNA was extracted from a pool of vegetative tissue taken from ten three-

week old plants using an adapted protocol (Healey et al., 2014) for Illumina Next-Generation

two x paired-end sequencing conducted at the Bristol Genomics Facility (University of

Bristol, UK). The Illumina TruSeq Nano LT gDNA kit (Illumina Inc) was used to generate a

genomic DNA sequencing library following the manufacturer's protocol. The final library

was diluted to a loading concentration of 1.4 pM for cluster generation and 2 x 150 bp paired-

end sequencing on the Illumina NextSeq500 system (Illumina Inc) alongside a 5% PhiX

spike-in control library. Read summary statistics were generated using the RTA 2.4.6

Primary Analysis Software (Illumina Inc). The read summaries were analysed in the

Sequencing Analysis Viewer (Illumina Inc). Filtered paired reads were subject to paired-end

alignment using the Bowtie2 2.3.4.2 aligner (Langmead and Salzberg, 2012). A bespoke

reference genome was produced that combined the TAIR10 Arabidopsis genome (Lamesch et

al., 2012) and the pROK2 vector sequence (Baulcombe et al., 1986). Alignments were

viewed in the Integrative Genomics Viewer IGV 2.3 (Robinson et al., 2011).

For Sanger sequencing, DNA was extracted from a pool of vegetative tissue taken

from two-week-old plants using a modified Edwards prep (Edwards et al., 1991). High

fidelity PCR was conducted on the *abcg43-1* line using the Q5 High-Fidelity 2X Master Mix

(NEB). PCR products were purified and extracted from a 1% agarose gel using the QIAquick

Gel Extraction Kit (QIAGEN), following the manufacturer's instructions. Purified PCR

product plus the *abcg43-1* forward, reverse or border genotyping primer (**Table S2**) was used

for Sanger sequencing (using the Mix2Seq overnight sequencing kit, Eurofins Genomics).

Chromatograms and FASTA files were obtained from Eurofin Genomics and following

manual low-quality end trimming, the final sequences were aligned to the pROK2 vector

sequence and *ABCG43* gene sequence using the MUSCLE alignment tool (Edgar, 2004).

Genotyping

To genotype the *abcg43* mutant alleles, genomic DNA was extracted from the vegetative

tissue of two-week-old plants using a modified Edwards prep (Edwards et al., 1991). T-DNA

border and gene-specific primers were used in PCR analyses to confirm the genotype (**Table**
S2).

172
173
174

Morphological Root Trait Analyses

Five-day-old Arabidopsis seedlings were imaged with a Leica MZ FLIII microscope

(Leica) with dark-field lighting. Root hair length and density were measured using

microscope images of five-day-old Arabidopsis seedlings and Fiji version 1.0 (Schindelin et

al., 2012) and the Bio-Formats Importer plugin. At least two experimental repeats were

conducted; in each experiment we imaged eight to ten individual plants from each line and

measured at least 30 root hairs per plant.

Statistical Analyses

All statistical analyses were conducted using RStudio, version 1.1453 (R Core Team,

2014) and all graphs were generated using the R package ggplot2 (Wickham, 2016).

A Cox proportional hazards (PH) regression model was used to statistically test for

differences between the detachment of experimental seedlings relative to the wild-type

control (Col-0) at increasing force (**Figure 1Ciii**). Cox PH regression models were conducted

using the coxph function within the survival package in R and in all cases, the assumption of

proportionality was satisfied. Comparisons of different candidate lines relative to the wild

type line were analysed *a priori*; therefore, a series of contrasts were set up using the R

function contr.treatment rather than using post-hoc testing methods. Seedling position and

individual plate number were incorporated as covariates to account for any heterogeneity;

when these covariates had no significant effect, they were removed from the model. We

censored seedlings that remained attached to the gel after the maximum centrifugal speed

because we did not determine the speed these seedlings would have detached from the gel.

For each Cox PH regression model run, we report the P value of the Wald Statistic (z-score)

and the hazard ratio with the upper and lower bound confidence intervals. An alpha level of

0.01 was used. The root-gel detachment of at least 70 plants from each line was assessed

within a single experiment.

Univariate analyses were conducted using the lm function in R to test for differences

in the mean root hair density or mean root hair length (of eight to ten individual plants from

each line) between candidate lines relative to wild type. In all cases, the assumptions of

normality and homoscedasticity were satisfied. Because comparisons of different mutant
lines relative to the wild type line were *a priori*, a series of contrasts were conducted using
the contr.treatment function in R. To prevent type I errors from multiple testing, we applied
the Bonferroni method and adjusted the alpha level to 0.025 (0.05/2).

RESULTS
Root Hair Morphological Effects on Root-Substrate Adhesion
211   We screened mutant lines with known root hair morphology defects to ask if the
centrifuge assay could determine if root hair shape affects root-substrate adhesion. The
mostly bald mutant *csld3*, which encodes the CESA-like 3D protein (Favery et al., 2001;
Yang et al., 2020) was included as a root hairless example. For lines that produce short or
malformed root hairs, we included the *gdi1-2* Rho GTPase GDP dissociation inhibitor
mutant, *rol1-2* rhamnose biosynthesis 1 mutant allele, *can of worms 1* (*cow1-3*)
phosphatidylinositol transferase protein mutant, and *lrx1-4* leucine rich extensin 1 mutant
(Baumberger et al., 2001; Böhme et al., 2004; Carol et al., 2005; Diet et al., 2006; Grierson et
al., 1997). The *rol1-2* and *gdi1-2* mutants have short root hairs and the root hairs on *gdi1-2*
plants can also branch (Parker et al., 2000; Ringli et al., 2008). *cow1-3* plants have short,
wide root hairs that can branch, whilst *lrx1-4* plants can develop root hairs that abort, swell,
branch, or collapse in a growth condition-dependent manner (Baumberger et al., 2001;
Grierson et al., 1997; Parker et al., 2000). Under our growth conditions, we confirmed the
reported phenotypes for each mutant line (**Figure 2A**). Wild type was used as the baseline
group for all centrifuge experiments, with a hazard ratio of one. Therefore, a candidate line
with a higher risk of detachment from the gel than wild type will have a hazard ratio above
one, while lines with a lower risk of detachment than wild type will have a hazard ratio below
one.
229   Consistent with previous results (De Baets et al., 2020), bald (*csld3*) seedlings
detached at lower centrifugal forces relative to wild-type seedlings, with 4.6 times the risk of
detachment than wild-type plants ($z = 8.644$, P<0.001, HR = 4.636, 95% CI = 3.274 – 6.564;
Figure 2B). Mutants with deformed root hair phenotypes including *gdi1-2, rol1-2, cow1-3*
and *lrx1-4* had risks of detachment that were 2.5, 4, 3.8 and 3.9 times that of wild-type plants,
respectively (*gdi1-2* – $z = 5.199$, P<0.001, HR = 2.459, 95% CI = 1.752 – 3.452; *rol1-2* – $z =$
7.927, P<0.001, HR = 4.036, 95% CI = 2.858 – 5.698; *cow1-3* – $z = 7.350$, P<0.001, HR =
3.790, 95% CI = 2.657 – 5.407; *lrx1-4* – $z = 8.049$, P<0.001, HR = 3.926, 95% CI = 2.814 –
5.478; **Figure 2B-D**). These results indicated that the centrifuge assay can quantify the

effects of root hair shape on root-substrate cohesion. Many of the root hair mutants tested are

associated with changes in root hair cell wall composition, suggesting that alterations in

secreted compounds could affect root-substrate interactions in addition to any physical or

surface area-related factors that are specific to root hairs.

Vesicle Trafficking Mutations Affect Root-Substrate Adhesion

245   We hypothesized that Arabidopsis mutants with defective trafficking pathways

might alter cell wall and apoplast characteristics that contribute to root-substrate interactions

and adhesion. We chose the secretory Soluble NSF (N-ethylmaleimide sensitive fusion

protein) Attachment proteins (SNAP) REceptor (SNARE) mutants *syp121* and *syp122-1*, and

the endocytic mutants, *chc1-2* and *chc2-3* as candidates for our assay. SYP121 is the primary

secretory SNARE found on the plasma membrane (Assaad et al., 2004; Geelen et al., 2002)

and has characterized aboveground phenotypes, including small rosettes and stomatal

mobility defects (Eisenach et al., 2012; Larson et al., 2017); however, the mutant does not

have a reported root hair phenotype. Although the related SNARE SYP122 is thought to

share partial functionality with SYP121, the *syp122-1* mutant does not share these

phenotypes with *syp121* and a recent proteomic analysis reported differences in their vesicle

cargoes, which could contribute to their functional independence (Waghmare et al., 2018).

The *chc1-2* and *chc2-3* mutants are defective in the heavy chain subunits of the clathrin coat

complex, which is required for vesicle traffic at the plasma membrane (Kitakura et al., 2011).

Endo- and exocytic rates are impaired in both mutants (Larson et al., 2017), but no root hair

phenotypes have been reported. For the centrifuge assay, it was important to select mutants

that do not have root hair phenotypes given the effect root hair morphology has on root-gel

adhesion. Therefore, we evaluated the root hairs of these trafficking mutants and found that

the root hairs of these lines did not significantly differ from wild-type plants (**Figure 3A**).

264   Overall, the risk of detachment for *syp121, syp122-1, chc1-2* and *chc2-3* plants was

3.6, 3.3, 4.5 and 3.9 times that of wild type, respectively (*syp121* – $z = 7.702$, P<0.001, HR =

3.593, 95% CI = 2.595 – 4975; *syp122-1* – $z = 7.199$, P<0.001, HR = 3.270, 95% CI = 2.369

– 4.515; *chc1-2* – $z = 8.238$, P<0.001, HR = 4.482, 95% CI = 3.137 – 6.405; *chc2-3* – $z =$

7.653, P<0.001, HR = 3.941, 95% CI = 2.774 – 5.599; **Figure 3B,C**). Given that all the

mutants detached from the gel at lower centrifugal forces than wild-type seedlings, these

results indicated that the corresponding genes contribute to root-substrate adhesion and that

the centrifuge assay can be used to probe for intracellular molecular machinery that affect

root-substrate interactions.

Exudate Composition Changes Root Detachment Rates

Over 20% of plant assimilated carbon is released as root exudates (Huang et al., 2016)

that modify the external environment in response to abiotic and biotic stimuli and contribute

to soil adhesion (Akhtar et al., 2018; Galloway et al., 2018). Root epidermal and hair cells

secrete compounds that contribute to root exudate profiles that can be plant species specific

(Naveed et al., 2017). Because mutants in both root hair development and vesicle trafficking

pathways showed root-substrate adhesion phenotypes (**Figure 2 and 3**), we asked if the

centrifuge assay could identify effects of exudate composition on root-substrate adhesion. We

selected Arabidopsis mutants reported to have altered exudate composition compared to wild

type, including *pft1-3*, *jin1-9*, and *pdr2* (Badri et al., 2009; Berger et al., 1996; Carvalhais et

al., 2015; Kidd et al., 2009). *PFT1* (*MED25*) encodes the MEDIATOR25 subunit of the

Mediator nuclear protein, and *JIN1* (*MYC2*) encodes a basic helix-loop-helix leucine zipper

transcription factor; both proteins are involved in the jasmonate signalling pathway (Kidd et

al., 2009; Lorenzo et al., 2004). *pft1-3* and *jin1-9* mutants are reported to have altered root

exudate composition, including lower amounts of the amino acids asparagine, ornithine and

tryptophan than wild-type plants (Carvalhais et al., 2015). *PDR2* (*ABCG30*) encodes a

pleiotropic drug resistance (PDR) full-length ABC transporter that is involved in ABA

transport and the exudation of secondary metabolites (Badri et al., 2008; Kang et al., 2015).

We did not observe root hair growth phenotypes in these mutant lines compared to wild type

(**Figure 4A**).

Compared to wild-type seedlings, *pdr2* seedlings resisted detachment, with 0.38 times

the risk of detaching from the gel (z = -5.765, P<0.001, HR = 0.379, 95% CI = 0.273 – 0.528;

Figure 4B). Conversely, *pft1-3* seedlings had an increased risk of detachment 1.7 times that

of wild-type plants (z = 3.315, P<0.001, HR = 1.698, 95% CI = 1.242 – 2.323), and there was

no difference in gel adhesion between *jin1-9* and wild-type plants (z = 0.737, P>0.05, HR =

1.124, 95% CI = 0.824 – 1.537; **Figure 4B**). These results show that exudate composition can

alter root-substrate interactions, although this may not be the case for all exudate

compositional changes, or the effects of the exudate composition of the *jin1-9* mutant lay

outside the sensitivity range of the assay. Since these mutant lines did not have root hair

phenotypes, our results suggest that changes in exudate composition alter root adhesive

properties without affecting plant root hair morphology, further indicating the power of the
centrifuge assay as an efficient mutant screening method.

Centrifuge Assay as a Method for Forward Genetic Screening

Having demonstrated that our centrifugation technique can identify different types of
traits that affect root-substrate adhesion, we used it in a forward genetic screen to find novel
genes involved in root-substrate adhesion. To conduct this screen, the root-gel adhesion
properties of individual plants from a pooled SALK T-DNA insertion mutant collection were
analysed (Alonso et al., 2003). Individual plants that detached at the lowest centrifugal speed
or remained attached to the gel after being spun at the highest centrifugal speed were
recovered from the screen and self-fertilized to obtain progeny and identify the putative
mutations. The centrifuge assay was used to rescreen the progeny (using a sample size of $\geq$
70 n) to confirm whether candidate lines had significantly increased or decreased root-gel
adhesion relative to wild-type plants. As an example, we identified a line with significantly
enhanced root-gel adhesion and conducted genomic Next-Generation sequencing followed by
Sanger sequencing of T-DNA flanking PCR products to confirm a T-DNA insertion in the
ABC transporter gene, *ABCG43,* whose function is unknown. We named this T-DNA
insertion line *abcg43-1* and identified two additional mutant alleles, *abcg43-2*
(SALK_201207c) and *abcg43-3* (SALKseq_30713) (**Figure 5A**).

We analysed the root-gel adhesion of the homozygous insertional *ABCG43* mutant
alleles and found *abcg43-1*, *abcg43-2* and *abcg43-3* had detachment risks 0.25, 0.35 and 0.2
times that of wild-type plants, respectively (*abcg43-1* – z = -7.823, P<0.001, HR = 0.245,
95% CI = 0.172 – 0.348; *abcg43-2* – z = -6.049, P<0.001, HR = 0.355, 95% CI = 0.253 –
0.496; *abcg43-3* – z = -7.179, P<0.001, HR = 0.203 – 0.403; **Figure 5B, C**). Collectively,
these results suggested that disruption of *ABCG43* expression affects root-gel adhesion.
These results also illustrate the use of the centrifuge assay as a powerful screening method for
identifying novel genes involved in root-substrate adhesion, as well as a tool for evaluating
the effects of known gene function on root-substrate interactions. Experiments are now being
conducted to identify mutant trait(s) in these lines and understand how *ABCG43* affects root-
substrate adhesion.

DISCUSSION

To date, most published methods that quantify the contribution of particular traits to
plant-soil interactions are time-intensive and expensive (Bailey et al., 2002; De Baets and

Poesen, 2010; De Baets et al., 2020; Lin and Ying, 1994; Toukura et al., 2006). In contrast,
our assay can produce data within a week, is available to any laboratory with access to a
bench top centrifuge and does not require speciality consumables. This method is a high-
throughput and quantitative way to test the effects of root morphology and cell function on
root-substrate adhesion and to screen for new biological and molecular factors that can alter
plant-substrate interactions. Using a defined model system like Arabidopsis and sterile
medium conditions permits the identification of plant-specific characteristics that can then be
probed in other substrate- or soil-based conditions.

346
347

Using the Centrifuge Assay to Investigate the Role of Root Hairs in Root-Substrate
Interactions

Results presented in this paper show that the centrifuge assay can quantify effects of
root hair morphology on plant-substrate adhesion. Previous work has shown that the presence
of root hairs can significantly enhance root-substrate adhesion compared to the absence of
root hairs (Choi and Cho, 2019; De Baets et al., 2020; Haling et al., 2013). We now show that
our centrifuge assay can distinguish the adherence strength of seedlings with altered root hair
shapes and sizes (**Figure 2**) and quantify the contribution these morphological traits make to
plant-substrate interactions. Some mutants with defective root hairs have mutations in cell
wall biosynthesis and modification proteins, so it is possible that altered root-substrate
interface compromises adhesion in these mutants, not just root hair shape. Using our method
to rapidly phenotype mutants with root hair growth defects could direct additional
experiments to characterize biochemical and morphological properties of genes of interest.
Indeed, current evidence for the importance of root hairs for the deposition of complex
polysaccharides with soil-binding properties (Galloway et al., 2020) supports the need for
assays such as ours to expedite the identification of genes and mechanisms that improve soil-
remediation, top soil maintenance, and erosion tolerance of at-risk land.

Identifying New Genes Involved in Root-Substrate Adhesion
Mutations in cellular functions that do not alter root hair growth but do affect
substrate adhesion can also be investigated using this assay. Vesicle trafficking pathways are
required for cell wall deposition and maintenance (De Caroli et al., 2011; Larson et al., 2014;
Rodriguez-Furlán et al., 2016); therefore, we hypothesized that mutants in trafficking
pathways, particularly those at the plasma membrane, could affect root-substrate adhesion.

Similarly, the composition and deposition of plant exudates help plants modify and optimize
their growth conditions (Carvalhais et al., 2015; Naveed et al., 2017), suggesting that
modification to plant exudate composition could affect root-substrate interactions. Consistent
with these hypotheses, mutants with vesicle trafficking defects and altered exudate
composition had root-substrate adhesion properties that differed from wild type, indicating
that this method can identify aspects of plant biology important for root-substrate interactions
that do not affect root or root hair morphology (**Figure 3 and 4**). Our results further indicate
that while root hairs provide significant adhesive and cohesive support both on solid gel and
soil media (De Baets et al., 2020), there are additional cellular factors contributing to plant-
substrate interactions that are not directly observable without the use of expensive reagents,
development of time-intensive transgenic lines, and microscopy equipment. Our centrifuge
assay provides an affordable and quick alternative to these current methodologies.

Forward genetic screening relies on distinctive phenotypes that distinguish differences
within large, genetically diverse populations. High throughput, creative assaying techniques
to identify phenotypes within morphologically similar populations, such as thermal imaging
(Plessis et al., 2011) and confocal imaging platforms (Drakakaki et al., 2011; Robert et al.,
2008) require specific equipment that not all laboratories have at their disposal. In contrast,
the centrifuge assay can be used as a screen to identify new genes affecting plant-substrate
interactions without specialist equipment, making it an accessible option for research groups
that may not otherwise be able to efficiently conduct genetic screens. Moreover, the
centrifuge assay does not require morphological or conditional differences for candidate
selection. As an example, we report the identification in a forward genetics screen of a
mutant population in which *abcg43* mutant seedlings were identified and shown to be more
adhesive to gel than wild type (**Figure 5**). The function of ABCG43 in Arabidopsis is
uncharacterised, indicating that this assay is successful at identifying novel candidate genes
that have no previously known association with root morphological defects, adhesion or
exudate composition for future study.

CONCLUSION
This assay has applications in plant cell biology and genetics to investigate how gene
activity, cell function and root structure affect plant-soil adhesion, and identify the molecular
pathways involved. Potential applications not tested in this report include the measurement of
the effects environmental conditions have on plant-substrate adhesion and how other plant

species or ecotypes respond to detachment forces. Given the recent discovery that root hair
presence is important for Arabidopsis cohesion between roots and gel, and compost or soil
(De Baets *et al.,* 2020), this assay can broadly contribute to plant and soil sciences through
the identification of plant traits that increase root-substrate cohesion and potentially stabilise
slopes or reduce soil erosion.
410

411 **SUPPLEMENTARY INFORMATION**
412 **Table S1** Centrifuge assay troubleshooting
413 **Table S2** Primers used in this study for genotyping *Arabidopsis thaliana* mutants
414
415

ACKNOWLEDGEMENTS

This work was supported by a Leverhulme Trust project grant RPG-2013-260 to CSG and
TBL. BME was supported by a BBSRC SWBio PhD studentship (grant BB/M009122/1). We
are thankful to Don Grierson FRS and Enrico Coen FRS who independently suggested the
centrifuge assay concept, Timothy Quine for providing insightful guidance during method
development, and Thomas Denbigh for providing technical advice. We are grateful to Jill
Harrison, Victoria Spencer, Zoe Nemec Venza, Sophie Carpenter and Nicholas H Fair for
their helpful comments during revision of the manuscript.

AUTHOR CONTRIBUTIONS
CSG designed the initial project, conducted proof-of-concept pilot experiments, obtained
funding, and recruited and advise the team. TBL and IC developed the theoretical parameters
for the centrifugal assay and the bounds on the errors from other mechanical forces. BME
and ERL planned and conducted all experiments apart from the initial genetic screen, which
was carried out by LW and KMS. BME and ERL analysed all of the data presented herein.
AS and BME optimised the centrifuge assay data collection and data analysis methods. BME
and ERL wrote the initial manuscript. BME, ERL, and CSG edited the manuscript. All
authors reviewed and approved the final manuscript.
434

REFERENCES

Akhtar, J., Galloway, A.F., Nikolopoulos, G., Field, K.J., and Knox, P. (2018). A quantitative
method for the high throughput screening for the soil adhesion properties of plant and
microbial polysaccharides and exudates. Plant Soil *428*, 57–65.

Alonso, J.M., Stepanova, A.N., Leisse, T.J., Kim, C.J., Chen, H., Shinn, P., Stevenson, D.K.,
Zimmerman, J., Barajas, P., Cheuk, R., et al. (2003). Genome-wide insertional mutagenesis
of Arabidopsis thaliana. Science *301*, 653–657.

Assaad, F.F., Qiu, J.-L., Youngs, H., Ehrhardt, D., Zimmerli, L., Kalde, M., Wanner, G.,
Peck, S.C., Edwards, H., Ramonell, K., et al. (2004). The PEN1 Syntaxin Defines a Novel
Cellular Compartment upon Fungal Attack and Is Required for the Timely Assembly of
Papillae. Mol. Biol. Cell *15*, 5118–5129.

Badri, D.V., Loyola-Vargas, V.M., Broeckling, C.D., De-la-Peña, C., Jasinski, M., Santelia,
D., Martinoia, E., Sumner, L.W., Banta, L.M., Stermitz, F., et al. (2008). Altered Profile of
Secondary Metabolites in the Root Exudates of Arabidopsis ATP-Binding Cassette
Transporter Mutants. Plant Physiology *146*, 762–771.

Badri, D.V., Quintana, N., El Kassis, E.G., Kim, H.K., Choi, Y.H., Sugiyama, A., Verpoorte,
R., Martinoia, E., Manter, D.K., and Vivanco, J.M. (2009). An ABC transporter mutation
alters root exudation of phytochemicals that provoke an overhaul of natural soil microbiota.
Plant Physiol. *151*, 2006–2017.

Bailey, P.H.J., Currey, J.D., and Fitter, A.H. (2002). The role of root system architecture and
root hairs in promoting anchorage against uprooting forces in Allium cepa and root mutants
of Arabidopsis thaliana. J. Exp. Bot. *53*, 333–340.

Baulcombe, D.C., Saunders, G.R., Bevan, M.W., Mayo, M.A., and Harrison, B.D. (1986).
Expression of biologically active viral satellite RNA from the nuclear genome of transformed
plants. Nature *321*, 446–449.

Baumberger, N., Ringli, C., and Keller, B. (2001). The chimeric leucine-rich repeat/extensin
cell wall protein LRX1 is required for root hair morphogenesis in Arabidopsis thaliana.
Genes Dev. *15*, 1128–1139.

Berger, S., Bell, E., and Mullet, J.E. (1996). Two Methyl Jasmonate-Insensitive Mutants
Show Altered Expression of AtVsp in Response to Methyl Jasmonate and Wounding. Plant
Physiology *111*, 525–531.

Böhme, K., Li, Y., Charlot, F., Grierson, C., Marrocco, K., Okada, K., Laloue, M., and
Nogué, F. (2004). The Arabidopsis COW1 gene encodes a phosphatidylinositol transfer
protein essential for root hair tip growth. Plant J. *40*, 686–698.

Burylo, M., Rey, F., Mathys, N., and Dutoit, T. (2012). Plant root traits affecting the
resistance of soils to concentrated flow erosion. Earth Surface Processes and Landforms *37*,
1463–1470.

Carol, R.J., Takeda, S., Linstead, P., Durrant, M.C., Kakesova, H., Derbyshire, P., Drea, S.,
Zarsky, V., and Dolan, L. (2005). A RhoGDP dissociation inhibitor spatially regulates
growth in root hair cells. Nature *438*, 1013–1016.

Carvalhais, L.C., Dennis, P.G., Badri, D.V., Kidd, B.N., Vivanco, J.M., and Schenk, P.M.
(2015). Linking Jasmonic Acid Signaling, Root Exudates, and Rhizosphere Microbiomes.
Mol. Plant Microbe Interact. *28*, 1049–1058.

Choi, H.-S., and Cho, H.-T. (2019). Root hairs enhance Arabidopsis seedling survival upon
soil disruption. Scientific Reports *9*, 11181.

Datta, S., kim, C.M., Pernas, M., Pires, N.D., Proust, H., Tam, T., Vijayakumar, P., and
Dolan, L. (2011). Root hairs: development, growth and evolution at the plant-soil interface.
Plant and Soil *346*, 1–14.

De Baets, S., and Poesen, J. (2010). Empirical models for predicting the erosion-reducing
effects of plant roots during concentrated flow erosion. Geomorphology *118*, 425–432.

De Baets, S., Poesen, J., Knapen, A., and Galindo, P. (2007). Impact of root architecture on
the erosion-reducing potential of roots during concentrated flow. Earth Surface Processes and
Landforms *32*, 1323–1345.

De Baets, S., Denbigh, T.D., Smyth, K.M., Eldridge, B.M., Weldon, L.M., Higgins, B.W.,
Matyjaszkiewicz, A.W., Meersmans, J., Larson, E.R., Chenchiah, I.V., et al. (2020). Micro-
scale interactions between Arabidopsis root hairs and soil particles influence soil erosion.
Nature Comms Biol *3*, 1–11.

De Caroli, M., Lenucci, M.S., Di Sansebastiano, G.-P., Dalessandro, G., De Lorenzo, G., and
Piro, G. (2011). Dynamic protein trafficking to the cell wall. Plant Signal Behav *6*, 1012–
1015.

Diet, A., Link, B., Seifert, G.J., Schellenberg, B., Wagner, U., Pauly, M., Reiter, W.-D., and
Ringli, C. (2006). The Arabidopsis root hair cell wall formation mutant lrx1 is suppressed by
mutations in the RHM1 gene encoding a UDP-L-rhamnose synthase. Plant Cell *18*, 1630–
1641.

Drakakaki, G., Robert, S., Szatmari, A.-M., Brown, M.Q., Nagawa, S., Van Damme, D.,
Leonard, M., Yang, Z., Girke, T., Schmid, S.L., et al. (2011). Clusters of bioactive
compounds target dynamic endomembrane networks in vivo. Proceedings of the National
Academy of Sciences *108*, 17850–17855.

Edwards, K., Johnstone, C., and Thompson, C. (1991). A simple and rapid method for the
preparation of plant genomic DNA for PCR analysis. Nucleic Acids Res *19*, 1349.

Eisenach, C., Chen, Z.-H., Grefen, C., and Blatt, M.R. (2012). The trafficking protein
SYP121 of Arabidopsis connects programmed stomatal closure and K$^+$ channel activity with
vegetative growth. Plant J. *69*, 241–251.

Favery, B., Ryan, E., Foreman, J., Linstead, P., Boudonck, K., Steer, M., Shaw, P., and
Dolan, L. (2001). KOJAK encodes a cellulose synthase-like protein required for root hair cell
morphogenesis in Arabidopsis. Genes Dev. *15*, 79–89.

Galloway, A.F., Pedersen, M.J., Merry, B., Marcus, S.E., Blacker, J., Benning, L.G., Field,
K.J., and Knox, J.P. (2018). Xyloglucan is released by plants and promotes soil particle
aggregation. New Phytol. *217*, 1128–1136.

Galloway, A.F., Akhtar, J., Marcus, S.E., Fletcher, N., Field, K., and Knox, P. (2020). Cereal
root exudates contain highly structurally complex polysaccharides with soil-binding
properties. The Plant Journal.

Geelen, D., Leyman, B., Batoko, H., Di Sansebastiano, G.-P., Moore, I., Blatt, M.R., and Di
Sansabastiano, G.-P. (2002). The abscisic acid-related SNARE homolog NtSyr1 contributes
to secretion and growth: evidence from competition with its cytosolic domain. Plant Cell *14*,
387–406.

Ghestem, M., Cao, K., Ma, W., Rowe, N., Leclerc, R., Gadenne, C., and Stokes, A. (2014). A
framework for identifying plant species to be used as "ecological engineers" for fixing soil on
unstable slopes. PLoS ONE *9*, e95876.

Grierson, C.S., Roberts, K., Feldmann, K.A., and Dolan, L. (1997). The COW1 locus of
arabidopsis acts after RHD2, and in parallel with RHD3 and TIP1, to determine the shape,
rate of elongation, and number of root hairs produced from each site of hair formation. Plant
Physiol *115*, 981–990.

Haling, R.E., Brown, L.K., Bengough, A.G., Young, I.M., Hallett, P.D., White, P.J., and
George, T.S. (2013). Root hairs improve root penetration, root-soil contact, and phosphorus
acquisition in soils of different strength. J. Exp. Bot. *64*, 3711–3721.

Healey, A., Furtado, A., Cooper, T., and Henry, R.J. (2014). Protocol: a simple method for
extracting next-generation sequencing quality genomic DNA from recalcitrant plant species.
Plant Methods *10*, 21.

Huang, Y., Wang, Y., Tan, L., Sun, L., Petrosino, J., Cui, M.-Z., Hao, F., and Zhang, M.
(2016). Nanospherical arabinogalactan proteins are a key component of the high-strength
adhesive secreted by English ivy. Proc. Natl. Acad. Sci. U.S.A. *113*, E3193-3202.

Kang, J., Yim, S., Choi, H., Kim, A., Lee, K.P., Lopez-Molina, L., Martinoia, E., and Lee, Y.
(2015). Abscisic acid transporters cooperate to control seed germination. Nat Commun *6*,
8113.

Kidd, B.N., Edgar, C.I., Kumar, K.K., Aitken, E.A., Schenk, P.M., Manners, J.M., and
Kazan, K. (2009). The mediator complex subunit PFT1 is a key regulator of jasmonate-
dependent defense in Arabidopsis. Plant Cell *21*, 2237–2252.

Kitakura, S., Vanneste, S., Robert, S., Löfke, C., Teichmann, T., Tanaka, H., and Friml, J.
(2011). Clathrin mediates endocytosis and polar distribution of PIN auxin transporters in
Arabidopsis. Plant Cell *23*, 1920–1931.

Lamesch, P., Berardini, T.Z., Li, D., Swarbreck, D., Wilks, C., Sasidharan, R., Muller, R.,
Dreher, K., Alexander, D.L., Garcia-Hernandez, M., et al. (2012). The Arabidopsis
Information Resource (TAIR): improved gene annotation and new tools. Nucleic Acids Res.
40, D1202-1210.

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat
Methods *9*, 357–359.

Larson, E.R., Domozych, D.S., and Tierney, M.L. (2014). SNARE VTI13 plays a unique role
in endosomal trafficking pathways associated with the vacuole and is essential for cell wall
organization and root hair growth in arabidopsis. Ann. Bot. *114*, 1147–1159.

Larson, E.R., Van Zelm, E., Roux, C., Marion-Poll, A., and Blatt, M.R. (2017). Clathrin
Heavy Chain Subunits Coordinate Endo- and Exocytic Traffic and Affect Stomatal
Movement. Plant Physiol. *175*, 708–720.

Lin, D.Y., and Ying, Z. (1994). Semiparametric Analysis of the Additive Risk Model.
Biometrika *81*, 61–71.

Lorenzo, O., Chico, J.M., Sánchez-Serrano, J.J., and Solano, R. (2004). JASMONATE-
INSENSITIVE1 encodes a MYC transcription factor essential to discriminate between
different jasmonate-regulated defense responses in Arabidopsis. Plant Cell *16*, 1938–1950.

Naveed, M., Brown, L.K., Raffan, A.C., George, T.S., Bengough, A.G., Roose, T., Sinclair,
I., Koebernick, N., Cooper, L., Hackett, C.A., et al. (2017). Plant exudates may stabilize or
weaken soil depending on species, origin and time. European Journal of Soil Science *68*,
806–816.

Parker, J.S., Cavell, A.C., Dolan, L., Roberts, K., and Grierson, C.S. (2000). Genetic
interactions during root hair morphogenesis in Arabidopsis. Plant Cell *12*, 1961–1974.

Plessis, A., Cournol, R., Effroy, D., Silva Pérez, V., Botran, L., Kraepiel, Y., Frey, A., Sotta,
B., Cornic, G., Leung, J., et al. (2011). New ABA-hypersensitive Arabidopsis mutants are
affected in loci mediating responses to water deficit and Dickeya dadantii infection. PLoS
ONE *6*, e20243.

Prentice, R.L., and Kalbfleisch, J.D. (2003). Mixed discrete and continuous Cox regression
model. Lifetime Data Anal *9*, 195–210.

R Core Team (2014). R: A language and environment for statistical computing (Vienna,
Austria: R Foundation for Statistical Computing).

Ringli, C., Bigler, L., Kuhn, B.M., Leiber, R.-M., Diet, A., Santelia, D., Frey, B., Pollmann,
S., and Klein, M. (2008). The Modified Flavonol Glycosylation Profile in the Arabidopsis
rol1 Mutants Results in Alterations in Plant Growth and Cell Shape Formation. The Plant
Cell *20*, 1470–1481.

Robert, S., Chary, S.N., Drakakaki, G., Li, S., Yang, Z., Raikhel, N.V., and Hicks, G.R.
(2008). Endosidin1 defines a compartment involved in endocytosis of the brassinosteroid
receptor BRI1 and the auxin transporters PIN2 and AUX1. PNAS *105*, 8464–8469.

Robinson, D.G., Scheuring, D., Naramoto, S., and Friml, J. (2011). ARF1 localizes to the
golgi and the trans-golgi network. Plant Cell *23*, 846–849; author reply 849-850.

Rodriguez-Furlán, C., Salinas-Grenet, H., Sandoval, O., Recabarren, C., Arraño-Salinas, P.,
Soto-Alvear, S., Orellana, A., and Blanco-Herrera, F. (2016). The Root Hair Specific
SYP123 Regulates the Localization of Cell Wall Components and Contributes to
Rizhobacterial Priming of Induced Systemic Resistance. Front Plant Sci *7*, 1081.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. Nat. Methods *9*, 676–682.

Stokes, A., Atger, C., Bengough, A.G., Fourcard, T., and Sidle, R.C. (2009). Desirable plant
root traits for protecting natural and engineered slopes against landslides. Plant and Soil *324*,
1–30.

Toukura, Y., Devee, E., and Hongo, A. (2006). Uprooting and shearing resistances in the
seedlings of four weedy species. Weed Biology and Management *6*, 35–43.

Waghmare, S., Lileikyte, E., Karnik, R., Goodman, J.K., Blatt, M.R., and Jones, A.M.E.
(2018). SNAREs SYP121 and SYP122 Mediate the Secretion of Distinct Cargo Subsets.
Plant Physiology *178*, 1679–1688.

Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis (Springer).

Yang, J., Bak, G., Burgin, T., Barnes, W.J., Mayes, H.B., Peña, M.J., Urbanowicz, B.R., and
Nielsen, E. (2020). Biochemical and Genetic Analysis Identify CSLD3 as a beta-1,4-Glucan
Synthase That Functions during Plant Cell Wall Synthesis. Plant Cell *32*, 1749–1767.

TABLES

Table 1 *Arabidopsis thaliana* mutant lines used in this study.

Mutant type	Mutant	AGI code[a]	Mutant line	Referenced in
Root hair shape	*cow1-3*	At4g34580	SALK_0021245	Böhme *et al.* (2004)
	csld3-1	At3g03050	CS899	Favery *et al.* (2001)
	lrx1-4	At1g12040	SALK_057038	Kim *et al.* (2006)
	rol1-2	At1g78570	CS16373	Diet *et al.* (2006)
	gdi1-2	At3g07880	SALK_035400	Kang *et al.* (2017)
Vesicle trafficking	*chc1-2*	At3g11130	SALK_103252	Kitakura *et al.* (2011)
	chc2-3	At3g08530	SALK_151638	Bourdais *et al.* (2016)
	syp121	At3g11820	-	Assaad *et al.* (2004)
	syp122-1	At3g52400	SALK_008617	Assaad *et al.* (2004)
Root exudate composition	*jin1-9*	At1g32640	SALK_017005	Anderson *et al.* (2004)
	pdr2	At4g15230	SAIL_811_F08	Badri *et al.* (2008)
	pft1-3	At1g25540	SALK_059316	Kidd *et al.* (2009)
Unknown	*abcg43-1*	At4g15236	N75206[b]	This study
	abcg43-2	At4g15236	SALK_201207	This study
	abcg43-3	At4g15236	SALKseq_30713	This study

FIGURE LEGENDS

FIGURE 1. Centrifuge assay workflow.

(A) i. Ten seeds (highlighted in pink) are sown onto the surface of sterile, solid gel growth

medium in a single Petri plate in two horizontal rows. ii. The plates are stacked in groups of

five and orientated vertically at approximately 80° to encourage the roots to grow down the

surface of the gel medium. Plates are grown in a growth chamber with constant light (120-

145 µmol m^{-2} s^{-1}) conditions at 22°C and 60% relative humidity.

(B) After 5-6 days, i. seedlings are visually analysed and numbered. ii. Petri plates are placed

into a swing-out-bucket centrifuge in an inverted orientation with their roots pointing inward

(indicated by the purple arrows). iii. After seedlings have been subjected to a one-minute

pulses of increasing centrifugal speeds, seedling detachment is recorded.

(C) i. The aerial tissue mass of each seedling is determined using ii. an analytical scale to iii.

determine the root-gel adhesion properties of candidate lines are compared to wild-type to

assess whether they have increased (e.g. line y) or decreased (e.g. line x) adhesion to the

sterile gel.

FIGURE 2 Physical root hair properties contribute to root-substrate adhesion in *Arabidopsis*

thaliana.

(A) Root hair phenotypes of 5-d-old wild-type (Col-0), *lrx 1-4, csld3-1, cow1-3, rol1-2,* and

gdi1-2 seedlings grown on gel medium and statistical comparisons of mean root hair density

(number per mm length of root) and mean root hair length (mm) for each mutant line relative

to wild type. White asterisks on the root hair images indicate the characteristic root hair

bulging phenotype in the *lrx1-4* mutant and the short, branching root hair phenotype in *gdi1-*

2. Scale bar = 0.25 mm. In the table, the mean ± standard error is given for root hair density

and root hair length of wild type and each mutant as well as a mean comparison to wild type,

which is listed in bold. "No difference" is stated when there was no statistically significant

difference between wild type and a mutant line. The statistical output of each univariate

linear model is given. Alpha = 0.025, statistical significance codes: '***' = < 0.001, '**' = $<$

0.01. The results shown are from a representative experiment for at least two independent

experiments. Survival curves displaying the proportion of seedlings that adhered to the gel at

increasing centrifugal force for **(B)** 92 wild type (Col-0 – black); 81 *lrx1-4* (light green); 70

csld3-1 (green); **(C)** 82 wild type (Col-0 - black); 83 *cow1-3* (green); and **(D)** 77 wild type

(Col-0- black); 85 *rol1-2* (light green); and 86 *gdi1-2* (dark green). Circled red crosses on the

survival curves represent seedlings that remained adhered to the gel after the maximum

centrifugal speed (1611 RPM). The results shown are from a representative experiment for at

least two independent experiments showing a statistically significant difference in adhesion

between mutant lines relative to wild type (Cox PH regression; alpha = 0.01). A single

experiment included $\geq$ 70 biological replicates for each candidate line.

FIGURE 3 Vesicle trafficking mechanisms that contribute to root-substrate adhesion in

Arabidopsis thaliana.

(A) Root hair phenotypes of 5-d-old wild-type (Col-0), *syp121, syp122-1, chc1-2* and *chc2-3*

seedlings grown on a gel medium and statistical comparisons of mean root hair density

(number per mm length of root) and mean root hair length (mm) for each mutant line relative

to wild type. Scale bar = 0.25 mm. In the table, the mean $\pm$ standard error is given for the root

hair density and root hair length of wild type and each mutant as well as a mean comparison

to wild type, which is listed in bold. "No difference" is stated when there was no statistically

significant difference between wild type and a mutant line. The statistical output of each

univariate linear model is given. Alpha = 0.025. The results shown are from a representative

experiment for at least two independent experiments. Survival curves displaying the

proportion of seedlings that adhered to the gel at increasing centrifugal force for **(B)** 87 wild

type (Col-0 – black); 91 *syp121* (turquoise); 83 *syp122-1* (blue/grey); and **(C)** 83 wild type

(Col-0 – black); 70 *chc1-2* (dark blue)*;* and 72 *chc2-3* (medium blue)*.* Red crosses circled on

the survival curves represent seedlings that remained adhered to the gel after the maximum

centrifugal speed (1611 RPM). The results shown are from a representative experiment for at

least two independent experiments showing a statistically significant difference in adhesion

between all mutant lines relative to wild type (Cox PH regression; alpha = 0.01). A single

experiment included $\geq$ 70 biological replicates for each candidate line.

FIGURE 4 Exudate composition changes root-substrate adhesion properties in *Arabidopsis*

thaliana.

(A) Root hair phenotypes of 5-d-old wild-type (Col-0), *jin1-9, pdr2* and *pft1-3* seedlings

grown on a gel medium and statistical comparisons of mean root hair density (number per

mm length of root) and mean root hair length (mm) for each mutant line relative to wild type.

Scale bar = 0.25 mm. In the table, the mean $\pm$ standard error is given for the root hair density

and root hair length of wild type and each mutant as well as a mean comparison to wild type,

which is listed in bold. "No difference" is stated when there was no statistically significant

difference between wild type and a mutant line. The statistical output of each univariate

linear model is given. Alpha = 0.025. The results shown are from a representative experiment

for at least two independent experiments. **(B)** Survival curves displaying the proportion of

seedlings that adhered to the gel at increasing centrifugal force for 80 wild type (Col-0 –

black), 81 *jin1-9* (light pink), 80 *pdr2* (light orange), and 81 *pft1-3* (dark orange). The results

shown are from a representative experiment for at least two independent experiments

showing a statistically significant difference in adhesion between mutant lines relative to wild

type (Cox PH regression; alpha = 0.01), except for *jin1-9*. A single experiment included $\geq$ 70

biological replicates for each candidate line.

FIGURE 5 Using the centrifuge assay in a forward genetic screen to identify *Arabidopsis*

thaliana root-substrate adhesion mutants.

(A) T-DNA insert locations in *ABCG43* for each *abcg43* mutant allele, with insertions

indicated by orange arrowheads. The *ABCG43* gene is located on chromosome four at

position Chr4: 8,969,677-8,702,727 and contains 23 exons (dark green) and 22 introns (white

gaps). *abcg43-1* has a T-DNA insert in the exon10/intron10 boundary, *abcg43-2*

(SALK_201207) has a T-DNA insert in exon 2 and *abcg43-3* (SALKseq_30713) has a T-

DNA insert in exon 3. **(B)** Genomic PCR confirming the homozygosity of each *abcg43*

mutant for the respective T-DNA inserts. Lanes **1** and **2** were loaded with the gene-specific

and T-DNA-border PCR products from *abcg43* mutant lines. Lanes **3** and **4** were loaded with

the gene-specific and T-DNA-border PCR products from a wild type (Col-0) genomic DNA

template. Lanes **5** and **6** were loaded with the water controls for the gene-specific and T-

DNA-border PCR reactions, respectively. '**L**' indicates the 100-bp ladder. The expected

product sizes for the gene-specific PCRs were ~350 nt, ~1025 nt and ~1090 nt; the T-DNA-

border PCR product sizes were ~400 nt, ~550 nt and ~750 nt for *abcg43-1, abcg43-2* and

abcg43-3 alleles, respectively. **(C)** Survival curves displaying the proportion of seedlings that

adhered to the gel at increasing centrifugal force for 79 wild type (Col-0 – black), 70 *abcg43-*

1 (dark pink), 85 *abcg43-2* (purple), and 73 *abcg43-3* (light purple). Red crosses circled on

the survival curves represent seedlings that remained adhered to the gel after the maximum

centrifugal speed (1611 RPM). The results are from a representative experiment for at least

two independent experiments showing a statistically significant difference in adhesion

between the mutant lines relative to wild type (Cox PH regression; alpha = 0.01). A single

experiment included $\geq$ 70 biological replicates for each candidate line.

A

Line	Root hair density (number per mm length of root)		Root hair length (mm)	
Col-0	30.556 ± 0.784	-	0.518 ± 0.009	-
lrx1-4	31.889 ± 0.935 **No difference**	t = 1.093, P > 0.05 d.f. = 16, n = 9	0.388 ± 0.008 **1.3 times shorter****	t = -2.819, P < 0.001 d.f. = 16, n = 9
csld3-1	8.778 ± 1.188 **3.5 times less*****	t = -15.30, P < 0.001 d.f. = 16, n = 9	0.045 ± 0.002 **10.1 times shorter*****	t = -11.790, P < 0.001 d.f. = 16, n = 9
cow1-3	29.200 ± 0.772 **No difference**	t = -1.23, P > 0.05 d.f. = 17, n = 10	0.049 ± 0.001 **10 times shorter*****	t = -12.37, P < 0.001 d.f. = 17, n = 10
rol1-2	30.700 ± 1.550 **No difference**	t = 0.08, P > 0.05 d.f. = 17, n = 9	0.195 ± 0.003 **2.7 times shorter*****	t = -8.417, P < 0.001 d.f. = 17, n = 10
gdi1-2	26.200 ± 1.041 **0.17 times less****	t = -3.282, P < 0.01 d.f. = 17, n = 9	0.135 ± 0.006 **3.8 times shorter*****	t = -9.04, P < 0.001 d.f. = 16, n = 9

B

C

D

Line	Root hair density (number per mm length of root)		Root hair length (mm)	
Col-0	32.000 ± 0.632	-	0.420 ± 0.005	-
syp121	31.200 ± 0.952 **No difference**	t = -0.700, P > 0.05 d.f. = 18, *n* = 10	0.461 ± 0.004 **No difference**	t = 1.974, P > 0.05 d.f. = 18, *n* = 10
syp122-1	32.200 ± 0.442 **No difference**	t = 0.259, P > 0.05 d.f. = 18, *n* = 10	0.452 ± 0.005 **No difference**	t = 1.379, P > 0.05 d.f. = 18, *n* = 10
chc1-2	30.889 ± 0.716 **No difference**	t = -1.168, P > 0.05 d.f. = 17, *n* =9	0.441 ± 0.006 **No difference**	t = 0.684, P > 0.05 d.f. = 17, *n* = 9
chc2-3	31.889 ± 0.790 **No difference**	t = -0.111, P > 0.05 d.f. = 17, *n* = 9	0.463 ± 0.006 **No difference**	t = 0.182, P > 0.05 d.f. = 18, *n* = 10

A

Line	Root hair density (number per mm length of root)		Root hair length (mm)	
Col-0	30.111 ± 1.612	-	0.416 ± 0.005	-
jin1-9	29.444 ± 1.271 **No difference**	t = -0.324, P > 0.05 d.f. = 16, *n* = 9	0.429 ± 0.006 **No difference**	t = 0.555, P > 0.05 d.f. = 16, *n* = 9
pdr2	31.125 ± 0.854 **No difference**	t = 0.533, P > 0.05 d.f. = 15, *n* = 8	0.428 ± 0.006 **No difference**	t = 0.521, P > 0.05 d.f. = 15, *n* = 8
pft1-3	29.000 ± 1.179 **No difference**	t = -0.555, P > 0.05 d.f. = 16, *n* = 9	0.424 ± 0.006 **No difference**	t = 0.295, P > 0.05 d.f. = 16, *n* = 9

B

A

abcg43-2
abcg43-3
abcg43-1

ABCG43
(At4g15236)

Chr4: 8,969,677   Chr4: 8,702,727

B

kb   L   1   2   3   4   5   6

abcg43-1
1.0
0.8
0.5
0.3

abcg43-2
1.2
1.0
0.8
0.5

abcg43-3
1.2
1.0
0.8
0.5

C

Percentage of seedlings adhered to the gel (%)

Centrifugal force (mN)

— Col-0
— *abcg43-1*
— *abcg43-2*
— *abcg43-3*