1   **When effort matters: Expectations of reward and efficacy guide cognitive control**

2   **allocation**

3   Frömer, R.[1*], Lin, H.[2*], Dean Wolf, C. K.[1], Inzlicht, M.[2,3], & Shenhav, A.[1]

5   *[1] Cognitive, Linguistic, and Psychological Sciences, Carney Institute for Brain*

6   *Science, Brown University, Providence, RI,*

7   *[2] Department of Psychology, University of Toronto*

8   *[3] Rotman School of Management, University of Toronto*

10   ** equal contributions*

13   Correspondence should be addressed to Romy Frömer (romy_fromer@brown.edu) or Hause Lin

14   (hauselin@gmail.com)

15   **Abstract**

The amount of mental effort we invest in a task is influenced by the reward we can expect if we

perform that task well. However, some of the rewards that have the greatest potential for driving

these efforts (e.g., jobs, grants) are partly determined by factors beyond one's control. In such

cases, effort has more limited *efficacy* for obtaining rewards. We have proposed that people

integrate information about the expected reward and efficacy for effort to determine the expected

value of control, and then adjust their control allocation (i.e. mental effort) accordingly. Here we

test this theory's key behavioral and neural predictions. We show that participants invest more

control when this control is more rewarding and more efficacious, and that these incentive

components separately modulate EEG signatures of incentive evaluation and proactive control

allocation. Our findings support the prediction that people combine worth and worthwhileness to

determine how much effort to invest.

Cognitive control is critical to one's ability to achieve most goals [1-3]–whether to complete

a paper in time for its deadline or sending that birthday message amidst a busy workday – but

exerting control appears to come at a cost. We experience cognitive control as mentally

effortful[4,5], and therefore require some form of incentive to justify investing control in a task[6,7].

For instance, a student is likely to study harder for an exam that has higher stakes (e.g., worth

half of their grade) than a lower-stakes exam. Accordingly, research has shown that participants

generally exert more mental effort on a cognitive control task (e.g., Stroop, flanker) when they

are offered higher rewards for performing well, as evidenced by improved task performance and

increased engagement of relevant control circuits [7-16].

In the real world, increased control may not translate to achieving desired outcomes (e.g.,

when *efficacy* is low). For instance, even when the stakes are high, that same student is likely to

exert less effort studying if they think that those efforts have little bearing on their score on that

exam, say if they felt that grading for the exam is driven mostly by factors out of their control

(e.g., random chance). While previous work has closely examined the mechanisms by which

people evaluate the potential rewards to expect for a certain control allocation, much less is

known about how they evaluate the efficacy of that control, nor how these two incentive

components (reward and efficacy) are integrated to determine how much control is invested.

We have recently developed a model that formalizes the roles of evaluation, decision-

making, and motivation in allocating and adjusting cognitive control [17,18] (Fig.1). Our model

describes how cognitive control can be allocated based on the overall worth of executing

different types and amounts of control, which we refer to as their Expected Value of Control

(EVC). The EVC of a given control allocation is determined by the extent to which the costs that

would need to be incurred (mental effort) are outweighed by the benefits. Critically, these

benefits are a function of both the expected outcomes for reaching one's goal (*reward*, e.g.,

money or praise) and the likelihood that this goal will be reached with a given investment of

control (*efficacy*) (Fig. 1A). The amount of control invested is predicted to increase

monotonically with a combination of these two incentive components (Fig. 1B).

Figure 1. The EVC model predicts that control should increase with expected reward and efficacy.
A. EVC model. Control intensity is chosen to optimize the trade-off between effort costs and payoff,
maximizing the expected value of control. The payoff of a given control signal is determined by the
expected reward and efficacy for a given control intensity. **B**. The EVC model proposes that higher
intensities of control (x-axis) are associated with greater effort costs (red) but that these effort costs can be
outweighed by the expected payoff for a given control intensity (blue). These payoffs typically increase as
a function of task performance, and task performance typically improves as a function of control
intensity. The EVC of each control intensity (purple) is calculated as the difference between its payoff
and its cost. The optimal level of control to invest is the one that maximizes EVC (vertical black arrows).
Left: When the expected reward for performing the task well is higher (from dashed to solid line), higher
control intensities achieve even higher payoffs. **Right**: When performance matters less for acquiring a
given reward (low efficacy [dashed line]) – here, simulated by having reward be unrelated to performance
on most trials (instead, occurring at a fixed high rate; see Fig. 2) – the relative payoff for high vs. low
control intensities decreases.

The EVC model integrates over and formalizes past theories that posit roles for

reward/utility and/or efficacy/controllability/agency in the motivation to engage in a particular

course of action [19-26]. In so doing, our model enables a description of the computational and

neural mechanisms of control allocation. For instance, past research has shown that behavioral

and neural markers of proactive control increase with increases in anticipated task difficulty[27-32].

Through the lens of the EVC theory (Fig. 1), these difficulty-related increases in control intensity

can be accounted for by changes in expected reward (i.e., the harder the task, the less likely you

are to achieve the rewards associated with performing the task well) and/or changes in expected

efficacy (i.e., the harder the task, the less helpful a given level of control is for achieving the

same level of performance). The latter explains why the relationship between control intensity

and task difficulty is nonmonotonic - once the task exceeds a certain difficulty (i.e., once effort is

no longer efficacious; [33,34]), a person stops intensifying their mental efforts and instead starts to

disengage from a task.

Our theory therefore makes the novel prediction that variability in efficacy (holding

expected reward and difficulty constant) should itself be sufficient to drive changes in behavioral

and neural signatures of control allocation. The theory makes the further prediction that reward

and efficacy should shape incentive processing and associated neural correlates at multiple

stages, including during the initial evaluation of each of these incentive components and at the

point when those components converge to determine control allocation based on their combined

value (EVC).

Here, we test these predictions across two studies using a novel paradigm that explicitly

dissociates expectations of reward and efficacy associated with a cognitive control task (the

Stroop task; Fig. 2), allowing us to isolate their individual and joint contributions to control

allocation. To further examine how reward and efficacy are encoded at different stages of

incentive processing, in Study 2 we measured EEG and pupillometry while participants

performed this task, allowing us to separately measure the extent to which reward and efficacy

are reflected in signals associated with the initial evaluation of the incentives available on a

given trial (putatively indexed by the post-cue P3b; [27,35]) versus those associated with proactive

allocation of the control deemed appropriate for the upcoming trial (putatively indexed by the

contingent negative variation [CNV] occurring prior to the presentation of the target stimulus;

[27,30,32,36-38]). Confirming our predictions, both studies found that participants adaptively increased

their control allocation (and thus performed better at the task) when expecting higher levels of

reward and efficacy. Study 2 showed that both incentive components amplify event-related

potentials (ERPs) associated with distinct stages of incentive processing: incentive evaluation

(indexed by the P3b following cue presentation) and control allocation (indexed by the

contingent negative variation [CNV] prior to Stroop target onset). Critically, only the CNV

reflected the integration of reward and efficacy. The amplitude of both ERPs, but more so the

CNV, predicted performance on the upcoming trial, supporting the prediction that these neural

signals index different stages in the evaluation and allocation of control.

Figure 2. Manipulation of expected reward and efficacy. On each trial, participants view an incentive
cue followed by a Stroop stimulus (target) and then feedback indicating how much reward they received.
Four different cues indicate whether a trial is high or low in reward and efficacy. The P3b and CNV
event-related potentials are measured during the interval between cue and target, as indices of cue
evaluation (P3b) and control allocation (CNV).

117   **Results**

To test the prediction that reward and efficacy together shape cognitive effort investment

and task performance, we developed and validated a novel paradigm that manipulates efficacy

independently from expected reward (Fig. 2). Specifically, prior to the onset of a Stroop stimulus

(the *target*), we cued participants with the amount of monetary reward they would receive if

successful on that trial ($0.10 vs. $1.00) and whether success would be determined by their

performance (being fast and accurate; *high efficacy*) or whether it would instead be determined

independently of their performance (based on a weighted coin flip; *low efficacy*). Using an

adaptive yoking procedure, we held expected reward constant across efficacy levels, while also

varying reward and efficacy independently of task difficulty (i.e., levels of congruency).

Participants performed this task in an experimental session that measured only behavior

(Study 1; N=21) or one that additionally measured EEG activity and pupillometry (Study 2;

N=44). Studies 1-2 had the same trial structure but differed slightly in the design of the incentive

cues, overall number of trials, and within-trial timing, and were run at different study sites (see

Methods). Predictions for Study 2 were preregistered based on findings from Study 1

(osf.io/35akg).

Performance improves with increasing expected reward and efficacy. We predicted that

reward and efficacy would together incentivize greater control allocation. Given that participants

needed to be fast and accurate to perform well on our task, we expected to find that participants

would be faster to respond correctly when they expected control to be more rewarding and more

efficacious. Replicating previous findings[27], across both studies we found that reaction times on

accurate trials (i.e., accurate RTs) were faster for high compared to low reward trials (Study 1: b

= -9.81, p = .002; Study 2: b = -5.03, p = .004). Critically, accurate RTs were *also* faster for high

compared with low efficacy trials (Study 1: b = -14.855, p <.001; Study 2: b = -5.89, p = .016).

We further found reward-efficacy interactions in the predicted direction – with the speeding

effect of reward being enhanced on high efficacy trials – but this interaction was only significant

in Study 2 (Study 1: b = -9.75, p = .116; Study 2: b = -9.23, p = .009; cf. Fig. 3).

Figure 3. Reward and efficacy improve performance. Participants in both Study 1 (**A**) and Study 2 (**B**)
were fastest to give a correct response when both reward and efficacy were high, suggesting that these
variables jointly determined control allocation. Error bars represent within-subject standard error of the
mean.

Additional analyses confirmed that these performance improvements were not driven by

speed-accuracy tradeoffs. Whereas participants were faster when reward or efficacy was high,

they were not less accurate (Tables S1-S2). In fact, their accuracies tended to improve when

reward or efficacy were high, though only the effect of efficacy on accuracy in Study 2 was

significant (b = 0.08, p = .033, Table S2). Together, the faster RTs and more accurate responses

suggest that the effects of reward and efficacy on response speed reflected increased control

allocation rather than speed-accuracy tradeoffs (lowering of response thresholds; i.e., increased

impulsivity).

All of the analyses above control for the effects of task difficulty (response congruence)

and trial (practice) effects on performance, which in both studies manifested as worse

performance (slower and less accurate responding) with increasing response incongruence, and

improved performance (faster and more accurate responding) over time (Table S1-S2, Fig. S1).

The effects of reward and efficacy on performance did not significantly differ between the two

studies (Table S3).

Reward and efficacy levels are reflected in neural signatures of cue evaluation and

control allocation. Our behavioral results suggest that participants adjust their mental effort

investment (allocation of cognitive control) based on the expected rewards and the degree to

which control is perceived to be efficacious; that is, they invest more effort when the expected

reward and efficacy are high. To examine the neural and temporal dynamics associated with the

processing of these two incentive components, we focused on two well-characterized event-

related potentials (ERPs): the P3b, which peaks around 250-550 ms following cue onset and is

typically associated with cue evaluation[27]; the CNV, which emerges about 500 ms prior to Stroop

target onset and is typically associated with preparatory attention or proactive

control[27,28,30,32,36,37]. Based on past research, we preregistered the predictions below for the CNV.

Additional predictions regarding the P3b were generated subsequent to preregistration based on

further review of the literature.

We found that reward and efficacy modulated both of these ERPs (Table 1, Fig. 4).

Replicating past work[27], cues signaling higher rewards were associated with significantly larger

amplitudes of both P3b (b = 0.34, p <.001) and CNV (b = -0.28, p = .001). Importantly, holding

reward constant, cues signaling high rather than low efficacy were likewise associated with

significantly larger amplitudes of P3b (b = 0.44, p <.001) and CNV (b = -0.30, p = .008).

Crucially, only the CNV tracked the interaction of reward and efficacy (b = -0.35, p

= .046), reflecting the finding that the effect of reward on CNV amplitude was enhanced when

efficacy was high. We did not find a significant reward-efficacy interaction with the P3b (b = -

0.01, p = .947). Thus, although reward and efficacy independently modulated the P3b and CNV

(i.e., main effects of reward and efficacy on both ERPs), only the CNV reflected their integration

(i.e., reward-efficacy interaction). This pattern of results is consistent with our prediction that

reward and efficacy are initially evaluated separately (reflected in the P3b), but are subsequently

integrated to determine EVC and thereby allocate control (reflected in the CNV).

Figure 4. P3b and CNV amplitudes track increases in expected reward and efficacy. A. ERP
averages for each incentive condition, separately for the P3b (**top**) and CNV (**bottom**). Shaded areas
indicate time windows used for quantification. Topographies show fixed-effect contrasts for reward and
efficacy, respectively. **B.** Average ERP amplitudes within the relevant time windows show that the
magnitude of the (positive-going) P3b and the (negative-going) CNV increase with greater reward and
efficacy. The CNV but not the P3b tracked the interaction of reward and efficacy, mirroring the
behavioral effects (cf. Fig. 3). **C-D.** Fixed effects of ERP amplitude on accuracy (**C**) and accurate RT (**D**)
show that increased P3b and CNV predicted better performance on the upcoming trial. Shaded error bars
represent standard error of the mean.

Table 1. *Effects of Reward and Efficacy on Neural Signals*

	P3b			CNV		
Predictors	*Estimates*	*CI*	*p*	*Estimates*	*CI*	*p*
(Intercept)	0.26	-0.27 – 0.78	0.339	-0.16	-0.63 – 0.31	0.507
Efficacy	0.44	0.24 – 0.65	**<0.001**	-0.30	-0.53 – -0.08	**0.008**
Reward	0.34	0.16 – 0.51	**<0.001**	-0.28	-0.45 – -0.11	**0.001**
Trial	-0.54	-0.62 – -0.46	**<0.001**	-0.03	-0.12 – 0.05	0.461
P3b Baseline	0.00	-0.01 – 0.01	0.950			
Efficacy:Reward	-0.01	-0.33 – 0.31	0.948	-0.35	-0.69 – -0.01	**0.046**
P3b				0.16	0.14 – 0.17	**<0.001**
CNV Baseline				-0.02	-0.03 – -0.00	**0.005**
Observations	22580			22580		

Note: Statistically significant p-values (< 0.05) are displayed in bold.

Neural signatures of incentive processing predict effort investment. We have shown

that reward and efficacy affect behavioral performance (accurate RT) and neural activity during

initial cue evaluation (P3b) and proactive control allocation (CNV), suggesting that these neural

signals reflect the transformation of incentives into effort allocation. To test this hypothesis more

directly, we included single-trial P3b and CNV amplitude (normalized within-subject) as

regressors in our models of accurate RT and accuracy, to test whether variability in these two

neural signals explained trial-by-trial variability in task performance (Table 2). We found that

both P3b and CNV were associated with better Stroop task performance on the upcoming trial:

Larger ERP magnitudes were associated with an increased probability of responding correctly

(P3b; b = 0.08, p <.001, CNV: b = -0.10, p <.001), and also with faster accurate RTs (P3b: b = -

7.04, p <.001, CNV: b = 15.58, p <.001). Crucially, the CNV's relationship with accurate RT was

significantly stronger than the P3b's (X^2 = 18.51, p < .001), providing evidence consistent with

our prediction that the CNV plays a more important role in allocating control than the P3b, and

with our observation that CNV's relationship with reward and efficacy more closely resembles

that found for accurate RT (i.e., both the CNV and accurate RT were modulated by the

interaction of reward and efficacy; compare Fig. 3 and Fig. 4B). However, CNV and P3b did not

differ reliably in their association with accuracy ($X^2 = 0.78$, p = .378). Together, these findings

suggest that the P3b and CNV index the transformation of incentive processing into effort

investment, a process that entails integration of the reward and efficacy expected on a given trial.

Table 2. *Effects of Neural Signals on Behavioral Performance*

Predictors	Accuracy			Accurate RT		
	Log-Odds	CI	p	Estimates	CI	p
(Intercept)	1.87	1.65 – 2.08	**<0.001**	647.66	630.60 – 664.71	**<0.001**
Efficacy	0.08	0.00 – 0.15	**0.044**	-4.00	-8.84 – 0.85	0.106
Reward	0.03	-0.05 – 0.10	0.468	-3.25	-6.85 – 0.34	0.076
P3b	0.08	0.04 – 0.11	**<0.001**	-7.04	-8.87 – -5.21	**<0.001**
CNV	-0.10	-0.14 – -0.06	**<0.001**	15.58	12.02 – 19.15	**<0.001**
Congruency (i-n)	0.65	0.44 – 0.85	**<0.001**	-64.00	-68.85 – -59.14	**<0.001**
Congruency (n-c)	0.34	0.18 – 0.49	**<0.001**	-15.86	-20.27 – -11.45	**<0.001**
Baseline	0.02	-0.02 – 0.07	0.326	2.23	0.05 – 4.42	**0.045**
Trial	0.04	0.00 – 0.08	**0.036**	-10.77	-12.58 – -8.95	**<0.001**
Efficacy:Reward				-9.96	-17.14 – -2.78	**0.007**
Observations	22580			18999		

223 Note: Statistically significant p-values (< 0.05) are displayed in bold. Congruency (n-i) refers to the comparison
224 between incongruent and neutral Stroop stimulus; Congruency (c-n) refers to the comparison between neutral and
225 congruent Stroop stimulus.
226

Opposing effects of expected reward and efficacy on pupil responses. As predicted,

pupil diameter was larger when expected reward was higher (Fig. 5). This effect was statistically

significant from 40 ms following cue onset and lasted throughout the trial (adjusted ps < .044).

Contrary to our predictions, however, pupil diameter was *smaller* when efficacy was higher and

this effect lasted from 220 ms to 2870 ms following cue onset (adjusted ps < .049). We did not

find a significant interaction between reward and efficacy on pupil diameter at any time point

within the 5000 ms window we tested (adjusted ps > .062). Although the opposing effects of

reward and efficacy on pupil responses might be difficult to reconcile with the behavioral and

EEG results (i.e., performance and ERP magnitudes both scaled positively with reward and

efficacy), they nevertheless provide evidence against the idea that our behavioral and EEG

results merely reflect changes in arousal. If this alternative explanation were true, pupil

diameter—a reliable index of arousal[39-41]—should have increased when either reward or efficacy

were high.

Figure 5. Effects of reward and efficacy on cue-related pupillary responses. **A.** Following an initial
cue-related constriction due to the pupillary light reflex, the pupil dilates and pupil responses then begin
to diverge depending on levels of expected reward and efficacy. **B.** Time-course of beta coefficients show
that reward and efficacy had opposing influences on pupil diameter, with reward positively associated and
efficacy negatively associated with pupil diameter. We did not find a significant interaction between these
two variables. The horizontal bars at the top of the plot show time points (in 10 ms steps) that survived
false-discovery rate (FDR) correction (p < .05). Shaded regions reflect 95% confidence intervals of the
parameter estimates.

Influences of incentives on EEG signatures of response and feedback monitoring.

While the focus of our study was on measures of incentive processing and proactive control

allocation, we preregistered secondary hypotheses regarding the potential influence reward and

efficacy might have on neural signatures of reactive control. Specifically, we predicted that these

incentive components might enhance monitoring of response accuracy and subsequent feedback.

Contrary to this hypothesis, when examining the error-related negativity (ERN) – a negative

deflection in response-locked activity for errors relative to correct responses [42,43] (though see [44]

for ERN elicited by partial errors on correct trials)– we did not find main effects of reward or

efficacy (ps > .444) but did find a significant interaction (b = 1.52, p = .001; Fig. S2A; Table S4),

whereby ERN amplitude on error trials was greatest (i.e., most negative) on trials with low

reward and low efficacy. While this pattern of activity was not predicted (we predicted largest

ERN amplitudes to errors on high reward, high efficacy trials), it is notable that it mirrors the

pattern we observed for error RTs (Fig. S2), where low-reward/low-efficacy trials were

associated with the fastest errors (consistent with effort minimization by, e.g., responding with

any key; b = 29.08, p = .001; Table S6). Speculatively, these findings may reflect a shift from

more effortful proactive to less effortful reactive control as the expected value of proactive

control decreased[45]. Thus, participants may have responded more impulsively, and then

evaluated those responses immediately after they were executed (see Supplementary Results and

Discussion, see also Table S7 for complementary analyses of midfrontal theta).

We found a different pattern of results when examining the feedback-related negativity

(FRN), which typically indexes the difference in feedback-locked activity for trials that resulted

in negative compared to positive feedback[46]. Consistent with previous findings[47-52], we found a

reliable effect of receipt vs omission of reward on FRN amplitude (b = 0.80, p <.001), and this

effect was enhanced for high reward trials (b = 0.81, p = .007; Table S7). However, in addition to

this, and contrary to the hypothesis we preregistered based on previous findings[53,54], we found

that effects of reward receipt vs omission on FRN amplitude were *reduced* for trials with high

efficacy compared to those with low efficacy (b = -0.83, p = .007; Table S8). This efficacy-

related FRN finding might reflect the fact that, under conditions of low efficacy, reward

outcomes are less predictable, thus weakening predictions about forthcoming reward (see

Supplementary Discussion).

280   **Discussion**

Cognitive control is critical but also costly. People must therefore choose what type and

how much control is worth exerting at a given time. Whereas previous studies have highlighted

the critical role of expected rewards when making these decisions, our studies highlight a further

determinant of control value that is equally critical: how efficacious one perceives their efforts to

be (i.e., how much does intensifying control increase their chances of obtaining their reward).

Across two studies, we showed that participants were sensitive to both expected reward and

efficacy when determining how much control to allocate, and therefore performed best when

expecting higher levels of reward and efficacy. Study 2 further demonstrated that both incentive

components increase distinct ERPs, separately related to cue evaluation and proactive control,

providing markers of different stages of incentive processing (evaluation vs. allocation of

control). Collectively, these findings lend support to our theory that participants integrate

information relevant to determining the Expected Value of Control (EVC), and adjust their

control allocation accordingly.

Previous research has shown that people often expend more effort on a task when it

promises greater reward[7-16], but this effort expenditure has its limits. If obtaining that reward also

requires greater effort (i.e., if higher reward is also associated with greater difficulty), the

individual may decide not to invest the effort[55]. Similarly, if difficulty remains constant, but

reward becomes less contingent on effort (i.e., efficacy decreases), the individual may again

decide to divest their efforts[33,34,56,57]. The EVC theory can account for all of these phenomena,

and predicts that expected reward and efficacy will jointly determine how mental effort is

allocated (in the form of cognitive control), and that the underlying evaluation process will

unfold from initial cue evaluation to eventual control allocation. Specifically, the theory predicts

that these incentive components will be processed sequentially over multiple stages that include

initial evaluation of each component, their integration, control allocation, and execution of the

allocated control. Our behavioral and neural findings validate the predictions of this theory:

When participants expected their control to have greater reward and efficacy, we saw increased

neural activity in consecutive ERPs associated with incentive evaluation (P3b) and control

allocation (CNV), followed by increases in control (reflected in improved performance).

Our EEG results extend and clarify previous findings. First, a previous study found (like

us) that the cue-locked P3b is sensitive to the expected reward but not the expected difficulty of

the task to be performed on the upcoming trial[27]. We varied expected efficacy while holding

expected difficulty constant, and show that varying efficacy alone is sufficient to generate

comparable increases in the cue-locked P3b as variability in expected reward. The difference

between our finding and the null result previously observed for task difficulty may be accounted

for by the fact that efficacy (like reward) has a monotonic relationship with motivational

salience, whereas difficulty does not.

Second, our results extend previous studies that linked the CNV with preparatory

attention and proactive control[58-60]. CNV amplitudes scale with a cue's informativeness about an

upcoming task[28,36,37,61,62], temporal expectations about an upcoming target[63-65], and an

individual's predicted confidence in succeeding on the upcoming trial[66]. Critically, CNV

amplitudes also scale with the expected reward for and difficulty of successfully performing an

upcoming task[27,30,32,38], suggesting that this component reflects adjustments to proactive control

in response to motivationally relevant variables. Here, we extend this body of work by showing

that the CNV not only varies with expected efficacy (when isolated from expected difficulty) but

that, unlike the P3b, it reflects an interaction between reward and efficacy (i.e., the expected

payoff for control; Fig. 1), and predicts trial-to-trial variability in performance, suggesting that it

may index the allocation and/or execution of control based on an evaluation of EVC.

With that said, we note that P3b amplitude was also associated with variability in

performance, though to a somewhat lesser degree. It is therefore possible that control allocation

was already being determined at this early stage of the trial but, in conjunction with past

findings[67], it is perhaps equally or more likely that the P3b indexed the initial evaluation of the

motivational relevance of cued incentives. Consistent with this alternate interpretation, we found

that the amplitude of the P3b (but not CNV) decreased over the course of the session, potentially

reflecting decreased attentiveness to the cues.

Our remaining findings provide evidence against alternative interpretations of these

neural results, for instance that they reflect increased arousal or overall engagement throughout

the trial. Pupil diameter, an index of arousal[40,41,68], was larger when participants were expecting

higher reward but also when expecting lower efficacy. Although this pattern was not predicted in

advance, these findings are consistent with evidence that pupil responses more closely track

arousal than proactive control[39]. In contrast, the magnitudes of the P3b and CNV increased

uniformly with reward and efficacy, suggesting that these two ERPs reflect processes related to

proactive control rather than changes in arousal.

Our response- and feedback-related results further suggest that reward and efficacy

specifically increased proactive control, but not reactive control (performance monitoring; [69]) or

overall engagement. Unlike the P3b and CNV, indices of performance monitoring (the ERN and

FRN) were not enhanced with greater reward and efficacy, suggesting that these incentive

conditions were not simply amplifying the motivational salience of errors and reward outcomes.

Rather than reflecting motivational influences on control, the unique patterns of ERN and FRN

amplitudes we observed across conditions may instead provide insight into how participants

formed and updated expectations of performance and outcomes across these different

conditions[70] (see Supplemental Discussion).

Our study builds on past approaches to studying interactions between motivation and

cognitive control[9,12,16,38] by examining changes in effort allocation in response to two incentive

components that are predicted to jointly increase one's motivation. Thus, unlike studies that only

vary the expected reward for an upcoming task, our behavioral and neural findings cannot be

accounted for by general increases in drive, vigor, or arousal. Further, unlike studies that vary the

expected difficulty of an upcoming task, resulting in non-monotonic allocation of effort (the

classic inverted U-shaped function of effort by difficulty; [71,72,73]) the incentive components we

varied should only engender monotonic increases in effort. The monotonic relationship between

these incentive components and the value of control (EVC) can in fact account for the nonlinear

effect of difficulty on effort allocation: at very high levels of difficulty, a given level of control

becomes less and less efficacious. Our study therefore provides the most direct insight yet into

the mechanisms and influences of EVC per se, rather than only some of its components.

One interesting feature of our results is that participants engaged in some reasonably high

level of effort even when their efforts were completely inefficacious (0% efficacy). There are

several plausible explanations for this, including an intrinsic bias towards accuracy (or against

error commission)[74] and potential switch costs associated with the interleaved trial structure[75].

For instance, switch costs associated with control adjustments may discourage a significant drop

in control following a high efficacy trial. An even more intriguing possibility is that experiences

in the real world drive participants to have strong priors that their efforts are generally

efficacious (and practice allocating control within a certain range of expected efficacies)[76],

making it difficult for them to adjust all the way to the expectation that reward is completely

unrelated to their performance on a task.

Individual differences in expectations of efficacy may also play a significant role in

determining one's motivation to engage in individual tasks or effortful behavior at large[19,21,77-79].

Forms of amotivation, like apathy and anhedonia, are common across a variety of psychiatric and

neurological disorders, and most likely reflect deficits in the process of evaluating potential

incentive components, determining the overall EVC of candidate control signals, specifying the

EVC-maximizing control allocation, and/or executing this control. Thus, to understand what

drives adaptive versus suboptimal control, we need to find new and better ways to assess what

drives these key processing stages underlying motivated effort. By highlighting the crucial role

efficacy plays in determining whether control is worthwhile, and identifying candidate neural

signatures of the process by which this is evaluated and integrated into decisions about control

allocation, our studies pave the way towards this goal.

 Method

 Study 1

      Participants. 21 individuals participated in Study 1 (age: M = 21.14, SD = 5.15; 17

 female). Participants gave informed consent and received partial course credits and cash ($5 to

 $10, depending on their performance and task contingencies) for completing the study. The study

 was approved by the university's institutional review board.

      Design and Procedure. We used a within-subject 2 (reward: high, low) x 2 (efficacy:

 high, low) design. On high and low reward Stroop trials, participants saw cues that informed

 them that they would receive $1.00 and $0.10 respectively on the upcoming trial (Fig. 2).

 Reward levels were crossed with efficacy. On high efficacy trials, whether participants were

 rewarded depended entirely on their performance (i.e., fast and accurate responses were always

 rewarded—100% performance-reward contingency). On low efficacy trials, rewards were not

 contingent on participants' performance; instead, rewards were sampled from a rolling window

 (size = 10) of reward rate in high efficacy trials to match reward rates across efficacy levels. This

 approach parallels and builds on recent work examining the influence of performance

 contingency in the domain of motor control (where individuals simply needed to respond quickly

 [56,57]), but importantly our task required participants engage cognitive control in order to be fast

 while also overcoming a prepotent bias to respond based on the color word.

      Participants first completed three practice blocks. In the first practice block (80 trials),

 participants learned the key-color mappings by indicating whether the stimulus XXXXX was

 displayed in red, yellow, green, or blue (using D, F, J, K keys; counterbalanced across

 participants). In the second practice block (16 trials), participants learned to associate cues with

 different reward and efficacy levels (Fig. 2). Finally, participants completed a practice block (64

trials) that resembled the actual task. Then, participants completed the main task. On an

individual trial, cues were presented for 1500 ms, followed by a 250 ms fixation cross, followed

by a target. To increase task difficulty, the response deadline for each trial was 750 ms but

reaction times were recorded as long as a response was made within 1000 ms after Stroop

stimulus onset. Immediately after a response was made, feedback was presented for 750 ms. If a

response was made before 1000 ms, the remaining time was added to the inter-trial-interval, in

which a fixation cross was displayed for 500 ms to 1000 ms. The main task consisted of four

blocks of 75 trials each (except for the first 14 participants, who completed 80 trials per block).

After completing the task, participants completed questionnaires. At the end of the

experiment, ten trials were randomly chosen and participants received a bonus that was the

summed outcomes of those trials.

Study 2

Participants. Before data collection, we conducted a sensitivity analysis, which indicated

that a sample size of $N = 50$ will provide 80% statistical power to detect effect sizes of $d = 0.3$ or

larger. We preregistered our sample size, task design, and analysis plan (osf.io/35akg) and

recruited 53 undergraduate students (age $M = 20.18$, $SD = 2.30$; 15 male; 38 female). We

excluded from all analyses 9 participants who performed poorly on the Stroop task (i.e., below

60% accuracy on high efficacy trials), leaving 44 participants in the final sample. Technical

issues also prevented us from recording clean pupil data from 7 participants in this final sample,

leaving 37 participants in the pupil analyses. Participants gave informed consent and received

partial course credits and cash ($5 to $10, depending on their performance and task

contingencies) for completing the study. The study was approved by the university's institutional

review board.

Design and Procedure. The behavioral paradigm and procedures were similar to those in

Study 1. In addition, we recorded EEG and pupillary responses and changed the following task

parameters: No fixation cross was presented during the cue-target interval to avoid inducing

visual evoked potentials that will influence the amplitude of the CNV; we added a post-response

blank screen (800 ms) to dissociate response evaluation and feedback processing; participants

performed eight blocks of 75 trials each.

EEG Recording and Preprocessing. EEG data were recorded from 32 Ag/AgCl

electrodes embedded in a stretched Lycra cap (Electro-Cap International, Eaton, OH) at a

sampling rate of 512 Hz. Impedances were kept below 5 kΩ during recording. Vertical

electrooculography (VEOG) was recorded from two electrodes placed above and below the right

eye, respectively. Signals were amplified using ANT TMSi Refa8 device (Advanced Neuro

Technology, Enschede, The Netherlands), grounded to the forehead, and referenced online to the

average of all electrodes. Offline, the EEG data were re-referenced to the average of electrodes

placed on the two earlobes. During preprocessing, continuous data were high-pass filtered at 0.1

Hz (12 dB/oct, zero phase-shift Butterworth filter) and decomposed into independent

components using the infomax independent component analysis algorithm implemented in

EEGLAB [80]. We inspected the independent components and used the ICLabel EEGLAB

extension [81] to help identify and remove blink and noise components.

Pre-processed EEG data were epoched relative to the onset of four events: cue (-200 to

1500 ms), stimulus (-200 to 800 ms), response (-200 to 800 ms), and feedback (-200 to 800 ms).

All epochs were baseline-corrected using the mean amplitude before event onset (-200 to 0 ms),

and single-trial baseline activity was included as covariates in the statistical models [82]. Epochs

containing artifacts, with amplitudes exceeding ±150 µV or gradients larger than 50 µV, were

excluded from further analysis. We focused our analyses on these event-related potentials: cue-

locked P3b (250 to 550 ms, averaged across Pz, P3 and P4), cue-locked early and late contingent

negative variations (CNV; early: 700 to 1000 ms; late: 1000 to 1500 ms, each averaged across

Fz, FCz, and Cz), stimulus-locked N2 (100 ms time-window centered on the peak at ~ 318 ms

determined in the grand average ERP), response-locked correct- and error-related negativities

(CRN/ERN; 0 to 100 ms), and feedback-locked feedback-related negativity (FRN; quantified

peak-to-peak at FCz as the difference between the negative peak between 250 to 350 ms and the

positive peak in the preceding 100 ms from the detected peak). For each ERP (except the FRN

that was quantified peak-to-peak), we averaged the amplitudes within the specified time window

separately for each epoch and exported these single-trial values for further analyses in R. All data

analyses were performed using custom MATLAB scripts using EEGLAB functions (cf.[83]).

Pupillometry Recording and Preprocessing. Pupil data were recorded using the

EyeLink 1000 Desktop Mount eye tracker (SR Research, Mississauga, Ontario, CA). The

EyeLink system was configured using a 35-mm lens, 5-point gaze location calibration,

monocular right-eye sampling at a rate of 500 Hz, and centroid fitting for pupil area recordings.

All data processing was performed using custom MATLAB and R scripts. Blink artifacts

detected using the EyeLink blink detection algorithm were removed using linear interpolation

from -200 ms prior to and post-blink onset.

Continuous pupil data were downsampled to 100 Hz and epoched relative to the onset of

four events (-1000 to 5000 ms around event onset): cue, stimulus, response, and feedback. All

epochs were baseline-corrected to the mean amplitude prior to cue onset (-300 to 0 ms). Single-

trial baseline activity (-300 to 0 ms) relative to stimulus, response, and feedback onset were also

included as covariates in the statistical models.

Analysis

Classical frequentist statistical analyses were performed in R. The design was entirely

within-subjects; unless stated otherwise, all estimates and statistics were obtained by fitting

mixed-effects (multilevel) single-trial regression models (two-levels: all factors and

neurophysiological responses for each condition were nested within participants) with random

intercepts and slopes (unstructured covariance matrix) using the R package lme4 [84]. Random

effects were modeled as supported by the data, determined using singular value decomposition,

to avoid overparameterization and model degeneration [85,86]. All continuous regressors were

mean-centered. Two-tailed probability values and degrees of freedom associated with each

statistic was determined using the Satterthwaite approximation implemented in lmerTest.

Behavioral. In both samples, accuracy was analyzed using generalized linear mixed

effects models with a binomial link function. The predictors or regressors were reward, efficacy,

their interaction, congruency, and trial number. Accurate RTs were modeled using linear mixed

effects models and the same predictors. For trial-by-trial predictions of performance with ERPs,

the behavioral models were extended by including P3b and CNV amplitudes z-scored within

participants as predictors.

EEG. Full linear mixed effects models for all ERPs included reward, efficacy, and their

interactions, as well as trial as predictors. For each ERP, we regressed out the baseline activity at

the same electrode sites [82]. In the analyses of CNV, we further controlled for variation in the

preceding P3b amplitude. For ERN analyses, we included as predictors target congruency,

response accuracy, and interactions with incentives. For FRN analyses, we included as predictors

the outcome (whether trials were rewarded or not), and interactions with incentives.

Pupillometry. For each event (cue, target, response, feedback), we fitted a mixed-effects

model to model trial-by-trial pupil responses at every timepoint from -1000 to 5000 ms around

event onset and controlled for error rates with false discovery rate correction [87]. We used this

data-driven mass univariate approach because pupil responses are more protracted than EEG

neural responses and it can be difficult to visually determine the delayed temporal dynamics of

experimental effects. Moreover, unlike ERPs that are similar across tasks, pupil responses vary

significantly across tasks. Predictors in the cue-locked models were reward, efficacy, reward-

efficacy interaction, and trial number; predictors in the stimulus-locked models were reward,

efficacy, congruency, all interactions between reward, efficacy, and congruency, trial number,

and pre-stimulus baseline; predictors in the response-locked models were reward, efficacy,

accuracy, all interactions between reward, efficacy, and accuracy, trial number, pre-response

baseline, and RT; predictors in the feedback-locked models were reward, efficacy, feedback, all

interactions between reward, efficacy, and feedback, trial number, and pre-feedback baseline. For

each event, we plotted the raw pupil time-series alongside the corresponding mixed-effects

model beta coefficients time-series. To avoid visual clutter, we visualized only effects pertinent

to our hypotheses.

Acknowledgments

The authors are grateful to Elizabeth Cory and Aravinth Jebanesan for assistance in data

collection.

Author Contributions

The authors conceived the study jointly. HL collected the data. RF and HL analyzed the data and

wrote the manuscript. All authors edited the manuscript.

Competing Interests

The authors declare no competing interests.

References

1   Inzlicht, M., Shenhav, A. & Olivola, C. Y. The Effort Paradox: Effort Is Both Costly and

Valued. *Trends Cogn Sci* **22**, 337-349, doi:10.1016/j.tics.2018.01.007 (2018).

2   Braver, T. S. *et al.* Mechanisms of motivation-cognition interaction: challenges and

opportunities. *Cogn Affect Behav Neurosci* **14**, 443-472, doi:10.3758/s13415-014-0300-0

(2014).

3   Botvinick, M. M. & Braver, T. Motivation and cognitive control: from behavior to neural

mechanism. *Annual review of psychology* **66**, 83-113, doi:10.1146/annurev-psych-

010814-015044 (2015).

4   Westbrook, A. & Braver, T. S. Cognitive effort: A neuroeconomic approach. *Cogn Affect*

Behav Neurosci **15**, 395-415, doi:10.3758/s13415-015-0334-y (2015).

5   Westbrook, A., Kester, D. & Braver, T. S. What is the subjective cost of cognitive effort?

Load, trait, and aging effects revealed by economic preference. *PLoS One* **8**, e68210,

doi:10.1371/journal.pone.0068210 (2013).

6   Smith, V. L. & Walker, J. M. Monetary Rewards and Decision Cost in Experimental

Economics. *Economic Inquiry* **31**, 245-261, doi:DOI 10.1111/j.1465-

7295.1993.tb00881.x (1993).

7   Kool, W. & Botvinick, M. A labor/leisure tradeoff in cognitive control. *J Exp Psychol*

Gen **143**, 131-141, doi:10.1037/a0031048 (2014).

8   Dixon, M. L. & Christoff, K. The decision to engage cognitive control is driven by

expected reward-value: neural and behavioral evidence. *PLoS One* **7**, e51637,

doi:10.1371/journal.pone.0051637 (2012).

9   Parro, C., Dixon, M. L. & Christoff, K. The neural basis of motivational influences on

cognitive control. *Hum Brain Mapp* **39**, 5097-5111, doi:10.1002/hbm.24348 (2018).

10   Croxson, P. L., Walton, M. E., O'Reilly, J. X., Behrens, T. E. & Rushworth, M. F. Effort-

based cost-benefit valuation and the human brain. *The Journal of neuroscience : the*

official journal of the Society for Neuroscience **29**, 4531-4541,

doi:10.1523/JNEUROSCI.4515-08.2009 (2009).

11   Vassena, E. *et al.* Overlapping neural systems represent cognitive effort and reward

anticipation. *PLoS One* **9**, e91008, doi:10.1371/journal.pone.0091008 (2014).

12   Krebs, R. M., Boehler, C. N., Roberts, K. C., Song, A. W. & Woldorff, M. G. The

involvement of the dopaminergic midbrain and cortico-striatal-thalamic circuits in the

integration of reward prospect and attentional task demands. *Cerebral cortex* **22**, 607-

615, doi:10.1093/cercor/bhr134 (2012).

13   Schmidt, L., Lebreton, M., Clery-Melin, M. L., Daunizeau, J. & Pessiglione, M. Neural

mechanisms underlying motivation of mental versus physical effort. *PLoS Biol* **10**,

e1001266, doi:10.1371/journal.pbio.1001266 (2012).

14   Padmala, S. & Pessoa, L. Reward reduces conflict by enhancing attentional control and

biasing visual cortical processing. *Journal of cognitive neuroscience* **23**, 3419-3432,

doi:10.1162/jocn_a_00011 (2011).

15   Hall-McMaster, S., Muhle-Karbe, P. S., Myers, N. E. & Stokes, M. G. Reward Boosts

Neural Coding of Task Rules to Optimize Cognitive Flexibility. *The Journal of*

neuroscience : the official journal of the Society for Neuroscience **39**, 8549-8561,

doi:10.1523/JNEUROSCI.0631-19.2019 (2019).

16   Yee, D. M., Krug, M. K., Allen, A. Z. & Braver, T. S. Humans Integrate Monetary and

Liquid Incentives to Motivate Cognitive Task Performance. *Front Psychol* **6**, 2037,

doi:10.3389/fpsyg.2015.02037 (2015).

17   Shenhav, A., Botvinick, M. M. & Cohen, J. D. The Expected Value of Control: An

Integrative Theory of Anterior Cingulate Cortex Function. *Neuron* **79**, 217-240,

doi:http://dx.doi.org/10.1016/j.neuron.2013.07.007 (2013).

18   Shenhav, A., Cohen, J. D. & Botvinick, M. M. Dorsal anterior cingulate cortex and the

value of control. *Nature neuroscience* **19**, 1286-1291, doi:10.1038/nn.4384 (2016).

19   Bandura, A. Self-efficacy: Toward a unifying theory of behavioral change. *Advances in*

Behaviour Research and Therapy **1**, 139-161, doi:https://doi.org/10.1016/0146-

6402(78)90002-4 (1978).

20   Vroom, V. H. *Work and motivation.* (Wiley, 1964).

21   Rotter, J. B. Generalized expectancies for internal versus external control of

reinforcement. *Psychological monographs: General and applied* **80**, 1 (1966).

22   Maier, S. F. & Seligman, M. E. Learned helplessness: theory and evidence. *Journal of*

experimental psychology: general **105**, 3 (1976).

23   Feather, N. T. Success probability and choice behavior. *Journal of Experimental*

Psychology **58**, 257 (1959).

24   Feather, N. T. Subjective probability and decision under uncertainty. *Psychological*

Review **66**, 150 (1959).

25   Atkinson, J. W. Motivational determinants of risk-taking behavior. *Psychol Rev* **64, Part**

1, 359-372, doi:10.1037/h0043445 (1957).

26   Wabba, M. A. & House, R. J. Expectancy Theory in Work and Motivation: Some Logical

594   and Methodological Issues. *Human Relations* **27**, 121-147,

595   doi:10.1177/001872677402700202 (1974).

596   27   Schevernels, H., Krebs, R. M., Santens, P., Woldorff, M. G. & Boehler, C. N. Task

597   preparation processes related to reward prediction precede those related to task-difficulty

598   expectation. *Neuroimage* **84**, 639-647, doi:10.1016/j.neuroimage.2013.09.039 (2014).

599   28   Alpay, G., Goerke, M. & Sturmer, B. Precueing imminent conflict does not override

600   sequence-dependent interference adaptation. *Psychol Res* **73**, 803-816,

601   doi:10.1007/s00426-008-0196-9 (2009).

602   29   Botvinick, M. M. & Rosen, Z. B. Anticipation of cognitive demand during decision-

603   making. *Psychol Res* **73**, 835-842, doi:10.1007/s00426-008-0197-8 (2009).

604   30   Frömer, R., Stürmer, B. & Sommer, W. (Don't) Mind the effort: Effects of contextual

605   interference on ERP indicators of motor preparation. *Psychophysiology* **53**, 1577-1586,

606   doi:10.1111/psyp.12703 (2016).

607   31   Strack, G., Kaufmann, C., Kehrer, S., Brandt, S. & Sturmer, B. Anticipatory regulation of

608   action control in a simon task: behavioral, electrophysiological, and FMRI correlates.

609   *Front Psychol* **4**, 47, doi:10.3389/fpsyg.2013.00047 (2013).

610   32   Frömer, R., Hafner, V. & Sommer, W. Aiming for the bull's eye: Preparing for throwing

611   investigated with event-related brain potentials. *Psychophysiology* **49**, 335-344,

612   doi:10.1111/j.1469-8986.2011.01317.x (2012).

613   33   Kukla, A. Foundations of an Attributional Theory of Performance. *Psychological Review*

614   **79**, 454-&, doi:DOI 10.1037/h0033494 (1972).

34   Brehm, J. W. & Self, E. A. The intensity of motivation. *Annual review of psychology* **40**,

109-131, doi:10.1146/annurev.ps.40.020189.000545 (1989).

35   Duncan-Johnson, C. C. & Donchin, E. The P300 component of the event-related brain

potential as an index of information processing. *Biological Psychology* **14**, 1-52,

doi:http://dx.doi.org/10.1016/0301-0511(82)90016-3 (1982).

36   Scheibe, C., Schubert, R., Sommer, W. & Heekeren, H. R. Electrophysiological evidence

for the effect of prior probability on response preparation. *Psychophysiology* **46**, 758-770,

doi:10.1111/j.1469-8986.2009.00825.x (2009).

37   Scheibe, C., Ullsperger, M., Sommer, W. & Heekeren, H. R. Effects of parametrical and

trial-to-trial variation in prior probability processing revealed by simultaneous

electroencephalogram/functional magnetic resonance imaging. *The Journal of*

neuroscience : the official journal of the Society for Neuroscience **30**, 16709-16717,

doi:10.1523/JNEUROSCI.3949-09.2010 (2010).

38   van den Berg, B., Krebs, R. M., Lorist, M. M. & Woldorff, M. G. Utilization of reward-

prospect enhances preparatory attention and reduces stimulus conflict. *Cogn Affect Behav*

Neurosci **14**, 561-577, doi:10.3758/s13415-014-0281-z (2014).

39   Trani, A. & Verhaeghen, P. Foggy windows: Pupillary responses during task preparation.

Quarterly journal of experimental psychology (2006) **71**, 2235-2248,

doi:10.1177/1747021817740856 (2018).

40   Lin, H., Saunders, B., Hutcherson, C. A. & Inzlicht, M. Midfrontal theta and pupil

dilation parametrically track subjective conflict (but also surprise) during intertemporal

choice. *Neuroimage* **172**, 838-852, doi:10.1016/j.neuroimage.2017.10.055 (2018).

41   Bradley, M. M., Miccoli, L., Escrig, M. A. & Lang, P. J. The pupil as a measure of

emotional arousal and autonomic activation. *Psychophysiology* **45**, 602-607,

doi:10.1111/j.1469-8986.2008.00654.x (2008).

42   Falkenstein, M., Hohnsbein, J., Hoormann, J. & Blanke, L. Effects of crossmodal divided

attention on late ERP components. II. Error processing in choice reaction tasks.

Electroencephalogr Clin Neurophysiol **78**, 447-455, doi:10.1016/0013-4694(91)90062-9

(1991).

43   Di Gregorio, F., Maier, M. E. & Steinhauser, M. Errors can elicit an error positivity in the

absence of an error negativity: Evidence for independent systems of human error

monitoring. *Neuroimage* **172**, 427-436, doi:10.1016/j.neuroimage.2018.01.081 (2018).

44   Maruo, Y., Schacht, A., Sommer, W. & Masaki, H. Impacts of motivational valence on

the error-related negativity elicited by full and partial errors. *Biological Psychology* **114**,

108-116, doi:https://doi.org/10.1016/j.biopsycho.2015.12.004 (2016).

45   Braver, T. S. The variable nature of cognitive control: a dual mechanisms framework.

Trends in Cognitive Sciences **16**, 106-113, doi:https://doi.org/10.1016/j.tics.2011.12.010

(2012).

46   Miltner, W. H., Braun, C. H. & Coles, M. G. Event-related brain potentials following

incorrect feedback in a time-estimation task: evidence for a "generic" neural system for

error detection. *Journal of cognitive neuroscience* **9**, 788-798,

doi:10.1162/jocn.1997.9.6.788 (1997).

47   Frömer, R., Nassar, M. R., Stürmer, B., Sommer, W. & Yeung, N. I knew that!

Confidence in outcome prediction and its impact on feedback processing and learning.

bioRxiv, 442822, doi:10.1101/442822 (2018).

48 Fischer, A. G. & Ullsperger, M. Real and fictive outcomes are processed differently but

converge on a common adaptive mechanism. *Neuron* **79**, 1243-1255,

doi:10.1016/j.neuron.2013.07.006 (2013).

49 Frömer, R., Stürmer, B. & Sommer, W. The better, the bigger: The effect of graded

positive performance feedback on the reward positivity. *Biol Psychol* **114**, 61-68,

doi:10.1016/j.biopsycho.2015.12.011 (2016).

50 Luft, C. D., Takase, E. & Bhattacharya, J. Processing graded feedback:

electrophysiological correlates of learning from small and large errors. *Journal of*

cognitive neuroscience **26**, 1180-1193, doi:10.1162/jocn_a_00543 (2014).

51 Meadows, C. C., Gable, P. A., Lohse, K. R. & Miller, M. W. The effects of reward

magnitude on reward processing: An averaged and single trial event-related potential

study. *Biol Psychol* **118**, 154-160, doi:10.1016/j.biopsycho.2016.06.002 (2016).

52 Ulrich, N. & Hewig, J. A miss is as good as a mile? Processing of near and full outcomes

in a gambling paradigm. *Psychophysiology* **51**, 819-823, doi:10.1111/psyp.12232 (2014).

53 Schiffer, A. M., Siletti, K., Waszak, F. & Yeung, N. Adaptive behaviour and feedback

processing integrate experience and instruction in reinforcement learning. *Neuroimage*

146, 626-641, doi:10.1016/j.neuroimage.2016.08.057 (2017).

54 Muhlberger, C., Angus, D. J., Jonas, E., Harmon-Jones, C. & Harmon-Jones, E. Perceived

control increases the reward positivity and stimulus preceding negativity.

Psychophysiology **54**, 310-322, doi:10.1111/psyp.12786 (2017).

55 Kool, W., McGuire, J. T., Rosen, Z. B. & Botvinick, M. M. Decision making and the

avoidance of cognitive demand. *J Exp Psychol Gen* **139**, 665-682, doi:10.1037/a0020198

(2010).

56   Manohar, S. G., Finzi, R. D., Drew, D. & Husain, M. Distinct Motivational Effects of

Contingent and Noncontingent Rewards. *Psychological science* **28**, 1016-1026,

doi:10.1177/0956797617693326 (2017).

57   Kohli, A. *et al.* Using Expectancy Theory to quantitatively dissociate the neural

representation of motivation from its influential factors in the human brain: An fMRI

study. *Neuroimage* **178**, 552-561, doi:10.1016/j.neuroimage.2018.05.021 (2018).

58   van Boxtel, G. J. & Brunia, C. H. Motor and non-motor aspects of slow brain potentials.

Biol Psychol **38**, 37-51 (1994).

59   Brunia, C. H. M., Hackley, S. A., van Boxtel, G. J. M., Kotani, Y. & Ohgami, Y. Waiting

to perceive: Reward or punishment? *Clinical Neurophysiology* **122**, 858-868, doi:DOI:

10.1016/j.clinph.2010.12.039 (2011).

60   Wascher, E., Verleger, R., Jaskowski, P. & Wauschkuhn, B. Preparation for action: An

ERP study about two tasks provoking variability in response speed. *Psychophysiology* **33**,

262-272, doi:10.1111/J.1469-8986.1996.Tb00423.X (1996).

61   Leuthold, H., Sommer, W. & Ulrich, R. Preparing for action: Inferences from CNV and

LRP. *Journal of Psychophysiology* **18**, 77-88, doi:10.1027/0269-880318.2-3.77 (2004).

62   Jentzsch, I., Leuthold, H. & Ridderinkhof, K. R. Beneficial effects of ambiguous precues:

parallel motor preparation or reduced premotoric processing time? *Psychophysiology* **41**,

231-244, doi:10.1111/j.1469-8986.2004.00155.x (2004).

63   Müller-Gethmann, H., Ulrich, R. & Rinkenauer, G. Locus of the effect of temporal

preparation: evidence from the lateralized readiness potential. *Psychophysiology* **40**, 597-

611, doi:10.1111/1469-8986.00061 (2003).

64   Ladanyi, M. & Dubrovsky, B. CNV and time estimation. *Int J Neurosci* **26**, 253-257,

706   doi:doi:10.3109/00207458508985622 (1985).

707   65   Macar, F. & Besson, M. Contingent negative variation in processes of expectancy, motor

708   preparation and time estimation. *Biol Psychol* **21**, 293-307, doi:10.1016/0301-

709   0511(85)90184-x (1985).

710   66   Boldt, A., Schiffer, A.-M., Waszak, F. & Yeung, N. Confidence Predictions Affect

711   Performance Confidence and Neural Preparation in Perceptual Decision Making.

712   *Scientific Reports* **9**, 4031, doi:10.1038/s41598-019-40681-9 (2019).

713   67   Cohen, M. A., Ortego, K., Kyroudis, A. & Pitts, M. Distinguishing the neural correlates

714   of perceptual awareness and post-perceptual processing. *bioRxiv*,

715   2020.2001.2015.908400, doi:10.1101/2020.01.15.908400 (2020).

716   68   Nassar, M. R. *et al.* Rational regulation of learning dynamics by pupil-linked arousal

717   systems. *Nature neuroscience* **15**, 1040-1046, doi:10.1038/nn.3130 (2012).

718   69   Ullsperger, M., Danielmeier, C. & Jocham, G. Neurophysiology of performance

719   monitoring and adaptive behavior. *Physiol Rev* **94**, 35-79,

720   doi:10.1152/physrev.00041.2012 (2014).

721   70   Holroyd, C. B. & Coles, M. G. The neural basis of human error processing:

722   reinforcement learning, dopamine, and the error-related negativity. *Psychol Rev* **109**, 679-

723   709, doi:10.1037/0033-295X.109.4.679 (2002).

724   71   Yerkes, R. M. & Dodson, J. D. The relation of strength of stimulus to rapidity of habit-

725   formation. *Journal of Comparative Neurology and Psychology* **18**, 459-482,

726   doi:10.1002/cne.920180503 (1908).

72   Broadhurst, P. L. The interaction of task difficulty and motivation: The Yerkes-Dodson

law revived. *Acta Psychologica* **16**, 321-338, doi:https://doi.org/10.1016/0001-

6918(59)90105-2 (1959).

73   Cools, R. & D'Esposito, M. Inverted-U–Shaped Dopamine Actions on Human Working

Memory and Cognitive Control. *Biological Psychiatry* **69**, e113-e125,

doi:https://doi.org/10.1016/j.biopsych.2011.03.028 (2011).

74   Hajcak, G. & Foti, D. Errors are aversive: defensive motivation and the error-related

negativity. *Psychological science* **19**, 103-108, doi:10.1111/j.1467-9280.2008.02053.x

(2008).

75   Nigbur, R., Schneider, J., Sommer, W., Dimigen, O. & Stürmer, B. Ad-hoc and context-

dependent adjustments of selective attention in conflict control: An ERP study with visual

probes. *NeuroImage* **107**, 76-84, doi:https://doi.org/10.1016/j.neuroimage.2014.11.052

(2015).

76   Moscarello, J. M. & Hartley, C. A. Agency and the Calibration of Motivated Behavior.

Trends Cogn Sci **21**, 725-735, doi:10.1016/j.tics.2017.06.008 (2017).

77   Grahek, I., Shenhav, A., Musslick, S., Krebs, R. M. & Koster, E. H. W. Motivation and

cognitive control in depression. *Neurosci Biobehav Rev* **102**, 371-381,

doi:10.1016/j.neubiorev.2019.04.011 (2019).

78   Huys, Q. J. M., Daw, N. D. & Dayan, P. Depression: A Decision-Theoretic Analysis.

Annual Review of Neuroscience **38**, 1-23, doi:10.1146/annurev-neuro-071714-033928

(2015).

79   Berwian, I. M. *et al.* Computational Mechanisms of Effort and Reward Decisions in

Patients With Depression and Their Association With Relapse After Antidepressant

Discontinuation. *JAMA Psychiatry*, doi:10.1001/jamapsychiatry.2019.4971 (2020).

80   Delorme, A. & Makeig, S.   (ELSEVIER SCIENCE BV, 2004).

81   Pion-Tonachini, L., Kreutz-Delgado, K. & Makeig, S. ICLabel: An automated

electroencephalographic independent component classifier, dataset, and website.

Neuroimage **198**, 181-197, doi:10.1016/j.neuroimage.2019.05.026 (2019).

82   Alday, P. M. How much baseline correction do we need in ERP research? Extended GLM

model can replace baseline correction while lifting its limits. *Psychophysiology* **56**,

e13451, doi:10.1111/psyp.13451 (2019).

83   Frömer, R., Maier, M. & Abdel Rahman, R. Group-Level EEG-Processing Pipeline for

Flexible Single Trial-Based Analyses Including Linear Mixed Models. *Frontiers in*

Neuroscience **12**, doi:10.3389/fnins.2018.00048 (2018).

84   Bates, D., Maechler, M., Bolker, B. M. & Walker, S. C. Fitting Linear Mixed-Effects

Models Using lme4. *Journal of Statistical Software* **67**, 1-48 (2015).

85   Bates, D., Kliegl, R., Vasishth, S. & Baayen, H. Parsimonious mixed models. *arXiv*

preprint arXiv:1506.04967 (2015).

86   Matuschek, H., Kliegl, R., Vasishth, S., Baayen, H. & Bates, D. Balancing Type I error

and power in linear mixed models. *Journal of Memory and Language* **94**, 305-315,

doi:10.1016/j.jml.2017.01.001 (2017).

87   Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B*

(Methodological) **57**, 289-300 (1995).