Mediterranean oaks harbor more specific soil microbes at the dry end of a precipitation gradient

Gemma Rutten[1,2,3*], Lorena Gómez-Aparicio[2], Beat Frey[1]

[1]Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), Research Unit Forest Soils and Biogeochemistry, 8903 Birmensdorf, Switzerland

[2]Instituto de Recursos Naturales y Agrobiología de Sevilla (IRNAS), CSIC, LINCGlobal. P.O. Box 1052, 41080 Sevilla, Spain

[3]Current address: German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103, Leipzig, Germany. / Martin-Luther-University Halle-Wittenberg, Institute of Biology/Geobotany and Botanical Garden, Am Kirchtor 1, 06108, Halle, Germany

*corresponding author: +49 34555 26199, ggpmrutten@gmail.com

Keywords

plant-soil interactions, precipitation, indicators, *Quercus* L. (oaks), soil microbiomes, gradient, host-specificity, Mediterranean, mycorrhiza, soil pathogens

Abstract

Background: Recent evidence suggests that soil microbial communities can regulate plant community dynamics. In addition, the drought tolerance of plants can be enhanced by soil microbes. So far, few studies have assessed the variation in the microbiome of specific plant species along environmental gradients. Yet understanding these dynamics is essential to improve predictions of plant-soil feedbacks and the consequences of ongoing climate changes. Here we characterized the soil microbiome of two co-occurring Mediterranean oaks along a precipitation gradient, using amplicon sequencing of phylogenetic marker genes for prokaryotes and fungi. Additionally, we identified tree-specific and locally-specific microbes potentially responsible for tree community dynamics.

Results: We show that two co-occurring, evergreen Mediterranean oak species harbor distinct microbiomes along a precipitation gradient. The soil microbial diversity increased along the precipitation gradient, for prokaryotic α and β diversity and for fungal β diversity. *Quercus ilex* harbored richer fungal communities than *Quercus suber*, and host-specific taxa more often belonged to fungi than to prokaryotes. Notably, the microbial communities at the dry end of the precipitation gradient harbored more locally-specific prokaryotic and fungal taxa than the microbial communities with a higher diversity, at the wet end of the gradient, suggesting higher specialization in drier areas.

Conclusions: Even congeneric tree species, belonging to the same functional group, can harbor distinct and specific soil microbiomes. These microbiomes become more similar and consist of more specialized taxa under drier compared with wetter conditions. With this, our study offers a step towards a better understanding of the context-dependency of plant-soil feedbacks by going beyond α and β diversities and focusing on specialized taxa potentially driving community changes along environmental gradients. We hope that our study will stimulate future research assessing the importance of context-dependency of interactions between plants and soil communities in a changing world.

Background

Interactions between plants and microbes affect ecosystem functioning [1–3], and plant-soil feedbacks (PSFs) have been shown to be relevant drivers of forest community dynamics [4,5]. Moreover, specific root-associated microbes have been found to enhance the drought tolerance and overall health of plants [6–8]. Few studies however have explored the variation in plant-microbiome interactions along environmental gradients, yet understanding these dynamics is essential for improving predictions of the consequences of ongoing climate changes.

At the global and regional scales, precipitation is among the most important drivers of soil microbial diversity [9–12], together with other variables like temperature [13,14] and soil pH [15–18]. However, the diverse life history strategies of different taxa can result in numerous responses along environmental gradients [18]. Reduced water availability and increased drought intensity lead to reduced microbial growth and shifts in microbial community composition [19,20]. In particular, decomposers, which contribute to the carbon cycle, have been found to have a lower activity in drier sites than in wetter sites [21]. Food webs with a fungal basis are generally more resistant to drier conditions than bacteria-based food webs [22] but small changes in water availability can result in shifts in fungal species dominance, whereas bacterial communities remain constant [23]. Repeated wet-dry cycles can result in adapted soil communities that show fewer shifts in community composition with fluctuations in water availability [24,25].

Soil microbial communities are not only shaped by abiotic factors such as precipitation and water availability, also biotic drivers such as plants play an important role. Plants can affect microbial communities both directly and indirectly. For example, mycorrhizal fungi and fungal pathogens can accumulate directly in the rhizosphere of the host plant. Moreover,

species-specific litter quality and root exudates might indirectly affect the microbial community through changes in nutrient availability and soil properties [26,27]. Prokaryotes are more likely related to differences in soil properties [28], potentially partly caused by plants, even though they can also be plant species-specific [7,29]. Most existing studies have focused on associations between plants and fungi, as fungi are more host-specific than prokaryotes [30].

Under dry conditions, communities are often dominated by highly specialized species [31] and the interactions among species become more facilitative as predicted by the stress gradient hypothesis [32,33]. For instance, under dry conditions, species-specific microbes can mitigate water stress for their hosts [6,7,29]. Particularly mycorrhiza can help increase the resource uptake of plants and can defend a plant against pathogens [34]. Therefore, both mutualists and pathogens can play an important role in facilitative interactions [35]. Conspecific seedlings experience more positive effects from the soil microbial community when growing in drier than in wetter conditions, suggesting more beneficial effects of the soil community under drier conditions [3,36,37]. However, if specialized mutualists or pathogens are responsible for these shifts along environmental gradients remains largely untested [6]. To disentangle how plant-microbiome interactions vary along climatic gradients, we need to combine the microbial perspective, often focusing on variations in nutrient cycling and decomposition, with the plant perspective, focusing mainly on the mutualistic and antagonistic soil community components.

In this study, we simultaneously tested the effects of precipitation and tree species on soil microbial communities in a Mediterranean climate. Specifically, we used high throughput amplicon sequencing to characterize the prokaryotic and fungal communities under the two dominant evergreen oak species in southern Spain, *Quercus ilex* and *Quercus suber*, along six

sites ranging in precipitation (637, 672, 751, 890, 912, 948 mm year^{-1}). We expected 1) a

reduced microbial community richness and diversity in areas with low precipitation,

particularly for prokaryotes. Additionally, we expected 2) stronger effects of tree species on

fungal than on prokaryotic community diversity. Moreover, we tested 3) the effects of local

variations in soil properties to account for indirect effects of trees on soil microbial

communities. Concerning the specialized soil community components, we expected 4) more

fungal than prokaryotic taxa to be tree-specific. Finally, following the stress gradient

hypothesis, we expected 5) more locally-specific microbes in drier areas and more generalist

in more humid areas. Testing these hypotheses will help to better understand the variation in

plant-microbiome interactions along environmental gradients and ultimately improve

predictions of the consequences of ongoing climate changes.

Results

Diversity of prokaryotes and fungi under two oak species along a precipitation gradient.

The α diversity was higher for prokaryotes than for fungi and differed across our

treatments (Table 1, Figure 1). For prokaryotes, an increased precipitation tended to result in

in more diverse communities, whereas their alpha diversity did not differ among tree species

(Table 1, Figure 1a). For fungi, a higher precipitation did not affect the α diversity, but more

diverse communities were found under *Q. ilex* than under *Q. suber* (Table 1, Figure 1b).

Specifically, the relative abundance of *Ascomycota*, *Basidiomycota*, *Verrucomicrobia* and

Gammaproteobacteria showed significant tree species effects (Table S1, Figure S1), whereas

Mortierellomycota, *Acidobacteria* and *Patescibacteria* showed significant interactions

between tree species and the precipitation gradient (Table S1, Figure S1).

The β diversity of fungal and prokaryotic communities changed along the precipitation

gradient (Table 2), with both prokaryotic (Figure 1c) and fungal (Figure 1d) communities becoming more dissimilar with increasing precipitation. Fungal communities were generally more variable whereas prokaryotic communities showed more overlap along the precipitation gradient and between tree species (Figure 1, Figure 2, Figure S2). Additionally, significant interactions between precipitation and tree species for prokaryotic (P x T interaction, P = 0.05, Table 2) and fungal communities (P x T interaction P = <0.001, Table 2), indicate that the dissimilarity of the soil microbiome under the two tree species is determined by the precipitation regime. Moreover, a significant tree species effect on the β diversity of fungi (P = <0.001, Table 2) indicates that the fungal communities under *Q. ilex* were less dissimilar than those under *Q. suber* (Table2; Figure 1).

The most influential edaphic conditions on prokaryotic and fungal communities were calcium (explaining 57% of prokaryotic structure and 60% of fungal structure), followed by pH (50% and 53%), and nitrate (24% and 45%), potassium (19% and 24%) and finally phosphorus (10% and 12, Table 3). Additionally, sand and ammonium explained 29% and 19% of the variation in prokaryotic communities, respectively (Table 3, Figure 2), whereas these variables did not affect fungal communities. Conversely, magnesium and SOM explained 23% and 15 % of the variation in fungal communities, respectively, whereas these variables did not impact the prokaryotic communities (Table 3, Figure 2).

Specialization of prokaryotes and fungi under two oak species along a precipitation gradient.

There was a clear difference in the taxa frequency distribution between the prokaryotic and fungal communities (Figure 3). Only a small proportion of taxa was found in all samples (Figure 3), namely 2.51% (115 OTUs) for prokaryotes and 0.65% (20 OTUs) for fungi respectively. These ubiquitous fungi taxa, included nine saprotrophs, six unassigned guilds, two animal pathogens, two plant pathogens (*Mycosphaerella tassiana* and *Trichoderma*

asperelluman) and one endophyte (*Clonostachys rosea*). In terms of abundance, these

ubiquitous OTUs accounted for 27% and 18% of the total reads for prokaryotes and fungi

respectively. Particularly the peaks in distribution of OTUs over the samples differed between

the prokaryotic and fungal community components. The prokaryotic distribution peaked

around 8 samples (>200 OTUs), corresponding to the number of samples per site and

suggesting an important role for locally-specific taxa in prokaryotic communities. On the

hand, the fungal distribution peaked around 4 samples (>300 OTUs) and exponentially

declined with occurrence, suggesting a role for both locally-specific and tree-specific taxa in

the fungal communities.

Both tree-specific prokaryotic and fungal taxa were found using logfold^2 changes in

abundance under *Q. ilex* versus *Q. suber* trees, independent of precipitation regime. Within

prokaryotes, 61 taxa belonging to unique 44 families were significantly associated to one of

the tree species (21 taxa to *Q. ilex,* 40 taxa to *Q. suber;* Figure 4a). Of those taxa, 23%

belonged to *Chloroflexi,* 11% to *Alphaproteobacteria, Planctomycetes* and *Patescibacteria,*

10% to *Actinobacteria* and *Gammaproteobacteria,* 7% to *Verrucomicrobia,* and 2-3 % to

each of the remaining phyla (*Firmicutes, Gemmatimonadetes*, *Cyanobacteria, Euryarchaeota,*

Acidobacteria, WPS-2, *Bacteroidetes*). Even more fungi (126 taxa), belonging to 62 unique

genera, were significantly associated to a particular tree species (72 taxa to *Q. ilex,* 54 taxa to

Q. suber; Figure 4b). Of those taxa, 71% belonged to *Ascomycota*, 20% to *Basidiomycota*, 4%

to *Chytridiomycota*, and 1% to each of the remaining phyla (*Mucoromycota*, *Rozellomycota*

Monoblepharomycota, Mortierellomycota). Another 2% remained unclassified. The OTUs

that could be classified to the genus level (75 of 126) mostly belonged to unique families and

genera. Except within the *Ascomycota*, many families (*Herpotrichiellaceae, Coniochaetaceae*

Dothioraceae,   *Helotiaceae,   Myxotrichaceae,   Nectriaceae,   Ophiostomataceae,*

Pyronemataceae, Sympoventuriaceae, Teratosphaeriaceae, Trichocomaceae) harbored

multiple tree-associated taxa, including mainly saprotrophic taxa. Within the Basidiomycota,

four families (*Bulleraceae, Cortinariaceae, Russulaceae, Thelephoraceae*) harbored multiple

tree-specific taxa, three of which were ectomycorrhizal. *Russulaceae* included the genera

Gymnomyces and *Russula*. The *Thelephoraceae*, included *Pseudotomentella* and *Tomentella*,

and the *Cortinariaceae* harbored more than one taxon (*Cortinarius* 4 OTUs). Finally, we

found one family within the *Chytridiomycota* (*Spizellomycetaceae*), that harbored the tree-

specific pathogenic genera (*Kochiomyces* (1 OTU) and *Spizellomyces* (2 OTUs).

Locally-specific taxa, defined as taxa which were more abundant than expected based

on random distributions over sites [38], were found in both prokaryotic and fungal

communities (Figure S3). The proportion of locally-specific taxa for each site decreased along

the precipitation gradient (prokaryotes and fungi analyzed together: $F_{1,8}$=8.26, R^2_{adj}=0.32,

P=0.021, Figure 5), indicating higher specialization rates for prokaryotes and fungi occurring

at the drier end of the precipitation gradient.

Discussion

Microbial community diversity under two oak species change along a precipitation gradient

Climate and precipitation in particular, can enhance overall prokaryotic and fungal

diversity [12] and biomass [11]. In line with this, most microbial diversity measures showed

positive trends along the precipitation gradient. However, fungal α diversity (richness) was

not affected by precipitation, suggesting that precipitation has stronger effects on prokaryotes

than fungi. Such differences in bacterial and fungal communities along precipitation

gradients, suggest that fungi are more resistant to drier conditions than bacteria [39,40],

however also opposing trends have been reported. Such differences could potentially be

explained by the copiotroph–oligotroph trade-off, also known as r- and K-selection theory

[17]. For example, Proteobacteria and Actinobacteria tended to be more ubiquitous in the

nutrient-rich soils in the wetter soils. On the contrary, Acidobacteria are known to be

ubiquitous, diverse, desiccation tolerant, and largely oligotrophic bacteria, abundant under

low resource availability [41]. Yet, it is known that subgroups within a phylum might show

different responses than the one observed at the phylum level, potentially leading to

misinterpretations. For example, different subgroups in Proteobacteria and Actinobacteria

might occupy different ecological niches and can show mixed responses to precipitation [20].

Former studies find fewer oligotrophs and more copiotrophs with increasing precipitation

[17,20]. In our study, the relative abundances of the most abundant fungal phyla like

Ascomycota and *Basidiomycota* did not vary significantly along the precipitation gradient.

However, the relative abundance of the prokaryotic *Acidobacteria* increased along the

precipitation gradient. Whereas changes in the relative abundances of *Patescibacteria* and

Mortierellomycota along the precipitation gradient depended on tree species. Such opposing

patterns between taxonomic groups have been reported before [10,42]. Notably, our results

are in line with those from earlier studies focusing on mycorrhizal community changes in

oaks across sites varying in water availability. For example, a long-term precipitation

reduction experiment resulted in modified EMF communities under *Q. ilex* [43], and the

mycorrhizal status of *Quercus agrifolia* shifted with soil moisture from predominantly EMF

in wet to almost exclusively arbuscular mycorrhizal fungi (AMF) during severe droughts [44].

Here, we show that besides ectomycorrhizal fungi, also prokaryotic and fungal communities

under several specific plant species vary along a precipitation gradient.

Host plant family and functional group determined the community composition of

ectomycorrhizal fungi (EMF) in a global analysis [45] and in dry areas [46]. In Mediterranean

ecosystems and temperate forests, even lower taxonomic levels (plant species identity) drove

fungal richness and fungal community structure [47,48]. Despite the close phylogenetic

relationship between the *Quercus* species, each species harbored distinct soil microbial

communities. Moreover, we show that even within the functional group of evergreen oaks,

differences in the relative abundance of *Basidiomycota* (which includes the EMF-rich

Agaricomycetes class) can be found. This adds to studies comparing the EMF communities of

evergreen and deciduous oaks [27,49]. It would be interesting to conduct further studies

including a larger pool of congeneric tree species to better understand what root or leaf traits

might drive this variation in soil communities. Particularly because it has been shown that

such variation in the soil microbiome can affect the performance of seedlings growing under

congeneric trees [36].

The setup of our study sites was intended to test for differences in the soil microbiome

between oak species along a precipitation gradient, but it also included a natural variation in

soil properties, possibly partly caused by the different tree species (e.g. litter input and root

exudates). The majority of earlier studies along environmental gradients have acknowledged

the importance of additional or indirect factors (e.g. edaphic conditions, soil properties and

plant communities) in structuring the soil microbiome [10–12]. Our study confirmed this, and

we found that soil calcium and pH were highly correlated and important variables for

explaining variation in both prokaryotic and fungal communities. At the global scale, these

same two variables emerged as important predictors of fungal richness [10], but pH did not

drive patterns in prokaryotic communities in drylands [12]. Future studies and experiments

are needed to better understand the separate effects of calcium and pH on soil geochemical

processes.

In line with our expectations, edaphic conditions affected prokaryotes more than

fungi. Notably, only prokaryotes were affected by variations in soil texture and ammonium,

whereas only fungi were impacted by changes in soil organic matter content. Whereas fungi

can spread their mycelium around, attaching to resource patches and plant roots, prokaryotes

are more constrained to the soil matrix and pores, which makes it more likely for the latter to

be affected by soil chemistry [17,28]. Taken together with differences in litter inputs and root

exudates among trees [26,27], this finding might also explain why we found fungal diversity,

in terms of richness and community structure, to be more related to tree species than

prokaryotic communities. Related to this, with increasing precipitation we found greater

phosphorus concentrations under *Q. suber*, whereas calcium concentrations decreased under

Q. ilex along the precipitation gradient (Table S3). Long-term field experiments designed to

separate precipitation from other tree-related processes, including litter input and root

exudation, would be particularly valuable for exploring possible explanations in more detail.

Specialized prokaryotes and fungi under two oak species along a precipitation gradient

Unravelling the individual contributions of soil microbial taxa facilitates an improved

understanding of the role that interactions between plants and soil microbes play in shaping

the responses of plant community composition and ecosystem processes to global

environmental changes [50]. Soil microbial components including enemies, symbionts and

decomposers, may show different responses resulting in different net magnitudes and

directions of plant–soil feedbacks along various environmental gradients [3,6,51], . Plant-soil

feedbacks require specific variations in the soil communities of a host species, leading to

asymmetric fitness differences in subsequent plant growth [52]. Our study provides evidence

for such mechanisms as we found proportion of ubiquitous taxa in the fungal communities

(0.65%) as well as in the prokaryotic communities (2.46%), suggesting that not all microbes

occur everywhere, but implying a certain degree of specialization, particularly within the

fungal communities. Particularly, we found more fungal taxa (173 OTUs) than prokaryotic taxa (40 OTUs) associated with a specific tree species. None of the fungal genera harbored specialized taxa under both tree species, except for the saprotrophic genera *Cladophialophora* and *Cryptococcus* and the plant pathogens *Spizellomyces*. Even on a family level, we only identified two families that harbored plant-specific taxa for both oak species. First, some ectomycorrhizas within *Russulaceae* (genus *Lactarius* and *Gymnomyces*) were specifically associated with *Q. suber,* whereas others from the confamiliar genus *Russula* were associated with *Q. ilex*. Second, in the family *Spizellomycetaceae* (*Chytridiomycota*), the plant pathogens *Kochiomyces* and an unclassified genus were associated with *Q. ilex*, whereas the confamiliar *Spizellomyces* was associated with *Q. suber*. These results suggest that a fraction of the soil microbiome harbors associated fungi, including mycorrhiza, saprotrophs and pathotrophs that are specific to a plant species, even when the plants are closely related.

The effects of abiotic conditions on interactions between plants and their soil microbes remain unclear. For example, a reduced water availability has been shown to result in smaller net negative effects of the soil microbial community on conspecific plant performance [36,37]. This suggests either a decrease in specific antagonistic interactions, an increase in mutualistic interactions or a combination of both [51–53]. Most studies so far have generally found negative effects of the conspecific soil community on subsequent plant growth [2,52]. A test with savanna tree species found positive heterospecific feedbacks suggesting an important role for mutualists in drier environments [53]. Droughts can reducing plant-microbe associations and neutralize plant soil feedbacks [54] and inconsistent abiotic conditions between training phase and feedback phase resulted in variable feedback effects [55]. We found that microbial communities harbored more locally-specific taxa in the drier sites than in the wetter sites, and particularly locally-specific prokaryotes decreased with increasing

precipitation. The occurrence of more specialized taxa at the dry end of our gradient, could be

driven by facilitation, whereas more generalist taxa in more humid areas, are possibly driven

by competition [3,36]. A similar pattern was observed for the locally-specific fungal taxa

along the precipitation gradient, although no significant relationship was detected. It is

possible that the difference between prokaryotic and fungal communities was found because

fungal communities are more buffered by host plants whereas bacteria are more strongly

dependent on edaphic conditions [17,28]. An additional explanation could be that fungi are

more resistant to drier conditions than bacteria [22], reinforcing our first hypothesis. Our

study serves as a proof of concept, which to our knowledge has not been demonstrated before.

It is bringing us one step closer to identify potential crucial habitat-specific prokaryotes and

fungi of particular ecological importance for plant-soil interactions along gradients.

Conclusion

In the last years, several studies have discussed the climate-dependency of microbial

communities, their interactions with and feedbacks on plants [55–57]. Here we show that

plants can harbor distinct associated microbes when climatic conditions change. Particularly,

drier conditions resulted in a higher proportion of locally-specific taxa as compared with soil

communities at the wetter end of the precipitation gradient. Moreover, we show that besides

frequently studied fungal associations such as mycorrhiza, also prokaryotes associate with

tree species and might therefore potentially play a prominent role generating plant-soil

feedbacks.

Our approach highlights taxa in the soil microbial community which are potentially

important in generating precipitation depended differences between host plant species. Such

approach goes beyond alpha and beta diversity measures of the microbiome and is not limited

to taxonomic or functional knowledge of the taxa. This can help to discover microbial taxa

which are important in generating context-dependent plant-soil feedbacks, which in turn help

to better understand underlying mechanisms of ecosystem functioning and vegetation

dynamics under global change.

Methods

Experimental design and site selection

We used the third Spanish National Forest Inventory (NFI3; Ministry of Environment, Madrid

2006) to select study sites where the two most abundant evergreen Mediterranean oak species

in the country (*Quercus suber* and *Quercus ilex*) co-occur. In November 2015, we selected six

sites covering a precipitation gradient (637, 672, 751, 890, 912, 948 mm year^{-1}) in south-

central Spain [36]. The sites were located between 100 and 594 m a.s.l and had mean annual

temperatures between 15°C and 18°C. Each of the six sites contained four *Q. ilex* and four *Q.*

suber adult trees (48 individual trees in total), which had no woody understory or overlapping

crowns. In January 2016, we collected soil samples (0-15 cm depth) from under the crown of

each tree using a soil corer (5 cm diameter). We pooled 9 to 15 cores to generate one

composite soil sample per tree. The 48 pooled soil samples were kept in a cooling container

until further processing in the lab.

Physical and chemical soil properties

In the lab, the soil samples were homogenized, sieved through a 2 mm sieve to remove

roots and mesofauna, and split into two parts. One part was air dried for three days 25°C and

subsequently analyzed for soil physical and chemical properties according to standardized

protocols [58]. As we could not separate indirect tree species effects, through leaf litter input,

nutrient depletion and root exudation, from inherent site differences, we assessed physical and

chemical soil properties as a whole. In short, we assessed soil texture, pH, total soil organic

matter (SOM) content, and concentrations of nitrate (NO_3^-), ammonium (NH_4^+), available

phosphorus (P), potassium (K), calcium (Ca) and magnesium (Mg). For additional

information see Supplement 1.

DNA extraction, PCR amplification and sequencing

We used the second part of the soil sample (0.25 g) to extract DNA using the

PowerSoil isolation kit (Mo-Bio Laboratories Inc., Carlsbad, CA, USA) according to the

manufacturer's instructions. Extracted DNA was quantified using Picogreen (Molecular

Probes, Eugene, OR, USA). We amplified the partial prokaryotic small-subunit ribosomal

RNA genes (region V3–V4 of 16S rRNA gene) and the fungal ribosomal internal transcribed

spacers (region ITS2), as described previously [59]. Each sample containing 40 ng of DNA

was amplified in triplicate and pooled before purification with Agencourt AMPure XP beads

(Beckman Colter, Berea, CA, USA). The amplicons were sent to the Genome Quebec

Innovation Centre (Montreal, Canada) for barcoding using Fluidigm Access Array technology

and paired-end sequencing on the Illumina MiSeq v3 platform (Illumina Inc., San Diego, CA,

USA).

Bioinformatics

For the quality control of prokaryotic and fungal reads, we used a customized pipeline

[59]. Briefly, we matched paired-end reads [60] correcting for substitution errors [61], and

trimmed the PCR primers, with one mismatch allowed [62]. We discarded prokaryotic

sequences (16S rRNA gene, region V3V4) shorter than 300 bp, fungal sequences (region

ITS2) shorter than 200 bp, and sequences without matching primers. The fastq_filter function

in USEARCH was used for quality filtering of the trimmed reads with a maximum expected

error threshold of one. All singletons were removed to reduce artificial OTU inflation while

clustering [63]. We used the cluster_otu function in USEARCH, which includes an 'on-the-

fly' chimera removal algorithm to cluster the sequences in OTUs applying a 97% similarity

threshold. Metaxa2 [64] and ITSx [65] were used to test the centroid OTU sequences for the

presence of ribosomal signatures, and unsupported centroid sequences were discarded.

Remaining reads that passed the quality filtering, were mapped with USEARCH

usearch_global algorithm (parameters: maxrejects 0, maxaccepts 0 and top_hit_only) against

the final OTU list. All raw sequences were deposited in the NCBI Sequence Read Archive

under the BioProject accession number PRJNA550539. We used MOTHUR with its

implemented naïve Bayesian classifier (minimum bootstrap support 60%) to query the

centroid sequences to the reference databases. SILVA Release 128 served as the reference

database for the prokaryotic sequences [66]. Eukaryotic sequences were queried against the

two databases in NCBI GenBank [67] and UNITE 8.0 [68]. We removed eukaryote sequences

(0.6%) and unknown sequences (1.4%) from the prokaryotic OTUs, as well as OTUs not

belonging to the fungal kingdom, including plants (2.2%), protists (0.4%) and unclassified

eukaryotes (0.9%). To increase the probability of assessing ecologically relevant taxa [69], we

pruned the prokaryotic and fungal datasets of rare OTUs (total abundance <10 reads;

occurring >3 samples).

Statistical analysis

To test for differences in alpha diversity between the oak species along the precipitation

gradient we first made rarefaction curves. These curves visualize sampling effort and help

determine the sample with the smallest number of reads (Figure S1). To correct for increasing

species numbers with an increasing number of reads, we rarefied the data by the smallest total

number of sequences per sample, resulting in 15,548 prokaryotic and 12,740 fungal reads per

sample. This step led to the removal of 10 fungal OTUs from the data. Of the many indexes

available to quantify alpha diversity, we chose to calculate the observed richness (S_{obs}), as it is

widely used and easily comparable between studies. We calculated other alpha diversity

measures for each sample, for both raw and rarified datasets [69], and found small differences

depending on the measure used (data not shown). To test for the effects of precipitation and

tree species on the diversity indexes, we used generalized linear mixed effects models

(GLMM) using *lme4* [70], and included site as a random factor to control for multiple

samples coming from the same site. In addition, we calculated the relative abundance of OTU

counts for each phylum (with a total abundance >1%) and for each treatment combination.

Finally, we assessed the effects of precipitation and tree species on the relative abundance of

the microbial sequences, using PERMANOVAs based on Euclidian distances with 9999

permutations, using *vegan* [71]. To account for site effects, we included this factor in the

random effects (strata).

To quantify differences in community structure (beta diversity) across the precipitation

gradient and between tree species, we used PERMANOVAs based on Bray-Curtis distances

with 9999 permutations, with site included in the strata. The Bray-Curtis dissimilarity

matrices were generated based on either relative or normalized OTU sequence counts of the

unrarefied data. To normalize the data, we performed a transformation by the weighted

trimmed mean of M-values [72,73]. Applying alternative transformations [69,74,75] yielded

similar results and are therefore only presented in the supplementary (Table S1). Differences

in the community structure among sites and between tree species were visualized with

unconstrained PCoA-ordinations and with site- and tree-species constrained CAP-ordinations

[71]. To determine which edaphic variables (pH, SOM, NO_3^-, NH_{4+}, P, K, Ca, Mg and

proportion of sand (as a measure of soil texture) significantly explained variation in the

prokaryotic and fungal communities, we used non-metric multidimensional scaling (NMDS)

in *vegan* [71].

To identify which OTUs led to changes in multivariate patterns between tree species

and along the precipitation gradient, we studied the relationship between OTUs and groups of

sites using indicator species analysis [76,77]. First, we investigated which taxa were

significantly associated to a tree species, hereafter referred to as species-specific taxa. We

combined an indicator species analysis with GLMMs and log-likelihood-ratio tests

implemented in the *EdgeR:glmLRT* and *EdgeR:topTag* [72,73]. To explore if more locally-

specific taxa exist in drier sites, we tested which taxa in the prokaryotic and the fungal

communities were more abundant than expected based on random distributions over the sites

[38]. Based on this analysis, we calculated the proportion of locally-specific taxa for each site

along the precipitation gradient, for prokaryotes and fungi separately. We then used a linear

model, with precipitation and tree species as fixed factors, to test whether the proportion of

locally-specific taxa decreased along the precipitation gradient. All statistics were performed

in R [78] using the cited packages, functions and their dependencies.

List of abbreviations

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

The datasets generated and/or analyzed during the current study are available in the NCBI
Sequence Read Archive repository, under the BioProject accession number PRJNA550539
[www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA550539]

Competing interests

The authors declare that they have no competing interests

Funding

This study was supported by the Swiss National Science Foundation (SNSF) in the context of

a mobility fellowship granted to GR (P2BEP3_162092). LGA acknowledges support from the

MICINN project INTERCAPA (CGL-2014-56739-R).

Authors' contributions

The study was initiated and performed by GR, who gained funding for the study. Laboratory

analyses were facilitated by LGA and BF. BF conducted the bioinformatics. GR analyzed the

resulting data and wrote the first draft, with the help of LGA and BF. All authors contributed

to revisions of the manuscript.

Acknowledgements

We thank P. Ruiz-Benito and L. Matías for help with third national forest inventory (NFI3)

and field site selection. We are grateful to E. Gutiérrez-González and B. Stierli for their help

in the laboratory, J. Domiguez, M. Renner, C. Saladin and A. Locher for help in the field, the

Genetic Diversity Center (GDC) at ETH Zürich and the McGill University and Génome

Québec Innovation Center in Montreal, Canada for performing Illumina MiSeq sequencing.

M. Dawes for textual improvements and A. Frossard and T. Wubet for useful comments on

earlier versions of the manuscript.

References

1. Schnitzer SA, Klironomos JN, HilleRisLambers J, Kinkel LL, Reich PB, Xiao K, et al. Soil
microbes drive the classic plant diversity–productivity pattern. Ecology. Wiley Online
Library; 2011;92:296–303.

2. Van der Putten WH, Bardgett RD, Bever JD, Bezemer TM, Casper BB, Fukami T, et al.
Plant–soil feedbacks: the past, the present and future challenges. J Ecol. Wiley Online
Library; 2013;101:265–76.

3. Lekberg Y, Bever JD, Bunn RA, Callaway RM, Hart MM, Kivlin SN, et al. Relative
importance of competition and plant–soil feedback, their synergy, context dependency and
implications for coexistence. Ecol Lett. Wiley Online Library; 2018;21:1268–81.

4. McCarthy☐Neumann S, Kobe RK. Conspecific and heterospecific plant–soil feedbacks
influence survivorship and growth of temperate tree seedlings. J Ecol. Wiley Online Library;
2010;98:408–18.

5. Bennett JA, Maherali H, Reinhart KO, Lekberg Y, Hart MM, Klironomos J. Plant-soil
feedbacks and mycorrhizal type influence temperate forest population dynamics. Science (80-
). American Association for the Advancement of Science; 2017;355:181–4.

6. Classen AT, Sundqvist MK, Henning JA, Newman GS, Moore JAM, Cregger MA, et al.
Direct and indirect effects of climate change on soil microbial and soil microbial☐plant
interactions: What lies ahead? Ecosphere. Wiley Online Library; 2015;6:1–21.

7. Fitzpatrick CR, Copeland J, Wang PW, Guttman DS, Kotanen PM, Johnson MTJ.
Assembly and ecological function of the root microbiome across angiosperm plant species.
Proc Natl Acad Sci. National Acad Sciences; 2018;115:E1157--E1165.

8. Delavaux CS, Smith☐Ramesh LM, Kuebbing SE. Beyond nutrients: a meta☐analysis of
the diverse effects of arbuscular mycorrhizal fungi on plants and soils. Ecology.
2017;98:2111–9.

9. Angel R, Soares MIM, Ungar ED, Gillor O. Biogeography of soil archaea and bacteria
along a steep precipitation gradient. ISME J. Nature Publishing Group; 2010;4:553.

10. Tedersoo L, Bahram M, Põlme S, Kõljalg U, Yorou NS, Wijesundera R, et al. Global
diversity and geography of soil fungi. Science (80-). American Association for the
Advancement of Science; 2014;346:1256688.

11. Chen D, Mi J, Chu P, Cheng J, Zhang L, Pan Q, et al. Patterns and drivers of soil
microbial communities along a precipitation gradient on the Mongolian Plateau. Landsc Ecol.
Springer; 2015;30:1669–82.

12. Maestre FT, Delgado-Baquerizo M, Jeffries TC, Eldridge DJ, Ochoa V, Gozalo B, et al.
Increasing aridity reduces soil microbial diversity and abundance in global drylands. Proc
Natl Acad Sci. National Acad Sciences; 2015;112:15684–9.

13. Zhou J, Deng Y, Shen L, Wen C, Yan Q, Ning D, et al. Temperature mediates
continental-scale diversity of microbes in forest soils. Nat Commun. Nature Publishing
Group; 2016;7:12083.

14. Větrovský T, Kohout P, Kopecký M, Machac A, Man M, Bahnmann BD, et al. A meta-
analysis of global fungal distribution reveals climate-driven patterns. Nat Commun. Nature
Publishing Group; 2019;10:1–9.

15. Rousk J, Bååth E, Brookes PC, Lauber CL, Lozupone C, Caporaso JG, et al. Soil bacterial
and fungal communities across a pH gradient in an arable soil. ISME J. Nature Publishing
Group; 2010;4:1340.

16. Lanzén A, Epelde L, Garbisu C, Anza M, Martín-Sánchez I, Blanco F, et al. The
community structures of prokaryotes and fungi in mountain pasture soils are highly correlated
and primarily influenced by pH. Front Microbiol. Frontiers; 2015;6:1321.

17. Fierer N. Embracing the unknown: disentangling the complexities of the soil microbiome.
Nat Rev Microbiol [Internet]. Nature Publishing Group, a division of Macmillan Publishers
Limited. All Rights Reserved.; 2017;15:579. Available from:
http://dx.doi.org/10.1038/nrmicro.2017.87

18. Peay KG, von Sperber C, Cardarelli E, Toju H, Francis CA, Chadwick OA, et al.
Convergence and contrast in the community structure of Bacteria, Fungi and Archaea along a
tropical elevation–climate gradient. FEMS Microbiol Ecol. Oxford University Press; 2017;93.

19. Felsmann K, Baudis M, Gimbel K, Kayler ZE, Ellerbrock R, Bruehlheide H, et al. Soil
bacterial community structure responses to precipitation reduction and forest management in
forest ecosystems across Germany. PLoS One. Public Library of Science; 2015;10:e0122539.

20. Hartmann M, Brunner I, Hagedorn F, Bardgett RD, Stierli B, Herzog C, et al. A decade of
irrigation transforms the soil microbiome of a semi☐arid pine forest. Mol Ecol. Wiley Online
Library; 2017;26:1190–206.

21. Yuste JC, Penuelas J, Estiarte M, Garcia☐Mas J, Mattana S, Ogaya R, et al.
Drought☐resistant fungi control soil organic matter decomposition and its response to
temperature. Glob Chang Biol. Wiley Online Library; 2011;17:1475–86.

22. de Vries FT, Griffiths RI, Bailey M, Craig H, Girlanda M, Gweon HS, et al. Soil bacterial
networks are less stable under drought than fungal networks. Nat Commun. Nature Publishing
Group; 2018;9:3033.

23. Kaisermann A, Maron PA, Beaumelle L, Lata JC. Fungal communities are more sensitive
indicators to non-extreme soil moisture variations than bacterial communities. Appl Soil Ecol.
Elsevier; 2015;86:158–64.

24. Evans SE, Wallenstein MD. Soil microbial community response to drying and rewetting
stress: does historical precipitation regime matter? Biogeochemistry. Springer; 2012;109:101–
16.

25. Lau JA, Lennon JT. Rapid responses of soil microorganisms improve plant fitness in
novel environments. Proc Natl Acad Sci [Internet]. 2012;109:14058–62. Available from:
http://www.pnas.org/cgi/doi/10.1073/pnas.1202319109

26. Wardle DA, Bardgett RD, Klironomos JN, Setälä H, Van Der Putten WH, Wall DH.
Ecological linkages between aboveground and belowground biota. Science (80-). American
Association for the Advancement of Science; 2004;304:1629–33.

27. Aponte C, García L V, Marañón T, Gardes M. Indirect host effect on ectomycorrhizal
fungi: Leaf fall and litter quality explain changes in fungal communities on the roots of co-
occurring Mediterranean oaks. Soil Biol Biochem. Elsevier; 2010;42:788–96.

28. Navrátilová D, Tláskalová P, Kohout P, Dřevojan P, Fajmon K, Chytrý M, et al. Diversity
of fungi and bacteria in species-rich grasslands increases with plant diversity in shoots but not
in roots and soil. FEMS Microbiol Ecol. Oxford University Press; 2018;95:fiy208.

29. Compant S, Van Der Heijden MGA, Sessitsch A. Climate change effects on beneficial
plant–microorganism interactions. FEMS Microbiol Ecol. Blackwell Publishing Ltd Oxford,
UK; 2010;73:197–214.

30. Urbanová M, Šnajdr J, Baldrian P. Composition of fungal and bacterial communities in
forest litter and soil is largely determined by dominant trees. Soil Biol Biochem. Elsevier;
2015;84:53–64.

31. Berdugo M, Maestre FT, Kéfi S, Gross N, Le Bagousse Pinguet Y, Soliveres S. Aridity
preferences alter the relative importance of abiotic and biotic drivers on plant species
abundance in global drylands. J Ecol. Wiley Online Library; 2019;107:190–202.

32. He Q, Bertness MD, Altieri AH. Global shifts towards positive species interactions with
increasing environmental stress. Ecol Lett. 2013;16:695–706.

33. Li H, Colica G, Wu P, Li D, Rossi F, De Philippis R, et al. Shifting Species Interaction in
Soil Microbial Community and Its Influence on Ecosystem Functions Modulating. Microb
Ecol [Internet]. 2013;65:700–8. Available from: https://doi.org/10.1007/s00248-012-0171-2

34. Lambers H, Albornoz F, Kotula L, Laliberté E, Ranathunge K, Teste FP, et al. How
belowground interactions contribute to the coexistence of mycorrhizal and non-mycorrhizal
species in severely phosphorus-impoverished hyperdiverse ecosystems. Plant Soil. Springer;
2018;1–23.

35. Wright AJ, Wardle DA, Callaway R, Gaxiola A. The overlooked role of facilitation in
biodiversity experiments. Trends Ecol Evol. Elsevier; 2017;32:383–90.

36. Rutten G, Gómez-aparicio L. Plant-soil feedbacks and root responses of two
Mediterranean oaks along a precipitation gradient. Plant and Soil; 2018;

37. Snyder AE, Harmon-Threatt AN. Reduced water-availability lowers the strength of
negative plant–soil feedbacks of two Asclepias species. Oecologia. Springer; 2019;1–8.

38. Mariadassou M, Pichon S, Ebert D. Microbial ecosystems are dominated by specialist
taxa. Ecol Lett. Wiley Online Library; 2015;18:974–82.

39. Si G, Lei T, Xia Y, Yuan Y, Zhang G. Microbial Nonlinear Response to a Precipitation
Gradient in the Northeastern Tibetan Plateau. Geomicrobiol J. Taylor & Francis; 2016;33:85–
97.

40. Vries FT de, Griffiths RI, Bailey M, Craig H, Girlanda M, Gweon HS, et al. Soil bacterial
networks are less stable under drought than fungal networks. Nat Commun. Nature Publishing
Group; 2018;9:3033.

41. Fierer N, Bradford MA, Jackson RB. Toward an ecological classification of soil bacteria.
Ecology. Wiley Online Library; 2007;88:1354–64.

42. Hendershot JN, Read QD, Henning JA, Sanders NJ, Classen AT. Consistently inconsistent
drivers of microbial diversity and abundance at macroecological scales. Ecology. Wiley
Online Library; 2017;98:1757–63.

43. Richard F, Roy M, Shahin O, Sthultz C, Duchemin M, Joffre R, et al. Ectomycorrhizal
communities in a Mediterranean forest ecosystem dominated by Quercus ilex: seasonal
dynamics and response to drought in the surface organic horizon. Ann For Sci. Springer;
2011;68:57–68.

44. Querejeta J, Egerton-Warburton LM, Allen MF. Topographic position modulates the
mycorrhizal response of oak trees to interannual rainfall variability. Ecology. Wiley Online

Library; 2009;90:649–62.

45. Tedersoo L, Bahram M, Toots M, Diedhiou AG, Henkel TW, Kjøller R, et al. Towards
global patterns in the diversity and community structure of ectomycorrhizal fungi. Mol Ecol.
Wiley Online Library; 2012;21:4160–70.

46. Bauman D, Raspe O, Meerts P, Degreef J, Ilunga Muledi J, Drouet T. Multiscale
assemblage of an ectomycorrhizal fungal community: the influence of host functional traits
and soil properties in a 10-ha miombo forest. FEMS Microbiol Ecol. Oxford University Press;
2016;92.

47. Saitta A, Anslan S, Bahram M, Brocca L, Tedersoo L. Tree species identity and diversity
drive fungal richness and community composition along an elevational gradient in a
Mediterranean ecosystem. Mycorrhiza. Springer; 2018;28:39–47.

48. Bahnmann B, Mašínová T, Halvorsen R, Davey ML, Sedlák P, Tomšovský M, et al.
Effects of oak, beech and spruce on the distribution and community structure of fungi in litter
and soils across a temperate forest. Soil Biol Biochem. Elsevier; 2018;119:162–73.

49. Morris MH, Smith ME, Rizzo DM, Rejmánek M, Bledsoe CS. Contrasting
ectomycorrhizal fungal communities on the roots of co□occurring oaks (Quercus spp.) in a
California woodland. New Phytol. Wiley Online Library; 2008;178:167–76.

50. van der Putten WH, Bradford MA, Pernilla Brinkman E, van de Voorde TFJ, Veen GF.
Where, when and how plant–soil feedback matters in a changing world. Funct Ecol [Internet].
John Wiley & Sons, Ltd; 2016;30:1109–21. Available from: https://doi.org/10.1111/1365-
2435.12657

51. Smith-Ramesh LM, Reynolds HL. The next frontier of plant-soil feedback research:
unraveling context dependence across biotic and abiotic gradients. Michalet R, editor. J Veg
Sci [Internet]. 2017 [cited 2018 Jan 15];28:484–94. Available from:
http://doi.wiley.com/10.1111/jvs.12519

52. Bever JD. Soil community feedback and the coexistence of competitors: conceptual
frameworks and empirical tests. New Phytol. Wiley Online Library; 2003;157:465–73.

53. Rutten G, Prati D, Hemp A, Fischer M. Plant–soil feedback in East-- African savanna
trees. Ecology. 2016;97:294–301.

54. Fry EL, Johnson GN, Hall AL, Pritchard WJ, Bullock JM, Bardgett RD. Drought
neutralises plant–soil feedback of two mesic grassland forbs. Oecologia. Springer;
2018;186:1113–25.

55. Duell EB, Zaiger K, Bever JD, Wilson GWT. Climate Affects Plant-Soil Feedback of
Native and Invasive Grasses: Negative Feedbacks in Stable but Not in Variable Environments
[Internet]. Front. Ecol. Evol. . 2019. p. 419. Available from:
https://www.frontiersin.org/article/10.3389/fevo.2019.00419

56. Pugnaire FI, Morillo JA, Peñuelas J, Reich PB, Bardgett RD, Gaxiola A, et al. Climate
change effects on plant-soil feedbacks and consequences for biodiversity and functioning of
terrestrial ecosystems. Sci Adv. American Association for the Advancement of Science;
2019;5:eaaz1834.

57. Bardgett RD, Caruso T. Soil microbial community responses to climate extremes:
resistance, resilience and transitions to alternative states. Philos Trans R Soc B. The Royal
Society; 2020;375:20190112.

58. Page AL, Miller RH, Keeney DR. Methods of soil analysis, part 2. Chem Microbiol Prop.
Madison: American Society of Agronomy; 1982;2.

59. Frey B, Rime T, Phillips M, Stierli B, Hajdas I, Widmer F, et al. Microbial diversity in
European alpine permafrost and active layers. FEMS Microbiol Ecol [Internet].
2016;92:fiw018–fiw018. Available from: http://dx.doi.org/10.1093/femsec/fiw018

60. Edgar RC, Flyvbjerg H. Error filtering, pair assembly and error correction for next-
generation sequencing reads. Bioinformatics. Oxford University Press; 2015;31:3476–82.

61. Nikolenko SI, Korobeynikov AI, Alekseyev MA. BayesHammer: Bayesian clustering for
error correction in single-cell sequencing. BMC Genomics. BioMed Central; 2013. p. S7.

62. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet J. 2011;17:10–2.

63. Edgar RC. UPARSE: highly accurate OTU sequences from microbial amplicon reads. Nat
Methods. Nature Publishing Group; 2013;10:996.

64. Bengtsson□Palme J, Hartmann M, Eriksson KM, Pal C, Thorell K, Larsson DGJ, et al.
METAXA2: improved identification and taxonomic classification of small and large subunit
rRNA in metagenomic data. Mol Ecol Resour. Wiley Online Library; 2015;15:1403–14.

65. Bengtsson□Palme J, Ryberg M, Hartmann M, Branco S, Wang Z, Godhe A, et al.
Improved software detection and extraction of ITS1 and ITS2 from ribosomal ITS sequences
of fungi and other eukaryotes for analysis of environmental sequencing data. Methods Ecol
Evol. Wiley Online Library; 2013;4:914–9.

66. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal
RNA gene database project: improved data processing and web-based tools. Nucleic Acids
Res. Oxford University Press; 2012;41:D590–6.

67. Benson DA, Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW. GenBank.
Nucleic Acids Res. Oxford University Press; 2015;43:D30.

68. Abarenkov K, Henrik Nilsson R, Larsson K, Alexander IJ, Eberhardt U, Erland S, et al.
The UNITE database for molecular identification of fungi–recent updates and future
perspectives. New Phytol. Wiley Online Library; 2010;186:281–5.

69. McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and
graphics of microbiome census data. PLoS One. Public Library of Science; 2013;8:e61217.

70. Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4.
arXiv Prepr arXiv14065823. 2014;

71. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan:
community ecology package. R package version 2.4–4. 2017. 2018.

72. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. Bioinformatics. Oxford University Press;
2010;26:139–40.

73. Hartman K, van der Heijden MGA, Wittwer RA, Banerjee S, Walser J-C, Schlaeppi K.
Cropping practices manipulate abundance patterns of root and soil microbiome members
paving the way to smart farming. Microbiome. BioMed Central; 2018;6:14.

74. Rime T, Hartmann M, Brunner I, Widmer F, Zeyer J, Frey B. Vertical distribution of the
soil microbiota along a successional gradient in a glacier forefield. Mol Ecol. Wiley Online

Library; 2015;24:1091–108.

75. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol. BioMed Central; 2014;15:550.

76. Cáceres M De, Legendre P. Associations between species and groups of sites: indices and
statistical inference. Ecology. Wiley Online Library; 2009;90:3566–74.

77. Cáceres M De, Legendre P, Moretti M. Improving indicator species analysis by
combining groups of sites. Oikos. Wiley Online Library; 2010;119:1674–84.

78. R Core Team. R: A language and environment for statistical computing. R Found Stat
Comput Vienna, Austria [Internet]. 2018; Available from: https://www.r-project.org/.

 Tables and Figures

669 **Table 1.** Alpha diversity of the prokaryotic and fungal communities, measured as Observed

670 Richness (S_{obs}). Linear models were performed on rarefied community data, with precipitation

671 (P), tree species (T) and their interaction (P x T) as fixed effects and site as a random factor.

672 Significant terms ($P < 0.1$) are shown in bold.

Observed richness	DF	Prokaryotic communities			Fungal communities		
		ΔAIC	Chi-Sq	*P*-value	ΔAIC	Chi-Sq	*P*-value
Precipitation (P)	4,5	1.59	3.59	**0.058**	-1.71	0.29	0.589
Tree species (T)	4,5	-1.89	0.11	0.736	5.44	7.44	**0.006**
P x T	5,6	-1.75	0.25	0.617	-1.14	0.86	0.354

674 **Table 2.** Beta diversity of prokaryotic and fungal communities. PERMANOVAs were

675 performed on *edgeR* normalized data, with precipitation (P) and tree species (T) and their

676 interactions as fixed factors. All models were run with 9999 permutations and site included in

677 strata to correct for site effects. Significant terms ($P < 0.1$) are shown in bold.

Terms	DF	Prokaryotic communities					Fungal communities				
		Sum Sq	Mean Sq	F_{model}	R^2	*P* value	Sum Sq	Mean Sq	F_{model}	R^2	*P* value
Precipitation (P)	1	0.57	0.57	3.69	0.07	**0.08**	0.79	0.79	2.52	0.05	**<0.001**
Tree species (T)	1	0.11	0.11	0.70	0.01	0.20	0.44	0.44	1.40	0.03	**<0.001**
P x T	1	0.16	0.16	1.06	0.02	**0.05**	0.41	0.41	1.31	0.03	**<0.001**
Residuals	44	6.81	0.15				13.77	0.31			

679 **Table 3.** Non-metric Multi-Dimensional Scaling (NMDS) of prokaryotic and fungal
680 communities. Significant correlations between measured soil properties and NMDS are
681 shown in bold and ordered by decreasing explanatory power, measured as R^2. Soil properties
682 significantly explaining variation in both prokaryotic and fungal communities are shown in
683 italics. Significant terms ($P <0.1$) are shown in bold.

Vectors	Prokaryotic communities				Fungal communities			
	NMDS1	NMDS2	R^2	*P* value	NMDS1	NMDS2	R^2	*P* value
Calcium	-0.98	0.18	0.57	**0.001**	0.52	0.85	0.60	**0.001**
pH	-1.00	0.10	0.50	**0.001**	0.49	0.87	0.53	**0.001**
Nitrate (NO_3^-)	-0.80	-0.60	0.24	**0.004**	0.86	0.52	0.45	**0.001**
Potassium (K)	-0.93	-0.36	0.19	**0.011**	0.47	0.88	0.23	**0.004**
Texture (% sand)	0.55	0.83	0.29	**0.001**	-0.99	0.17	0.07	0.191
Magnesium (Mg)	-0.98	-0.18	0.03	0.445	-0.59	0.81	0.23	**0.006**
Phosphorus (P)	0.65	-0.76	0.10	**0.076**	0.60	0.80	0.12	**0.043**
SOM	0.99	-0.14	0.06	0.252	-0.01	1.00	0.15	**0.023**
Ammonium (NH_4^+)	0.89	-0.46	0.19	**0.008**	0.72	0.70	0.02	0.665

 Figure legends

Figure 1. Diversity of microbial communities under *Quercus ilex* trees (triangles) and
Quercus suber trees (circles) along a precipitation gradient. Alpha diversity (a, b) measured as
observed richness of rarefied a) prokaryotic and b) fungal OTUs. Beta diversity (c, d)
measured as Euclidean distance in principle coordinate between a sample and the group
centroid for c) prokaryotic and d) fungal communities. Dissimilarity indices were calculated
using Bray-Curtis distances of relative abundance per sample.

Figure 2. Non-metric multi-dimensional scaling (NMDS) based on the Bray-Curtis distances.
Community structure was based on the squared-root relative abundances of (**a**) prokaryotic
OTUs and (**b**) fungal OTUs, along the precipitation gradient (indicated by the different
colors) and between *Quercus ilex* trees (triangles) and *Quercus suber* trees (circles). Stress
values indicate the goodness of fit. Vectors indicate the strength and direction of significant
environmental variables ($P<0.01$).

Figure 3. The prokaryotic (**a**) and fungal (**b**) frequency distribution of OTUs over the
samples, where each OTU is assigned to the number of samples it occurs in. Taxa occurring
in all samples ('ubiquitous taxa') are shown in gray, whereas most OTUs only occurred in
fewer than 10 samples. Total number of OTUs per domain is indicated in brackets behind the
title.

Figure 4. Tree-specific prokaryotic (a) and fungal (b) taxa, grouped by order (prokaryotes)
and genus (fungi) and colored by phyla (legends). Log$_2$fold change values indicate the
strength and direction of the association to *Quercus ilex* (<0) and *Quercus suber* (>0).
Significance ($P<0.01$) based on FDR corrected likelihood-ratio tests. Legends show phyla
sorted by log$_2$fold change abundance, with OTU richness per phyla given in brackets.
Number of host-specific taxa per domain is indicated in brackets behind the title.

Figure 5. Proportion of locally-specific taxa for prokaryotes (filled circles) and fungi (open
circles) along the precipitation gradient. More local indicator taxa were found in drier than in
wetter areas. The trend line corresponds to the linear model including both community types
($F_{1,8} = 8.26$, $R^2_{adj} = 0.32$, $P = 0.021$). Separate linear model for prokaryotes ($F_{1,4} = 7.37$, $R^2_{adj} =$
0.56, $P = 0.053$) and for fungi ($F_{1,4} = 2.27$, $R^2_{adj} = 0.20$, $P = 0.20$) respectively.

Fig. 1

Figure 1. Diversity of microbial communities under *Quercus ilex* trees (triangles) and *Quercus suber* trees (circles) along a precipitation gradient. Alpha diversity (a, b) measured as observed richness of rarefied a) prokaryotic and b) fungal OTUs. Beta diversity (c, d) measured as Euclidean distance in principle coordinate between a sample and the group centroid for c) prokaryotic and d) fungal communities. Dissimilarity indices were calculated using Bray-Curtis distances of relative abundance per sample.

Fig. 2

Figure 2. Non-metric multi-dimensional scaling (NMDS) based on the Bray-Curtis distances.

Community structure was based on the squared-root relative abundances of (**a**) prokaryotic

OTUs and (**b**) fungal OTUs, along the precipitation gradient (indicated by the different

colors) and between *Quercus ilex* trees (triangles) and *Quercus suber* trees (circles). Stress

values indicate the goodness of fit. Vectors indicate the strength and direction of significant

environmental variables ($P<0.01$).

Fig.

Figure 3. The prokaryotic **(a)** and fungal **(b)** frequency distribution of unique OTUs over the

samples, where each OTU is assigned to the number of samples it occurs in. Taxa occurring

in all samples ('ubiquitous taxa') are shown in gray, whereas most OTUs only occurred in

fewer than 10 samples. Total number of OTUs per domain is indicated in brackets behind the

title.

Figure 4. Tree-specific prokaryotic (a) and fungal (b) taxa, grouped by order (prokaryotes) and genus (fungi) and colored by phyla (legends). Log$_2$fold change values indicate the strength and direction of the association to *Quercus ilex* (<0) and *Quercus suber* (>0). Significance (*P*<0.01) based on FDR corrected likelihood-ratio tests. Legends show phyla sorted by log$_2$fold change abundance, with OTU richness per phyla given in brackets. Number of host-specific taxa per domain is indicated in brackets behind the title.

Fig. 5

Figure 5. Proportion of locally-specific taxa for prokaryotes (filled circles) and fungi (open

circles) along the precipitation gradient. More local indicator taxa were found in drier than in

wetter areas. The trend line corresponds to the linear model including both community types

($F_{1,8} = 8.26$, $R^2_{adj} = 0.32$, $P = 0.021$). Separate linear model for prokaryotes ($F_{1,4} = 7.37$, $R^2_{adj} =$

0.56, $P = 0.053$) and for fungi ($F_{1,4} = 2.27$, $R^2_{adj} = 0.20$, $P = 0.20$), respectively.