Transcriptional profiling reveals TRPM5-expressing cells involved in viral infection in the

olfactory epithelium

B. Dnate' Baxter[1,2,‡], Eric D. Larson[3 ‡], Paul Feinstein[4], Arianna Gentile Polese[1,2], Andrew N.

Bubak[5], Christy S. Niemeyer[5], James Hassell, Jr.[5], Laetitia Merle[1,2], Doug Shepherd[6], Vijay R.

Ramakrishnan[3], Maria A. Nagel[5], and Diego Restrepo[1,2,*]

[1]Neuroscience Graduate Program, University of Colorado Anschutz Medical Campus, Aurora,

CO 80045, USA

[2]Department of Cell and Developmental Biology, University of Colorado Anschutz Medical

Campus, Aurora, CO 80045, USA

[3]Department of Otolaryngology, University of Colorado Anschutz Medical Campus, Aurora, CO

80045, USA

[4]Department of Biological Sciences, Hunter College, CUNY, New York, NY, 10065, USA

[5]Department of Neurology, University of Colorado Anschutz Medical Campus, Aurora, CO

80045, USA

[6]Department of Pharmacology, University of Colorado Anschutz Medical Campus and

Center for Biological Physics and Department of Physics, Arizona State University, USA

[‡]Co-first authors

[*]Corresponding author: Diego Restrepo, diego.restrepo@cuanschutz.edu

Abstract

Background: Understanding viral infection of the olfactory epithelium is essential because

smell loss can occur with coronavirus disease 2019 (COVID-19), caused by severe acute

respiratory syndrome coronavirus clade 2 (SARS-CoV-2), and because the olfactory nerve is an

important route of entry for viruses to the central nervous system. Specialized chemosensory

epithelial cells that express the transient receptor potential cation channel subfamily M member 5

(TRPM5) are found throughout the airways and intestinal epithelium and are involved in

responses to viral infection.

Results: Herein we performed deep transcriptional profiling of olfactory epithelial cells sorted

by flow cytometry based on the expression of fluorescent protein markers for olfactory sensory

neurons and TRPM5 in the mouse (*Mus musculus*). We find profuse expression of transcripts

involved in inflammation, immunity and viral infection in TRPM5-expressing microvillous cells

and olfactory sensory neurons. These cells express the *Tmprss2* transcript that encodes for a

serine protease that primes the SARS-CoV-2 spike protein before entry into host cells. Intranasal

infection with herpes simplex virus type 1 (HSV-1) elicited a decrease in olfactory sensory

neurons.

Conclusion: Our study provides new insights into a potential role for TRPM5-expressing cells in

viral infection of the olfactory epithelium. We find that, as found for solitary chemosensory cells

(SCCs) and brush cells in the airway epithelium, and for tuft cells in the intestine, the

transcriptome of TRPM5-expressing microvillous cells and olfactory sensory neurons indicates

that they are likely involved in the inflammatory response elicited by viral infection of the

olfactory epithelium.

Keywords: Olfactory sensory neurons, Microvillous cells, Viral infection, Immunity,

Inflammation, Mouse

Background

Self-reported loss of smell in a large fraction of patients with coronavirus disease 2019 (COVID-19), caused by severe acute respiratory syndrome coronavirus clade 2 (SARS-CoV-2), (Giacomelli et al., 2020; Parma et al., 2020; Yan et al., 2020a; Yan et al., 2020b) raises the question of how SARS-CoV-2 affects olfaction in a subset of patients. Entry of SARS-CoV-2 into cells is mediated by spike protein attachment to the SARS-CoV receptor ACE2 followed by spike protein priming by the serine protease TMPRSS2 (Hoffmann et al., 2020). Chemosensory cells found in the airway (SCCs/brush cells) and intestinal epithelium (tuft cells) express the transient receptor potential cation channel subfamily M member 5 (TRPM5) and other elements of the taste transduction pathway and have been implicated in immune and inflammatory responses to bacterial, viral and parasitic infection (Luo et al., 2019; Maina et al., 2018; O'Leary et al., 2019; Perniss et al., 2020; Rane et al., 2019; Saunders et al., 2014; Tizzano et al., 2010). In the olfactory epithelium TRPM5 and other proteins involved in taste transduction are also expressed in SCC-like microvillous cells (MVCs)(Genovese and Tizzano, 2018; Lin et al., 2008), which have been proposed to be involved in a protective response to high concentrations of odorants (Fu et al., 2018; Lemons et al., 2017). The ACE2 receptor and TMPRSS2 are expressed in support cells, stem cells, and MVCs (Brann et al., 2020; Fodoulian et al., 2020). However, whether MVCs play a role in viral infection or viral infection defense of the olfactory epithelium is unknown.

Herein, we performed transcriptional profiling of MVCs and a subset of olfactory sensory neurons (OSNs) expressing eGFP under control of the TRPM5 promoter (OSN_eGFP+ cells)(Lin et al., 2007; Lopez et al., 2014). In order to profile these low abundance cells we used

a modified version of Probe-Seq, which allows deep transcriptional profiling of specific cell

types identified by fluorescent markers as the defining feature (Amamoto et al., 2019). We

crossed a mouse expressing mCherry in the nuclei of OSNs under control of the OMP promoter

(OMP-H2B::mCherry mice) with TRPM5-eGFP transgenic mice (Clapp et al., 2006) (OMP-

H2B::mCherry/TRPM5-eGFP mice). We isolated cells from the olfactory epithelium and used

fluorescence-activated cell sorting (FACS) to sort MVC_eGFP cells (mCherry negative and

eGFP positive) and cells labeled by OMP-driven mCherry that did or did not express eGFP

(OSN_eGFP+ and OSN_eGFP- cells) followed by transcriptional profiling by RNA sequencing

(RNAseq).

Results

Fluorescence-activated cell sorting of cells isolated from the main olfactory epithelium. The

olfactory epithelium of OMP-H2B::mCherry/TRPM5-eGFP mice expressed nuclear mCherry

driven by the OMP promoter in the intermediate layer of the olfactory epithelium (Figures 1a), as

expected for the location of nuclei of mature OSNs (Farbman and Margolis, 1980). eGFP

expression driven by the TRPM5 promoter was found in MVCs, with cell bodies located mostly

in the apical layer of the epithelium (asterisks), and at lower expression levels in a subset of

OSNs double-labeled with mCherry (Figure 1a), consistent with earlier publications (Lin et al.,

2008; Lin et al., 2007; Pyrski et al., 2017).

We proceeded to isolate cells from the main olfactory epithelium of OMP-

H2B::mCherry/TRPM5-eGFP mice (see Methods, Figure 1b). Figure 1c shows two isolated

OSNs with differential expression of eGFP. Using flow cytometry we found that fluorescence

intensity of individual cells for mCherry and eGFP spanned several orders of magnitude (Figure

1d). We proceeded to sort three groups of cells: high mCherry-expressing cells with low and

high eGFP fluorescence (presumably mature OSNs, these cells are termed OSN_eGFP- and

OSN_eGFP+ cells respectively) and cells with low mCherry and high eGFP expression

(MVC_eGFP, presumably MVCs). Reverse transcription quantitative PCR (RT-qPCR) showed

that, as expected the OSN_eGFP- and OSN_eGFP+ cells have higher levels of OMP transcript

than MVC_eGFP cells (Figure 1e,i), and OSN_eGFP+ cells and MVC_eGFP cells have higher

levels of eGFP transcript compared to OSN_eGFP- cells (Figure 1e,ii). Furthermore, compared

to OSN_eGFP- cells both the MVC_eGFP cells and OSN_eGFP+ cells expressed higher levels

of TRPM5 transcript (Figure 1e,iii) and choline acetyl transferase (ChAT)(Figure 1e,iv), a

protein involved in acetylcholine neurotransmission that is expressed in MVCs (Ogura et al.,

2011). The asterisks in Figure 1e denote significant differences tested with either t-test or

ranksum with p-values below the p-value of significance corrected for multiple comparisons

using the false discovery rate (pFDR)(Curran-Everett, 2000) (pFDR is 0.033 for OMP, 0.05 for

TRPM5, 0.05 for EGFP and 0.03 for ChAT, n=8 for OMP OSN_eGFP-, 4 for OMP

OSN_eGFP+ and 4 for MVC_eGFP cells).

The number of OSN_eGFP+ cells sorted by FACS is decreased when OMP-

H2B::mCherry/TRPM5-eGFP mice are placed in ventilated cages. In our vivarium we have

ventilated cages (HV cages) where air is mechanically exchanged with fresh air once every

minute and static cages (LV cages) where air is exchanged passively through a filter in the cover.

When we moved the OMP-H2B::mCherry/TRPM5-eGFP to HV cages we noticed a decrease in

the number of OSN_eGFP+ cells sorted per mouse (Figures 2a,b and c), suggesting that changes

in ventilation conditions affect TRPM5 promoter-driven expression of eGFP. Following this

observation, mice were moved back to LV cages. We proceeded to study the dependence of the

number of OSN_eGFP+ cells sorted on the number of days in LV vs. HV cages. The number of

OSN_eGFP+ cells is positively correlated with the number of days the animal spends in LV

cages (Figure 2d) and negatively correlated to the number of days the animals spend in the HV

cages (Figure 2e). Generalized linear model (GLM) analysis found significant differences for the

number of OSN_eGFP+ cells sorted as a function of the number of days in LV cages ($p<0.05$, 26

observations, 24 d.f., F-statistic = 5.64, p-value for GLM <0.05) and the number of days in HV

cages ($p<0.05$, 26 observations, 24 d.f., F-statistic = 5.99, p-value for GLM <0.05). For RNAseq

experiments one FACS sort was done using cells from mice born and maintained in HV housing,

and the OSN_eGFP+ yield was low. Subsequently, we performed all FACS with cells isolated

from the olfactory epithelium of mice raised in LV cages.

Coverage of TRPM5 transcript by RNAseq encompasses the full transcript in MVC_eGFP

cells and OSN_eGFP+ cells. Pyrski and co-workers did not find full-length TRPM5 transcript

in reverse transcriptase polymerase chain reactions with mRNA extracted from isolated OSNs

from the adult mouse and did not find *in situ* signal in the OSN layer of the olfactory epithelium

for the full-length TRPM5 transcript (Pyrski et al., 2017). These investigators found full-length

TRPM5 transcript and strong *in situ* signal in MVCs. We find strong *in situ* signal for TRPM5 in

MVC_eGFP cells located in the apical layer of the olfactory epithelium (Figure 3aii, asterisks).

In addition, we find sparse TRPM5 *in situ* labeling in the nuclear OSN layer (Figure 3aii,

arrows). In order to gain a better understanding of which TRPM5 transcript is expressed in OSNs

we performed an analysis of TRPM5 transcript coverage for the RNAseq performed with RNA

from the different groups of cells sorted by FACS. Consistent with *in situ* labeling, TRPM5

transcript was significantly higher in OSN_eGFP+ cells compared to OSN_eGFP- cells and in

MVC_eGFP cells compared to OSN_eGFP- cells in both male and female adult mice (Figures

3b,c). To explore RNA sequencing coverage of individual *Trpm5* exons, we computed read

depth over the sequence of the *Trpm5* gene for each sample. We found coverage for all exons in

OSN_eGFP+ cells and for all exons except exon 1 and 5'UTR1 in MVC_eGFP cells, but there

was no coverage in OSN_eGFP- cells (Figures 3b,c). GLM analysis found statistically

significant differences for exons ($p < 0.001$, 450 observations, 446 d.f., F-statistic = 33.8, p-value

for GLM < 0.001), but no overall significance between MVC and OSN_eGFP+ groups ($p > 0.05$,

450 observations, 446 d.f., F-statistic = 33.8, p-value for GLM <0.001). However, post-hoc

ranksum tests did yield significant differences between MVC and OSN_eGFP+ groups for exon

1 and 5'UTR1 (p<pFDR=0.0033). We observed no differences in coverage between male and

female mice (within MVC_eGFP cells: p>0.05, 210 observations, 206 d.f., F-statistic = 22.8, p

value for GLM <0.001; within OSN_GFP+: p>0.05, 240 observations, 236 d.f., F-statistic =

11.5, p value for GLM <0.001). Together, these data suggest that both OSN_eGFP+ cells and

MVC_eGFP cells are capable of expressing full-length TRPM5 transcript.

RNAseq indicates that OSN_eGFP+, OSN_eGFP- and MVC are three distinct groups of

chemosensory cells in the mouse olfactory epithelium. Differential gene expression analysis of

the RNAseq data was used to compare the three olfactory epithelium cell groups sorted by

FACS. We found that expression of 2000 genes was significantly higher in OSN_eGFP+

compared to OSN_eGFP-, and expression of 1821 genes was lower in OSN_eGFP+ cells (Figure

4 -figure supplement 1 shows the results of RNAseq and Figure 4 -figure supplement 2 shows

the metadata). Figure 4a shows expression levels for the transcripts that showed the largest

differences between OSN_eGFP+ and OSN_eGFP- cells. The transcripts for TRPM5 and eGFP

were among the top 10 genes whose transcription was higher in OSN_eGFP+ compared to

OSN_eGFP- with 105-fold and 42-fold increases respectively. This top 10 OSN_eGFP+

upregulated group also includes *Avil* and *Adgrg6* that are involved in remodeling processes after

peripheral nerve injury (Chuang et al., 2018; Jablonka-Shariff et al., 2020) and *Espn*, encoding

for espin, a protein playing a structural role in microvilli of chemosensory cells (Sekerkova et al.,

2004). Interestingly, the olfactory activity-dependent protein S100a5 (Fischl et al., 2014) is

found among the top 10 OSN_eGFP+ downregulated transcripts suggesting that in these mice

the OSN_eGFP+ are not stimulated by the odorants in their housing environment (Figure 4a).

Additionally, a majority of the olfactory receptors show decreased transcription in OSN_eGFP+

compared to OSN_eGFP- cells (Figures 4a,b) and the volcano plot for olfactory receptor

transcript expression shows only 25 olfactory receptors show increased expression in

OSN_eGFP+ cells (Figure 4c, based on fold change > 4 and average expression > 100 counts,

Table 1).

Expression of 4386 genes was significantly higher in MVC_eGFP cells compared to

OSN_eGFP- cells, and expression of 5630 genes was lower in MVC_eGFP cells (Figure 5 –

figure supplement 1). Transcripts for 550 olfactory receptors were lower in MVC_eGFP cells

(Figure 5 – figure supplement 1), and no olfactory receptors were upregulated in MVC_eGFP

cells compared to OSN_eGFP-. Figure 5a shows expression levels for the transcripts with the

largest differences between MVC_eGFP cells and OSN_eGFP- cells. Six of the transcripts that

are within the top 10 upregulated genes found in MVC_eGFP cells compared to OSN_eGFP-

cells (Figure 5a) are also found within the 10 top upregulated transcripts when OSN_eGFP+ cells

are compared to OSN_eGFP- cells (Figure 4a)(*Adgrg6*, *Avi*l, *Cd24a*, *eGFP*, *Espn* and *Trpm5*).

TRPM5 and eEGFP were among the top 10 genes whose transcription was higher in

MVC_eGFP cells compared to OSN_eGFP- cells with 1471-fold and 75-fold differences

respectively. Interestingly, *Pou2f3*, a transcription factor important in differentiation of MVCs

(Yamaguchi et al., 2014; Yamashita et al., 2017), is found within the top 10 upregulated genes

found in MVC_eGFP cells compared to OSN_eGFP- (Figure 5a) and is also significantly higher

in OSN_eGFP+ cells compared to OSN_eGFP- cells (Figure 5 – figure supplement 1). Finally,

OMP and *s100a5*, genes for two proteins expressed in mature OSNs (Farbman and Margolis,

1980; Fischl et al., 2014), were among the top 10 downregulated transcripts in MVC_eGFP cells

compared to OSN_eGFP- cells (Figure 5a).

We found expression of 3068 genes which was significantly higher in MVC_eGFP cells

compared to OSN_eGFP+ cells, and expression of 4060 genes was lower in MVC_eGFP cells

(Figure 5 – figure supplement 2). Figure 5b shows expression levels for the genes that showed

the largest differences between MVC and OSN_eGFP+ cells. Among the 10 genes that are

highly expressed in OSN_eGFP+ cells compared to MVC_eGFP cells we find *Pde4a,* a gene

expressed in mature OSNs (Juilfs et al., 1997) (*OMP* is also significantly higher in OSN_eGFP+

cells, but is not among the top 10 genes, Figure 5 – figure supplement 2). Interestingly, *Hcn2*, a

gene that encodes for a hyperpolarization-activated cAMP channel that has been postulated to

participate in OSN axon growth and glomerular innervation (Mobley et al., 2010), is found in the

top 10 upregulated OSN_eGFP+ genes. Finally, the gene encoding for the synaptic protein

Snap25 is found in the top 10 upregulated lists for both OSN_eGFP+ and OSN_eGFP- indicating

that both OSNs are involved in synaptic transmission (Figures 5a and b).

We did not find major differences in transcriptome profiling between males and females for

genes that were differentially expressed between the three cell groups (Figure 5 – figure

supplement 3,4). We found a substantial number of olfactory receptor genes that were

differentially expressed between males and females (Figure 5 – figure supplement 4).

Surprisingly, the differentially expressed olfactory receptors differed from receptors identified by

van der Linden et al. (van der Linden et al., 2018). Finally, we compared expression of

transcripts involved in taste transduction, canonical olfactory transduction, and non-canonical

OSNs (Figure 4d). The non-canonical OSNs considered here included guanilyl-cyclase D (GC-

D) OSNs (Juilfs et al., 1997), Trpc2 OSNs (Omura and Mombaerts, 2014) and Cav2.1 OSNs

(Pyrski et al., 2018). Both OSN_eGFP+ and OSN_eGFP- expressed low levels of *Cancna1a*

encoding for Cav2.1 and *Trpc2*. OSN_eGFP- expressed higher levels of Trace amine-associated

receptors (Liberles, 2015) than OSN_eGFP+ cells. Both OSN_eGFP+ and OSN_eGFP-

expressed transcripts for OMP, BBS1 and 2 and proteins involved in the canonical olfactory

transduction pathway, markers of canonical OSNs.

Gene ontology enrichment analysis reveals differences in chemosensory transduction and

synaptic vesicle function between the three groups of cells. Perusal of the top differences

between the three cell groups suggested that these are distinct chemosensory cell types found in

the olfactory epithelium. Both OSN_eGFP+ and OSN_eGFP- share expression of OSN-specific

transcripts, express distinct subsets of olfactory receptors and differ in expression of the activity-

dependent transcript *S100a5*, and MVC_eGFP cells differ from both OSN groups in expression

of transcripts for synaptic transmission and for markers of mature OSNs and microvillous cells.

In order to perform a thorough analysis of the differences between these chemosensory cell

groups we performed an analysis of gene ontology (GO) enrichment for lists of genes related to

chemosensory perception. When compared with either OSN_eGFP+ or OSN_eGFP- we found

that MVC_eGFP cells were enriched for transcripts for the gene ontology list of sensory

perception of sweet/umami taste (GO:0050916 and GO:0050917) (Figure 5d, Figure 5 – figure

supplements 3,4) involving taste detection/transduction proteins that have been reported to be

expressed in MVCs (Genovese and Tizzano, 2018; Hegg et al., 2010): *Gnat3*, encoding for

gustducin, the G protein mediating sweet and umami taste transduction (McLaughlin et al.,

1992), *Itpr3*, encoding for the inositol-1,4,5-triphosphate receptor type 3 and *Tas1r3*, encoding

for a gustducin-coupled receptor involved in umami and sweet taste (Damak et al., 2003; Zhang

et al., 2003). Interestingly, the GO lists for sensory perception of sweet/umami taste

(GO:0050916 and GO:0050917) and other lists for sensory and taste perception (GO:0050906,

GO:0050912) are enriched in OSN_eGFP+ cells compared to OSN_eGFP- cells (Figure 4e,

Figure 4 – figure supplement 3, including *Gnat3, Itpr3, Tas1r3*). Furthermore, gene ontology

analysis for OSN_eGFP+ cells compared to OSN_eGFP- cells finds decreased enrichment for

sensory perception of smell (GO:0007608) and G protein-coupled receptor signaling pathways

(GO:0007186) that include a large number of olfactory receptors and transcripts encoding for

proteins involved in peripheral olfaction such as *Gfy*, *Omp*, *Pde1c* and *Pde4a* (GO:0007608) and

Dgkg, Gng13, Itgb1, Nsg1 (GO:0007186). Finally, enrichment of gene ontology lists for synaptic

vesicle function were decreased for MVC_eGFP cells compared with either OSN_eGFP+ or

OSN_eGFP- cells (Figure 5c,d). Results of this gene ontology analysis of chemosensation and

synaptic vesicle function reinforces the finding that the three cell groups in this study are distinct

chemosensory cell types of the olfactory epithelium. OSN_eGFP+ cells are related to

MVC_eGFP cells because of expression of taste perception gene ontology, but differ from

MVC_eGFP cells in expression of olfactory receptors and transcripts related to synaptic function

as expected for an OSN.

Gene ontology analysis finds enrichment of lists of viral-related, inflammation and immune

transcripts in MVC_eGFP cells and OSN_eGFP+ cells. SCCs, tuft and brush cells have been

implicated in responses to bacterial and viral infection, immunity and inflammation (Luo et al.,

2019; Maina et al., 2018; O'Leary et al., 2019; Perniss et al., 2020; Rane et al., 2019; Saunders et

al., 2014; Tizzano et al., 2010; Ualiyeva et al., 2020). The fact that MVCs are closely related to

these cells (Fu et al., 2018; Genovese and Tizzano, 2018; Ogura et al., 2011) lead us to search for

gene ontology enrichment related to bacterial and viral infection, immunity and inflammation for

MVC_eGFP cells. We found robust enrichment of these gene ontologies in MVC_eGFP cells

and OSN_eGFP+ cells (Figures 4e, 5c,d). Transcripts related to viral infection that were higher

in MVC_eGFP cells and OSN_eGFP+ cells compared to OSN_eGFP- cells (Figure 6) including

those involved in viral entry into host cells, viral transcription and regulation of viral

transcription, negative regulation of viral genome replication and negative regulation of viral

process (Figures 4e, 5c, Figure 6 – figure supplements 1-3). We also found gene ontology

enrichment in MVC_eGFP cells and OSN_eGFP+ cells compared to OSN_eGFP- cells for

defense response to bacterium (Figure 6 – figure supplements 1-3).

Importantly, we also find enrichment for transcript expression for immunity and inflammation

(Figures 4e, 5c,d and Figure 6 – figure supplements 1-3). Genes related to inflammation and

immunity that were higher in MVC_eGFP cells and OSN_eGFP+ cells compared to

OSN_eGFP- cells are shown in Figure 6 – figure supplements 4-7. Among these transcripts *IL25*

and its receptor *Il17rb* are enriched in both MVC_eGFP cells and OSN_eGFP+ cells. In SCCs,

brush cells and tuft cell generation of IL25 leads to a type 2 inflammation and stimulates

chemosensory cell expansion in a sequence of events that also involves cysteinyl leukotrienes

(Bankova et al., 2018; Luo et al., 2019; von Moltke et al., 2016). The presence of both *Il25* and

Il17rb suggests an autocrine effect. Furthermore, both cell types displayed increased expression

of transcripts encoding for enzymes involved in eicosanoid biosynthesis such as *Alox5*, *Ptgs1*

and *Ptgs2* that are found in brush cells in the airways (Bankova et al., 2018) and tuft cells in the

intestine (McGinty et al., 2020) where they drive type 2 immune responses.

Acute infection with herpes simplex virus-1 (HSV-1) elicits a decrease in the fraction of

OSNs. HSV-1 infects the olfactory epithelium in mice (Shivkumar et al., 2013). We found that

acute (5 DPI) high titer ($1x\ 10^6$ PFU/naris; McKrae strain) intranasal infection with HSV-1

elicited a decrease in the fraction of OSN_eGFP- and OSN_eGFP+ cells (Figure 7). GLM

analysis yielded significant effects for the fraction of OSN_eGFP- compared to MVCs ($p<0.001$)

and for the interaction between the fraction of OSN_eGFP- compared to MVCs and HSV-1 and

treatment ($p<0.001$, 75 observations, 69 d.f., GLM F-statistic=111 and p-value < 0.001, n=14

mice for control, n=11 mice for HSV-1 inoculation). Post-hoc t-test p-value was significant for

OSN_eGFP- and OSN_eGFP+ cells ($p < pFDR = 0.047$), but not for MVCs.

Discussion

We performed transcriptional profiling of three chemosensory cells in the mouse olfactory epithelium: MVC_eGFP cells and two types of OSNs: OSN_eGFP+ and OSN_eGFP-. We found that while the transcriptome of each of these cell types is distinct they share common features across groups. The two groups of OSNs share transcript expression for proteins expressed in OSNs such as OMP, olfactory transduction proteins, and proteins involved in synaptic function. Yet, they differ in olfactory receptor expression and OSN_eGFP+ express taste transduction transcripts and other transcripts found in SCCs such as *Il25* and *Pou2f3*. On the other hand, MVC_eGFP cells express transcripts encoding for taste transduction proteins and other transcripts found in SCCs such as *Pou2f3* but they do not express transcripts for proteins involved in olfactory transduction and synaptic function, and they do not express olfactory receptors. Finally, we found that MVC_eGFP cells and OSN_eGFP+ cells express a substantial number of transcripts involved in viral infection, inflammation and immunity.

Here we find that OSN_eGFP+ are OSNs expressing full-length TRPM5 in the adult mouse (Figure 3). This raises the question why our results differ from Pyrski and co-workers who did not find full-length TRPM5 in OSNs (Pyrski et al., 2017). Likely, this is due to differences in environmental conditions that alter the number of OSNs expressing TRPM5 in the adult mouse (Figure 2). Interestingly, consistent with our finding of TRPM5 expression in OSNs, analysis of scRNA data in the literature indicates that there are 8 OSNs expressing TRPM5 among 3209 OSNs in the data set from Ziegler and co-workers (Ziegler et al., 2020) and 3 OSNs expressing TRPM5 among 2113 OSNs in the data set from Wu and co-workers (Wu et al., 2018). Finally, OSN_eGFP+ express low levels of the activity-dependent transcript s100a5 (Fischl et al., 2014)

(Figure 4a) suggesting that these cells express olfactory receptors that are not stimulated by

odorants present in the cage.

Gene ontology analysis revealed that MVC_eGFP cells (and OSN_eGFP+ cells to a lesser

extent) are enriched in viral-related transcripts compared to OSN_eGFP- (Figures 4e, 5c,d, and

Figure 6 – figure supplements 1-3). To infect cells, viruses must interact with host cell

membranes to trigger membrane fusion and viral entry. Membrane proteins at the surface of the

host cell are thus key elements promoting or preventing viral infection. Here we find that

transcripts for several membrane proteins and cell adhesion molecules involved in viral entry are

enriched in MVC_eGFP cells. *Plscr1* encodes a phospholipid scramblase which has been shown

to promote herpes simplex virus (HSV) entry in human cervical or vaginal epithelial cells and

keratinocytes (Cheshenko et al., 2018), and hepatitis C virus entry into hepatocytes (Gong et al.,

2011). In contrast with its role in viral entry, PLSCR1 impairs the replication of other types of

viruses in infected cells (influenza A virus (Luo et al., 2018), hepatitis B virus (Yang et al.,

2012)). IFTM2 is another transmembrane protein that mediates viral entry. In contrast with

PLSCR1, IFTM2 inhibits viral entry of human immunodeficiency virus (HIV, (Yu et al., 2015)),

hepatitis C virus (Narayana et al., 2015), influenza A H1N1 virus, West Nile virus, and dengue

virus (Brass et al., 2009). IFTM2 also inhibits viral replication (Brass et al., 2009) and protein

synthesis (Lee et al., 2018). Nectins are transmembrane glycoproteins and constitute cell surface

receptors for numerous viruses. There is wide evidence that HSV can enter host cells through

Nectin-1 dependent mechanisms, particularly for neuronal entry (Kopp et al., 2009; Petermann et

al., 2015; Sayers and Elliott, 2016; Shukla et al., 2012), and Nectin-4 appears essential for

measles virus epithelial entry (Noyce and Richardson, 2012; Singh et al., 2015; Singh et al.,

2016). In addition to cell surface molecules, the mucus contains secreted proteins that confer

protection against viruses to the underlying cells. Glycoproteins are major constituents of mucus

and exhibit multiple pathogens binding-sites. We found the *Ltf* transcript in MVC_eGFP cells,

which encodes for lactotransferrin. Lactotransferrin is a globular glycoprotein widely represented

in the nasal mucus with anti-viral activity against Epstein-Barr virus (Zheng et al., 2014; Zheng

et al., 2012), HSV (Shestakov et al., 2012; Valimaa et al., 2009)) and Hepatitis C virus (Allaire et

al., 2015).

Viruses have developed numerous strategies to overcome barrier mechanisms to enter the cells.

After viral entry infected cells have other resources to fight against viral infection by disrupting

the production of new viral particles, limiting inflammation processes and activating innate

immune responses. For example, TRIM25 is an ubiquitin ligase that activates retinoic acid-

inducible gene I (RIG-I) to promote the antiviral interferon response (Gack et al., 2007).

Furthermore, influenza A virus targets TRIM25 to evade recognition by the host cell (Gack et al.,

2009). In addition, TRIM25 displays a nuclear role in restricting influenza A virus replication

(Meyerson et al., 2017). Zc3h12a, also known as MCPIP-1, inhibits hepatitis B and C virus

replication, reduces virus-induced inflammation (Li et al., 2020; Lin et al., 2014), and exerts

antiviral effects against influenza A virus (Dong et al., 2017). Finally, *Pou2f3* also called *Skn1a*,

encodes for a key regulator for the generation of TRPM5-expressing cells in various epithelial

tissues (Yamashita et al., 2017). *Pou2f3* transcript was increased in MVC_eGFP cells (and to a

lesser extent in OSN_eGFP+) compared to OSN_eGFP-. *Skn1a/Pou2f3*-deficient mice lack

intestinal tuft cells and have defective mucosal type 2 responses to helminth infection in the

intestine (Gerbe et al., 2016). Finally, both OSN_eGFP+ and MVC_eGFP cells express Il25, an

interleukin that is involved in the inflammatory response of TRPM5-expressing epithelial cells in

the airway epithelium and the gut (O'Leary et al., 2019), and in the skin Il25 expression leads to

disruption of the epithelium and enhances HSV-1 and vaccinia virus replication (Kim et al.,

2013).

Our findings of expression of virally relevant transcripts in MVC_eGFP cells complement

published studies on the role of MVC-related SCCs in viral infection. In the trachea, viral-

associated formyl peptides activate SCCs to release acetylcholine and activate mucocilliary

clearance by ciliated cells (Perniss et al., 2020). This activation is mediated by the TRPM5

transduction pathway in the SCC and muscarinic acetylcholine receptors in the ciliated cell. In a

similar manner in the olfactory epithelium MVCs respond to ATP, which is involved in

activating mucociliary movement by releasing acetylcholine and activating adjacent

sustentacular cells through a muscarinic receptor (Fu et al., 2018). Therefore, viral infection

could result in activation of MVCs resulting in activation of mucociliary clearance by adjacent

sustentacular cells. In addition, in the anterior olfactory epithelium, where there is a higher

density of MVCs, mice exposed to mild odorous irritants exhibited a time-dependent increase in

apoptosis and a loss of mature OSNs without a significant increase in proliferation or

neurogenesis (Lemons et al., 2020). This MVC-induced apoptosis could contribute to the

decrease in the fraction of OSNs elicited by intranasal inoculation of HSV-1 (Figure 7). Future

experiments are necessary to determine whether activation of MVCs by viruses could lead to loss

of mature OSNs contributing to smell loss after viral infection. We did not find changes in MVC

cell number after acute infection with HSV-1 (Figure 7c,iii). Interestingly, in the mouse distal

lung, where there is no expression of SCCs, there was de novo generation of SCCs after infection

with A/H1N1/PR/8 influenza virus (Rane et al., 2019) raising the question whether longer virus

exposure could alter MVC number in the olfactory epithelium. Finally, because we find high

expression of transcripts involved in Type 2 immune response in MVCs viral activation of these

cells could result in activation of cytokine-induced inflammation by long-term horizontal basal

cells that activate type 1 immune responses within the olfactory epithelium (Chen et al., 2019).

The olfactory epithelium provides direct viral access to the brain through the olfactory nerve.

Whether this olfactory path constitutes route of entry for viruses to the brain is a matter of

intense discussion, especially because some viruses are postulated to be involved in

encephalopathy and neurodegenerative disorders (Dando et al., 2014; Doty, 2008). Our findings

that these TRPM5-bearing OSN_eGFP+ cells are enriched in virally-related genes suggests that

these cells may be involved in or prevention of viral entry into the brain (and these two

alternatives are not exclusive since they may be different for different viruses). On the one hand,

we identified transcripts encoding for viral receptors in OSN_eGFP+ cells, suggesting that

viruses can enter these OSNs. If viral particles were to enter the OSNs they could reach the

olfactory bulb through anterograde transport along the olfactory nerve and from the olfactory

bulb, viruses can spread throughout the brain along the olfactory bulb-hippocampus route. On

the other hand, we found enrichment for transcripts encoding for proteins involved in limiting

viral infection and promoting immune and anti-inflammatory responses in OSNs_eGFP+ and

MVC_eGFP cells. In this case, viral spread to the brain would be prevented. Finally, the

olfactory epithelium is innervated by the trigeminal nerve, and substance P immunostaining is

closely associated with subsets of MVCs (Lin et al., 2008). This raises the question whether an

interaction between MVCs and trigeminal nerve fibers could participate in local inflammation as

found for SCCs (Saunders et al., 2014), and could modulate the entry of virus to the brain stem

through the trigeminal nerve. Future experiments are necessary to study the potential role of

MVC_eGFP cells and OSNs_eGFP+ in viral infection of the olfactory epithelium and the brain.

Recently, due to the current COVID-19 pandemic, researchers have focused their attention on

investigating SARS-CoV-2 mechanism of entry into cells. SARS-CoV-2 targets mainly cells of

the respiratory pathway where viral entry is mediated by ACE2 and TMPRSS2 (Hoffmann et al.,

2020). Because numerous patients reported loss of smell (Giacomelli et al., 2020; Parma et al.,

2020; Yan et al., 2020a; Yan et al., 2020b), researchers wondered about the mechanism for

SARS-CoV-2 infection of the olfactory epithelium. In our study, we found the *Tmprss2*

transcript was significantly increased in MVC_eGFP cells and OSN_eGFP+ compared to

OSN_eGFP- (Figure 6). We did not find *Ace2* enrichment in these cells, but this may be due to

inefficiency in finding with RNAseq low abundance transcripts like *Ace2* (Ziegler et al., 2020).

Transcriptional profiling of single cells in the olfactory epithelium from other laboratories found

expression of transcripts for both *Tmprss2* and *Ace2* in in sustentacular cells and stem cells, and

at lower levels in MVCs (Brann et al., 2020; Fodoulian et al., 2020). Viral infection of

sustentacular cells may explain loss of smell because these cells play a key role in supporting

olfactory function by providing glucose for the energy necessary for olfactory transduction in the

OSN cilia (Villar et al., 2017). Importantly, type I interferons, and to a lesser extent type II

interferons induced by response of the host to SARS-CoV-2, and infection by other viruses

inducing the interferon pathway increases *Ace2* expression in the nasal epithelium (Ziegler et al.,

2020). MVCs may play a role in SARS-CoV-2 infection of the olfactory epithelium because

these cells may participate in activating inflammation of the epithelium that elicits type 1

immune response (Chen et al., 2019). Finally, our finding of transcripts involved in viral entry,

replication and defense in a subset of OSNs raises the question whether viruses enter the central

nervous system through the olfactory nerve (Bilinska et al., 2020). This is relevant to the

potential long term effect of SARS-CoV-2 and other viruses in neurological and

neurodegenerative disorders (De Felice et al., 2020). Our study provides new insights into a

potential role for TRPM5-expressing cells in viral infection of the main olfactory epithelium.

Conclusion

Here we compared the transcriptome of OSNs that do not express TRPM5 (OSN_eGFP-) to a

small fraction of cells that express TRPM5 (MVC_eGFP cells and OSN_eGFP+ cells) in the

olfactory epithelium. We find that transcript expression in these three types of cells is distinct.

OSN_eGFP- and OSN_eGFP+ cells share transcripts expressed in OSNs, while OSN_eGFP+

and MVCs share expression of transcripts found in SCCs and brush cells in the airways and tuft

cells of the intestine. Interestingly, OSN_eGFP+ and MVCs express transcriptomes within the

gene ontology transcript groups for immunology, inflammation and viral infection.

Abbreviations

COVID-19: Coronavirus disease 2019

DPI: Days post infection

eGFP: Enhanced green fluorescent protein

FACS: Fluorescence-activated cell sorting

FDR: False discovery rate

GLM: Generalized linear model

GO: Gene ontology

MVCs: Microvillous cells

OMP: Olfactory marker protein

OSNs: Olfactory sensory neurons

PSF: Point spread function

SARS-CoV-2: Severe acute respiratory syndrome coronavirus clade 2

SSC: Saline-sodium citrate

SSCT: SSC with tween

TRPM5: Transient receptor potential cation channel subfamily M member 5

Methods

Key Resources Table

REAGENT TYPE	REAGENT or RESOURCE	SOURCE	IDENTIFIER	ADDITIONAL INFORMATION
Chemical compound, drug	BrainPhys Neuronal Medium	Stemcell Technologies		Product # 05791
Chemical compound, drug	Dispase II	Sigma		Product # D4693
Chemical compound, drug	AcGFP1/eGFP calibration beads	Takara		Flow cytometry calibration beads Product # 632594
Chemical compound, drug	mCherry calibration beads	Takara		Flow cytometry calibration beads Product # 632595
Chemical compound, drug	RQ1 RNase-free DNase	Promega		Product # M6101
Chemical compound, drug	Papain	Sigma		Product # P3125
Chemical compound, drug	Paraformaldehyde (32%)	Electron Microscopy Sciences		Product # 157145
Chemical compound, drug	RNAprotect Tissue Reagent	Qiagen		Product # 76526
Chemical compound, drug	RNeasy Plus Micro Kit	Qiagen		Product # 74034
Chemical compound, drug	High Capacity c-DNA Reverse Transcription kit	ABI		
Chemical compound, drug	18s rRNA	PE ABI		
Strain, strain background	TRPM5-eGFP	Dr. Robert Margolskee (Clapp et al., 2006)		
Strain, strain background	OMP-H2B::Cherry	Generated for this publication		This mouse will be deposited in Jackson Laboratories
Software, algorithm	MATLAB_R2018a	Mathworks	RRID: SCR_001622	
Software, algorithm	Illustrator	Adobe	RRID: SCR_010279	
Software, algorithm	Photoshop	Adobe	RRID: SCR_014199	

Software, algorithm	InDesign	Adobe		
Software, algorithm	MoFlo Astrios Summit Software (6.3.1.16945).	Beckman Coulter		
Software, algorithm	BBMap (BBDuk)		RRID:SCR_016968	
Software, algorithm	Salmon v1.2.1	https://combine-lab.github.io/salmon/	RRID:SCR_017036	(Patro et al., 2017)
Software, algorithm	DeSEQ2 v1.28.0	bioconductor.org https://bioconductor.org/packages/release/bioc/html/DESeq2.html	RRID:SCR_015687	(Love et al., 2014)
Software, algorithm	TopGO, v2.40.0		RRID:SCR_014798	
Software, algorithm	pHeatmap, 1.0.12		RRID:SCR_016418	
Software, algorithm	Ensembl GRCm38, v99			
Software, algorithm	R, v4.0		RRID:SCR_001905	
Software, algorithm	Tximport, v1.16.0		RRID:SCR_016752	
Software, algorithm	SAMtools	SAMtools http://samtools.sourceforge.net/	RRID:SCR_002105	(Li et al., 2009)
Software, algorithm	Bedtools		RRID:SCR_006646	
Software, algorithm	STAR v2.5.3a	https://github.com/alexdobin/STAR	RRID:SCR_015899	
Software, algorithm	Sigmaplot, v12.5	Systat Software	RRID:SCR_003210	
Software, algorithm	Custom code for bioinformatics analysis	https://github.com/eric-d-larson/OE_TRPM5		

Overview of the method for transcriptional profiling of low abundance cell populations. For

transcriptional profiling of TRPM5-bearing MVC_eGFP cells and OSN_eGFP+ cells that

constitute a small fraction of the cells in the epithelium, we used FACS to separate the cell

populations targeted for RNAseq (Amamoto et al., 2019). In our experiments, we isolated the

cells from mice that expressed fluorescent marker proteins appropriate for cell sorting. OSNs

were expressing mCherry under the control of OMP promoter. eGFP was expressed in MVCs

and a subset of OSNs (OSN_eGFP+ cells) under control of the TRPM5 promoter.

Generation of OMP-H2B::Cherry mice. A PacI cassette containing PacI-H2B::mCherry-pA

PGK-puro-pA-PacI was inserted into an OMP gene-targeting vector (pPM9)(Mombaerts et

al., 1996), which replaces the OMP coding sequence with the PacI cassette and expresses a

H2B::mCherry fusion protein. Animals are maintained in a mixed 129/B6 background.

Animals. Mice with TRPM5-driven eGFP expression (Clapp et al.,2006) were crossed with

OMP-H2B::Cherry mice. The TRPM5-eGFP mice were obtained with written informed consent

from Dr. Robert Margolskee. Both lines were maintained separately as homozygous and

backcrossed regularly. Experiments were performed on mice from the F1 generation cross of

TRPM5-eGFP and OMP-H2B::Cherry mice (OMP-H2B::mCherry/TRPM5-eGFP). PCR was

used to verify genotype of experimental mice for eGFP and mCherry expression. Both male and

female mice were used for experiments with ages ranging from 3- 8 months. Estrous and cage

mate information was collected for all female mice in conjunction with experimental use. Mice

were housed in passive air exchange caging under a 12:12 light/dark cycle and were given food

and water *ad libitum*. Mice were housed in the National Institutes of Health approved Center for

Comparative Medicine at the University of Colorado Anschutz Medical Campus. All procedures

were performed in compliance with University of Colorado Anschutz Medical Campus

Institutional Animal Care and Use Committee (IACUC) that reviews the ethics of animal use.

Tissue dissociation of the olfactory epithelium. Following euthanasia via CO2 inhalation, the

olfactory epithelium was immediately removed from the nasal cavity and epithelial tissue was

separated from the bone in the turbinates. Care was taken not to include respiratory epithelium.

The epithelium was dissociated enzymatically with Dispase II (2 mg/ml) diluted in Ringer's

solution (145mM NaCl, 5mM KCL, 20mM HEPES, 1mM $MgCL_2$, 1mM $CaCl_2$, 1mM Ny-

Pyruvate, 5mM Glucose) (~25 minutes at 37^0C) followed by an incubation in a papain plus

Ca/Mg++ free Ringer's solution (Ca/Mg++ free Ringer's: 145mM NaCl, 5mM KCL, 20mM

HEPES, 1mM Ny-Pyruvate, 1mM EDTA , L-cysteine: 1mg L-cysteine /1.5mL Ca/Mg++ free

Ringer's, Papain:1-3ul/1mL Ca/Mg++ free Ringer's), for ~40-45 minutes at 37^0C. Following

incubation, DNase I (Promega) at $0.05U/\mu l$ and RNAse free 10x Reaction buffer (1:20) were

added to solution and the tissue was gently triturated using a ~1mm opening pipette. Isolated

OSNs were collected from supernatants via centrifugation and resuspended in cell sorting

medium of 1x PBS (diluted from commercial 10x PBS, pH 7.4) and BrainPhys Neuronal

Medium (Stemcell Technologies). Initially, isolated cells were examined with a confocal

microscope to confirm efficacy of dissociation methods, and examine cell types and

fluorescence. For RNAseq, cells were strained through a 40 μm cell strainer and kept on ice until

sorted via flow cytometry.

Flow cytometry. Fluorescence activated cell sorting was performed in the University of

Colorado Cancer Center Flow Cytometry Core on a Beckman Coulter MoFlo Astrios EQ using

MoFlo Astrios Summit Software (6.3.1.16945). eGFP signal was detected using a 488 nm laser

and a bandpass 526/52nm collection filter. mCherry signal was detected using a 561 nm laser

and a bandpass 614/20 nm collection filter. The 488nm laser was also used to detect light

scatter. The threshold was set at 3%. Gating was set to exclude doublets and optimized as cell

populations emerged based on fluorescent markers. Flow cytometry calibration beads for

AcGFP1/eGFP and mCherry (Takara, 632594, 632595) were used as fluorescence intensity

controls. Olfactory epithelium cell suspensions from wild type and OMP-H2B::Cherry mice or

TRPM5-eGFP mice were sorted as controls for auto fluorescence for eGFP and mCherry

populations respectively. Cells were sorted into RNAprotect Tissue Reagent (Qiagen).

RNA-extraction. Total RNA was extracted from sorted, pooled cells from each cell population

using the RNeasy Plus Micro Kit (Qiagen) according to the manufacturers recommended

protocol.

RT-qPCR. Quantitative reverse transcription polymerase chain reaction (RT-qPCR) was used to

assess and confirm identities of cell types from each of the sorted cell populations. Following

total RNA extraction, RT-qPCR was performed in the PCR core at University of Colorado

Anschutz Medical Campus for the following markers: OMP, TRPM5, eGFP and ChAT. Primers

and probes used for eGFP, TRPM5 and OMP were described in (Oshimoto et al., 2013).

Predesigned primers and probes for ChAT were purchased from Life Technologies. The mRNA

for these targets was measured by RT-qPCR using ABI QuantStudio 7 flex Sequence detector.

1µg total RNA was used to synthesize cDNA using the High Capacity c-DNA Reverse

Transcription kit (ABI-P/N 4368814). cDNA was diluted 1: 2 before PCR amplification.

The TaqMan probes were 5'labeled with 6-carboxyfluorescein (FAM). Real time PCR reactions

were carried out in MicroAmp optical tubes (PE ABI) in a 25 µl mix containing 8 % glycerol,

1X TaqMan buffer A (500 mM KCl, 100 mM Tris-HCl, 0.1 M EDTA, 600 nM passive

reference dye ROX, pH 8.3 at room temperature), 300 µM each of dATP, dGTP, dCTP and 600

µM dUTP, 5.5 mM MgCl2, 1X primer-probe mix, 1.25 U AmpliTaq Gold DNA and 5 µl

template cDNA. Thermal cycling conditions were as follows: Initiation was performed at 50ºC

for 2 min followed by activation of TaqGold at 95ºC for 10 min. Subsequently 40 cycles of

amplification were performed at 95ºC for 15 secs and 60ºC for 1 min. Experiments were

performed with duplicates for each data point. Each PCR run included the standard curve (10

fold serially diluted pooled cDNA from control and experimental samples), test samples, no-

template and NORT controls. The standard curve was then used to calculate the relative

amounts of targets in test samples. Quantities of targets in test samples were normalized to the

corresponding 18s rRNA (PE ABI, P/N 4308310).

RNA sequencing and pre-processing. RNA quality control, library preparation, and sequencing

were performed at the University of Colorado Genomics and Microarray core. Extracted RNA

was used as the input for the Nugen Universal Plus mRNA-seq kit (Redwood City, CA) to build

stranded sequencing libraries. Indexed libraries were sequenced using an Illumina

NovaSEQ6000. Library preparation and sequencing was performed in two batches, separated by

gender. 11 female samples were sequenced with an average depth of 37.3 million +/- SD of 6.5

million read pairs, and 25 male samples were sequenced with an average depth of 34.8 million

+/- SD of 3.5 million read pairs. Metadata for the samples submitted are shown in Figure 4 –

figure supplement 2. Raw BCL files were demultiplexed and converted to FASTQ format.

Trimming, filtering, and adapter contamination removal was performed using BBDuk

(Bushnell).

RNA Sequencing Analysis. Transcript abundance was quantified from trimmed and filtered

FASTQ files using Salmon v1.2.1(Patro et al., 2017) and a customized Ensembl GRCm38

(release 99) transcriptome (Zerbino et al., 2018). A customized version of the transcriptome was

prepared by appending FASTA sequences of eGFP and mCherry to the GRCm38 FASTA file.

The corresponding gene transfer format (GTF) file was modified accordingly to incorporate the

new transcripts. Transcript abundance was summarized at the gene level using the TxImport

(Soneson et al., 2015) package in R. Differential gene expression was quantified using DESeq2

(Love et al., 2014) with default parameters after removing genes with an average count of < 5

reads in each group. Significance was determined by FDR-adjusted p-value < 0.05. TopGO was

used for gene ontology analysis (Alexa and Rahnenfuhrer, 2020). The input to TopGO was a list

of significant DEGs and a list of all detected genes in the dataset. Enrichment was calculated by

dividing the number of detected genes by the number of expected genes within each ontology of

the TopGO output. To make the bar graphs in Figures 4 and 5, enrichment scores of

downregulated GO terms were multiplied by -1 for visualization. Heatmap visualization was

performed using *pHeatmap* in R (Kolde, 2019).

TRPM5 exon coverage. To visualize exon coverage of *Trpm5*, trimmed and filtered FASTQ

files were mapped to the Ensembl GRCm38 (version 99) using STAR (params, version

2.5.3a)(Dobin et al., 2013). The genome FASTA and GTF files were modified as described

above. Read coverage was summarized by *Trpm5* exon using a combination of Samtools ('depth'

and 'bedcov')(Li et al., 2009) and Bedtools (Quinlan and Hall, 2010). Data were compared using

a Scheirer-Ray-Hare test in R (*rcompanion* package).

Tissue Preparation for Fluorescence Microscopy and *in situ*. For euthanasia, mice were

anesthetized with ketamine/xylazine (20–100 _g/g of body weight), perfused transcardially with

0.1 M phosphate buffer (PBS) followed by a PBS-buffered fixative (EMS 32%

Paraformaldehyde aqueous solution diluted to 4% with 1x PBS) . The nose was harvested and

postfixed for 12 h before being transferred for cryoprotection into PBS with 20% sucrose

overnight. The olfactory epithelium was cryosectioned coronally into 16 μm -thick sections

mounted on Superfrost Plus slides (VWR, West Chester, PA).

In situ. *In situ* hybridization was performed with the hybridization chain reaction method (Choi

et al., 2018) using HCR v3.0 Probe Sets, Amplifiers, and Buffers from Molecular Instruments,

Inc. Frozen slides were allowed to thaw and dry, then immersed in 70% ethanol overnight at

4°C, and allowed to dry again completely. Slides were inverted and placed on a Plexiglas

platform inside a humidified chamber; subsequent steps were performed using this setup. Slides

were incubated in 10 μg/μl proteinase K for 10 minutes at 37°C, then pre-hybridized with HCR

hybridization buffer (30% formamide buffer from Molecular Instruments) for 10 minutes at

37°C. *Trpm5-B3* probes and *OMP-B2* probes (0.4 pmol of each probe in 100 μl HCR

hybridization buffer per slide) were added, and slides were hybridized overnight at 37°C. Slides

were briefly incubated in undiluted HCR Wash Buffer (30% formamide buffer from Molecular

Instruments) at 37°C. Excess probes were removed by incubating slides for 15 minutes each at

37°C in solutions of 75% HCR Wash Buffer / 25% SSCT (5X SSC, 0.1% Tween, diluted in

RNAse free water), 50% Buffer / 50% SSCT, 25% Buffer / 75% SSCT, and 100% SSCT. Slides

were incubated in Amplification Buffer (Molecular Instruments) at room temperature for 30

minutes. B3 hairpins labeled with Alexa Fluor 647 and B2 hairpins labeled with Alexa Fluor 546

were prepared (6 pmol of each hairpin were heat shocked, then cooled for 30 minutes, and added

to 100μl of Amplification Buffer) added to slides, and incubated overnight at room temperature.

Excess hairpins were removed with three washes in SSCT at room temperature. Tissue was

counterstained with DAPI, and slides were mounted using Invitrogen SlowFade Diamond

Antifade Mountant (Thermo Fisher Scientific).

Fluorescence microscopy. Microscopy was performed with confocal microscopes (Nikon A1R

or 3i Marianas). Images shown are flattened Z stacks. In situ images in Figure 3 were

deconvolved in DeconvolutionLab 2 (Sage et al., 2017) using 10 rounds of Lucy-Richardson

deconvolution (Lucy, 1974; Richardson, 1972) and a theoretical point spread function (PSF).

The theoretical PSF was calculated using the Gibson-Lanni model (Gibson and Lanni, 1989) for

325 nm lateral and 800 nm axial pixel size, wavelength of the reporter dye molecule, and an air-

immersion 20X NA 0.75 objective. Image was maximum projected and Gamma of 0.8 applied to

the TRMP5 channel for viewing.

Viral stocks. HSV-1, McKrae strain (gift from Dr. David Bloom, University of Florida), was

used for all experiments. Viral stocks were prepared in Vero cells; infectious particles were

purified from the supernatant, aliquoted, and stored at -80C. Viral titers were determined via

plaque forming unit (PFU) assays.

Viral infection. 3-6 month old OMP-H2B::mCherry/TRPM5-eGFP mice were lightly

anesthetized with isoflurane and inoculated intranasally with 10 μL of solution per naris (20 μL

total). The solution contained either HSV-1 ($1x\ 10^6$ PFU/naris; McKrae strain), or PBS (7.4 pH)

(sham control). Mice were randomly assigned to these groups. Mice were sacrificed at 5 days

post-infection (5 DPI). Controls included sham-treated mice (5), mice that were lightly

anesthetized (3), untreated mice housed in the BSL-2 facility (3) or the regular room (3). All

controls were used because ranksum tests corrected for multiple comparisons did not find

differences between these groups.

Statistical analysis. Statistical analysis was performed in Matlab (Mathworks, USA). Statistical

significance was estimated using a generalized linear model (GLM), with post-hoc tests for all

data pairs corrected for multiple comparisons using false discovery rate (Curran-Everett, 2000).

The post hoc comparisons between pairs of data were performed either with a t-test, or a ranksum

test, depending on the result of an Anderson-Darling test of normality. 95% CIs shown in the

figures as vertical black lines or shading bounding the lines were estimated by bootstrap analysis

of the mean by sampling with replacement 1000 times using the bootci function in MATLAB.

Declarations

Ethics approvals. Mouse experiments were carried out under guidelines of the National Institutes of Health in compliance with University of Colorado Anschutz Medical Campus Institutional Animal Care and Use Committee (IACUC).

Consent for publication. Not applicable.

Availability of data and materials. All data sequencing data are available in NCBI SRA https://www.ncbi.nlm.nih.gov/sra/PRJNA632936. The code used for bioinformatics analysis is found in https://github.com/eric-d-larson/OE_TRPM5

Competing interests. The authors declare no competing interests.

Funding. This work was supported by NIDCD DC014253 and NIA DC014253-04S1 (DR) and by the RNA Bioscience Initiative of the University of Colorado Anschutz Medical Campus (DS and DR). A Starr Stem Cell Grant (JA, AKH and PF) supported the production and characterization of the OMP-H2B::mCherry mouse strain. The funding bodies had no role in the experimental design or collection, analysis and interpretation of data or in writing the manuscript.

Authors' contributions. D.R., B.D.B., M.N. and V.R. conceptualized the project. B.D.B. performed FACS, qPCR and RNAseq experiments. E.D.L. performed genomic analysis. P.F.

generated the OMP-H2B::Cherry mice. D.S. designed and analyzed in situ experiments. M.N.,

A.N.B. and C.N. designed the HSV-1 inoculation experiments. C.N. and J.H.Jr. performed the

viral inoculation. M.L. performed literature search and wrote the section on viral infection in the

discussion. All authors contributed to writing and editing the manuscript.

Acknowledgements: We would like to acknowledge the support of Nicole Arevalo for laboratory

support, Jerome Artus and Anna-Katerina Hadjantonakis for the construction of the targeting

vector and production of OMP-H2B::Cherry mice.

Declaration of interests: The authors declare no competing interests.

References

Alexa, A., and Rahnenfuhrer, J. (2020). topGO: Enrichment Analysis for Gene Ontology (R

Package).

Allaire, A., Picard-Jean, F., and Bisaillon, M. (2015). Immunofluorescence to Monitor the

Cellular Uptake of Human Lactoferrin and its Associated Antiviral Activity Against the Hepatitis

C Virus. J Vis Exp.

Amamoto, R., Garcia, M.D., West, E.R., Choi, J., Lapan, S.W., Lane, E.A., Perrimon, N., and

Cepko, C.L. (2019). Probe-Seq enables transcriptional profiling of specific cell types from

heterogeneous tissue by RNA-based isolation. Elife *8*.

Bankova, L.G., Dwyer, D.F., Yoshimoto, E., Ualiyeva, S., McGinty, J.W., Raff, H., von Moltke,

J., Kanaoka, Y., Frank Austen, K., and Barrett, N.A. (2018). The cysteinyl leukotriene 3 receptor

regulates expansion of IL-25-producing airway brush cells leading to type 2 inflammation. Sci

Immunol *3*.

Bilinska, K., Jakubowska, P., Von Bartheld, C.S., and Butowt, R. (2020). Expression of the

SARS-CoV-2 Entry Proteins, ACE2 and TMPRSS2, in Cells of the Olfactory Epithelium:

Identification of Cell Types and Trends with Age. ACS Chemical Neuroscience.

Brann, D.H., Tsukahara, T., Weinreb, C., Lipovsek, M., Van den Berge, K., Gong, B., Chance,

R., Macaulay, I.C., Chou, H.-j., Fletcher, R.*, et al.* (2020). Non-neuronal expression of SARS-

CoV-2 entry genes in the olfactory system suggests mechanisms underlying COVID-19-

associated anosmia. bioRxiv, 2020.2003.2025.009084.

Brass, A.L., Huang, I.C., Benita, Y., John, S.P., Krishnan, M.N., Feeley, E.M., Ryan, B.J.,

Weyer, J.L., van der Weyden, L., Fikrig, E.*, et al.* (2009). The IFITM proteins mediate cellular

resistance to influenza A H1N1 virus, West Nile virus, and dengue virus. Cell *139*, 1243-1254.

Bushnell, B. BBMap (SourceForge).

Chen, M., Reed, R.R., and Lane, A.P. (2019). Chronic Inflammation Directs an Olfactory Stem

Cell Functional Switch from Neuroregeneration to Immune Defense. Cell Stem Cell *25*, 501-513

e505.

Cheshenko, N., Pierce, C., and Herold, B.C. (2018). Herpes simplex viruses activate

phospholipid scramblase to redistribute phosphatidylserines and Akt to the outer leaflet of the

plasma membrane and promote viral entry. PLoS Pathog *14*, e1006766.

Choi, H.M., Schwarzkopf, M., Fornace, M.E., Acharya, A., Artavanis, G., Stegmaier, J., Cunha,

A., and Pierce, N.A. (2018). Third-generation in situ hybridization chain reaction: multiplexed,

quantitative, sensitive, versatile, robust. Development *145*, dev165753.

Chuang, Y.C., Lee, C.H., Sun, W.H., and Chen, C.C. (2018). Involvement of advillin in

somatosensory neuron subtype-specific axon regeneration and neuropathic pain. Proc Natl Acad

Sci U S A *115*, E8557-E8566.

Clapp, T.R., Medler, K.F., Damak, S., Margolskee, R.F., and Kinnamon, S.C. (2006). Mouse

taste cells with G protein-coupled taste receptors lack voltage-gated calcium channels and

SNAP-25. BMC Biol *4*, 7.

Curran-Everett, D. (2000). Multiple comparisons: philosophies and illustrations. AmJPhysiol

RegulIntegrComp Physiol *279*, R1-R8.

Damak, S., Rong, M., Yasumatsu, K., Kokrashvili, Z., Varadarajan, V., Zou, S., Jiang, P.,

Ninomiya, Y., and Margolskee, R.F. (2003). Detection of sweet and umami taste in the absence

of taste receptor T1r3. Science *301*, 850-853.

Dando, S.J., Mackay-Sim, A., Norton, R., Currie, B.J., St John, J.A., Ekberg, J.A.K., Batzloff,

M., Ulett, G.C., and Beacham, I.R. (2014). Pathogens Penetrating the Central Nervous System:

Infection Pathways and the Cellular and Molecular Mechanisms of Invasion. Clinical

Microbiology Reviews *27*, 691-726.

De Felice, F.G., Tovar-Moll, F., Moll, J., Munoz, D.P., and Ferreira, S.T. (2020). Severe Acute

Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) and the Central Nervous System. Trends in

Neurosciences.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,

M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*,

15-21.

Dong, C., Sun, X., Guan, Z., Zhang, M., and Duan, M. (2017). Modulation of influenza A virus

replication by microRNA-9 through targeting MCPIP1. J Med Virol *89*, 41-48.

Doty, R.L. (2008). The olfactory vector hypothesis of neurodegenerative disease: is it viable?

Ann Neurol *63*, 7-15.

Farbman, A.I., and Margolis, F.L. (1980). Olfactory marker protein during ontogeny:

immunohistochemical localization. Developmental biology *74*, 205-215.

Fischl, A.M., Heron, P.M., Stromberg, A.J., and McClintock, T.S. (2014). Activity-dependent

genes in mouse olfactory sensory neurons. Chem Senses *39*, 439-449.

Fodoulian, L., Tuberosa, J., Rossier, D., Landis, B.N., Carleton, A., and Rodriguez, I. (2020).

SARS-CoV-2 receptor and entry genes are expressed by sustentacular cells in the human

olfactory neuroepithelium. bioRxiv, 2020.2003.2031.013268.

Fu, Z., Ogura, T., Luo, W., and Lin, W. (2018). ATP and Odor Mixture Activate TRPM5-

Expressing Microvillous Cells and Potentially Induce Acetylcholine Release to Enhance

Supporting Cell Endocytosis in Mouse Main Olfactory Epithelium. Front Cell Neurosci *12*, 71.

Gack, M.U., Albrecht, R.A., Urano, T., Inn, K.S., Huang, I.C., Carnero, E., Farzan, M., Inoue,

S., Jung, J.U., and Garcia-Sastre, A. (2009). Influenza A virus NS1 targets the ubiquitin ligase

TRIM25 to evade recognition by the host viral RNA sensor RIG-I. Cell Host Microbe *5*, 439-

449.

Gack, M.U., Shin, Y.C., Joo, C.H., Urano, T., Liang, C., Sun, L., Takeuchi, O., Akira, S., Chen,

Z., Inoue, S., *et al.* (2007). TRIM25 RING-finger E3 ubiquitin ligase is essential for RIG-I-

mediated antiviral activity. Nature *446*, 916-920.

Genovese, F., and Tizzano, M. (2018). Microvillous cells in the olfactory epithelium express

elements of the solitary chemosensory cell transduction signaling cascade. PLoS One *13*,

e0202754.

Gerbe, F., Sidot, E., Smyth, D.J., Ohmoto, M., Matsumoto, I., Dardalhon, V., Cesses, P.,

Garnier, L., Pouzolles, M., Brulin, B., *et al.* (2016). Intestinal epithelial tuft cells initiate type 2

mucosal immunity to helminth parasites. Nature *529*, 226-230.

Giacomelli, A., Pezzati, L., Conti, F., Bernacchia, D., Siano, M., Oreni, L., Rusconi, S.,

Gervasoni, C., Ridolfo, A.L., Rizzardini, G., *et al.* (2020). Self-reported olfactory and taste

disorders in SARS-CoV-2 patients: a cross-sectional study. Clin Infect Dis.

Gibson, S.F., and Lanni, F. (1989). Diffraction by a circular aperture as a model for three-

dimensional optical microscopy. J Opt Soc Am A *6*, 1357-1367.

Gong, Q., Cheng, M., Chen, H., Liu, X., Si, Y., Yang, Y., Yuan, Y., Jin, C., Yang, W., He, F., *et*

al. (2011). Phospholipid scramblase 1 mediates hepatitis C virus entry into host cells. FEBS Lett

585, 2647-2652.

Hegg, C.C., Jia, C., Chick, W.S., Restrepo, D., and Hansen, A. (2010). Microvillous cells

expressing IP3 receptor type 3 in the olfactory epithelium of mice. Eur J Neurosci *32*, 1632-

1645.

Hoffmann, M., Kleine-Weber, H., Schroeder, S., Kruger, N., Herrler, T., Erichsen, S.,

Schiergens, T.S., Herrler, G., Wu, N.H., Nitsche, A*., et al.* (2020). SARS-CoV-2 Cell Entry

Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. Cell

181, 271-280 e278.

Jablonka-Shariff, A., Lu, C.Y., Campbell, K., Monk, K.R., and Snyder-Warwick, A.K. (2020).

Gpr126/Adgrg6 contributes to the terminal Schwann cell response at the neuromuscular junction

following peripheral nerve injury. Glia *68*, 1182-1200.

Juilfs, D.M., Fulle, H.J., Zhao, A.Z., Houslay, M.D., Garbers, D.L., and Beavo, J.A. (1997). A

subset of olfactory neurons that selectively express cGMP-stimulated phosphodiesterase (PDE2)

and guanylyl cyclase-D define a unique olfactory signal transduction pathway. Proc Natl Acad

Sci U S A *94*, 3388-3395.

Kim, B.E., Bin, L., Ye, Y.M., Ramamoorthy, P., and Leung, D.Y.M. (2013). IL-25 enhances

HSV-1 replication by inhibiting filaggrin expression, and acts synergistically with Th2 cytokines

to enhance HSV-1 replication. J Invest Dermatol *133*, 2678-2685.

Kolde, R. (2019). pheatmap: Pretty Heatmaps (cran.r-project.org).

Kopp, S.J., Banisadr, G., Glajch, K., Maurer, U.E., Grunewald, K., Miller, R.J., Osten, P., and

Spear, P.G. (2009). Infection of neurons and encephalitis after intracranial inoculation of herpes

simplex virus requires the entry receptor nectin-1. Proc Natl Acad Sci U S A *106*, 17916-17920.

Lee, W.J., Fu, R.M., Liang, C., and Sloan, R.D. (2018). IFITM proteins inhibit HIV-1 protein

synthesis. Sci Rep *8*, 14551.

Lemons, K., Fu, Z., Aoude, I., Ogura, T., Sun, J., Chang, J., Mbonu, K., Matsumoto, I.,

Arakawa, H., and Lin, W. (2017). Lack of TRPM5-Expressing Microvillous Cells in Mouse

Main Olfactory Epithelium Leads to Impaired Odor-Evoked Responses and Olfactory-Guided

Behavior in a Challenging Chemical Environment. eNeuro *4*.

Lemons, K., Fu, Z., Ogura, T., and Lin, W. (2020). TRPM5-expressing Microvillous Cells

Regulate Region-specific Cell Proliferation and Apoptosis During Chemical Exposure.

Neuroscience *434*, 171-190.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,

Durbin, R., and Genome Project Data Processing, S. (2009). The Sequence Alignment/Map

format and SAMtools. Bioinformatics *25*, 2078-2079.

Li, M., Yang, J., Zhao, Y., Song, Y., Yin, S., Guo, J., Zhang, H., Wang, K., Wei, L., Li, S*., et al.*

(2020). MCPIP1 inhibits Hepatitis B virus replication by destabilizing viral RNA and negatively

regulates the virus-induced innate inflammatory responses. Antiviral Res *174*, 104705.

Liberles, S.D. (2015). Trace amine-associated receptors: ligands, neural circuits, and behaviors.

Current opinion in neurobiology *34*, 1-7.

Lin, R.J., Chu, J.S., Chien, H.L., Tseng, C.H., Ko, P.C., Mei, Y.Y., Tang, W.C., Kao, Y.T.,

Cheng, H.Y., Liang, Y.C.*, et al.* (2014). MCPIP1 suppresses hepatitis C virus replication and

negatively regulates virus-induced proinflammatory cytokine responses. J Immunol *193*, 4159-

4168.

Lin, W., Ezekwe, E.A., Jr., Zhao, Z., Liman, E.R., and Restrepo, D. (2008). TRPM5-expressing

microvillous cells in the main olfactory epithelium. BMC Neurosci *9*, 114.

Lin, W., Margolskee, R., Donnert, G., Hell, S.W., and Restrepo, D. (2007). Olfactory neurons

expressing transient receptor potential channel M5 (TRPM5) are involved in sensing

semiochemicals. Proc Natl Acad Sci U S A *104*, 2471-2476.

Lopez, F., Delgado, R., Lopez, R., Bacigalupo, J., and Restrepo, D. (2014). Transduction for

Pheromones in the Main Olfactory Epithelium Is Mediated by the Ca2+-Activated Channel

TRPM5. J Neurosci *34*, 3268-3278.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.

Lucy, L.B. (1974). An iterative technique for the rectification of observed distributions. The

Astronomical Journal *79*, 745.

Luo, W., Zhang, J., Liang, L., Wang, G., Li, Q., Zhu, P., Zhou, Y., Li, J., Zhao, Y., Sun, N., *et al.*

(2018). Phospholipid scramblase 1 interacts with influenza A virus NP, impairing its nuclear

import and thereby suppressing virus replication. PLoS Pathog *14*, e1006851.

Luo, X.C., Chen, Z.H., Xue, J.B., Zhao, D.X., Lu, C., Li, Y.H., Li, S.M., Du, Y.W., Liu, Q.,

Wang, P., *et al.* (2019). Infection by the parasitic helminth Trichinella spiralis activates a Tas2r-

mediated signaling pathway in intestinal tuft cells. Proc Natl Acad Sci U S A *116*, 5564-5569.

Maina, I.W., Workman, A.D., and Cohen, N.A. (2018). The role of bitter and sweet taste

receptors in upper airway innate immunity: Recent advances and future directions. World J

Otorhinolaryngol Head Neck Surg *4*, 200-208.

McGinty, J.W., Ting, H.-A., Billipp, T.E., Nadjsombati, M.S., Khan, D.M., Barrett, N.A., Liang,

H.-E., Matsumoto, I., and von Moltke, J. (2020). Tuft-Cell-Derived Leukotrienes Drive Rapid

Anti-helminth Immunity in the Small Intestine but Are Dispensable for Anti-protist Immunity.

Immunity *52*, 528-541.e527.

McLaughlin, S.K., McKinnon, P.J., and Margolskee, R.F. (1992). Gustducin is a taste-cell-

specific G protein closely related to the transducins. Nature *357*, 563-569.

Meyerson, N.R., Zhou, L., Guo, Y.R., Zhao, C., Tao, Y.J., Krug, R.M., and Sawyer, S.L. (2017).

Nuclear TRIM25 Specifically Targets Influenza Virus Ribonucleoproteins to Block the Onset of

RNA Chain Elongation. Cell Host Microbe *22*, 627-638 e627.

Mobley, A.S., Miller, A.M., Araneda, R.C., Maurer, L.R., Muller, F., and Greer, C.A. (2010).

Hyperpolarization-activated cyclic nucleotide-gated channels in olfactory sensory neurons

regulate axon extension and glomerular formation. The Journal of neuroscience : the official

journal of the Society for Neuroscience *30*, 16498-16508.

Mombaerts, P., Wang, F., Dulac, C., Chao, S.K., Nemes, A., Mendelsohn, M., Edmondson, J.,

and Axel, R. (1996). Visualizing an olfactory sensory map. Cell *87*, 675-686.

Narayana, S.K., Helbig, K.J., McCartney, E.M., Eyre, N.S., Bull, R.A., Eltahla, A., Lloyd, A.R.,

and Beard, M.R. (2015). The Interferon-induced Transmembrane Proteins, IFITM1, IFITM2,

and IFITM3 Inhibit Hepatitis C Virus Entry. J Biol Chem *290*, 25946-25959.

Noyce, R.S., and Richardson, C.D. (2012). Nectin 4 is the epithelial cell receptor for measles

virus. Trends Microbiol *20*, 429-439.

O'Leary, C.E., Schneider, C., and Locksley, R.M. (2019). Tuft Cells-Systemically Dispersed

Sensory Epithelia Integrating Immune and Neural Circuitry. Annu Rev Immunol *37*, 47-72.

Ogura, T., Szebenyi, S.A., Krosnowski, K., Sathyanesan, A., Jackson, J., and Lin, W. (2011).

Cholinergic microvillous cells in the mouse main olfactory epithelium and effect of acetylcholine

on olfactory sensory neurons and supporting cells. J Neurophysiol *106*, 1274-1287.

Omura, M., and Mombaerts, P. (2014). Trpc2-Expressing Sensory Neurons in the Main

Olfactory Epithelium of the Mouse. Cell Reports *8*, 583-595.

Oshimoto, A., Wakabayashi, Y., Garske, A., Lopez, R., Rolen, S., Flowers, M., Arevalo, N., and

Restrepo, D. (2013). Potential role of transient receptor potential channel M5 in sensing putative

pheromones in mouse olfactory sensory neurons. PLoS One *8*, e61990.

Parma, V., Ohla, K., Veldhuizen, M.G., Nim, M.Y., Kelly, C.E., Reed, D.R., Hummel, T.,

Munger, S., and Hayes, J.E. (2020). More than just smell - COVID-19 is associated with severe

impairment of smell, taste, and chemesthesis. medRxiv, 2020.2005.2004.20090902.

Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., and Kingsford, C. (2017). Salmon provides fast

and bias-aware quantification of transcript expression. Nature methods *14*, 417-419.

Perniss, A., Liu, S., Boonen, B., Keshavarz, M., Ruppert, A.L., Timm, T., Pfeil, U., Soultanova,

A., Kusumakshi, S., Delventhal, L.*, et al.* (2020). Chemosensory Cell-Derived Acetylcholine

Drives Tracheal Mucociliary Clearance in Response to Virulence-Associated Formyl Peptides.

Immunity *52*, 683-699 e611.

Petermann, P., Rahn, E., Thier, K., Hsu, M.J., Rixon, F.J., Kopp, S.J., and Knebel-Morsdorf, D.

(2015). Role of Nectin-1 and Herpesvirus Entry Mediator as Cellular Receptors for Herpes

Simplex Virus 1 on Primary Murine Dermal Fibroblasts. J Virol *89*, 9407-9416.

Pyrski, M., Eckstein, E., Schmid, A., Bufe, B., Weiss, J., Chubanov, V., Boehm, U., and Zufall,

F. (2017). Trpm5 expression in the olfactory epithelium. Mol Cell Neurosci *80*, 75-88.

Pyrski, M., Tusty, M., Eckstein, E., Oboti, L., Rodriguez-Gil, D.J., Greer, C.A., and Zufall, F.

(2018). P/Q Type Calcium Channel Cav2.1 Defines a Unique Subset of Glomeruli in the Mouse

Olfactory Bulb. Frontiers in Cellular Neuroscience *12*, 295.

Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing

genomic features. Bioinformatics *26*, 841-842.

Rane, C.K., Jackson, S.R., Pastore, C.F., Zhao, G., Weiner, A.I., Patel, N.N., Herbert, D.R.,

Cohen, N.A., and Vaughan, A.E. (2019). Development of solitary chemosensory cells in the

distal lung after severe influenza injury. Am J Physiol Lung Cell Mol Physiol *316*, L1141-

L1149.

Richardson, W.H. (1972). Bayesian-Based Iterative Method of Image Restoration*. J Opt Soc

Am *62*, 55-59.

Sage, D., Donati, L., Soulez, F., Fortun, D., Schmit, G., Seitz, A., Guiet, R., Vonesch, C., and

Unser, M. (2017). DeconvolutionLab2: An open-source software for deconvolution microscopy.

Methods *115*, 28-41.

Saunders, C.J., Christensen, M., Finger, T.E., and Tizzano, M. (2014). Cholinergic

neurotransmission links solitary chemosensory cells to nasal inflammation. Proc Natl Acad Sci U

S A *111*, 6075-6080.

Sayers, C.L., and Elliott, G. (2016). Herpes Simplex Virus 1 Enters Human Keratinocytes by a

Nectin-1-Dependent, Rapid Plasma Membrane Fusion Pathway That Functions at Low

Temperature. J Virol *90*, 10379-10389.

Sekerkova, G., Zheng, L., Loomis, P.A., Changyaleket, B., Whitlon, D.S., Mugnaini, E., and

Bartles, J.R. (2004). Espins are multifunctional actin cytoskeletal regulatory proteins in the

microvilli of chemosensory and mechanosensory cells. J Neurosci *24*, 5445-5456.

Shestakov, A., Jenssen, H., Nordstrom, I., and Eriksson, K. (2012). Lactoferricin but not

lactoferrin inhibit herpes simplex virus type 2 infection in mice. Antiviral Res *93*, 340-345.

Shivkumar, M., Milho, R., May, J.S., Nicoll, M.P., Efstathiou, S., and Stevenson, P.G. (2013).

Herpes simplex virus 1 targets the murine olfactory neuroepithelium for host entry. J Virol *87*,

10477-10488.

Shukla, N.D., Tiwari, V., and Valyi-Nagy, T. (2012). Nectin-1-specific entry of herpes simplex

virus 1 is sufficient for infection of the cornea and viral spread to the trigeminal ganglia. Mol Vis

18, 2711-2716.

Singh, B.K., Hornick, A.L., Krishnamurthy, S., Locke, A.C., Mendoza, C.A., Mateo, M., Miller-

Hunt, C.L., Cattaneo, R., and Sinn, P.L. (2015). The Nectin-4/Afadin Protein Complex and

Intercellular Membrane Pores Contribute to Rapid Spread of Measles Virus in Primary Human

Airway Epithelia. J Virol *89*, 7089-7096.

Singh, B.K., Li, N., Mark, A.C., Mateo, M., Cattaneo, R., and Sinn, P.L. (2016). Cell-to-Cell

Contact and Nectin-4 Govern Spread of Measles Virus from Primary Human Myeloid Cells to

Primary Human Airway Epithelial Cells. J Virol *90*, 6808-6817.

Soneson, C., Love, M.I., and Robinson, M.D. (2015). Differential analyses for RNA-seq:

transcript-level estimates improve gene-level inferences. F1000Res *4*, 1521.

Tizzano, M., Gulbransen, B.D., Vandenbeuch, A., Clapp, T.R., Herman, J.P., Sibhatu, H.M.,

Churchill, M.E., Silver, W.L., Kinnamon, S.C., and Finger, T.E. (2010). Nasal chemosensory

cells use bitter taste signaling to detect irritants and bacterial signals. Proc Natl Acad Sci U S A

107, 3210-3215.

Ualiyeva, S., Hallen, N., Kanaoka, Y., Ledderose, C., Matsumoto, I., Junger, W.G., Barrett,

N.A., and Bankova, L.G. (2020). Airway brush cells generate cysteinyl leukotrienes through the

ATP sensor P2Y2. Science Immunology *5*, eaax7224.

Valimaa, H., Tenovuo, J., Waris, M., and Hukkanen, V. (2009). Human lactoferrin but not

lysozyme neutralizes HSV-1 and inhibits HSV-1 replication and cell-to-cell spread. Virol J *6*, 53.

van der Linden, C., Jakob, S., Gupta, P., Dulac, C., and Santoro, S.W. (2018). Sex separation

induces differences in the olfactory sensory receptor repertoires of male and female mice. Nat

Commun *9*, 5081.

Villar, P.S., Delgado, R., Vergara, C., Reyes, J.G., and Bacigalupo, J. (2017). Energy

Requirements of Odor Transduction in the Chemosensory Cilia of Olfactory Sensory Neurons

Rely on Oxidative Phosphorylation and Glycolytic Processing of Extracellular Glucose. J

Neurosci *37*, 5736-5743.

von Moltke, J., Ji, M., Liang, H.E., and Locksley, R.M. (2016). Tuft-cell-derived IL-25 regulates

an intestinal ILC2-epithelial response circuit. Nature *529*, 221-225.

Wu, Y., Ma, L., Duyck, K., Long, C.C., Moran, A., Scheerer, H., Blanck, J., Peak, A., Box, A.,

Perera, A*., et al.* (2018). A Population of Navigator Neurons Is Essential for Olfactory Map

Formation during the Critical Period. Neuron *100*, 1066-1082 e1066.

Yamaguchi, T., Yamashita, J., Ohmoto, M., Aoude, I., Ogura, T., Luo, W., Bachmanov, A.A.,

Lin, W., Matsumoto, I., and Hirota, J. (2014). Skn-1a/Pou2f3 is required for the generation of

Trpm5-expressing microvillous cells in the mouse main olfactory epithelium. BMC Neurosci *15*,

13.

Yamashita, J., Ohmoto, M., Yamaguchi, T., Matsumoto, I., and Hirota, J. (2017). Skn-1a/Pou2f3

functions as a master regulator to generate Trpm5-expressing chemosensory cells in mice. PLoS

One *12*, e0189340.

Yan, C.H., Faraji, F., Prajapati, D.P., Boone, C.E., and DeConde, A.S. (2020a). Association of

chemosensory dysfunction and Covid-19 in patients presenting with influenza-like symptoms.

Int Forum Allergy Rhinol.

Yan, C.H., Faraji, F., Prajapati, D.P., Ostrander, B.T., and DeConde, A.S. (2020b). Self-reported

olfactory loss associates with outpatient clinical course in Covid-19. Int Forum Allergy Rhinol.

Yang, J., Zhu, X., Liu, J., Ding, X., Han, M., Hu, W., Wang, X., Zhou, Z., and Wang, S. (2012).

Inhibition of Hepatitis B virus replication by phospholipid scramblase 1 in vitro and in vivo.

Antiviral Res *94*, 9-17.

Yu, J., Li, M., Wilkins, J., Ding, S., Swartz, T.H., Esposito, A.M., Zheng, Y.M., Freed, E.O.,

Liang, C., Chen, B.K.*, et al.* (2015). IFITM Proteins Restrict HIV-1 Infection by Antagonizing

the Envelope Glycoprotein. Cell Rep *13*, 145-156.

Zerbino, D.R., Achuthan, P., Akanni, W., Amode, M.R., Barrell, D., Bhai, J., Billis, K.,

Cummins, C., Gall, A., Giron, C.G.*, et al.* (2018). Ensembl 2018. Nucleic Acids Res *46*, D754-

D761.

Zhang, Y., Hoon, M.A., Chandrashekar, J., Mueller, K.L., Cook, B., Wu, D., Zuker, C.S., and

Ryba, N.J. (2003). Coding of sweet, bitter, and umami tastes: different receptor cells sharing

similar signaling pathways. Cell *112*, 293-301.

Zheng, Y., Qin, Z., Ye, Q., Chen, P., Wang, Z., Yan, Q., Luo, Z., Liu, X., Zhou, Y., Xiong, W.*,

et al.* (2014). Lactoferrin suppresses the Epstein-Barr virus-induced inflammatory response by

interfering with pattern recognition of TLR2 and TLR9. Lab Invest *94*, 1188-1199.

Zheng, Y., Zhang, W., Ye, Q., Zhou, Y., Xiong, W., He, W., Deng, M., Zhou, M., Guo, X.,

Chen, P.*, et al.* (2012). Inhibition of Epstein-Barr virus infection by lactoferrin. J Innate Immun

4, 387-398.

Ziegler, C.G.K., Allon, S.J., Nyquist, S.K., Mbano, I.M., Miao, V.N., Tzouanas, C.N., Cao, Y.,

Yousif, A.S., Bals, J., Hauser, B.M.*, et al.* (2020). SARS-CoV-2 receptor ACE2 is an interferon-

stimulated gene in human airway epithelial cells and is detected in specific cell subsets across

tissues. Cell.

Figure legends

Figure 1. Fluorescence activated sorting (FACS) of cells isolated from the olfactory

epithelium.

a. TRPM5 promoter driven expression of eGFP and OMP promoter driven expression of

mCherry in the olfactory epithelium. Expression of eGFP is found both in MVCs that do not

express mCherry (asterisk) and in OSNs double labeled with eGFP and mCherry (arrow). **i.**

Composite, **ii.** eGFP, **iii.** mCherry, **iv**. Composite magnification. Magenta: mCherry, green:

eGFP. Scale bar: i-iii, 50 mm, iv, 10 mm.

b. Schematic of RNA-seq process from tissue to RNA extraction. Mouse OE was dissociated

into single cells and sorted via FACS. RNA was extracted from each of the resulting cell

populations.

c. Two isolated OSNs differing in eGFP expression. Magenta: mCherry, green: eGFP. Scale bar:

10 mm.

d. Distribution of mCherry and eGFP fluorescence intensity for FACS-sorted cells. Three cell

populations were isolated for RNAseq: Cells with low OMP promoter-driven mCherry

expression and high TRPM5 promoter-driven eGFP expression (MVC_eGFP cells), cells with

high OMP promoter-driven mCherry and low eGFP expression (OSN_eGFP- cells) and cells

with eGFP expression of the same magnitude as MVC_eGFP cells and high OMP promoter-

driven mCherry expression (OSN_eGFP+ cells). The number of cells collected for this FACS

run were: OSN_eGFP-s 1,500,000, OSN_eGFP+s 5336 and MVC_eGFP cells 37,178.

e. qPCR levels (normalized to levels 18s RNA) for expression of transcripts encoding for OMP

(**i**), TRPM5 (**ii**), eGFP (**iii**) and ChAT (**iv**). The asterisks denote significant differences tested

with either t-test or ranksum with p-values below the significance p-value corrected for multiple

comparisons using the false discovery rate (pFDR)(Curran-Everett, 2000). pFDR is 0.033 for

OMP, 0.05 for TRPM5, 0.05 for eGFP and 0.03 for ChAT, n=8 for OMP OSN_eGFP-s, 4 for

OMP OSN_eGFP+s and 4 for MVC_eGFP cells.

1015

Figure 2. Decreased yield of OSN_eGFP+ cells when mice are moved from low ventilation

(LV) to high ventilation (HV) cages.

a and b. Distribution of mCherry and eGFP fluorescence intensity for FACS-sorted cells that

either were not transferred to HV cages (**a**) or were transferred to HV cages for 22 days before

sorting (**b**).

c. Time course showing change in the number of sorted OSN_eGFP+s after mice were

transferred to HV cages.

d and e. Dependence of the yield of OSN_eGFP+ cells after sorting on the number of days in

LV cages (**d**) or the number of days in HV cages (**e**).

1025

Figure 3. Coverage of TRPM5 transcript by RNAseq in the different cell groups sorted

from the olfactory epithelium

a. In situ for TRPM5 and OMP transcripts in the olfactory epithelium shows strong label for

TRPM5 in MVCs (asterisks) and sparse labeling in the OSN nuclear layer (arrows). **i.** OMP, **ii.**

TRPM5, **iii.** Composite. The scale bar is 100 μm.

b and c. Coverage of the reads for the TRPM5 transcript for the different cell groups. Read

coverage was computed with 'Samtools depth' over the region corresponding to the Trpm5 gene.

b. Coverage per exon. All samples normalized to TRPM5 reads show distribution among exons

with little difference between MVC and OSN_eGFP+ cells.

c. Coverage shown per base pair as a function of location in the genomic DNA. Plots show read

depth per base pair for each cell type normalized to total number of mapped reads. Gray shading

indicates 95% confidence interval.

1038

Figure 4. RNAseq comparison of OSN_eGFP- and OSN_eGFP+ cells.

a. Heatmap showing the top 10 upregulated and top 10 downregulated genes identified by

DESeq2.

b. Heatmap showing all *Olfr* genes detected in the data.

For both a and b, row and column order were determined automatically by the *pHeatmap*

package in R. For each data point relative expression was calculated by subtracting the average

row value from each individual value.

c. Volcano plot of all Olfactory receptors, demonstrating the small number of enriched olfactory

receptors in the OSN_eGFP+ population.

d. Hierarchical clustering of transcripts for taste transduction and transcripts expressed in

canonical and non-canonical OSNs identified by RNAseq as significantly different in expression

between the cell groups. We compared expression of transcripts involved in taste transduction,

canonical olfactory transduction, and non-canonical OSNs. The non-canonical OSNs considered

here included guanilyl-cyclase D (GC-D) OSNs (Juilfs et al., 1997), Trpc2 OSNs (Omura and

Mombaerts, 2014), Cav2.1 OSNs (Pyrski et al., 2018), and OSNs expressing trace amine-

associated receptors (Taars) (Liberles, 2015). Transcripts identified by DESeq2.

e. Gene ontology (GO) term enrichment was calculated from differentially expressed genes using

TopGO in R. An enrichment value for genes with Fischer p value <0.05 was calculated by

dividing the number of expressed genes within the GO term by the number expected genes (by

random sampling, determined by *TopGO*).

1059

Figure 5. RNAseq comparison of OSN_eGFP- and OSN_eGFP+ vs. MVC_eGFP cells.

a and b. Heatmaps showing hierarchical clustering of the top 10 upregulated and top 10

downregulated genes identified by DESeq2. **a.** OSN_eGFP- vs. MVC_eGFP cells. **b.**

OSN_eGFP+ vs. MVC_eGFP cells. For both a and b, row and column order were determined

automatically by the *pHeatmap* package in R. For each data point relative expression was

calculated by subtracting the average row value from each individual value.

c and d. Gene ontology (GO) term enrichment was calculated from differentially expressed

genes using *TopGO* in R. **c.** OSN_eGFP- vs. MVC_eGFP cells. **d**. OSN_eGFP+ vs.

MVC_eGFP cells. An enrichment value for genes with Fischer p value <0.05 was calculated by

dividing the number of expressed genes within the GO term by the number expected genes (by

random sampling, determined by *TopGO*).

1071

1072   **Figure 6. Significant differences in virally-related gene ontology.** Heatmaps show hierarchical

1073   clustering of significantly differentially expressed genes identified by DESeq2. **a.** MVC_eGFP

1074   cells compared to OSN_eGFP-. **b.** OSN_eGFP+ compared to OSN_eGFP-.

1075

Figure 7. Intranasal inoculation with HSV-1 elicits a decrease in OSNs.

a and b. Distribution of mCherry and eGFP fluorescence intensity for FAC-sorted cells that

either were not infected (**a**) or were inoculated intranasally with 10 μL of solution per naris (20

μL total with HSV-1 (1x 10^6 PFU/naris; McKrae strain) 5 DPI (**b**).

c. Fraction of total cells for OSN_eGFP- (**i**), OSN_eGFP+ (**ii**) or MVC (**iii**) cell groups. GLM

analysis yielded significant effects for the fraction of OSN_eGFP- compared to MVCs (p<0.001)

and for the interaction between the fraction of OSN_eGFP- compared to MVCs and HSV-1

treatment (p<0.001, 75 observations, 69 d.f., GLM F-statistic=111 and p-value < 0.001, n=14

mice for control, n=11 mice for HSV-1 inoculation). Asterisks denote a significant difference

post-hoc with t-test p-value < pFDR = 0.047.

1086

1087

Name	OSN_eGFP-	OSN_eGFP+	MVC_eGFP	p-value adjusted
Olfr292	3.61	959	4.29	6.35E-09
Olfr282	2.05	486	0	8.01E-05
Olfr1434	53.3	7730	0	9.71E-16
Olfr390	101	10800	43.1	1.56E-16
Olfr305	6.14	612	0	6.3E-12
Olfr293	6.96	664	16.9	1.42E-07
Olfr378	3.41	322	0	1.1E-06
Olfr128	39.6	3660	12.7	7.33E-14
Olfr344	16.2	1050	0	1.4E-11
Olfr307	7.59	393	0	3.76E-06
Olfr391	156	8000	9.79	1.01E-15
Olfr299	13.1	651	0	4.58E-09
Olfr142	36.4	1720	3.77	1.62E-08
Olfr1	147	5720	52.7	3.08E-10
Olfr1279	16.4	552	10.9	3.52E-07
Olfr39	13.8	388	21	2.81E-06
Olfr1447	64.1	1610	0	1.23E-07
Olfr728	2150	45700	320	6.13E-22
Olfr727	560	11000	179	1.64E-07
Olfr1555-ps1	10.1	175	0	0.0397
Olfr346	27.6	465	0	3.85E-05
Olfr1228	533	5320	35.4	3.09E-08
Olfr1181	87.4	766	4.61	0.000766
Olfr943	97.1	844	2.89	0.000886
Olfr298	60.1	509	0	0.00132

1088

1089 **Table 1. Levels of expression and adjusted p-value for the olfactory receptor genes whose**

1090 **levels are significantly higher in OSN_eGFP+ compared to OSN_eGFP-.** These olfactory

1091 receptors had an adjusted p-value for expression level difference between OSN_eGFP+

1092 compared to OSN_eGFP- and had a fold change > 4 and average expression > 100 counts.

1093

1094

1095

1096

Figure 1. Fluorescence activated sorting (FACS) of cells isolated from the olfactory

epithelium.

a. TRPM5 promoter driven expression of eGFP and OMP promoter driven expression of

mCherry in the olfactory epithelium. Expression of eGFP is found both in MVCs that do not

express mCherry (asterisk) and in OSNs double labeled with eGFP and mCherry (arrow). **i.**

Composite, **ii.** eGFP, **iii.** mCherry, **iv**. Composite magnification. Magenta: mCherry, green:

eGFP. Scale bar: i-iii, 50 μm, iv, 10 μm.

b. Schematic of RNA-seq process from tissue to RNA extraction. Mouse OE was dissociated

into single cells and sorted via FACS. RNA was extracted from each of the resulting cell

populations.

c. Two isolated OSNs differing in eGFP expression. Magenta: mCherry, green: eGFP. Scale bar:

10 μm.

d. Distribution of mCherry and eGFP fluorescence intensity for FACS-sorted cells. Three cell

populations were isolated for RNAseq: Cells with low OMP promoter-driven mCherry

expression and high TRPM5 promoter-driven eGFP expression (MVC_eGFP cells), cells with

high OMP promoter-driven mCherry and low eGFP expression (OSN_eGFP- cells) and cells

with eGFP expression of the same magnitude as MVC_eGFP cells and high OMP promoter-

driven mCherry expression (OSN_eGFP+ cells). The number of cells collected for this FACS

run were: OSN_eGFP-s 1,500,000, OSN_eGFP+s 5336 and MVC_eGFP cells 37,178.

e. qPCR levels (normalized to levels 18s RNA) for expression of transcripts encoding for OMP

(**i**), TRPM5 (**ii**), eGFP (**iii**) and ChAT (**iv**). The asterisks denote significant differences tested

with either t-test or ranksum with p-values below the significance p-value corrected for multiple

comparisons using the false discovery rate (pFDR)(Curran-Everett, 2000). pFDR is 0.033 for

OMP, 0.05 for TRPM5, 0.05 for eGFP and 0.03 for ChAT, n=8 for OMP OSN_eGFP-s, 4 for

OMP OSN_eGFP+s and 4 for MVC_eGFP cells.

1122

1123

1124

Figure 2. Decreased yield of OSN_eGFP+ cells when mice are moved from low ventilation

(LV) to high ventilation (HV) cages.

a and b. Distribution of mCherry and eGFP fluorescence intensity for FACS-sorted cells that

either were not transferred to HV cages (**a**) or were transferred to HV cages for 22 days before

sorting (**b**).

c. Time course showing change in the number of sorted OSN_eGFP+s after mice were

transferred to HV cages.

d and e. Dependence of the yield of OSN_eGFP+ cells after sorting on the number of days in

LV cages (**d**) or the number of days in HV cages (**e**).

Figure 3. Coverage of TRPM5 transcript by RNAseq in the different cell groups sorted

from the olfactory epithelium

a. In situ for TRPM5 and OMP transcripts in the olfactory epithelium shows strong label for

TRPM5 in MVCs (asterisks) and sparse labeling in the OSN nuclear layer (arrows). **i.** OMP, **ii.**

TRPM5, **iii.** Composite. The scale bar is 100 μm.

b and c. Coverage of the reads for the TRPM5 transcript for the different cell groups. Read

coverage was computed with 'Samtools depth' over the region corresponding to the Trpm5 gene.

b. Coverage per exon. All samples normalized to TRPM5 reads show distribution among exons

with little difference between MVC and OSN_eGFP+ cells.

c. Coverage shown per base pair as a function of location in the genomic DNA. Plots show read

depth per base pair for each cell type normalized to total number of mapped reads. Gray shading

indicates 95% confidence interval.

1148

1149

Figure 4. RNAseq comparison of OSN_eGFP- and OSN_eGFP+ cells.

a. Heatmap showing the top 10 upregulated and top 10 downregulated genes identified by

DESeq2.

b. Heatmap showing all *Olfr* genes detected in the data.

For both a and b, row and column order were determined automatically by the *pHeatmap*

package in R. For each data point relative expression was calculated by subtracting the average

row value from each individual value.

c. Volcano plot of all Olfactory receptors, demonstrating the small number of enriched olfactory

receptors in the OSN_eGFP+ population.

d. Hierarchical clustering of transcripts for taste transduction and transcripts expressed in

canonical and non-canonical OSNs identified by RNAseq as significantly different in expression

between the cell groups. We compared expression of transcripts involved in taste transduction,

canonical olfactory transduction, and non-canonical OSNs. The non-canonical OSNs considered

here included guanilyl-cyclase D (GC-D) OSNs (Juilfs et al., 1997), Trpc2 OSNs (Omura and

Mombaerts, 2014), Cav2.1 OSNs (Pyrski et al., 2018), and OSNs expressing trace amine-

associated receptors (Taars) (Liberles, 2015). Transcripts identified by DESeq2.

e. Gene ontology (GO) term enrichment was calculated from differentially expressed genes using

TopGO in R. An enrichment value for genes with Fischer p value <0.05 was calculated by

dividing the number of expressed genes within the GO term by the number expected genes (by

random sampling, determined by *TopGO*).

1170

1171

1172

1173

Figure 5. RNAseq comparison of OSN_eGFP- and OSN_eGFP+ vs. MVC_eGFP cells.

a and b. Heatmaps showing hierarchical clustering of the top 10 upregulated and top 10

downregulated genes identified by DESeq2. **a.** OSN_eGFP- vs. MVC_eGFP cells. **b.**

OSN_eGFP+ vs. MVC_eGFP cells. For both a and b, row and column order were determined

automatically by the *pHeatmap* package in R. For each data point relative expression was

calculated by subtracting the average row value from each individual value.

1180   **c and d.** Gene ontology (GO) term enrichment was calculated from differentially expressed

1181   genes using *TopGO* in R. **c.** OSN_eGFP- vs. MVC_eGFP cells. **d**. OSN_eGFP+ vs.

1182   MVC_eGFP cells. An enrichment value for genes with Fischer p value <0.05 was calculated by

1183   dividing the number of expressed genes within the GO term by the number expected genes (by

1184   random sampling, determined by *TopGO*).

1185

1186

1187

Figure 6. Significant differences in virally-related gene ontology. Heatmaps show hierarchical

clustering of significantly differentially expressed genes identified by DESeq2. **a.** MVC_eGFP

cells compared to OSN_eGFP-. **b.** OSN_eGFP+ compared to OSN_eGFP-.

1191

1192

Figure 7. Intranasal inoculation with HSV-1 elicits a decrease in OSNs.

a and b. Distribution of mCherry and eGFP fluorescence intensity for FAC-sorted cells that

either were not infected (**a**) or were inoculated intranasally with 10 μL of solution per naris (20

μL total with HSV-1 (1x 10^6 PFU/naris; McKrae strain) 5 DPI (**b**).

c. Fraction of total cells for OSN_eGFP- (**i**), OSN_eGFP+ (**ii**) or MVC (**iii**) cell groups. GLM

analysis yielded significant effects for the fraction of OSN_eGFP- compared to MVCs (p<0.001)

and for the interaction between the fraction of OSN_eGFP- compared to MVCs and HSV-1

treatment (p<0.001, 75 observations, 69 d.f., GLM F-statistic=111 and p-value < 0.001, n=14

1201   mice for control, n=11 mice for HSV-1 inoculation). Asterisks denote a significant difference

1202   post-hoc with t-test p-value < pFDR = 0.047.

1203

1204 **Supplemental Information**

1205

1206

1207 **Transcriptional profiling reveals TRPM5-expressing cells involved in viral infection in the**

1208 **olfactory epithelium**

1209

1210 B. Dnate' Baxter[1,2,‡], Eric D. Larson[3 ‡], Paul Feinstein[4], Arianna Gentile Polese[1,2], Andrew N.

1211 Bubak[5], Christy S. Niemeyer[5], Laetitia Merle[1,2], Doug Shepherd[6], Vijay R. Ramakrishnan[3],

1212 Maria A. Nagel[5], and Diego Restrepo[1,2,*]

1213

1214

1215   **Figure 4 – figure supplement 1. Excel worksheet with the results of comparison of gene**

1216   **transcription between OSN_EGFP+ and OSN_EGFP-.**

1217

1218   **Figure 4 – figure supplement 2. Metadata for the RNAseq.**

1219

1220   **Figure 5 - figure supplement 1. Excel worksheet with the results of comparison of gene**

1221   **transcription between MVC_eGFP cells and OSN_EGFP-.**

1222

1223   **Figure 5 - figure supplement 2. Excel worksheet with the results of comparison of gene**

1224   **transcription between MVC_eGFP cells and OSN_EGFP+.**

1225

1226

1227

1228

1229 **Figure 5 – figure supplement 3. Hierarchical clustering of transcripts identified by RNAseq**

1230 **as significantly differentially expressed between male and female. a.** OSN_eGFP-. **b.**

1231 MVC_eGFP cells **c.** OSN_eGFP+. Transcripts identified by DESeq2.

1232

1233

Figure 5 – figure supplement 4. Hierarchical clustering of olfactory receptor transcripts

identified by RNAseq as significantly differentially expressed between male and female. a.

OSN_eGFP-. **b.** OSN_eGFP+. Transcripts identified by DESeq2.

1237

1238 **Figure 6 – figure supplement 1. Excel worksheet with the results of gene ontology**

1239 **enrichment analysis for OSN_eGFP+ compared to OSN_eGFP-.**

1240

1241 **Figure 6 - figure supplement 2. Excel worksheet with the results of gene ontology**

1242 **enrichment analysis for MVC_eGFP cells compared to OSN_EGFP-.**

1243

1244 **Figure 6- figure supplement 3. Excel worksheet with the results of gene ontology**

1245 **enrichment analysis for MVC_eGFP cells compared to OSN_EGFP+.**

1246

1247

1248 **Figure 6- figure supplement 4. Significant differences in inflammation gene ontology for**

1249 **OSN_eGFP+ compared to OSN_eGFP-.**

1250

1251

1252   **Figure 6- figure supplement 5. Significant differences in inflammation gene ontology for**

1253   **MVC_eGFP+ compared to OSN_eGFP-.**

1254

1255

Figure 6- figure supplement 6. Significant differences in immune gene ontology for

OSN_eGFP+ compared to OSN_eGFP-.

1256

1257

1258

1259

1260

1261

Figure 6- figure supplement 7. Significant differences in immune gene ontology for

MVC_eGFP+ compared to OSN_eGFP-.

1264

1265