Recurrent evolution of two competing haplotypes in an insect DNA virus

Tom Hill[1]*, Robert L. Unckless[1]

1. 4055 Haworth Hall, The Department of Molecular Biosciences, University of Kansas, 1200 Sunnyside Avenue, Lawrence, KS 66045.

* Corresponding author - Email: tom.hill@ku.edu

Keywords: nudivirus, *Drosophila innubila,* DiNV, virulence, coevolution

Abstract

Hosts and viruses are constantly evolving in response to each other: as hosts attempt to suppress the virus,

the virus attempts to evade and suppress the host's immune system. This arms race results in the evolution

of novel pathways in both the host and virus to gain the upper hand. Here we describe the coevolution

between *Drosophila* species and a common and virulent DNA virus. We identify two distinct viral types

that differ 100-fold in viral titer in infected individuals, with similar effects across multiple species. Our

analysis suggests that one of the viral types appears to have recurrently evolved at least 4 times in the past

~30,000 years, including in another geographically distinct species, due to the high effective mutation rate

which increases with titer. The higher titer viral type is associated with suppression of the host immune

system and an increased transmission rate compared to the low viral titer type. Both types are maintained

in all populations, likely due to an increased virulence in the high titer type creating a trade-off between

effective transmission and virulence and resulting in nearly equal reproduction rates (R_0) in both types.

Together these results suggest that the reciprocal selective pressures caused by the co-evolution between

host and virus has resulted in this recurrently evolving relationship.

Introduction

Antagonistic coevolution between hosts and their parasites is nearly ubiquitous across the tree of life. As a

result, genes involved in immune defense are among the fastest evolving genes in host genomes (NIELSEN

et al. 2005; SACKTON *et al.* 2007; ENARD *et al.* 2016; SHULTZ AND SACKTON 2019). Viruses are a particular

fitness burden on hosts; for viruses to persist within populations, they must successfully invade the host

organism, contend with the host immune system, replicate and then transmit the newly produced particles

to a new host (HOLMES 2007; GIFFORD 2012). Once successfully established in a population, natural

selection acts to modulate the rate the virus propagates relative to its virulence, optimizing the ratio of viral

virulence to transmission (WILLIAMS AND NESSE 1991; MAY AND NOWAK 1995; LIPSITCH *et al.* 1996).

Thanks to their elevated mutation rate and high population sizes, viruses can easily do all of this, and even

co-opt or manipulate host-pathways in the process (BURGYAN AND HAVELDA 2011; DAVEY *et al.* 2011;

PALMER *et al.* 2019). As a result, proteins that interact with viruses often show the fastest rates of evolution,

and the highest rates of adaptation compared to the rest of the genome, as the host attempts to suppress the

pathogen, or escape infection (OBBARD *et al.* 2006; OBBARD *et al.* 2009a; MUKHERJEE *et al.* 2013; ENARD

et al. 2016; PALMER *et al.* 2018a).

Hosts have evolved numerous immune pathways to reduce viral burden and enhance survival after

infection (MERKLING AND VAN RIJ 2013; WEST AND SILVERMAN 2018). Interestingly, several of these

pathways (IMD, Toll, JAK-STAT) are generally associated with other pathogens, yet also respond to

infection by viruses, though the specific mechanisms are not yet known (HOFFMANN 2003; HULTMARK

2003; COCCIA *et al.* 2004; ZAMBON *et al.* 2005b; COSTA *et al.* 2009; FERREIRA *et al.* 2014a; WEST AND

SILVERMAN 2018). In *Drosophila melanogaster*, the RNA interference (RNAi) pathway is involved in

resisting viral infection by generating suppressive RNAs complementary to the viral sequence (HULTMARK

2003). As might be expected, this antiviral RNAi pathway is also rapidly evolving in many species

(OBBARD *et al.* 2006; WANG *et al.* 2006; OBBARD *et al.* 2009b; MERKLING AND VAN RIJ 2013).

Drosophila innubila Nudivirus (DiNV), which infects the host *D. innubila*, is the one of the few

DNA viruses to naturally infect *Drosophila,* and the only documented case of a DNA virus infection at high

frequency in natural *Drosophila* populations (UNCKLESS 2011; HILL *et al.* 2019). *Drosophila innubila* is a

mushroom feeding species of the *Drosophila* subgenus found inhabiting woodlands found on mountains

across Arizona & New Mexico, separated by large expanses of desert. *D. innubila* radiated north from

Mexico during the last glaciation period and came to inhabit these forests after the glacial retreat, creating

a subdivided population with high rates of gene flow between locations (DYER AND JAENIKE 2005; HILL

AND UNCKLESS 2020).

During this period *D. innubila* likely became infected with DiNV, suggesting a long-lasting host-

pathogen relationship in multiple populations. This could lead to opportunities to study the coevolution of

DiNV and *D. innubila* in replicate (HILL AND UNCKLESS 2020), which could potentially result in parallel

or divergent evolution of the virus and interacting host pathways (ANDERSON AND MAY 1982; KALTZ AND

SHYKOFF 1998).

A pair of previous studies examined the rates of evolution of DiNV and *D. innubila,* finding the

envelope and replication machinery to be rapidly evolving in DiNV, suggesting its importance in viral

propagation (HILL AND UNCKLESS 2018). In *D. innubila,* the antiviral RNAi machinery is not rapidly

evolving, possibly as DNA viruses interact with different immune pathways to RNA viruses (the primary

burden of *D. melanogaster*) (WEBSTER *et al.* 2015). Consistent with this, the Toll pathway is both rapidly

evolving in *D. innubila* and suppressed by a related nudivirus upon infection in *D. melanogaster* (HILL *et*

al. 2019; PALMER *et al.* 2019).

DNA viruses, and nudiviruses such as DiNV in particular, have much larger genomes (100kbp or

larger) than RNA viruses, with much more complicated replication cycles (ROHRMANN 2013). RNA viruses

also have much higher mutation rates than DNA viruses, yet much lower levels of diversity due to lower

recombination rates and efficient selection on variation in the genome (PENNINGS 2012; PENNINGS *et al.*

2014; WILSON *et al.* 2016; FEDER *et al.* 2019). As a result, we expect the evolutionary dynamics of DNA

viruses to differ dramatically from RNA viruses (ROHRMANN 2013; HILL AND UNCKLESS 2017). In fact,

because of their large genomes, high recombination rate and low mutation rate, we expect that DNA virus

coevolution with hosts will be qualitatively different than RNA viruses or bacteria. Together this paints a

picture suggesting that hosts can have different relationships with different pathogens (such as RNA viruses

or DNA viruses), and the pathogens themselves can behave differently within the host. Characterizing the

relationships between different species and their long-term pathogens, such as DiNV and *D. innubila*, will

help broaden and expand our understanding of how hosts and pathogens evolve in response to each other.

Here, we survey the genetic variation in DiNV to infer its co-evolutionary history with *D. innubila*

and two other associated hosts. We identify two viral multilocus genotypes (considered to be haplotypes)

that differ by 11 focal SNPs and show that these viral types are maintained within the same host population

and across multiple isolated host populations. Despite high rates of recombination, these SNPs are tightly

linked likely due to extremely strong selection and possibly incompatibilities between types. One viral type

is associated with 100-fold higher viral titer and increased virulence compared to the other. Further, we find

evidence that the high titer type evolved independently in at least four geographically-isolated host

populations. Together, these results suggest rapid evolutionary dynamics of host-virus interactions, due to

the multiple competing viral types that interact with different host pathways.

Results

DiNV segregates for linked variants strongly associated with viral titer

To characterize the evolutionary dynamics of wild Drosophila innubila Nudivirus (DiNV) in its host (*D.*

innubila), we sequenced wild-caught individuals from four populations with the expectation that some

(~40% in previous samples) individuals would be infected (HILL AND UNCKLESS 2020). We considered

strains to be infected with DiNV if they had at least 10x coverage for 95% of the genome. In total, we used

sequencing information for 57, 92, 92, and 92 individuals from the Huachucas (HU), Santa Ritas (SR),

Chiricahuas (CH), and Prescott (PR) populations with infection rates 26%, 44%, 63% and 79%, respectively

(Supplementary Table 1). We also used 35 individuals collected in the Chiricahuas in 2001 (52% infected

with DiNV) and 80 individuals collected in the Chiricahua's in 2018 (Supplementary Table 2, pre-selected

using PCR, 40 infected with DiNV and 40 uninfected).

We isolated and sequenced DNA from these samples and, after filtering and mapping to the genome

we called variation in the viral genomes to assess the extent of adaptation in each viral population.

Consistent with an arms race between host and virus, most envelope and novel virulence (GrBNV-like)

genes show strong signatures of adaptive evolution in each population compared to background viral genes

(using McDonald-Kreitman based statistic Direction of Selection (STOLETZKI AND EYRE-WALKER 2011)

and Selection Effect (EILERTSON *et al.* 2012), Supplementary Figure 1, DoS > 0, GLM *p*-value < 0.05).

Given these potential signatures of an arms race between *D. innubila* and DINV, we first attempted

to determine if any host or viral genetic variation is associated with within-host viral titer, which we use

here as a measure of virulence. For each virus-infected individual, we quantified viral titer (as viral genome

coverage normalized to host autosomal genome coverage) and identified both host and viral

polymorphisms. We then performed an association study across both host and virus variable sites to identify

variants significantly associated with viral titer using PLINK (PURCELL *et al.* 2007).

103   Of 5,283 viral SNPs in the 155kbp DiNV genome, 1,403 SNPs are segregating in at least 5 infected

host individuals. Of those 1,403 SNPs, 78 are significantly associated with viral titer after multiple testing

correction (FDR < 0.01, Figure 1A). Of these, 16 are within 2000bp of the start site of a gene, 18 are coding

nonsynonymous, 11 are coding synonymous and 33 are intergenic. The most significantly associated SNP

is a non-synonymous polymorphism in the active site of *Helicase-2* (Figure 1A, Supplementary Figure 2).

The *Helicase-2* polymorphism is the only significantly associated polymorphism found segregating at a

range of frequencies *within* individuals (Supplementary Figure 2). The frequency of this derived

polymorphism has a negative relationship between viral titer and the derived SNPs frequency (GLM t-value

= -20.516, *p*-value = 5.55e-62). However, when ranking samples by viral titer, the SNP frequency does not

fit this expectation and several samples fixed for the ancestral allele also have a lower viral titer

(Supplementary Figure 2).

114   We also identified a striking association between viral titer and eleven nearly perfectly linked

polymorphisms found across the DiNV genome (Figure 1A, highlighted SNPs, Table 1, Supplementary

Figures 2-4, Sig. SNPs). We refer to these two types as the 'High Type' and 'Low Type' (Figure 1B).

This multilocus genotype includes three non-synonymous SNPs and five SNPs in the UTRs of known

virulence factor genes and three intergenic SNPs. Viral titer is, on average, 100-fold higher in individuals

infected with High type virus compared to the ancestral Low type (Figure 1B). Though we found few

strains with an intermediate number of SNPs, viral titer increases as the number of High type SNPs

increases (Figure 1C, GLM t-value = 34.971, *p*-value = 5.912e-16), though the rate of increase slows as

the number of High type SNPs increases suggesting diminishing returns (Figure 1C). Some of these

polymorphisms are associated with known virulence factors, or are related to the formation of the viral

envelope co-opting the host vesicle trafficking system and are rapidly evolving in nudiviruses (e.g. *VLF-*

1, ODV-E56, PIF-3) (ROHRMANN 2013; HILL AND UNCKLESS 2017; HILL AND UNCKLESS 2018).

Additionally, several are associated with orphan genes thought to be novel virulence factors, including a

gp83, a gene that downregulates Toll-induced antimicrobial peptides (AMPs) and upregulates those

induced by IMD (*gp83*) (PALMER *et al.* 2019). Both pathways may interact with DNA viruses (ZAMBON

et al. 2005a; COSTA *et al.* 2009; MERKLING AND VAN RIJ 2013; FERREIRA *et al.* 2014b; LAMIABLE *et al.*

2016; PALMER *et al.* 2019).

131   Among populations there is a positive correlation between the frequency of the High type and

132   overall DiNV infection frequency (Figure 1D, GLM logistic regression z-value = 6.104, *p*-value =

133   0.00883), suggesting that the High type may have a higher effective transmission rate, resulting in a

higher number of new individuals infected, per DiNV infected individual. The transmission rate appears to be higher for the High type, despite a possibly higher death rate than the Low type. We also find both viral types in collections from 2001 and 2017, with the High type significantly more common in the 2017 collection (Fisher Exact Test p-value = 0.0167, Figure 1D).

Table 1: Candidate viral SNPs associated with viral titer, their loci, associated genes and the functional category of that gene.

SNP loci	Nearest gene	SNP functional annotation	Nearest gene functional annotation
G14249T	*VLF-1*	Non-Synonymous	Required for late stage virus activity and particle assembly
C41210T	*LEF-4*	Non-Synonymous	RNA polymerase subunit for RNA modification
G42389T	*gp83*	Upstream	Suspected virulence factor which suppresses Toll activity
A59194G	*gp51*	Intergenic	Suspected virulence factor
C59275A	*PIF-6*	Upstream	Per OS Infectivity factor envelope protein, required for oral infection
T59276C	*PIF-6*	Upstream	Per OS Infectivity factor envelope protein, required for oral infection
G66615A	*gp19*	Non-Synonymous	Suspected virulence factor
C78978T	*gp94*	Intergenic	Suspected virulence factor
G78991A	*gp94*	Intergenic	Suspected virulence factor
C126118A	*ODV-E56-2*	Upstream	Occlusion-derived virus envelope protein required for particle formation
A132593C	*Helicase-2*	Non-Synonymous	Unwinds DNA and is critical for DNA replication
T140117C	*PIF-3*	Upstream	Per OS Infectivity factor envelope protein, required for oral infection

Further, the expression of the virally encoded suppressor of Toll signaling, *gp83,* per viral particle is greater in strains containing at least one derived SNP of the high type increase (Figure 1E, GLM t-value = 10.32, p-value = 7.34e-12), suggesting enhanced virulence in the high type. For these SNPs, few are found at intermediate frequencies within samples (Supplementary Figure 2), and no samples contain more than two SNPs at a similar frequency (Supplementary Figure 2). This suggests that hosts are either

infected completely with Low type or High type virus particles and the two types may be partially

incompatible with each other. Together, these results suggest that the high viral type is more virulent

possibly because it is better able to suppress host Toll signaling.

Figure 1: Viral genome-wide association study for DiNV titer in wild *D. innubila*. **A.** Manhattan plot for

each DiNV SNP and the significance of its association with DiNV titer. SNPs are colored if they are

upstream, synonymous, non-synonymous or nonsense mutations. 12 named SNPs are either part of the

significantly associated viral haplotype or are in *Helicase-2*. The FDR corrected *p*-value cutoff of 0.01 is

shown as a dashed line (multiple testing correction for 1,403 tests). **B.** Viral titer for individual wild

caught flies infected with Low and High DiNV haplotypes (containing all high-type SNPs). The middle

bar represents median value, upper and lower bars represent 25th and 75th percentile and whiskers

represent a 95% confidence interval. **C.** Association between the number of significant SNPs and the viral

titer of a sample. **D.** Across five populations, the frequency of the High DiNV haplotype is correlated with

the frequency of the virus infection. **E.** Expression (in FPKM per viral particle) of *gp83* increases with the

number of High DiNV haplotype SNPs.

The High DiNV viral type suppresses the Drosophila immune system and has elevated virulence

Given the striking difference in viral titer and infection frequencies across populations between High and

Low type viruses, we sought to further characterize the differences in infection dynamics between the types.

We sequenced mRNA from 80 wild *D. innubila* males collected in 2018 (Supplementary Table 2, 40

infected with DiNV, 40 uninfected) and performed a differential expression analysis between infected and

uninfected individuals. Few genes were differentially expressed (DE) between infection states in *D.*

innubila (Supplementary Figure 5A, *p*-value < 0.01 after FDR multiple testing correction, above the dotted

line), but these DE genes were enriched for several interesting categories. Specifically, we found IMD-

induced antimicrobial peptides (AMPs) were upregulated upon DiNV infection, while one Toll-induced

AMP, and several chorion genes and heat shock proteins were downregulated (Supplementary Figure 5A).

We also compared these results to a laboratory experiment in *D. melanogaster* of differential expression

after infection with a close relative of DiNV (PALMER *et al.* 2018c). These same genes are also differentially

expressed in *D. melanogaster*. Of the 12 genes which are differentially expressed in the same direction in

both species, five are AMPs and five are chorion genes (Supplementary Figure 5B).

We then compared gene expression between *D. innubila* infected with High or Low type. We find

17 DE host genes and 9 DE viral genes between types (after controlling for virus copy number as

FPKM/titer, Figure 2, FDR corrected *p*-value < 0.01, GLM t-value = -4.6239413, *p*-value = 9.876143e-05).

Specifically, four Toll-mediated immune peptides (*Listericin*, *IM33*, Bomanins *BomBC2* and *BomT2*) have

reduced expression in High Type infected individuals compared to the Low Type (Figure 2, Supplementary

Figure 6). Finally, viral genes of interest (*PIF-3, VLF-1, gp83*) have higher expression per viral particle

(FPKM/titer) in the High type compared to the Low type, all also increase in expression per viral particle

(FPKM/titer) as the number of High type alleles increases (Figure 1E & 2, t-value = 13.732, *p*-value 3.36e-

15). This pattern may be driven by the allele of the non-synonymous SNP in *VLF-1* (GLM t-value = 2.13,

p-value = 0.04272) and the alleles of the SNPs upstream of *gp83*, *PIF-3*, *gp51* and *ODV-E56* (GLM t-value

= 3.518, *p*-value = 0.00162). Together these results suggest that the high viral type has increased expression

of key virulence factors, which in turn, manipulate the expression of host genes involved in immune defense

to result in the observed differences in viral titer. These results suggest that higher *gp83* expression may

cause the lower Toll-mediated AMP expression, possibly due to lowering *Myd88* expression, which in turn

prevents the host from enacting a proper immune response to DiNV infection (Figure 2, Supplementary

Figure 5).

Experimental infections recapitulate differences in viral type virulence

To assess if the virulence of the virus types differs in experimental infections, we performed experimental

infections of *D. innubila* males using viral filtrate of strains infected with one of the two types of DiNV.

As infectious viral titer is increased, survival decreases regardless of viral type (Supplementary Figure 7 &

8, ANOVA residual deviance = 3.536, *p*-value = 2.454e-07, Cox Hazard Ratio z-value > 2.227, *p*-value <

0.02592), with survival decreasing as titer increases (Supplementary Figure 7, Cox Hazard Ratio z-value >

5.428, *p*-value < 5.69e-08). In both cases viral titer also increases for the first 3 days of infection (GLM t-

value = 9.817, *p*-value = 3.6e-14).

Figure 2: Differential expression of *D. innubila* and DiNV genes between *D. innubila* infected with either

the Low type or High type DiNV multilocus genotypes. For host genes, the log-fold change of mRNA

fragments per million fragments is compared, while for viral genes the log-fold change of viral mRNA

fragments per million fragments per viral particle is compared. Genes are colored/labelled by categories of

interest, specifically antimicrobial peptides (AMPs), proteins involved in the extracellular matrix and viral

proteins. Specific genes of interest, such as *Myd88*, are also named. The FDR-correct significance cut-off

of 0.01 (10,320 tests) is shown as a dashed line.

For a set of independent viral isolates, four High type- and four Low type-infected samples, we

diluted samples to roughly equal concentrations of viral particles (about 0.5 particles per host genome prior

to filtration) and performed infections for replicates of 10 males with microneedles dipped in one of the
filtrate samples. Survival is significantly lower for flies infected with High type viruses when compared to
either flies pricked with sterile media (Figure 3A, Cox Hazard Ratio z-value = 3.671 p-value = 0.000242)
or those pricked with Low type virus (Figure 3A, Cox Hazard Ratio z-value = 4.611 p-value = 4e-06). Flies
pricked with Low type virus do not show a significant reduction in survival compared to control flies (Cox
Hazard Ratio z-value = 1.353, p-value = 0.176). We also measured viral titer over time using qPCR, and
find titer increases through time in flies infected with either type (Figure 3B, GLM Log_{10}(titer) ~ days +
type, days t-value = 9.912, p-value = 1.76e-14). Flies infected with High type virus have significantly higher
viral titer compared to flies infected with Low type virus (Figure 3B, GLM Log_{10}(titer) ~ days + type, type
t-value = 3.934, p-value = 0.000211). These results suggest that while the High type strain has higher viral
titer and potentially higher transmission rate in wild flies, it also has higher virulence, even after controlling
for initial infection titer.

Figure 3: Effect of viral type in experimental infections. **A.** Survival curves of *D. innubila* infected with
high and low viral types compared to control flies pricked with sterile media, for 15 days post infection.
Survival 5 days post-infection separated by strain is shown in Supplementary Figure 8. **B.** qPCR copy
number of viral *p47* relative to *tpi* in *D. innubila* infected with DiNV filtrate of high and low types.

DiNV types are under strong selection in the host
Recombination is required during nudivirus replication and recombination start sites can be at any point
in the single chromosome circular genome (KELLY 1982; ROHRMANN 2013). These factors likely cause
the incredibly high recombination rates observed in nudiviruses (BLISSARD AND ROHRMANN 1990;
WANG AND JEHLE 2009; ROHRMANN 2013). In DiNV, the eleven key SNPs that distinguish the High and
Low haplotypes are spread across the genome yet are nearly perfectly linked. In contrast, other SNPs in

the genome have relatively low linkage disequilibrium, suggesting that selection to maintain each

haplotype is strong (Supplementary Figure 4 and 9).

Using McDonald-Kreitman based statistics for detecting selection (MCDONALD AND KREITMAN

1991; STOLETZKI AND EYRE-WALKER 2011; EILERTSON *et al.* 2012), we tested whether genes that are

associated with the High and Low haplotypes exhibited different signatures of natural selection compared

to other viral genes. Envelope and virulence proteins show significantly elevated signatures of adaptation

(Figure 4, envelope & virulence versus background paired T-test t-value = 2.1761, *p*-value = 0.03814).

Genes found in the initial GWAS for virulence, which defined the High and Low types (such as VLF-1,

PIF-3, LEF-4 and LEF-5) have significantly higher rate of substitutions being fixed due to selection than

background genes (Figure 4, type-associated genes versus all other, t-value = 2.718, *p*-value = 0.00068).

We also performed a GWAS using the host polymorphism and find few associated SNPs, after

controlling for the viral haplotype (Supplementary Figure 10). Consistent with the arms race model, host

genes we suspect are interacting with DiNV (such as the GWAS hits, AMPs, chorion genes, piRNA genes

and extracellular genes) show elevated levels of substitutions fixed by selection compared to background

genes in *D. innubila* (Figure 4 & Supplementary Figure 1). Finally, DE chorion genes, extracellular genes

and AMPs have significantly more adaptive substitutions than similar non-DE genes (Figure 4, blue dots,

differentially expressed versus all other T-test: *D. innubila* t-value = 4.755, *p*-value = 0.000671). Overall

these results suggest strong selection is acting on both the host to suppress viral activity and the virus to

escape this suppression.

The High viral type of DiNV evolved repeatedly in three D. innubila populations

We next sought to understand the evolutionary origin of the two types. Given that both types are found in

all populations surveyed (Figure 1D), we hypothesized that this could occur one of three ways: First, the

derived haplotype was present ancestrally and has been maintained since before geographic isolation

occurred. Second, the derived haplotype evolved following geographic isolation and has spread via

migration between locations. Third, the derived haplotype has recurrently evolved in each location.

To distinguish between these possibilities and determine the timeframe of divergence, we used the

site frequency spectrum of silent DiNV polymorphism to estimate effective population size backwards in

time for all populations (LIU AND FU 2015). We find three populations (CH, HU and SR) expand from a

single viral particle (N_e= 1) to millions of particles during the last glacial maximum (30-100 thousand years

ago) when *D. innubila* settled its current range (Supplementary Figure 11). This supports a single invasion

event during a host-range change. PR appears to expand between 1 and 10 thousand years ago, suggesting

a much more recent bottleneck during the range expansion in PR (Supplementary Figure 11) (HILL AND

UNCKLESS 2020).

We aligned genomic regions containing SNPs to two related nudiviruses, Kallithea virus and

Oryctes rhinoceros Nudivirus (OrNV) (WANG *et al.* 2008; HILL AND UNCKLESS 2018; PALMER *et al.*

2018b). The High haplotype alleles are not present in either Kallithea or OrNV, and are not found in short

read information for wild *D. melanogaster* infected with Kallithea virus (WEBSTER *et al.* 2015), suggesting

they are derived in DiNV.

Figure 4: Genes implicated in host/virus interaction are rapidly evolving by positive selection in the

Chiricahua population. Difference in selection effect for viral and host gene categories of interest, and

nearby background genes, as indicated by the proportion of substitutions fixed by adaptation, weighted by

mutations in SnIPRE (EILERTSON *et al.* 2012). Genes that have associated SNPs from the GWAS are

highlighted in red, while genes which are differentially expressed upon infection, or between viral types

are labelled in blue. All GWAS hits are also differentially expressed and labelled in red. Genes of interest

are named.

We generated consensus DiNV sequences for each infected *D. innubila* individual and created a

whole genome phylogeny to infer geographic diffusion of samples using BEAST (BOUCKAERT *et al.* 2014).

We then performed ancestral reconstruction of the presence of the High type across the phylogeny using

APE (PARADIS *et al.* 2004). Our samples group as three populations (with HU and SR forming one

population) and, consistent with our expectation, the Low type is the ancestral state (Figure 5A).

Surprisingly, the High type appears to have evolved repeatedly and convergently within each population,

forming separate groups within each population (Figure 5A). The High type also clusters within each

population in a principal component analysis of all viral SNPs (Supplementary Figure 9), and when

repeating these analyses while excluding the eleven focal SNPs.

We next surveyed each background SNP (not associated with the High or Low type) to determine

if the general background supports one of the three outlined ways in which the High type evolved and

spread in each location. We grouped SNPs by their presence in just the High type or Low type (supporting

a single origin and spread by migration) or if they were unique to a single population but shared between

the High and Low types (supporting recurrent evolution with recombination).

In total, 391 SNPs (28% of SNPs surveyed) are unique to a single population yet are still shared

between both High and Low types (Figure 5 and Supplementary Figure 9), compared to 23 SNPs shared

between locations but exclusive to High type samples.

These unique SNPs (161 for CH, 127 for HU and SR, and 53 for PR), are present in all High type

samples of a population but a variable proportion of Low types for that population (between 19% and 94%)

and are unique to that population. This pattern fits with the High type recurrently evolving on a single

background (a different background in each location), supporting recurrent evolution of the High type. The

population-specific background SNPs are spread throughout the DiNV genome, with little evidence of

recombination with the high type SNPs, making it unlikely that these SNPs recombined onto different

backgrounds (Supplementary Figure 4).

Though there is strong linkage between the High type SNPs, they are not perfectly associated with

each other (Supplementary Figure 4). Using this slight disassociation and APE (PARADIS *et al.* 2004), we

performed ancestral reconstruction of SNP origins in each population (assuming recurrent evolution) and

find that, excluding three variable SNPs, the evolution of these SNPs was the same order in each population

(Figure 5B).

To determine if this recurrent evolution is plausible in our estimated timeframe (~10,000 years),

we simulated viral populations using a modified discrete SIR model using deSolve (SOETAERT *et al.* 2010)

with estimated baculovirus mutation rates, ranges of viral titer taken our samples and estimated population

sizes for each viral population (parameters described in the methods). In this model we used an effective

mutation rate scaled to viral titer, considering the mutation rate per particle, so total mutations per

generation increase with viral titer. The simulations suggest that waiting time for the first mutation that

increases titer is highly variable between replicates but usually occurs within 1000 generations (~200 years

assuming 5 generations per year, in >99.1% of replicates, Figure 5C). The average wait time for each

subsequent mutation decreases monotonically (GLM t-value = -2.389, *p*-value = 0.03686). In each case,

the next mutation appears in the background of the previous high titer mutation (Figure 5C) due to the

elevated effective mutation rate and increased basic reproduction number (R_0). The accumulation of

mutations therefore occurs at a geometric (approximately exponential) rate. Additionally, the standard

deviation of time wait times also decreases with each new mutation (GLM t-value = -2.441, *p*-value =

0.04241), increasing the certainty that the entire multilocus genotype will appear in a population rapidly

once the initial mutations appear. This chain reaction of adaptation facilitates the repeated evolution of the

virulent High type independently in three populations, with all eleven mutations fixing in a population

within 6000 generations (~1200 years) in all replicates (3372 generations on average, ~675 years), a

plausible amount of time given our estimated timeframe.

Both viral types are found in two other Drosophila species and have also evolved in a geographically

distinct population

Since we find two types of DiNV are maintained, and that other species are infected with DiNV, we

hypothesized that another species could be a reservoir for the less effective Low type. We chose to study

D. azteca from the Chiricahuas since it is frequently infected with DiNV (~33% infection), overlaps with

D. innubila, and is genetically divergent (40-60 million years) which could mean a very different genetic

interaction between host and virus (UNCKLESS 2011). We also examined DiNV-infected *D. falleni*

(collected in Georgia) as an outgroup. In all, we sequenced 36 *D. azteca* and 56 *D. falleni*. Both types are

present in all examined species, but the high type is rare in *D. azteca* (Figure 6B). The High type has a

significantly higher titer than the low type in all cases (Figure 6A). Viral titer is not significantly different

across species for either High or Low type (Figure 6A, GLM t-value = -1.351, *p*-value = 0.179). We also

find the *D. azteca* samples cluster with CH *D. innubila* samples and contain the CH background SNPs

(Supplementary Figure 9C), suggesting no differentiation in the virus infecting different species.

Interestingly, *D. falleni* DiNV clusters completely separately from the other samples, likely due to its

geographic separation, but still has a derived cluster of High type virus, suggesting a fourth separate

evolution of the High type in Georgia. Despite the lack of difference between species samples in Arizona,

a lower proportion of the *D. azteca* population is infected with DiNV, and the High Type is less common

than the Low type DiNV (Figure 6B). Thus perhaps, even though the relative differences in titer are

preserved between the two species, the Low Type is favored in *D. azteca* because this reduced virulence

leads to a greater R_0 in *D. azteca*. Thus, the two types of the virus may be maintained in both host species

because though they have become specialized to maximize fitness in one host, messy transmission between

host species could lead to their continued presence in both hosts.

We also repeated the GWAS for viral titer in DiNV infecting *D. azteca* and *D. falleni.* In both cases

we again found the 11 High type SNPs associated with viral titer (GLM t-value > 4.28, *p*-value > 0.0001 in

both cases), but not the *Helicase-2* SNP (despite its presence in *D. falleni* DiNV samples). After controlling

for the High type, we find no other significant DiNV SNPs in *D. azteca* associated with viral titer. For

DiNV infecting *D. falleni*, we find 478 significant SNPs (FDR corrected *p*-value < 0.01), though none of
them with as large an effect as the High type associated SNPs.
355

Figure 5: The evolution and maintenance of two viral types. **A.** Phylogeographic reconstruction of the
spread of DiNV through *D. innubila*, rooted on the Kallithea virus reference sequence, including a
reconstruction of the High type evolution (with strains containing all 11 High type variants shown in red,
strains with an intermediate number of high type variants are shown in pink, and strains with no high type
variants are shown in black). Branches are colored when the SNPs found in the background for each High
haplotype are present in the population, showing that the background differs per population. Black
branches show states where branch tips do not contain all the shared High/Low population specific
background SNPs. **B.** Order of mutations in the viral haplotype appearing in each population. Apart from
three mutations the order is consistent between locations. **C.** The number of generations needed for 'High
titer' mutations to evolve in simulated populations, given that each mutation increases the mutation rate.
The number of generations between each mutation appearing decreases as titer increases.

367
368

Two DiNV types may be maintained due to a trade-off between transmission and virulence

Given the ease that the High type appears to evolve recurrently in populations (Figure 5C), its apparent association with increased infection frequency, and its apparent inability to coinfect with the low type, it is surprising that the High type has not outcompeted the Low type. There are several possible explanations for the maintenance of the two types. First, a soft selective sweep may be occurring on the High type, where recurrent mutation followed by a change in environment changes the fitness of the High type that will eventually result in its fixation (HERMISSON AND PENNINGS 2005). Second, both types may be maintained due to a trade-off (ALIZON AND VAN BAALEN 2008). Such a trade-off might even be associated with frequency-dependent selection and cycling frequencies over time. This trade-off could involve different transmission and virulence strategies or might be related to specific adaptation to different hosts.

In a simple model of viral infection dynamics, the success of the virus is measured by its basic reproduction number (R_0) which is the ratio of the instantaneous transmission rate (β) to the virulence of the virus (γ). If there is a trade-off between transmission and virulence, we might expect that although the

High type has a higher instantaneous rate of transmission, it also has higher virulence, killing infected
individuals before they can infect other possible hosts. In contrast, those infected with the Low type persist
with the infection and can therefore transmit proportionally more virus due to more interactions with
susceptible individuals, despite a lower instantaneous transmission rate. To test this, we simulated
populations with two viral types using a modified SIR model in deSolve (SOETAERT *et al.* 2010). We varied
transmission and virulence rates and estimated sets of parameters in which types are maintained within
populations. We find that a stable infection frequency depends on both the magnitude of instantaneous
transmission rate, and the R_0, with higher transmission rates increasing the infection frequency, to a
maximum of $1 - (\gamma / \beta)$ (Figure 7A). The total infected proportion effectively saturates due to the increased
death rate (virulence) of infected individuals, suggesting a trade-off between transmission and virulence as
titer increases (Figure 7A). This is consistent with our results in experimental infections in DiNV
(Supplementary Figures 7 & 8, Cox Hazard Ratio z-value > 2.227, *p*-value < 0.02592), and other theoretical
treatments (MAY AND NOWAK 1995; ALIZON AND VAN BAALEN 2008). Two types are only maintained
when the R_0 is equal for each type ($\gamma_1/\beta_1 = \gamma_2/\beta_2$). As the transmission rate of the High type increases, it
infects a larger proportion of the population and outcompetes the Low type (with High and Low types at
equal proportions when the transmission rates equal), the High type proportion saturates due to the equally
increasing virulence rate which keeps the R_0 equal to the Low type (Figure 7A). Given the requirement for
an equal R_0 for maintenance, the actual proportion of individuals infected with each type depends on the
starting infection frequencies of each type and the difference in absolute transmission rate (Figure 7B).
Based on the infection frequencies of our sampled populations, we suspect that the High type was able to
evolve earlier in the recently bottlenecked PR population (consistent with the High type background being
shared with 94% of PR Low types), or the absolute transmission rate (and virulence rate) may have
increased in PR population, which is likely what has occurred in CH over time (Figure 7B). Together this
suggests that differences in population infection frequencies may depend on a combination of demographic
factors, host genetic factors and the instantaneous transmission rate in each population (with lower
transmission rates in HU and SR compared to PR and CH). This also implies there is a limit to how virulent
a strain can become before it becomes detrimental, as even with higher transmission rates per individual,
DiNV may kill the host before it can transmit, reducing its basic reproduction number (R_0).

Discussion

Viruses are constantly evolving not just to better infect their host, but also to optimize their infection, to
infect as many individuals as possible without preventing the transmission to new hosts (MAY AND NOWAK
1995; LIPSITCH *et al.* 1996; ALIZON AND VAN BAALEN 2008). Since the host is also evolving in response
to the virus, an evolutionary arms-race often ensues (DAWKINS AND KREBS 1979; KALTZ AND SHYKOFF

1998; DAUGHERTY AND MALIK 2012). Here, to work towards expanding our understanding of the co-
evolution of viruses and their hosts, we examine the population dynamics of Drosophila innubila Nudivirus
(DiNV), a DNA virus infecting *D. innubila* (UNCKLESS 2011). DNA viruses have large genomes and often
recombination, placing them as a somewhat transitionary pathogen between RNA viruses, bacteria and
eukaryotic pathogens and parasites. Within our set of viral samples, we find two DiNV haplotypes which
differ by 11 SNPs (Figure 1, named High and Low types). One haplotype (the High type) is associated with
higher viral titer, likely due to an increased manipulation of the host immune system and increased
expression of viral factors. This derived (High type) has likely recurrently evolved in each population since
the last glacial maximum (~10,000 years ago). The two types appear to be incompatible to some degree, as
we find little evidence of co-infections, and mutations appear in a similar order as if navigating an epistatic
fitness landscape (DOBZHANSKY 1937; KONDRASHOV *et al.* 2002; GAVRILETS 2004). Finally, despite the
higher titer and transmission rate of the High type, we find that the two types are maintained in all
populations sampled, possibly because the increased viral titer also increases the virulence, leading to
similar basic reproduction numbers in the High and Low type (ALIZON AND VAN BAALEN 2008).

If the reproduction rate, R_0, was equal between the two viral types we would expect the infection
frequencies to be equal in our sampled populations, which is not the case. This could be caused by the
starting frequencies of each type not being equal (with the Low type starting at a higher frequency, Figure
7). There could also be time or host dependent variation, so over time (and between locations), changes in
the environment could alter transmission rates of each type, also altering the ratio of High type to Low type.
We also do not consider frequency dependent selection in our model, where the transmission rate depends
on an interaction between the infection frequencies of the two types.

The R_0 could be similar between types, but not identical, resulting in slow changes in the ratio of
types over time (as seen between 2001 and 2017, Figures 1D & 7). As we only have two time points, we
could be witnessing a selective sweep of the High type spreading to fixation (NIELSEN 2005), with
recombination causing the observed differences in the background. As we find the High type appears to
have evolved recurrently, it would be unlikely that we have caught these sweeps partway through in all four
populations sampled (Figure 1D). Using the frequency of the High type between 2001 and 2017 CH
samples, we can calculate the selection coefficient for the High type if increasing at an exponential rate
(which is likely given the intermediate frequency at both time points, Figure 7B). If we assume 5
generations per year and an increase among infected individuals from 39% on 2001 to 71% in 2017, the
selection coefficient = 0.007, which suggests the High type would take ~275 years to fix in a population
once it has arisen. Given the coalescence time of the two types is close to the expansion time of the two
viruses (2-30 thousand years), this does not fit with our results, suggesting the two types are being

maintained and not sweeping. Further, if the High type was sweeping, it would be remarkable for us to
catch these sweeps occurring in all four populations sampled, given our estimated time to fixation.
454

Figure 7: Simulated infected populations to determine parameters of stable infection frequencies of two
viral types. **A.** The frequency of stable infection, given the ratio of virulence to transmission, for different
magnitudes of transmission but the same starting frequency (0.001%). **B.** Stable infection frequencies given
the difference in transmission rates (β) between two viral strains, as strain 2 transmission rate (starting
frequency 0.001%) increases relative to strain 1 transmission rate (starting frequency 25%). The difference
in infection frequencies of each strain given differing transmission rates of both strains. Stacked bars show
the observed frequencies of High type infection (Strain 2, Blue), Low type infection (Strain 1, Red) and
Uninfected (Grey) for four sampled populations, positioned on the X-axis to show the estimated ratio of
transmission rates for High and Low type, based on the infection frequencies.

Previous surveys of nudivirus evolution found that a few replication-related genes (including *VLF-*
1 and *ODV-E56*), likely key targets for host suppression, are under recurrent positive selection (HILL AND
UNCKLESS 2017; HILL AND UNCKLESS 2018). Given that these repeatedly rapidly evolving genes are also
associated with the High and Low types and show the highest levels of adaptation (Figure 1, Figure 4), it is
entirely possible that these genes are key factors for infection in nudiviruses. They may also be associated
with virulence in other nudiviruses or baculoviruses. It is interesting that we find the same SNPs recurrently
evolving in each population, opposed to different SNPs affecting the same genes in each population.

One important factor for DNA virus replication is the *Helicase* gene (BLISSARD AND ROHRMANN

1990; ROHRMANN 2013), and we find variation in *Helicase-2* is associated with viral titer in our survey

(Figure 3A & C). As in other nudiviruses, there is extensive adaptive evolution in the *Helicase* gene (Figure

4A) (HILL AND UNCKLESS 2017; HILL AND UNCKLESS 2018) which has previously been posited to be

associated with change in host range (MAEDA *et al.* 1993; CROIZIER *et al.* 1994; ARGAUD *et al.* 1998;

AFONSO *et al.* 2001). It is possible the *Helicase-2* variants allow for optimized infection of different species

(CROIZIER *et al.* 1994). In fact, DiNV infects several *Drosophila* species in the New World at varying

frequencies (including high prevalence in *D. munda*) (UNCKLESS 2011), and so could have an alternate

variant reservoir in any species, with some migration into *D. innubila* (but not *D. azteca* or *D. falleni*),

resulting in the appearance of two competing strains. However, this hypothesis ignores the existence of the

high and low types which appear to explain the difference in viral titer across multiple species much more

convincingly than the *Helicase-2* polymorphism (Supplementary Figure 2).

Nudiviruses have extremely high rates of recombination, as they require at least one crossover

during their replication (ROHRMANN 2013). Given this high rate of recombination, it is interesting we don't

find more intermediate strains with a mix of High and Low SNPs, supporting the idea of an incompatibility

or negative epistatic interactions between SNPs of the two types, similar to a Dobzhansky-Muller

incompatibility (GAVRILETS 2003).

Some models suggest that viruses are under constant selection to maintain an optimum ratio of

virulence to transmission (MAY AND NOWAK 1995; LIPSITCH *et al.* 1996; ALIZON AND VAN BAALEN 2008).

This delicate balance of transmission to virulence could be disrupted with the evolution of a second, high-

titer, viral type. In the case of DiNV, the High type has radically higher virulence and appears to compete

with the first type impacting the persistence of the virus in a host population. Additionally, the High type

appears to recurrently evolve, frequently affecting the persistence of the virus (Figure 5 & 6, Supplementary

Figure 9). This posits a situation where the optimum strategy for the ancestral viral type is a reduced

transmission rate and the fixation of mutations that are incompatible with High type mutations. Further, this

fits with the observed consistent order of fixation for mutations that form the High type (Figure 5B), similar

to navigating the neutral adaptive landscape between two incompatible forms in a Bateson-Dobzhansky-

Muller incompatibilities (DOBZHANSKY 1937; ORR 1995; ORR AND TURELLI 2001; GAVRILETS 2003; ORR

2004). This type of viral type interaction is rarely considered in models used for studying infection and

could lead to a better understanding of viral dynamics and host-virus co-evolution (JACKSON *et al.* 2005;

JACKSON 2009).

DNA viruses such as DiNV have complicated replication cycles and large genomes. This makes

them a sort of evolutionary intermediate between RNA-viruses (small genomes, high mutation rates) and

eukaryotes (large genomes, low mutation rates) and tangential to bacteria and archaea (intermediate

genomes, low recombination rates). However, adaptation appears to occur through changes in a few key

proteins. Here we find the evolution of two competing viral types that differ in these few key genes. These

viral types are maintained in populations likely due to a trade-off between transmission and virulence.

Overall our results suggest that the high mutation rates and extremely high levels of selection can result in

the repeated and convergent evolution of novel host-virus interactions. Additionally, we find that these

host-virus interactions for large DNA viruses can be much more complicated than previous models suggest

(DOLAN *et al.* 2018; FEDER *et al.* 2019).

Materials and Methods

Fly collection, DNA isolation and sequencing

In this study we used previously collected and sequenced *D. innubila* (HILL AND UNCKLESS 2020). Briefly

we collected these flies across the four mountainous locations in Arizona between the 22nd of August and

the 11th of September 2017. Specifically, we collected at the Southwest research station in the Chiricahua

mountains (~5,400 feet elevation, 31.871 latitude -109.237 longitude), Prescott National Forest (~7,900

feet elevation, 34.540 latitude -112.469 longitude), Madera Canyon in the Santa Rita mountains (~4,900

feet elevation, 31.729 latitude -110.881 longitude) and Miller Peak in the Huachuca mountains (~5,900 feet

elevation, 31.632 latitude -110.340 longitude). Baits consisted of store-bought white button mushrooms

(*Agaricus bisporus*) placed in large piles about 30cm in diameter, at least 5 baits per location. A sweep net

was used to collected flies over the baits in either the early morning or late afternoon between one and three

days after the bait was set. Flies were sorted by sex and species at the University of Arizona and were flash

frozen at -80°C before being shipped on dry ice to the University of Kansas in Lawrence, KS. During these

collections we also obtained *D. azteca* during collections which we also sorted by species and sex and flash

froze. *D. falleni* were collected using a similar method in the Smoky Mountains (~6,600 feet elevation) in

Georgia in 2017 by Kelly Dyer, these flies were then sorted at the University of Georgia in Athens GA and

shipped on dry ice to the University of Kansas in Lawrence, KS.

For collected *D. falleni* and *D. azteca*, we attempted to assess the frequency of DiNV infection

using PCR, looking for amplification of the viral gene *p47*. Using primers from (UNCKLESS 2011), P47F:

5′–TGAAACCAGAATGACATATATAACGC and P47R: 5′–TCGGTTTCTCAATTAACTTGATAGC.

We used the following conditions: 95°C 30 seconds, 55°C 30 seconds, 72°C 60 seconds per cycle for 35

cycles.

We sorted 343 *D. innubila* flies, 60 DiNV positive *D. falleni* and 40 DiNV positive *D. azteca* which

we then homogenized and used to extract DNA using the Qiagen Gentra Puregene Tissue kit (USA Qiagen

Inc., Germantown, MD, USA). We prepared a genomic DNA library of these 343 DNA samples using a

modified version of the Nextera DNA library prep kit (~ 350bp insert size, Illumina Inc., San Diego, CA,

USA) meant to conserve reagents. We sequenced the *D. innubila* libraries on two lanes of an Illumina

HiSeq 4000 run (150bp paired-end) (Data to be deposited in the SRA). We sequenced the *D. falleni* and *D.*

azteca libraries on a separate run of a lane of an Illumina HiSeq 4000 (150bp paired-end).

For 80 male *Drosophila innubila* collected in 2018 (indicated in Supplementary Table 2), we split

the sample homogenate in half, isolated DNA from half as described above and isolating RNA using the

Direct-zol RNA Microprep protocol (R2061, ZymoResearch, Irvine, CA, USA). We then prepared a cDNA

library for each of these 80 RNA samples using a modified version of the Nextera TruSeq library prep kit

meant to conserve reagents and sequenced these samples on a NovaSeq NS6K SP 100SE (100bp single

end). We also sequenced DNA for these samples, with DNA isolated and prepared as above, also sequenced

on a NovaSeq NS6K SP 100SE (100bp single end) (Data to be deposited in the SRA).

Sample filtering, mapping and alignment

Following sequencing, we removed primer and adapter sequences using cutadapt (MARTIN 2011) and

Scythe (BUFFALO 2018) and trimmed all data using Sickle (-t sanger -q 20 -l 50) (JOSHI AND FASS 2011).

We masked the *D. innubila* reference genome (HILL *et al.* 2019), using *D. innubila* TE sequences and

RepeatMasker (SMIT AND HUBLEY 2008; SMIT AND HUBLEY 2013-2015). We then mapped short reads to

the masked genome and the Drosophila innubila Nudivirus genome (DiNV) (HILL AND UNCKLESS 2018)

using BWA MEM (LI AND DURBIN 2009) and sorted using SAMtools (LI *et al.* 2009). Following this we

added read groups, marked and removed sequencing and optical duplicates, and realigned around indels

in each mapped BAM file using GATK and Picard (HTTP://BROADINSTITUTE.GITHUB.IO/PICARD ;

MCKENNA *et al.* 2010; DEPRISTO *et al.* 2011). We considered lines to be infected with DiNV if at least

95% of the viral genome is covered to at least 10-fold coverage. We then filtered for low coverage and

mis-identified species by removing individuals with low coverage of the *D. innubila* genome (less than 5-

fold coverage for 80% of the non-repetitive genome), and individuals we suspected of being misidentified

as *D. innubila* following collection. This left us with 318 *D. innubila* wild flies with at least 5-fold

coverage across at least 80% of the euchromatic genome, of which 254 are infected with DiNV

(Supplementary Table 1). We also checked for read pairs which were split mapped between the DiNV

genome and the *D. innubila* genome using SAMtools.

For *D. falleni* we used a previously generated *D. innubila* genome with *D. falleni* variants

inserted (HILL *et al.* 2019). We masked the genome with Repeatmasker (SMIT AND HUBLEY 2013-2015)

and mapped short reads to the masked genome, the repeat sequences and the DiNV genome using BWA

MEM and SAMtools (LI AND DURBIN 2009; LI *et al.* 2009). Then, as with *D. innubila* we filtered for low

coverage and mis-identified species by removing individuals with low coverage (less than 5-fold

coverage for 80% of the non-repetitive genome) leaving us with 56 *D. falleni* samples infected with

DiNV.

For *D. azteca*, we downloaded the genome from NCBI (Accession: GCA_005876895.1) which

we then called repeats from with RepeatModeler (SMIT AND HUBLEY 2008). We masked the genome with

Repeatmasker (SMIT AND HUBLEY 2013-2015) and mapped short reads to the masked genome, the repeat

sequences and the DiNV genome using BWA MEM and SAMtools (LI AND DURBIN 2009; LI *et al.*

2009). As with *D. innubila* we then filtered for low coverage and mis-identified species by removing

individuals with low coverage of the *D. azteca* genome (less than 5-fold coverage for 80% of the non-

repetitive genome), which left us with 37 *D. azteca* samples infected with DiNV. We then called DiNV

variation using LoFreq (WILM *et al.* 2012).

Calling nucleotide polymorphisms across the population samples

For the 318 sequenced samples with reasonable coverage, for host polymorphism, we used the previously

generated multiple strain VCF file, generated using a standard GATK HaplotypeCaller/BCFTools pipeline.

We used LoFreq (WILM *et al.* 2012) to call polymorphic viral SNPs within each of the 254 DiNV infected

samples, following filtering using BCFtools to remove sites below a quality of 950 and a frequency less

than 5%. We then merged each VCF to create a multiple strain VCF file, containing 5,283 SNPs in the

DiNV genome. The LoFreq VCF (WILM *et al.* 2012) output contains estimates of the frequency of each

SNP in DiNV in each sample, to confirm these frequencies, in SAMtools (LI *et al.* 2009) we generated

mPileups for each sample and for SNPs of interest (related to viral titer), we counted the number of each

nucleotide to confirm the estimated frequencies of these nucleotides at each position in each sample. To

confirm that there are no coinfections of types, we also subsampled samples and randomly merged low and

high type samples and again generated mPileup files, for SNPs of interest we again counted the number of

each nucleotide at each position and confirmed these matched our expected counts in the merged files. We

then compared these artificial coinfections to actual samples to confirm the presence or absence of

coinfections, finding no samples consistent with coinfections. We then used SNPeff to identify the

annotation of each SNP and label synonymous and non-synonymous (CINGOLANI *et al.* 2012). We extracted

the synonymous site frequency spectrum to estimate the effective population size backwards in time using

StairwayPlot (LIU AND FU 2015).

Identifying differentially expressed genes between DiNV infected and uninfected Drosophila innubila

For 100 male *Drosophila innubila* collected in 2018 (indicated in Supplementary Table 2), we

homogenized each fly separately in 100μL of PBS. We then split the sample homogenate in half, isolated

DNA from half as described above and isolating RNA using the Direct-zol RNA Microprep protocol

(R2061). Using the isolated DNA, we tested each sample for DiNV using PCR for *P47* as described

previously, using 40 DiNV infected samples and 40 uninfected samples. We then prepared a cDNA

library for each of these 80 RNA samples using a modified version of the Nextera TruSeq library prep kit

meant to conserve reagents and sequenced these samples on a NovaSeq NS6K SP 100SE (100bp single

end). We also sequenced DNA for these samples, with DNA isolated and prepared as above, also

sequenced on a NovaSeq NS6K SP 100SE (100bp single end). The DNA sequenced here was mapped as

described above, with variation called as described above for other DNA samples.

611   Following trimming and filtering the data as described in the methods, we mapped all mRNA

sequencing data to a database of rRNA (QUAST *et al.* 2013) to remove rRNA contaminants. Then we

mapped the short read data to the masked *D. innubila* genome and DiNV genome using GSNAP (-N 1 -o

sam) (WU AND NACU 2010). We estimated counts of reads uniquely mapped to *D. innubila* or DiNV

genes using HTSEQ (ANDERS *et al.* 2015) for each sample. Using EdgeR (ROBINSON *et al.* 2009) we

calculated the counts per million (CPM) of each gene in each sample and counted the number of samples

with CPM > 1 for each gene. We find that over 70.3% of genes have a CPM > 1 in at least 70 samples.

For the remaining genes, we find these genes are expressed in all samples of a subset of the strains (e.g.

DiNV uninfected, DiNV infected, DiNV high infected, DiNV low infected). This supports the validity of

the annotation of *D. innubila*, given most genes are expressed in some manner, and suggests our RNA

sequencing samples show expression results consistent with the original annotation of the *D. innubila*

genome.

623   We attempted to improve the annotation of the *D. innubila* genome to find genes expressed only

under infection. We extracted reads that mapped to unannotated portions of the genome and combined these

for uninfected samples, samples infected with high type DiNV and samples infected with low type DiNV

as three separate samples. We then generated a *de novo* assembly for each of these three groups using

Trinity and Velvet (SCHULZ *et al.* 2012; HAAS *et al.* 2013). We then remapped these assemblies to the

genome to identify other transcripts and found the consensus of these two for each sample. Using the

Cufflinks pipeline (GHOSH AND CHAN 2016), we mapped reads to the *D. innubila* genome and counted the

number of reads mapping to each of these putative novel transcript regions, identifying 15,676 regions of

at least 100bp, with at least 1 read mapping in at least 1 sample. Of these, 717 putative genic regions have

at least 1 CPM in all 80 samples, or in all samples of one group (DiNV uninfected, DiNV infected, DiNV-

low infected, DiNV-high infected). We next attempted to identify if any of these genes are differentially

expressed between types, specifically between uninfected strains and DiNV infected strains, and between

low-type infected and high-type infected strains. Using a matrix of CPM for each putative transcript region

in each sample, we calculated the extent of differential expression between each type using EdgeR

(ROBINSON *et al.* 2009), after removing regions that are under expressed, normalizing data and estimating

the dispersion of expression. We find that 26 putative genes are differentially expressed between infected

and uninfected types, and 69 putative genes are differentially expressed between high and low types. We

took these regions and identified any homology to *D. virilis* transcripts using blastn (ALTSCHUL *et al.* 1990).

We find annotations for 37 putative genes are either expressed in all samples, or differentially expressed

between samples. Of the 14 putative genes expressed in all samples, nine have the closest blast hit to an

rRNA gene, and five have hits to unknown genes. For 23 differentially expressed putative genes with blast

hits, 3 genes are like antimicrobial peptides (*IM1, IM14, IM3*), these genes are significantly downregulated

upon infection, like other Toll regulated AMPs, and have significantly lower expression in strains infected

with high type DiNV compared to low types. The remaining 20 genes all have similarity to genes associated

with cell cycle regulation, actin regulation and tumor suppression genes.

Identifying genes associated with viral titer in Drosophila innubila

As the logarithm of viral titer was normally distributed (Shapiro-Wilk test W = 0.05413, *p*-value = 0.342),

we used PLINK (PURCELL *et al.* 2007) to associate nucleotide polymorphism to logarithm of viral titer in

infected samples. We fit a linear model in PLINK including population, sex, *Wolbachia* presence, the date

of collection and the relationship matrix for relationship of each sample (inferred using PLINK).

We first fit this model for all 5,283 viral polymorphisms, before performing the association study,

we also pruned viral SNPs for both the total population and each subpopulation leaving 1,403 SNPs. For

the total sample we identified associations between the logarithm of viral titer and the frequency of the viral

polymorphism in each individual sample, resulting in the following model:

$$Log_{10}(viral\ titre) \sim SNP + hs + w + p + dc + (SNP * hs) + (SNP * p) + (SNP * w)$$

$$+ relationship[strain]$$

Where hs = host sex, p = location of collection, w = *Wolbachia* presence, dc = date collected

Following model fitting, we found factors which seemed to show little or no effect on viral titre (*p*-value >

0.1) using an ANOVA in R (TEAM 2013), and removed these, refitting the model. This was done step-wise,

leaving the following model by the end:

$$Log_{10}(viral\ titre) \sim SNP + hs + p + (SNP * hs) + relationship[strain]$$

Following this we also performed a GWAS using PLINK (PURCELL *et al.* 2007) in the host, using

previously called host variation, and considering viral haplotype as an additional covariate.

$$Log_{10}(viral\ titre) \sim SNP + hs + p + (SNP * hs) + vh + relationship[strain]$$

Where hs = host sex, p = location of collection, w = *Wolbachia* presence, dc = date collected, vh = viral

haplotype. We found no convincing significant associations (Supplementary Figure 11).

We repeated this analysis for DiNV variants in *D. azteca* and *D. falleni* separately. We performed

the GWAS twice, first using the original model, then including viral haplotype as an additional covariate.

Estimating viral titre using qPCR

Following the identification of the viral haplotype associated with viral titre we sought to determine the

effect of viral haplotypes in actual infections. For 20 samples with fly homogenate, we determine the viral

titer and haplotype following filtration with a 0.22µM filter.

We performed qPCR for the viral gene *p47* (Forward 5-TCGTGCCGCTAAGCATATAG-3,

Reverse 5-AAAGCTACATCTGTGCGAGG-3) on 1 µL of fly filtrate per sample and compared the

estimated Cq values across 3 replicates to estimated viral copy number to confirm viral concentration

(protocol: 2 minutes at 95°C, 40 cycles of 95°C for 30 seconds and 59°C for 20 seconds, followed by 2

minutes at 72°C). Following this we diluted samples to similar Cq values, relative to the sample with the

highest Cq value. We confirmed this by repeating qPCR with *p47* primers of 1 µL of each sample.

For each filtrate sample we performed infections on 30 *D. innubila* males 4-5 days following

emergence using pricks with sterile needles dipped in viral filtrate. We recorded survival of each fly each

day and removed dead flies. Finally, we took samples 1, 3 and 5-days post infection and measured viral

copies of *p47* relative to *tpi* at each time point.

Phylogeography of DiNV infection and the evolution of the different viral haplotypes

For each DiNV infected *D. innubila* sample, we reconstructed the consensus DiNV genome infecting them

using GATK AlternateReferenceMaker and the VCF generated for each strain (MCKENNA *et al.* 2010;

DEPRISTO *et al.* 2011). We then converted these genomes into Phylip format and used BEAST2 to build

the phylogeny of DiNV genomes using 100 million iterations with a burn in of 5 million (BOUCKAERT *et*

al. 2014). We considered phylogeography by providing the longitude and latitude of each samples

collection. We then generated a final consensus phylogeny using Tracer with at least 90% majority

consensus and removed trees that had not converged to the same joint density (BOUCKAERT *et al.* 2014).

To reconstruct the evolution of the high type viral type we used APE (PARADIS *et al.* 2004) to infer the

appearance order of the six perfectly linked SNPs on the phylogeny using the all different rates (ARD)

discrete model (PARADIS *et al.* 2004). We also confirmed these recurrent mutations across the phylogeny

using TreeTime to identify recurrently evolving SNPs (SAGULENKO *et al.* 2018). Finally, we also created

a matrix of SNPs present in at least 2 viral samples and used this matrix in a principle component analysis

in R (TEAM 2013), labelling each sample by their viral type in the PCA.

Simulating the evolution of the high and low viral haplotypes.

We sought to simulate the infection of DiNV in *D. innubila* when considering the evolution of a high titer

viral haplotype, specifically if two viral types can be maintained against each other at stable frequencies,

and if the high viral haplotype with 5 shared mutations could evolve recurrently in the given time period

given realistic parameters. We used the R package DeSolve (SOETAERT *et al.* 2010) to simulate infection

dynamics in a modified SIR model. Specifically, we removed a resistant class, under the assumption that

flies won't live long enough to shed the infection. Therefore, the proportion of population infected per

generation is as follows:

$$p_{Infected} = (Susceptible * Infected * \beta) - (Infected * \gamma)$$

Where β represents an infection parameter and γ represents a virulence parameter (e.g. the increased

likelihood an infected individual has of dying before it can spread its infection). This equation can be

rearranged to show the stabilized maximum frequency of infection:

$$p_{Infected} = 1 - \frac{\gamma}{\beta}$$

Which is maintained while γ is greater than 0.01. The average frequency decreases as absolute transmission

rate decreases past this point. We then extended this to include two competing infection types, to represent

the two viral types, with the total proportion of population infected per generation as follows:

$$p_{Susceptible} = 1 - (Susceptible * Infected_1 * \beta_1) - (Infected_1 * \gamma_1)$$

$$+ (Susceptible * Infected_2 * \beta_2) - (Infected_2 * \gamma_2)$$

This equation can be rearranged as before to show the stabilized frequency of infection for two viral types:

$$p_{Infected} = 1 - \frac{\gamma_1 + \gamma_2}{\beta_1 + \beta_2}$$

In this case, the two viral haplotypes can both be maintained within a population when the ratio of infection

to virulence are the same:

$$\frac{\gamma_1}{\beta_1} = \frac{\gamma_2}{\beta_2}$$

We assessed how starting frequency and difference in γ and β affects the evolution of each type and their

stable frequencies but repeating these simulations for 10,000 generations for each set of parameters. We

repeated simulations with transmission rates varying between 0.00001 and 1, and virulence rates set as the

transmission rate, divided by a scaling factor between 0.01 and 10 (to vary virulence at multiple rates higher

and lower than transmission rate), with all pairwise combinations across all parameters.

Next, we attempted to assess if the 'high titer' viral haplotype could possibly evolve recurrently in

each population in the time scale seen in our findings. We again used the modified SIR model, this time

discrete with population sizes set to 1 million individuals, based on *StairwayPlot* estimates (LIU AND FU

2015). For each infected individual in the population, we tracked the viral titer and also recorded the

presence of absence of five mutations, with each mutation increasing the titer of infection, but each further

mutation having successively smaller increases in viral titer ($titer^{\sqrt{no.muts}}$), representing the epistatic

interaction of high titer associated mutations seen in the viral haplotype. We multiplied the infection

parameter, mutation parameter and virulence parameter by viral titer, under the assumption that viral titer

increases both infection and death rate, and the mutation rate is per viral particle. We considered a per site

mutation rate of 10^{-6}, based on estimated baculovirus mutation rate (ROHRMANN 2013; CHATEIGNER *et al.*
2015), with a specific mutation rate for the five haplotype mutations of 6.4e-12 (1e-6/155kbp) * viral titer.
We then simulated populations in replicate 1000 times for 100,000 generations with a starting infection
frequency of 10% for the 'low titer' haplotype, recording the frequency of the virus in a population, the
frequency of the haplotype and the time that each 'high titer' mutation reaches high enough frequency to
escape stochastic behavior and behave deterministically under selection (GILLESPIE 2004).

To estimate the possible selection coefficient for DiNV in the CH population, we assumed an
exponential distribution and 5 viral generations per year (80 generations between 2001 and 2017). We then
solved the following equation:

$$P_{2017} = P_{2001} * (1 + s)^t$$

Where P_{2017} = the frequency of the High type among viral samples in 2017 (71%), P_{2001} = the frequency of
the High type among viral samples in 2001 (39%), s = the selection coefficient and t = the number of
generations (80). We then used this estimated selection coefficient in the same equation to find the number
of generations (t) to go from $1/2N_e s$ (0.0000714, assuming an N_e of 1000000) to fixation (0.99):

$$0.99 = 0.0000714 * (1 + 0.007)^t$$

Experimental infections of Drosophila innubila with DiNV
We chose *D. innubila* samples infected with DiNV and with sequenced genomes, 4 infected with the high
type DiNV and 4 infected with the low type. For these samples we estimated their viral copy number per
host genome as described previously. We used qPCR on *p47* and *tpi* to find the differences in Cq values to
calculate the concentration of each sample relative to the lowest concentration sample and diluted 50µL of
filtrate for each sample to match the concentration of each sample to the samples with the lowest titer
(IPR07). For a separate 50µL of the IPR01 sample, we performed 1 in 10 serial dilutions to give 45µL of
filtrate at full concentration, 1 in 10 concentration, 1 in 100 concentration and 1 in 1000 concentration.
Using these sets of samples (matched titer and serial dilutions) we next performed experimental infections.

We transferred 50 *D. innubila* (of roughly equal sex ratio) to new food and let them lay eggs for 1
week, following this we collected male offspring aged 2-5 days for experimental infections.

Across 4 separate days in the mornings (between 9am and 11am), we infected the collected male
flies with each sample. For flies in batches of 10, we performed pricks with microneedles dipped in the
prepared viral filtrate. For each day we also had 2 control replicates of 10 flies pricked with microneedles
dipped in sterile media. Following infections, we checked on each vial of 10 flies one-hour post infection
and removed dead flies (likely killed by the needle instead of the virus). We also checked each vial each
morning for 15 days, removing dead flies (freezing to determine the viral titer), and flipping flies to new

food every 3-4 days. Checking at 10am each day, we recorded the day that each fly died, what filtrate they

had been infected with, and what replicate/infection day set they belonged to. We next looked for

differences in survival over time compared to sterile wound controls using a Fit proportional hazards

regression model in R (TEAM 2013; KASSAMBARA *et al.* 2017), considering titer, infection sample and

replicate as co-variates (day of death ~ [titer or strain] + infection date).

For a second set of experimental infections (performed as described above, stabbed with diluted

filtrate from different strains), we also removed 3 living flies 1 hour, 1 day and 5 days post infection. Using

qPCR, we found the difference in *p47* log-Cq and *tpi* log-Cq to estimate the viral copy number for each

sample over time.

Acknowledgements

This work was completed with helpful discussion from Justin Blumenstiel, Joanne Chapman, John Kelly,

Stuart MacDonald, Andrew Mongue and Carolyn Wessinger. We would especially like to thank Maria

Orive, Kelly Dyer and Paul Ginsberg for helpful feedback in the writing of the manuscript and framing of

the discussion. Collections were completed with assistance from Todd Schlenke, Paul Ginsberg, Kelly Dyer,

Brandon Cooper, John Jaenike and the Southwest Research Station. We thank Brittny Smith and Jenny

Hackett at the KU CMADP Genome Sequencing Core (NIH Grant P20 GM103638) and K-INBRE

Bioinformatics Core for assistance in genome isolation, library preparation, sequencing and computational

resources. This work was supported by a K-INBRE postdoctoral grant to TH (NIH Grant P20 GM103418).

This work was also funded by NIH Grants R00 GM114714 and R01 AI139154 to RLU. *D. falleni* collection

was funded by NSF grant DEB-1737824.

Supplementary Figures

789 **Supplementary Figure 1:** McDonald-Kreitman based statistics for each gene in population of Drosophila

790 innubila Nudivirus, with viral envelope and GrBNV potential virulence factors shown separately. DoS =

791 direction of selection, Selection Effect = SnIPRE estimated weighted DoS. Boxplots marked with a * are

792 significantly higher than background/other viral genes (GLM *p*-value < 0.05).

 Supplementary Figure 2: Frequency of each significant SNP within each sample, ranked by the viral titer
 in each sample (left = lowest, right = highest), to show the strong linkage of SNPs and little evidence of co-
 infection, also highlights the association between SNP frequency and Helicase-2.

798 **Supplementary Figure 3:** Linkage disequilibrium between SNPs in DiNV. The labelled SNPs

799 (significant SNPs found in the GWAS) are strongly linked. Points are colored by the estimated linkage

800 between SNPs, from red (highly linked, $r^2 = 1$) to white (unlinked, $r^2 = 0$)

Supplementary Figure 4: Linkage (r^2) between different types of SNPs in each population of DiNV, and across all samples. Other = SNPs which are not significantly associated with DiNV titer and do not form the viral haplotype. Sig = SNPs which are significantly associated with DiNV titer and do not form the viral haplotype. BG = SNPs which are in the background which the viral haplotype evolved on in each population.

Supplementary Figure 5: Volcano plot of changes in gene expression between *D. innubila* infected with
DiNV and uninfected controls. Gene categories of interest, such as enriched categories, are highlighted in
color. The FDR-correct significance cut-off of 0.01 (10,320 tests) is shown as a dashed line. **B.** Comparison
of gene expression changes upon infection for *D. innubila* and *D. melanogaster*. Significantly differentially
expressed genes (*p*-value < 0.01, FDR-corrected) are colored, genes differentially expressed in both species
are colored blue, genes differentially expressed in just *D. melanogaster* are colored yellow and genes
differentially expressed in just *D. innubila* are colored red.

Supplementary Figure 6: Expression changes (shown as transcript fragments per 1 million reads per 1kbp of exon) of antimicrobial peptides between strains infected with high type DiNV, low type DiNV or not infected. Black stars above low samples show significant differential expression between DiNV infected strains and uninfected strains (multiple testing corrected *p*-value < 0.05). Red stars above high samples show significant differential expression between low type infected strains and high type infected strains (multiple testing corrected *p*-value < 0.05).

Supplementary Figure 7: Effect of differences in viral type and titer in experimental infections. **A.**
Survival curves of *D. innubila* infected with DiNV filtrate of different dilutions compared to control flies
pricked with sterile media, for 15 days post infection. **B.** qPCR copy number of viral *p47* relative to *tpi* in
samples of *D. innubila* infected with DiNV filtrate of different dilutions, between 1 and 1000 viral particles
per host genome copy.

Supplementary Figure 8 A: Survival of *D. innubila* reference strain 5 days post infection, using filtrate

from different samples versus uninfected control, colored by high type virus or low type virus. **B.** Survival

of *D. innubila* reference strain 5 days post infection using serial dilutions of IPR01 filtrate versus control.

C. Viral titer estimated per viral genotype at 5 days post-infection, colored by high type virus or low type

virus.   **D.** Viral titer of DiNV infecting *D. innubila* reference strain 5 days post infection using serial

dilutions of IPR01 filtrate versus control.

Supplementary Figure 9. A. Frequency of samples with different numbers of SNPs in the viral
haplotype, there ae very few intermediate types. **B.** Frequency of each SNP in samples infected with the
virus, showing there is little evidence of co-infections. **C.** Principle component analysis of DiNV strains
using variation of strains. Strains are colored by the viral type, showing its recurrent evolution. Point
shape denotes species in which DiNV was found (*D. azteca, D. falleni or D. innubila*). Strains cluster by
collection location. **D.** Linkage between SNPs in the viral haplotype (r^2) and other SNPs in the haplotype,
to other SNPs in the viral genome.

Supplementary Figure 10. A: Manhattan plot of significance of SNP on viral titer after factoring in
interaction with the viral haplotype. The significance cut offs are labelled (*p*-value < 0.05 after multiple
testing correction dotted, *p*-value < 0.05 after permutations dashed). **B**: Manhattan plot of SNP x viral
haplotype interaction for viral titer GWAS in *D. innubila*, calculated using *PLINK*. The significance cut
offs are labelled (*p*-value < 0.05 after multiple testing correction dotted, *p*-value < 0.05 after permutations
dashed). **C:** Manhattan plot of SNP x sex interaction for viral titer GWAS in *D. innubila*, calculated using
PLINK.

Supplementary Figure 11: Effective population size backwards for each population of DiNV going

backwards in time, estimated using StairwayPlot. Dotted lines indicate the error windows for N_e at a given

time point. Lines are colored by population.

Supplementary Tables

Supplementary Table 1: Summary of *Drosophila innubila* and *D. azteca* fly samples collected and

sequenced for this study, table includes summary of coverage for X chromosome, , Muller B, other

autosomes, virus and *Wolbachia*. Also contains SRA accessions for each strain.

Supplementary Table 2: Summary of *Drosophila innubila* fly RNA and DNA collected and sequenced

for this study, including if infected with DiNV.

Supplementary Data

Supplementary Data 1: FPKM of each gene in each sample across the whole *D. innubila* genome,

formatted for use in fitting a generalized linear model. Table include the gene name, gene flybase

annotation, *D. innubila* name, if the strains is infected with DiNV and the FPKM.

Supplementary Data 2: FPKM of each gene in each sample across the whole *D. innubila* genome,

formatted for differential gene expression analysis. Table include the gene name, gene flybase annotation,

D. innubila name, if the strains is infected with DiNV and the FPKM.

Supplementary Data 3: Differential gene expression analysis results between viral infected and
uninfected strains for both *D. innubila* and *D. melanogaster*. Genes are labelled as if differentially
expressed in one of the two species, or if differentially expressed in both.
Supplementary Data 4: VCF file for SNPs in DiNV, used in estimation of population genetic statistics
and in GWAS using PLINK.
Supplementary Data 5: VCF file for SNPs in *D. innubila*, used in estimation of population genetic
statistics and in GWAS.
Supplementary Data 6: Population genetic statistics calculated for each gene in *D. innubila* using
VCFtools for each population.
Supplementary Data 7: McDonald-Kreitman statistics calculated for each gene in *D. innubila* using
VCFtools for each population.
Supplementary Data 8: Population genetic statistics calculated for each gene in DiNV using VCFtools
for each population.
Supplementary Data 9: McDonald-Kreitman statistics calculated for each gene in DiNV using VCFtools
for each population.

References

Afonso, C. L., E. R. Tulman, Z. Lu, C. a. Balinsky, B. a. Moser *et al.*, 2001 Genome sequence of a
baculovirus pathogenic for *Culex nigripalpus.* Journal of virology 75**:** 11157-11165.
Alizon, S., and M. van Baalen, 2008 Transmission-virulence trade-offs in vector-borne diseases. Theor
Popul Biol 74**:** 6-15.
Altschul, S. F., W. Gish, W. Miller, E. W. Myers and D. J. Lipman, 1990 Basic local alignment search tool.
Journal of Molecular Biology 215**:** 403-410.
Anders, S., P. T. Pyl and W. Huber, 2015 HTSeq-A Python framework to work with high-throughput
sequencing data. Bioinformatics 31**:** 166-169.
Anderson, R. M., and R. M. May, 1982 Coevolution of hosts and parasites. Parasitology 85**:** 411-426.
Argaud, O., L. Croizier, M. López-Ferber and G. Croizier, 1998 Two key mutations in the host-range
specificity domain of the p143 gene of Autographa californica nucleopolyhedrovirus are required
to kill *Bombyx mori* larvae. Journal of General Virology 79**:** 931-935.
Blissard, G. W., and G. F. Rohrmann, 1990 Baculovirus diversity and molecular biology. Annual Review
of Entomology 35**:** 127-155.
Bouckaert, R., J. Heled, D. Kühnert, T. Vaughan, C. H. Wu *et al.*, 2014 BEAST 2: A Software Platform
for Bayesian Evolutionary Analysis. PLoS Computational Biology 10**:** 1-6.
Buffalo, V., 2018 *Scythe*.
Burgyan, J., and Z. Havelda, 2011 Viral suppressors of RNA silencing. Trends Plant Sci 16**:** 265-272.
Chateigner, A., A. Bezier, C. Labrousse, D. Jiolle, V. Barbe *et al.*, 2015 Ultra deep sequencing of a
baculovirus population reveals widespread genomic variations. Viruses 7**:** 3625-3646.
Cingolani, P., A. Platts, L. L. Wang, M. Coon, T. Nguyen *et al.*, 2012 A program for annotating and
predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of
Drosophila melanogaster strain w1118; iso-2; iso-3. Fly 6**:** 80-92.
Coccia, E. M., M. Severa, E. Giacomini, D. Monneron, M. E. Remoli *et al.*, 2004 Viral infection and Toll-
like receptor agonists induce a differential expression of type I and lambda interferons in human
plasmacytoid and monocyte-derived dendritic cells. European journal of immunology 34**:** 796-805.

Costa, A., E. Jan, P. Sarnow and D. Schneider, 2009 The Imd pathway is involved in antiviral immune
responses in Drosophila. PLoS ONE 4.
Croizier, G., L. Croizier, O. Argaud and D. Poudevigne, 1994 Extension of Autographa californica nuclear
polyhedrosis virus host range by interspecific replacement of a short DNA sequence in the p143
helicase gene. Proceedings of the National Academy of Sciences of the United States of America
91: 48-52.
Daugherty, M. D., and H. S. Malik, 2012 Rules of engagement: molecular insights from host-virus arms
races. Annual review of genetics 46: 677-700.
Davey, N. E., G. Trave and T. J. Gibson, 2011 How viruses hijack cell regulation. Trends Biochem Sci 36:
159-169.
Dawkins, R., and J. R. Krebs, 1979 Arms Races between and within Species. Proceedings of the Royal
Society of London B: Biological Sciences 205: 489-511.
DePristo, M. A., E. Banks, R. Poplin, K. V. Garimella, J. R. Maguire et al., 2011 A framework for variation
discovery and genotyping using next-generation DNA sequencing data. Nature genetics 43: 491-
498.
Dobzhansky, T., 1937 Genetics and the origin of species. 364.
Dolan, P. T., Z. J. Whitfield and R. Andino, 2018 Mechanisms and Concepts in RNA Virus Population
Dynamics and Evolution. Annual Review of Virology 5: 69-92.
Dyer, K. a., and J. Jaenike, 2005 Evolutionary dynamics of a spatially structured host-parasite association:
Drosophila innubila and male-killing *Wolbachia*. Evolution; international journal of organic
evolution 59: 1518-1528.
Eilertson, K. E., J. G. Booth and C. D. Bustamante, 2012 SnIPRE: Selection Inference Using a Poisson
Random Effects Model. PLoS Computational Biology 8.
Enard, D., L. Cai, C. Gwennap and D. A. Petrov, 2016 Viruses are a dominant driver of protein adaptation
in mammals. eLife 5: 1-25.
Feder, A. F., P. S. Pennings, J. Hermisson and D. A. Petrov, 2019 Evolutionary Dynamics in Structured
Populations Under Strong Population Genetic Forces. G3 (Bethesda) 9: 3395-3407.
Ferreira, Á. G., H. Naylor, S. S. Esteves, I. S. Pais, N. E. Martins et al., 2014a The Toll-Dorsal Pathway Is
Required for Resistance to Viral Oral Infection in Drosophila. PLoS Pathogens 10.
Ferreira, Á. G., H. Naylor, S. S. Esteves, I. S. Pais, N. E. Martins et al., 2014b The Toll-Dorsal Pathway Is
Required for Resistance to Viral Oral Infection in *Drosophila.* PLoS Pathogens 10.
Gavrilets, S., 2003 Perspective: models of speciation: what have we learned in 40 years? Evolution;
international journal of organic evolution 57: 2197-2215.
Gavrilets, S., 2004 Fitness Landscapes and the Origin of Species. 117-147.
Ghosh, S., and C. K. Chan, 2016 Analysis of RNA-Seq Data Using TopHat and Cufflinks. Methods Mol
Biol 1374: 339-361.
Gifford, R. J., 2012 Viral evolution in deep time: lentiviruses and mammals. Trends Genet 28: 89-100.
Gillespie, J., 2004 Population Genetics: A Concise Guide. 232.
Haas, B. J., A. Papanicolaou, M. Yassour, M. Grabherr, P. D. Blood et al., 2013 De novo transcript sequence
reconstruction from RNA-seq using the Trinity platform for reference generation and analysis.
Nature protocols 8: 1494-1512.
Hermisson, J., and P. S. Pennings, 2005 Soft sweeps: molecular population genetics of adaptation from
standing genetic variation. Genetics 169: 2335-2352.
Hill, T., B. Koseva and R. L. Unckless, 2019 The genome of *Drosophila innubila* reveals lineage-specific
patterns of selection in immune genes. Molecular Biology and Evolution: 1-36.
Hill, T., and R. L. Unckless, 2017 Baculovirus molecular evolution via gene turnover and recurrent positive
selection of key genes. Journal of Virology 91: e01319-01317.
Hill, T., and R. L. Unckless, 2018 The dynamic evolution of Drosophila innubila Nudivirus. Infection,
Genetics and Evolution: 1-25.
Hill, T., and R. L. Unckless, 2020 Selection and demography shape genomic variation in a 'Sky Island'
species. Biorxiv: 1-35.

Hoffmann, J. A., 2003 The immune response of *Drosophila*. Nature 426**:** 33-38.

Holmes, E. C., 2007 Viral evolution in the genomic age. PLoS Biol 5**:** e278.

Http://broadinstitute.github.io/picard, *Picard*.

Hultmark, D., 2003 *Drosophila* immunity: Paths and patterns. Current Opinion in Immunology 15**:** 12-19.

Jackson, T., 2009 The Use of Oryctes Virus for Control of Rhinoceros Beetle in the Pacific Islands, pp. 133-140 in *Use of Microbes for Control and Eradication of Invasive Arthropods SE - 8*, edited by A. Hajek, T. Glare and M. O'Callaghan. Springer Netherlands.

Jackson, T. A., A. M. Crawford and T. R. Glare, 2005 Oryctes virus - Time for a new look at a useful biocontrol agent. Journal of Invertebrate Pathology 89**:** 91-94.

Joshi, N., and J. Fass, 2011 Sickle: A sliding window, adaptive, quality-based trimming tool for fastQ files. 1.33.

Kaltz, O., and J. A. Shykoff, 1998 Local adaptation in host–parasite systems. Heredity 81**:** 361-370.

Kassambara, A., M. Kosinski and P. Biecek, 2017 survminer: Drawing Survival Curves using'ggplot2'. R package version 0.3 1.

Kelly, D. C., 1982 Baculovirus Replication. Journal of General Virology 63**:** 1-13.

Kondrashov, A. S., S. Sunyaev and F. A. Kondrashov, 2002 Dobzhansky–Muller incompatibilities in protein evolution. PNAS 99**:** 14878-14883.

Lamiable, O., C. Kellenberger, C. Kemp, L. Troxler, N. Pelte *et al.*, 2016 Cytokine Diedel and a viral homologue suppress the IMD pathway in *Drosophila*. Proceedings of the National Academy of Sciences 113**:** 698-703.

Li, H., and R. Durbin, 2009 Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics (Oxford, England) 25**:** 1754-1760.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan *et al.*, 2009 The sequence alignment/map format and SAMtools. Bioinformatics (Oxford, England) 25**:** 2078-2079.

Lipsitch, M., S. Siller and M. Nowak, 1996 The evolution of virulence in pathogens with veritcal and horizontal transmission. Evolution 50**:** 1729-1741.

Liu, X., and Y.-X. Fu, 2015 Exploring population size changes using SNP frequency spectra. Nature genetics 47**:** 555-559.

Maeda, S., S. G. Kamita and A. Kondo, 1993 Host range expansion of Autographa californica nuclear polyhedrosis virus (NPV) following recombination of a 0.6-kilobase-pair DNA fragment originating from Bombyx mori NPV. Journal of virology 67**:** 6234-6238.

Martin, M., 2011 Cutadapt removes adapter sequences from high-throughput sequencing reads. Technical Notes**:** 1-12.

May, R. M., and M. Nowak, 1995 Coinfection and the Evolution of Parasite Virulence. Proceedings: Biological Sciences 261**:** 209-215.

McDonald, J. H., and M. Kreitman, 1991 Adaptive protein evolution at the Adh locus in *Drosophila*. Nature 351**:** 652-654.

McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis *et al.*, 2010 The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. Proceedings of the International Conference on Intellectual Capital, Knowledge Management & Organizational Learning 20**:** 1297-1303.

Merkling, S. H., and R. P. van Rij, 2013 Beyond RNAi: Antiviral defense strategies in Drosophila and mosquito. Journal of Insect Physiology 59**:** 159-170.

Mukherjee, K., H. Campos and B. Kolaczkowski, 2013 Evolution of animal and plant dicers: early parallel duplications and recurrent adaptation of antiviral RNA binding in plants. Mol Biol Evol 30**:** 627-641.

Nielsen, R., 2005 Molecular signatures of natural selection. Annu Rev Genet 39**:** 197-218.

Nielsen, R., C. Bustamante, A. G. Clark, S. Glanowski, T. B. Sackton *et al.*, 2005 A scan for positively selected genes in the genomes of humans and chimpanzees. PLoS Biol 3**:** e170.

Obbard, D. J., K. H. J. Gordon, A. H. Buck and F. M. Jiggins, 2009a The evolution of RNAi as a defence against viruses and transposable elements. Philosophical transactions of the Royal Society of London. Series B, Biological sciences 364: 99-115.

Obbard, D. J., F. M. Jiggins, D. L. Halligan and T. J. Little, 2006 Natural Selection Drives Extremely Rapid Evolution in Antiviral RNAi Genes. Current biology 16: 580-585.

Obbard, D. J., J. J. Welch, K. W. Kim and F. M. Jiggins, 2009b Quantifying adaptive evolution in the *Drosophila* immune system. PLoS Genetics 5: e1000698.

Orr, H. A., 1995 Population genetics of speciation: the evolution of hybrid incompatibilities. Genetics 139: 1805-1813.

Orr, H. A., 2004 Dobzhansky, Bateson, and the genetics of speciation. Genetics 168: 1097-1104.

Orr, H. A., and M. Turelli, 2001 The evolution of postzygotic isolation: accumulating Dobzhansky-Muller incompatibilities. Evolution; international journal of organic evolution 55: 1085-1094.

Palmer, W. H., J. D. Hadfield and D. J. Obbard, 2018a RNA Interference Pathways Display High Rates of Adaptive Protein Evolution in Multiple Invertebrates. Genetics: genetics.300567.302017.

Palmer, W. H., J. Joosten, G. J. Overheul, P. W. Jansen, M. Vermeulen *et al.*, 2019 Induction and suppression of NF- κB signalling by a DNA virus of *Drosophila*. Journal of Virology 93: e01443-01418.

Palmer, W. H., N. Medd, P. M. Beard and D. J. Obbard, 2018b Isolation of a natural DNA virus of Drosophila melanogaster, and characterisation of host resistance and immune responses. PLoS Pathogens 4: e1007050.

Palmer, W. H., N. Medd, P. M. Beard and D. J. Obbard, 2018c Isolation of a natural DNA virus of *Drosophila melanogaster*, and characterisation of host resistance and immune responses. PLoS Pathogens 14: 1-26.

Paradis, E., J. Claude and K. Strimmer, 2004 APE: analyses of phylogenetics and evolution in R language. Bioinformatics 20: 289-290.

Pennings, P. S., 2012 Standing genetic variation and the evolution of drug resistance in HIV. PLoS Computational Biology 8.

Pennings, P. S., S. Kryazhimskiy and J. Wakeley, 2014 Loss and Recovery of Genetic Diversity in Adapting Populations of HIV. PLoS Genetics 10.

Purcell, S., B. Neale, K. Todd-Brown, L. Thomas, M. A. R. Ferreira *et al.*, 2007 PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. The American Journal of Human Genetics 81: 559-575.

Quast, C., E. Pruesse, P. Yilmaz, J. Gerken, T. Schweer *et al.*, 2013 The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic Acids Res 41: D590-596.

Robinson, M. D., D. J. McCarthy and G. K. Smyth, 2009 edgeR: A Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics 26: 139-140.

Rohrmann, G. F., 2013 Baculovirus Molecular Biology. 211.

Sackton, T. B., B. P. Lazzaro, T. A. Schlenke, J. D. Evans, D. Hultmark *et al.*, 2007 Dynamic evolution of the innate immune system in *Drosophila*. Nature Genetics 39: 1461-1468.

Sagulenko, P., V. Puller and R. A. Neher, 2018 TreeTime: Maximum-likelihood phylodynamic analysis. Virus Evol 4: vex042.

Schulz, M. H., D. R. Zerbino, M. Vingron and E. Birney, 2012 Oases: Robust de novo RNA-seq assembly across the dynamic range of expression levels. Bioinformatics 28: 1086-1092.

Shultz, A., and T. B. Sackton, 2019 Immune genes are hotspots of shared positive selection across birds and mammals. eLife 8: e41815.

Smit, A. F. A., and R. Hubley, 2008 RepeatModeler Open-1.0.

Smit, A. F. A., and R. Hubley, 2013-2015 RepeatMasker Open-4.0, pp. RepeatMasker.

Soetaert, K., T. Petzoldt and R. W. Setzer, 2010 Solving Differential Equations in R: Package deSolve. Journal of Statistical Software 33: 1-24.

Stoletzki, N., and A. Eyre-Walker, 2011 Estimation of the neutrality index. Molecular Biology and Evolution 28: 63-70.

Team, R. C., 2013 R: A Language and Environment for Statistical Computing, pp. R Foundation for
Statistical Computing, Vienna, Austria.
Unckless, R. L., 2011 A DNA Virus of *Drosophila*. PLoS ONE 6**:** e26564.
Wang, X.-H. X.-H., R. Aliyari, W.-X. Li, H.-W. Li, K. Kim *et al.*, 2006 RNA interference directs innate
immunity against viruses in adult *Drosophila*. Science 312**:** 452-454.
Wang, Y., and J. a. Jehle, 2009 Nudiviruses and other large, double-stranded circular DNA viruses of
invertebrates: new insights on an old topic. Journal of invertebrate pathology 101**:** 187-193.
Wang, Y., R. G. Kleespies, M. B. Ramle and J. A. Jehle, 2008 Sequencing of the large dsDNA genome of
Oryctes rhinoceros nudivirus using multiple displacement amplification of nanogram amounts of
virus DNA. Journal of Virological Methods 152**:** 106-108.
Webster, C. L., F. M. Waldron, S. Robertson, D. Crowson, G. Ferrari *et al.*, 2015 The discovery,
distribution, and evolution of viruses associated with *Drosophila melanogaster.* PLoS biology 13**:**
e1002210.
West, C., and N. Silverman, 2018 p38b and JAK-STAT signaling protect against invertebrate iridescent
virus 6 infection in *Drosophila*. Plant Pathology 14**:** e1007020.
Williams, G. C., and R. M. Nesse, 1991 The Dawn of Darwinian Medicine. The Quarterly Review of
Biology 66**:** 1-22.
Wilm, A., P. Poh, K. Aw, D. Bertrand, G. Hui *et al.*, 2012 LoFreq: a sequence-quality aware, ultra-sensitive
variant caller for uncovering cell-population heterogeneity from high-throughput sequencing
datasets. Nucleic Acids Research 40**:** 11189-11201.
Wilson, B. A., N. R. Garud, A. F. Feder, Z. J. Assaf and P. S. Pennings, 2016 The population genetics of
drug resistance evolution in natural populations of viral, bacterial and eukaryotic pathogens. Mol
Ecol 25**:** 42-66.
Wu, T. D., and S. Nacu, 2010 Fast and SNP-tolerant detection of complex variants and splicing in short
reads. Bioinformatics (Oxford, England) 26**:** 873-881.
Zambon, R. A., M. Nandakumar, V. N. Vakharia and L. P. Wu, 2005a The Toll pathway is important for
an antiviral response in *Drosophila*. Proceedings of the National Academy of Sciences 102**:** 7257-
7262.
Zambon, R. A., M. Nandakumar, V. N. Vakharia and L. P. Wu, 2005b The Toll pathway is important for
an antiviral response in *Drosophila*. Proceedings of the National Academy of Sciences 102**:** 7257-
7262.
1093