Topography of Hippocampal Connectivity with Sensorimotor Cortex Revealed

by Optimizing Smoothing Kernel and Voxel Size

Douglas D. Burman[1*]

[1]Department of Radiology, NorthShore University HealthSystem, 2650 Ridge Ave.,

Evanston, Illinois 60201, United States of America

*Corresponding author:

Email: DBurman2@northshore.org (DB)

Abstract

Studies of the hippocampus use smaller voxel sizes and smoothing kernels than cortical activation studies, typically using a multivoxel seed with specified radius for connectivity analysis. This study identified optimal processing parameters for evaluating hippocampal connectivity with sensorimotor cortex (SMC), comparing effectiveness by varying parameters during both activation and connectivity analysis. Using both 3mm and 4mm isovoxels, smoothing kernels of 6-10mm were evaluated on the amplitude and extent of motor activation and hippocampal connectivity with SMC. Psychophysiological interactions identified hippocampal connectivity with SMC during volitional movements, and connectivity effects from multivoxel seeds were compared with alternate methods; a structural seed represented the mean connectivity map from all voxels within a region, whereas a functional seed represented the regional voxel with maximal SMC connectivity. With few exceptions, the same parameters were optimal for activation and connectivity. Larger isovoxels showed larger activation volumes in both SMC and the hippocampus; connectivity volumes from structural seeds were also larger, except from the posterior hippocampus. Regardless of voxel size, the 10mm smoothing kernel generated larger activation and connectivity volumes from structural seeds, as well as larger beta estimates at connectivity maxima; structural seeds also produced larger connectivity volumes than multivoxel seeds. Functional seeds showed lesser effects from voxel size and smoothing kernels. Optimal parameters revealed topography in structural seed connectivity along both the longitudinal axis and mediolateral axis of the hippocampus. These results indicate larger voxels and smoothing kernels improve sensitivity for detecting both cortical activation and hippocampal connectivity.

30   *Keywords*: fMRI, connectivity, hippocampus, sensorimotor cortex, smoothing kernel,
31   topography

32   *Declarations of interest*: none

Introduction

Noise can obscure the weak blood oxygen-level dependent (BOLD) response used to detect activation and functional connectivity in functional magnetic resonance imaging (fMRI). A number of approaches have been tried over the years to improve signal detection. Two fundamental considerations are voxel size and the size of the smoothing kernel (1). Optimal voxel size should generally be commensurate with activation volume, although due to susceptibility field gradients, smaller voxel sizes may sometimes increase sensitivity despite less total signal intensity (2, 3). Larger voxel sizes improve signal strength but provide lower spatial resolution, whereas hippocampal studies generally prefer better resolution due its small size; during analysis, voxel dimensions within any plane typically range from 1mm to 3mm (4-12).

Spatial smoothing is used to increase signal-to-noise in BOLD signals and provide smoothness to imaged data (13, 14). Smoothing expands the size of detected activation by reducing noise; generally, the optimal smoothing kernel is 2-3 times the voxel size, both for individual analysis (15, 16) and for large group analysis, with larger smoothing kernels suggested for group analysis in studies with fewer subjects (17). Smoothing kernels may degrade the resolution of signal (18), so some studies of the hippocampus avoid smoothing (12), whereas others use a small Gaussian smoothing kernel ranging from 3-6mm full width at half maximum (FWHM) (6, 8, 10, 19, 20). Nonetheless, several hippocampal studies have used larger smoothing kernels of 8mm (4, 5, 9, 11, 21).

The selection of appropriate processing parameters raises challenges for connectivity studies, especially those designed to examine the influence of hippocampal activity on cortical areas that differ in optimal processing parameters. Generally, the optimal smoothing kernel for connectivity analysis is 2-3 times the voxel size (22), as it is for activation analysis. With few

exceptions (23, 24), smoothing kernels used in hippocampal connectivity studies are small,

ranging from 2-6mm (25).

The effects of voxel and smoothing kernel size in the hippocampus have not been explored

empirically, so their precise relationship to activation and connectivity is unknown. Although

effective in activating the hippocampus and its connections with prefrontal cortex (PFC),

memory tasks may be less than optimal to identify effects of processing parameters, as the actual

size, intensity, and location of memory-related PFC activation may depend on memory content

(26-31). The sensorimotor cortex (SMC), by contrast, shows robust activation based on

movements of the represented body part, known from studies of topography (32-35);

furthermore, a recent study showed hippocampal-SMC connectivity restricted to the hand

representation during a hand movement task (36).

Building on that work, the current study systematically explores the effect of voxel size,

smoothing kernel, and seed selection on task-specific activation and connectivity from

hippocampus to sensorimotor cortex (SMC). A larger voxel size (4mm vs. 3mm isovoxels)

generated a modest increase in cortical activation and hippocampal connectivity volumes, except

from posterior hippocampus, whereas larger smoothing kernels consistently increased both

cortical activation and connectivity, particularly connectivity from structural seeds. Structural

seeds, representing the mean connectivity from all voxels in a specified region of the

hippocampus, proved superior to conventional multivoxel seeds, providing larger connectivity

volumes and a demonstrable topographic organization.

Materials and methods

Data availability

Subjects, task, data acquisition, and processing procedures have previously been described in

detail (36, 37), with the dataset available for download (38).

Subjects

82 Thirteen normal right-handed adults participated in the study (ages 24-59, mean=42.3, five

83 females) following written consent; informed consent procedures complied with the Code of

84 Ethics set forth in the Declaration of Helsinki, and were approved by the Institutional Review

85 Board at the NorthShore University HealthSystem / Research Institute.

Experimental task

Each subject performed a visual/motor task, consisting of 6 cycles of a specified sequence of

visual and motor conditions over a period of 6m 4s. Each cycle included a block of sequential

tapping and a block of repetitive tapping, each separated by a passive visual condition. The task

ended with 10s of passive fixation on a central cross.

During the sequential tapping block, a 4-button sequence to be remembered was displayed while

a metronome ticked at 2 Hz; the subject pressed the remembered sequence of buttons in time

with the metronome once the onscreen instructions were replaced with a cross. Subjects repeated

the 4-button sequence throughout the 16s block, which ended when a circular checkerboard

pattern flickered onscreen for 9s. During this visual block, subjects fixated the center of the

pattern and refrained from moving.

During the repetitive tapping block, the subject was instructed to tap the same finger on both

hands in synchrony with the metronome. Once the instruction screen was replaced with the

number '1', the subject tapped the index finger from both hands; every 4s, the onscreen number

increased by one, and the subject changed finger. This motor condition was also followed by the

passive visual condition.

Button presses were recorded from the right hand during both motor tasks to verify accurate

performance. Behavioral analysis demonstrated motor learning and recall effects during the

sequence learning task; anticipatory movements prior to the metronome demonstrated cognitive

control of movements during both repetitive tapping and sequence learning (36).

MRI data acquisition

Images were acquired using a 12-channel head coil in a 3 Tesla Siemens scanner. Blood-oxygen

level dependent (BOLD) functional images were acquired with the echo planar imaging (EPI),

using the following parameters: time of echo (TE) = 25 ms, flip angle = 90^o, matrix size = 64 x

64, field of view = 22x22 cm, slice thickness = 3 mm, number of slices = 32; time of repetition

(TR) = 2000 ms; and the number of repetitions = 182. These parameters produced in-plane

voxel dimensions of 3.4 x 3.4mm. A structural T1 weighted 3D image (TR = 1600 ms, TE =

3.46 ms, flip angle = 9^o, matrix size = 256 x 256, field of view = 22x22 cm, slice thickness = 1

mm, number of slices = 144) was acquired in the same orientation as the functional images.

fMRI data processing

Data were processed and analyzed using SPM12 software (http://www.fil.ion.ucl.ac.uk/spm).

Images were spatially aligned to the first volume to correct for small movements. After

removing one cycle from a subject's data set due to excessive head movements, maximal RMS

movement for any subject was less than 1.5mm (mean = 0.850 ± 0.291mm). Sinc interpolation

minimized timing-errors between slices; functional images were coregistered to the anatomical

image, normalized to the standard T1 Montreal Neurological Institute (MNI) template, and

resliced for two sets of analyses. For one dataset, functional images were resliced as 3mm

isovoxels, potentially improving in-plane image resolution by incorporating differences in signal

intensity associated with small head movements. For the other dataset, functional images were

resliced as 4mm isovoxels, potentially increasing sensitivity while decreasing image resolution.

To identify task-related activation, copies of images from each dataset were smoothed with a

6mm, 8mm, or 10mm isotropic Gaussian kernel and filtered with a high pass cutoff frequency of

128s. Conditions of interest were specified for sequence learning, visual, and repetitive tapping,

then modeled for block analysis using a canonical hemodynamic response function.

Activation analysis

A parameter estimate of the BOLD response to each condition was generated; motor activation

was identified for each subject by contrasting mean BOLD responses to motor vs. passive visual

conditions. Analyses used an intensity threshold of $p<0.05$ with a family-wise error (FWE)

correction for multiple comparisons, applied to a sensorimotor region of interest (ROI).

Typically, this ROI was specified as the overlap between TD and aal atlas labels for post- plus

precentral gyrus in the WFUPickatlas toolbox for SPM. An additional analysis used the hand

representation as the ROI (36) in order to evaluate whether effects of kernel size and smoothing

kernel had a similar effect on activation parameters within the region known to be active.

For each voxel size and smoothing kernel combination, the mean and standard error of activation

parameters from all individual subjects were recorded and quantitatively compared; these

parameters included t-value threshold (corresponding to that required by the FWE correction),

maxima t-values, and contrast amplitudes. Additionally, paired t-tests applied to individual

subject data allowed statistical evaluation of different voxel sizes with each smoothing kernel,

and of different smoothing kernels with the same voxel size.

For analysis of group effects, BOLD contrasts from individual subjects during the motor memory

and repetitive tapping conditions were entered into a 1-sample t-test for each voxel size /

smoothing kernel combination, recording the same parameters as for individuals (t-threshold,

maxima t-values, and contrast amplitudes). In addition, paired t-tests looked for significant

group differences in activation between 6mm and 10mm smoothing kernels.

Activation maps from different conditions were overlapped to allow visual comparisons for

differences in the size and location of activation.

Connectivity analysis

PPI analysis

Psychophysiological interactions (PPI) were used to identify task-specific connectivity of the

hippocampus with sensorimotor cortex (39-41). The general approach was described previously

(36). For 4mm isovoxels, 78 voxels were identified from the left hippocampus of the normalized

brain and 78 from the right, as delimited by the aal atlas in the WFU PickAtlas toolbox

(http://fmri.wfubmc.edu/software/PickAtlas); each seed was wholly contained within the

hippocampus. For 3mm isovoxels, 278 voxels were identified in the left hippocampus.

At each hippocampal voxel, a contrast was selected and specified to create eigenvariates for all

conditions in the statistical model; an interaction term then specified greater effect on activity

from the motor condition than from the visual condition. After adjustments for regional

differences in timing and baseline activity, a regression analysis showed the magnitude of the

BOLD signal that correlated with this interaction term elsewhere in the brain. The magnitude of

SMC connectivity was quantified from each hippocampal voxel.

Seed selection

Single- and multivoxel approaches were both used to select seeds for connectivity analysis. A

3x3 matrix was created by dividing the hippocampus into thirds of equal distance along the A-P

plane, then dividing each section into equal thirds across the medial-lateral plane. Because the

head of the hippocampus is enlarged, anterior seeds contained more voxels than middle or

posterior seeds, the number of voxels in each structural seed was the same across the medial-

lateral plane. Anatomical seeds were labelled by their position within the matrix (A to C from

anterior to posterior, 1 to 3 from medial to lateral) and the sign of connectivity (positive or

negative).

A "structural" seed's connectivity map was calculated as the mean connectivity from all voxels

within the region. Alternatively, a single "multivoxel" seed was selected from the center of the

anatomical location with a 6mm radius; this provided comparable spatial evaluation of

connectivity from 3mm isovoxels (a diameter of 4 voxels) and 4mm isovoxels (a diameter of 3

voxels). To minimize asymmetry effects, comparisons in connectivity across voxel sizes were

limited to the left hippocampus.

Single-voxel seeds could be based either on anatomical location or function. Anatomical

selection was based on spaced intervals within the structural seeds, providing finer-grain analysis

that was useful for examining topography. "Functional" seed selection was restricted to voxels

within (or adjacent to) a structural seed showing significant connectivity; the single voxel was

selected that showed maximal connectivity within the SMC mask. By reflecting individual

variability in functional localization, functional seeds sought to provide a better estimate of the

magnitude and extent of hippocampal influence on SMC activity. Functional seeds were

identified from the left and right hippocampus for both sequence learning ("memseed") and

repetitive tapping ("tapseed"), using global analysis from both sides to improve sensitivity (36).

PPI Group analysis

For the sequence learning task, beta estimates of connectivity from each subject's left

hippocampus were entered into a 1-sample random effects analysis; the hand representation

delineated from activation analysis was used as the ROI, applying an intensity threshold of

$p<0.05$ with FWE correction. A separate analysis was run for multivoxel and structural seeds at

4 combinations of voxel size (3mm and 4mm) with smoothing kernel (6mm and 10mm).

To improve sensitivity for repetitive tapping, beta estimates of connectivity from each subject's

left and right hippocampus were entered into an ANOVA model for global analysis; as before,

the hand representation delineated from activation analysis was used as the ROI. This ROI

provides optimal sensitivity for detecting effects, as hippocampal connectivity with SMC during

volitional finger movements is selective for the hand representation (36).

Results

Effects of voxel size and smoothing kernels on SMC activation

Group and individual analyses showed similar trends of voxel size and smoothing kernels on

patterns of SMC activation, as described below. During group analyses, differences in group

activation between 3mm and 4mm isovoxels appeared minor, as were most smoothing kernel

effects *except* for activation volume. At the individual level, however, effects from different

voxel sizes and smoothing kernels were both significant

For both 3mm and 4mm isovoxels, maxima values generally decreased incrementally with larger

smoothing kernels. The maximal t-value decreased (Table 1, Fig 1A), a pattern observed during

both group and individual analyses; lower maxima t-values were observed during group analysis.

Estimates for the magnitude of contrast effects also decreased, especially for individual analyses;

the magnitude of effects nearly converged for individual and group analyses for 10mm

smoothing (see bottom of graphs). In addition, the threshold to identify significant activation

dropped, resulting from less local variability in signal amplitude with increasing kernel size.

During both individual and group analyses, the net effect was larger activation volumes with

larger smoothing kernels, with 4mm isovoxels always generating larger activation volumes than

3mm isovoxels (Fig 1B). With a 10mm smoothing kernel, group differences in SMC activation

between the two voxel sizes overlapped extensively, but with a slightly larger activation volume

for 4mm isovoxels during sequence learning (Fig 1C).

Larger smoothing kernels thus provide greater sensitivity for detecting activation by reducing

noise (random variability in signal). Comparing changes in activation volume with different

ROIs provides further insight (Table 1). With the SMC mask, large increases in activation

volume were observed as the smoothing kernel increased 6mm to 10mm, volume increases of

84.2 - 440.0%. These increases were much smaller within the hand representation, never

exceeding 33.3% and nearly non-existent for 3mm isovoxels (volume increases of 1.9 – 1.6% for

repetitive tapping and sequence learning, respectively). This suggests larger smoothing kernels

are particularly effective for improving statistical detection by reducing signal variability outside

the activated region, thus requiring a smaller mean difference in signal to detect activation.

Because data from the same individuals were processed with different voxel size and smoothing

kernels, differences in activation parameters resulting from different processing methods could

be measured directly for each individual. The standard error for mean differences in parameters

across the group of subjects often differed greatly for both repetitive tapping and sequence

learning; nonetheless, paired t-tests showed that individuals consistently had lower activation

thresholds and greater activation volumes with larger (4mm) isovoxels, regardless of smoothing

kernel (Table 2).

For each task, the effects of different smoothing kernels on activation parameters were also

evaluated with paired t-tests during each motor task (Tables 3-4). With 3mm isovoxels, larger

smoothing kernels during both motor tasks generated higher activation volumes, but significantly

lower maxima t-values, contrast magnitudes, and activation thresholds. Often, these patterns

were also observed with 4mm isovoxels; however, no differences were observed between 8mm

and 10mm smoothing kernels for activation threshold during repetitive tapping (Table 3), or for

the maxima t-values and contrast magnitudes during sequence learning (Table 4).

To summarize, 4mm isovoxels provided greater sensitivity to activation than 3mm isovoxels

(lower threshold and higher activation volume), as did 10mm compared with 6mm smoothing

kernels (for both 3mm and 4mm isovoxels). For 4mm isovoxels, differences between 8mm and

10mm smoothing kernels were less differentiated for some activation parameters.

Effect of smoothing kernels on hippocampal motor activation

Using the hippocampal ROI mask, group analysis failed to demonstrate motor activation for any

combination of voxel size and smoothing kernel. Using paired t-tests to compare activation

following 10mm vs. 6mm smoothing, however, more intense activation was observed from the

larger smoothing kernel during both motor tasks (Fig 2). For both 3mm and 4mm isovoxels, the

most intense activation (orange) was observed along the border between structural seeds B1 and

C1; mildly elevated activation (yellow) was observed throughout B1.

Hippocampal connectivity: Structural seeds

Effects of voxel size and smoothing on hippocampal connectivity

Inverse connectivity from many regions of the hippocampus was observed during sequence

learning (Fig 3 and Table 5). For a given combination of voxel size and smoothing kernel, a

structural seed (striped and dotted bars) generated a larger volume of SMC connectivity than the

corresponding multivoxel seed (dashed and solid bars), even when the parameter estimate at the

postcentral (black dot) or precentral maximum (red dot) was larger for the multivoxel seed. In

some cases, differences in methodology could determine whether connectivity remained

significant following an additional correction for testing these nine regions (stars). The

connectivity from A1 and A2 seeds, for example, were significant only for structural seeds

derived from 4mm isovoxels with 10mm smoothing.

In anterior (A1-A3) and middle hippocampus (B1-B2), the volume of connectivity during

random effects analysis was typically greater for 4mm isovoxels, both for structural and

multivoxel seeds; only in posterior hippocampus (C1-C3) was the volume of connectivity greater

for 3mm isovoxels (see also Table 6). In paired t-tests, beta parameter estimates in structural

seeds were significantly larger for 4mm voxels and also with the larger smoothing kernel

(10mm), especially at postcentral maxima (Table 6). Overall, the volume of connectivity was

larger for structural seeds than multivoxel seeds (invariably so when effects survived the FWE

correction), even when maxima parameter estimates were significantly smaller (observed only

with the smaller 6mm smoothing kernel).

Connectivity demonstrated through different methodologies overlapped extensively, particularly

when the volume of connectivity was similar (see Fig 3, seed B1- at bottom left). When one

method generated a much larger volume, however, an additional locus of connectivity could

appear (in bottom right of Fig 3, see precentral connectivity for S3 method at seed C1-). The

volume of connectivity thus reflected the sensitivity of the method.

To summarize, structural seeds improved volume and sensitivity for detecting connectivity

compared to traditional multivoxel seeds. For structural seeds, larger voxels (4mm vs. 3mm)

improved connectivity except in posterior hippocampus; a larger smoothing kernel (10mm vs.

6mm) improved connectivity for both 3mm and 4mm voxels.

Effects of smoothing connectivity during repetitive tapping

For repetitive tapping, connectivity was examined using global analysis from both the left and

right hippocampus, which improves sensitivity for detecting connectivity (36). We further

examined smoothing kernel effects on connectivity from structural seeds during repetitive

tapping, comparing effects of 6mm and 10mm smoothing kernels on 4mm isovoxels.

Structural seeds showed similar maxima locations in pre- and postcentral gyri with both

smoothing kernels, but the extent of connectivity was invariably greater with the larger kernel

(Table 7). Depending on the size of the smoothing kernel, a topographic arrangement of

connectivity was discernible (Fig 4). In the central third of the hippocampus, a topography was

observed along the medial / lateral axis with the 6mm smoothing kernel (Fig 4A left);

connectivity from more lateral seeds (cyan and navy blue) extended progressively further

posterior and superior. This topography was largely obscured with 10mm smoothing (Fig 4A,

right), because larger clusters of connectivity from the middle seed (cyan) produced extensive

overlap.

Along the anterior / posterior axis of the hippocampus, however, the topographic organization

was more apparent with the 10mm smoothing kernel (Fig 4B). Connectivity for each seed was

centered in the precentral gyrus, barely extending across the central sulcus into the postcentral

gyrus. From the anterior hippocampal seed A1 (red), intense connectivity projected further

anterolateral; the middle seed B1 projected superomedial (yellow), whereas the posterior seed C1

projected further medial and posterior (green). Because its clusters were smaller and spotty, this

topography was indiscernible with the 6mm smoothing kernel.

Topography was also demonstrable from individual voxels with 10mm smoothing along both the

medial-lateral and longitudinal axes (Fig 5). In the medial-lateral plane, the organization

described above is seen in the pattern of overlapping connectivity maps (Fig 5A, left), as well as

differential connectivity maps that illustrate which seeds generated the greatest amplitude of

connectivity (Fig 5A, right). Along the longitudinal axis, less overlap was observed from

individual voxels (Fig 5B), perhaps because of the greater distance between sampled voxels

along this axis. From voxels in both planes, connectivity included both pre- and postcentral

gyrus (Table 6).

Precentral connectivity was more extensive in the right hemisphere during repetitive tapping (Fig

5), whereas postcentral connectivity was more extensive in the left hemisphere during sequential

learning (Fig 4).

Overall, the 10mm smoothing kernel provided better sensitivity for structural seeds to

demonstrate both hippocampal connectivity and topography.

317 **Hippocampal connectivity: Functional seeds**

Functional seeds showed unilateral connectivity during group analysis. Tapseed⁻ generated

right-hemispheric connectivity during the repetitive tapping task (Fig 6A, red), whereas

memseed⁻ generated connectivity in the left hemisphere (Fig 6A, yellow). Smoothing kernels of

6mm and 10mm showed no differences in connectivity during repetitive tapping or in postcentral

gyrus during sequence, whereas the extent of precentral connectivity during sequence learning

was larger with the 10mm kernel for both memseed⁻ (2880 vs. 1600 mm^3) and memseed⁺ (1728

vs. 320 mm^3).

The 10mm smoothing kernel increased the intensity of connectivity as well as its extent. As

noted earlier, a few hippocampal voxels individually showed greater motor activation with a

10mm vs. 6mm smoothing kernel; these voxels also showed significantly greater connectivity

with 10mm smoothing (orange voxels, arrows) interspersed within the area of connectivity

shown by functional seeds in both motor tasks (Fig 6B).

Discussion

This study evaluated the effects of seed selection (structural vs. multivoxel vs. functional), voxel

size (3mm vs. 4mm isovoxels), and smoothing kernels (6-10mm) on hippocampal connectivity

with sensorimotor cortex; similarities to the effects of voxel size and smoothing kernels on SMC

motor activation were also evaluated. Larger voxels modestly improved sensitivity for detecting

activation in SMC; larger voxels also improved detection of connectivity from anterior regions

of the hippocampus, although showed lesser sensitivity from the posterior third of the

hippocampus. Larger smoothing kernels improved sensitivity for detecting both hippocampal

activation and connectivity; this sensitivity allowed the topography of hippocampal connectivity

to be mapped from single voxels as well as structural seeds. Although structural seeds showed

greater sensitivity for connectivity than multivoxel seeds, they were more affected by the size of

the smoothing kernel than functional seeds, which addressed individual variability by selecting

voxels with maximal SMC connectivity within a hippocampal region.

Effects of voxel size and smoothing kernel on cortical activation

Voxel size had modest effects on activation in this study. In SMC, larger voxels (4mm) showed

a smaller statistical threshold and a larger activation volume during both repetitive tapping and

sequence learning. These effects were small, detected through direct within-subject

comparisons, whereas group comparisons of the mean showed no effect. Random-effects

activation revealed only minor differences from voxel size in the area of activation along cluster

borders; furthermore, no significant differences in contrast magnitude were observed, regardless

of smoothing kernel. Although the larger 4mm isovoxels provide slightly better sensitivity for

detecting cortical activation, this effect was modest.

The effect of smoothing kernel on cortical activation depended on the ROI. With the SMC

mask, increases in the smoothing kernel from 6mm to 10mm resulted in the detection of much

greater activation volumes; within the hand representation, this increase in activation volume

was much less. This suggests the larger smoothing kernel more effectively reduced random

variability (noise) from inactive regions within the SMC mask, reducing the mean difference in

signal required to detect activation. This interpretation is consistent with observed decreases in

the maxima t-value, contrast magnitude, and threshold.

During both individual and group analysis, the largest (10mm) smoothing kernel applied to 4mm

isovoxels was most sensitive for detecting activation. This was not simply an overestimation of

activation size from a large smoothing kernel, as suggested by others (15); global analysis across

both tasks demonstrated an even larger, bilateral area of activation consistent with task behavior

(36). With these processing parameters, the contrast estimate at the maxima from group analysis

approached the mean value from individual analyses, suggesting this combination of voxel size

and smoothing kernel provided the most accurate group representation of sensorimotor

activation.

Effect of smoothing kernel on hippocampal activation

Using conventional group analysis, the hippocampus did not show motor activation during

repetitive movements or sequence learning. By using paired t-tests to compare motor activation

across different smoothing kernels, greater activation was demonstrated with the 10mm

smoothing kernel (compared with 6mm), evident with both 3mm and 4mm isovoxels. This

differential activation occurred in the medial hippocampus, mostly within the central third of the

hippocampus but extending slightly into the posterior third, most intense at the boundary

between the B1 (central-medial) and C1 (posterior-medial) structural seeds.

During smoothing, local variability is minimized by recalculating the BOLD signal based upon

peaks and their surrounding signal, fit to a Gaussian curve. When activity surrounding a 4mm

isovoxel showing peak activity is symmetrical, both a 6mm and 10mm smoothing kernel utilize

the magnitude of signal from a single voxel on each side in its calculation, differing only in the

weighting provided from the adjacent voxel (3mm on each side of the peak 4mm-voxel using a

10mm kernel, and 1mm on each side of the peak 4mm-voxel with a 6mm kernel). Activity

surrounding a peak voxel near the edge of gray matter is asymmetrical; the calculated peak

coordinate is shifted toward the higher activity within gray matter, away from the edge (42, 43).

Thus, differential activation from the larger smoothing kernel did not result from extending the

smoothed signal outside hippocampal boundaries; instead, the 10mm kernel provided a better

estimate of local activation by more fully incorporating information from nearby activity.

The region showing differential activation also showed more intense and widespread SMC

connectivity following 10mm smoothing. Improved sensitivity for detecting local hippocampal

activity with the larger smoothing kernel is thus useful for detecting its connectivity with other

regions.

Effect of voxel size and analytical approach on hippocampal

connectivity

Because direct point-to-point comparisons is not possible, effects of voxel size were identified by

comparing the volume of significant connectivity and maximal parameter estimates, using

different methods and smoothing kernels. Using a 3x3 matrix, all hippocampal regions were

sampled, thereby allowing detection of regional effects.

Traditionally, either the entire hippocampus or a multivoxel subregion is specified as a seed for

connectivity analysis (44-47). Multivoxel seeds average hippocampal activity before looking for

correlations elsewhere in the brain; this differs from our structural seed alternative, which

identified the magnitude of correlations from each voxel within a hippocampal region, then

averaged these connectivity maps. The method of analysis showed inconsistent effects on the

maximal parameter estimate from a region; however, greater connectivity volumes were

consistently observed from structural seeds in all regions, indicating this approach improves

sensitivity for detecting connectivity. Indeed, multivoxel seeds failed to demonstrate significant

connectivity in anterior hippocampus. Furthermore, the topographic organization of

hippocampal connectivity only emerged with structural seed analysis.

In posterior hippocampus, 3mm voxels generated greater connectivity volumes than 4mm

voxels, particularly in posteromedial hippocampus. This likely reflects both anatomical and

functional properties of the hippocampus. The posterior hippocampus is thinner (four 4mm-

voxels wide, compared with six in middle and anterior regions); furthermore, posteromedial

hippocampus may be functionally more uniform. More lateral regions include positive

connectivity, particularly in middle-to-anterior regions (e.g., see Fig S2 from Burman, 2019

(36)). A sharp transition between functionally distinct regions may thus be requisite for smaller

voxels to demonstrate greater connectivity. By contrast, 4mm isovoxels generated greater

connectivity volumes in anterior and central regions of the hippocampus, as well as generating

larger parameter estimates at structural seed maxima (Fig 3 and Table 6).

In summary, structural seeds were more effective for demonstrating hippocampal connectivity.

Except in posterior areas, perhaps due to a transition to a functionally distinct area, 4mm

generated more extensive connectivity maps than 3mm voxels. In this study, a topographical

organization to hippocampal connectivity was only evident with the structural seed analysis

using 4mm voxels.

Effect of smoothing kernel on hippocampal connectivity

This study showed different patterns of inverse hippocampal connectivity with SMC during

sequence learning and repetitive tapping: the left hippocampus generated both pre- and post-

central connectivity maxima in left SMC during sequence learning, whereas bilateral

hippocampal seeds generated a single connectivity maximum in the right pericentral gyrus

during repetitive tapping. The size of the smoothing kernel showed similar effects on

connectivity during both tasks.

Structural seeds demonstrated larger clusters of SMC connectivity with the larger kernel in both

motor tasks. A connectivity map from a structural seed represents the mean connectivity from

every voxel within the designated region of hippocampus, so larger connectivity clusters reflect

the effectiveness of the smoothing kernel in reflecting local regional activity (see also (48)).

With the 6mm smoothing kernel, a topographic pattern from structural seeds was evident along

the medial-lateral axis of the hippocampus, but not the longitudinal axis, where small and

disparate clusters were generated with no apparent topography. With the 10mm kernel, a

topographic pattern was partially obscured along the medial-lateral axis, due to extensive overlap

in connectivity from adjacent seeds, yet a topography along this axis was still evident from

connectivity of individual voxels. Furthermore, the 10mm smoothing kernel was needed to

demonstrate topography along the longitudinal axis of the hippocampus.

The size of the smoothing kernel may be a factor in detecting functional heterogeneity within the

hippocampus. A resting state connectivity study using a small (3mm) smoothing kernel failed to

find functional heterogeneity along the anterior-posterior axis (49), whereas a study of

connectivity from the hippocampus to cortical regions during episodic memory found functional

heterogeneity along the same axis using a larger (5mm) smoothing kernel (25).

In summary, the 10mm smoothing kernel was advantageous for demonstrating hippocampal

topography in its connectivity with SMC.

Topography of hippocampal connectivity

Similar to previous studies, hippocampal connectivity in this study showed a topographical

organization along both the long anterior-posterior axis (50-53) and medial-lateral axis (50).

In the current study, hippocampal connectivity with SMC was preferentially seen within the

SMC hand representation for both motor tasks, with location and laterality similar to group

activation by the tasks. Group activation and inverse connectivity were both limited to the left

SMC during sequence learning (reflecting right-handed movements) and the right SMC during

repetitive tapping (despite bilateral hand movements). Furthermore, activation and connectivity

both covered the breadth of the postcentral gyrus within the hand representation during sequence

learning, whereas activation and connectivity were both centered in precentral gyrus during

repetitive tapping, with postcentral connectivity limited to the region adjacent to the central

sulcus. Whether these task differences arose from differences in behavioral requirements, hand

dominance (right-handed subjects), or hemispheric differences in motor function is unknown.

The role of hippocampal topography to SMC function is unclear. Within the hand area of the

precentral gyrus, overlapping but distinguishable regions are involved in wrist and individual

finger movements (32, 54-56); multiple representations for each digit may differentially reflect

flexion and extension movements (35) or different degrees of movement complexity (57).

Similarly, somatotopy for individual fingers in postcentral gyrus are overlapping but

distinguishable, with multiple representations across the anterior-posterior width of the gyrus

(58-60). Methods here did not allow mapping onto individual finger representations, or even

flexion vs. extension, so the functional significance of the hippocampal topography observed

here is unclear.

Different cognitive functions have been suggested for anterior vs. posterior hippocampus,

differences such as encoding vs. recall (61, 62), internal vs. external attention (63),

gist/conceptual vs. detailed/spatial information (64-66), and pattern completion vs. separation

(67). In the current study, the topographical pattern of connectivity observed from anterior vs.

posterior hippocampus is inconsistent with these roles, particularly as the repetitive tapping task

required paced, volitional movements without learning or memory recall (36). Topographical

connectivity with different regions within the pre- or postcentral hand representation suggest a

role more directly involved in motor control, perhaps coordinating cognitive and motor functions

to facilitate and suppress muscle actions required for the correct timing of finger movements (37,

56).

478   **Practical considerations for future studies**

Many hippocampal studies have used small voxels and smoothing kernels, believing these

processing parameters more accurately reflect true hippocampal activity.   The current study

suggests this may not always be the case, particularly when examining hippocampal connectivity

with cortical areas. This study specifically examined the effects of voxel size and smoothing

kernel on hippocampal activation and connectivity with SMC; however, results should apply to

other hippocampal studies as well, as observed effects from changes in processing parameters are

consistent with previous studies (as noted earlier).

Results also indicate greater sensitivity can be obtained with structural seeds, rather than

conventional multivoxel seeds. For structural seed analysis, connectivity is calculated from each

voxel of the hippocampus, then a mean connectivity map is calculated from all voxels in a

region. This approach can map connectivity from the entire hippocampus, particularly

advantageous when the precise location of a functional region is unknown.

Paired t-tests comparing cortical activation with 4mm vs. 3mm isovoxels showed lower

thresholds and larger volumes of activation from larger voxels, with the additional volume

expanding the edges of an activation cluster. Similarly, larger smoothing kernels decreased the

threshold and increased the volume of cortical activation. Even small differences in activation

between 8mm and 10mm smoothing kernels were significant during sequence learning, although

these differences did not reach significance for 4mm voxels during repetitive tapping.

Parameters optimal for detecting SMC cortical activation also improved hippocampal

connectivity (10mm smoothing kernels and 4mm voxels, except from the posterior third of the

hippocampus). This similarity suggests better detection of cortical activity might allow better

correlation with hippocampal activity. Larger smoothing kernels also improved the signal-to-

noise ratio with local hippocampal activity, however, as shown in direct comparison between

10mm vs. 6mm smoothing kernels. Furthermore, localized areas with elevated hippocampal

activity from the 10mm smoothing kernel also showed elevated connectivity with SMC,

indicating this activation was functionally relevant. The effectiveness of these parameters thus

reflect both cortical and local hippocampal properties.

Because the size and structure of the hippocampus is affected by a wide array of factors,

including age, exercise, depression, and stress (68-73), its functional localization must be

variable. Functional seeds sought to reduce effects of individual variability during group

analysis. Restricted to structural regions showing significant connectivity during group analysis,

the voxel showing maximal connectivity within the entire SMC was identified. Using this ROI

during seed selection, the selection of functional seeds did not bias the outcome; connectivity

maxima used for seed selection were scattered, located in both hemispheres and often outside the

hand representation (36). These functional seeds nonetheless showed selective connectivity

within the hand region during group analysis. The current study showed that the sensitivity of

functional seeds was minimally affected by the size of the smoothing kernel, suggesting

smoothing mitigates the effects of individual variability during group analysis, but also that

sensitivity for demonstrating connectivity with non-optimal parameters can be improved by

identifying functional seeds.

Conclusions

This study shows that processing fMRI data with a larger voxel size (4mm) and smoothing

kernel (8-10mm) can improve sensitivity to hippocampal activity and its connectivity with

cortical areas; optimizing these processing parameters uncovered topography in hippocampal

connectivity along two axes. Structural seeds that represent the mean connectivity from all

voxels in a hippocampal region provide better sensitivity than traditional multivoxel seeds, and

have the additional advantage of mapping connectivity throughout the hippocampus.

Acknowledgments

527 The author wishes to thank the Center for Advanced Imaging (CAI) at NorthShore University

528 HealthSystem for its administrative support.

529 Funding: This research did not receive any specific grant from funding agencies in the public,

530 commercial, or not-for-profit sectors.

References

1. Hopfinger JB, Büchel C, Holmes AP, Friston KJ. A study of analysis parameters that influence the sensitivity of event-related fMRI analyses. NeuroImage 2000;11(4):326-333.

2. Chen N-K, Dickey CC, Yoo S-S, Guttmann CRG, Panych LP. Selection of voxel size and slice orientation for fMRI in the presence of susceptibility field gradients: application to imaging of the amygdala. NeuroImage 2003;19(3):817-825.

3. Thulborn KR, Chang SY, Shen GX, Voyvodic JT. High-resolution echo-planar fMRI of human visual cortex at 3.0 tesla. NMR in Biomedicine 1997;10(4-5):183-190.

4. Danker JF, Tompary A, Davachi L. Trial-by-trial hippocampal encoding activation predicts the fidelity of cortical reinstatement during subsequent retrieval. Cerebral Cortex 2017;27(7):3515-3524.

5. Forcato C, Bavassi L, De Pino G, Fernández RS, Villarreal MF, Pedreira ME. Differential left hippocampal activation during retrieval with different types of reminders: An fMRI study of the reconsolidation process. PLoS One 2016;11(3):e0151381.

6. Lech RK, Suchan B. Involvement of the human medial temporal lobe in a visual discrimination task. Behavioural Brain Research 2014;268:22-30.

7. Liu Z-X, Shen K, Olsen RK, Ryan JD. Visual sampling predicts hippocampal activity. Journal of Neuroscience 2017;37(3):599-609.

8. Malivoire BL, Girard TA, Patel R, Monson CM. Functional connectivity of hippocampal subregions in PTSD: relations with symptoms. BMC psychiatry 2018;18(1):129.

9.   Mayes AR, Montaldi D, Roper A, Migo EM, Gholipour T, Kafkas A. Amount, not

strength of recollection, drives hippocampal activity: A problem for apparent word

familiarity-related hippocampal activation. Hippocampus 2019;29(1):46.

10.   Qin S, Cho S, Chen T, Rosenberg-Lee M, Geary DC, Menon V. Hippocampal-

neocortical functional reorganization underlies children's cognitive development. Nature

Neuroscience 2014;17(9):1263-1269.

11.   Sperling R, Chua E, Cocchiarella A, Rand-Giovannetti E, Poldrack R, Schacter DL, et al.

Putting names to faces: Successful encoding of associative memories activates the

anterior hippocampal formation. NeuroImage 2003;20(2):1400-1410.

12.   Zeidman P, Lutti A, Maguire EA. Investigating the functions of subregions within

anterior hippocampus. Cortex 2015;73:240-256.

13.   Acton PD, Friston KJ. Statistical parametric mapping in functional neuroimaging: beyond

PET and fMRI activation studies. European Journal of Nuclear Medicine 1998(25):663-

667.

14.   Worsley KJ. Spatial smoothing of autocorrelations to control the degrees of freedom in

fMRI analysis. NeuroImage 2005;26:635-641.

15.   Liu P, Calhoun V, Chen Z. Functional overestimation due to spatial smoothing of fMRI

data. Journal of Neuroscience Methods 2017;291:1.

16.   Pajula J, Tohka J. Effects of spatial smoothing on inter-subject correlation based analysis

of FMRI. Magnetic Resonance Imaging 2014;32(9):1114-1124.

17.   Mikl M, Marecek R, Hlustík P, Pavlicová M, Drastich A, Chlebus P, et al. Effects of

spatial smoothing on fMRI group inferences. Magnetic Resonance Imaging

2008;26(4):490-503.

18. Murphy JE, Yanes JA, Kirby LAJ, Reid MA, Robinson JL. Left, right, or bilateral

amygdala activation? How effects of smoothing and motion correction on ultra-high

field, high-resolution functional magnetic resonance imaging (fMRI) data alter

inferences. Neuroscience Research 2020;150:51-59.

19. Adcock RA, Thangavel A, Whitfield-Gabrieli S, Knutson B, Gabrieli JDE. Reward-

motivated learning: mesolimbic activation precedes memory formation. Neuron

2006;50(3):507-517.

20. Selmeczy D, Fandakova Y, Grimm KJ, Bunge SA, Ghetti S. Longitudinal trajectories of

hippocampal and prefrontal contributions to episodic retrieval: Effects of age and

puberty. Developmental Cognitive Neuroscience 2019;36:100599-100611.

21. Rigoli F, Michely J, Friston KJ, Dolan RJ. The role of the hippocampus in weighting

expectations during inference under uncertainty. Cortex 2019;115:1-14.

22. Chen Z, Calhoun V. Effect of spatial smoothing on task fMRI ICA and functional

connectivity. Frontiers in Neuroscience 2018;12:15.

23. Smith APR, Stephan KE, Rugg MD, Dolan RJ. Task and content modulate amygdala-

hippocampal connectivity in emotional retrieval. Neuron 2006;4(49):631-638.

24. Wang L, LaViolette P, O'Keefe K, Putcha D, Bakkour A, Van Dijk KRA, et al. Intrinsic

connectivity between the hippocampus and posteromedial cortex predicts memory

performance in cognitively intact older individuals. NeuroImage 2010;2(51):910-917.

25. Geng F, Redcay E, Riggins T. The influence of age and performance on hippocampal

function and the encoding of contextual information in early childhood. NeuroImage

2019;195:433-443.

26.   Ankudowich E, Pasvanis S, Rajah MN. Age-related differences in prefrontal-

hippocampal connectivity are associated with reduced spatial context memory.

Psychology and aging 2019;34(2):251.

27.   Cohen MX. Hippocampal-prefrontal connectivity predicts midfrontal oscillations and

long-term memory performance. Current Biology 2011;21.

28.   Courtney SM, Petit L, Haxby JV, Ungerleider LG. The role of prefrontal cortex in

working memory: examining the contents of consciousness. Philosophical Transactions

of the Royal Society of London. Series B: Biological Sciences 1998;353(1377):1819-

1828.

29.   Grady CL, McIntosh AR, Craik FI. Age-related differences in the functional connectivity

of the hippocampus during memory encoding. Hippocampus 2003;13(5):572-86.

30.   Eichenbaum H. Prefrontal–hippocampal interactions in episodic memory. Nature

Reviews Neuroscience 2017;18(9):547-558.

31.   Preston AR, Eichenbaum H. Interplay of Hippocampus and Prefrontal Cortex in Memory.

Current Biology 2013;23(17):R764-R773.

32.   Burman DD, Lie-Nemeth T, Brandfonbrener AG, Parisi T, Meyer JR. Altered finger

representations in sensorimotor cortex of musicians with focal dystonia: Precentral

cortex. Brain Imaging and Behavior 2009;3(1):10-23.

33.   De Lange FP, Hagoort P, Toni I. Neural topography and content of movement

representations. Journal of Cognitive Neuroscience 2005;17(1):97-112.

34.   Pascual-Leone A, Grafman J, Hallett M. Modulation of cortical motor output maps

during development of implicit and explicit knowledge.[see comment]. Science

1994;263(5151):1287-9.

35.   Zartl M, Kapfer T, Muellbacher W. Functional topography of cortical thumb movement

621   representations in human primary motor cortex. Brain Topography 2014;27(2):228-239.

622   36.   Burman DD. Hippocampal connectivity with sensorimotor cortex during volitional finger

623   movements: Laterality and relationship to motor learning. PloS One

624   2019;14(9):e0222064.

625   37.   Burman DD. Hippocampal connectivity with sensorimotor cortex during volitional finger

626   movements. II. Spatial and temporal selectivity. bioRxiv 2018;doi:

627   https://doi.org/10.1101/479436.

628   38.   Burman DD. Functional MRI dataset during passive visual stimulation and two motor

629   conditions (sequence learning, paced repetitive tapping). NITRC 2019.

630   39.   Friston KJ, Buechel C, Fink GR, Morris J, Rolls E, Dolan RJ. Psychophysiological and

631   modulatory interactions in neuroimaging. NeuroImage 1997;6(3):218-29.

632   40.   Gitelman DR, Penny WD, Ashburner J, Friston KJ. Modeling regional and

633   psychophysiologic interactions in fMRI: The importance of hemodynamic deconvolution.

634   Neuroimage 2003;19(1):200-207.

635   41.   O'Reilly JX, Woolrich MW, Behrens TEJ, Smith SM, Johansen-Berg H. Tools of the

636   trade: Psychophysiological interactions and functional connectivity. Social Cognitive and

637   Affective Neuroscience 2012;7(5):604-609.

638   42.   Jo HJ, Lee J-M, Kim J-H, Choi C-H, Gu B-M, Kang D-H, et al. Artificial shifting of

639   fMRI activation localized by volume- and surface-based analyses. NeuroImage

640   2008;40(3):1077-1089.

641   43.   Sacchet MD, Knutson B. Spatial smoothing systematically biases the localization of

642   reward-related brain activity. NeuroImage 2013;66:270-277.

44.   Blankenship SL, Redcay E, Dougherty LR, Riggins T. Development of hippocampal

functional connectivity during childhood. Human Brain Mapping 2017;1(38):182-201.

45.   Chang J, Yu R. Hippocampal connectivity in the aftermath of acute social stress.

Neurobiology of Stress 2019;11:100195.

46.   Haneef Z, Lenartowicz A, Yeh HJ, Levin HS, Engel Jr J, Stern JM. Functional

connectivity of hippocampal networks in temporal lobe epilepsy. Epilepsia

2014;55(1):137-145.

47.   Zeidman P, Mullally SL, Maguire EA. Constructing, Perceiving, and Maintaining

Scenes: Hippocampal Activity and Connectivity. Cerebral cortex (New York, NY: 1991)

2015;25(10):3836-3855.

48.   Wu CW, Chen C-L, Liu P-Y, Chao Y-P, Biswal BB, Lin C-P. Empirical evaluations of

slice-timing, smoothing, and normalization effects in seed-based, resting-state functional

magnetic resonance imaging analyses. Brain Connectivity 2011;1(5):401-410.

49.   Wang S-F, Ritchey M, Libby LA, Ranganath C. Functional connectivity based

parcellation of the human medial temporal lobe. Neurobiology of Learning and Memory

2016(134):123-134.

50.   de Wael RV, Larivière S, Caldairou Bt, Hong S-J, Margulies DS, Jefferies E, et al.

Anatomical and microstructural determinants of hippocampal subfield functional

connectome embedding. Proceedings of the National Academy of Sciences of the United

States of America 2018;115(40):10154.

51.   Qin S, Duan X, Supekar K, Chen H, Chen T, Menon V. Large-scale intrinsic functional

network organization along the long axis of the human medial temporal lobe. Brain

Structure and Function 2016;221(6):3237-3258.

52.   Robinson JL, Barron DS, Kirby LAJ, Bottenhorn KL, Hill AC, Murphy JE, et al.

667   Neurofunctional topography of the human hippocampus. Human Brain Mapping

668   2015;36(12):5018-5037.

669   53.   Strange BA, Witter MP, Lein ES, Moser EI. Functional organization of the hippocampal

670   longitudinal axis. Nature Reviews Neuroscience 2014;15(10):655.

671   54.   Beisteiner R, Windischberger C, Lanzenberger R, Edward V, Cunnington R, Erdler M, et

672   al. Finger somatotopy in human motor cortex.[see comment]. NeuroImage 2001;13(6 Pt

673   1):1016-26.

674   55.   Kleinschmidt A, Nitschke MF, Frahm J. Somatotopy in the human motor cortex hand

675   area. A high-resolution functional MRI study. European Journal of Neuroscience

676   1997;9(10):2178-86.

677   56.   Sanes JN, Donoghue JP, Thangaraj V, Edelman RR, Warach S. Shared neural substrates

678   controlling hand movements in human motor cortex. Science 1995;268(5218):1775-1778.

679   57.   Lotze M, Erb M, Flor H, Huelsmann E, Godde B, Grodd W. fMRI evaluation of

680   somatotopic representation in human primary motor cortex. NeuroImage 2000;11(5 Pt

681   1):473-81.

682   58.   Blankenburg F, Ruben J, Meyer R, Schwiemann J, Villringer A. Evidence for a rostral-

683   to-caudal somatotopic organization in human primary somatosensory cortex with mirror-

684   reversal in areas 3b and 1. Cerebral Cortex 2003;13(9):987-93.

685   59.   Kurth R, Villringer K, Mackert BM, Schwiemann J, Braun J, Curio G, et al. fMRI

686   assessment of somatotopy in human Brodmann area 3b by electrical finger stimulation.

687   NeuroReport 1998;9(2):207-12.

688   60.   van Westen D, Fransson P, Olsrud J, Rosen B, Lundborg G, Larsson EM. Finger

689   somatotopy in area 3b: an fMRI-study. BMC Neuroscience 2004;5:28.

690   61.   Lepage M, Habib R, Tulving E. Hippocampal PET activations of memory encoding and

691   retrieval: the HIPER model. Hippocampus 1998;8(4):313-322.

692   62.   Rombouts SA, Barkhof F, Witter MP, Machielsen WC, Scheltens P. Anterior medial

693   temporal lobe activation during attempted retrieval of encoded visuospatial scenes: an

694   event-related fMRI study. NeuroImage 2001;14(1 Pt 1):67-76.

695   63.   Kim H. Encoding and retrieval along the long axis of the hippocampus and their

696   relationships with dorsal attention and default mode networks: the HERNET model.

697   Hippocampus 2015;25(4):500-510.

698   64.   Brunec IK, Bellana B, Ozubko JD, Man V, Robin J, Liu Z-X, et al. Multiple scales of

699   representation along the hippocampal anteroposterior axis in humans. Current Biology

700   2018;28(13):2129-2135. e6.

701   65.   Nadel L, Hoscheidt S, Ryan LR. Spatial cognition and the hippocampus: The anterior-

702   posterior axis. Journal of Cognitive Neuroscience 2013;25(1):22-28.

703   66.   Sheldon S, McAndrews MP, Pruessner J, Moscovitch M. Dissociating patterns of anterior

704   and posterior hippocampal activity and connectivity during distinct forms of category

705   fluency. Neuropsychologia 2016;90:148-158.

706   67.   Poppenk J, Evensmoen HR, Moscovitch M, Nadel L. Long-axis specialization of the

707   human hippocampus. Trends in Cognitive Sciences 2013;17(5):230-240.

708   68.   Ahmed-Leitao F, Spies G, van den Heuvel L, Seedat S. Hippocampal and amygdala

709   volumes in adults with posttraumatic stress disorder secondary to childhood abuse or

710   maltreatment: a systematic review. Psychiatry Research: Neuroimaging 2016;256:33-43.

711   69.   Chao LL, Yaffe K, Samuelson K, Neylan TC. Hippocampal volume is inversely related

712   to PTSD duration. Psychiatry Research: Neuroimaging 2014;222(3):119-123.

713   70.   Erickson KI, Miller DL, Roecklein KA. The aging hippocampus: interactions between

714   exercise, depression, and BDNF. The Neuroscientist 2012;18(1):82-97.

715   71.   Erickson KI, Voss MW, Prakash RS, Basak C, Szabo A, Chaddock L, et al. Exercise

716   training increases size of hippocampus and improves memory. Proceedings of the

717   National Academy of Sciences 2011;108(7):3017-3022.

718   72.   Fotuhi M, Do D, Jack C. Modifiable factors that alter the size of the hippocampus with

719   ageing. Nature Reviews Neurology 2012;8(4):189-202.

720   73.   Taylor WD, McQuoid DR, Payne ME, Zannas AS, MacFall JR, Steffens DC.

721   Hippocampus atrophy and the longitudinal course of late-life depression. The American

722   Journal of Geriatric Psychiatry 2014;22(12):1504-1512.

A

Repetitive tapping

Smooth6

Smooth10

Sequence learning

Smooth6

Smooth10

B

Repetitive tapping

Sequence learning

Repetitive tapping

Sequence learning

A

individual maxima t-value

group maxima t-value

individual threshold t-value

contrast estimates

B

3mm, individual analysis
4mm, individual analysis
3mm, group analysis
4mm, group analysis

Thousands mm^3

C

3mm isovoxels

4mm isovoxels

3mm isovoxels

4mm isovoxels

A

B