1

The legacy of C_4 evolution in the hydraulics of C_3 and C_4 grasses

Haoran Zhou[1,2,3], Erol Akçay[1], Erika Edwards[2], Brent Helliker[1]

[1]Department of Biology, University of Pennsylvania, Philadelphia, PA 19104, USA

[2]Department of Ecology and Evolutionary Biology, Yale University, New Haven, CT 06511,

USA

[3]University Corporation for Atmospheric Research, Boulder, CO 80301, USA

Correspondence: Haoran Zhou

Phone: 1-215-808-7042

Email: haoran.zhou@yale.edu

Erol Akçay: *eakcay@sas.upenn.edu*

Erika Edwards: *erika.edwards@yale.edu*

Brent R. Helliker: helliker@sas.upenn.edu

Abstract

The anatomical reorganization required for optimal C_4 photosynthesis should also impact plant hydraulics. Most C_4 plants possess large bundle-sheath cells and high vein density, which should also lead to higher leaf hydraulic conductance (K_{leaf}) and capacitance. Paradoxically, the C_4 pathway reduces water demand and increases water-use-efficiency, creating a potential mis-match between supply capacity and demand in C_4 plant water relations. We use phylogenetic analyses, physiological measurements, and models to examine the reorganization of hydraulics in closely-related C_4 and C_3 grasses. Evolutionarily young C_4 lineages have higher K_{leaf}, capacitance, turgor-loss-point, and lower stomatal conductance than their C_3 relatives. In contrast, species from older C_4 lineages show decreased K_{leaf} and capacitance, indicating that over time, C_4 plants have evolved to optimize hydraulic investments while maintaining C_4 anatomical requirements. The initial "over-plumbing" of C_4 plants disrupts the positive correlation between maximal assimilation rate and K_{leaf}, decoupling a key relationship between hydraulics and photosynthesis generally observed in vascular plants.

Introduction

The evolution of C_4 photosynthesis in the grasses— and the attendant fine-tuning of both anatomical and biochemical components across changing selection landscapes[1,2,3]— likely impacted leaf hydraulics and hydraulics-photosynthesis relationships, both within the grass lineages in which C_4 evolved independently > 20 times[4], and as compared to closely-related C_3[5,6]. C_4 plants typically exhibit lower stomatal conductance (g_s) and consequently greater water-use efficiency than C_3, because the concentration of CO_2 inside bundle sheath cells permits reduced intercellular CO_2 concentrations and conservative stomatal behavior[7,8,9]. At the same time, C_4 plants require high bundle sheath to mesophyll ratios (BS:M), which are accomplished with increased vein density and bundle sheath size as compared to C_3 plants. In C_3 species, leaf hydraulic conductance (K_{leaf}) has a positive relationship with vein density[10,11,12,13]. The decreased inter-veinal distance and consequently higher vein density in C_4 species has been predicted to lead to a higher K_{leaf} than closely-related C_3 species[14,15]. Further, increased bundle sheath size was proposed to lead to a higher leaf capacitance in C_4 species[15,16,] This would lead to a potential physiological "mis-match", where the evolution of the C_4 pathway simultaneously increases a plant's hydraulic capacity while reducing its transpirational demand.

The significance of such a potential physiological mismatch depends on the potential costs and

tradeoffs associated with the building of an 'over-plumbed' leaf. If the costs are high[12,17], then

one would expect to see a reduction of K_{leaf} over evolutionary time, as continued selection works

to optimize the C_4 metabolism[5,18]. Alternatively, a maintenance of high K_{leaf} over time could

result from either a lack of strong selection to reduce K_{leaf}, or a strong evolutionary constraint

imposed by the anatomical requirements of C_4 photosynthesis. In other words, the high BS:M

ratio required for an efficient C_4 system may directly limit the ability of C_4 plants to optimize

their hydraulic architecture.

The evolution of a new photosynthetic pathway that results in multiple potential changes to the

plant hydraulic system represents the ideal platform to expand our understanding of the

relationship between photosynthesis and water transport. It is generally thought that maximum

photosynthetic rate (A_{max}) and hydraulic capacity (K_{leaf}) are tightly linked, because the ability to

transport water through leaves to the sites of evaporation at a high rate allows for the

maximization of carbon gain. Studies have documented a positive correlation between A_{max} and

K_{leaf} across many scales, from a broad phylogenetic spectrum of species spanning vascular

plants[11], to smaller clades of closely related species[13]. Grasses are largely absent from previous

efforts to examine this relationship, which is unfortunate because of the parallel venation found

in grasses and other monocots. With over 20 origins of C_4 photosynthesis with ages that span ~

30 million years, grasses also present a unique opportunity to examine the influence of C_4

evolution on A_{max}-K_{leaf} relationships. Using a broad sampling of grasses (Fig. 1), we determined

whether anatomical differences associated with C_4 evolution result in greater K_{leaf} and leaf

capacitance compared to their C_3 relatives. We then compared these properties between closely

related C_3 and C_4 clades to determine how C_4 evolution alters the predicted A_{max}-K_{leaf}

relationships. Finally, we then quantified evolutionary trends in K_{leaf}, capacitance and turgor loss

point after the evolution of C_4 within a lineage by asking whether more recent origins of C_4 are

represented by higher K_{leaf} and a greater K_{leaf}-A_{max} mismatch.

Results

Within each phylogenetic cluster, there were no clear patterns between C_3 and C_4 hydraulic traits

by conducting ANOVA tests only. C_4 grasses had higher or equivalent K_{leaf}, leaf capacitance

leaf turgor loss point, A_{max} and lower or equivalent g_s than their closest C_3 relatives (Fig. 2). The

one C_3-C_4 intermediate species, *Steinchisma decipiens*, in our analysis had K_{leaf} similar or

equivalent to C_4, but leaf capacitance, leaf turgor loss point, g_s and A_{max} equivalent to C_3 (Fig. 2).

By analyzing our data in the context of the evolutionary models (Supplementary Table S1),

however, we found clear C_3-C_4 differences in most measured traits. We first fitted evolutionary

models of Brownian motion and Ornstein-Uhlenbeck processes to the hydraulic traits based on a

reliable dated phylogenetic tree[19]. The best fitting evolutionary model to the data for K_{leaf}, leaf

turgor loss point, A_{max} and g_s was Ornstein-Uhlenbeck model, while the Brownian model is the

best-fitting model for leaf capacitance, as determined by the AICc and Akaike weights and LRT

test (Table 1, Supplementary Tables S2-S6). Higher K_{leaf}, higher A_{max}, lower leaf turgor loss

point, and lower g_s are detected C_4 species compared to C_3 (LRT test, all $p<0.01$; all $\Delta AICc<-3$).

For leaf capacitance, there is no significant difference for C_3 and C_4 species.

We also looked for evolutionary trends in hydraulic traits after the evolution of a C_4 system to

probe for an extended 'optimization' phase of C_4 evolution[3, 20]. Identifying directional trends in

continuous character evolution is difficult without fossil taxa, and it is impossible to directly

measure hydraulic traits for fossils; however, we can test for trends indirectly using extant

species. For example, if reduction in K_{leaf} is selected for subsequent to C_4 evolution we expect

older C_4 lineages to have lower K_{leaf} values than younger C_4 lineages. We extracted the

evolutionary age of C_4 origin for each of our lineages from the dated phylogeny[19]. Regressions

of evolutionary age versus hydraulic traits provide strong evidence for a long-term directional

trend in hydraulic evolution following the origin of C_4 photosynthesis (Fig. 3). K_{leaf}, leaf turgor

loss point and capacitance showed significant negative correlations with evolutionary age, while

A_{max} had a significant positive correlation. In contrast, there was no significant relationship

between g_s and evolutionary age. No evolutionary relationships were detected in C_3 species,

which indicated the correlations between evolutionary age and hydraulic traits were unique to C_4

species. We also tested for an evolutionary trend by modelling hydraulic trait evolution using a

phylogeny with branch lengths scaled to molecular substitutions/site, which provides an estimate

of differences in evolutionary rates between lineages[4]. While the second approach requires

many assumptions that are likely violated, the results also provide additional support to a

directional trend in K_{leaf} and capacitance in C_4 lineages: comparing 12 different types of models

with or without evolutionary trends (supplementary Table S7), we found K_{leaf} and leaf

capacitance were best fitted by the Brownian motion model with a significant negative trend for

C_4 (Supplementary Table S8, Table S9-13).

Fig. 1 Phylogenetic sampling of the species for measuring physiological traits and the

independent evolutionary lineages corresponding to grass lineages. The figure on the left was a

grass phylogeny adapted from GPWGII (2012), on which the tags represent the recommended

independent evolution of C_4 for comparative studies in grasses (numbers represent there are

multiple origins within a lineage). The figure on the right is the phylogeny for our species,

extracted from a dated phylogeny [19] for species sampled in our experiments. We sampled nine

independent evolution of C_4 in total.

Fig. 2 Hydraulic conductance (K_{leaf}, mmol m^{-2} s^{-1} MPa^{-1}), leaf capacitance (mmol m^{-2} MPa^{-1}), maximal stomatal conductance (g_s, mmol m^{-2} s^{-1}), maximal assimilation rate (A_{max}, µmol m^{-2} s^{-1}), and leaf turgor loss points (Turgor loss, -MPa) of closely related C_3 and C_4 species. Different colored clusters of bars show nine different origins of closely-related C_3 and C_4 species. C_3 species are colored black. Error bars indicated standard errors.

133 Table 1 Phylogenetic results of the best-fitted models and their parameters for hydraulic
134 conductance (K_{leaf}), leaf capacitance (Capacitance), stomatal conductance (g_s), and leaf turgor
135 loss point (Turgor loss) (summarizing Table S2-S6; model description: Table S1).

Property	Model	Model type	AICw	Root/Theta	
				C_3	C_4
K_{leaf}	Model 6*	OU2	0.984	2.682	4.295
Capacitance	Model 2	BM1	0.323	0.523	
g_s	Model 6*	OU2	0.980	0.183	0.102
Turgor loss	Model 6*	OU2	0.996	-1.522	-1.192
A_{max}	Model 6*	OU2	0.5292	13.66	17.34

* indicates the model fit significantly better than all the other models. Different root or theta values for C_3
and C_4 indicates that the evolutionary model with two different values of the root or theta for C_3 and C_4
species is a significantly better fit than the evolutionary model with the same root or theta.

We next explored how A_{max} and hydraulic traits are correlated across the phylogeny, and whether
this relationship is different for C_3 and C_4 lineages. The correlations between A_{max} and K_{leaf} were
different between C_3 and C_4 (Fig. 4, Table 2, Table S13). A_{max} was significantly positively
correlated with K_{leaf} for C_3, but not for C_4 (Fig. 4, Table 2, Table S13). A_{max} was weakly
positively correlated with leaf capacitance and g_s and the correlations were not significantly
different for C_3 and C_4 (Fig. 4, Table 2, Supplementary Table S21, S22). A_{max} was negatively,
but not significantly related with leaf turgor loss point in C_3 and C_4 species (Supplementary
Table S23).

Fig. 3 The regression for hydraulic conductance (K_{leaf}), leaf capacitance, leaf turgor loss point,

stomatal conductance (g_s) and maximal assimilation rate (A_{max}) vs. the evolutionary age for the

nine origins of C_4 to show the evolutionary trend within C_4 and within their closely-related C_3

species. The evolutionary age for each sampled origin is derived from the dated phylogeny[19].

Table 2 Phylogenetic Correlations between maximal assimilation rates (A_{max}) and hydraulic traits

for C_3 and C_4 species (summarizing Table S20-S23; model description: Table S19).

Parameter 1	Parameter 2	Best Model	r for C_3	r for C_4	*p* value
A_{max}	K_{leaf}	CorModel 3	0.695	0.129	0.012/0.51
A_{max}	Capacitance	CorModel 2	0.259		0.027
A_{max}	g_s	CorModel 1	0.533		0.003

	A_{max}	leaf turgor loss	CorModel 1	-0.223	0.256

Different r means the best fitted model assuming different correlations for C_3 and C_4. One r means the

best fitted model assuming similar correlations for C_3 and C_4. *p* values indicated whether the correlation

coefficients are significant.

We used our mechanism-based physiological model[32] to consider how the evolution of higher

K_{leaf} would affect the optimal g_s and photosynthesis in C_3 and C_4 plants. An increase in K_{leaf} in

the C_3 ancestor selects for higher g_s and increases the steady-state leaf water potential to a

limited extent (Fig. 5, S1). Changing K_{leaf} has a smaller effect on the photosynthesis rate of C_4

than that of C_3 (Fig. 6, Table S25), Decreasing K_{leaf} by half or doubling it changes the

photosynthesis rate of a C_4 plant by an average of -4.27% and 3.48%, respectively. In contrast,

the same shifts in K_{leaf} has average effects of -10.07% and 9.14% on the assimilation rate of a C_3

plant. The sensitivity of the assimilation rate to changes in K_{leaf} decreases with increasing CO_2

concentration and increasing water-limitation for both C_3 and C_4 plants (Table S25). These

differences in sensitivity to K_{leaf} were robust to differences in physiological properties between

C_3 and C_4 (specifically, the temperature response properties and J_{max}/V_{cmax} ratio; Table S25). The

assimilation rate of C_4 plants was still less sensitive to K_{leaf} than that of C_3 species under

different CO_2 concentration and water-limited conditions (Table S25). The physiological

modeling results indicates that C_4 species maintain lower g_s and higher leaf water potential

compared to closely related C_3 species because the CCM reduces transpirational demand. The

modeling effects of varying K_{leaf} on photosynthesis confirmed the diminished returns for high-

efficiency water transport in C_4 species mentioned above.

Fig. 4 Phylogenetic correlation for C_3 and C_4 between A_{max} and other hydraulic traits (K_{leaf}, leaf

capacitance and g_s). Different/same correlation values on the figure mean C_3 and C_4 have

significantly different/same correlations. Detailed phylogenetic correlation models and analysis

results are shown in Table 2. Dashed black line: C_3; dashed red line: C_4; solid black line: C_3 and

C_4 have the same correlation; grey lines indicate the phylogeny.

Fig. 5 The effect of changing K_{leaf} on stomatal resistance (the inverse of g_s) and leaf water

potential under VPD=1.25 kPa, ψ_s =-1 MPa and CO_2 concentration of 200 ppm for the C_3 model.

Solid black line: measured K_{leaf}; dashed black line: K_{leaf} doubled; dashed grey line: K_{leaf} reduced

by 50%.

To see if C_4 subtypes varied in hydraulic traits and their evolutionary rates or variance, we also

considered evolutionary models where we allowed each variable to have a subtype-specific value

(Supplementary Table S1).   We found no significant differences in K_{leaf}, leaf capacitance, g_s,

leaf turgor loss point and A_{max} among C_4 subtypes (all ΔAICc>0, ΔAICc obtained by AICc of

subtype models minus AICc model not considering subtypes; Supplementary Tables S14-18).

Although different decarboxylation enzymes are utilized by the three major subtypes (NADP-

ME, NAD-ME and PCK), there does not seem to be an evolutionary effect on hydraulic traits.

However, a previous study documenting PCK species from the Chloridoideae and Panicoideae

lineages with lower leaf turgor loss point[23]. Such differences were not apparent when we

compared C_4 subtypes with multiple lineages. Our current representation of different subtypes is,

however, somewhat limited. It would be advantageous to increase both lineage and species

diversity and to balance subtypes within lineages to more deeply examine C_4 subtypes.

Fig. 6 Modeling results of photosynthesis rates along with different CO_2 concentration, different

temperatures and different water limited conditions for C_3 (black lines) and for C_4 (red lines).

Solid lines: modeling results for C_3 and C_4 with measured leaf hydraulic conductance; dashed

lines: modeling results for C_3 and C_4 with twice of the regular leaf hydraulic conductance; dotted

lines: modeling results for C_3 and C_4 with half of the regular leaf hydraulic conductance. C_3 and

C_4 parameters are kept the same except for C_4 has the carbon concentration mechanism.

Discussion

The evolution of the C_4 pathway in the grasses caused a series of shifts in hydraulic properties as

compared to closely-related C_3 grasses. The anatomical requirements of C_4 initially increased

K_{leaf} and leaf capacitance, as predicted by previous studies[14,15,16]; however, K_{leaf} and leaf

capacitance appear to decline over evolutionary time, suggesting a long period of physiological

optimization after the initial assembly of a new photosynthetic system. Previous examination of

leaf hydraulic traits in grasses focused on investigating single species or were not developed

within a phylogenetic framework when comparing multiple species[21,22], and phylogenetic

studies have assumed trait evolution as simple Brownian motion[23,24]. Hydraulic traits, however,

may have evolved along different trajectories before and after the evolution of the C_4 pathway

and associated anatomical reorganization, resulting in more complicated evolutionary dynamics.

Our evolutionary models indicated C_4 grasses initially had higher K_{leaf}, leaf capacitance, turgor

loss point than corresponding C_3, and a lower stomatal conductance (g_s) than grasses consistent

with previous studies[25,26]. Decreased vein distance and increased bundle sheath size are thought

to be anatomical precursors to the evolution of C_4[27,28], and both are thought to increase K_{leaf}

and/or leaf capacitance[14,15]. Therefore, the shifts of K_{leaf} and leaf capacitance likely occurred

before, or at the initial formation of, the C_4 CCM. After the full formation of C_4, K_{leaf} and/or leaf

capacitance started to decrease, which led to higher or equivalent K_{leaf} and leaf capacitance in the

current C_3 and C_4 species (Fig. 2). Liu et al. (2019) found that K_{leaf} in C_4 grasses overlapped with

C_3 values[24]. The positive correlation between A_{max} and the evolutionary age also supports an

extended optimization phase for C_4. Previous studies have indicated that species from the oldest

C_4 lineages (*Chloridoideae* and *Andropogoneae* for example) contain the most productive crops

(Sage, 2016), while some recent C_4 lineages are not more productive than C_3 (Ripley et al., 2008;

Lundgren et al., 2016). In contrast, the significant decrease of g_s and the increase of leaf turgor

loss point occurred with the evolution of a fully operational C_4 CCM, as suggested by our

physiological models discussed below. Consistent with this prediction, in clades that possess a

range of C_3, C_3-C_4 intermediate and C_4 physiologies, the increased water use efficiency,

decreased g_s, and a broadened ecological niche are observed only in plants with a full C_4

CCM[29,30].

The evolution of C_4 significantly alters the widely-accepted A_{max}-K_{leaf} relationships existing in

vascular plants. A_{max} is limited by the efficient transport of water through leaves to replace water

loss through open stomata, which is the likely cause of a positive correlation between K_{leaf} and

A_{max} across and within plant taxa[11,13,31]. We found that A_{max} and K_{leaf} are positively correlated

in our C_3 species but not in C_4 (Fig 4). Ocheltree et al. (2016)[22] similarly found no relationship

between K_{leaf} and A_{max} in a set of nine C_4 species. We see possible explanations that are not

necessarily mutually exclusive. First, the positive relationship of A_{max} and K_{leaf} is weakened

under high K_{leaf}, possibly due to diminished returns of further increasing the efficiency of water

transport[11,31], a conclusion supported by our physiological modeling results below. As K_{leaf}

tends to be lower in grasses than in other species, it is possible that the diminishing returns from

increasing K_{leaf} manifest at lower values in grasses, and the initial high K_{leaf} resulting from C_4

anatomy could be in the A_{max} "saturation" zone. Lastly, we see evidence here that the time-since-

C_4-evolution affects several hydraulic traits across and within lineages, and it could be that a

walk towards A_{max}–K_{leaf} optimality is slowly occurring within C_4 grass lineages in relatively

newfound ecological niches. However, the similar correlations of g_s vs. A_{max} in C_3 and C_4 and

lack of evolutionary trend in g_s indicated the evolutionary processes of g_s might be already near

the optimal condition or stabilized quickly. Other hydraulic traits of leaf capacitance and leaf

turgor loss point do not seem to contribute to the A_{max} directly because of weak correlations.

We identified the mode and direction of evolution for hydraulic traits in C_3 and C_4 lineages and

found evidence that different traits followed different evolutionary processes. Hydraulic

conductance and leaf capacitance could therefore evolve with directions in a step-wise fashion

due to anatomical constraints, but g_s and leaf turgor loss point might have a more quick process

of readjustments, which allows them to stabilize soon. This suggests that there could be greater

diversification of K_{leaf} and leaf capacitance in the existing C_4 species and maybe in the future.

Also, these rearrangements of hydraulic properties interacted with each other throughout the

evolutionary trajectory. For example, increased K_{leaf} and leaf capacitance would lead to an

increased water transport efficiency, which enabled greater g_s of the C_4 ancestor (either a C_3

grass or a C_3-C_4 intermediate), but the formation of the full C_4 CCM enables a decrease of g_s.

Therefore, observed g_s in C_4 grasses reflects a balance of these two contrasting physiologies

playing out in a given ecological and phenological background, which may explain why although

C_4 g_s was lower than the C_3, the difference was not large. This line of reasoning might also

explain the inconsistent observations of g_s comparisons between C_3 and C_4. Most previous

studies found that C_4 grasses had lower g_s than C_3 grasses in both closely related and unrelated

species[25,33], yet Taylor et al. (2014) found that C_4 grasses maintained a higher or equivalent g_s

to closely-related C_3 grasses[34]. Likewise, artificial selection or genetic engineering might have

more success in adjusting these hydraulic traits in advance. Consciously selecting or

manipulating narrower xylem, decreasing the expression of aquaporins, or other mechanisms of

decreasing leaf conductance while maintain high bundle sheath to mesophyll ratio, together with

CCM may increase the water use efficiency of C_4 species further. Our phylogenetic analyses can

thus inform both the evolutionary history of C_4 plants and future efforts to modify C_4 crops.

By capitalizing on the multiple origins of C_4 photosynthesis in grasses, we have shown that the

vascular organization that is a hallmark of C_4 plants also impacts leaf hydraulics, and disrupts the

established link between hydraulic and photosynthetic capacity demonstrated in C_3 plants. C_4

grasses are "overplumbed" relative to their C_3 counterparts, suggesting that the costs associated

with the production of an extensive leaf vasculature require re-evaluation in plants with C_4

photosynthetic systems. The gradual decline in K_{leaf} in C_4 lineages over millions of years also

requires an explanation. The C_4-K_{leaf} conundrum provides an opportunity to examine what we

mean by "evolutionary constraint" and highlights the very dynamic nature of evolutionary trade-

offs and functional optimization. First, we assume that the costs of building and maintaining a

high K_{leaf} are still significant in C_4 plants[12,35,36,37,38]. The most efficient way to reduce K_{leaf} costs

would be to reduce venation density, as veins come with high construction costs[12,17], and also

reduce the leaf area that is available for carbon fixation. Yet the anatomical requirements of the

C_4 system preclude this option: reducing vein density would result in a highly inefficient C_4

system[15], which would negatively impact the plant's carbon budget, presumably to a much

greater extent than the cost of an overbuilt venation system. As vein construction is a primary

contribution to the cost of a high K_{leaf}, and high vein densities are now linked to a new function

(C_4 carbon fixation), the cost-benefit calculations in optimizing K_{leaf} have shifted, and the

tradeoff is in favor of overplumbing in order to maintain a highly efficient new carbon fixation

system. In evolutionary vocabulary, what emerges is a new constraint – and in this example, it is

clear that the emergence of a new constraint to organismal evolution is simply due to a shift in

the tradeoffs associated with characters that influence multiple aspects of organismal function. In

other words, we assume a low vein density is a phenotype that is still developmentally

achievable for C_4 grasses; what has prevented its emergence is the shift in functional costs

associated with reduced vein densities.

And yet, we documented a gradual reduction in K_{leaf} over time, which we presume was

accomplished via changes in other factors that influence leaf hydraulic capacity– perhaps by

changing xylem conduit diameters, shifts in extra-xylary mesophyll conductance, decreased

expression of aquaporins, and reorganization of internal air spaces[6,12,37,39,40]. It is possible that

these changes resulted from a continued and direct selection pressure to reduce investment in an

underutilized hydraulic system. An alternative explanation is that all of the traits that influence

K_{leaf} also play important roles in other aspects of leaf function – and the emergent of a new

constraint (a high vein density to maintain C_4 function) has **released** still other constraints on

other traits so that they may be optimized for their other functions. A striking pattern in our data

is that older C_4 lineages have achieved both lower K_{leaf} and higher A_{max} – suggesting that they

are continuing to optimize their photosynthetic capacity, long after the initial origin of C_4. We

suspect that the slow evolutionary decline in K_{leaf} is due in large part to the optimization of traits

to increase A_{max} at the expense of K_{leaf}, which is possible only because hydraulic capacity was

already "buffered" by the vein density requirements of C_4 – allowing for continued reductions of

K_{leaf} at no functional cost. Increased suberization of bundle sheath cells is one example of a

potential release of constraint[22]: it allows C_4 plants to gain higher A_{max} through reducing bundle

sheath leakiness, but it likely simultaneously reduces water flow from veins out into the

mesophyll. Since C_4 plants are already operating in hydraulic excess, bundle sheath suberization

may be optimized for C_4 function without any negative repercussions for plant water relations.

This hypothesis could also explain the opposing trends in A_{max} and K_{leaf} when viewed as a

function of evolutionary age. The examination of C_4 evolution in grasses provides an exciting

system to study the evolutionary dynamics of constraints highlighted by the interplay between

photosynthesis and plant hydraulics.

Methods

Plant material

We collected seeds of 39 closely related C_3 (9 species), C_4 species (29 species), representing

three C_4 subtypes, nine C_4 origins, and one C_3-C_4 intermediate species. The selected C_3 and C_4

species fall into nine identified C_4 lineages belong to the 11 recommended grass lineages for C_3

and C_4 study (11 out of the total 24 grass lineages have clear C_3 sister species and are

recommended for comparative studies in GPWGII, 2012[4]): *Aristida*, *Stipagrostis*,

Chloridoideae (*Eragrostideae*), *Eriachne*, *Tristachyideae*, Arthropogoninae, *Otachyrinae*

(Anthaenantia), *Panicinae*, *Melinidinae*, and *Cenchrinae* (Fig. 1). In 2015, seeds were surface

sterilized before germination and the seedlings were transferred to 6 inch pots with the soil of

Fafard #52 (Sungro, Ajawam, MA). Six replicates of each species were randomized in the

greenhouse of the University of Pennsylvania supplemented with artificial lighting. The plants

were watered twice daily. Daytime/night temperature was controlled at 23.9-29.4/18.3-23.8 $^{\circ}$C;

relative humidity was around 50-70%. Plants were fertilized once per week with 300 ppm

Nitrogen solution (Jacks Fertilizer; JR Peters, Allentown, PA) and 0.5 tsp of 18-6-8 slow release

Nutricote Total (Arysta LifeScience America Inc, NY) per pot was applied when plants were

potted into 6 inch pots. To maintain optimal plant growth a 15-5-15 cal-mg fertilizer was used

every third week. All measurements were performed on the most-recent fully expanded leaves.

Hydraulic traits

Leaf hydraulic conductance (K_{leaf}) was measured using the evaporative flux method[41], with

some adjustments to maintain stability of the evaporative environment to which the leaf was

exposed (Supplementary Methods). The evening before measurements, potted plants were

brought to the laboratory, watered, and then covered by black plastic bags filled with wet paper

towels to rehydrate overnight. For the leaf gasket, a 1 cm diameter, ~ 1 cm long solid silicone

rubber cylinder was cut nearly in two, leaving a hinge on one end. The cylinder was placed

around the leaf blade near the ligule and glued shut with superglue[42]. The leaf was cut from the

plant with a razor blade while submerged in a 15 mmol L^{-1} KCl solution; the rubber gasket was

then attached to tubing filled with the same KCl solution. The other end of the tubing was inside

a graduated cylinder that sat on a digital balance (Mettler-Toledo). The leaf was then placed

inside a custom, environmentally controlled cuvette that allowed for the measurement of entire

grass blades. Throughout measurements, cuvette temperature was controlled at 25 $^{\circ}$C and the

humidity was 55-65% (VPD range of 1.1-1.4 kPa) across measurements, but remained constant

during a particular measurement. Photosynthetically active radiation in the system is 1000 μmol

m^{-2} s^{-1}. Flow from the balance was monitored for 45 m to 1h until the flow rates reach steady

state. After the measurements, the leaf was detached and was put into a plastic bag to equilibrate

for 20 minutes to measure the leaf water potential (Model 1000, PMS Instrument, USA). K_{leaf}

values were further standardized to 25 $^{\circ}$C and leaf area to make the K_{leaf} comparable among

studies and across species. Data indicating a sudden change of flow and whose leaf water

potential was an obvious outlier were deleted.

We measured pressure-volume (PV) curves for six leaves per species using the bench-drying

method[43,44]. A leaf was cut directly from the same plants rehydrated in the lab (as described

above) using a razor blade and leaf water potential was measured immediately. Then, the leaf

weight was recorded. The leaf was initially allowed to dry on the bench for 2-minute intervals

and put into a ziplock bag and under darkness for 10-minute equilibration before measuring the

leaf water potential and leaf weight again. Then, the waiting intervals could be adjusted based on

the decrease of the leaf water potential (from 2 minutes-1h). Ideally, a decreasing gradient of -

0.2MPa for leaf water potential was obtained for the curves, until the leaf weight reached a

steady state. At the end of the experiment, leaves were dried in the oven at 70°C for 48h to obtain

the dry weight. The PV curves were used in curve fitting to obtain leaf capacitance, and leaf

turgor loss point using an excel program from Sack and Pasquet-Kok (2010)[44].

Maximal assimilation rate (A_{max}) and stomatal conductance (g_s) were measured under saturated

light intensity. A_{max} and g_s were obtained using a standard 2 x 3 cm^2 leaf chamber with a red/blue

LED light source of LI-6400XT (LI-COR Inc., Lincoln, NE, USA). Light curves were measured

with light intensities of 2000, 1500, 1200, 1000, 800, 500, 300, 200, 150, 100, 75, 50, 20, 0 μmol

m^{-2} s^{-1} under CO_2 of 400 ppm. Then, A_{max} was estimated from the light curve[45,46]. All the

measurements were made under the temperature of 25°C and the leaf temperature to air vapor

pressure deficit was controlled around 2kPa. g_s at the saturated light intensity of 2000 μmol m^{-2}

s^{-1} was recorded for each plant. The cuvette opening was covered by Fun-Tak to avoid and

correct for the leakiness.

Phylogenetic analysis

Phylogenetic analysis for C_3 and C_4. We pruned the dated phylogeny from a published grass

phylogeny to include only the species in our physiological experiments[19](Fig. 1). Using the

dated phylogeny, for each of the hydraulic traits, we fitted evolutionary models to test which

evolutionary model best explains observed distribution of traits along the phylogeny and how

these models differ between C_3 and C_4 (Table S1). We fitted evolutionary models belonging

Brownian Motion model and Ornstein-Uhlenbeck Model using the package "mvMORPH" in

R[47]. To determine the best fitted evolutionary model, we compared two criteria, the small-

sample-size corrected version of Akaike information criterion (AICc, the lower AICc, the better

fit) and Akaike weights (AICw, the higher AICw, the better fit)[48,49,50]. The evolutionary models

have nested variants (Models 1-4; Models 5-6), varying in whether C_3 and C_4 species had the

same or different fluctuation rates, root states for Brownian motion model and optima for

Ornstein-Uhlenbeck model. We used likelihood-ratio test (LRT) to verify whether a specific

model variant performs significantly better. The AICc, AICw and LRT allowed us to test

evolutionary hypotheses, for instance, if the model in which C_3 and C_4 have different root states

fit significantly better than model in which C_3 and C_4 have the same root states, it means there is

a shift of physiological trait along with the formation of C_4. To examine the further evolution of

hydraulic traits after a full C_4 evolved, we extracted the evolutionary ages for each represented

C_4 origin from the dated phylogenetic trees. Then, we regressed the hydraulic traits with

evolutionary age. A significant negative correlation between evolutionary age and hydraulic trait

will indicate a further decreasing evolutionary direction after C_4 evolved. We also performed an

additional analysis to test the original states and further direction together. We extracted

molecular phylogeny for all the species from Edwards, GPWG II (2012)[4]. Except for the six

evolutionary models mentioned above, the molecular phylogeny allows us to fit for additional six

Brownian motion models with trend (Supplementary Table S7). Likewise, if Brownian motion

model with trend fits the phylogenetic patterns better than Brownian motion model without trend

it means there is an evolutionary trend, and a significant LRT test for a two-trend model suggests

that C_3 and C_4 lineages differ in the speed or direction of hydraulic evolution. We also mapped

the traits on the phylogeny for potential further references (Fig. S2-S5).

To further test whether there are significant differences among C_4 subtypes, evolutionary models

with subtypes (Table S1) were used to fit the data. We again used AICc, AICw and LRT

methods to find the best model variants: whether there are significant differences for hydraulic

shifts and evolutionary trends among three different subtypes. For the leaf capacitance analysis,

Dichanthelium clandestinum is deleted as it is an obvious outlier.

Phylogenetic analysis for correlations among traits. Multivariate analysis in "mvMORPH"

was used to estimate the correlations between A_{max} and each of the hydraulic traits and to test the

hypotheses that whether such correlations are different between C_3 and C_4. The process of

brownian motion with different root for C_3 and C_4 was used for K_{leaf}, g_s and leaf turgor loss and

brownian motion with the same root was used for leaf capacitance. Since the Ornstein-

Uhlenbeck process is difficult to take the root state difference into consideration, here we used

Brownian motion assumptions as approximation for leaf turgor loss. Seven different correlation

models are fitted (Table S19). We used LRT for the seven correlation models to test whether the

correlation of the two traits is significantly different from 0 and whether the correlation of two

traits is significantly different between C_3 and C_4. Such correlation analysis is similar to PGLS

considering C_3 and C_4, but with more varieties on the setting of variance and covariance matrix.

Physiological Modeling

Furthermore, we used physiological models that couples the photosynthesis systems and

hydraulic systems to predict the effect of changing K_{leaf} on assimilation rate[32]. The change of

K_{leaf} was assumed to change the plant hydraulic conductance (K_{plant}) proportionally in the

modeling process. We double or reduce by half K_{leaf} relative to the original value to predict the

effects on assimilation rates for C_3 and C_4 pathways. We assumed C_4 had the same

photosynthetic properties with C_3 species (e.g., Rubisco affinity and specificity, Supplementary

Table S24) other than the carbon concentration mechanism, which mimics the initial evolution of

C_4 and the closely-related C_3-C_4 system. We also model the additional scenarios in which C_4 had

different photosynthetic properties to support the above condition further (Supplementary Table

S25).

Acknowledgements

HZ and this research is supported by the NOAA Climate and Global Change Postdoctoral

Fellowship Program, administered by UCAR's Cooperative Programs for the Advancement of

Earth System Science (CPAESS) under award #NA18NWS4620043B and is also supported by

the Dissertation Completion Fellowship provided by the Graduate Division of School of Arts and

Sciences, University of Pennsylvania. BH is supported by NSF-IOS award 1856587.

Data availability

The data that support the findings of this study are available from the corresponding author upon
request.
470
471 **Code availability**
472 All source code is available upon request.
473
474
475 **References**
476

1. Sage, R.F., 2017. A portrait of the C4 photosynthetic family on the 50th anniversary of its
discovery: species number, evolutionary lineages, and Hall of Fame. Journal of
experimental botany, 68(2), pp.e11-e28.
2. Ehleringer JR, Monson RK (1993). Evolutionary and ecological aspects of
photosynthesis pathway variation. Annu Rev Ecol Syst 24: 411-439.
3. Edwards, E. J. (2019). Evolutionary trajectories, accessibility and other metaphors: the
case of C_4 and CAM photosynthesis. *New Phytol. 223*(4), 1742-1755.
4. Grass Phylogeny Working Group II. (2012). New grass phylogeny resolves deep
evolutionary relationships and discovers C4 origins. New Phytol 193: 304-312.
5. Christin PA, Osborne CP (2014) Tansley Review. The evolutionary ecology of
C_4 photosynthesis. New Phytol 204: 765-781.
6. Kocacinar F, Sage RF (2003) Photosynthetic pathway alters xylem structure and
hydraulic function in herbaceous plants. Plant, Cell & Environment 26(12): 2015-2026.
7. Pearcy RW, Ehleringer J (1984) Comparative ecophysiology of C_3 and C_4 plants. Plant,
Cell & Environment 7(1): 1-13.
8. Huxman TE, Monson RK (2003) Stomatal responses of C_3, C_3-C_4 and C_4 Flaveria species
to light and intercellular CO_2 concentration: implications for the evolution of stomatal
behaviour. Plant, Cell & Environment 26(2): 313-322.
9. Way DA, Katul GG, Manzoni S, Vico G (2014) Increasing water use efficiency along the
C_3 to C_4 evolutionary pathway: a stomatal optimization perspective. Journal of
Experimental Botany 65(13): 3683-3693.
10. Sack L, Frole K (2006) Leaf structural diversity is related to hydraulic capacity in
tropical rain forest trees. Ecology 87: 483–491.
11. Brodribb TJ, Feild TS, Jordan GJ (2007) Leaf maximum photosynthetic rate and venation
are linked by hydraulics. Plant Physiol 144: 1890–1898.
12. McKown AD, Cochard H, Sack L (2010) Decoding leaf hydraulics with a spatially
explicit model: principles of venation architecture and implications for its evolution. Am
Nat 175: 447–460.
13. Scoffoni C, Chatelet DS, Pasquet-kok J, Rawls M, Donoghue MJ, Edwards EJ, Sack L
(2016) Hydraulic basis for the evolution of photosynthetic productivity. Nature
plants 2:16072.
14. Osborne CP, Sack L (2012). Evolution of C_4 plants: a new hypothesis for an interaction of
CO_2 and water relations mediated by plant hydraulics. Philos Trans R Soc Lond B Biol
Sci 367: 583-600.

15. Griffiths H, Weller G, Toy L, Dennis RJ (2012) You're so vein: bundle sheath physiology, phylogeny and evolution in C_3 and C_4 plants. Plant Cell Environ 36: 249–261.

16. Sage RF (2001) Environmental and evolutionary preconditions for the origin and diversification of the C_4 photosynthetic syndrome. Plant Biology 3: 202-213.

17. McCulloh, KA, Sperry, JS, Adler, FR (2003) Water transport in plants obeys Murray's law. Nature 421: 939-942.

18. Sage RF, Sage TL, Kocacinar F (2012) Photorespiration and the evolution of C_4 photosynthesis. Annual review of plant biology 63: 19-47.

19. Spriggs EL, Christin PA and Edwards EJ (2014) C_4 photosynthesis promoted species diversification during the Miocene grassland expansion. Plos one 9(5): p.e97722.

20. Heyduk ., Moreno-Villena JJ, Gilman IS, Christin PA, and Edwards EJ (2019). The genetics of convergent evolution: insights from plant photosynthesis. Nature Review Genetics, 20, 485-493.

21. Martre P, Cochard H, Durand JL (2001) Hydraulic architecture and water flow in growing grass tillers (Festuca arundinacea Schreb.). Plant, Cell and Environment 24: 65–76.

22. Ocheltree TW, Nippert JB, Prasad PV (2016) A safety vs efficiency trade-off identified in the hydraulic pathway of grass leaves is decoupled from photosynthesis, stomatal conductance and precipitation. New Phytologist 210(1): 97-107.

23. Liu H, Osborne CP (2015) Water relations traits of C_4 grasses depend on phylogenetic lineage, photosynthetic pathway, and habitat water availability. Journal of Experimental Botany 66, 761–773.

24. Liu H, Taylor SH, Xu Q, Lin Y, Hou H, Wu G, Ye, Q (2019) Life history is a key factor explaining functional trait diversity among subtropical grasses, and its influence differs between C_3 and C_4 species. Journal of experimental botany 70(5): 1567-1580.

25. Taylor SH, Hulme SP, Rees M, Ripley BS, Woodward FI, Osborne CP (2010). Ecophysiological traits in C_3 and C_4 grasses: a phylogenetically controlled screening experiment New Phytol 185: 780-791

26. Taylor SH, Franks PJ, Hulme SP, Spriggs E, Christin PA, Edwards EJ, Woodward FI, Osborne CP (2012) Photosynthetic pathway and ecological adaptation explain stomatal trait diversity amongst grasses. New Phytol 193: 387-396.

27. Sage RF (2004) The evolution of C_4 photosynthesis. New Phytol 161(2):341-370.

28. Christin PA, Osborne CP, Chatelet DS, Columbus JT, Besnard G, Hodkinson TR, Garrison LM, Vorontsova MS, Edwards EJ (2013) Anatomical enablers and the evolution of C_4 photosynthesis in grasses. Proc Natl Acad Sci USA 110: 1381-1386.

29. Lundgren MR, Besnard G, Ripley BS, Lehmann CE, Chatelet DS, Kynast RG, Namaganda M, Vorontsova MS, Hall RC, Elia J, Osborne CP (2015) Photosynthetic innovation broadens the niche within a single species. Ecology Letters 18(10): 1021-1029.

30. Sage RF, Monson RK, Ehleringer JR, Adachi S, Pearcy RW (2018) Some like it hot: the physiological ecology of C4 plant evolution. Oecologia 187(4):941-966.

31. Brodribb TJ, Holbrook NM, Zwieniecki MA, Palma B (2005) Leaf hydraulic capacity in ferns, conifers and angiosperms: impacts on photosynthetic maxima. New Phytol 165: 839–846

32. Zhou H, Helliker BR, Huber M, Dicks A, Akçay E (2018) C_4 photosynthesis and climate through the lens of optimality. Proceedings of the National Academy of Sciences 115:

12057-12062.

33. Ripley BS, Cunniff J, Osborne CP (2013) Photosynthetic acclimation and resource use by
the C_3 and C_4 subspecies of *Alloteropsis semialata* in low CO_2 atmospheres. Glob
Change Biol 19: 900–910.

34. Taylor SH, Ripley BS, Martin T, De-Wet L-A, Woodward FI, Osborne CP (2014).
Physiological advantages of C_4 grasses in the field: a comparative experiment
demonstrating the importance of drought. Glob Change Biol 20: 1922-2003.

35. Tyree MT, Sperry JS (1989) Vulnerability of xylem to cavitation and embolism. Annu
Rev Plant Physiol Plant Mol Biol 40(1):19–36.

36. Niinemets Ü, Portsmuth A, Tobias M (2007) Leaf shape and venation pattern alter the
support investments within leaf lamina in temperate species: a neglected source of leaf
physiological differentiation? Functional Ecology 21: 28-40.

37. Scoffoni C, Rawls M, McKown A, Cochard H, Sack L (2011) Decline of leaf hydraulic
conductance with dehydration: relationship to leaf size and venation architecture. Plant
Physiology 156(2): 832-843.

38. Wolf A, Anderegg WR, Pacala SW (2016) Optimal stomatal behavior with competition
for water and risk of hydraulic impairment. Proceedings of the National Academy of
Sciences 113(46): E7222-E7230.

39. Scoffoni C, Albuquerque C, Brodersen CR, Townes SV, John GP, Bartlett MK, Buckley
TN, McElrone AJ, Sack L (2017) Outside-xylem vulnerability, not xylem embolism,
controls leaf hydraulic decline during dehydration. Plant Physiology 173: 1197-1210.

40. Pathare VS, Sonawane BV, Koteyeva N, Cousins AB (2020) C_4 grasses adapted to low
precipitation habitats show traits related to greater mesophyll conductance and lower leaf
hydraulic conductance. Plant Cell & Environment doi: 10.1111/pce.13807.

41. Sack L, Scoffoni C (2012) Measurement of Leaf Hydraulic Conductance and Stomatal
Conductance and Their Responses to Irradiance and Dehydration Using the Evaporative
Flux Method (EFM). J Vis Exp (70), e4179, doi:10.3791/4179.

42. Ocheltree T, Gleason S, Cao KF, Jiang GF (2020) Loss and recovery of leaf hydraulic
conductance: Root pressure, embolism, and extra-xylary resistance. Journal of Plant
Hydraulics 7

43. Tyree MT, Hammel HT (1972) The measurement of the turgor pressure and the water
relations of plants by the pressure-bomb technique. Journal of Experimental
Botany 23(1): 267-282.

44. Sack L, Pasquet-Kok J and PrometheusWiki contributors. Leaf pressure-volume curve
parameters. PrometheusWiki. May 20, 2010, 17:08 UTC. Available at: /tiki-
pagehistory.php?page=Leaf pressure-volume curve parameters&preview=16.

45. Marshall B, Biscoe PV (1980) A model for C_3 leaves describing the dependence of net
photosynthesis on irradiance. Journal of Experimental Botany 31:29-39.

46. Thornley JHM (1998) Dynamic model of leaf photosynthesis with acclimation to light
and nitrogen. Annals of Botany 81(3): 421-430.

47. Clavel J, Escarguel G, Merceron G (2015) mvMORPH: an R package for fitting
multivariate evolutionary models to morphometric data. Methods in Ecology and
Evolution, 6(11):1311-1319.<doi:10.1111/2041-210X.12420>

48. Akaike H (1974) A new look at the statistical model identification. In Selected Papers of
Hirotugu Akaike (pp. 215-222). Springer, New York, NY.

49. Cavanaugh JE (1997) Unifying the derivations for the Akaike and corrected Akaike

information criteria. Statistics & Probability Letters 33(2): 201-208.
50. Burnham KP and Anderson DR (1998) Practical use of the information-theoretic
approach. In Model Selection and Inference (pp. 75-117). Springer, New York, NY.
606