PREFERENTIAL SYNTHESIS OF VERY LONG CHAIN POLYUNSATURATED FATTY ACIDS IN *EUTREPTIELLA* SP. (EUGELNOZOA) REVEALED BY CHROMATOGRAPHIC AND TRANSCRIPTOMIC ANALYSES[1]

Rita C. Kuo[2]

Department of Marine Sciences, University of Connecticut, Groton, CT 06340, USA

Huan Zhang

Department of Marine Sciences, University of Connecticut, Groton, CT 06340, USA

James D. Stuart

Department of Chemistry, University of Connecticut, Storrs, CT 06269, USA

Anthony A. Provatas

Center of Environmental Sciences and Engineering, University of Connecticut, Storrs, CT 06269, USA

Linda Hannick

SAIC-Frederick, Inc., Frederick National Laboratory for Cancer Research, Rockville, MD 20852, USA

Senjie Lin[3]

Department of Marine Sciences, University of Connecticut, Groton, CT 06340, USA

[2]Current address: Joint Genome Institute, Lawrence Berkeley National Laboratory, Walnut Creek, CA 94598, USA. The work is part of her dissertation.

[3]Address for correspondence: Phone, 860-405-9168; Email, senjie.lin@uconn.edu

Running Title: Fatty acid and lipid synthesis in alga *Eutreptiella*

Abstract

Algal lipids are a fundamental energy source in the ecosystem and material for biofuel production, but their production and regulation remains unexplored in many species including the common coastal phytoplankton *Eutreptiella* spp. Using gas chromatography-tandem mass spectrometry (GC/MS/MS), we discovered 24 types of fatty acids (FAs) in *Eutreptiella* sp. with unsaturated C24 FAs and a relatively high proportion of long chain unsaturated FAs. The abundances of C16, C18 and saturated FAs decreased when phosphate in the culture medium was depleted. Among the 24 FAs, docosahexaenoic acid (22:6) and eicosapentaenoic acid (20:5) were the most abundant, suggesting that *Eutreptiella* sp. preferentially invests in the synthesis of very long chain polyunsaturated fatty acids (VLCPFA), making this alga nutritionally valuable to grazers. Further transcriptomic analysis revealed that *Eutreptiella* sp. likely synthesizes VLCPFA via Δ8 pathway and uses type I and II fatty acid synthases. Using RT-qPCR, we found that some of the lipid production genes, such as β –ketoacyl-ACP reductase, fatty acid desaturase, acetyl-CoA carboxylase, acyl carrier protein, Δ8 desaturase, and Acyl-ACP thioesterase, were more actively expressed during light period. In addition, were more highly expressed in the high lipid-producing light period, suggesting a linkage between photosynthesis and lipid production.

Keywords: *Eutreptiella,* fatty acid biosynthesis, lipid biosynthesis, unsaturated fatty acids

Introduction

Eutreptiella is a genus of the euglenoid algae, which have excellent nutritive value (*i.e.* high vitamin and lipid content, Takeyama et al., 1996; Yamane et al., 2001) and are rather abundant in some marine ecosystems (Henriksen et al. 2002). Some *Eutreptiella* spp. can seasonally be a dominant group of phytoplankton (Álvarez-Góngora and Herrera-Silveira 2006, Bates and Strain 2006, Olli et al. 1996, Rodríguez-Graña et al. 2008, Seong et al. 2006), forming blooms in nutrient-rich coastal or brackish waters (Anderson et al. 2000, Lindholm 1993, Olli et al. 1996, Stonik & Selina 200, Stonik 2007). In some areas, *Eutreptiella braarudii* alone can make up to 46% of the phytoplankton population (Stonik, 2007). As primary producers, these algae are important in energy flow and nutrient cycling in the coastal marine ecosystem. Meanwhile, as solar energy converters, these algae are potentially candidates of biofuel species.

Lipid content is one important indicator of quality of an alga as prey and as source of biofuel. In general, lipids are essential for structural constituents of cellular membranes in all organisms and protection of delicate internal organs and hormones in animals (Singh 2002). In the marine ecosystem, fatty acids are a fundamental energy source for growth and cell membrane fluidity (especially in cold water) in animals such as zooplankton and fish (*e.g.* Arzel et al. 1994, Cossins et al. 1977, Klein Breteler 2004, Rainuzzo et al. 1997, Tang and Taal 2005), which consume phytoplankton as food. To gain better understanding on the energy flow in marine food chains, it is important to study fatty acid and lipid biosynthesis pathways in dominant phytoplankton (prey) species. As the type and abundance of lipids in the cells also determine the potential of algae as source of biofuel, understanding the regulatory pathways of lipid production in algae is also important for evaluating or genetically enhancing algal species for biofuel production.

In land plants, photosynthesis produces small precursors for lipid production in

chloroplasts. The small precursors are later converted into long chain fatty acids via two enzyme

systems, acetyl-CoA carboxylase (ACCase) and fatty acid synthase (FAS). ACCase is essential

for lipid metabolism as it catalyzes the first step of fatty acid synthesis and is found in both

bacteria and eukaryotes (Cronan and Waldrop 2002, Harwood 1988). FAS consists of 6 enzyme

activities and an acyl-carrier protein (CA) to catalyze a 2 carbon elongation process (Bloch and

Vance 1977, Nelson and Cox 2008). It produces palmitate and stearate and these fatty acids are

subject to elongation, desaturation or further modification by elongase or desaturase in

chloroplasts (Gunstone et al. 2007). There are two major types of fatty acid synthases. Fatty acid

synthase I (FASI), found in vertebrates and fungi, consists of a multi-enzyme complex contained

in a single polypeptide chain (Nelson and Cox 2008, Schweizer and Hofmann 2004). In contrast,

fatty acid synthase II (FASII), found in plants and bacteria, contains discrete enzymes (Nelson

and Cox 2008, White et al. 2005). Among euglenoid algae, *Euglena gracilis* composes both

FASI and plastid FASII for *de novo* fatty acid synthesis (Delo et al. 1971, Worsham et al. 1993).

In addition to FASII, Goldberg and Bloch (1972) reported another ACP-dependent fatty acid

synthase (FASIII) that elongates acyl-CoA derivatives from C10 to C18 to longer chain ACP

thioesters in plastids.

Similar to land plants, microalgae initiate fatty acid synthesis in chloroplasts (Guschina

and Hardwood, 2006) and lipid biosynthesis is probably linked directly to photosynthesis.

Studies have shown that the activities of ACCase are related to the light reactions of

photosynthesis, as NADPH and ATP produced in the photochemical reaction are required for the

synthesis of palmitic acid (Sasaki et al. 1997). Furthermore, fatty acid synthesis in photosynthetic

organisms obviously relies on carbon fixation for carbon precursors for fatty acid synthesis (Bao

et al. 2000). The synthesized fatty acids are in turn used for triacylglycerol (TAG) synthesis

(Thelen and Ohlrogge 2002). In addition, dihydroxyacetone phosphate, the precursor for glycerol

3-phosphate needed in the synthesis of lipid TAG, can be produced from Calvin-Benson Cycle

(Nelson and Cox 2008).

It is well understood that lipid production in microalgae is influenced by nutrient and

other environmental conditions (e.g. Gouveia and Oliveira 2009, Griffiths and Harrison 2009, Hu

et al. 2008, Illman et al. 2000, Liu et al. 2008). Nutrient limitation can enhance the production of

lipid in algal cells. For example, the lipid content of some *Chlorella* species (*e.g. C. emersonil*

and C. pyrenoldosa) can increase to more than 60% of the dry weight under nitrogen-deprived

conditions (Griffiths and Harrison 2009, Illman et al. 2000). However, our knowledge of the

genetic regulatory mechanisms of lipid biosynthesis in marine microalgae is limited and

fragmentary (Guchina and Harwood 2006, Hu et al., 2008). Current understanding of lipid

biosynthesis pathways is mainly based on land plants.

In this study, we used 454 high-throughput sequencing to study transcriptomic profiles

and gas chromatography-tandem mass spectrometry (GC/MS/MS) to identify fatty acids

produced under phosphate-depleted and phosphate-replete conditions, and conducted reverse-

transcription qPCR (RT-PCR) to quantify expression of several key enzyme coding genes to

investigate molecular mechanisms that regulate lipid biosynthesis in *Eutreptiella* sp.

Materials and Methods

Culture preparation and RNA isolation. Two sets of cultures were maintained in phosphate-

depleted medium (to produce high-lipid culture) and phosphate-replete f/2 medium (to produce

low-lipid culture); cells were harvested, total RNA and mRNA were isolated as reported

previously (Kuo et al. 2013)

Lipid measurement. Nile Red (9-diethylamino-5H-benzo-α-phenoxazine-5-one, a lipid-soluble

fluorescent probe) staining was used to measure relative abundance of neutral lipids as reported

(Lee et al., 1998, Hu *et al*., 2008). Forty μl of Nile Red solution in acetone (250 mg/l) were

added to 10 ml of algal suspension at room temperature for 10 min. A spectrophotometer

(HITACHI, Tokyo, Japan) was then used to measure the fluorescence with excitation at the

wavelength of 490 nm and emission in the wavelength band of 580 to 590 nm, as reported in

Kuo and Lin (2012).

Lipid extraction and fatty acid identification by gas chromatography-tandem mass

spectrometry (GC/MS/MS). A gravimetric method was applied to determine actual lipid contents

by using chloroform-methanol method according to Bligh and Dyer (1959). In short, lipids were

extracted from freeze-dried cells with chloroform/methanol (1:2, v/v) and the residue was

extracted one more time with 1:1 chloroform/methanol. The chloroform layer was collected and

evaporated under a gentle flow of nitrogen gas. Derivatization for GC/MS/MS analysis followed

the method summarized by (Carvalho and Malcata 2005). Briefly, samples were dissolved in 1

ml of a freshly prepared mixture of dry acetyl chloride and methanol, at a ratio of 5:100 (v/v),

and kept at 100 °C under pure nitrogen for 1 h. After cooling, 1 ml of hexane wad added and

mixed by vortexing. Purification of the solution was achieved by adding 1 ml of saturated

sodium chloride solution. The prepared solution of fatty acid methyl esters was filtered using

0.45 μm Millipore filter. Analysis of the solution was performed on a Waters Quattro Micro

GC/MS/MS system equipped with a Rxi-5Sil MS column (30 m x 0.25 mm, 0.25 μm film

thickness, Restek Co., PA). Helium was used as the carrier gas with a flow rate of 1 ml/min.

Initial oven temperature was 70 °C for 1 min and the temperature gradient was 5 °C/min from

70 °C to 270 °C with no holding time. Sample was introduced in splitless injection mode.

Injector temperature was 270 °C and purged for 1 min with a purge flow of 25 ml/min. Total run

time was 41 min. The MS source temperature was set at 250 °C and the GC interface was 275 °C.

A fatty acids methyl ester mixture (FAMQ-005 FAME Reference Standard, AccuStandard Inc.,

CT, USA) was used as external standards. The mass spectrum and the corresponding retention

time of each component were exported to NIST MS Search 2.0 software to identify compounds

by means of library and standard comparisons.

454 sequencing of the SL-based transcriptomes. mRNA samples isolated as described above

from the four culture conditions were used to synthesize cDNA, which was subsequently used

for 454 sequencing as reported previously (Kuo et al., under peer review process). Briefly, a

modified random oligo named $454AT_7N_9$ (5'-

CGTATCGCCTCCCTCGCGCCATCAGTAATACGACTCACTATAGGGAGNNNNNNNNN-

3', where N is any of the 4 nucleotides) was used to synthesize the 1st strand cDNAs. The

cDNAs were then used as templates for PCR amplification of the 5'-end of the cDNAs of the

trans-spliced transcripts using ExTaq with a *Eutreptiella* spliced leader-based primer 454BEutSL,

5'-

GAGACTATGCGCCTTGCCAGCCCGCTCAGACACTTTCTGAGTGTCTATTTCTTTTCG-

3'), paired with $454AT_7$ (5'-

CGTATCGCCTCCCTCGCGCCATCAGTAATACGACTCACTATAGGGAG-3'). After PCR

we selected the amplicons from agarose gel in the size range of 300-700 bp as the template for

emulsion PCR (emPCR) using GS Titanium SV emPCR Kit. Sequencing was carried out using

GS Titanium Sequencing Kit on the GS FLX System at the Center for Applied Genetics and

Technology, University of Connecticut.

Sequence processing and annotation. Sequence reads were processed and analyzed as recently

reported (Kuo et al. 2012). In short, raw sequencing reads from all of the 4 samples were pooled

together and trimmed using CLC Genomics Workbench (CLC Bio, Aarhus, Demark). After

quality and primer trimming, sequences shorter than 150 nt were discarded. In order to filter off

454 sequencing errors and create reduced-redundancy sequence dataset, USEARCH (Edgar 2010)

was used for sequence clustering. The resulting unique transcripts were then annotated using

Blast2Go V.2.5.0 (Götz et al. 2008) against NCBI's non-redundant (nr) database using BLASTx

algorithm (Altschul et al. 1990), with a cut-off-E-value $\leq 10^{-3}$.

Reverse-transcription quantitative PCR (RT-qPCR). In order to further investigate the

expression patterns of genes potentially regulating lipid production in *Eutreptiella*, some of the

genes identified from the transcriptomic data as related to carbon fixation and lipid synthesis

were selected and further analyzed by RT-qPCR. First strand cDNA libraries were prepared by

the methods mentioned earlier. It is suggested that at least two reference genes are needed for

proper normalization of gene expression levels (Guo and Ki 2012, Vandesompele et al. 2002).

Tubulin, actin, and elongation factor 1α were selected as candidates, as they had been shown to

be good reference genes in some microalgae and plants (Guo and Ki 2012, Le Bail et al. 2008).

We used the geometric average of the expression levels of these reference genes to normalize the

measured expression levels of target genes as reported (Vandesompele et al. 2002). For each

gene investigated, two sets of primers were designed (Table 3). To prepare standards, the primer

combination that specifies longer gene fragment was used to PCR amplify each target gene from

Eutreptiella cDNA. The amplicons of the target genes were checked with gel electrophoresis to

assure the absence of primer dimmers and then purified using DNA Clean & Concentrate[TM]

column (Zymo Research, Orange, CA). The purified DNA was serially diluted to 10^2, 10^3, 10^4,

10^5, and 10^6 gene copies to generate standard curves to analyze the amplification efficiency and

primer specificity for every primer pair. qPCR was performed using the iCycler iQ[TM] real-time

PCR Detection System (Bio-Rad Laboratories, Inc., Hercules, CA) with SYBR Green supermix.

The qPCR reactions included a single denaturation cycle of 95 °C for 3 min, 40 cycles of 95 °C

for 20 sec, 58 °C for 30 sec, and 72 °C for 15 sec, followed by a melt curve analysis from 55 to

100 °C.

Results

Physiological conditions of the cultures. Two sets of cultures were grown under phosphate-

depleted (to produce high-lipid culture) and phosphate-replete (to produce low-lipid culture)

conditions. On the 11th day, the phosphate-replete cultures were in exponential growth phase

whereas phosphate-depleted cultures already reached stationary phase. When normalized to per

cell basis, the lipid contents of the cultures grown in phosphate-depleted medium were about 2-

fold higher than that of the cultures grown in phosphate-replete medium (Figure 1). This

observed difference in lipid content prompted cell harvesting from both cultures in the light and

dark periods, which yielded four samples referred to as high-lipid-light, high-lipid-dark, low-

lipid-light, and low-lipid-dark cultures.

Comparison of gene diversity among samples. The four samples were subjected to RNA

extraction, cDNA library construction, and 454 sequencing (Kuo et al. 2012). The Venn diagram

in Figure 2 summarizes the numbers of unique genes, total genes and shared genes among the 4

cDNA libraries. The analysis revealed that each library contained different numbers of total

genes as well as different unique genes that were not shared by any other libraries. Between the

two light cultures, the high-lipid cultures had more unique genes and total genes. Between the

cultures in the same lipid content categories, cultures harvested under the light condition

displayed more unique genes (i.e. light-specific) and total genes. Overall, high-lipid-light

cultures expressed the highest number of unique and total genes among the 4 samples,

suggesting that a more complex system was involved in lipid production and light reaction in

Eutreptiella sp.

Fatty acid synthesis. Based on the functional annotation, we have identified genes encoding key

enzymes involved in the lipid biosynthesis and catabolism (Table 1). Fatty acid biosynthesis

from acetyl-CoA involved in two enzymes: ACCase and FAS. Eukaryotic type ACCase (72%

identical to ACCase of *Thalassiosira pseudonana*; GenBank accession number: XP_002296083)

and type I FAS (67% identical to FASI of *Caenorhabditis elegans*; NP_492417) were found in

our dataset. Figure 3 illustrates a potential lipid biosynthesis pathway of *Eutreptiella* sp. ACCase

catalyzes carboxylation of acetyl-CoA to produce malonyl-CoA, which is catalyzed to malonyl-

ACP by malonyl-CoA transacylase. Next, long-chain saturated fatty acids are synthesized by

type I fatty acid synthase, which have 7 active sites within a single large polypeptide complex

(Nelson and Cox, 2008). The active enzyme domains include β-ketoacyl-ACP synthase (KAS),

malonyl/acetyl-CoA-ACP transferase (MAT), β –hydroxyacyl-ACP dehydratase (HAD), enoyl-

ACP reductase (EAR), β –ketoacyl-ACP reductase (KAR), and Acyl-ACP thioesterase (TE).

Synthesized palmitate (C16:0) and stearate (C18:0) via the activity of FASI are further subjected

to elongation, desaturation, or other modifications (see next section for more details). Besides

FASI, individual genes encoding β –ketoacyl-ACP reductase and Acyl-ACP thioesterase related

to type II fatty acid synthase (FASII) were also identified (Table 1). Malonyl-CoA transacylase,

which functions in the fatty acid synthesis pathway in other plants, was missing in our dataset.

Fatty acid identification and biosynthesis pathways. The stepwise desaturation and elongation

of C18 acid lead to the extension of fatty acids and synthesis of unsaturated fatty acids (Nelson

and Cox, 2008). The major fatty acids found in *Eutreptiella* sp. were the very long chain

polyunsaturated fatty acids, docosahexaenoic acid (DHA, 22:6n-3). Various desaturases and

elongases were found in our study (Table 1), including Δ5 desaturase, Δ6 desaturase, Δ8

desaturase, Δ9 desaturase, Δ12 desaturase and Δ15 desaturase. The elongases were: Δ5 elongase,

Δ6 elongase, and Δ9 elongase.

In order to link the identified genes to their expected roles in *Eutreptiella* sp., the fatty

acid (FA) profiles of *Eutreptiella* sp. were analyzed by screening C16-C24 fatty acids via

GC/MS/MS. We identified 24 fatty acids (including C16, C20, C22, and C24 fatty acids) from

the cultures of *Eutreptiella* sp. (Figure 4 and Table 2). In both of the phosphate-depleted and

phosphate-replete cultures, *Eutreptiella* sp. was relatively rich in polyunsaturated fatty acids

(PUFAs), especially very long chain polyunsaturated ω3 fatty acids (VLCPFA),

docosahexaenoic acid (DHA, 22:6) and eicosapentaenoic acid (EPA, 20:5). DHA had the highest

amount in both cultures (Figure 4), so we used it as a standard to estimate relative abundance of

other fatty acids (Table 2). Trace amount of tetracosahexaenoic acid, a C24 VLCPFA, was found.

Three saturated fatty acids, palmitic acid (16:0), stearic acid (18:0), and behenic acid (20:0),

were also identified. The abundance of saturated fatty acids was relatively low in the phosphate-

depleted cultures. The analysis also revealed the presence of C16 unsaturated fatty acids: 7-

hexadeceonoid acid (16:1), 9-hexadecenoid acid (16:1), 7,10-hexadecadienoic acid (16:2), 7, 10,

13-hecadecatrienoic acid (16:3), and 4, 7, 10, 13-hexadecatetraenoic acid (16:4) in both of the

cultures. In the phosphate-depleted cultures, the relative abundance of C16 and C18 fatty acids

was lower than the fatty acids in the phosphate-replete cultures (Figure 4 and Table 2),

suggesting that phosphate availability in the culture medium affected the lipid composition in

Eutreptiella sp.

Based on the transcriptome and GC/MS/MS results, *Eutreptiella* sp. is predicted to be

capable of producing various intermediates via ω3 and ω6 pathways, as well as producing

VLCPFA EPA and DHA by using the Δ8 alternative pathway, as reported in *Euglena* (Meyer et

al. 2003; Qi et al. 2004; Wallis and Browse 1999). Figure 5 shows the potential fatty acid

elongation pathways for *Eutreptiella* sp. as constructed from the transcriptome results. The first

desaturation for unsaturated fatty acids synthesis is catalyzed by Δ9 desaturase to introduce a

double bond into stearic acid resulting in 18:1 –ACP. Then a series of elongation and

desaturation takes place to produce the 24 types of unsaturated fatty acid mentioned before

(Figure 5). The elongation procedure is terminated when the acyl group is removed by acyl-ACP

thioesterase (TE) or oleoyl-ACP hydrolase (OAH), or when the fatty acids are transferred to the

biosynthesis of triacylglycerol.

Triacylglycerol (TAG) biosynthesis. Microalgal TAG biosynthesis is proposed to take place via

transfer of fatty acids from CoA to glycerol-3-phosphate using the direct glycerol pathway

(Ratledge 1988). However, the existing knowledge on the pathways and enzymes involved in

TAG biosynthesis in most of the microalgae (e.g. euglenophytes) are limited. In this study, four

transcripts coding for the enzymes involved in TAG biosynthesis were found in our *Eutreptiella*

sp. transcriptomic dataset. They were glycerol 3-phosphate dehydrogenase (GPD), glycerol 3-

phsphate acyltransferase (GPAT), and acylglycerophosphate acyltransferase (AGPAT) (Table 1).

For TAG biosynthesis, the precursor glycerol 3-phosphate is produced during glycolysis from

dihydroxyacetone phosphate by the action of glycerol 3-phosphate dehydrogenase.

Based on the data, a potential pathway for *Eutreptiella* sp. TAG biosynthesis is proposed

(Figure 3). The first step is the acylation of the two hydroxyl groups of glycerol3-phosphate by

two fatty acyl-CoA to yield phosphatidic acid, which is hydrolyzed by phosphatidate

phosphatase to form 1,2 diacylglycerol. Diacylglycerols are then converted to triacylglycerols by

transferring a third fatty acyl-CoA. Phosphatidic acid phosphohydrolase (PAP) and

diacylglycerol acyltransferase (DGAT) were missing in our cDNA libraries.

Gene expression levels as measured using reverse transcription quantitative PCR. We further

confirmed the differential expression patterns of several genes using reverse transcription

quantitative PCR (RT-qPCR). Nine genes involved in lipid synthesis were chosen for the

analysis, including type I fatty acid synthase (FASI), 3 different types of β –ketoacyl-ACP

reductase (KAR1, KAR2, KAR3, 12-33% identity at amino acid level), acetyl-CoA carboxylase

(ACCase), acyl carrier protein (ACP), thioesterase (TE), Δ8 desaturase, and an unclassified fatty

acid desaturase (FAD). Seven of the nine genes, KAR2, KAR3, FAD, ACCase, ACP, Δ8, and

TE had higher expression levels in the high-lipid-light cultures (Figure 6 and Figure 7). The

expression of FASI showed an opposite trend between high-lipid and low-lipid cultures. In high-

lipid cultures, FASI was expressed at a higher level in the light period, whereas in the low-lipid

cultures, FASI was expressed at a lower level in the light period (Figure 6). As for genes

involved in triacylglycerol synthesis, glycerol 3-phosphate dehydrogenase (GPD) and 1-acyl-sn-

glycero3-phosphate o-acyltransferase (AGPAT) exhibited slightly higher expression (transcript

abundance) in high-lipid-light cultures (Figure 7).

We also examined the expression of genes potentially involved in carbon fixation in

Eutreptiella sp.: phosphoenolpyruvate carboxylase (PEPCase), 2 variants of

phosphoenolpyruvate carboxykinase (PEPCK1 and PEPCK2, 54% identical at amino acid level),

pyruvate-phosphate dikinase (PPDK), and phosphoribulokinase (PRK). Three out of the five

genes (i.e. PPDK, PRK, and PEPCK1) showed positive correlation of transcript abundance with

lipid production. In particular, PEPCK1 and PRK expression levels in the high-lipid-light

cultures were over 6- and 4-fold higher than in the low-lipid-light cultures, respectively (Figure

8).

Discussion

Unique lipid and fatty acid production patterns. While the effects of depleting nitrate on lipid

production has been studied in many algae (Griffiths and Harrison 2009, Hu et al. 2008), effects

of phosphate are less documented. Our preliminary experiments showed that phosphate depletion

more strongly promotes lipid content in *Eutreptiella* sp. cells than nitrate depletion (data not

shown). The study reported here also showed that phosphate stress markedly increased

Eutreptiella lipid content. Among the cultures grown in parallel, lipid content was 2 fold as

much in the phosphate-depleted cultures as that in the phosphate-replete, f/2 medium-grown,

cultures (Figure 1). The type of main energy reserve (carbohydrates or lipids) in algae depends

on which carbon fixation product is utilized in the Calvin-Benson cycle. Among those products,

hexose phosphates are used in the synthesis of carbohydrates, while 3-phophoglycerate is used

for lipid synthesis (Raven 1974). In addition, the availability of inorganic nitrogen influences

carbon partition (Huppe and Turpin 1994, Raven 1974). The presence of inorganic nitrogen

source favors 3-phosphoglycerate and increases the flux of photosynthetically fixed carbon into

citrate cycle (Raven 1974), which is one of the sources for acetyl CoA (Nelson and Cox 2008).

Interestingly, *Eutreptiella* sp. has abundant polyunsaturated fatty acids. DHA was the

most abundant fatty acid from both the phosphate-depleted and phosphate-replete cultures.

Polyunsaturated fatty acids are valuable for cold adaptation (Wallis et al. 2002 and references

therein). The relatively high abundance of unsaturated fatty acids in *Eutreptiella* sp. may be due

to the adaptation to their native environment in Long Island Sound, where the temperature can be

below 0 °C in the winter (McCarthy 2008). All the fatty acids that we identified in this study

were also present in *Euglena gracilis* (Hulanicka et al. 1964, Korn 1964, Meyer et al. 2003),

except C24:1 and C24:6 fatty acids. Different from *E. gracilis*, *Eutreptiella* sp. has a very high

proportion of very long chain unsaturated fatty acids (C22 and longer, i.e. DHA and EPA),

indicating that this species may be a nutritional food resource for zooplankton and other

heterotrophic organisms in the marine ecosystem. Moreover, the proportions of saturated fatty

acids (i.e. C16:0, C18:0, and C20:0) and unsaturated C16 and C18 fatty acids decreased when

phosphate was depleted (Figure 4 and Table 2), suggesting that it is possible to enhance the

production of desired components of fatty acids from *Eutreptiella* sp. by manipulating nutrient

composition in the culture medium. Although it is not clear why the abundance of C16:0 and

C16: 4 fatty acids were much lower in the phosphate-depleted cultures, phosphate stress might

alter the composition of fatty acids in the algal cells, as plants and algae can respond to stress by

adjusting membrane fluidity and by releasing unsaturated fatty acids from the membrane

(Maréchal et al. 1997, Upchurch 2008).

Expressed genes linked to lipid synthesis in* Eutreptiella *sp. Fatty acids are synthesized via two

enzyme systems, ACCase and FAS (Nelson and Cox 2008). ACCase plays a critical role in

regulating fatty acid synthesis and is essential for lipid metabolism as it catalyzes the first step of

fatty acid synthesis and is found in all kingdoms of life except archaea (Cronan Jr and Waldrop

2002, Harwood 1988). It comprises three functional constituents: biotin carrier protein, biotin

carboxylase, and transcarboxylase. There are two distinct types of ACCase in nature: one

consists of three separate subunits (prokaryote-type), and the other is a single multi-domain

multifunctional polypeptide (eukaryote-type) (Sasaki et al. 1995). The eukaryote-type occurs in

mammals, fungi or yeast, while both the prokaryote- and the eukaryote- type can be found in

plants (Huang et al. 2002, Sasaki et al. 1995). Nine 454 sequencing reads encoding an ACCase

were found in our dataset, which shared ~72% amino acid identity to an eukaryotic type found in

the diatom *T. pseudonana*, suggesting *Eutreptiella* sp. ACCase to be of the eukaryotic type. In

the qPCR analysis, ACCase showed highest expression level in the high-lipid-light cultures,

indicating that this gene is related to photosynthetic lipid production in *Eutreptiella* sp.

Our results suggest that *Eutreptiella* sp. may have both type I fatty acid synthase (FASI)

and type II fatty acid synthase (FASII), which are the two major types of fatty acid synthases that

naturally occur in different groups of organisms (Nelson and Cox 2008). FASI, found in

vertebrate and fungi, is a single multifunctional polypeptide with 7 active sites, whereas FASII,

found in plants and bacteria, compose discrete and freely diffusible enzymes. Interestingly,

Euglena gracilis, which is another euglenoid species, has different types of fatty acid synthesis

pathways and enzymes to synthesize lipids under different environmental conditions (Delo et al.

1971, Goldberg and Bloch 1972, Inui et al. 1984). ACP-dependent FASI is located in the cytosol

and mainly produces C16 fatty acids (Goldberg and Bloch 1972). When grown in the light, *E.*

gracilis utilizes ACP-dependent FAS II in plastids to produce fatty acids (Ernst-Fonberg and

Bloch 1971). Inui et al. (1984) also reported a malonyl-CoA independent fatty acid synthesis

system to form wax ester from paramylon as energy storages in *E. gracilis* under anaerobic

conditions. We found genes coding for FASI and for β –ketoacyl-ACP reductase and Acyl-ACP

thioesterase of FASII in *Eutreptiella* sp. Our failure to retrieve the rest of FASII gene complex in

our cDNA libraries could be due to their being plastid-coded genes, which would not contain

Eut-SL that we used to construct the cDNA libraries, or just simply were not expressed. Whether

Eutreptiella sp. contains the other type of ACP-dependent FAS similar to that in *E. gracilis*

remains to be further investigated in the future.

Based on the fatty acid composition and possession of genes coding for enzymes

involved in fatty acid elongation and desaturation in *Eutreptiella* sp., it is evident that the ω3, ω6

and alternative Δ8 pathways occur in the euglenoid algae. Different from most of plants and most

algae, we found that *Eutreptiella* is similar to *Euglena* in using Δ8 desaturase to introduce the

third double bond on 20:2 (11, 14) to form 20:3 (8, 11, 14) (Meyer et al. 2003, Qi et al. 2004).

The 20:3 fatty acid is then converted to 22:5 EPA via stepwise Δ5 desaturase, Δ5 elongase and

Δ4 desaturase (Figure 5). In order to produce DHA, *Eutreptiella* may also utilize another Δ8

pathway to introduce the 4[th] double bound on 20:3 (11,14,17), resulting in 20:4 (8,11,14,17) for

later stepwise process of elongation and desaturation. Although Δ4 desaturase was not identified

in our dataset, it is very likely that *Eutreptiella* sp. has this gene since it was identified and

involved in DHA synthesis in *E. gracilis* (Meyer et al. 2003). In our study, both EPA would need

Δ4 desaturase to form their last double bond. We also detected trace amount of 15-tetracosenoic

acid (24:1) and 6,9,12,15,18,21-tetracosahexaenoic acid (24:6). The 24:6 fatty acid can be found

in some algae (Mansour et al. 2005), but the synthesis pathway is not well studied. We proposed

that the 24:6 fatty acid could potentially be synthesized by elongation from DHA or elongation

and desaturation from other C22 fatty acids.

387   The 16:4 (4,7,10,13) fatty acid was also abundant in phosphate-replete cultures of

Eutreptiella sp. Highly unsaturated stearidonic acid (18:4) and hexadecatetraenoic acid (16:4) are

produced in some macroalgae (Dembitsky et al. 1991, Ishihara et al. 2000) and microalgae

(Wallis and Browse 1999). Unsaturated C16 fatty acids have been identified in *E. gracilis*

(Constantopoulos and Bloch 1967). Increasing light intensity can enhance the production of 16:4

acid and 18:3 α-linolenic acid in *E. gracilis* (Constantopoulos and Bloch 1967). Korn (1964)

suggested a possible C16 pathway for *E. gracilis*: 16:1 (7) → 16:2 (7, 10) → 16:3 (7, 10, 13)

→16:4 (4, 7, 10, 13). Although we identified 16:4 (4, 7, 10, 13) in *Eutreptiella* sp., the genes

responsible for the synthesis of this abundant fatty acid remain uncertain.

Regulation of genes associated with lipid production. Our RT-qPCR revealed that many of the

genes examined had different expression levels between the dark and light periods. Genes

involved in high lipid biosynthesis showed higher expression levels under light condition,

suggesting that these genes promoted lipid production in *Eutreptiella* sp. during photosynthesis.

Those genes were ACCase, ACP, KAR2, KAR3, FAD, Δ8 desaturase, TE, AGPAT, and GPD.

The results of RT-qPCR verified our physiological observation from a 24h diel experiment,

showing that lipids in *Eutreptiella* sp. cells start to accumulate during the light period after cells

have actively divided (Kuo and Lin, 2012). Other evidence showed that the activity of ACCase is

correlated to that of the Calvin-Benson Cycle enzymes, with the greatest activities under light

(Sasaki et al. 1997) and negligible in the dark (Bao et al. 2000).

Three genes encoding carbon fixation enzymes showed higher expression levels under

the high-lipid-light condition, indicating that those enzymes are related to the biosynthesis of

precursors for lipid production. Phosphoribulokinase (PRK) and one copy of

phosphoenolpyruvate carboxykinase (i.e. PEPCK1) exhibited the highest level of relative gene

expression among these genes. PEPCK activity is linked to CO_2 fixation in algae (Reiskind and

Bowes 1991). It converts oxaloacetate into CO_2 and phosphoenolpyruvate, bridging C4 to C3

pathway in providing CO_2 for photosynthesis. It is noteworthy that C4 enzymes have been

identified in this species (Kuo et al. 2012). The phosphoenolpyruvate can later be converted into

pyruvate, which is an important precursor for Acetyl-CoA. PRK catalyzes the reaction of ATP

and ribulose 5-phosphate to produce ADP and ribulose 1,5-bisphosphate, the substrate needed to

capture CO_2 in photosynthesis. The concomitant up-regulation of the fatty acid synthesis and

photosynthesis genes indicates the linkage of the two carbon metabolic processes.

Acknowledgements

We thank the Center of Environmental Science and Engineering at UConn for providing the

facility of GC/MS/MS, and Yunyun Zhuang for helping with 454 sequencing. The research was

supported by the Predoctoral Awards of the Department of Marine Sciences at UConn (to RCK),

Multidisciplinary Environmental Research Award from the Center of Environmental Science and

Engineering at UConn (to RCK), and National Science Foundation "Assembling the Tree of Life

(AToL)" grant EF-0629624 (to SL and collaborators).

References

Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. 1990. Basic local alignment search tool. *J. Mol. Biol.* 215: 403-410

Álvarez-Góngora, C. & Herrera-Silveira J. A. 2006. Variations of phytoplankton community structure related to water quality trends in a tropical karstic coastal zone. *Mar. Pollut. Bull.* 52: 48-60

Anderson, D. M., Hoagland, P., Kaoru, Y. & White, A. W. 2000. Estimated annual economic impacts from harmful algal blooms (HABs) in the Unites States. Woods Hole, MA, USA: Woods Hole Oceanographic Isntitution

Arzel J., Martinez Lopez, F. X., Métailler, R., Stéphan, G., Viau, M., Gandemer, G. & Guillaume, J. 1994. Effect of dietary lipid on growth performance and body composition of brown trout (*Salmo trutta*) reared in seawater. Aquaculture. 123: 361-375

Bao, X., Focke, M., Pollard, M. & Ohlrogge, J. 2000. Understanding in vivo carbon precursor supply for fatty acid synthesis in leaf tissue. The Plant Journal 22: 39-50

Bates, S. S., Strain & P. M. 2006. Nutrients and phytoplankton in Prince Edward Island Inlets during late summer to fall: 2001-2003. Canadian Technical Report of Fisheries and Aquatic Sceicnes 2668. Canada: Fisheries and Oceans

Bloch, K. & Vance, D. 1977. Control mechanisms in the synthesis of saturated fatty acids. Annu. Rev. Biochem. 46: 263-298

Carvalho, A. P. & Malcata F. X. 2005. Preparation of fatty acid methyl esters for gas-chromatographic analysis of marine lipids: insight studies. J. Agric. Food Chem. 53: 5049-5059

Constantopoulos, G. & Bloch, K. 1967. Effect of Light Intensity on the Lipid Composition of Euglena gracilis. J. Biol. Chem. 242: 3538-3542

Cossins, A. R., Friedlander, M. J. & Prosser, C. L. 1977. Correlations between behavioral temperature adaptations of goldfish and the viscosity and fatty acid composition of their synaptic membranes. J. Comp. Physiol. A Neuroethol. Sens. Neural. Behav. Physiol. 120: 109-121

Cronan, J. E. & Waldrop, G. L. 2002. Multi-subunit acetyl-CoA carboxylases. Prog. Lipid Res. 41: 407-435

Delo, J., Ernst-Fonberg, M. L. & Bloch, K. 1971. Fatty acid synthetases from Euglena gracilis. Arch. Biochem. Biophys. 143: 384-391

Dembitsky, V. M., Pechenkina-Shubina, E. E. & Rozentsvet, O. A. 1991. Glycolipids and fatty acids of some seaweeds and marine grasses from the black sea. Phytochemistry 30: 2279-2283

Edgar, R. C. 2010. Search and clustering orders of magnitude faster than BLAST. Bioinformatics 26: 2460-2461

Ernst-Fonberg, M. L & Bloch, K. 1971. A chloroplast-associated fatty acid synthetase system in Euglena. Arch. Biochem. Biophys. 143: 392-400

Goldberg, I. & Bloch, K. 1972. Fatty acid synthetases in *Euglena gracilis*. J. Biol. Chem. 247: 7349-7357

Götz, S., García-Gómez, J. M., Terol, J., Williams, T. D., Nagaraj, S. H., Nueda, M. J., Robles,
M., Talón, M., Dopazo, J. & Conesa, A. 2008. High-throughput functional annotation and
data mining with the Blast2GO suite. Nucleic Acids Res 36: 3420-3435
Gouveia, L. & Oliveira, A. 2009. Microalgae as a raw material for biofuels production. J. Ind.
Microbiol. Biotechnol. 36: 269-274
Griffiths, M. & Harrison, S. 2009. Lipid productivity as a key characteristic for choosing algal
species for biodiesel production. J. Appl. Phycol. 21: 493-507
Gunstone, F. D., Harwood, J. L. & Dijkstra A. J. 2007. *The Lipid Handbook ,* 2nd ed. Taylor &
Francis, FL, USA
Guo, R.& Ki, J. S. 2012. Evaluation and validation of internal control genes for studying gene
expression in the dinoflagellate Prorocentrum minimum using real-time PCR. Eur. J.
Protistol. 48(3): 199-206
Guschina, I. A & Harwood, J. L. 2006. Lipids and lipid metabolism in eukaryotic algae. Prog
Lipid Res 45: 160-186
Harwood, J. L. 1988. Fatty acid metabolism. Annu. Rev. Plant Physiol. Plant Mol. Biol. 39: 101-
138
Henriksen, P., Riemann, B., Kaas, H., Sørensen, H. M. & Sørensen, H. L. 2002. Effects of
nutrient-limitation and irradiance on marine phytoplankton pigments. J. Plankton Res. 24:
835-858
Hu, Q., Sommerfeld, M., Jarvis, E., Ghirardi, M., Posewitz, M., Seibert, M. & Darzins, A. 2008.
Microalgal triacylglycerols as feedstocks for biofuel production: perspectives and
advances. The Plant J. 54: 621-639
Huang, S., Sirikhachornkit. A., Faris, J. D., Su, X., Gill, B. S., Haselkorn, R. & Gornicki, P. 2002.
Phylogenetic analysis of the acetyl-CoA carboxylase and 3-phosphoglycerate kinase loci
in wheat and other grasses. Plant Mol. Biol. 48: 805-820
Hulanicka, D., Erwin, J & Bloch, K. 1964. Lipid metabolism of Euglena gracilis. J. Biol. Chem.
239: 2778-2786
Huppe, H. C.& Turpin, D. H. 1994. Intergration of carbon and nitrogen metabolism in plant and
algal cells. Annu. Rev. Plant Physiol. Plant Mol. Biol. 45: 577-607
Illman, A. M., Scragg, A. H. & Shales, S. W. 2000. Increase in Chlorella strains calorific values
when grown in low nitrogen medium. Enzyme Microb. Technol. 27: 631-635
Inui, H., Miyatake, K., Nakano, Y. & Kitaoka, S. 1984. Fatty acid synthesis in mitochondria of
Euglena gracilis. Eur. J. Biochem. 142: 121-126
Ishihara, K., Murata, M., Kaneniwa, M., Saito, H., Komatsu, W. & Shinohara, K. 2000.
Purification of stearidonic Acid (18:4(n-3)) and hexadecatetraenoic Acid (16:4(n-3))
from algal fatty acid with lipase and medium pressure liquid chromatography. Biosci.
Biotechnol. Biochem. 64: 2454-2457
Klein Breteler, W. C. M., Koski, M. & Rampen, S. 2004. Role of essential lipids in copepod
nutrition: no evidence for trophic upgrading of food quality by a marine ciliate. Mar. Ecol.
Prog. Ser. (Halstenbek) 274: 199
Korn, E. D. 1964. The fatty acids of Euglena gracilis. J. Lipid Res. 5: 352-362
Kuo, R. C & Lin, S. 2012. Ectobiotic and endobiotic bacteira associated with *Eutreptiella* sp.
isolated from Long Island Sound. Protist 164 (1): 60-74
Kuo, R. C., Zhang, H., Zhuang, Y., Hannick, L.& Lin, S. 2013 Transcriptomic study reveals
widespread spliced leader *trans*-splicing, short 5'-UTRs and potential complex carbon

fixation mechanisms in the euglenoid alga *Eutreptiella* sp. PLoS ONE. 8(4): e60826. doi:10.1371/journal.pone.0060826.

Le Bail, A., Dittami, S., De Franco, P. O., Rousvoal, S., Cock, M., Tonon, T.& Charrier, B. 2008. Normalisation genes for expression analyses in the brown alga model Ectocarpus siliculosus. BMC Mol. Biol. 9: 75

Lee S., Yoon, B. D. & Oh, H. M. 1998. Rapid method for the determination of lipid from the green alga Botryococcus braunii. Biotechnol Tech. 12: 553-556

Lindholm, T. 1995. Green water caused by *Eutreptiella gymnastica* (Euglenophyceae) in a stratified Baltic Sea inlet. In *Lassus, P., Arzul G., Denn EE-L., Gentien, P., Baut CM-L., [Eds.] Harmful Marine Alagae Blooms (Proceedings of the Sixth International Conference on Toxic Marine Phytoplankton)*. Nantes, France.

Liu, Z. Y., Wang, G. C.& Zhou, B. C. 2008. Effect of iron on growth and lipid accumulation in Chlorella vulgaris. Bioresour. Technol. 99: 4717-4722

Mansour, M. P., Frampton, D. M. F., Nichols, P. D., Volkman, J. K.& Blackburn, S. I. 2005. Lipid and fatty acid yield of nine stationary-phase microalgae: Applications and unusual C24–C28; polyunsaturated fatty acids. J. Appl. Phycol. 17: 287-300

Maréchal, E., Block, M. A., Dorne, A. J., Douce, R.& Joyard, J. 1997. Lipid synthesis and metabolism in the plastid envelope. Physiol. Plant 100: 65-77

Mccarthy, G. 2008. Monitoring Long Island Sound 2008. Connecticut Department of Environmental Protection, Connecticut

Meyer, A., Cirpus, P., Ott, C., Schlecker, R., Zähringer, U. & Heinz, E. 2003. Biosynthesis of docosahexaenoic acid in *Euglena gracilis*: biochemical and molecular evidence for the involvement of a Δ4-fatty acyl group desaturase. Biochemistry 42: 9779-9788

Nelson, D. L. &Cox, M. M. 2008. Lipid biosynthesis. In *Nelson, D. L. & Cox, M. M. [Eds.] Lehninger Principles of Biochemistry*. W. H. Freeman and Company, New York, pp 805-845

Olli, K., Heiskanen, A. S. & Seppälä, J. 1996. Development and fate of *Eutreptiella gymnastica* bloom in nutrient-enriched enclosures in the coastal Baltic Sea. J. Plankton Res. 18: 1587-1604

Qi, B., Fraser, T., Mugford, S., Dobson, G., Sayanova, O., Butler, J., Napier, J. A., Stobart, A. K. & Lazarus, C. M. 2004. Production of very long chain polyunsaturated omega-3 and omega-6 fatty acids in plants. Nat. Biotech. 22: 739-745

Rainuzzo, J. R., Reitan, K. I. & Olsen, Y. 1997. The significance of lipids at early stages of marine fish: a review. Aquaculture 155: 103-115

Ratledge, C. 1988. An overview of microbial lipids. In C Ratledge, SG Wilkinson, eds, Microbial Lipids. Academic Press, New York, pp 3-21

Raven, J. A. 1974. Carbon dioxide fixation. In *Stewart, W. D. P. [Ed.] Algal Physiology and Biochemistry: Biotenical Monographs, Vol 10*. University of California Press, California, pp 434-455

Reiskind, J. B. & Bowes, G. 1991. The role of phosphoenolpyruvate carboxykinase in a marine macroalga with C4-like photosynthetic characteristics. PNAS 88: 2883-2887

Rodríguez-Graña, L., Calliari, D., Conde, D., Sellanes, J. & Urrutia, R. 2008. Food web of a SW Atlantic shallow coastal lagoon: spatial environmental variability does not impose substantial changes in the trophic structure. Mar. Ecol. Prog. Ser. 362: 69-83

Sasaki, Y., Konishi, T. & Ngano, Y. 1995. The compartmentation of Acetyl-Coenzyme A carboxylase in plants. Plant Physiol. 108: 445-449

Sasaki, Y., Kozaki, A. & Hatano, M. 1997. Link between light and fatty acid synthesis:
Thioredoxin-linked reductive activation of plastidic acetyl-CoA carboxylase. PNAS 94:
11096-11101
Schweizer, E. & Hofmann, J. 2004. Microbial type I fatty acid synthases (FAS): major players in
a network of cellular FAS systems. Microbiol Mol. Biol. Rev. 68: 501-517
Seong, K. A., Jeong, H. J., Kim, S., Kim, G. H. & Kang, J. H. 2006. Bacterivory by co-
occurring red-tide algae, heterotrophic nanoflagellates, and ciliates. Mar. Ecol. Prog. Ser.
322: 85-97
Singh, S. C. 2002. Role of lipids and fatty acids in stress tolerance in cyanobacteria. Acta.
Protozool. 41: 297
Stonik, I. 2007. Species of the genus *Eutreptiella* (Euglenophyceae) from Russian waters of
East/Japan Sea. Ocean Sci. J. 42: 81-88
Stonik, I. V. & Selina, M. S. 2001. Species composition and seasonal dynamics of density and
biomass of euglenoids in Peter the Great Bay, Sea of Japan. Russ. J. Mar. Biol. 27: 174-
176
Takeyama, H., Kanamaru, A., Yoshino, Y., Kakuta, H., Kawamura, Y. & Matsunaga, T. 1997.
Production of antioxidant vitamins, β-carotene, vitamin C, and vitamin E, by two-step
culture of *Euglena gracilis* Z. Biotechnol. Bioeng. 53: 185-190
Tang, K. W. & Taal, M. 2005. Trophic modification of food quality by heterotrophic protists:
species-specific effects on copepod egg production and egg hatching. J. Exp. Mar. Biol.
Ecol. 318: 85-98
Thelen, J. J. & Ohlrogge, J. B. 2002. Metabolic engineering of fatty acid biosynthesis in plants.
Metab. Eng. 4: 12-21
Upchurch, R. 2008. Fatty acid unsaturation, mobilization, and regulation in the response of
plants to stress. Biotechnol. Lett. 30: 967-977
Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A. & Speleman,
F. 2002. Accurate normalization of real-time quantitative RT-PCR data by geometric
averaging of multiple internal control genes. Genome Biology 3: research0034.0031 -
research0034.0011
Wallis, J. G. & Browse, J. 1999. The Δ8-Desaturase of *Euglena gracilis*: an alternate pathway
for synthesis of 20-carbon polyunsaturated fatty acids. Arch. Biochem. Biophys. 365:
307-316
Wallis, J. G, Watts, J. L. & Browse, J. 2002. Polyunsaturated fatty acid synthesis: what will they
think of next? Trends Biochem. Sci. 27: 467-473
White, S. W., Jie, Z., Yong-Mei, Z. & Rock, C. O. 2005. The structural biology of type II fatty
acid biosynthesis. Annu. Rev. Biochem. 74: 791-831
Worsham, L. M. S., Williams, S. G. & Ernst-Fonberg, M. L. 1993. Early catalytic steps of
Euglena gracilis chloroplast type II fatty acid synthase. Biochimica et Biophysica Acta
(BBA) - Lipids and Lipid Metabolism 1170: 62-71
Yamane, Y. I., Utsunomiya, T., Watanabe, M. & Sasaki, K. 2001. Biomass production in
mixotrophic culture of Euglena gracilis under acidic condition and its growth energetics.
Biotechnol. Lett. 23: 1223-1228

Figure legends

Figure 1. Growth and lipid production of *Eutreptiella* cultures under phosphate-replete and phosphate-depleted conditions. (A) Growth curves of *Eutreptiella* cultures. (B) Growth rates and lipid fluorescence levels at the time when cultures were harvested.

Figure 2. Common and unique expressed genes between different culture conditions. HL: high-lipid-light cultures. HD: high-lipid-dark cultures. LL: low-lipid-light cultures. LD: low-lipid-dark cultures. Numbers outside the diagram represent total number of genes found in each library.

Figure 3. Putative lipid biosynthesis pathway in *Eutreptiella* sp. inferred from our transcriptomic data. GPD: glycerol 3-phosphate dehydrogenase (EC: 1.1.1.53); GPAT: glycerol-3-phosphate acyltransferase (EC: 2.3.1.15); AGPAT: acylglycerophosphate acyltransferase(EC: 2.3.1.51); PP: phosphatidate phosphatase (EC: 3.1.3.4); DGAT: diacylglycerol acyltransferase (EC: 2.3.1.20); FA: fatty acids; FA-CoA: fatty acyl-CoAs. FA: fatty acids. Dashed boxes indicate enzymes that were not found in our dataset. The process of FA elongation and desaturation is shown in Figure 5.

Figure 4. Chromatograms of total fatty acid esters from *Eutreptiella* sp. Note that DHA (peak 19) was the most abundant fatty acid. Corresponding fatty acids (labeled by length of carbon, e.g. C16) were listed in Table 3. (A) Chromatogram of fatty acid esters from the phosphate–replete cultures. (B) Chromatogram of fatty acid esters from the phosphate-depleted cultures. Insets are closer-ups of some of the small peaks of esters. The first number (bold-typed) by the peak is peak ID, the second number retention time, and the third number peak area (only trace elements are shown). Note that retention times between the two samples are slightly different.

Figure 5. Putative pathways of fatty acids elongation and desaturation inferred from *Eutreptiella*

sp. transcriptome annotation and GC/MS/MS analyses. Boxes with dashed lines indicate the

enzymes or substrates that were not found in our study.

Figure 6. Expression levels of genes involved in fatty acids biosynthesis in *Eutreptiella* sp. HD:

high-lipid-dark cultures. HL: high-lipid-light cultures. LL: low-lipid-light cultures. LD: low-

lipid-dark cultures.

Figure 7. Expression levels of genes involved in fatty acids biosynthesis (FAD, Δ 8, and TE) and

triacylglycerol biosynthesis (GPD, GPAT, and AGPT). HD: high-lipid-dark cultures. HL: high-

lipid-light cultures. LL: low-lipid-light cultures. LD: low-lipid-dark cultures.

Figure 8. Expression levels of genes involved in carbon fixation. HD: high-lipid-dark cultures.

HL: high-lipid-light cultures. LL: low-lipid-light cultures. LD: low-lipid-dark cultures.

641 Table 1. Genes potentially involved in fatty lipid biosynthesis in *Eutreptiella* sp. Sequences and BLAST
642 results were listed in the supporting file "lipid_biosynthesis_genes".

Gene	EC code	Number of transcripts
Fatty acid biosynthesis		
Acetyl-CoA carboxylase	EC 6.4.1.2	2
Fatty acid synthase (animal type)	EC 2.3.1.85	1
Acyl-ACP thioesterase	EC 3.1.2.14	1
Acyl carrier protein		1
β-ketoacyl-ACP reductase	EC 1.1.1.100	3
Fatty acid desaturation		
Δ^5 desaturase	EC 1.14.19.-	2
Δ^6 desaturase	EC 1.14.19.-	5
Δ^8 desaturase	EC 1.14.19.-	2
Δ^9 desaturase (Acyl-ACP desaturase)	EC 1.14.19.2	1
Δ^{12} desaturase	EC 1.14.19.-	1
Δ^{15} desaturase	EC 1.14.19.-	1
Fatty acid elongation		
Δ^5 elongase		1
Δ^6 elongase	EC 1.14.19.-	1
Δ^9 elongase	EC 1.14.19.2	1
Palmitoyl-CoA hydrolase	EC 3.1.2.2	1
Palmitoyl hydrolase	EC 3.1.2.22	1
acetyl-CoA C-acyltransferase	EC 2.3.1.16	1
3-hydroxyacyl-CoA dehydrogenase	EC 1.1.1.35	2
Triacylglycerol biosynthesis		
Glyerol-3-phosphate dehydrogenase	EC 1.1.5.3	1
Glycerol3-phosphate acyltransferase	EC 2.3.1.15	2
Acylglycerophosphate acyltransferase	EC 2.3.1.51	2
Fatty acid metabolism		
Acyl-CoA dehydrogenase	EC 1.3.99.3	3
Long-chain-fatty-acid-CoA ligase	EC 6.2.1.3	2
Acetyl-CoA C-acyltransferase	EC 2.3.1.16	1
3-hydroxyacyl-CoA dehydrogenase	EC 1.1.1.35	2
Carnitine o-palmitoyltransferase	EC 2.3.1.21	1
Enoyl-CoA hydratase	EC 4.2.1.17	1
Acetyl-CoA thiolase	EC 2.3.1.9	1
Aldehyde dehydrogenase (AND+)	EC 1.2.1.3	2

643 Table 2. Types and relative abundance of fatty acids detected in *Eutreptiella* sp.

Peak ID	Retention time	Systematic name	Abbreviation	Relative abundance (%)F/2*	Relative abundance (%)F/2-P*
1	26.19	4,7,10,13-Hexadecatetraenoic acid	16:4(n-3)	98.04	11.44
2	26.37	7,10-Hexadecadienoic acid	16:2(n-6)	2.59	0.08
3	26.47	7,10,13-Hexadecatrienoic acid	16:3(n-3)	12.63	0.54
4	26.62	7-Hexadecenoid acid	16:1(n-9)	11.89	0.27
5	26.97	9-Hexadecenoid acid	16:1(n-7)	13.54	0.79
6	27.1	Hexadecanoic acid	16:0	90.45	11.28
7	29.11	6,9,12-Octadecatrienoic acid	18:3(n-6)	3.90	0.13
8	30.04	6.9.12.15-Octadecatetraenoic acid	18:4(n-3)	41.00	12.40
9	30.23	9,12-Octadecadienoic acid	18:2(n-6)	3.20	0.53
10	30.36	9,12,15-Octadecatrienoic acid	18:3(n-3)	72.63	23.58
11	30.47	9-Octadecenoic acid	18:1(n-9)	7.97	0.58
12	30.85	Octadecanoic acid	18:0	8.43	2.12
13	33.1	5,8,11,14-Eicosatetraenoic acid	20:4(n-6)	6.44	1.88
14	33.22	5,8,11,14,17-Eicosapentaenoic acid	20:5(n-3)	78.69	40.02
15	33.53	8,11,14,17-Eicosatetraenoic acid	20:4(n-3)	8.39	4.02
16	33.77	8,11-Eicosadienoic acid	20:2(n-9)	1.13	0.71
17	33.87	11,14,17-Eicosatrienoic acid	20:3(n-3)	3.16	1.02
18	36.17	4,7,10,13,16-Docosapentaenoic acid	22:5(n-6)	68.78	54.93
19	36.3	4,7,10,13,16,19-Docosahexaenoic acid	22:6(n-3)	100.00	100.00
20	36.4	7,10,13,16-Docosatetraenoic acid	22:4(n-6)	43.11	47.18
21	36.49	7,10,13,16,19-Docosapentaenoic acid	22:5(n-3)	3.56	6.03
22	37.54	Docosanoic acid	22:0	5.94	1.01
23	38.79	6,9,12,15,18,21-Tetracosahexaenoic acid	24:6 (n-3)	5.32	0.85
24	40.14	15-Tetracosenoic acid	24:1(n-9)	0.46	-

645 *The numbers of relative abundance were obtained by dividing the peak area of the fatty acid to the peak
646 area of DHA; F/2 and F/2-P depict phosphate-replete and phosphate-depleted F/2 medium, respectively.

 Table 3. Primers used in reverse-transcription quantitative PCR (RT-qPCR) analysis in this study.

Primer name	Sequences (5´-3´)	RT-qPCR for *Eutreptiella* sp. gene
ACPF1	GCAGCAAAGCAAATCAAACCA	acyl carrier protein forward
ACPF2	CAACCCAACTCTCACTTCTACAT	acyl carrier protein forward
ACPR	GCATCAGCGACCGATACAATA	acyl carrier protein reverse
AGPCF	TTACGAACTTGCGATGCTTCA	1-acyl-sn-glycerol-3-phosphate acyltransferase forward
AGPCR1	GGAGGTATGATTGGAGATGATGAG	1-acyl-sn-glycerol-3-phosphate acyltransferase reverse
AGPCR2	CGATGACTTGACACCACCAA	1-acyl-sn-glycerol-3-phosphate acyltransferase reverse
ACCASE-F	GTGTTGGCATGTTGTCTCTTG	ACCase forward
ACCASE-R1	GGATACCGCCATTGGAACC	ACCase reverse
ACCASE-R2	GAAGTCGGCTTGAATCTGTGA	ACCase reverse
DELTA8FADF	GACATCATTGAGAATTATCGGAACC	delta-8 fatty acid desaturase forward
DELTA8FADR1	GCGTCACCATCAGCATAGC	delta-8 fatty acid desaturase reverse
DELTA8FADR2	CGGAAGTCCTCAGCAACCT	delta-8 fatty acid desaturase reverse
PRKF	TTCCAAGCCAATCTACAACCAT	Phosphoribulokinase forward
PRKR1	CGAATCTGAGACGGAAGAACTT	Phosphoribulokinase reverse
PRKR2	GCAACACGGTCATCCAACA	Phosphoribulokinase reverse
ThioesteraseF	AGCAATCGGCAGGAGCAA	Thioesterase forward
ThioesteraseR1	ACAGGCAAGCGGCATCT	Thioesterase reverse
ThioesteraseR2	TGGAACAACGGCAGCAATG	Thioesterase reverse
KAR1-F	AAGCCATCGCTGTTGGATTG	β-Ketoacyl-ACP reductase1 forward
KAR1-R1	ACCGCATCCCAATCAGCAA	β-Ketoacyl-ACP reductase1 reverse
KAR1-R2	CTGATTCGTAATGTCTGCCTGAA	β-Ketoacyl-ACP reductase1 forward
KAR2-F1	TTCAACGACGCAGCCTGTA	β-Ketoacyl-ACP reductase2 forward
KAR2-F2	CCGTCAACTCGCTCATTGTT	β-Ketoacyl-ACP reductase2 forward
KAR2-R	GCCTCTTCTGTCATCTTGTCAA	β-Ketoacyl-ACP reductase2 reverse
KAR3-F1	CGGTGCTTCCAGAGGTGTT	β-Ketoacyl-ACP reductase3 forward
KAR3-F2	GGAGCCTCTGTGGTTGTCA	β-Ketoacyl-ACP reductase3 forward
KAR3-R1	TTCATCTGCTTGTCCATCATCTC	β-Ketoacyl-ACP reductase3 reverse
FAD-F	AACATTCTAACGCCACACCAT	FAD family gene forward
FAD-R1	ACTGCCACATTGAGACTCCT	FAD family gene reverse
FAD-R2	ACAGTCGGTGCGTTGTAATC	FAD family gene reverse
GPD-F	CTTGGCAGGCTTCTGTAATCG	glycerol-3-phosphate dehydrogenase forward
GPD-R1	GCTTCGTGGATCGTGAGGA	glycerol-3-phosphate dehydrogenase reverse

GPD-R2	ATCACCACAACATCGTACTCCT	glycerol-3-phosphate dehydrogenase reverse
GPAT-F	CCAGATGAGGCTTCTTCTACAA	glycerol-3-phosphate acyltransferase forward
GPAT-R1	CAACAGCGATGGCGTTCAT	glycerol-3-phosphate acyltransferase reverse
GPAT-R2	TCACCGCATCCACGAACT	glycerol-3-phosphate acyltransferase reverse
PEPCASE-F	GCCTGTCTTGAATGACTCCTT	Phosphoenolpyruvate carboxylase forward
PEPCASE-R1	GCATACTTGACCAGCAGTGA	Phosphoenolpyruvate carboxylase reverse
PEPCASE-R2	CCGTGGTTCTGGATCAGGTT	Phosphoenolpyruvate carboxylase reverse
PEPCK1-F1	AAGGCTCTGGCGACATTAAG	Phosphoenolpyruvate carboxykinase1 forward
PEPCK1-F2	TGGCGACATTAAGAACAGCATA	Phosphoenolpyruvate carboxykinase1 forward
PEPCK1-R	TCTGGTTGAGAGGAATGAGTGT	Phosphoenolpyruvate carboxykinase1 reverse
PEPCK2-F	TGGTATTGATTGAGCACAACAATG	Phosphoenolpyruvate carboxykinase2 forward
PEPCK2-R1	GAGGTCTTGATTGTGGATTGGAAT	Phosphoenolpyruvate carboxykinase2 reverse
PEPCK2-R2	TACTTGCGGATCTCGGTGATA	Phosphoenolpyruvate carboxykinase2 reverse
PEPCK3-F	GCAACGGACGATGTGAACC	Phosphoenolpyruvate carboxykinase3 forward
PEPCK3-R1	CGCATGTGAGGAACCAACTG	Phosphoenolpyruvate carboxykinase2 reverse
PEPCK3-R2	GTCTGATTGCGGTCCAACTTAT	Phosphoenolpyruvate carboxykinase2 reverse
PPDK-F1	CTTATCTCCCACAAATGACTACCC	phosphate dikinase forward
PPDK-F2	ACTGGTTCATCTTCGCCTCAT	phosphate dikinase forward
PPDK-R	TCAGCCAAGTTCGCACCTT	phosphate dikinase reverse
EF-1aF	GCACCTCTCACCATCTTACAAT	elongation factor 1 alpha forward
EF-1aR1	GTTCCGCCTTCAGCTTGTC	elongation factor 1 alpha reverse
EF-1aR2	TTTCCTTTCGCTTTCCCTTCTT	elongation factor 1 alpha reverse
TubulinF	CCTTGGYCAGGCTGGTATCC	tubulin forward
TubulinR1	CGGCGTTGAAGCACAAGAA	tubulin reverse
TubulinR2	AGTTGTTGGCGGCATCCT	tubulin reverse
ActinF	AAGCGAAGCGTGGTATCTTG	actin forward
ActinR1	ACATAATCTGCGTCATCTTCTCC	actin reverse
ActinR2	GGTTCATAGGAGCCTCAGTGA	actin reverse

A

B

Kuo et al. Figure 1

Kuo et al. Figure 2

FAT biosynthesis pathway   TAG biosynthesis pathway

Kuo et al. Figure 3

Kuo et al. Figure 4

Kuo et al. Figure 5

Kuo et al. Figure 6

Kuo et al. Figure 7

Kuo et al. Figure 8