Contrasting functions of *Arabidopsis* SUMO1/2 isoforms with SUMO3

intersect to modulate innate immunity and global SUMOylome responses

Kishor Dnyaneshwar Ingole[1,2], Mritunjay Kasera[1], Harrold A. van den Burg[3], and Saikat

Bhattacharjee[1*]

[1]Laboratory of Signal Transduction and Plant Resistance, UNESCO-Regional Centre for

Biotechnology (RCB), NCR Biotech Science Cluster, 3[rd] Milestone, Faridabad-Gurgaon

Expressway, Faridabad- 121 001, Haryana, India.

[2]Kalinga Institute of Industrial Technology (KIIT) University, Bhubaneswar- 751 024, Odisha,

India.

[3]Molecular Plant Pathology, Swammerdam Institute for Life Sciences, Faculty of Science,

University of Amsterdam, Science Park 904, 1098 XH, Amsterdam, Netherlands.

* Corresponding author:

Saikat Bhattacharjee

E-mail: saikat@rcb.res.in

Short Title: Intersecting functions of *Arabidopsis* SUMO isoforms

Abstract

Reversible covalent attachment of SMALL UBIQUITIN-LIKE MODIFIERS (SUMOs) on target proteins regulate diverse cellular process across all eukaryotes. In *Arabidopsis thaliana*, most mutants with perturbed global SUMOylome display severe impairments in growth and adaptations to physiological stresses. Since SUMOs self-regulate activities of SUMOylation-associated proteins, existence of multiple isoforms introduces possibilities of their functional intersections which remain unexplored especially in plant systems. Using well-established defense responses elicited against virulent and avirulent *Pseudomonas syringae* pv. *tomato* strains, we investigated crosstalks in individual and combinatorial Arabidopsis *sum* mutants. Here we report that while *SUM1* and *SUM2* additively, but not equivalently suppress basal and TNL-specific immunity via down-regulation of salicylic acid (SA)-dependent responses, *SUM3* promotes these defenses genetically downstream of SA. Remarkably, the expression of *SUM3* is transcriptionally suppressed by SUMO1 or SUMO2. The loss of *SUM3* not only lowers basal or post-bacterial challenge responsive enhancements of SUMO1/2-congugates but also reduces upregulation dynamics of defensive proteins and SUMOylation-associated transcripts. Combining a *sum3* mutation partially attenuates heightened immunity of *sum1* or *sum2* mutants suggesting intricate functional impingements among these isoforms in optimizing immune amplitudes. Similar *SUM1-SUM3* intersections also affect global SUMOylome responses to heat-shock affecting most notably the induction of selective heat-shock transcription factors. Overall, our investigations reveal novel insights into auto-regulatory mechanisms among SUMO isoforms in host SUMOylome maintenance and adjustments to environmental challenges.

Author Summary

In plants, similar to animals, protein functions are regulated at multiple levels. One prevalent mode is to allow covalent linkage of small proteins to specific amino acids on targets thereby affecting its fate and function. One such kind of modification named as SUMOylation involves attachment of SUMO proteins. A plant maintains strict control over its pool of SUMOylated proteins (termed SUMOylome) which upon biotic or abiotic stresses are altered to facilitate appropriate responses, returning back to steady-state when the threat subsides. In mutants of the model plant *Arabidopsis thaliana* having disturbed steady-state SUMOylome, growth and developmental defects ensue. These mutants are auto-immune showing more resistance to infection by the bacterial pathogen *Pseudomonas syringae*. However, Arabidopsis SUMO-family are comprised of multiple members raising the question about their specificity or functional crosstalks. We discovered that two SUMO members function in coordination to suppress immunity including the repression of a third member which supports defenses. The expression of this third member during pathogen attack or heat-shock influences the responsive changes in the host SUMOylome likely suggesting SUMOs themselves play vital role in these adaptations. Overall, our work highlights novel intersections of SUMO members in mounting stress-specific responses.

Introduction

Post-translational modifications (PTMs) dynamically and reversibly modulate target proteins properties either by affecting their fate, location, or function and facilitate transitory adaptations often necessary for developmental passages and responding to biotic/abiotic cues. Of special importance in this category is the reversible attachment of the SMALL UBIQUITIN-LIKE MODIFIER (SUMO) through a process known as SUMOylation. This highly conserved mode of covalently modifying proteins is utilized primarily to regulate nuclear functions such as DNA

replication, chromatin remodeling, transcription, and post-transcriptional processes in eukaryotes

[1–3]. In plants, temperature perturbations, salinity changes and application of defensive hormones

such as salicylic acid (SA) cause massive changes on global SUMOylome [4–6]. Interestingly, a

vast majority of differentially SUMOylated proteins are nuclear, perhaps indicative of a rapid

mechanism to modulate transcriptome in response to stimulus. Computational studies identify

SUMOylation-annotated proteins as central relay players of protein-protein interaction webs,

especially for transcriptional processes [7].

SUMOylation cascade recruits processed SUMOs with C-terminus di-glycine (GG) residues, a

product of catalysis by specialized SUMO or Ubiquitin-like proteases (ULPs), to covalently attach

via an isopeptide bond to ε-amino group of lysine residues on target proteins. The modified lysine

often is a part of a partially conserved motif, ψ-K-X-D/E (ψ = hydrophobic amino acid, X= any

amino acid, D/E= aspartate/glutamate). A conjugatable SUMO, subsequently forms a thiol-ester

bond with the SUMO E1 ACTIVATING ENZYME (SAE). A trans-esterification reaction further

shuttles SUMO to the SUMO E2 CONJUGATING ENZYME 1 (SCE1) and then to target lysine

on SUMOylation substrates. Enrichment of SUMOylated proteins upon heat shock reveal that

SUMO1 can conjugate to itself and form poly-SUMO1 chains [5]. Although SCE1 is capable of

catalyzing polySUMO-chain formation, members of SUMO E4 ligases the PROTEIN

INHIBITOR OF ACTIVATED STAT LIKE protein (PIAL1/2 in Arabidopsis) enhance

polySUMOylation [2,8,9]. Substrate specificities both for mono- or polySUMOylation are further

regulated by SUMO E3 LIGASES such as HIGH PLOIDY2/METHYL METHANE

SULFONATE21 (HPY2/MMS21) and SAP and MIZ1 (SIZ1) [10]. Fate of covalently-attached

SUMOs may either include de-conjugation and recycling by SUMO proteases or targeted

proteolysis of the substrate through ubiquitin-mediated pathway. Indeed, proteasome components

are enriched in poly-SUMO1 pull-downs [5]. These interactions are non-covalent in nature

facilitated by hydrophobic amino acid-rich SUMO-interaction motifs (SIMs) present in the

recipient. A group of moderately conserved SUMO-targeted ubiquitin ligases (STUbLs) bind poly-

SUMO chains via internal SIMs to ubiquitinylate the substrate [11]. Interestingly, SIMs are also

only enriched in the SUMOylation-associated proteins such as SIZ1, SCE1, SAE2, SUMO-

protease EARLY IN SHORT DAYS 4 (ESD4), PIAL1/2, STUbLs implying strong auto-

regulatory mechanisms [8,12]. Not the least, increasing evidences that reciprocal influences of

phosphorylation, ubiquitination, and acetylation that often compete with or modulate the efficacy

of SUMOylation highlights the complexity of crosstalks among PTMs [13–15].

Unlike fruit fly, worm, or yeast, humans and plants express multiple SUMO isoforms [16]. In

Arabidopsis thaliana, 4 SUMO isoforms are expressed namely SUMO1, SUMO2, SUMO3 and

SUMO5 [4,17]. SUMO1 and SUMO2 share 89% sequence identity, whereas SUMO3 and SUMO5

are considerably diverged from SUMO1 (48% and 35% identical, respectively). The homolog pair

SUMO1/2 are a result of genomic duplication event of a SUMO clade that preceded the evolution

of monocots and eudicots from common ancient angiosperms [18]. Tandem organization of *SUM2*

and *SUM3* genes in Arabidopsis are a result of gene duplication subsequently followed by

diversification of only *SUM3* sequences. At relative expression levels, *SUM1/2* are more abundant

than *SUM3* or *SUM5* [19]. Partial overlaps in tissue-specific expression patterns and biochemical

properties taken together with embryonic lethality of *sum1 sum2* double mutant indicate that plants

require at least one functional copy of either of these redundant isoforms [20–22]. The *Arabidopsis*

sum3 mutant is viable with mild late-flowering phenotype [21]. Acutely different SUMO3

however, unlike SUMO1/2, cannot form poly-SUMO chains *in vitro*, and shows little or no change

to heat-shock treatment [4,17,19,21,23]. Not the least, SUMO1/2 but not SUMO3-modified

targets, are efficiently deconjugated by SUMO proteases *in vitro* [17,24].

Host SUMOylome readjustments play a vital role in regulation of plant immunity and been

comprehensively highlighted in several excellent reviews [25–27]. This was first evident in the

Arabidopsis SUMO E3 ligase *SIZ1* mutant (*siz1-2*) and subsequently in SUMO protease mutants

of *OVERLY TOLERANT TO SALT 1/2 (OTS1/2)* and *ESD4* [6,20,28,29]. Increased accumulation

of defensive hormone salicylic acid (SA) and constitutive expression of PATHOGENESIS-

RELATED (PR) proteins in these mutants conferred enhanced resistance when challenged with

the bacterial pathogen *Pseudomonas syringae* pv. *tomato* strain DC3000 (*PstDC3000*). In this

context, it is therefore not surprising that pathogens attempt to manipulate host SUMOylome to

increase their colonization efficiencies [30–32]. Several bacterial phytopathogenic effectors

interfere with host SUMOylation as a mode to suppress immunity [33,34]. XopD, a secreted

effector from *Xanthomonas campestris pv. vesicatoria* (*Xcv*) de-conjugates SUMO from unknown

targets in plants [35]. Mutations that disrupt SUMO-protease functions not only render the cognate

strain deficient in virulence but also lower host defense induction.

Suggestively, both SUM1/2 jointly suppress SA-dependent defenses since plants null for *SUM1*

and expressing microRNA-silenced *SUM2* (*sum1-1 amiR-SUM2*) have drastically reduced total

SUMO1/2-conjugates compared to wild-type and display heightened anti-bacterial immunity [21].

The SUMO E3 ligase mutant *siz1-2*, also with lower levels of SUMO1/2 conjugates are similarly

enhanced resistant to *PstDC3000* [20,28]. On the contrary, *esd4-1* or *ots1ots2* plants with deficient

SUMO protease ESD4 or OTS1/2 functions, respectively have increased SUMO1/2-conjugates

yet display elevated SA-dependent defenses [1,6,29]. Constitutive activation of SA-dependent

defenses also occur in *SUM1/2*-overexpressing transgenic plants regardless of whether the over-

expressed SUMO isoforms are conjugation-proficient or deficient [21]. Thus, it is likely that

perturbations rather than increase or decrease in SUMO1/2-conjugates *per se* regulate immune

responses. Unlike *SUM1/2*, *SUM3* is SA-inducible, upregulated in *sum1-1 amiR-SUM2* plants, and

upon over-expression enhance the resistance to *PstDC3000* [21]. SUMO3-mediated SUMOylation of NPR1 (NONEXPRESSOR OF PATHOGENESIS-RELATED 1), a master transducer of SA-signaling is essential for defenses thus placing *SUM3* as a *bona fide* positive immune regulator [36]. Whether increased SUMO3 activities during defenses intersect to alter SUMOylome outcomes of a host remains unexplored. Role of *SUM3* in other stress responses similarly is also unknown. Evidently, perturbing host SUMOylome during biotic/abiotic stresses or through loss/gain of a specific isoform function initiates complex signaling more especially because the SUMOs themselves moderate SUMOylation proficiencies of targets. SAE2, SCE1, SIZ1, and EDS4 were identified as candidates whose SUMOylation footprints change upon stress exposure [5]. Animal studies also elegantly demonstrate 'SUMO-preference/switching' wherein the isoform choice for substrate conjugation is modulated not only by their relative levels but also by its influence on SUMOylation machinery [37–39]. Hence, it is beyond doubt that SUMO isoforms functionally intersect when the host SUMOylome-equilibrium is disturbed. However, such evidences from plant systems are completely lacking.

In this study, we utilized defense responses to various strains of *PstDC3000* as a measure to test the contribution of individual and combinatorial *Arabidopsis* SUMO isoforms in immunity. Our data suggest that *SUM1/2* function additively, but not equally, as negative regulators of SA-driven basal and TNL (Toll-Interleukin-1 receptor-like domain)-type immunity. In contrast, *SUM3* has a more positive immune role potentiating auto-immunity that occur due to loss of *SUM1* or *SUM2*. We demonstrate that accumulation kinetics of not only defense-associated markers but also of SUMOylation-associated genes are regulated by SUMO1-SUMO3 crosstalks. We further report that global change in a host SUMOylome in response to immune activation or heat-stress is also influenced by intersections of SUMO isoform activities. Overall, our results open newer avenues to unravel role of SUMO isoforms functions in maintenance and alterations of a host SUMOylome.

Results

SUM1/2 genetically function as negative immune regulators whereas *SUM3* promotes defense responses

To study individual *Arabidopsis* SUMO isoform influences on defenses, we obtained previously characterized knockout mutant lines of *sum1-1*, *sum2-1* and *sum3-1* [20,21]. Wild-type (Col-0) and mutant plants were propagated either in short day (SD) or long day conditions (LD) as indicated for respective assays. Interestingly, in all propagation regimes and especially in SD conditions, we noticed clear growth defects in *sum1-1*, but not *sum2-1* or *sum3-1* plants. The *sum1-1* mutant was developmentally dwarf with elongated leaves, reduced fresh tissue mass and increased trichomes density although with normal architecture compared to Col-0, *sum2-1* or *sum3-1* plants (Figs 1A,B; S1 Fig). Although identical *sum1-1* mutant has been utilized previously the morphological defects we observed have not been reported earlier. The reason for this discrepancy is not clear and we speculate differences in soil compositions or growth variations as possible contributing factors. Nevertheless, the phenotypic attributes of *sum1-1* were novel and we present evidences in subsequent sections that genetically link these defects solely to loss of *SUM1*.

We then investigated defense responses in the *sum* mutants utilizing the standard bacterial growth assays with virulent and avirulent *PstDC3000* strains. In Col-0, *PstDC3000* is virulent and triggers basal or PAMP-triggered immunity (PTI). In contrast, *PstDC3000* harboring *avrRps4, avrRpm1, or hopA1* effector is avirulent, triggering effector-triggered immunity (ETI), mediated by the cognate resistance gene *RPS4*, *RPM1* or *RPS6*, respectively [40]. The virulent *PstDC3000* strain we used harbors the plasmid backbone into which the avirulent effectors were cloned and hence named as *PstDC3000* (*empty vector; EV*) in subsequent

sections. Fully expanded leaves of 3-4 weeks-old SD-grown plants were infiltrated with the indicated *PstDC3000* strains and the growth of the bacteria measured at 0- and 3-days post-infiltration (dpi). When compared to Col-0, total bacterial growth was reduced almost 10-fold in the *sum1-1* and ~8-fold in *sum2-1* plants whereas *sum3-1* allowed more bacterial colonization (~8-fold higher) (Fig 1C). These results implied that *SUM1/2* and *SUM3* function as negative and positive regulators of PTI, respectively. When challenged with avirulent *PstDC3000* expressing *avrRps4* or *hopA1*, two TNL-specific ETI-eliciting effectors, lower bacterial accumulation than Col-0 persisted in the *sum1-1* and *sum2-1* plants (Fig 1D; S2A Fig). The *sum3-1* plants were hyper-susceptible to these avirulent infections. Curiously, modest but consistent difference with less bacterial accumulation in *sum1-1* than *sum2-1* was observed to virulent or *PstDC3000(hopA1)* but not to *PstDC3000(avrRps4)* challenges. Further on, increased callose deposits, a well-established defensive phenomenon [41], were more pronounced in *sum1-1* (~3-fold) and *sum2* (~2-fold) than Col-0 in response to both virulent as well as avirulent (*avrRps4*) *PstDC3000* (Fig 1E; S3 Fig). Although previous studies suggest redundant roles of *SUM1/2* in defenses [21], as our results implicate that their degree of contribution however may slightly vary. Likely indicative of enhanced susceptibility to *PstDC3000*, lower levels of callose than Col-0 accumulated in *sum3-1*. Remarkably, ETI in all plants to avirulent *PstDC3000* (*avrRpm1)* remained comparable suggesting that CNL-type responses are not affected by the loss of individual SUMO isoforms (S2B Fig). Indeed, as reported earlier *avrRpm1*-mediated HR is not affected in any *sum* mutants [21]. Overall, we identify a partial redundancy between *SUM1/2* with a concomitant antagonism to *SUM3* functions in immune responses to *PstDC3000* strains.

Basal levels of SA and SA-responsive defense markers are upregulated in *sum1-1* and *sum2-1*

Significant increases in salicylic acid, both free (SA) and glucose-conjugated (SAG), mediate

signal responses against *PstDC3000* [42]. A SUMOylation-deficient and enhanced resistant

siz1-2 plants have elevated SA/SAG due to upregulated expression of SA-biosynthesis gene

SID2/ICS1 (*SALICYLIC ACID INDUCTION DEFICIENT2/ ISOCHORISMATE SYNTHASE1*)

[21,43]. To determine whether SA/SAG perturbations reflect the immune outcomes, we

measured their relative levels in the different *sum* mutants. Remarkably, both *sum1-1* and *sum2-*

1 plants contained significantly elevated whereas the hyper-susceptible *sum3-1* had lower

SA/SAG, respectively than Col-0 (Figs 2A,B). We noted that SA/SAG levels in *sum1-1* plants

were slightly higher than *sum2-1*, perhaps indicative of its modestly higher degree of basal and

TNL-type immunity. Transcripts of *SID2/ICS1*, defense-associated markers *PR1* and *PR2,* and

PR2 protein levels were upregulated in both *sum1-1* and *sum2-1* whereas in *sum3-1* plants these

were significantly lower than Col-0 (Figs 2C,D). Likewise, increased expressions of PTI

markers *FLG22-INDUCED RECEPTOR LIKE KINASE 1* (*FRK1*) and *WRKY*

TRANSCRIPTION FACTOR 29 (*WRKY29*) [44] than Col-0 were detected in both *sum1-1* and

sum2-1 plants (Fig 2E). Relative to Col-0, *FRK1* expression remained unaltered in *sum3-1*,

whereas *WRKY29* was drastically reduced suggesting that *SUM3* promotes expression of only

a subset of PTI-markers. Accumulation of a well-known SA-responsive TNL-type R protein

SUPPRESSOR OF npr1-1 CONSTITUTIVE 1 (*SNC1*) is enhanced in several auto-immune

mutants [45,46]. Remarkably, both *sum1-1* and *sum2-1* displayed upregulated *SNC1*

expressions whereas in *sum3-1* the transcript levels remained comparable to Col-0 (Fig 2E).

Since *FRK1*, *WRKY29* and *SNC1* are SA-inducible, upregulated SA-signaling sectors likely

contribute to the enhanced basal and TNL-type immunity in *sum1-1* and *sum2-1* while these

defenses are deficient in *sum3-1*. Also taking into account that *SUM3* but not *SUM1* or *SUM2,*

is SA-inducible [21], we reasoned that enhanced resistance in *sum1-1* or *sum2-1* may be due to

elevated *SUM3* expression and its role as a positive immune regulator. Indeed, we detected ~3-

and 2-fold higher *SUM3* transcripts than Col-0 in *sum1-1* and *sum2-1*, respectively (Fig 2F).

Relative transcript levels of *SUM1* in *sum2-1*, or *SUM2* in *sum1-1* however remain unaltered.

Overall our results identify redundant but unequal roles of *SUM1* and *SUM2* in suppressing SA-

dependent defenses that include expression of PTI markers, *SNC1* and *SUM3*.

Expression of genomic copies of *SUM1* or *SUM3* complement immune response

alterations in the respective *sum* mutants

Phenotypic abnormalities of *sum1-1* plants necessitated us to genetically link the defects to loss

of *SUM1*. Although our *sum1-1* mutant is identical to earlier reports [20,21], to validate our

observations, we utilized the transgenic *His-H89R-SUM1* line that express a His$_6$-tagged

genomic *SUM1* variant (H89R) from its native promoter in a *sum1-1 sum2-1* background [5].

These plants as reported earlier not only complement the lethality of double homozygous *sum1-*

1 sum2-1 mutant but also functionally mirror wild-type SUMO1 SUMOylome changes upon

heat stress. From the *His-H89R-SUM1* plants, the *sum2-1* mutation was segregated out by

generating F2 populations upon crossing with *sum1-1*. Henceforth, we termed these plants as

sum1-1:His-H89R-SUM1. We constantly observed that growth defects we noted earlier were

always linked to homozygous *sum1-1* genotypes in absence of the *His-H89R-SUM1* transgene.

Thus, a functional *SUM1* abolishes the growth defects apparent for *sum1-1* plants (S4 Fig). We

noted that the *sum1-1:His-H89R-SUM1* plants had slightly elevated *SUM1* (but not *SUM2*)

transcripts compared to Col-0 (Fig 3A). Implicatively, as reported earlier for *SUM1* over-

expression [21], SA/SAG levels although considerably reduced in comparison to the *sum1-1*

parent, were still maintained at slightly higher levels than Col-0 (Fig 3B). Increased SA/SAG

levels in *sum1-1:His-H89R-SUM1* plants also resulted in upregulated *SUM3* transcripts and

higher PR1 proteins in comparison to Col-0 (Figs 3A,C). In pathogen growth assays using the

virulent *PstDC3000(EV)*, enhanced defenses noticeable for *sum1-1* was abrogated to Col-0

levels in the *sum1-1:His-H89R-SUM1* plants (Fig 3D). Interestingly, the avirulent

PstDC3000(hopA1) challenge although diminished enhanced resistance relative to the *sum1-1*

parent, nevertheless these plants displayed stronger defenses than Col-0 (Fig 3D). This likely

is attributed to higher than Col-0 levels of SA-regulated defense networks. These results

unambiguously support the loss of *SUM1* as the cause of enhanced resistance and phenotypic

defects in *sum1-1*.

We also generated two independent complemented lines of *sum3-1* to validate immune

deficiencies due to the loss of *SUM3*. These transgenic plants express native promoter-driven

His$_6$-StrepII-tagged genomic fragment of *SUM3* (*sum3-1:HS-SUM3*). Although owing to its

low abundance [21] the expression of His$_6$-StrepII-SUMO3 proteins remained undetectable in

immunoblots, transcripts of *SUM3* drastically reduced in the *sum3-1*, were restored to

equivalent or slightly lower (~0.5X) than Col-0 levels in Line #1 and Line #2, respectively (Fig

3A). Relative abundance of *SUM1* or *SUM2* remained unaffected in both lines. For further

assays, we therefore continued with *sum3-1:HS-SUM3*#1. We observed that deficiencies in

SA/SAG or PR1 protein accumulations inherent to *sum3-1* were reinstated to Col-0 levels in

the *sum3-1:HS-SUM3#1* plants suggesting functional complementation by the transgene (Figs

3B,C). Pathogen growth assays with either virulent *PstDC3000(EV)* or avirulent

PstDC3000(hopA1) strains regained Col-0 levels of resistance in *sum3-1:HS-SUM3#1* (Fig

3D). These data lead us to conclude that defensive deficiencies in *sum3-1* is indeed due to loss

of *SUM3*.

Constitutively active defenses in *sum1-1* and *sum2-1* caused rapid induction of immunity

To further establish that upregulated or dampened defense marker expressions in *sum1-1*, *sum2-*

1 or *sum3-1*, respectively contribute to corresponding immune outcomes we challenged these

plants with virulent *PstDC3000(EV)* or avirulent *PstDC3000(avrRps4)* strains. Leaf extracts

harvested at 6-, 12-, and 24-hpi were used to compare accumulation kinetics of *PR1/PR2*

transcripts and proteins (Fig4; S5 Fig). Basal level (0-hpi) of PR1 in Col-0, *sum2-1*, or *sum3-1*

was below detection limit whereas in *sum1-1* some enhancements were noticeable (Fig 4A). At

6- or 12-hpi although Col-0 or *sum3-1* barely accumulated sufficient PR1/PR2 in response to

both pathogen challenges, both *sum1-1* and *sum1-2* had markedly elevated levels of these

proteins (Figs 4B,C). A similar trend continued at 24-hpi wherein in Col-0, PR2 but not PR1

proteins levels, almost matched *sum1-1* or *sum2-1*. Most significantly, even at the 24-hpi *sum3-*

1 plants were deficient in accumulating wild-type levels of PR1 or PR2. Real-time accumulation

kinetics of *PR1*/*PR2* transcripts to both bacterial infiltrations in general mirrored the

corresponding protein levels although in some instances a direct correlation was not apparent

(S5 Fig). Although the reason for this discrepancy is not clear, suggested role of SUMOylation

in selective translation and/or its crosstalk with other PTMs that modulate protein synthesis or

stability may likely be the cause [47,48]. Nevertheless, these data provide strong molecular

evidence of constitutive SA-regulated defenses in *sum1-1* and *sum2-1* conferring their enhanced

resistance to *PstDC3000* strains. Contrastingly, delayed induction of defenses upon a pathogen

challenge likely makes *sum3-1* hypersusceptible.

Enhanced defenses in *sum1-1* is SA-dependent

To genetically determine whether constitutive SA-signaling routes impart increased defenses

to *sum1-1*, we generated *sum1-1sid2-1* and *sum1-1eds1-2* double mutants. A *sid2-1* plant

harbors a null mutation in *SID2/ICS1* whereas an *eds1-2* expresses a non-functional EDS1

(ENHANCED DISEASE SUSCEPTIBILITY1), a central player orchestrating SA-mediated

defenses [43,49]. In segregating F2 populations, we noted that phenotypic defects always

associated with homozygous *sum1-1* allele provided at least one functional copy of *EDS1* or

SID2 were present (data not shown). Remarkably, the reduced leaf mass apparent in *sum1-1*

was improved in *sum1-1eds1-2* and *sum1-1sid2-1* plants (Fig 5A; S1 Fig). While *sum1-1eds1-*

2 achieved wild-type mass, abolishing *SID2/ICS1* in *sum1-1* although ameliorated growth

defects but these plants still retained *sid2-1*-characterstics including smaller and paler leaves

than Col-0 (Fig 5A; S1 Fig). Interestingly, both *sum1-1sid2-1* or *sum1-1eds1-2* had wild-type

trichome densities unlike the *sum1-1* parent thus associating these defects to perturbations in

SA-regulated networks (S6A Fig).

Loss of either *EDS1* or *SID2/ICS1* remarkably abolished accumulated PR2 proteins and

upregulated *PR1*, *PR2*, *FRK1* and *WRKY29* transcripts in *sum1-1* (Figs 4A, 5B; S6B Fig).

Further on, pathogen growth assays using virulent and avirulent *PstDC3000* strains

demonstrated that *sid2-1* or *eds1-2* mutations were epistatic to *sum1-1* abolishing not only the

increased defenses but also conferring hyper-susceptibility to the respective double mutants

towards virulent *PstDC3000(EV)* or avirulent (*avrRps4-* or *hopA1*-expressing) strains (Figs

5C,D; S6C Fig). Neither *eds1-2* nor *sid2-1* mutation affected resistance towards avirulent

PstDC3000 (*avrRpm1*) which remained comparable to *sum1-1* or Col-0 (S6D Fig). As

expected, loss of *EDS1* or *SID2/ICS1* prevented accumulation of PR1 or PR2 proteins during

PstDC3000 infections (Figs 4B,C). With these assays, we identify that enhanced resistance in

sum1-1 is undoubtedly due to constitutive SA-routed defense signaling. A *sum2-1eds1-2* double

mutant, we also generated, mimicked *sum1-1eds1-2* hyper-susceptible outcomes in disease

assays (S7 Fig). Overall, our results provide several parallel lines of evidences suggesting

SUM1/2 intersections on SA-mediated immune signaling.

sum3-1 alleviates enhanced defenses in sum1/2 mutants

Since our data suggested antagonistic involvement of *SUM1/2* and *SUM3* as negative or

positive defense regulators, respectively we surmise a crosstalk between these isoforms. To

investigate this, we generated *sum1-1sum3-1* double mutant by genetic crossing. Unlike *sum1-*

1eds1-2 or *sum1-1sid2-1*, only partial restoration of *sum1-1* growth defects was observed in

sum1-1sum3-1 plants (Fig 6A; S8A,B Figs). These observations hinted that *SUM3* is mildly

responsible for *sum1-1* growth anomalies. Increased trichome densities noticed in *sum1-1*

however remained unaffected by the loss of *SUM3* (S6A Fig). Because *SUM3* affects basal SA

accumulation (Figs 2A,B), we tested defense outputs are altered in *sum1-1 sum3-1*. Elevated

SA levels in *sum1-1* were marginally reduced in *sum1-1 sum3-1* plants. Indeed, the expression

of *SID2/ICS1* remained comparable between *sum1-1* and *sum1-1 sum3-1* plants (Fig 2C).

Additionally, *PR1* or *PR2* transcripts in *sum1-1 sum3-1*, were intermediate between *sum1-1* and

Col-0. These data imply *SUM3* does not affect SA-biosynthesis but modulates positive

feedback loop of SA-signaling wherein it intersects with *SUM1* role as a transcriptional

repressor of *SID2/ICS1*, *PR1* or *PR2*. As is therefore expected, loss of *SUM3* reduced PR1 or

PR2 levels in *sum1-1* (Fig 4A). Not the least, we also demonstrate that increased SNC1

accumulation in *sum1-1* plants although reduced in *sum1-1 sum3-1*, are still maintained higher

than Col-0 levels (Fig 6B).

In pathogen growth assays, either virulent *PstDC3000(EV)* or avirulent *PstDC3000 (hopA1)*

strains accumulated to intermediate levels in *sum1-1 sum3-1* leaves, particularly higher than

sum1-1, but significantly lower than in Col-0 or *sum3-1* (Fig 6C; S8C Fig). Interestingly, wild-

type level of resistance was observed in *sum1-1 sum3-1* to the avirulent effector *avrRps4* (Fig

6D). Growth of avirulent *PstDC3000 (avrRpm1)* was not affected in *sum1-1sum3-1* and

remained comparable in all plants (S8D Fig). As investigated earlier, in response to virulent

and avirulent *PstDC3000(avrRps4)* infections and unlike *sum1-1*, *sum1-1 sum3-1* remained

deficient in induction of PR1 or PR2 at 6-hpi (Figs 4B,C; S9A,B Figs). However, we observed

that at later time points (12- and 24-hpi) loss of *SUM3* did not affect the rapid induction of

PR1/PR2 in *sum1-1*. This is a sharp contrast to *sum3-1* plants which even at 24-hpi accumulated

very less PR1/PR2 proteins. Thus, although *SUM3* is deemed essential for upregulation of

PR1/PR2 upon a pathogen attack, *sum1-1* plants eventually overcome this requirement. Taken

together, we principally support that *SUM3* promotes *PR1/PR2* transcription downstream of SA

as suggested earlier [21]. We provide further molecular evidence that *SUM1* suppression of

defenses likely is upstream of SA and may involve transcriptional repression of *SID2/ICS1* and

its subsequent consequences on SA-responsive markers such as *PR1*, *PR2*, *FRK1*, *WRKY29* and

SNC1 among others.

Adjacent chromosomal arrangement of *SUM2* and *SUM3* in Col-0 genome suggest that they

likely arose by a tandem duplication event [18]. Hence, unlike a *sum1-1 sum3-1* double mutant,

a *sum2-1 sum3-1* is difficult to obtain by genetic crossing. To test whether *SUM3* also intersects

on *SUM2* function as a negative immune regulator, we generated *amiR-SUM2 sum3-1* plants

by crossing the two parental lines. The *amiR-SUM2* plants reported earlier contains specific

knockdown of *SUM2* and mimics the null mutant *sum2-1* [21]. *SUM1* or *SUM3* transcripts

remain unaffected in these lines. Plants homozygous for *sum3-1* and containing *amiR-SUM2*

transgene have significantly downregulated *SID2/ICS1, PR1* and *PR2* levels in comparison to

a *sum2* mutant (Fig 2C). These transcripts however are still elevated than Col-0 plants

implicating that *SUM3* functions impinge partially on *SUM2* role as a transcriptional repressor.

This was also supported in pathogen challenges wherein growth of both virulent and avirulent

PstDC3000 strains were intermediate between resistant Col-0 and hyper-resistant *sum2* mutant

(S7B,C Figs). Taken together, whether the functional antagonism in immune regulatory roles

of *SUM1/2* and *SUM3* affect immune outcomes in plants.

SUMO3, but not SUMO1, can form non-covalent homo-dimers

Conjugation proficiencies *in planta* of SUMO isoforms are clearly different. Upon SA application, drastic increase in SUMO1/2 conjugates are observed whereas target modifications by SUMO3 is barely detectable even though *SUM3* expression, unlike *SUM1/2* is SA-inducible [6,21]. A distinct difference in the protein sequence of these isoforms is the presence of a predicted SIM motif in SUMO3 but not SUMO1/2 (S10 Fig). We utilized Bi-molecular Fluorescence Complementation (BiFC) assays to test homo- and hetero- non-covalent associations between SUMO1 and SUMO3. The BiFC vectors introduced via *Agrobacterium*-mediated transient transformation of *N. benthamiana* leaves expressed split YFP fusions of either SUMOylation-proficient (GG) or -deficient (AA) SUMOs wherein the C-terminal di-glycine (GG) residues were kept intact or mutated to di-alanine (AA), respectively (Fig 7A). Co-expression of only SUMO1GG/SUMO1GG, but not SUMO1GG/SUMO1AA or SUMO1AA/SUMO1AA showed reconstitution of the split YFP protein. Although positive BiFC indicate non-covalent interactions among protein partners, the lack of fluorescence in SUMO1AA/SUMO1AA suggests that SUMO1GG/SUMO1GG combinations and not homo-oligomers and likely reflect covalent polySUMO1-conjugates wherein the split fluorescent fusion proteins are in allowed proximity for reconstitution. Remarkably, all BiFC combinations of SUMO3 with itself, regardless of GG- or AA-forms, showed YFP fluorescence. Since SUMO3 lacks poly-SUMOylation properties [22], we reason that the observed associations are non-covalent SUMO3 oligomers. To test whether SUMO3 binds SUMO1 non-covalently, combinations of SUMO1GG or AA were co-expressed with SUMO3GG or AA. While SUMO1GG showed clear fluorescence when expressed with either SUMO3GG or AA, similar combinations of SUMO1AA did not. These observations are suggestive that only a conjugable-SUMO1 likely present on a target may achieve permissible molecular proximity *in vivo* to a

SUMO3 that may be either bound to a SIM (as in SUMO1GG/SUMO3AA pair) or attached

covalently to the same target at a separate SUMOylation motif (as in SUMO1GG/SUMO3GG

pair). Evidences from human cell lines demonstrate that ortholog of *Arabidopsis* SUMO3,

HsSUMO1 'cap' poly-SUMO chains of the SUMO1 ortholog HsSUMO2 [50,51]. Whether

Arabidopsis SUMO3 regulates poly-SUMO1/2 chain lengths remain a promising possibility to

explore further. Since no interaction was detected with SUMO1AA/SUMO3AA, it is clear that

SUMO3 and SUMO1 do not interact non-covalently. To validate further, we performed *in vitro*

binding assays with tagged recombinant SUMO1AA/SUMO3AA proteins expressed in *E. coli.*

Enrichment of His-SUMO1AA via Ni^{2+} beads failed to co-elute either Strep-SUMO1AA or

Strep-SUMO3AA suggesting that SUMO1-SUMO1 homo- and SUMO1-SUMO3 hetero-

dimers cannot form *in vitro* (Fig 7B). Homodimers of SUMO3AA however were detected by

the presence of Strep-SUMO3AA in His-SUMO3AA-enriched eluates. These results support

our hypothesis that BiFC interactions observed for SUMO1GG with SUMO3GG/AA may

likely suggest close proximity, but not direct binding, of SUMO1 and SUMO3 *in planta*.

SUMO3 regulates SUMO1/2 conjugation efficiencies upon various stress exposures

With the above results, it is encouraging to speculate that SUMOylation efficiencies may be

modulated by intersecting SUMO1-SUMO3 functions. Since *SUM3* partially modulates *sum1-*

1 defenses, we investigated dynamic changes in SUMO1/2-SUMOylome in Col-0, *sum1-1*,

sum3-1, or *sum1-1 sum3-1* upon a pathogen challenge. Leaf tissues either mock-treated or

infiltrated with *PstDC3000*(*EV*), were collected at 24-hpi and processed for selective qPCRs or

immunoblots for SUMO1/2-conjugates. The polyclonal anti-SUMO1 antibody we used cannot

distinguish SUMO1/2 isoforms and the lack of anti-SUMO3 antibodies prevented us for

analyzing SUMO3-conjugates. We observed that while a pathogen exposure caused remarkable

increase in SUMO1/2-conjugates in Col-0, for *sum1-1* these was considerably lower (Fig 8A).

These results suggested that SUMO1- rather than SUMO2-conjugates are not only more

prevalent but also undergo massive increment upon pathogen treatment. Surprisingly, the

conjugates are also distinctly lower in uninfected *sum3-1* and are considerably less upregulated

than Col-0 in response to the pathogen infection. As is expected, in *sum1-1 sum3-1* extracts

barely any SUMO-conjugates are detectable pre- or post-infection. With these results, we

identify clear intersection of SUMO3 functions on SUMO1/2-conjugation efficiencies and its

perturbations during a pathogen attack. We also observed that both *sum1-1:His-H89R-SUM1*

and *sum3-1:HS-SUM3* plants achieved Col-0 levels of SUMO1/2-cojugates upon

PstDC3000(*EV*) infection thus further validating the complete functional complementation of

sum1-1 or *sum3-1*, respectively (S11 Fig).

To determine whether changes in conjugation efficiencies are due to differential expression of

SUMO isoforms and/or SUMOylation-associated genes, we investigated their relative

expressions in Col-0 and the *sum* mutants (Figs 8B,C; S11B Fig). We observed that *SUM3*, but

not *SUM1/2*, was upregulated upon a *PstDC3000*(*EV*) challenge. This is in accordance with

SUM3 being SA-inducible [21]. Additionally, we noticed that the enhanced *SUM3* expression

in *sum1-1* was further boosted in pathogen-exposed samples. Curiously, for the investigated

SUMOylation-associated genes although *sum1-1* mutation did not significantly alter their basal

expression levels, several of these (*SAE2*, *SIZ1*, *HYP2*, *ESD4*, *ELS1*, and *OTS1*) were down-

regulated in *sum3-1* plants. Thus, the modest reduction in global SUMOylome in *sum3-1* we

observe may be attributed to *SUM3*-dependent modulation of expression of these genes.

Interestingly, *sum1-1 sum3-1* plants showed slight upregulation of *SAE2*, *SCE1*, *SIZ1*, *HPY2*,

ELS1 and *OTS2* in mock-treatment supporting our earlier claim that *sum1-1* plants with

constitutively active defenses and elevated SA levels, overcome the requirement of *SUM3* for

the expression of these genes. We also noted that upregulation of *SAE2*, *SIZ1*, *ELS1* or *OTS1*

that occurred only modestly in Col-0 upon *PstDC3000*(*EV*) infection were further aggravated

in *sum1-1*, but intermediate in *sum3-1* (Figs 8B,C). These enhancements are maintained in

sum1-1 sum3-1 plants and are likely *SUM3*-independent. Remarkably, *sum3-1* plants down-

regulated the expression of *SUM1* and *SUM2* only upon pathogen-treatment highlighting that

the positive immune function of *SUM3* may involve transcriptional repression of negative

defense regulators such as *SUM1/2*. Overall, our data lead us to speculate that immune

responses recruit *SUM1-SUM3* crosstalks to modulate the expression of these genes at a

transcriptional level thereby influencing host SUMOylome changes.

SAE2, SCE1, SIZ1 or EDS4 are direct SUMO1 targets and both SUMO1/3 interact *in planta*

with SCE1 and SIZ1 [5,52]. These led us to test whether SUMO1-SUMO3 intersections directly

influence the efficiency of SUMO1/2-conjugate formation through modulation of

SUMOylation-associated protein functions. We utilized the *E. coli* SIZ1-independent

Arabidopsis SUMOylation reconstitution system [53]. Co-expression of His-SUMO1, and

Strep-SUMO3 (GG or AA) along with SCE1, and SAE1/2 showed modest but clear increase in

SUMO1-conjugates in comparison to extracts that lacked Strep-SUMO3 (S12 Fig).

Remarkably, a more dramatic increase in SUMO3-conjugates when co-expressed with either

Strep-SUMO1 GG or AA was also noted. From these results, it is undeniably clear that SUMO1

and SUMO3 have a mutually beneficial role in reciprocal SUMOylation process likely through

modulation of SCE1 or SAE1/2 functions. The significant decrease in SUMO1/2 conjugates in

absence of *SUM3* (Fig 8A) provides reasonable endorsement to our hypothesis.

SUMO1/2-conjugates are enhanced in response to heat shock [20,21]. To test whether *SUM3*

role intersect on these responses, we subjected Col-0, *sum1-1*, *sum3-1*, or *sum1-1 sum3-1* plants

to heat shock and analyzed SUMO1/2-conjugates as well as relative expression levels of

SUM1/2/3 and several previously characterized heat stress-responsive genes [54] (S13 Fig). In

accordance with [20], we observe strong induction of SUMO1/2-conjugates in heat-treated Col-

0 extracts (S13A Fig). These conjugates were low in heat-stressed *sum1-1* plants reinstating

that SUMO1-mediated SUMOylation is most predominant during heat stress. Surprisingly, in

sum3-1 considerably reduced conjugates than Col-0 accumulated upon heat shock. We

especially noted that unlike *PstDC3000* challenges, *SUM1/2* transcripts were significantly

upregulated by heat-exposure, while *SUM3* induction was almost negligible (S13B Fig).

Further on, transcriptional upregulation of *SUM1/2* was *SUM3*-independent. Therefore,

reduced accumulation of SUMO1/2-conjugates in heat-treated *sum3-1* suggests that *SUM3*

affects SUMO1-SUMOylation efficiencies at a post-transcriptional level. With these results,

we infer that although different physiological stresses may cause apparently similar effects on

global SUMO1/2-conjugates, the responsive routes to condition these are not only distinct but

also stress-specific. Interestingly, increased expression of several heat stress-responsive

markers such as *HsfA2*, *HSP22.0-ER* that are upregulated in Col-0 upon heat-treatment, were

hyper-elevated in *sum1-1* suggesting *SUM1* suppresses their expression (S13C Fig). In *sum3-1*

plants, their fold-induction upon heat stress was relatively lower than Col-0 implying that *SUM3*

is partially responsible for their upregulation. Induction of other tested markers such as

HSP18.2 or *HSP23.5-P* were unaffected in *sum1-1* or *sum3-1* plants (S13D Fig). Taken

together, our results clearly identify *SUM1-SUM3* crosstalks in responses to multiple stresses.

Discussion

Functional overlaps between *Arabidopsis* SUMO1 and SUMO2 isoforms, likely prevent

noticeable phenotypic defects in the individual mutants under non-stressed conditions

[17,20,21,23]. Interestingly, reduction in SUMO-conjugates both basally as well as upon heat-

stress is more prominent in *sum1-1* than *sum2-1* suggesting a *SUM1* predominance in these

responses [20]. Our observations therefore of phenotypic defects in *sum1-1* but not *sum2-1*

plants may reflect a similar developmental process preferentially regulated by *SUM1*. We

convincingly establish that constitutive upregulation of SA-regulated networks is the primary

cause of these defects in *sum1-1* (S6A Fig). This is unlike *siz1-2* where the associated growth

abnormalities are EDS1- or SID2/ICS1-independent [55]. We deduce that reduction in global

SUMO1/2- conjugates apparently similar between *siz1-2* and *sum1-1* affect downstream

responses differently. At one instance, loss of *SIZ1* may affect SUMO-conjugation for all

isoforms, whereas the *sum1-1* clearly restricts only SUMO1 functions. Previous studies propose

SA antagonism or jasmonic acid (JA) promotion in increased trichome formation [56]. We

reason that while direct application of SA or JA may affect trichome production as reported,

the loss of *SUM1* whose substrates include both JA and SA signaling regulators such as SIZ1,

TPL, or JAZs among others may impact trichome density through a more complex SA-JA

crosstalk [5,57]. Further investigations into relative SA-JA signaling routes perturbed in *sum1-*

1 may unravel this mystery.

In context of defense responses, our data provide substantial molecular support not only to

individual *SUM1/2* or *SUM3* but also to their intersecting contributions as negative or positive

regulators, respectively of anti-bacterial basal and TNL-specific immunity. Firstly, bacterial

accumulations are strongly reduced in both *sum1-1* and *sum2-1*, with *sum1-1* displaying

stronger immunity than *sum2-1*, for both virulent as well as avirulent *PstDC3000*(*hopA1*)

infections. Although comparable immunity between *sum1-1* and *sum2-1* is observed for

avirulent *PstDC3000*(*avrRps4*) challenges, this likely is due to relatively weaker ETI responses

elicited by *AvrRps4* in comparison to *HopA1*-expresssing *PstDC3000* (compare Fig 1D and

S2A Fig). In contrast to *sum1/2* mutants, immunity in *sum3-1* is compromised implicating

SUM3 is essential for these defenses. Secondly, we demonstrate that impairment of SA-

signaling routes due to loss of *EDS1* (*eds1-2*) or *SID2/ICS1* (*sid2-1*) abolish enhanced resistance

in *sum1-1* and *sum2-1* plants providing support to genetic placement of *SUM1/2* as suppressors

of these defenses [21] (Fig 8D). In this pathway, *SUM3* partially delegates SA-defenses likely

through positive feedback mechanisms. Indeed, both *sum1-1* and *sum2-1* plants accumulate

increased SA than Col-0, while in *sum3-1*, these are relatively lower. Correspondingly,

transcripts of SA-biosynthesis *SID2/ICS1* and responsive markers such as *FRK1*, *PR1*,

WRKY29, and *SNC1* are upregulated in *sum1-1* or *sum2-1*, and lower in *sum3-1*. Accumulation

kinetics of PR1/PR2 proteins or transcripts (Fig 4; S5 Fig) further validate primed or deficient

SA-mediated defenses in *sum1-1* or *sum2-1* and *sum3-1*, respectively. The modest difference

we note between *sum1-1* and *sum2-1*, in the upregulation of some of these SA-inducible

markers, is suggestive of additive but unequal roles of *SUM1* and *SUM2* as negative immune

regulators. Only ~17% common targets were identified between human SUMO1 and SUMO2

isoforms suggesting that they are partially redundant at best [58]. Whether SUMOylation-

targets distinct between *Arabidopsis* SUMO1 and SUMO2 isoforms affect immune amplitudes

differentially awaits further studies.

We provide the first *in planta* evidence of partial SUMO3 moderation on SUMO1/2 functions

in biotic, abiotic and developmental responses. Introducing *sum3-1* mutation not only attenuates

reduced tissue mass but also subdues enhanced resistance of *sum1-1* supporting *SUM3* role in

promoting defenses downstream of SA. Thus, elevated SA and *PR1*, *PR2*, or *SNC1* expressions

are partially dampened in either *sum1-1* or in *amiR-SUM2* plants when *SUM3* is mutated (Fig

2). Intersections of *SUM3* functions on SUMOylation-proficiencies, are also best noted on

SUMO1/2-conjugate intensities in response to different stresses. Although, enhancement upon

heat shock or SA-treatments have been previously reported [5,6], we first report their

upregulation in *PstDC3000* exposures. Induction of several SUMOylation-associated genes

such as *SAE2*, *SIZ1*, *ELS1*, or *OTS1* during *PstDC3000* challenges are influenced by *SUM1-*

SUM3 intersections and contribute to the corresponding defense outcomes (Fig 8). Recently,

the activity of TPR1, a transcriptional co-repressor involved in suppression of negative defense

regulators *DND1/2* was reported to be regulated in a partial SA-dependent manner via

SUMOylation by SIZ1 [59]. SUMOylation-deficient TPR1 not only is more enhanced in

repressing *DND1/2* expressions, but also its over-expression cause enhanced resistance than the

SUMOylation-proficient TPR1. Overall, our results raised the possibility that increased

SUMO1/2 conjugates may reflect negative feedback mechanisms to maintain immune

responses as transitory thus regulating *SUM3* functions in promoting defenses.

Functional intersections among SUMO isoforms are anticipated at multiple post-transcriptional

events. Amino acid distinctions at key conserved positions among the SUMO isoforms

influence their *in vivo* conjugation proficiencies [60] (Schematically summarized in S10 Fig).

Aspartate63 (D^{63}) in *Arabidopsis* SUMO1 (D^{75} in SUMO2) is replaced by Asparagine63 (N^{63})

in SUMO3, reducing its relative interaction strength than SUMO1 with SCE1, thus lowering

its poly-SUMO formation efficiencies. Other residues also diverged in SUMO3 weaken

thioester-bond formation and interaction with SAE1. Whether *SUM3* upregulation during

defences improve these propensities and hence alters host SUMOylome outcomes although

remains unknown, evidential support is obtained from several studies. Mutants with decreased

SUMOylation including *siz1-2* have increased SCE1 protein abundance [15,20,61]. And *siz1-2*

impairment in accumulating SUMO-conjugates during heat stress, is completely recovered in

pial1 pial2 siz1-2 plants [8]. Considering PIAL1/2 improves SCE1 ability to form otherwise

less efficient polySUMO3 chains, defense responses with upregulated *SUM3* likely introduces

competition between SUMO1 and SUMO3 for substrate polySUMOylation thereby altering

their fates via respective isoform-specific SUMO-targeted E3 Ubiquitin ligases (STUbLs) [8].

Indeed, *pial1/2* double mutants although less stress-tolerant, have upregulated levels of proteins

related to biotic stress [8,15]. In *sum1-1*, the reduced global SUMO1/2-conjugates upon

pathogen or heat-shock challenges taken together with elevated *SUM3* transcripts is suggestive

of this occurrence and need to be explored further.

As substrates of Arabidopsis SCE1 and ESD4, candidates predominantly involved in RNA-

related processes such as nucleocytoplasmic transport, splicing, or turnover and chromatin-

modification including transcriptional activation/repression-associated proteins were identified

[62]. A majority of these candidates possessed predicted SUMOylation motifs and were

covalently modified by both SUMO1 or SUMO3. In *Arabidopsis*, a SUMOylation-proficient

SUMO1 is essential to covalently charge SCE1 at the catalytic site, promote its association with

SIZ1, and form the ternary complex (SUMO-SCE1-SIZ1) [52]. Interestingly, the subcellular

localization of this complex is determined by the specific SUMO isoform bound non-covalently

at the SIM site distinct from the catalytic pocket of SCE1. Since SCE1 possess ability to

distinguish substrates for SUMOylation [9], selection of substrates and formation of

polySUMO chains therefore may be affected by fluctuations in relative SUMO1/3 levels, as

has been previously suggested in animal studies [63,64]. Our data showing reciprocal

improvements in SUMOylation efficiencies *in vitro* regardless of conjugation-proficiencies of

the influencing SUMO isoform may suggest towards one such consequence of SUMO1-

SUMO3 crosstalks (S12 Fig). Likewise, constitutive activation of defenses due to over-

expression of either conjugation-proficient (GG) or -deficient (ΔGG) SUMO isoforms may at

least be partially attributed to this phenomenon [21].

SUMOylation machineries such as SAE2, SIZ1 and ESD4 also bind SUMOs non-covalently

owing to their intrinsic SIMs [12,65]. It is postulated that spatio-temporal regulations of

SUMOylation/de-SUMOylation coordination primarily by SIZ1 and ESD4 regulate steady-

state levels of host proteins conjugated to SUMOs which surprisingly are fewer in numbers [2].

Dramatic changes in this minimal SUMOylome across various stress conditions reveal that

instead of newer targets undergoing covalent modifications by SUMOs, SUMOylation levels

on prior-SUMOylated proteins pools are more altered [66]. Likely, functional inactivation of

SUMO proteases especially noted post-stress in mammalian systems [67–69] and enhanced

pool of SUMOylated pool SIZ1 [66] coordinate to achieve this. Taking this into account, a host

SUMOylome output is likely to be influenced by relative changes in SUMO isoforms. Indeed,

we demonstrate that transcripts of *SIZ1*, *EDS4* or *SAE2* are up-regulated basally in *sum1-1*, and

in both Col-0 or *sum1-1* upon pathogen challenge in a *SUM3*-dependent manner (Fig 8). Hence

the requirement of *SUM3* to maintain and adjust optimal level of SUMO1/2 SUMOylome is

clearly evident. Non-covalent associations with SUMOs also affect chromatin architecture via

their interaction with DNA methyltransferases and demethylases [11]. Expression of

FLOWERING LOCUS C (FLC), a central transcription factor of flowering is dynamically

regulated by DNA methylation status while the protein function is modulated by HPY2, SIZ1

and SUMO proteases [70,71]. FLC regulation presents a classical example of this versatility of

SUMO-influences on a developmental process.

Stimulus-driven SUMO isoform switches and subsequent fate of substrates have been widely

documented in animal systems [37–39]. The mammalian GTPase activating protein RanGAP1

although is equally modified by SUMO1/2/3 *in vitro*, conjugation *in vivo* to SUMO1, but not

SUMO2, imparts more stability from isopeptidases thereby facilitating its association with

Nup358 [39]. Similarly, HDAC1 is targeted for degradation upon SUMOylation by SUMO1,

but not SUMO2, in cancerous cell lines thus potentiating invasive properties of the tumour [72].

Specific SUMO-proteases also regulate stimulus-dependent SUMO isoform switching [73].

Upon arsenic exposure, SUMO2 at Lys[65] of PML (Promyelocytic leukemia protein) is replaced

with SUMO1 to direct its ubiquitinylation. Clearly with the isoform selectivity regulated at

multiple levels, a host SUMOylome undergoes dynamic changes in response to physiological

perturbations. Evidences on SUMO isoform switches however are completely lacking from

plant systems. Nonetheless several reports suggest the likelihood of this phenomenon. The

effector XopD from *Xcv* is a SUMO3-specific isopeptidase [17]. The replication protein AL1

from Tomato Golden Mosaic Virus (TGMV) and the RNA-dependent RNA polymerase NIb

from Turnip mosaic virus (TuMV) interact with and require SCE1 for replication [74,75].

While our results undoubtedly place SUMO3 as a strong contributor in regulating dynamics

changes in a host SUMOylome, the loss of *SUM3* in many *Brassicaeae* and other higher

eudicots still remain an enigma [18]. Whether functions of *SUM3* has been incorporated in

SUM1/2 roles in these plants remains to be explored further. We foresee the immediate need of

a plant system that would facilitate enrichment via distinct affinity tags on individual SUMO

isoforms in order to obtain evidences of SUMO preference, switches and intersections. To

summarize, our data lay the foundation on functional impingements of *Arabidopsis* SUMO

isoforms and their adjustments on global SUMOylome in response to physiological changes.

Methods

Plant materials and growth conditions

Arabidopsis thaliana mutant lines *sum1-1* (SAIL_296_C12), *sum2-1* (SALK_029775C) and

sum3-1 (SM_3_2707/SM_3_21645) were obtained from Arabidopsis stock centre

(https://www.arabidopsis.org). Seeds of *amiR-SUM2 sum3-1* were generated by Dr. Harrold

van den Burg. All plants were grown at 22^0C with 70% humidity under Long Days (LD; 16 h:

8 h, light: dark) or Short Days (SD; 8 h: 16 h, light : dark) having light intensity 100 μmol μm⁻

²s⁻¹ light. Specific growth conditions are indicated in respective legends. To generate

combinatorial mutants, *sum1-1* was genetically crossed with *eds1-2*, *sid2-1* or *sum3-1* mutant

plants. The *sum2-1* mutant was similarly crossed with *eds1-2* to generate *sum2-1 eds1-2*.

Double mutants (*sum1-1 eds1-2, sum1-1 sid2-1* and *sum1-1 sum3-1*) were identified in F2

population by PCR based genotyping. To generate *sum1-1:His-H89R-SUM1*, *His-H89R-SUM1*

[5] was crossed with *sum1-1*. Segregating F2 populations were PCR-based genotyped for the

presence of homozygous *sum1-1*, wild-type *SUM2* and homozygous copies of *His-H89R-SUM1*

transgene. All primers used are listed in S1 Table.

Trichome visualisation and quantification

For trichome visualization, leaves from 4-weeks old SD-grown plants were harvested and

washed in Acetic acid: Ethanol (1:3) solution for overnight to bleach all pigments then

visualised under bright field in a fluorescence microscope. The images were captured as 4848

X 3648 pixels from 4.1 mm² leaf area. The trichome numbers were counted in 10 randomly

selected images. Experiment was repeated twice with similar observations.

Salicylic acid (SA) measurements

Free salicylic acid (SA) and glucose-conjugated SA (SAG) measurements were done using the

Acinetobacter sp. ADPWH_*lux* biosensor system [76]. In brief, 100 mg leaf tissues were

collected and frozen in liquid nitrogen. The tissue was homogenized in 250μl of acetate buffer

(0.1 M, pH 5.6). Samples were centrifuged at 12000 rpm for 15 minutes. One aliquot (100μl)

of the supernatant was used for free SA measurements, and another was incubated with 6U of

β-glucosidase (Sigma-Aldrich) for 90 min at 37°C for total SA measurement. 20μl aliquot of

each plant extract were added to 50μl of secondary culture of *Acinetobacter* sp. ADPWH_*lux*

(OD_{600} of 0.4) with additional 60 μl of LB. Standard SA solutions (prepared in *sid2-1* extract)

were also taken to generate standard curve to calculate the amount of SA present in samples.

Plate was incubated at 37°C for 1 h, and luminescence detected using the POLARStar Omega

Luminometer (BMG Labtech). Data shown is representative of three biological replicates with

SD.

***In planta* assays for bacterial growth and kinetics of defense-associated marker**

expressions

Bacterial growth assays were performed according to [46]. Briefly, *PstDC3000* strains were

infiltrated with a needleless syringe at a density of 5×10^4 cfu ml^{-1}into fully expanded rosette

leaves of 3-4-weeks-old SD grown plants. Leaf discs harvested from infiltrated area were

macerated in 10mM $MgCl_2$, serially diluted and plated onto selective medium plates. Bacterial

growth was determined at 0 and 3-days post-infiltration. Bacterial infiltrations for kinetics of

defensive markers were carried as above except a 5×10^6 cfu ml^{-1} bacterial inoculum was used.

Infiltrated samples harvested at 6, 12 and 24 hpi were processed separately for total RNA

extraction and qPCR analysis or for immunoblots with indicated antibodies.

Callose deposition assay and image analysis

The callose deposition assays were performed according to [77]. In brief, 4-week-old leaves of

SD grown Col-0, *sum1-1, sum2-1* and *sum3-1* plants were infected with the indicated

PstDC3000 strains. At 24 hpi, 3-4 random infiltrated leaves were harvested and washed first in

Acetic acid: Ethanol (1:3) solution for overnight to bleach all pigments and then washed for 30

min in 150 mM K_2HPO_4 solution. The leaves were then incubated in dark for 2 hours in 150

mM K_2HPO_4 containing 0.01% aniline blue in a 16-well tray. Samples were then embedded in

50% glycerol and observed under a Nikon fluorescence microscope with DAPI filter. Images

were analysed using IMARIS 8.0 software for quantifying the number of callose deposits/ mm^2

leaf area.

RNA extraction and gene expression analysis by qRT-PCR

Total RNA extraction and cDNA synthesis was performed as described later. All qPCR primers

used in this study are listed in S1 Table. qPCRs were performed in QuantStudio 6 Flex Real-

Time PCR system (Applied Biosystems) with 5X HOT FIREPol$^®$ EvaGreen$^®$ qPCR Mix Plus

(ROX) (Solis BioDyne) according to the manufacturer's instructions. All qPCR experiments

were replicated thrice with three biological and technical replicates (n=3). *SAND* (At2g28390)

expression was used as internal control [46]. Relative expression was calculated according to

the PCR efficiency$^{\text{-}\Delta\Delta Ct}$ formula. Expression differences were normalised to Col-0 and plotted

as fold change.

Protein extraction and western blotting

For immunoblotting, leaf tissues collected at indicated time points/treatments were

homogenised in protein extraction buffer [50 mM Tris HCl (pH 8.0), 8M Urea, 50 mM NaCl,

1% v/v NP-40, 0.5% Sodium deoxycholate, 0.1% SDS and 1 mM EDTA] containing 20 mM

N-ethylmaleimide (NEM), 1X plant protease inhibitors cocktail (Sigma Aldrich) and 2% w/v

Polyvinylpyrrolidone (PVPP). The homogenates were clarified by centrifugation, mixed with

2X Laemmli buffer, proteins separated by SDS-PAGE and then transferred onto polyvinylidene

fluoride (PVDF) membrane by wet transfer method. The membrane was blocked with 5% non-

fat skim milk and western blots performed with indicated primary antibodies [anti-SNC1

(Abiocode), anti-PR1 or anti-PR2 (Agrisera), or anti-SUMO1 (Abcam), or anti-Actin C3

antibodies (Abiocode)] in 1X TBST at 4^{0}C for overnight. Comparable protein loading was

determined by Ponceau S staining of Rubisco subunit. Blots were washed thrice with 1X TBST

and then incubated at RT for one hour with appropriate horse-radish peroxidase (HRP)-

conjugated secondary antibodies. The blots developed using ECL[TM] Prime western blotting

system (GE Healthcare) and visualised in ImageQuant[TM] LAS 4000 biomolecular imager (GE

Healthcare).

SUMOylome changes in response to SA or heat shock treatments

755   For salicylic acid (SA)-induced SUMOylation changes, extracts were obtained from 2-weeks

756   old seedlings treated with 2 mM SA or buffer alone for 1 hour [6]. For heat shock treatments,

757   2-weeks old seedlings were incubated at 37^0C or at RT for 30 minutes. Total protein extracts

758   were immunoblotted as described earlier.

Construction of Plasmid clones

To generate cDNA clones of Arabidopsis *SUM1* and *SUM3* genes, total RNA isolated from

Col-0 plants (RNAiso Plus; Takara) was reverse transcribed (iScript[TM] cDNA Synthesis Kit;

Bio-Rad) according to manufacturer's instructions. Specific *SUM1* and *SUM3* (GG and AA

forms) sequences were amplified (Phusion High-Fidelity DNA Polymerase; ThermoFisher)

from the cDNA using the primers listed (S1 Table). PCR products were cloned into the Gateway

entry vector *p*DONR201 and subsequently into *p*MDC-cCFP and *p*MDC-nVenus (BiFC

destination vectors; [40] using the Clonase[TM] Recombination system (ThermoFisher).

Confirmed BiFC clones were then electroporated into *A. tumefaciens* GV3101 strain.

For dimerization studies, cDNAs of *SUM1AA* or *SUM3AA* were cloned as EcoRI-SalI fragment

into *p*ASK-IBA16 vector (N-terminal Strep-tag II affinity tag; Neuromics). For *in vitro* isoform

influence on SUMOylation assays *SUM1GG* or *SUM3GG* were cloned as EcoRI-SalI fragment

into *p*ASK-IBA16 vector. These clones were named accordingly as Strep-SUMO1GG, Strep-

SUMO1AA, Strep-SUMO3GG, or Strep-SUMO3AA, respectively. For His-tagged versions,

*p*CDFDuet vector harbouring *SUM1GG* (*p*KT-973), *SUM1AA* (*p*KT-1017), *SUM3GG* (*p*KT-

975), or *SUM3AA* (*p*KT-1788) were used [53]. These plasmids were a kind gift from Prof.

Katsunori Tanaka, Kwansei Gakuin University, Japan).

Cloning and generation of *SUM3p-His-StrepII-SUM3g* transgenic plants

Genomic DNA extracted from Col-0 plants were used for PCR of a ~1.3kbp genomic fragment

with KpnI-SUM3p For/XbaI-SUM3-UTR Rev primers. Using the amplicon as a template, two

independent PCRs were performed by using KpnI-SUM3p For/His-StrepII Rev or His-StrepII

For/ XbaI-SUM3-UTR Rev primers combinations respectively. Individual PCR amplifications

were used for overlap PCR with KpnI-SUM3p For/XbaI-SUM3-UTR Rev primers. The product

was cloned into XbaI-KpnI site of the binary vector *p*BIB-Hyg [78]. Generated clones were

sequenced and confirmed. The binary vector was introduced in *Agrobacterium* strain GV3101

via electroporation. Pool of *sum3-1* plants were transformed via floral-dip method [79].

Transgenic *sum3-1:HS-SUM3* plants were selected on Hygromycin containing medium,

propagated through T3 generations to identify lines containing single locus but homozygous

sum3-1:HS-SUM3 transgenes. Subsequently, the plants were used for assays as indicated.

Bimolecular Fluorescence Complementation (BiFC) assays

All BiFC assays were performed according to [40]. Briefly, *Agrobacterium* cells harbouring

the indicated BiFC vectors were induced with 150 μM Acetosyringone for 4 hours, equal

bacterial density suspensions of desired cCFP and nVenus BiFC combinations made and then

infiltrated in leaves of 4-weeks old *N. benthamiana* plants. At 48-hpi, a small section of

infiltrated area was imaged under a SP8 Leica confocal microscopy system using FITC filter

(488-nm Argon Laser).

In vitro SUMO binding assays

SUMO-binding assays were performed with recombinant protein expressed in *E. coli.*

Expression was induced with 200 μg/L Anhydrotetracycline (for *p*ASK-IBA16 clones) or with

0.5 mM IPTG (for *p*CDFDuet clones) in BL21 (DE3) for overnight at 25^0C. Cell pellet was

resuspended in lysis buffer (50 mM NaH_2PO4, 300 mM NaCl, 10 mM Imidazole; pH 8.0) and

sonicated. Equal volumes of His- and Strep-II tagged-SUMO combination supernatants were

mixed and incubated at 4^0C with rotation for 2-3 hours to allow binding. Ni^{2+}-NTA beads

(Qiagen) were added to the mix and incubated further for 2 hrs. His-tagged protein pull down

was performed under native conditions according to manufacturer's instructions. Immunoblots

were performed with anti-His-HRP (Santa Cruz Biotech) or anti-StrepII-HRP (IBA

Lifesciences) antibodies.

In vitro SUMOylation reconstitution assays

The His-tagged constructs used were according to [53]. Generation of Strep-tagged SUMO

isoform clones have been described earlier. *E. coli* BL21 (DE3) cells containing both *p*KT-973

(SUMO1GG + SCE1) and *p*KT-978 (SAE2 + SAE1) was transformed with either empty vector

(*p*ASK-IBA16), Strep-SUMO3GG, or Strep-SUMO3AA plasmids. Similarly, BL21 (DE3)

cells containing *p*KT-975 (SUMO3GG + SCE1) and *p*KT-978 (SAE2 + SAE1) was either

transformed with either empty vector (*p*ASK-IBA16), Strep-SUMO1GG, or Strep-SUMO1AA

plasmids. As controls, conjugation-deficient SUMO1AA (*p*KT-1017) or SUMO3AA (*p*KT-

1788) in combination with *p*KT-978 was used. Transformed cells were induced with 200 μg/L

Anhydrotetracycline and 0.5 mM IPTG for overnight at 25^0C. After harvesting, cell pellets were

lysed and immunoblotted with anti-His or anti-Strep antibodies.

Statistical Analysis

For all gene expression experiments, Student's t-test was performed to check significance and denoted by one, two and three asterisks (*) indicating *p-value* <0.05, <0.01, and 0.001, respectively. For growth curve and other assays, ANOVA was performed to check statistical significance in growth of bacteria among different genotypes and indicated by alphabets e.g. a, b, c, d, e etc. which depict statistical difference from each other at *p-value* <0.001.

Acknowledgements

IKD, MK and SB deeply acknowledge DST-SERB (Grant No: EMR/2016/001899) and Regional Centre for Biotechnology (RCB), Faridabad for providing financial support and central instrumental facilities. We express our gratitude to Prof. Walter Gassmann, Missouri University, USA, and Prof. Ashis Kumar Nandi, Jawaharlal Nehru University (JNU), New Delhi, India for providing *PstDC3000* strains and *sid2-1* lines and SA measurements, respectively used in this study. We also thank Dr. Souvik Bhattacharjee (JNU) for providing *p*ASK-IBA16 plasmid and Prof. Katsunori Tanaka, (Kwansei Gakuin University, Japan) for providing *E. coli* SUMOylation reconstitution system constructs. IKD acknowledges Department of Biotechnology (DBT), Government of India for providing fellowships during his PhD tenure and KIIT University, Bhubaneswar, India for his PhD registration. MK thanks UGC for his PhD fellowship.

Author Contributions

SB conceived the research. IKD, MK and SB designed the research. IKD and MK performed all the experiments. HvdB generated *amiR-SUM2 sum3-1* line used in this study. SB helped in plasmid constructions and supervised the experiments. IKD, MK and SB analyzed the data and wrote the manuscript.

References

1.   Morrell R, Sadanandom A. Dealing With Stress: A Review of Plant SUMO Proteases.

Front Plant Sci. 2019;10: 1–19. doi:10.3389/fpls.2019.01122

2.   Elrouby N. Extent and significance of non-covalent SUMO interactions in plant

development. Plant Signal Behav. 2014;9: e27948. doi:10.4161/psb.27948

3.   Zhao X. SUMO-Mediated Regulation of Nuclear Functions and Signaling Processes.

Mol Cell. 2018;71: 409–418. doi:10.1016/j.molcel.2018.07.027

4.   Kurepa J, Walker JM, Smalle J, Gosink MM, Davis SJ, Durham TL, et al. The small

ubiquitin-like modifier (SUMO) protein modification system in Arabidopsis.

Accumulation of sumo1 and -2 conjugates is increased by stress. J Biol Chem. 2003;278:

6862–6872. doi:10.1074/jbc.M209694200

5.   Miller MJ, Barrett-Wilt G a, Hua Z, Vierstra RD. Proteomic analyses identify a diverse

array of nuclear processes affected by small ubiquitin-like modifier conjugation in

Arabidopsis. Proc Natl Acad Sci U S A. 2010;107: 16512–16517.

doi:10.1073/pnas.1004181107

6.   Bailey M, Srivastava A, Conti L, Nelis S, Zhang C, Florance H, et al. Stability of small

ubiquitin-like modifier (SUMO) proteases OVERLY TOLERANT to SALT1 and -2

modulates salicylic acid signalling and SUMO1/2 conjugation in Arabidopsis thaliana.

J Exp Bot. 2015;67: 353–363. doi:10.1093/jxb/erv468

7.   Duan G, Walther D. The Roles of Post-translational Modifications in the Context of

Protein Interaction Networks. PLoS Comput Biol. 2015;11: 1–23.

doi:10.1371/journal.pcbi.1004049

8.   Tomanov K, Zeschmann A, Hermkes R, Eifler K, Ziba I, Grieco M, et al. Arabidopsis

PIAL1 and 2 promote SUMO chain formation as E4-type SUMO ligases and are

involved in stress responses and sulfur metabolism. Plant Cell. 2014;26: 4547–60.

doi:10.1105/tpc.114.131300

9.   Tomanov K, Nehlin L, Ziba I, Bachmair A. SUMO chain formation relies on the amino-

terminal region of SUMO-conjugating enzyme and has dedicated substrates in plants.

Biochem J. 2018;475: 61–74. doi:10.1042/BCJ20170472

10.   Gareau JR, Lima CD. The SUMO pathway: Emerging mechanisms that shape

specificity, conjugation and recognition. Nat Rev Mol Cell Biol. 2010;11: 861–871.

doi:10.1038/nrm3011

11.   Elrouby N, Bonequi MV, Porri A, Coupland G. Identification of Arabidopsis SUMO-

interacting proteins that regulate chromatin activity and developmental transitions. Proc

Natl Acad Sci U S A. 2013;110: 19956–61. doi:10.1073/pnas.1319985110

12.   Wang J, Taherbhoy AM, Hunt HW, Seyedin SN, Miller DW, Miller DJ, et al. Crystal

Structure of UBA2ufd-Ubc9: Insights into E1-E2 Interactions in Sumo Pathways. PLoS

One. 2010;5. doi:10.1371/journal.pone.0015805

13.   Park BS, Song JT, Seo HS. Arabidopsis nitrate reductase activity is stimulated by the

E3 SUMO ligase AtSIZ1. Nat Commun. 2011;2: 400. doi:10.1038/ncomms1408

14.   Hendriks IA, D'Souza RCJ, Yang B, Verlaan-de Vries M, Mann M, Vertegaal ACO.

Uncovering global SUMOylation signaling networks in a site-specific manner. Nat

Struct Mol Biol. 2014;21: 927–936. doi:10.1038/nsmb.2890

15.   Nukarinen E, Tomanov K, Ziba I, Weckwerth W, Bachmair A. Protein sumoylation and

phosphorylation intersect in Arabidopsis signaling. Plant J. 2017;91: 505–517.

doi:10.1111/tpj.13575

16.   Flotho A, Melchior F. Sumoylation: A Regulatory Protein Modification in Health and

Disease. Annu Rev Biochem. 2013;82: 357–385. doi:10.1146/annurev-biochem-

061909-093311

17.   Colby T, Matthäi A, Boeckelmann A, Stuible HP. SUMO-conjugating and SUMO-

899   deconjugating enzymes from Arabidopsis. Plant Physiol. 2006;142: 318–332.

900   doi:10.1104/pp.106.085415

901   18.   Hammoudi V, Vlachakis G, Schranz ME, van den Burg HA. Whole-genome

902   duplications followed by tandem duplications drive diversification of the protein

903   modifier SUMO in Angiosperms. New Phytol. 2016;1. doi:10.1111/nph.13911

904   19.   Budhiraja R, Hermkes R, Müller S, Schmidt J, Colby T, Panigrahi K, et al. Substrates

905   related to chromatin and to RNA-dependent processes are modified by Arabidopsis

906   SUMO isoforms that differ in a conserved residue with influence on desumoylation.

907   Plant Physiol. 2009;149: 1529–40. doi:10.1104/pp.108.135053

908   20.   Saracco SA, Miller MJ, Kurepa J, Vierstra RD. Genetic analysis of SUMOylation in

909   Arabidopsis: Conjugation of SUMO1 and SUMO2 to nuclear proteins is essential. Plant

910   Physiol. 2007;145: 119–134. doi:10.1104/pp.107.102285

911   21.   van den Burg HA, Kini RK, Schuurink RC, Takken FLW. Arabidopsis small ubiquitin-

912   like modifier paralogs have distinct functions in development and defense. Plant Cell.

913   2010;22: 1998–2016. doi:10.1105/tpc.109.070961

914   22.   Castaño-Miquel L, Seguí J, Lois LM. Distinctive properties of Arabidopsis SUMO

915   paralogues support the in vivo predominant role of AtSUMO1/2 isoforms. Biochem J.

916   2011;436: 581–590. doi:10.1042/BJ20101446

917   23.   Lois LM, Lima CD, Chua NH. Small ubiquitin-like modifier modulates abscisic acid

918   signaling in Arabidopsis. Plant Cell. 2003;15: 1347–1359. doi:10.1105/tpc.009902

919   24.   Chosed R, Tomchick DR, Brautigam CA, Mukherjee S, Negi VS, Machius M, et al.

920   Structural analysis of Xanthomonas XopD provides insights into substrate specificity of

921   ubiquitin-like protein proteases. J Biol Chem. 2007;282: 6773–6782.

922   doi:10.1074/jbc.M608730200

25. Augustine RC, Vierstra RD. SUMOylation: re-wiring the plant nucleus during stress

and   development.   Curr   Opin   Plant   Biol.   2018;45:   143–154.

doi:10.1016/j.pbi.2018.06.006

26. de Vega D, Newton AC, Sadanandom A. Post-translational modifications in priming the

plant immune system: ripe for exploitation? FEBS Lett. 2018;592: 1929–1936.

doi:10.1002/1873-3468.13076

27. van den Burg HA, Takken FLW. SUMO-, MAPK- and resistance protein-signaling

converge at transcription complexes that regulate plant innate immunity. Plant Signal

Behav. 2010;5: 1597–1601. doi:10.4161/psb.5.12.13913

28. Lee J, Nam J, Park HC, Na G, Miura K, Jin JB, et al. Salicylic acid-mediated innate

immunity in Arabidopsis is regulated by SIZ1 SUMO E3 ligase. 2006; 79–90.

doi:10.1111/j.1365-313X.2006.02947.x

29. Villajuana-Bonequi M, Elrouby N, Nordström K, Griebel T, Bachmair A, Coupland G.

Elevated salicylic acid levels conferred by increased expression of ISOCHORISMATE

SYNTHASE 1 contribute to hyperaccumulation of SUMO1 conjugates in the

Arabidopsis   mutant   early   in   short   days   4.   Plant   J.   2014;79:   206–219.

doi:10.1111/tpj.12549

30. Wimmer P, Schreiner S. Viral mimicry to usurp ubiquitin and sumo host pathways.

Viruses. 2015;7: 4854–4877. doi:10.3390/v7092849

31. Ribet D, Cossart P. Ubiquitin, SUMO, and NEDD8: Key Targets of Bacterial Pathogens.

Trends Cell Biol. 2018;28: 926–940. doi:10.1016/j.tcb.2018.07.005

32. Verma V, Croley F, Sadanandom A. Fifty shades of SUMO: its role in immunity and at

the fulcrum of the growth–defence balance. Mol Plant Pathol. 2018;19: 1537–1544.

doi:10.1111/mpp.12625

33. Hotson A, Chosed R, Shu H, Orth K, Mudgett MB. Xanthomonas type III effector XopD

targets SUMO-conjugated proteins in planta. Mol Microbiol. 2003;50: 377–389.

doi:10.1046/j.1365-2958.2003.03730.x

34.   Roden J, Eardley L, Hotson A, Cao Y, Mudgett MB. Characterization of the

Xanthomonas AvrXv4 effector, a SUMO protease translocated into plant cells. Mol

Plant-Microbe Interact. 2004;17: 633–643. doi:10.1094/MPMI.2004.17.6.633

35.   Tan CM, Li MY, Yang PY, Chang SH, Ho YP, Lin H, et al. Arabidopsis HFR1 is a

potential nuclear substrate regulated by the xanthomonas type III effector

XopDXcc8004. PLoS One. 2015;10: 1–18. doi:10.1371/journal.pone.0117067

36.   Saleh A, Withers J, Mohan R, Marqués J, Gu Y, Yan S, et al. Posttranslational

modifications of the master transcriptional regulator NPR1 enable dynamic but tight

control of plant immune responses. Cell Host Microbe. 2015;18: 169–182.

doi:10.1016/j.chom.2015.07.005

37.   Meulmeester E, Kunze M, Hsiao HH, Urlaub H, Melchior F. Mechanism and

Consequences for Paralog-Specific Sumoylation of Ubiquitin-Specific Protease 25. Mol

Cell. 2008;30: 610–619. doi:10.1016/j.molcel.2008.03.021

38.   Zhu J, Zhu S, Guzzo CM, Ellis NA, Ki SS, Cheol YC, et al. Small ubiquitin-related

modifier (SUMO) binding determines substrate recognition and paralog-selective

SUMO   modification.   J   Biol   Chem.   2008;283:   29405–29415.

doi:10.1074/jbc.M803632200

39.   Zhu S, Goeres J, Sixt KM, Békés M, Zhang XD, Salvesen GS, et al. Protection from

Isopeptidase-Mediated Deconjugation Regulates Paralog-Selective Sumoylation of

RanGAP1. Mol Cell. 2009;33: 570–580. doi:10.1016/j.molcel.2009.02.008

40.   Bhattacharjee S, Halane MK, Kim SH, Gassmann W. Pathogen effectors target

Arabidopsis EDS1 and alter its interactions with immune regulators. Science. 2011;334:

1405–1408. doi:10.1126/science.1211592

41.   Voigt CA. Callose-mediated resistance to pathogenic intruders in plant defense-related

974   papillae. Front Plant Sci. 2014;5: 1–6. doi:10.3389/fpls.2014.00168

975   42.   Dempsey DA, Vlot AC, Wildermuth MC, Klessig DF. Salicylic Acid Biosynthesis and

976   Metabolism. Arab B. 2011;9: e0156. doi:10.1199/tab.0156

977   43.   Cui H, Gobbato E, Kracher B, Qiu J, Bautor J, Parker JE. A core function of EDS1 with

978   PAD4 is to protect the salicylic acid defense sector in Arabidopsis immunity. New

979   Phytol. 2017; 213(4):1802-1817. doi:10.1111/nph.14302

980   44.   Sardar A, Nandi AK, Chattopadhyay D. CBL-interacting protein kinase 6 negatively

981   regulates immune response to Pseudomonas syringae in Arabidopsis. J Exp Bot.

982   2017;68: 3573–3584. doi:10.1093/jxb/erx170

983   45.   Li Y, Li S, Bi D, Cheng YT, Li X, Zhang Y. SRFR1 negatively regulates plant NB-LRR

984   resistance protein accumulation to prevent autoimmunity. PLoS Pathog. 2010;6: 1–11.

985   doi:10.1371/journal.ppat.1001111

986   46.   Kim SH, Gao F, Bhattacharjee S, Adiasor JA, Nam JC, Gassmann W. The arabidopsis

987   resistance-like gene SNC1 is activated by mutations in SRFR1 and contributes to

988   resistance   to   the   bacterial   effector   AvrRps4.   PLoS   Pathog.   2010;6.

989   doi:10.1371/journal.ppat.1001172

990   47.   Chen LZ, Li XY, Huang H, Xing W, Guo W, He J, et al. SUMO-2 promotes mRNA

991   translation by enhancing interaction between eIF4E and eIF4G. PLoS One. 2014;9.

992   doi:10.1371/journal.pone.0100457

993   48.   Watts FZ, Baldock R, Jongjitwimol J, Morley SJ. Weighing up the possibilities :

994   Controlling translation by ubiquitylation and sumoylation. Translation. 2014; 30;2(2).

995   doi: 10.4161/2169074X.2014.959366.

996   49.   Wildermuth MC, Dewdney J, Wu G, Ausubel FM. Isochorismate synthase is required

997   to   synthesize   salicylic   acid   for   plant   defence.   Nature.   2001;414:   562–565.

doi:10.1038/417571a

50.   Matic I, Hagen M Van, Schimmel J, Macek B, Ogg SC. In Vivo Identification of Human

Small Ubiquitin-like Modifier Polymerization Sites by High Accuracy Mass

Spectrometry and an in Vitro to in Vivo Strategy. Mol Cell Proteomics. 2008;7(1):132-

44. doi: 10.1074/mcp.M700173-MCP200

51.   Sharma P, Yamada S, Lualdi M, Dasso M, Kuehn MR. Senp1 Is Essential for

Desumoylating Sumo1-Modified Proteins but Dispensable for Sumo2 and Sumo3

Deconjugation in the Mouse Embryo. Cell Rep. 2013;3: 1640–1650.

doi:10.1016/j.celrep.2013.04.016

52.   Mazur MJ, Kwaaitaal M, Mateos MA, Maio F, Kini RK, Prins M, et al. The SUMO

conjugation complex self-assembles into nuclear bodies independent of SIZ1 and COP1.

Plant Physiol. 2019;179: 168–183. doi:10.1104/pp.18.00910

53.   Okada S, Nagabuchi M, Takamura Y, Nakagawa T, Shinmyozu K, Nakayama JI, et al.

Reconstitution of Arabidopsis thaliana SUMO pathways in E. coli: Functional

evaluation of SUMO machinery proteins and mapping of SUMOylation sites by mass

spectrometry. Plant Cell Physiol. 2009;50: 1049–1061. doi:10.1093/pcp/pcp056

54.   Ohama N, Kusakabe K, Mizoi J, Zhao H, Kidokoro S, Koizumi S, et al. The

transcriptional cascade in the heat stress response of arabidopsis is strictly regulated at

the level of transcription factor expression. Plant Cell. 2016;28: 181–201.

doi:10.1105/tpc.15.00435

55.   Hammoudi V, Fokkens L, Beerens B, Vlachakis G, Chatterjee S, Arroyo-Mateos M, et

al. The Arabidopsis SUMO E3 ligase SIZ1 mediates the temperature dependent trade-

off between plant immunity and growth. Quint M, editor. PLOS Genet. 2018;14:

e1007157. doi:10.1371/journal.pgen.1007157

56.   An L, Zhou Z, Yan A, Gan Y. Progress on trichome development regulated by

phytohormone signaling. Plant Signal Behav. 2011;6: 1959–1962.
doi:10.4161/psb.6.12.18120

57. Srivastava AK, Orosa B, Singh P, Cummins I, Walsh C, Zhang C, et al. Sumo suppresses
the activity of the jasmonic acid receptor CORONATINE INSENSITIVE1. Plant Cell.
2018;30: 2099–2115. doi:10.1105/tpc.18.00036

58. Vertegaal ACO, Andersen JS, Ogg SC, Hay RT, Mann M, Lamond AI. Distinct and
overlapping sets of SUMO-1 and SUMO-2 target proteins revealed by quantitative
proteomics. Mol Cell Proteomics. 2006;5: 2298–2310. doi:10.1074/mcp.M600212-
MCP200

59. Niu D, Lin XL, Kong X, Qu GP, Cai B, Lee J, et al. SIZ1-Mediated SUMOylation of
TPR1 Suppresses Plant Immunity in Arabidopsis. Mol Plant. 2019;12: 215–228.
doi:10.1016/j.molp.2018.12.002

60. Castaño-Miquel L, Seguí J, Lois LM. Distinctive properties of Arabidopsis SUMO
paralogues support the in vivo predominant role of AtSUMO1/2 isoforms. Biochem J.
2011;436: 581–90. doi:10.1042/BJ20101446

61. Castaño-Miquel L, Mas A, Teixeira I, Seguí J, Perearnau A, Thampi BN, et al.
SUMOylation Inhibition Mediated by Disruption of SUMO E1-E2 Interactions Confers
Plant Susceptibility to Necrotrophic Fungal Pathogens. Mol Plant. 2017;10: 709–720.
doi:10.1016/j.molp.2017.01.007

62. Elrouby N, Coupland G. Proteome-wide screens for small ubiquitin-like modifier
(SUMO) substrates identify Arabidopsis proteins implicated in diverse biological
processes. Proc Natl Acad Sci U S A. 2010;107: 17415–17420.
doi:10.1073/pnas.1005452107

63. Capili AD, Lima CD. Structure and Analysis of a Complex between SUMO and Ubc9
Illustrates Features of a Conserved E2-Ubl Interaction. J Mol Biol. 2007;369: 608–618.

doi:https://doi.org/10.1016/j.jmb.2007.04.006

64.   Duda DM, van Waardenburg RCAM, Borg LA, McGarity S, Nourse A, Waddell MB,

1050   et al. Structure of a SUMO-binding-motif Mimic Bound to Smt3p–Ubc9p: Conservation

1051   of a Non-covalent Ubiquitin-like Protein–E2 Complex as a Platform for Selective

1052   Interactions within a SUMO Pathway. J Mol Biol. 2007;369: 619–630.

1053   doi:https://doi.org/10.1016/j.jmb.2007.04.007

1054   65.   Cheong MS, Park HC, Hong MJ, Lee J, Choi W, Jin JB, et al. Specific domain structures

1055   control abscisic acid-, salicylic acid-, and stress-mediated SIZ1 phenotypes. Plant

1056   Physiol. 2009;151: 1930–1942. doi:10.1104/pp.109.143719

1057   66.   Miller MJ, Scalf M, Rytz TC, Hubler SL, Smith LM, Vierstra RD. Quantitative

1058   proteomics reveals factors regulating RNA biology as dynamic targets of stress-induced

1059   SUMOylation in Arabidopsis. Mol Cell Proteomics. 2013;12: 449–63. Available:

1060   http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3567865&tool=pmcentrez

1061   &rendertype=abstract%5Cnhttp://www.ncbi.nlm.nih.gov/pubmed/23197790

1062   67.   Bossis G, Melchior F. Regulation of SUMOylation by reversible oxidation of SUMO

1063   conjugating enzymes. Mol Cell. 2006;21: 349–357. doi:10.1016/j.molcel.2005.12.019

1064   68.   Sydorskyy Y, Srikumar T, Jeram SM, Wheaton S, Vizeacoumar FJ, Makhnevych T, et

1065   al. A Novel Mechanism for SUMO System Control: Regulated Ulp1 Nucleolar

1066   Sequestration. Mol Cell Biol. 2010;30: 4452–4462. doi:10.1128/mcb.00335-10

1067   69.   Pinto MP, Carvalho AF, Grou CP, Rodríguez-Borges JE, Sá-Miranda C, Azevedo JE.

1068   Heat shock induces a massive but differential inactivation of SUMO-specific proteases.

1069   Biochim   Biophys   Acta   -   Mol   Cell   Res.   2012;1823:   1958–1966.

1070   doi:10.1016/j.bbamcr.2012.07.010

1071   70.   Hepworth J, Dean C. Flowering locus C's lessons: Conserved chromatin switches

1072   underpinning developmental timing and adaptation. Plant Physiol. 2015;168: 1237–

1245. doi:10.1104/pp.15.00496

71. Kong X, Luo X, Qu GP, Liu P, Jin JB. Arabidopsis SUMO protease ASP1 positively

regulates flowering time partially through regulating FLC stability. J Integr Plant Biol.

2017;59: 15–29. doi:10.1111/jipb.12509

72. Citro S, Jaffray E, Hay RT, Seiser C, Chiocca S. A role for paralog-specific sumoylation

in histone deacetylase 1 stability. J Mol Cell Biol. 2013;5: 416–427.

doi:10.1093/jmcb/mjt032

73. Fasci D, Anania V, Lill J, Salvesen G. Response to comment on "sumo deconjugation

is required for arsenic-triggered ubiquitylation of PML." Sci Signal. 2015;9: 1–12.

doi:10.1126/scisignal.aad9777

74. Sanchez-Duran MA, Dallas MB, Ascencio-Ibanez JT, Reyes MI, Arroyo-Mateos M,

Ruiz-Albert J, et al. Interaction between Geminivirus Replication Protein and the

SUMO-Conjugating Enzyme Is Required for Viral Infection. J Virol. 2011;85: 9789–

9800. doi:10.1128/jvi.02566-10

75. Xiong R, Wang A. SCE1, the SUMO-conjugating enzyme in plants that interacts with

NIb, the RNA-dependent RNA polymerase of Turnip mosaic virus, is required for viral

infection. J Virol. 2013;87: 4704–15. doi:10.1128/JVI.02828-12

76. Defraia CT, Schmelz EA, Mou Z. A rapid biosensor-based method for quantification of

free and glucose-conjugated salicylic acid. Plant Methods. 2008;4: 1–11.

doi:10.1186/1746-4811-4-28

77. Schenk ST, Hernández-Reyes C, Samans B, Stein E, Neumann C, Schikora M, et al. N-

acyl-homoserine lactone primes plants for cell wall reinforcement and induces resistance

to bacterial pathogens via the salicylic acid/oxylipin pathway. Plant Cell. 2014;26:

2708–2723. doi:10.1105/tpc.114.126763

78. Becker D. Binary vectors which allow the exchange of plant selectable markers and

reporter genes. Nucleic Acids Res. 1990;18: 203. doi:10.1093/nar/18.1.203

79.   Clough SJ, Bent AF. Floral dip: A simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. Plant J. 1998;16: 735–743. doi:10.1046/j.1365-313X.1998.00343.x

Supporting information

Additional supporting information may be found in the online version of this article.

S1 Fig: *sum3-1* **partially alleviates, whereas** *eds1-2* **and** *sid2-1* **abolishes growth deficiencies of** *sum1-1***.**

S2 Fig: *sum1-1* **and** *sum2-1* **plants are enhanced resistant whereas** *sum3-1* **is hypersusceptible to TNL-specific** *PstDC3000(hopA1)* **but not to CNL-specific** *PstDC3000(avrRpm1)* **avirulent strains.**

S3 Fig: *sum1-1* **and** *sum2-1* **leaves display increased whereas** *sum3-1* **is deficient in callose deposition in response to virulent** *PstDC3000(EV)* **or avirulent** *PstDC3000(avrRps4)* **infections.**

S4 Fig: Developmental phenotype of 4-week-old SD grown plants of indicated genotypes.

S5 Fig: *sum1-1* **and** *sum2-1* **display rapid whereas** *sum3-1* **is delayed than Col-0 in induction of** *PR1* **and** *PR2* **transcripts in response to virulent** *PstDC3000(EV)* **or avirulent** *PstDC3000(avrRps4)* **infections.**

1122 **S6 Fig: Increased trichome density, elevated expression of PTI markers *FRK1* or**

1123 ***WRKY29*, and enhanced resistance to avirulent *PstDC3000(hopA1)* in *sum1-1* is SA-**

1124 **modulated.**

1125

1126 **S7 Fig: Enhanced basal immunity and ETI defences in *sum2-1* to TNL-specific *PstDC3000***

1127 **strains is EDS1-dependent.**

1128

1129 **S8 Fig: Developmental defects and enhanced TNL-specific immunity in *sum1-1* is partially**

1130 ***SUM3* regulated.**

1131

1132 **S9 Fig: *SUM3* buffers increased induction of *PR1* in *sum1-1* in response to *PstDC3000***

1133 **challenges.**

1134

1135 **S10 Fig: Schematic representation of key amino acid conservation and divergences among**

1136 **three *Arabidopsis* SUMO isoforms suggest their functional overlaps/distinctions.**

1137

1138 **S11 Fig: Complemented *sum1-1* or *sum3-1* lines have wild-type levels of SUMO1/2**

1139 **conjugates upon *PstDC3000* (EV) infection.**

1140

1141 **S12 Fig: SUMO1 and SUMO3 cause reciprocal enhancements of SUMO-conjugates in *E.***

1142 ***coli* SUMOylation-reconstitution system.**

1143

1144 **S13 Fig: *SUM3* mutation reduces SUMO1/2-conjugate enhancements during heat-shock.**

1145

1146 **S1 Table: List of primers used in this study.**

Figures

Fig 1: *sum1* **and** *sum2* **mutants display enhanced whereas** *sum3* **is deficient in defenses against** *Pseudomonas syringae* **pv.** *tomato* **(***PstDC3000***) strains.** (A) Developmental phenotypes of 4-week-old Col-0, *sum1-1, sum2-1,* and *sum3-1* mutants propagated under short day (SD) and long day (LD) growth conditions. (B) Whisker Box plot of trichome densities (#Trichomes/4.1 mm^2) in Col-0 and *sum1-1, sum2-1,* and *sum3-1* mutants grown under SD conditions. Trichomes in 4-week-old plants were counted after imaging leaves under bright field in a fluorescence microscope. Error bars indicate standard deviation of trichome densities in 10 random images of

different leaf sections (n=10). *** indicates p <0.001, ns= non-significant. (C-D) Growth of (C) virulent *PstDC3000(EV)* or (D) avirulent *PstDC3000*(*avrRps4*) strains, respectively in Col-0 and *sum1-1, sum2-1,* and *sum3-1* mutants. Three to four expanded leaves from 4-week-old plants of each genotype were infiltrated with the indicated bacterial suspension at a density of 5 X 10^4 cfu ml^{-1}. Leaf discs (of predefined diameter) were punched from the infiltrated area, macerated in 10mM $MgCl_2$, serially diluted and plated on appropriate antibiotic plates. Bacterial titer was calculated at 0- and 3-dpi (days post-infiltration) for the indicated bacterial infection. Whisker Box plot with Tukey test; n= 12; ANOVA was performed to measure statistical significant differences (at *p-value* <0.001) in growth of bacteria in Log_{10} scale. (E) Whisker Box plot of callose deposits/mm^2 area of infected leaves. Infiltrated leaves at 24-hpi (hours post-infiltration) were bleached in acetic acid: ethanol solution, stained with Aniline blue and were observed under DAPI filter in a fluorescence microscope. The images were analyzed in Imaris 8.0 software. The data is representative of callose deposits median from independent random sections area (n=10). Statistical significance was determined by Student's t-test (*** indicates at p <0.001.

Fig 2: Basal SA levels and expression of several defense-associated markers are elevated in *sum1-1* or *sum2-1* whereas downregulated in *sum3-1* plants. (A) Total SA (SA + SAG) and (B) free SA levels in *sum* and combinatorial mutants in 4-week-old plants measured by biosensor *Acinetobacter* method. Data is representative of three biological replicates. Statistical significance calculated by Student's t-test is denoted by alphabet on top of each bar. Different alphabets indicate significance at *p*<0.001. Relative transcript abundance of (C) *PR1*, *PR2*, *ICS1* (E) *FRK1*, *WRKY29*, *SNC1* (F) *SUM1*, *SUM2*, and *SUM3* in 4-week-old SD grown Col-0, *sum* and indicated combinatorial mutants was determined by qRT-PCR and normalized to *SAND* expression. The values are represented as fold change relative to Col-0 (WT). Data is representative of mean of three biological replicates (n=3). Error bars indicate SD. Student's t-test was performed to calculate statistical significance; *=*p*<0.05; **=*p*<0.01; ***=*p*<0.001, ns= not significant. (D) Total protein extracts from 4-week-old SD grown Col-0, *sum1-1*, *sum2-1*, and *sum3-1* plants were immunoblotted with anti-PR2 antibodies. Ponceau S stain of the Rubisco subunit indicative of equal protein loading between samples is shown. Migration positions of molecular weight standards (in kDa) are indicated.

Fig 3. Altered defense responses to *PstDC3000* strains in *sum1-1* or *sum3-1* plants are restored to wild-type levels in the respective complemented lines. (A) Relative transcript abundance of *SUM1, SUM2* and *SUM3* in 4-week-old SD grown Col-0, *sum* mutants and the corresponding complemented line(s) was determined by qRT-PCR and normalized to *SAND* expression. The values are represented as fold change relative to Col-0 (WT). Data is representative of mean of three biological replicates (n=3). Error bars indicate SD. Statistical significance was determined by Student's t-test; **=$p<0.01$; ***=$p<0.001$, ns= not significant. (B) Total (*left panel*) and free SA (*right panel*) levels in *sum* mutants and the complemented lines. Data is representative of three biological replicates. Statistical significance was determined by Student's t-test and represented by alphabets with significance at $p<0.001$. (C) Endogenous PR1 protein levels in *sum* and complemented lines. Immunoblot with anti-Actin antibodies and Ponceau S staining of membrane show equal protein loading among the extracts. (D) Growth of (*left panel*) virulent *PstDC3000(EV)* or (*right panel*) avirulent *PstDC3000(hopA1)* strains, respectively in the indicated plants. Leaves from 4-week-old plants of each line were infiltrated with the indicated bacterial suspension at a density of 5 X 10^4 cfu ml^{-1}. Leaf discs from the infiltrated area was macerated in 10mM MgCl$_2$, serially diluted and plated on appropriate antibiotic plates. Bacterial titers calculated at 0- and 3-dpi (days post-infiltration) for the indicated infection are shown. Different alphabets denote statistical significance at *p-value* <0.001.

Fig 4: *sum1-1* and *sum2-1* display enhanced basal and rapid induction of PR1 and PR2 proteins upon pathogen challenge whereas *sum3-1* plants are deficient in these responses. Total protein from 4-week-old SD grown Col-0, *sum1-1*, *sum2-1*, *sum3-1*, *sum1-1 sum3-1*, *sum1-1 eds1-2*, *sum2-1 eds1-2*, *eds1-2*, *sum1-1 sid2-1*, and *sid2-1* was extracted from (A) un-infiltrated or at 6-, 12-, and 24-hpi (hours post-infiltration) with either (B) virulent *PstDC3000(EV)* or (C) avirulent *PstDC3000(avrRps4)* strains and immunoblotted with anti-PR1 or anti-PR2 antibodies. The membranes were also probed with anti-Actin antibodies or stained with Ponceau S for Rubisco subunit to indicate comparable loading between extracts. Migration position of protein molecular weight standards (in kDa) are indicated.

Fig 5: Growth retardation and enhanced defenses in *sum1-1* are SA-dependent. (A) Developmental phenotypes of 4-week-old Col-0, *sum1-1*, *sum1-1 eds1-2*, *eds1-2*, *sum1-1 sid2-1*, and *sid2-1* plants grown under SD and LD growth conditions. (B) Relative transcript abundance of *PR1* and *ICS1* in 4-week-old SD grown indicated plants was determined by qRT-PCR and normalized to *SAND* expression. The values are represented as fold change relative to Col-0 (WT). Data is representative of independent experiments (n=3). Error bars indicate SD. Student's t-test significances are mentioned (*=$p<0.05$; **=$p<0.01$; ***=$p<0.001$). Growth of (C) virulent *PstDC3000(EV)* and (D) avirulent *PstDC3000 (avrRps4)* strains, respectively in indicated plants are shown. Three to four expanded leaves from 4-week-old SD grown plants of each genotype were infiltrated with the indicated bacterial suspension at a density of 5 X 10^4 cfu ml^{-1}. Leaf discs were punched from the infiltrated area, macerated in 10mM MgCl$_2$, serially diluted and plated on appropriate antibiotic plates. Bacterial titer was calculated at 0- and 3-dpi for the indicated infiltration. Whisker Box plot with Tukey test; n= 12-18; ANOVA was performed to measure statistical significant differences (at *p-value* <0.001) in growth of bacteria in Log$_{10}$ scale.

Fig 6: *sum3-1* **alleviates developmental defects and enhanced defense responses in** *sum1-1*. (A) Developmental phenotypes of 4-week-old SD grown Col-0, *sum1-1, sum3-1* and *sum1-1 sum3-1* plants. (B) Total protein extracts from indicated plants were immunoblotted with anti-SNC1 antibodies. Ponceau S stain of the Rubisco subunit and anti-Actin immunoblot indicative of equal protein loading between samples are shown. Migration positions of molecular weight standards (in kDa) are indicated. Growth of (C) virulent *PstDC3000(EV)* or (D) avirulent *PstDC3000(avrRps4)* strains, respectively in indicated plants. Three expanded leaves from 4-week-old SD grown plants of each genotype were infiltrated with the respective bacterial suspension at a density of 5×10^4 cfu ml^{-1}. Leaf discs of predefined diameter were punched from the infiltrated area, macerated in 10mM MgCl$_2$, serially diluted and plated on appropriate antibiotic plates. Bacterial titer was calculated at 0- and 3-dpi with for indicated infiltration. Whisker Box plot with Tukey test; n= 12; ANOVA was performed to measure statistical significance (at *p-value* <0.001) in growth of bacteria in Log$_{10}$ scale.

Fig 7: SUMO3, but not SUMO1, can form oligomers. (A) Green fluorescence indicative of YFP reconstitution between BiFC (Bi-molecular Fluorescence complementation) combinations of SUMOylation proficient (GG) or deficient (AA) forms of SUMO1 and SUMO3 isoforms are shown. *Agrobacterium* GV3101 strains expressing the indicated BiFC constructs were co-infiltrated in expanded *N. benthamiana* leaves. Infiltrated leaf sections were visualized under a confocal microscope at 2-days post-infiltration (dpi). (Scale bar = 50 μm). Images are representative of pattern observed in two independent experiments. (B) *In vitro* binding assays between SUMO1 and SUMO3 isoforms. His-or Strep-II-tagged SUMO1 or SUMO3 isoform as indicated were expressed in *E. coli.* Bacterial lysates from indicated combinations were mixed and enriched through the Ni^{2+}-NTA matrix. Enrichment of the mentioned His-SUMO and the co-eluting isoform was investigated via immunoblotting with anti-His or anti-Strep antibodies, respectively (pull-down panel). The input panel shows the protein levels in the extracts used for the enrichments. Position of protein molecular weight standards (in kDa) are indicated.

Fig 8: Loss of *SUM3* reduces SUMO1/2-conjugate enhancements in response to *PstDC3000(EV)* infection. (A) Fully expanded 3-4 leaves from 4-week-old SD grown Col-0, *sum1-1, sum3-1* and *sum1-1 sum3-1* plants were infiltrated either with buffer alone (- lanes) or with the virulent *PstDC3000(EV)* strain (+ lanes) at a density of 5 X 10^6 cfu ml^{-1}. Total protein isolated from the infected leaves at 24 hrs post-infiltration (hpi) was used for immunoblotting with anti-SUMO1/2 antibodies. Approximate positions of SUMO1/2-conjugates are shown. Anti-Actin immunoblot or Ponceau S straining demonstrate comparable protein loadings from the extracts. Relative positions of protein molecular weight standards (in kDa) are indicated. Relative transcript abundance at 24-hpi of (B) *SAE1, SAE2, SCE1, SIZ1, HPY2* and (C) *ESD4, ELS1, OTS1, OTS2* in mock versus *PstDC3000(EV)* infiltrated samples of Col-0, *sum1-1* and *sum3-1* was determined by qRT-PCR and normalized to *SAND* expression. The values are represented as fold change relative to Col-0 (WT). Data is representative of mean of three biological replicates (n=3). Error bars indicate SD. Student's t-test was performed to calculate statistical significance; *=p<0.05;

**=$p<0.01$; ***=$p<0.001$, ns= not significant. (D) Genetic model for *SUM1/2* crosstalks with *SUM3* in regulation of SA-dependent defences and heat-shock responses in *Arabidopsis*. *SUM1/2* function additively but not equally as negative regulators of defences via modulation of SA-signaling routes and expression of defense-associated markers. Pathogen exposure induce SA-dependent *SUM3* role as a positive regulator leading to increase in SUMO1/2-conjugation proficiencies, and expression of response-appropriate markers such as PR proteins, PTI markers, and callose deposits. Induced SUMO1/2-conjugates modulate SUMO3 functions and prevents overshooting of responses. *SUM1* also regulates via *SUM3*-independent but SA-dependent developmental aspects such as trichome production and leaf textures.