Corresponding Author: Saikat Bhattacharjee, saikat@rcb.res.in

Article Type: Regular Manuscript

Contrasting functions of *Arabidopsis* SUMO1/2 isoforms with SUMO3 intersect to
modulate innate immunity and global SUMOylome responses

Ingole Kishor Dnyaneshwar[1,2], Mritunjay Kasera[1], Harrold A. van den Burg[3], and Saikat
Bhattacharjee[1*]

[1]Laboratory of Signal Transduction and Plant Resistance, UNESCO-Regional Centre for
Biotechnology (RCB), NCR Biotech Science Cluster, 3[rd] Milestone, Faridabad-Gurgaon
Expressway, Faridabad- 121 001, Haryana, India.

[2]Kalinga Institute of Industrial Technology (KIIT) University, Bhubaneswar- 751 024, Odisha,
India.

[3]Molecular Plant Pathology, Swammerdam Institute for Life Sciences, Faculty of Science,
University of Amsterdam, Science Park 904, 1098 XH, Amsterdam, Netherlands.

Author for Correspondence:
Saikat Bhattacharjee
Tel: (+91) 0129-2848837
email: saikat@rcb.res.in

ORCID
Saikat Bhattacharjee https://orcid.org/0000-0003-1369-7700
Ingole Kishor Dnyaneshwar https://orcid.org/0000-0003-1946-6639
Mritunjay Kasera
Harrold A. van den Burg https://orcid.org/0000-0003-4142-374X

Word counts
-    Abstract: 150
-    Introduction: 1455

ABSTACT

Reversible covalent attachment of SMALL UBIQUITIN-LIKE MODIFIERS (SUMOs) on

target proteins regulate diverse cellular process across all eukaryotes. In *Arabidopsis thaliana*,

most mutants with perturbed SUMOylation display severe impairments in growth and

adaptations to physiological stresses. Since SUMOs self-regulate SUMOylation efficiencies,

existence of multiple isoforms introduces possibilities of their functional intersections which

remain unexplored. Using well-established defense responses elicited against virulent and

avirulent *Pseudomonas syringae* pv. *tomato* strains, we investigated crosstalks in individual

and combinatorial Arabidopsis *sum* mutants. We report that while *SUM1* and *SUM2* additively,

but not equivalently suppress basal and TNL-specific immunity, *SUM3* promotes these

defenses. Remarkably, the loss of *SUM3* partially attenuates heightened immunity of *sum1* or

sum2 mutants suggesting intricate crosstalks among these isoforms in optimizing immune

amplitudes. Similar *SUM1-SUM3* intersections also affect global SUMOylome responses to

heat-shock. Overall, our investigations reveal novel insights into auto-regulatory mechanisms

of host SUMOylome maintenance and adjustments to environmental challenges.

Key words: *Arabidopsis thaliana,* crosstalks, defense, *PstDC3000*, salicylic acid (SA), SUMO

isoforms, SUMO1/2-conjugates.

INTRODUCTION

Post-translational modifications (PTMs) dynamically and reversibly modulate target proteins

properties either by affecting their fate, location, or function and facilitate transitory adaptations

often necessary for developmental passages and responding to biotic/abiotic cues. Of special

importance in this category is the reversible attachment of the SMALL UBIQUITIN-LIKE

MODIFIER (SUMO) through a process known as SUMOylation. This highly conserved mode of

covalently modifying proteins is utilized primarily to regulate nuclear functions such as DNA replication, chromatin remodeling, transcription, and post-transcriptional processes in eukaryotes (reviewed in Morell and Sadanandom 2019; Elrouby 2014; Zhao, 2018). In plants, temperature perturbations, salinity changes and application of defensive hormones such as salicylic acid (SA) cause massive changes on global SUMOylome (Bailey et al., 2015; Kurepa et al., 2003; Miller et al., 2010). Interestingly, a vast majority of differentially SUMOylated proteins are nuclear, perhaps indicative of a rapid mechanism to modulate transcriptome in response to stimulus. Computational studies identify SUMOylation-annotated proteins as central relay players of protein-protein interaction webs, especially for transcriptional processes (Duan and Walther, 2015).

SUMOylation cascade recruits processed SUMOs with C-terminus di-glycine (GG) residues, a product of catalysis by specialized SUMO or Ubiquitin-like proteases (ULPs), to covalently attach via an isopeptide bond to ε-amino group of lysine residues on target proteins. The modified lysine often is a part of a partially conserved motif, ψ-K-X-D/E (ψ = hydrophobic amino acid, X= any amino acid, D/E= aspartate/glutamate). A conjugatable SUMO, subsequently forms a thiol-ester bond with the SUMO E1 ACTIVATING ENZYME (SAE). A trans-esterification reaction further shuttles SUMO to the SUMO E2 CONJUGATING ENZYME 1 (SCE1) and then to target lysine on SUMOylation substrates. Enrichment of SUMOylated proteins upon heat shock reveal that SUMO1 can conjugate to itself and form poly-SUMO1 chains (Miller et al., 2010). Although SCE1 is capable of catalyzing polySUMO-chain formation, members of SUMO E4 ligases the PROTEIN INHIBITOR OF ACTIVATED STAT LIKE protein (PIAL1/2 in Arabidopsis) enhance polySUMOylation (Elrouby, 2014; Tomanov et al., 2014, 2018). Substrate specificities both for mono- or polySUMOylation are further regulated by SUMO E3 LIGASES such as HIGH PLOIDY2/METHYL METHANE SULFONATE21 (HPY2/MMS21) and SAP and MIZ1 (SIZ1) (Gareau and Lima, 2010). Fate of covalently-attached SUMOs may either include de-conjugation and recycling by SUMO proteases or targeted proteolysis of the substrate through ubiquitin-mediated pathway. Indeed, proteasome components are enriched in poly-SUMO1 pull-downs (Miller et al., 2010). These interactions are non-covalent in nature facilitated by hydrophobic amino acid-rich SUMO-interaction motifs (SIMs) present in the recipient. A group of moderately conserved SUMO-targeted ubiquitin ligases (STUbLs) bind poly-SUMO chains via internal SIMs to ubiquitinylate the substrate (Elrouby et al., 2013). Interestingly, SIMs are also only enriched in the SUMOylation-associated proteins such as SIZ1, SCE1, SAE2, SUMO-protease EARLY IN SHORT DAYS 4 (ESD4), PIAL1/2, STUbLs implying strong auto-regulatory mechanisms (Tomanov et al., 2014; Wang et al., 2010). Not the least, increasing evidences that reciprocal

influences of phosphorylation, ubiquitination, and acetylation that often compete with or modulate

the efficacy of SUMOylation highlights the complexity of crosstalks among PTMs (Park et al.,

2011; Hendriks *et al.*, 2014; Nukarinen et al., 2017)

Unlike fruit fly, worm, or yeast, humans and plants express multiple SUMO isoforms (Flotho and

Melchior, 2013). In *Arabidopsis thaliana*, 4 SUMO isoforms are expressed namely SUMO1,

SUMO2, SUMO3 and SUMO5 (Colby et al., 2006; Kurepa et al., 2003). SUMO1 and SUMO2

share 89% sequence identity, whereas SUMO3 and SUMO5 are considerably diverged from

SUMO1 (48% and 35% identical, respectively). The homolog pair SUMO1/2 are a result of

genomic duplication event of a SUMO clade that preceded the evolution of monocots and eudicots

from common ancient angiosperms (Hammoudi et al., 2016). Tandem organization of *SUM2* and

SUM3 genes in Arabidopsis are a result of gene duplication subsequently followed by

diversification of only *SUM3* sequences. At relative expression levels, *SUM1/2* are more abundant

than *SUM3* or *SUM5* (Budhiraja et al., 2009). Partial overlaps in tissue-specific expression patterns

and biochemical properties taken together with embryonic lethality of *sum1sum2* double mutant

indicate that plants require at least one functional copy of either of these redundant isoforms

(Saracco *et al.*, 2007; van den Burg *et al.*, 2010; Castaño-Miquel *et al.*, 2011). The *Arabidopsis*

sum3 mutant is viable with mild late-flowering phenotype (van den Burg et al., 2010). Acutely

different SUMO3 however, unlike SUMO1/2, cannot form poly-SUMO chains *in vitro*, and shows

little or no change to heat-shock treatment (Budhiraja et al., 2009; Colby et al., 2006; Kurepa et

al., 2003; Lois et al., 2003; van den Burg et al., 2010). Not the least, SUMO1/2 but not SUMO3-

modified targets, are efficiently deconjugated by SUMO proteases *in vitro* (Chosed et al., 2007;

Colby et al., 2006).

Host SUMOylome readjustments play a vital role in regulation of plant immunity and been

comprehensively highlighted in several excellent reviews (Augustine and Vierstra, 2018; de Vega

et al., 2018; van den Burg and Takken, 2010). This was first evident in the Arabidopsis SUMO E3

ligase *SIZ1* mutant (*siz1-2*) and subsequently in SUMO protease mutants of *OVERLY TOLERANT*

TO SALT 1/2 (OTS1/2) and *ESD4* (Lee *et al.*, 2006; Saracco *et al.*, 2007; Bailey *et al.*, 2015;

Villajuana-Bonequi *et al.*, 2014). Increased accumulation of defensive hormone salicylic acid (SA)

and constitutive expression of PATHOGENESIS-RELATED (PR) proteins in these mutants

conferred enhanced resistance when challenged with the bacterial pathogen *Pseudomonas*

syringae pv. *tomato* strain DC3000 (*PstDC3000*). In this context, it is therefore not surprising that

pathogens attempt to manipulate host SUMOylome to increase their colonization efficiencies

(Wimmer & Schreiner, 2015; Ribet & Cossart, 2018; Verma *et al.*, 2018). Several bacterial

phytopathogenic effectors interfere with host SUMOylation as a mode to suppress immunity

(Hotson et al., 2003; Roden et al., 2004). XopD, a secreted effector from *Xanthomonas campestris*

pv. vesicatoria (*Xcv*) de-conjugates SUMO from unknown targets in plants (Tan et al., 2015).

Mutations that disrupt SUMO-protease functions not only render the cognate strain deficient in

virulence but also lower host defense induction.

Suggestively, both SUM1/2 jointly suppress SA-dependent defenses since plants null for *SUM1*

and expressing microRNA-silenced *SUM2* (*sum1-1 amiR-SUM2*) have drastically reduced total

SUMO1/2-conjugates compared to wild-type and display heightened anti-bacterial immunity (van

den Burg et al., 2010). The SUMO E3 ligase mutant *siz1-2*, also with lower levels of SUMO1/2

conjugates are similarly enhanced resistant to *PstDC3000* (Lee et al., 2006; S. A. Saracco et al.,

2007). On the contrary, *esd4-1* or *ots1ots2* plants with deficient SUMO protease ESD4 or OTS1/2

functions, respectively have increased SUMO1/2-conjugates yet display elevated SA-dependent

defenses (Bailey et al., 2015; Morrell and Sadanandom, 2019; Villajuana-Bonequi et al., 2014).

Constitutive activation of SA-dependent defenses also occur in *SUM1/2*-overexpressing transgenic

plants regardless of whether the over-expressed SUMO isoforms are conjugation-proficient or

deficient (van den Burg et al., 2010). Thus, it is likely that perturbations rather than increase or

decrease in SUMO1/2-conjugates *per se* regulate immune responses. Unlike *SUM1/2*, *SUM3* is

SA-inducible, upregulated in *sum1-1 amiR-SUM2* plants, and upon over-expression enhance the

resistance to *PstDC3000* (van den Burg et al., 2010). SUMO3-mediated SUMOylation of NPR1

(NONEXPRESSOR OF PATHOGENESIS-RELATED 1), a master transducer of SA-signaling is

essential for defenses thus placing *SUM3* as a *bona fide* positive immune regulator (Saleh et al.,

2015). Whether increased SUMO3 activities during defenses intersect to alter SUMOylome

outcomes of a host remains unexplored. Role of *SUM3* in other stress responses similarly is also

unknown. Evidently, perturbing host SUMOylome during biotic/abiotic stresses or through

loss/gain of a specific isoform function initiates complex signaling more especially because the

SUMOs themselves moderate SUMOylation proficiencies of targets. Miller *et al.* (2010) indeed

identified SAE2, SCE1, SIZ1, and EDS4 as candidates whose SUMOylation footprints change

upon stress exposure. Animal studies also elegantly demonstrate 'SUMO-preference/switching'

wherein the isoform choice for substrate conjugation is modulated not only by their relative levels

but also by its influence on SUMOylation machinery (Meulmeester *et al.*, 2008; Zhu *et al.*, 2008,

Zhu *et al.*, 2009). Hence, it is beyond doubt that SUMO isoforms functionally intersect when the

host SUMOylome-equilibrium is disturbed. However, such evidences from plant systems are

completely lacking.

In this study, we utilized defense responses to various strains of *PstDC3000* as a measure to test

the contribution of individual and combinatorial *Arabidopsis* SUMO isoforms in immunity. Our

data suggest that *SUM1/2* function additively, but not equally, as negative regulators of SA-driven

basal and TNL (Toll-Interleukin-1 receptor-like domain)-type immunity. In contrast, *SUM3* has a

more positive immune role potentiating auto-immunity that occur due to loss of *SUM1* or *SUM2*.

We demonstrate that accumulation kinetics of not only defense-associated markers but also of

SUMOylation-associated genes are regulated by SUMO1-SUMO3 crosstalks. We further report

that global change in a host SUMOylome in response to immune activation or heat-stress is also

influenced by intersections of SUMO isoform activities. Overall, our results open newer avenues

to unravel role of SUMO isoforms functions in maintenance and alterations of a host SUMOylome.

MATERIALS AND METHODS

Plant materials and growth conditions

Arabidopsis thaliana mutant lines *sum1-1* (SAIL_296_C12), *sum2-1* (SALK_029775C) and

sum3-1 (SM_3_2707/SM_3_21645) were obtained from Arabidopsis stock centre

(https://www.arabidopsis.org). Seeds of *amiR-SUM2 sum3-1* were generated by Dr. Harrold

van den Burg. All plants were grown at 22^0C with 70% humidity under Long Days (LD; 16 h:

8 h, light: dark) or Short Days (SD; 8 h: 16 h, light : dark) having light intensity 100 μmol μm^{-2}s^{-1}

light. Specific growth conditions are indicated in respective legends. To generate

combinatorial mutants, *sum1-1* was genetically crossed with *eds1-2*, *sid2-1* or *sum3-1* mutant

plants. The *sum2-1* mutant was similarly crossed with *eds1-2* to generate *sum2-1 eds1-2*.

Double mutants (*sum1-1 eds1-2, sum1-1 sid2-1* and *sum1-1 sum3-1*) were identified in F2

population by PCR based genotyping. To generate *sum1-1:His-H89R-SUM1*, *His-H89R-SUM1*

(Miller et al., 2010) was crossed with *sum1-1*. Segregating F2 populations were PCR-based

genotyped for the presence of homozygous *sum1-1*, wild-type *SUM2* and homozygous copies

of *His-H89R-SUM1* transgene. All primers used are listed in Table supplement 1.

Trichome visualisation and quantification

For trichome visualization, leaves from 4-weeks old SD-grown plants were harvested and washed in Acetic acid: Ethanol (1:3) solution for overnight to bleach all pigments then visualised under bright field in a fluorescence microscope. The images were captured as 4848 X 3648 pixels from 4.1 mm^2 leaf area. The trichome numbers were counted in 10 randomly selected images. Experiment was repeated twice with similar observations.

Salicylic acid (SA) measurements

Free salicylic acid (SA) and glucose-conjugated SA (SAG) measurements were done using the *Acinetobacter* sp. ADPWH_*lux* biosensor system (Defraia et al., 2008). In brief, 100 mg leaf tissues were collected and frozen in liquid nitrogen. The tissue was homogenized in 250μl of acetate buffer (0.1 M, pH 5.6). Samples were centrifuged at 12000 rpm for 15 minutes. One aliquot (100μl) of the supernatant was used for free SA measurements, and another was incubated with 6U of β-glucosidase (Sigma-Aldrich) for 90 min at 37°C for total SA measurement. 20μl aliquot of each plant extract were added to 50μl of secondary culture of *Acinetobacter* sp. ADPWH_*lux* (OD$_{600}$ of 0.4) with additional 60 μl of LB. Standard SA solutions (prepared in *sid2-1* extract) were also taken to generate standard curve to calculate the amount of SA present in samples. Plate was incubated at 37 °C for 1 h, and luminescence detected using the POLARStar Omega Luminometer (BMG Labtech). Data shown is representative of three biological replicates with SD.

***In planta* assays for bacterial growth and kinetics of defense-associated marker expressions**

Bacterial growth assays were performed according to Kim *et al.* (2010). Briefly, *PstDC3000* strains were infiltrated with a needleless syringe at a density of 5 X 10^4 cfu ml^{-1}into fully expanded rosette leaves of 3-4-weeks-old SD grown plants. Leaf discs harvested from infiltrated area were macerated in 10mM MgCl$_2$, serially diluted and plated onto selective medium plates. Bacterial growth was determined at 0 and 3-days post-infiltration. Bacterial infiltrations for kinetics of defensive markers were carried as above except a 5 X 10^6 cfu ml^{-1} bacterial inoculum was used. Infiltrated samples harvested at 6, 12 and 24 hpi were processed separately for total RNA extraction and qPCR analysis or for immunoblots with indicated antibodies.

Callose deposition assay and image analysis

The callose deposition assays were performed according to (Schenk et al., 2014). In brief, 4-week-old leaves of SD grown Col-0, *sum1-1, sum2-1* and *sum3-1* plants were infected with the indicated *PstDC3000* strains. At 24 hpi, 3-4 random infiltrated leaves were harvested and washed first in Acetic acid: Ethanol (1:3) solution for overnight to bleach all pigments and then washed for 30 min in 150 mM K_2HPO_4 solution. The leaves were then incubated in dark for 2 hours in 150 mM K_2HPO_4 containing 0.01% aniline blue in a 16-well tray. Samples were then embedded in 50% glycerol and observed under a Nikon fluorescence microscope with DAPI filter. Images were analysed using IMARIS 8.0 software for quantifying the number of callose deposits/ mm^2 leaf area.

RNA extraction and gene expression analysis by qRT-PCR

Total RNA extraction and cDNA synthesis was performed as described later. All qPCR primers used in this study are listed in Table supplement 1. qPCRs were performed in QuantStudio 6 Flex Real-Time PCR system (Applied Biosystems) with 5X HOT FIREPol® EvaGreen® qPCR Mix Plus (ROX) (Solis BioDyne) according to the manufacturer's instructions. All qPCR experiments were replicated thrice with three biological and technical replicates (n=3). *SAND* (At2g28390) expression was used as internal control (Kim et al., 2010). Relative expression was calculated according to the PCR efficiency$^{-\Delta\Delta Ct}$ formula. Expression differences were normalised to Col-0 and plotted as fold change.

Protein extraction and western blotting

For immunoblotting, leaf tissues collected at indicated time points/treatments were homogenised in protein extraction buffer [50 mM Tris HCl (pH 8.0), 8M Urea, 50 mM NaCl, 1% v/v NP-40, 0.5% Sodium deoxycholate, 0.1% SDS and 1 mM EDTA] containing 20 mM *N*-ethylmaleimide (NEM), 1X plant protease inhibitors cocktail (Sigma Aldrich) and 2% w/v Polyvinylpyrrolidone (PVPP). The homogenates were clarified by centrifugation, mixed with 2X Laemmli buffer, proteins separated by SDS-PAGE and then transferred onto polyvinylidene fluoride (PVDF) membrane by wet transfer method. The membrane was blocked with 5% non-fat skim milk and western blots performed with indicated primary antibodies [anti-SNC1 (Abiocode), anti-PR1 or anti-PR2 (Agrisera), or anti-SUMO1 (Abcam), or anti-Actin C3 antibodies (Abiocode)] in 1X TBST at 4^0C for overnight. Comparable protein loading was determined by Ponceau S staining of Rubisco subunit. Blots were washed thrice with 1X TBST

266 and then incubated at RT for one hour with appropriate horse-radish peroxidase (HRP)-
267 conjugated secondary antibodies. The blots developed using ECL™ Prime western blotting
268 system (GE Healthcare) and visualised in ImageQuant™ LAS 4000 biomolecular imager (GE
269 Healthcare).
270

271 **SUMOylome changes in response to SA or heat shock treatments**
272 For salicylic acid (SA)-induced SUMOylation changes, extracts were obtained from 2-weeks
273 old seedlings treated with 2 mM SA or buffer alone for 1 hour (Bailey et al., 2015). For heat
274 shock treatments, 2-weeks old seedlings were incubated at 37^0C or at RT for 30 minutes. Total
275 protein extracts were immunoblotted as described earlier.
276

Construction of Plasmid clones
To generate cDNA clones of Arabidopsis *SUM1* and *SUM3* genes, total RNA isolated from
Col-0 plants (RNAiso Plus; Takara) was reverse transcribed (iScript™ cDNA Synthesis Kit;
Bio-Rad) according to manufacturer's instructions. Specific *SUM1* and *SUM3* (GG and AA
forms) sequences were amplified (Phusion High-Fidelity DNA Polymerase; ThermoFisher)
from the cDNA using the primers listed (Table supplement 1). PCR products were cloned into
the Gateway entry vector *p*DONR201 and subsequently into *p*MDC-cCFP and *p*MDC-nVenus
(BiFC destination vectors; (Bhattacharjee et al., 2011) using the Clonase™ Recombination
system (ThermoFisher). Confirmed BiFC clones were then electroporated into *A. tumefaciens*
GV3101 strain.
For dimerization studies, cDNAs of *SUM1AA* or *SUM3AA* were cloned as EcoRI-SalI fragment
into *p*ASK-IBA16 vector (N-terminal Strep-tag II affinity tag; Neuromics). For *in vitro* paralog
influence on SUMOylation assays *SUM1GG* or *SUM3GG* were cloned as EcoRI-SalI fragment
into *p*ASK-IBA16 vector. These clones were named accordingly as Strep-SUMO1GG, Strep-
SUMO1AA, Strep-SUMO3GG, or Strep-SUMO3AA, respectively. For His-tagged versions,
*p*CDFDuet vector harbouring *SUM1GG* (*p*KT-973), *SUM1AA* (*p*KT-1017), *SUM3GG* (*p*KT-
975), or *SUM3AA* (*p*KT-1788) were used (Okada et al., 2009). These plasmids were a kind gift
from Prof. Katsunori Tanaka, Kwansei Gakuin University, Japan).
295

296 **Cloning and generation of *SUM3p-His-StrepII-SUM3g* transgenic plants**
297 Genomic DNA extracted from Col-0 plants were used for PCR of a ~1.3kbp genomic fragment
298 with KpnI-SUM3p For/XbaI-SUM3-UTR Rev primers. Using the amplicon as a template, two

independent PCRs were performed by using KpnI-SUM3p For/His-StrepII Rev or His-StrepII For/ XbaI-SUM3-UTR Rev primers combinations respectively. Individual PCR amplifications were used for overlap PCR with KpnI-SUM3p For/XbaI-SUM3-UTR Rev primers. The product was cloned into XbaI-KpnI site of the binary vector *pBIB-Hyg* (Becker, 1990). Generated clones were sequenced and confirmed. The binary vector was introduced in *Agrobacterium* strain GV3101 via electroporation. Pool of *sum3-1* plants were transformed via floral-dip method (Clough and Bent, 1998). Transgenic *sum3-1:HS-SUM3* plants were selected on Hygromycin containing medium, propagated through T3 generations to identify lines containing single locus but homozygous *sum3-1:HS-SUM3* transgenes. Subsequently, the plants were used for assays as indicated.

Bimolecular Fluorescence Complementation (BiFC) assays

All BiFC assays were performed according to (Bhattacharjee et al., 2011). Briefly, *Agrobacterium* cells harbouring the indicated BiFC vectors were induced with 150 µM Acetosyringone for 4 hours, equal bacterial density suspensions of desired cCFP and nVenus BiFC combinations made and then infiltrated in leaves of 4-weeks old *N. benthamiana* plants. At 48-hpi, a small section of infiltrated area was imaged under a SP8 Leica confocal microscopy system using FITC filter (488-nm Argon Laser).

In vitro **SUMO binding assays**

SUMO-binding assays were performed with recombinant protein expressed in *E. coli*. Expression was induced with 200 µg/L Anhydrotetracycline (for *pASK-IBA16* clones) or with 0.5 mM IPTG (for *pCDFDuet* clones) in BL21 (DE3) for overnight at 25^0C. Cell pellet was resuspended in lysis buffer (50 mM NaH_2PO_4, 300 mM NaCl, 10 mM Imidazole; pH 8.0) and sonicated. Equal volumes of His- and Strep-II tagged-SUMO combination supernatants were mixed and incubated at 4^0C with rotation for 2-3 hours to allow binding. Ni^{2+}-NTA beads (Qiagen) were added to the mix and incubated further for 2 hrs. His-tagged protein pull down was performed under native conditions according to manufacturer's instructions. Immunoblots were performed with anti-His-HRP (Santa Cruz Biotech) or anti-StrepII-HRP (IBA Lifesciences) antibodies.

In vitro **SUMOylation reconstitution assays**

The His-tagged constructs used were according to (Okada et al., 2009). Generation of Strep-
tagged SUMO paralog clones have been described earlier. *E. coli* BL21 (DE3) cells containing
both *p*KT-973 (SUMO1GG + SCE1) and *p*KT-978 (SAE2 + SAE1) was transformed with
either empty vector (*p*ASK-IBA16), Strep-SUMO3GG, or Strep-SUMO3AA plasmids.
Similarly, BL21 (DE3) cells containing *p*KT-975 (SUMO3GG + SCE1) and *p*KT-978 (SAE2
+ SAE1) was either transformed with either empty vector (*p*ASK-IBA16), Strep-SUMO1GG,
or Strep-SUMO1AA plasmids. As controls, conjugation-deficient SUMO1AA (*p*KT-1017) or
SUMO3AA (*p*KT-1788) in combination with *p*KT-978 was used. Transformed cells were
induced with 200 μg/L Anhydrotetracycline and 0.5 mM IPTG for overnight at 25^0C. After
harvesting, cell pellets were lysed and immunoblotted with anti-His or anti-Strep antibodies.

Statistical Analysis
For all gene expression experiments, Student's t-test was performed to check significance and
denoted by one, two and three asterisks indicating *p-value* <0.05, <0.01, and 0.001,
respectively. For growth curve and other assays, ANOVA was performed to check statistical
significance in growth of bacteria among different genotypes and indicated by alphabets e.g. a,
b, c, d, e etc. which depict statistical difference from each other at *p-value* <0.001.

RESULTS

SUM1/2 **genetically function as negative immune regulators whereas** *SUM3* **promotes**
defense responses

To study individual *Arabidopsis* SUMO paralog influences on defenses, we obtained previously
characterized knockout mutant lines of *sum1-1*, *sum2-1* and *sum3-1* (Saracco *et al.*, 2007; van
den Burg *et al.*, 2010). Wild-type (Col-0) and mutant plants were propagated either in short day
(SD) or long day conditions (LD) as indicated for respective assays. Interestingly, in all
propagation regimes and especially in SD conditions, we noticed clear growth defects in *sum1-*
1, but not *sum2-1* or *sum3-1* plants. The *sum1-1* mutant was developmentally dwarf with
elongated leaves, reduced fresh tissue mass and increased trichomes density although with
normal architecture compared to Col-0, *sum2-1* or *sum3-1* plants (Figure 1A,B-figure
supplement 1). Although identical *sum1-1* mutant has been utilized previously the
morphological defects we observed have not been reported earlier. The reason for this
discrepancy is not clear and we speculate differences in soil compositions or growth variations

as possible contributing factors. Nevertheless, the phenotypic attributes of *sum1-1* were novel
and we present evidences in subsequent sections that genetically link these defects solely to
loss of *SUM1*.

We then investigated defense responses in the *sum* mutants utilizing the standard bacterial
growth assays with virulent and avirulent *PstDC3000* strains. In Col-0, *PstDC3000* is virulent
and triggers basal or PAMP-triggered immunity (PTI). In contrast, *PstDC3000* harboring
avrRps4, avrRpm1, or hopA1 effector is avirulent, triggering effector-triggered immunity
(ETI), mediated by the cognate resistance gene *RPS4*, *RPM1* or *RPS6*, respectively
(Bhattacharjee *et al*., 2011). The virulent *PstDC3000* strain we used harbors the plasmid
backbone into which the avirulent effectors were cloned and hence named as *PstDC3000*
(*empty vector; EV*) in subsequent sections. Fully expanded leaves of 3-4 weeks-old SD-grown
plants were infiltrated with the indicated *PstDC3000* strains and the growth of the bacteria
measured at 0- and 3-days post-infiltration (dpi). When compared to Col-0, total bacterial
growth was reduced almost 10-fold in the *sum1-1* and ~8-fold in *sum2-1* plants whereas *sum3-
1* allowed more bacterial colonization (~8-fold higher) (Figure 1C). These results implied that
SUM1/2 and *SUM3* function as negative and positive regulators of PTI, respectively. When
challenged with avirulent *PstDC3000* expressing *avrRps4* or *hopA1*, two TNL-specific ETI-
eliciting effectors, lower bacterial accumulation than Col-0 persisted in the *sum1-1* and *sum2-
1* plants (Figure 1D- figure supplement 2A). The *sum3-1* plants were hyper-susceptible to these
avirulent infections. Curiously, modest but consistent difference with less bacterial
accumulation in *sum1-1* than *sum2-1* was observed to virulent or *PstDC3000(hopA1)* but not
to *PstDC3000(avrRps4)* challenges. Further on, increased callose deposits, a well-established
defensive phenomenon (Voigt, 2014), were more pronounced in *sum1-1* (~3-fold) and *sum2*
(~2-fold) than Col-0 in response to both virulent as well as avirulent (*avrRps4*) *PstDC3000*
(Figure 1E- figure supplement 3). Although previous studies suggest redundant roles of
SUM1/2 in defenses (van den Burg et al., 2010), as our results implicate that their degree of
contribution however may slightly vary. Likely indicative of enhanced susceptibility to
PstDC3000, lower levels of callose than Col-0 accumulated in *sum3-1*. Remarkably, ETI in all
plants to avirulent *PstDC3000* (*avrRpm1)* remained comparable suggesting that CNL-type
responses are not affected by the loss of individual SUMO paralogs (Figure supplement 2B).
Indeed, as reported earlier *avrRpm1*-mediated HR is not affected in any *sum* mutants (van den
Burg et al., 2010). Overall, we identify a partial redundancy between *SUM1/2* with a
concomitant antagonism to *SUM3* functions in immune responses to *PstDC3000* strains.

Basal levels of SA and SA-responsive defense markers are upregulated in *sum1-1* and *sum2-1*

Significant increases in salicylic acid, both free (SA) and glucose-conjugated (SAG), mediate signal responses against *PstDC3000* (Dempsey et al., 2011). A SUMOylation-deficient and enhanced resistant *siz1-2* plants have elevated SA/SAG due to upregulated expression of SA-biosynthesis gene *SID2/ICS1* (*SALICYLIC ACID INDUCTION DEFICIENT2/ ISOCHORISMATE SYNTHASE1*) (Cui et al., 2016; van den Burg et al., 2010). To determine whether SA/SAG perturbations reflect the immune outcomes, we measured their relative levels in the different *sum* mutants. Remarkably, both *sum1-1* and *sum2-1* plants contained significantly elevated whereas the hyper-susceptible *sum3-1* had lower SA/SAG, respectively than Col-0 (Figure 2A,B). We noted that SA/SAG levels in *sum1-1* plants were slightly higher than *sum2-1*, perhaps indicative of its modestly higher degree of basal and TNL-type immunity. Transcripts of *SID2/ICS1*, defense-associated markers *PR1* and *PR2,* and PR2 protein levels were upregulated in both *sum1-1* and *sum2-1* whereas in *sum3-1* plants these were significantly lower than Col-0 (Figure 2C,D). Likewise, increased expressions of PTI markers *FLG22-INDUCED RECEPTOR LIKE KINASE 1* (*FRK1*) and *WRKY TRANSCRIPTION FACTOR 29* (*WRKY29*) (Sardar et al., 2017) than Col-0 were detected in both *sum1-1* and *sum2-1* plants (Figure 2E). Relative to Col-0, *FRK1* expression remained unaltered in *sum3-1*, whereas *WRKY29* was drastically reduced suggesting that *SUM3* promotes expression of only a subset of PTI-markers. Accumulation of a well-known SA-responsive TNL-type R protein *SUPPRESSOR OF npr1-1 CONSTITUTIVE 1* (*SNC1*) is enhanced in several auto-immune mutants (Kim et al., 2010; Li et al., 2010). Remarkably, both *sum1-1* and *sum2-1* displayed upregulated *SNC1* expressions whereas in *sum3-1* the transcript levels remained comparable to Col-0 (Figure 2E). Since *FRK1*, *WRKY29* and *SNC1* are SA-inducible, upregulated SA-signaling sectors likely contribute to the enhanced basal and TNL-type immunity in *sum1-1* and *sum2-1* while these defenses are deficient in *sum3-1*. Also taking into account that *SUM3* but not *SUM1* or *SUM2,* is SA-inducible (van den Burg et al., 2010), we reasoned that enhanced resistance in *sum1-1* or *sum2-1* may be due to elevated *SUM3* expression and its role as a positive immune regulator. Indeed, we detected ~3- and 2-fold higher *SUM3* transcripts than Col-0 in *sum1-1* and *sum2-1*, respectively (Figure 2F). Relative transcript levels of *SUM1* in *sum2-1*, or *SUM2* in *sum1-1* however remain unaltered. Overall our results identify redundant but unequal roles of *SUM1* and *SUM2* in suppressing SA-dependent defenses that include expression of PTI markers, *SNC1* and *SUM3*.

Expression of genomic copies of *SUM1* or *SUM3* complement immune response alterations in the respective *sum* mutants

Phenotypic abnormalities of *sum1-1* plants necessitated us to genetically link the defects to loss of *SUM1*. Although our *sum1-1* mutant is identical to earlier reports (Saracco et al., 2007; van den Burg et al., 2010), to validate our observations, we utilized the transgenic *His-H89R-SUM1* line that express a His$_6$-tagged genomic *SUM1* variant (H89R) from its native promoter in a *sum1-1 sum2-1* background (Miller et al., 2010). These plants as reported earlier not only complement the lethality of double homozygous *sum1-1 sum2-1* mutant but also functionally mirror wild-type SUMO1 SUMOylome changes upon heat stress. From the *His-H89R-SUM1* plants, the *sum2-1* mutation was segregated out by generating F2 populations upon crossing with *sum1-1*. Henceforth, we termed these plants as *sum1-1:His-H89R-SUM1*. We constantly observed that growth defects we noted earlier were always linked to homozygous *sum1-1* genotypes in absence of the *His-H89R-SUM1* transgene. Thus, a functional *SUM1* abolishes the growth defects apparent for *sum1-1* plants (Figure supplement 4). We noted that the *sum1-1:His-H89R-SUM1* plants had slightly elevated *SUM1* (but not *SUM2*) transcripts compared to Col-0 (Figure 3A). Implicatively, as reported earlier for *SUM1* over-expression (van den Burg et al., 2010), SA/SAG levels although considerably reduced in comparison to the *sum1-1* parent, were still maintained at slightly higher levels than Col-0 (Figure 3B). Increased SA/SAG levels in *sum1-1:His-H89R-SUM1* plants also resulted in upregulated *SUM3* transcripts and higher PR1 proteins in comparison to Col-0 (Figure 3A,C). In pathogen growth assays using the virulent *PstDC3000(EV)*, enhanced defenses noticeable for *sum1-1* was abrogated to Col-0 levels in the *sum1-1:His-H89R-SUM1* plants (Figure 3D). Interestingly, the avirulent *PstDC3000(hopA1)* challenge although diminished enhanced resistance relative to the *sum1-1* parent, nevertheless these plants displayed stronger defenses than Col-0 (Figure 3D). This likely is attributed to higher than Col-0 levels of SA-regulated defense networks. These results unambiguously support the loss of *SUM1* as the cause of enhanced resistance and phenotypic defects in *sum1-1*.

We also generated two independent complemented lines of *sum3-1* to validate immune deficiencies due to the loss of *SUM3*. These transgenic plants express native promoter-driven His$_6$-StrepII-tagged genomic fragment of *SUM3* (*sum3-1:HS-SUM3*). Although owing to its low abundance (van den Burg et al., 2010) the expression of His$_6$-StrepII-SUMO3 proteins remained undetectable in immunoblots, transcripts of *SUM3* drastically reduced in the *sum3-1*,

were restored to equivalent or slightly lower (~0.5X) than Col-0 levels in Line #1 and Line #2,

respectively (Figure 3A). Relative abundance of *SUM1* or *SUM2* remained unaffected in both

lines. For further assays, we therefore continued with *sum3-1:HS-SUM3*#1. We observed that

deficiencies in SA/SAG or PR1 protein accumulations inherent to *sum3-1* were reinstated to

Col-0 levels in the *sum3-1:HS-SUM3#1* plants suggesting functional complementation by the

transgene (Figure 3B,C). Pathogen growth assays with either virulent *PstDC3000(EV)* or

avirulent *PstDC3000(hopA1)* strains regained Col-0 levels of resistance in *sum3-1:HS-SUM3#1*

(Figure 3D). These data lead us to conclude that defensive deficiencies in *sum3-1* is indeed due

to loss of *SUM3*.

Constitutively active defenses in *sum1-1* and *sum2-1* caused rapid induction of immunity

To further establish that upregulated or dampened defense marker expressions in *sum1-1*, *sum2-*

1 or *sum3-1*, respectively contribute to corresponding immune outcomes we challenged these

plants with virulent *PstDC3000(EV)* or avirulent *PstDC3000(avrRps4)* strains. Leaf extracts

harvested at 6-, 12-, and 24-hpi were used to compare accumulation kinetics of *PR1/PR2*

transcripts and proteins (Figure 4- figure supplement 5). Basal level (0-hpi) of PR1 in Col-0,

sum2-1, or *sum3-1* was below detection limit whereas in *sum1-1* some enhancements were

noticeable (Figure 4A). At 6- or 12-hpi although Col-0 or *sum3-1* barely accumulated sufficient

PR1/PR2 in response to both pathogen challenges, both *sum1-1* and *sum1-2* had markedly

elevated levels of these proteins (Figure 4B,C). A similar trend continued at 24-hpi wherein in

Col-0, PR2 but not PR1 proteins levels, almost matched *sum1-1* or *sum2-1*. Most significantly,

even at the 24-hpi *sum3-1* plants were deficient in accumulating wild-type levels of PR1 or

PR2. Real-time accumulation kinetics of *PR1/PR2* transcripts to both bacterial infiltrations in

general mirrored the corresponding protein levels although in some instances a direct

correlation was not apparent (Figure supplement 5). Although the reason for this discrepancy

is not clear, suggested role of SUMOylation in selective translation and/or its crosstalk with

other PTMs that modulate protein synthesis or stability may likely be the cause (Chen et al.,

2014; Watts et al., 2014). Nevertheless, these data provide strong molecular evidence of

constitutive SA-regulated defenses in *sum1-1* and *sum2-1* conferring their enhanced resistance

to *PstDC3000* strains. Contrastingly, delayed induction of defenses upon a pathogen challenge

likely makes *sum3-1* hypersusceptible.

Enhanced defenses in *sum1-1* is SA-dependent

To genetically determine whether constitutive SA-signaling routes impart increased defenses

to *sum1-1*, we generated *sum1-1sid2-1* and *sum1-1eds1-2* double mutants. A *sid2-1* plant

harbors a null mutation in *SID2/ICS1* whereas an *eds1-2* expresses a non-functional EDS1

(ENHANCED DISEASE SUSCEPTIBILITY1), a central player orchestrating SA-mediated

defenses (Wildermuth *et al.*, 2001; Cui *et al.*, 2016). In segregating F2 populations, we noted

that phenotypic defects always associated with homozygous *sum1-1* allele provided at least one

functional copy of *EDS1* or *SID2* were present (data not shown). Remarkably, the reduced leaf

mass apparent in *sum1-1* was improved in *sum1-1eds1-2* and *sum1-1sid2-1* plants (Figure 5A-

figure supplement 1). While *sum1-1eds1-2* achieved wild-type mass, abolishing *SID2/ICS1* in

sum1-1 although ameliorated growth defects but these plants still retained *sid2-1*-characterstics

including smaller and paler leaves than Col-0 (Figure 5A- figure supplement 1). Interestingly,

both *sum1-1sid2-1* or *sum1-1eds1-2* had wild-type trichome densities unlike the *sum1-1* parent

thus associating these defects to perturbations in SA-regulated networks (Figure supplement

6A).

Loss of either *EDS1* or *SID2/ICS1* remarkably abolished accumulated PR2 proteins and

upregulated *PR1*, *PR2*, *FRK1* and *WRKY29* transcripts in *sum1-1* (Figure 4A; 5B-figure

supplement 6B). Further on, pathogen growth assays using virulent and avirulent *PstDC3000*

strains demonstrated that *sid2-1* or *eds1-2* mutations were epistatic to *sum1-1* abolishing not

only the increased defenses but also conferring hyper-susceptibility to the respective double

mutants towards virulent *PstDC3000(EV)* or avirulent (*avrRps4-* or *hopA1*-expressing) strains

(Figure 5C,D- figure supplement 6C). Neither *eds1-2* nor *sid2-1* mutation affected resistance

towards avirulent *PstDC3000* (*avrRpm1*) which remained comparable to *sum1-1* or Col-0

(Figure supplement 6D). As expected, loss of *EDS1* or *SID2/ICS1* prevented accumulation of

PR1 or PR2 proteins during *PstDC3000* infections (Figure 4B,C). With these assays, we

identify that enhanced resistance in *sum1-1* is undoubtedly due to constitutive SA-routed

defense signaling. A *sum2-1eds1-2* double mutant, we also generated, mimicked *sum1-1eds1-*

2 hyper-susceptible outcomes in disease assays (Figure supplement 7). Overall, our results

provide several parallel lines of evidences suggesting *SUM1/2* intersections on SA-mediated

immune signaling.

sum3-1 alleviates enhanced defenses in sum1/2 mutants

Since our data suggested antagonistic involvement of *SUM1/2* and *SUM3* as negative or

positive defense regulators, respectively we surmise a crosstalk between these paralogs. To

investigate this, we generated *sum1-1sum3-1* double mutant by genetic crossing. Unlike *sum1-1eds1-2* or *sum1-1sid2-1*, only partial restoration of *sum1-1* growth defects was observed in *sum1-1sum3-1* plants (Figure 6A-figure supplement 8A,B). These observations hinted that *SUM3* is mildly responsible for *sum1-1* growth anomalies. Increased trichome densities noticed in *sum1-1* however remained unaffected by the loss of *SUM3* (Figure supplement 6A). Because *SUM3* affects basal SA accumulation (Figure 2A,B), we tested defense outputs are altered in *sum1-1 sum3-1*. Elevated SA levels in *sum1-1* were marginally reduced in *sum1-1 sum3-1* plants. Indeed, the expression of *SID2/ICS1* remained comparable between *sum1-1* and *sum1-1 sum3-1* plants (Figure 2C). Additionally, *PR1* or *PR2* transcripts in *sum1-1 sum3-1*, were intermediate between *sum1-1* and Col-0. These data imply *SUM3* does not affect SA-biosynthesis but modulates positive feedback loop of SA-signaling wherein it intersects with *SUM1* role as a transcriptional repressor of *SID2/ICS1*, *PR1* or *PR2*. As is therefore expected, loss of *SUM3* reduced PR1 or PR2 levels in *sum1-1* (Figure 4A). Not the least, we also demonstrate that increased SNC1 accumulation in *sum1-1* plants although reduced in *sum1-1 sum3-1*, are still maintained higher than Col-0 levels (Figure 6B).

In pathogen growth assays, either virulent *PstDC3000(EV)* or avirulent *PstDC3000 (hopA1)* strains accumulated to intermediate levels in *sum1-1 sum3-1* leaves, particularly higher than *sum1-1*, but significantly lower than in Col-0 or *sum3-1* (Figure 6C- figure supplement 8C). Interestingly, wild-type level of resistance was observed in *sum1-1 sum3-1* to the avirulent effector *avrRps4* (Figure 6D). Growth of avirulent *PstDC3000 (avrRpm1)* was not affected in *sum1-1sum3-1* and remained comparable in all plants (Figure supplement 8D). As investigated earlier, in response to virulent and avirulent *PstDC3000(avrRps4)* infections and unlike *sum1-1*, *sum1-1 sum3-1* remained deficient in induction of PR1 or PR2 at 6-hpi (Figure 4B,C- figure supplement 9A,B). However, we observed that at later time points (12- and 24-hpi) loss of *SUM3* did not affect the rapid induction of PR1/PR2 in *sum1-1*. This is a sharp contrast to *sum3-1* plants which even at 24-hpi accumulated very less PR1/PR2 proteins. Thus, although *SUM3* is deemed essential for upregulation of PR1/PR2 upon a pathogen attack, *sum1-1* plants eventually overcome this requirement. Taken together, we principally support that *SUM3* promotes *PR1*/*PR2* transcription downstream of SA as suggested earlier (van den Burg et al., 2010). We provide further molecular evidence that *SUM1* suppression of defenses likely is upstream of SA and may involve transcriptional repression of *SID2/ICS1* and its subsequent consequences on SA-responsive markers such as *PR1*, *PR2*, *FRK1*, *WRKY29* and *SNC1* among others.

Adjacent chromosomal arrangement of *SUM2* and *SUM3* in Col-0 genome suggest that they likely arose by a tandem duplication event (Hammoudi et al., 2016). Hence, unlike a *sum1-1 sum3-1* double mutant, a *sum2-1 sum3-1* is difficult to obtain by genetic crossing. To test whether *SUM3* also intersects on *SUM2* function as a negative immune regulator, we generated *amiR-SUM2 sum3-1* plants by crossing the two parental lines. The *amiR-SUM2* plants reported earlier contains specific knockdown of *SUM2* and mimics the null mutant *sum2-1* (van den Burg et al., 2010). *SUM1* or *SUM3* transcripts remain unaffected in these lines. Plants homozygous for *sum3-1* and containing *amiR-SUM2* transgene have significantly downregulated *SID2/ICS1, PR1* and *PR2* levels in comparison to a *sum2* mutant (Figure 2C). These transcripts however are still elevated than Col-0 plants implicating that *SUM3* functions impinge partially on *SUM2* role as a transcriptional repressor. This was also supported in pathogen challenges wherein growth of both virulent and avirulent *PstDC3000* strains were intermediate between resistant Col-0 and hyper-resistant *sum2* mutant (Figure supplement 7B,C). Taken together, whether the functional antagonism in immune regulatory roles of *SUM1/2* and *SUM3* affect immune outcomes in plants.

SUMO3, but not SUMO1, can form non-covalent homo-dimers

Conjugation proficiencies *in planta* of SUMO paralogs are clearly different. Upon SA application, drastic increase in SUMO1/2 conjugates are observed whereas target modifications by SUMO3 is barely detectable even though *SUM3* expression, unlike *SUM1/2* is SA-inducible (Bailey et al., 2015; van den Burg et al., 2010). A distinct difference in the protein sequence of these paralogs is the presence of a predicted SIM motif in SUMO3 but not SUMO1/2 (Figure supplement 10). We utilized Bi-molecular Fluorescence Complementation (BiFC) assays to test homo- and hetero- non-covalent associations between SUMO1 and SUMO3. The BiFC vectors introduced via *Agrobacterium*-mediated transient transformation of *N. benthamiana* leaves expressed split YFP fusions of either SUMOylation-proficient (GG) or -deficient (AA) SUMOs wherein the C-terminal di-glycine (GG) residues were kept intact or mutated to di-alanine (AA), respectively (Figure 7A). Co-expression of only SUMO1GG/SUMO1GG, but not SUMO1GG/SUMO1AA or SUMO1AA/SUMO1AA showed reconstitution of the split YFP protein. Although positive BiFC indicate non-covalent interactions among protein partners, the lack of fluorescence in SUMO1AA/SUMO1AA suggests that SUMO1GG/SUMO1GG combinations and not homo-oligomers and likely reflect covalent polySUMO1-conjugates wherein the split fluorescent fusion proteins are in allowed proximity for reconstitution.

Remarkably, all BiFC combinations of SUMO3 with itself, regardless of GG- or AA-forms, showed YFP fluorescence. Since SUMO3 lacks poly-SUMOylation properties (Castaño-Miquel *et al.*, 2011), we reason that the observed associations are non-covalent SUMO3 oligomers. To test whether SUMO3 binds SUMO1 non-covalently, combinations of SUMO1GG or AA were co-expressed with SUMO3GG or AA. While SUMO1GG showed clear fluorescence when expressed with either SUMO3GG or AA, similar combinations of SUMO1AA did not. These observations are suggestive that only a conjugable-SUMO1 likely present on a target may achieve permissible molecular proximity *in vivo* to a SUMO3 that may be either bound to a SIM (as in SUMO1GG/SUMO3AA pair) or attached covalently to the same target at a separate SUMOylation motif (as in SUMO1GG/SUMO3GG pair). Evidences from human cell lines demonstrate that ortholog of *Arabidopsis* SUMO3, HsSUMO1 'cap' poly-SUMO chains of the SUMO1 ortholog HsSUMO2 (Matic et al., 2013; Sharma et al., 2013). Whether *Arabidopsis* SUMO3 regulates poly-SUMO1/2 chain lengths remain a promising possibility to explore further. Since no interaction was detected with SUMO1AA/SUMO3AA, it is clear that SUMO3 and SUMO1 do not interact non-covalently. To validate further, we performed *in vitro* binding assays with tagged recombinant SUMO1AA/SUMO3AA proteins expressed in *E. coli*. Enrichment of His-SUMO1AA via Ni^{2+} beads failed to co-elute either Strep-SUMO1AA or Strep-SUMO3AA suggesting that SUMO1-SUMO1 homo- and SUMO1-SUMO3 hetero-dimers cannot form *in vitro* (Figure 7B). Homodimers of SUMO3AA however were detected by the presence of Strep-SUMO3AA in His-SUMO3AA-enriched eluates. These results support our hypothesis that BiFC interactions observed for SUMO1GG with SUMO3GG/AA may likely suggest close proximity, but not direct binding, of SUMO1 and SUMO3 *in planta*.

SUMO3 regulates *SUMO1/2* conjugation efficiencies upon various stress exposures

With the above results, it is encouraging to speculate that SUMOylation efficiencies may be modulated by intersecting SUMO1-SUMO3 functions. Since *SUM3* partially modulates *sum1-1* defenses, we investigated dynamic changes in SUMO1/2-SUMOylome in Col-0, *sum1-1*, *sum3-1*, or *sum1-1 sum3-1* upon a pathogen challenge. Leaf tissues either mock-treated or infiltrated with *PstDC3000*(*EV*), were collected at 24-hpi and processed for selective qPCRs or immunoblots for SUMO1/2-conjugates. The polyclonal anti-SUMO1 antibody we used cannot distinguish SUMO1/2 paralogs and the lack of anti-SUMO3 antibodies prevented us for analyzing SUMO3-conjugates. We observed that while a pathogen exposure caused remarkable increase in SUMO1/2-conjugates in Col-0, for *sum1-1* these was considerably lower (Figure

8A). These results suggested that SUMO1- rather than SUMO2-conjugates are not only more prevalent but also undergo massive increment upon pathogen treatment. Surprisingly, the conjugates are also distinctly lower in uninfected *sum3-1* and are considerably less upregulated than Col-0 in response to the pathogen infection. As is expected, in *sum1-1 sum3-1* extracts barely any SUMO-conjugates are detectable pre- or post-infection. With these results, we identify clear intersection of SUMO3 functions on SUMO1/2-conjugation efficiencies and its perturbations during a pathogen attack. We also observed that both *sum1-1:His-H89R-SUM1* and *sum3-1:HS-SUM3* plants achieved Col-0 levels of SUMO1/2-cojugates upon *PstDC3000*(*EV*) infection thus further validating the complete functional complementation of *sum1-1* or *sum3-1*, respectively (Figure supplement 11).

To determine whether changes in conjugation efficiencies are due to differential expression of SUMO paralogs and/or SUMOylation-associated genes, we investigated their relative expressions in Col-0 and the *sum* mutants (Figure 8B,C- figure supplement 11B). We observed that *SUM3*, but not *SUM1/2*, was upregulated upon a *PstDC3000*(*EV*) challenge. This is in accordance with *SUM3* being SA-inducible (van den Burg et al., 2010). Additionally, we noticed that the enhanced *SUM3* expression in *sum1-1* was further boosted in pathogen-exposed samples. Curiously, for the investigated SUMOylation-associated genes although *sum1-1* mutation did not significantly alter their basal expression levels, several of these (*SAE2*, *SIZ1*, *HYP2*, *ESD4*, *ELS1*, and *OTS1*) were down-regulated in *sum3-1* plants. Thus, the modest reduction in global SUMOylome in *sum3-1* we observe may be attributed to *SUM3*-dependent modulation of expression of these genes. Interestingly, *sum1-1 sum3-1* plants showed slight upregulation of *SAE2*, *SCE1*, *SIZ1*, *HPY2*, *ELS1* and *OTS2* in mock-treatment supporting our earlier claim that *sum1-1* plants with constitutively active defenses and elevated SA levels, overcome the requirement of *SUM3* for the expression of these genes. We also noted that upregulation of *SAE2*, *SIZ1*, *ELS1* or *OTS1* that occurred only modestly in Col-0 upon *PstDC3000*(*EV*) infection were further aggravated in *sum1-1*, but intermediate in *sum3-1* (Figure 8B,C). These enhancements are maintained in *sum1-1 sum3-1* plants and are likely *SUM3*-independent. Remarkably, *sum3-1* plants down-regulated the expression of *SUM1* and *SUM2* only upon pathogen-treatment highlighting that the positive immune function of *SUM3* may involve transcriptional repression of negative defense regulators such as *SUM1/2*. Overall, our data lead us to speculate that immune responses recruit *SUM1-SUM3* crosstalks to modulate the expression of these genes at a transcriptional level thereby influencing host SUMOylome changes.

SAE2, SCE1, SIZ1 or EDS4 are direct SUMO1 targets and both SUMO1/3 interact *in planta*

with SCE1 and SIZ1 (Miller et al., 2010; Mazur *et al.*, 2019). These led us to test whether

SUMO1-SUMO3 intersections directly influence the efficiency of SUMO1/2-conjugate

formation through modulation of SUMOylation-associated protein functions. We utilized the

E. coli SIZ1-independent *Arabidopsis* SUMOylation reconstitution system (Okada et al., 2009).

Co-expression of His-SUMO1, and Strep-SUMO3 (GG or AA) along with SCE1, and SAE1/2

showed modest but clear increase in SUMO1-conjugates in comparison to extracts that lacked

Strep-SUMO3 (Figure supplement 12). Remarkably, a more dramatic increase in SUMO3-

conjugates when co-expressed with either Strep-SUMO1 GG or AA was also noted. From these

results, it is undeniably clear that SUMO1 and SUMO3 have a mutually beneficial role in

reciprocal SUMOylation process likely through modulation of SCE1 or SAE1/2 functions. The

significant decrease in SUMO1/2 conjugates in absence of *SUM3* (Figure 8A) provides

reasonable endorsement to our hypothesis.

SUMO1/2-conjugates are enhanced in response to heat shock (Saracco *et al.*, 2007; van den

Burg *et al.*, 2010). To test whether *SUM3* role intersect on these responses, we subjected Col-

0, *sum1-1*, *sum3-1*, or *sum1-1 sum3-1* plants to heat shock and analyzed SUMO1/2-conjugates

las well as relative expression levels of *SUM1/2/3* and several previously characterized heat

stress-responsive genes (Ohama et al., 2016) (Figure supplement 13). In accordance with

Saracco *et al.* (2007) we observe strong induction of SUMO1/2-conjugates in heat-treated Col-

0 extracts (Figure supplement 13A). These conjugates were low in heat-stressed *sum1-1* plants

reinstating that SUMO1-mediated SUMOylation is most predominant during heat stress.

Surprisingly, in *sum3-1* considerably reduced conjugates than Col-0 accumulated upon heat

shock. We especially noted that unlike *PstDC3000* challenges, *SUM1/2* transcripts were

significantly upregulated by heat-exposure, while *SUM3* induction was almost negligible

(Figure supplement 13B). Further on, transcriptional upregulation of *SUM1/2* was *SUM3*-

independent. Therefore, reduced accumulation of SUMO1/2-conjugates in heat-treated *sum3-1*

suggests that *SUM3* affects SUMO1-SUMOylation efficiencies at a post-transcriptional level.

With these results, we infer that although different physiological stresses may cause apparently

similar effects on global SUMO1/2-conjugates, the responsive routes to condition these are not

only distinct but also stress-specific. Interestingly, increased expression of several heat stress-

responsive markers such as *HsfA2*, *HSP22.0-ER* that are upregulated in Col-0 upon heat-

treatment, were hyper-elevated in *sum1-1* suggesting *SUM1* suppresses their expression (Figure

supplement 13C). In *sum3-1* plants, their fold-induction upon heat stress was relatively lower

than Col-0 implying that *SUM3* is partially responsible for their upregulation. Induction of other

tested markers such as *HSP18.2* or *HSP23.5-P* were unaffected in *sum1-1* or *sum3-1* plants

(Figure supplement 13D). Taken together, our results clearly identify *SUM1-SUM3* crosstalks

in responses to multiple stresses.

DISCUSSION

Functional overlaps between *Arabidopsis* SUMO1 and SUMO2 isoforms, likely prevent

noticeable phenotypic defects in the individual mutants under non-stressed conditions (Lois *et*

al., 2003; Colby *et al.*, 2006; Saracco *et al.*, 2007; van den Burg *et al.*, 2010). Interestingly,

reduction in SUMO-conjugates both basally as well as upon heat-stress is more prominent in

sum1-1 than *sum2-1* suggesting a *SUM1* predominance in these responses (S. A. Saracco et al.,

2007). Our observations therefore of phenotypic defects in *sum1-1* but not *sum2-1* plants may

reflect a similar developmental process preferentially regulated by *SUM1*. We convincingly

establish that constitutive upregulation of SA-regulated networks is the primary cause of these

defects in *sum1-1* (Figure supplement 6A). This is unlike *siz1-2* where the associated growth

abnormalities are EDS1- or SID2/ICS1-independent (Hammoudi et al., 2018). We deduce that

reduction in global SUMO1/2- conjugates apparently similar between *siz1-2* and *sum1-1* affect

downstream responses differently. At one instance, loss of *SIZ1* may affect SUMO-conjugation

for all isoforms, whereas the *sum1-1* clearly restricts only SUMO1 functions. Previous studies

propose SA antagonism or jasmonic acid (JA) promotion in increased trichome formation (An

et al., 2011). We reason that while direct application of SA or JA may affect trichome

production as reported, the loss of *SUM1* whose substrates include both JA and SA signaling

regulators such as SIZ1, TPL, or JAZs among others may impact trichome density through a

more complex SA-JA crosstalk (Miller et al., 2010; Srivastava et al., 2018). Further

investigations into relative SA-JA signaling routes perturbed in *sum1-1* may unravel this

mystery.

In context of defense responses, our data provide substantial molecular support not only to

individual *SUM1/2* or *SUM3* but also to their intersecting contributions as negative or positive

regulators, respectively of anti-bacterial basal and TNL-specific immunity. Firstly, bacterial

accumulations are strongly reduced in both *sum1-1* and *sum2-1*, with *sum1-1* displaying

stronger immunity than *sum2-1*, for both virulent as well as avirulent *PstDC3000*(*hopA1*)

infections. Although comparable immunity between *sum1-1* and *sum2-1* is observed for avirulent *PstDC3000*(*avrRps4*) challenges, this likely is due to relatively weaker ETI responses elicited by *AvrRps4* in comparison to *HopA1*-expresssing *PstDC3000* (compare Figure 1D and Figure supplement 2A). In contrast to *sum1/2* mutants, immunity in *sum3-1* is compromised implicating *SUM3* is essential for these defenses. Secondly, we demonstrate that impairment of SA-signaling routes due to loss of *EDS1* (*eds1-2*) or *SID2/ICS1* (*sid2-1*) abolish enhanced resistance in *sum1-1* and *sum2-1* plants providing support to genetic placement of *SUM1/2* as suppressors of these defenses (van den Burg et al., 2010) (Figure 8D). In this pathway, *SUM3* partially delegates SA-defenses likely through positive feedback mechanisms. Indeed, both *sum1-1* and *sum2-1* plants accumulate increased SA than Col-0, while in *sum3-1*, these are relatively lower. Correspondingly, transcripts of SA-biosynthesis *SID2/ICS1* and responsive markers such as *FRK1*, *PR1*, *WRKY29*, and *SNC1* are upregulated in *sum1-1* or *sum2-1*, and lower in *sum3-1*. Accumulation kinetics of PR1/PR2 proteins or transcripts (Figure 4- figure supplement 5) further validate primed or deficient SA-mediated defenses in *sum1-1* or *sum2-1* and *sum3-1*, respectively. The modest difference we note between *sum1-1* and *sum2-1*, in the upregulation of some of these SA-inducible markers, is suggestive of additive but unequal roles of *SUM1* and *SUM2* as negative immune regulators. Vertegaal *et al.*, (2006) identified only ~17% common targets between human SUMO1 and SUMO2 paralogs suggesting that they are partially redundant at best. Whether SUMOylation-targets distinct between *Arabidopsis* SUMO1 and SUMO2 isoforms affect immune amplitudes differentially awaits further studies.

We provide the first *in planta* evidence of partial SUMO3 moderation on SUMO1/2 functions in biotic, abiotic and developmental responses. Introducing *sum3-1* mutation not only attenuates reduced tissue mass but also subdues enhanced resistance of *sum1-1* supporting *SUM3* role in promoting defenses downstream of SA. Thus, elevated SA and *PR1*, *PR2*, or *SNC1* expressions are partially dampened in either *sum1-1* or in *amiR-SUM2* plants when *SUM3* is mutated (Figure 2). Intersections of *SUM3* functions on SUMOylation-proficiencies, are also best noted on SUMO1/2-conjugate intensities in response to different stresses. Although, enhancement upon heat shock or SA-treatments have been previously reported (Bailey et al., 2015; Miller et al., 2010), we first report their upregulation in *PstDC3000* exposures. Induction of several SUMOylation-associated genes such as *SAE2*, *SIZ1*, *ELS1*, or *OTS1* during *PstDC3000* challenges are influenced by *SUM1-SUM3* intersections and contribute to the corresponding defense outcomes (Figure 8). Recently, the activity of TPR1, a transcriptional co-repressor involved in suppression of negative defense regulators *DND1/2* was reported to be regulated in

a partial SA-dependent manner via SUMOylation by SIZ1 (Niu et al., 2019). SUMOylation-

deficient TPR1 not only is more enhanced in repressing *DND1/2* expressions, but also its over-

expression cause enhanced resistance than the SUMOylation-proficient TPR1. Overall, our

results raised the possibility that increased SUMO1/2 conjugates may reflect negative feedback

mechanisms to maintain immune responses as transitory thus regulating *SUM3* functions in

promoting defenses.

Functional intersections among SUMO isoforms are anticipated at multiple post-transcriptional

events. Amino acid distinctions at key conserved positions among the SUMO isoforms

influence their *in vivo* conjugation proficiencies (Castaño-Miquel *et al.*, 2011; schematically

summarized in Figure supplement 10). Aspartate63 (D^{63}) in *Arabidopsis* SUMO1 (D^{75} in

SUMO2) is replaced by Asparagine63 (N^{63}) in SUMO3, reducing its relative interaction

strength than SUMO1 with SCE1, thus lowering its poly-SUMO formation efficiencies. Other

residues also diverged in SUMO3 weaken thioester-bond formation and interaction with SAE1.

Whether *SUM3* upregulation during defences improve these propensities and hence alters host

SUMOylome outcomes although remains unknown, evidential support is obtained from several

studies. Mutants with decreased SUMOylation including *siz1-2* have increased SCE1 protein

abundance (Castaño-Miquel et al., 2017; Nukarinen et al., 2017; S. A. Saracco et al., 2007).

And *siz1-2* impairment in accumulating SUMO-conjugates during heat stress, is completely

recovered in *pial1 pial2 siz1-2* plants (Tomanov et al., 2014). Considering PIAL1/2 improves

SCE1 ability to form otherwise less efficient polySUMO3 chains, defense responses with

upregulated *SUM3* likely introduces competition between SUMO1 and SUMO3 for substrate

polySUMOylation thereby altering their fates via respective isoform-specific SUMO-targeted

E3 Ubiquitin ligases (STUbLs) (Tomanov et al., 2014). Indeed, *pial1/2* double mutants

although less stress-tolerant, have upregulated levels of proteins related to biotic stress

(Nukarinen et al., 2017; Tomanov et al., 2014). In *sum1-1*, the reduced global SUMO1/2-

conjugates upon pathogen or heat-shock challenges taken together with elevated *SUM3*

transcripts is suggestive of this occurrence and need to be explored further.

As substrates of Arabidopsis SCE1 and ESD4, candidates predominantly involved in RNA-

related processes such as nucleocytoplasmic transport, splicing, or turnover and chromatin-

modification including transcriptional activation/repression-associated proteins were identified

(Elrouby and Coupland, 2010). A majority of these candidates possessed predicted

SUMOylation motifs and were covalently modified by both SUMO1 or SUMO3. In

Arabidopsis, a SUMOylation-proficient SUMO1 is essential to covalently charge SCE1 at the catalytic site, promote its association with SIZ1, and form the ternary complex (SUMO-SCE1-SIZ1) (Mazur et al., 2019). Interestingly, the subcellular localization of this complex is determined by the specific SUMO paralog bound non-covalently at the SIM site distinct from the catalytic pocket of SCE1. Since SCE1 possess ability to distinguish substrates for SUMOylation (Tomanov et al., 2018), selection of substrates and formation of polySUMO chains therefore may be affected by fluctuations in relative SUMO1/3 levels, as has been previously suggested in animal studies (Capili and Lima, 2007; Duda et al., 2007). Our data showing reciprocal improvements in SUMOylation efficiencies *in vitro* regardless of conjugation-proficiencies of the influencing SUMO isoform may suggest towards one such consequence of SUMO1-SUMO3 crosstalks (Figure supplement 12). Likewise, constitutive activation of defenses due to over-expression of either conjugation-proficient (GG) or -deficient (ΔGG) SUMO isoforms may at least be partially attributed to this phenomenon (van den Burg et al., 2010).

SUMOylation machineries such as SAE2, SIZ1 and ESD4 also bind SUMOs non-covalently owing to their intrinsic SIMs (Cheong et al., 2009; Wang et al., 2010). It is postulated that spatio-temporal regulations of SUMOylation/de-SUMOylation coordination primarily by SIZ1 and ESD4 regulate steady-state levels of host proteins conjugated to SUMOs which surprisingly are fewer in numbers (reviewed in Elrouby, 2014). Dramatic changes in this minimal SUMOylome across various stress conditions reveal that instead of newer targets undergoing covalent modifications by SUMOs, SUMOylation levels on prior-SUMOylated proteins pools are more altered (Miller et al., 2013). Likely, functional inactivation of SUMO proteases especially noted post-stress in mammalian systems (Bossis and Melchior, 2006; Pinto et al., 2012; Sydorskyy et al., 2010) and enhanced pool of SUMOylated pool SIZ1 (Miller et al., 2013) coordinate to achieve this. Taking this into account, a host SUMOylome output is likely to be influenced by relative changes in SUMO isoforms. Indeed, we demonstrate that transcripts of *SIZ1*, *EDS4* or *SAE2* are up-regulated basally in *sum1-1*, and in both Col-0 or *sum1-1* upon pathogen challenge in a *SUM3*-dependent manner (Figure 8). Hence the requirement of *SUM3* to maintain and adjust optimal level of SUMO1/2 SUMOylome is clearly evident. Non-covalent associations with SUMOs also affect chromatin architecture via their interaction with DNA methyltransferases and demethylases (Elrouby et al., 2013). Expression of FLOWERING LOCUS C (FLC), a central transcription factor of flowering is dynamically regulated by DNA

methylation status while the protein function is modulated by HPY2, SIZ1 and SUMO proteases
(Hepworth and Dean, 2015; Kong et al., 2017). FLC regulation presents a classical example of
this versatility of SUMO-influences on a developmental process.

Stimulus-driven SUMO isoform switches and subsequent fate of substrates have been widely
documented in animal systems (Meulmeester *et al.*, 2008; Zhu *et al.*, 2008; Zhu *et al.,* 2009).
The mammalian GTPase activating protein RanGAP1 although is equally modified by
SUMO1/2/3 *in vitro*, conjugation *in vivo* to SUMO1, but not SUMO2, imparts more stability
from isopeptidases thereby facilitating its association with Nup358 (Zhu et al., 2009). Similarly,
HDAC1 is targeted for degradation upon SUMOylation by SUMO1, but not SUMO2, in
cancerous cell lines thus potentiating invasive properties of the tumour (Citro et al., 2013).
Specific SUMO-proteases also regulate stimulus-dependent SUMO isoform switching (Fasci
et al., 2015). Upon arsenic exposure, SUMO2 at Lys^{65} of PML (Promyelocytic leukemia
protein) is replaced with SUMO1 to direct its ubiquitinylation. Clearly with the isoform
selectivity regulated at multiple levels, a host SUMOylome undergoes dynamic changes in
response to physiological perturbations. Evidences on SUMO isoform switches however are
completely lacking from plant systems. Nonetheless several reports suggest the likelihood of
this phenomenon. The effector XopD from *Xcv* is a SUMO3-specific isopeptidase (Colby et al.,
2006). The replication protein AL1 from Tomato Golden Mosaic Virus (TGMV) and the RNA-
dependent RNA polymerase NIb from Turnip mosaic virus (TuMV) interact with and require
SCE1 for replication (Sanchez-Duran et al., 2011; Xiong and Wang, 2013).

While our results undoubtedly place SUMO3 as a strong contributor in regulating dynamics
changes in a host SUMOylome, the loss of *SUM3* in many *Brassicaeae* and other higher
eudicots still remain an enigma (Hammoudi et al., 2016). Whether functions of *SUM3* has been
incorporated in *SUM1/2* roles in these plants remains to be explored further. We foresee the
immediate need of a plant system that would facilitate enrichment via distinct affinity tags on
individual SUMO paralogs in order to obtain evidences of SUMO preference, switches and
intersections. To summarize, our data lay the foundation on functional impingements of
Arabidopsis SUMO isoforms and their adjustments on global SUMOylome in response to
physiological changes.

Acknowledgements

IKD, MK and SB deeply acknowledge DST-SERB (Grant No: EMR/2016/001899) and

Regional Centre for Biotechnology (RCB), Faridabad for providing financial support and

central instrumental facilities. We express our gratitude to Prof. Walter Gassmann, Missouri

University, USA, and Prof. Ashis Kumar Nandi, Jawaharlal Nehru University (JNU), New

Delhi, India for providing *PstDC3000* strains and *sid2-1* lines and SA measurements,

respectively used in this study. We also thank Dr. Souvik Bhattacharjee (JNU) for providing

*p*ASK-IBA16 plasmid and Prof. Katsunori Tanaka, (Kwansei Gakuin University, Japan) for

providing *E. coli* SUMOylation reconstitution system constructs. IKD acknowledges

Department of Biotechnology (DBT), Government of India for providing fellowships during

his PhD tenure and KIIT University, Bhubaneswar, India for his PhD registration. MK thanks

UGC for his PhD fellowship.

Author Contributions

SB conceived the research. IKD, MK and SB designed the research. IKD and MK performed

all the experiments. HvdB generated *amiR-SUM2 sum3-1* line used in this study. SB helped in

plasmid constructions and supervised the experiments. IKD, MK and SB analyzed the data and

wrote the manuscript.

References

An L, Zhou Z, Yan A, Gan Y. 2011. Progress on trichome development regulated by

phytohormone signaling. *Plant Signal Behav* **6**:1959–1962. doi:10.4161/psb.6.12.18120

Augustine RC, Vierstra RD. 2018. SUMOylation: re-wiring the plant nucleus during stress and

development. *Curr Opin Plant Biol* **45**:143–154. doi:10.1016/j.pbi.2018.06.006

Bailey M, Srivastava A, Conti L, Nelis S, Zhang C, Florance H, Love A, Milner J, Napier R,

Grant M, Sadanandom A. 2015. Stability of small ubiquitin-like modifier (SUMO)

proteases OVERLY TOLERANT to SALT1 and -2 modulates salicylic acid signalling

and SUMO1/2 conjugation in Arabidopsis thaliana. *J Exp Bot* **67**:353–363.

doi:10.1093/jxb/erv468

Becker D. 1990. Binary vectors which allow the exchange of plant selectable markers and

reporter genes. *Nucleic Acids Res* **18**:203. doi:10.1093/nar/18.1.203

Bhattacharjee S, Halane MK, Kim SH, Gassmann W. 2011. Pathogen effectors target

Arabidopsis EDS1 and alter its interactions with immfune regulators. *Science* **334**:1405–

1408. doi:10.1126/science.1211592

Bossis G, Melchior F. 2006. Regulation of SUMOylation by reversible oxidation of SUMO

conjugating enzymes. *Mol Cell* **21**:349–357. doi:10.1016/j.molcel.2005.12.019

Budhiraja R, Hermkes R, Müller S, Schmidt J, Colby T, Panigrahi K, Coupland G, Bachmair

A. 2009. Substrates related to chromatin and to RNA-dependent processes are modified

by Arabidopsis SUMO isoforms that differ in a conserved residue with influence on

desumoylation. *Plant Physiol* **149**:1529–40. doi:10.1104/pp.108.135053

Capili AD, Lima CD. 2007. Structure and Analysis of a Complex between SUMO and Ubc9

Illustrates Features of a Conserved E2-Ubl Interaction. *J Mol Biol* **369**:608–618.

doi:https://doi.org/10.1016/j.jmb.2007.04.006

Castaño-Miquel L, Mas A, Teixeira I, Seguí J, Perearnau A, Thampi BN, Schapire AL, Rodrigo

N, La Verde G, Manrique S, Coca M, Lois LM. 2017. SUMOylation Inhibition Mediated

by Disruption of SUMO E1-E2 Interactions Confers Plant Susceptibility to Necrotrophic

Fungal Pathogens. *Mol Plant* **10**:709–720. doi:10.1016/j.molp.2017.01.007

Castaño-Miquel L, Seguí J, Lois LM. 2011. Distinctive properties of Arabidopsis SUMO

paralogues support the in vivo predominant role of AtSUMO1/2 isoforms. *Biochem J*

436:581–90. doi:10.1042/BJ20101446

Chen LZ, Li XY, Huang H, Xing W, Guo W, He J, Sun ZY, Luo AX, Liang HP, Hu J, Xu X,

Xu YS, Wang ZG. 2014. SUMO-2 promotes mRNA translation by enhancing interaction

between eIF4E and eIF4G. *PLoS One* **9**. doi:10.1371/journal.pone.0100457

Cheong MS, Park HC, Hong MJ, Lee J, Choi W, Jin JB, Bohnert HJ, Lee SY, Bressan RA,

Yun DJ. 2009. Specific domain structures control abscisic acid-, salicylic acid-, and stress-

mediated SIZ1 phenotypes. *Plant Physiol* **151**:1930–1942. doi:10.1104/pp.109.143719

Chosed R, Tomchick DR, Brautigam CA, Mukherjee S, Negi VS, Machius M, Orth K. 2007.

Structural analysis of Xanthomonas XopD provides insights into substrate specificity of

ubiquitin-like protein proteases. *J Biol Chem* **282**:6773–6782.

doi:10.1074/jbc.M608730200

Citro S, Jaffray E, Hay RT, Seiser C, Chiocca S. 2013. A role for paralog-specific sumoylation

in histone deacetylase 1 stability. *J Mol Cell Biol* **5**:416–427. doi:10.1093/jmcb/mjt032

Clough SJ, Bent AF. 1998. Floral dip: A simplified method for Agrobacterium-mediated

transformation of Arabidopsis thaliana. *Plant J* **16**:735–743. doi:10.1046/j.1365-

313X.1998.00343.x

Colby T, Matthäi A, Boeckelmann A, Stuible HP. 2006. SUMO-conjugating and SUMO-

deconjugating enzymes from Arabidopsis. *Plant Physiol* **142**:318–332.

doi:10.1104/pp.106.085415

Cui H, Gobbato E, Kracher B, Qiu J, Bautor J, Parker JE. 2016. A core function of EDS1 with

PAD4 is to protect the salicylic acid defense sector in Arabidopsis immunity.
doi:10.1111/nph.14302

de Vega D, Newton AC, Sadanandom A. 2018. Post-translational modifications in priming the
plant immune system: ripe for exploitation? *FEBS Lett* **592**:1929–1936.
doi:10.1002/1873-3468.13076

Defraia CT, Schmelz EA, Mou Z. 2008. A rapid biosensor-based method for quantification of
free and glucose-conjugated salicylic acid. *Plant Methods* **4**:1–11. doi:10.1186/1746-
4811-4-28

Dempsey DA, Vlot AC, Wildermuth MC, Klessig DF. 2011. Salicylic Acid Biosynthesis and
Metabolism. *Arab B* **9**:e0156. doi:10.1199/tab.0156

Duan G, Walther D. 2015. The Roles of Post-translational Modifications in the Context of
Protein   Interaction   Networks.   *PLoS   Comput   Biol*   **11**:1–23.
doi:10.1371/journal.pcbi.1004049

Duda DM, van Waardenburg RCAM, Borg LA, McGarity S, Nourse A, Waddell MB, Bjornsti
M-A, Schulman BA. 2007. Structure of a SUMO-binding-motif Mimic Bound to Smt3p–
Ubc9p: Conservation of a Non-covalent Ubiquitin-like Protein–E2 Complex as a Platform
for Selective Interactions within a SUMO Pathway. *J Mol Biol* **369**:619–630.
doi:https://doi.org/10.1016/j.jmb.2007.04.007

Elrouby N. 2014. Extent and significance of non-covalent SUMO interactions in plant
development. *Plant Signal Behav* **9**:e27948. doi:10.4161/psb.27948

Elrouby N, Bonequi MV, Porri A, Coupland G. 2013. Identification of Arabidopsis SUMO-
interacting proteins that regulate chromatin activity and developmental transitions. *Proc
Natl Acad Sci U S A* **110**:19956–61. doi:10.1073/pnas.1319985110

Elrouby N, Coupland G. 2010. Proteome-wide screens for small ubiquitin-like modifier
(SUMO) substrates identify Arabidopsis proteins implicated in diverse biological
processes. *Proc Natl Acad Sci U S A* **107**:17415–17420. doi:10.1073/pnas.1005452107

Fasci D, Anania V, Lill J, Salvesen G. 2015. Response to comment on "sumo deconjugation is
required for arsenic-triggered ubiquitylation of PML." *Sci Signal* **9**:1–12.
doi:10.1126/scisignal.aad9777

Flotho A, Melchior F. 2013. Sumoylation: A Regulatory Protein Modification in Health and
Disease. *Annu Rev Biochem* **82**:357–385. doi:10.1146/annurev-biochem-061909-093311

Gareau JR, Lima CD. 2010. The SUMO pathway: Emerging mechanisms that shape
specificity, conjugation and recognition. *Nat Rev Mol Cell Biol* **11**:861–871.
doi:10.1038/nrm3011

Hammoudi V, Fokkens L, Beerens B, Vlachakis G, Chatterjee S, Arroyo-Mateos M, Wackers
PFK, Jonker MJ, van den Burg HA. 2018. The Arabidopsis SUMO E3 ligase SIZ1
mediates the temperature dependent trade-off between plant immunity and growth. *PLOS*
Genet **14**:e1007157. doi:10.1371/journal.pgen.1007157

Hammoudi V, Vlachakis G, Schranz ME, van den Burg HA. 2016. Whole-genome duplications
followed by tandem duplications drive diversification of the protein modifier SUMO in
Angiosperms. *New Phytol* **1**. doi:10.1111/nph.13911

Hendriks IA, D'Souza RCJ, Yang B, Verlaan-de Vries M, Mann M, Vertegaal ACO. 2014.
Uncovering global SUMOylation signaling networks in a site-specific manner. *Nat Struct*
Mol Biol **21**:927–936. doi:10.1038/nsmb.2890

Hepworth J, Dean C. 2015. Flowering locus C's lessons: Conserved chromatin switches
underpinning developmental timing and adaptation. *Plant Physiol* **168**:1237–1245.
doi:10.1104/pp.15.00496

Hotson A, Chosed R, Shu H, Orth K, Mudgett MB. 2003. Xanthomonas type III effector XopD
targets SUMO-conjugated proteins in planta. *Mol Microbiol* **50**:377–389.
doi:10.1046/j.1365-2958.2003.03730.x

Kim SH, Gao F, Bhattacharjee S, Adiasor JA, Nam JC, Gassmann W. 2010. The arabidopsis
resistance-like gene SNC1 is activated by mutations in SRFR1 and contributes to
resistance to the bacterial effector AvrRps4. *PLoS Pathog* **6**.
doi:10.1371/journal.ppat.1001172

Kong X, Luo X, Qu GP, Liu P, Jin JB. 2017. Arabidopsis SUMO protease ASP1 positively
regulates flowering time partially through regulating FLC stability. *J Integr Plant Biol*
59:15–29. doi:10.1111/jipb.12509

Kurepa J, Walker JM, Smalle J, Gosink MM, Davis SJ, Durham TL, Sung DY, Vierstra RD.
2003. The small ubiquitin-like modifier (SUMO) protein modification system in
Arabidopsis. Accumulation of sumo1 and -2 conjugates is increased by stress. *J Biol*
Chem **278**:6862–6872. doi:10.1074/jbc.M209694200

Lee J, Nam J, Park HC, Na G, Miura K, Jin JB, Yoo CY, Baek D, Kim DH, Jeong JC, Kim D,
Lee SY, Salt DE, Mengiste T, Gong Q, Bohnert HJ, Kwak S, Bressan RA, Hasegawa PM,
Yun D. 2006. Salicylic acid-mediated innate immunity in Arabidopsis is regulated by
SIZ1 SUMO E3 ligase 79–90. doi:10.1111/j.1365-313X.2006.02947.x

Li Y, Li S, Bi D, Cheng YT, Li X, Zhang Y. 2010. SRFR1 negatively regulates plant NB-LRR
resistance protein accumulation to prevent autoimmunity. *PLoS Pathog* **6**:1–11.
doi:10.1371/journal.ppat.1001111

Lois LM, Lima CD, Chua NH. 2003. Small ubiquitin-like modifier modulates abscisic acid
signaling in Arabidopsis. *Plant Cell* **15**:1347–1359. doi:10.1105/tpc.009902

Matic I, Hagen M Van, Schimmel J, Macek B, Ogg SC. 2013. Europe PMC Funders Group In
Vivo Identification of Human Small Ubiquitin-like Modifier Polymerization Sites by
High Accuracy Mass Spectrometry and an in Vitro to in Vivo Strategy *, , S **7**:1–24.
doi:10.1074/mcp.M700173-MCP200.In

Mazur MJ, Kwaaitaal M, Mateos MA, Maio F, Kini RK, Prins M, van den Burg HA. 2019.
The SUMO conjugation complex self-assembles into nuclear bodies independent of SIZ1
and COP1. *Plant Physiol* **179**:168–183. doi:10.1104/pp.18.00910

Meulmeester E, Kunze M, Hsiao HH, Urlaub H, Melchior F. 2008. Mechanism and
Consequences for Paralog-Specific Sumoylation of Ubiquitin-Specific Protease 25. *Mol*
Cell **30**:610–619. doi:10.1016/j.molcel.2008.03.021

Miller MJ, Barrett-Wilt G a, Hua Z, Vierstra RD. 2010. Proteomic analyses identify a diverse
array of nuclear processes affected by small ubiquitin-like modifier conjugation in
Arabidopsis. *Proc Natl Acad Sci U S A* **107**:16512–16517. doi:10.1073/pnas.1004181107

Miller MJ, Scalf M, Rytz TC, Hubler SL, Smith LM, Vierstra RD. 2013. Quantitative
proteomics reveals factors regulating RNA biology as dynamic targets of stress-induced
SUMOylation in Arabidopsis. *Mol Cell Proteomics* **12**:449–63.

Morrell R, Sadanandom A. 2019. Dealing With Stress: A Review of Plant SUMO Proteases.
Front Plant Sci **10**:1–19. doi:10.3389/fpls.2019.01122

Niu D, Lin XL, Kong X, Qu GP, Cai B, Lee J, Jin JB. 2019. SIZ1-Mediated SUMOylation of
TPR1 Suppresses Plant Immunity in Arabidopsis. *Mol Plant* **12**:215–228.
doi:10.1016/j.molp.2018.12.002

Nukarinen E, Tomanov K, Ziba I, Weckwerth W, Bachmair A. 2017. Protein sumoylation and
phosphorylation intersect in Arabidopsis signaling. *Plant J* **91**:505–517.
doi:10.1111/tpj.13575

Ohama N, Kusakabe K, Mizoi J, Zhao H, Kidokoro S, Koizumi S, Takahashi F, Ishida T,
Yanagisawa S, Shinozaki K, Yamaguchi-Shinozaki K. 2016. The transcriptional cascade
in the heat stress response of arabidopsis is strictly regulated at the level of transcription
factor expression. *Plant Cell* **28**:181–201. doi:10.1105/tpc.15.00435

Okada S, Nagabuchi M, Takamura Y, Nakagawa T, Shinmyozu K, Nakayama JI, Tanaka K.
2009. Reconstitution of *Arabidopsis thaliana* SUMO pathways in *E. coli:* Functional
evaluation of SUMO machinery proteins and mapping of SUMOylation sites by mass
spectrometry. *Plant Cell Physiol* **50**:1049–1061. doi:10.1093/pcp/pcp056

Park HJ, Kim WY, Park HC, Lee SY, Bohnert HJ, Yun DJ. 2011. SUMO and SUMOylation
in plants. *Mol Cells* **32**:305–316. doi:10.1007/s10059-011-0122-7

Pinto MP, Carvalho AF, Grou CP, Rodríguez-Borges JE, Sá-Miranda C, Azevedo JE. 2012.
Heat shock induces a massive but differential inactivation of SUMO-specific proteases.
Biochim   Biophys   Acta   -   Mol   Cell   Res   **1823**:1958–1966.
doi:10.1016/j.bbamcr.2012.07.010

Ribet D, Cossart P. 2018. Ubiquitin, SUMO, and NEDD8: Key Targets of Bacterial Pathogens.
Trends Cell Biol **28**:926–940. doi:10.1016/j.tcb.2018.07.005

Roden J, Eardley L, Hotson A, Cao Y, Mudgett MB. 2004. Characterization of the
Xanthomonas AvrXv4 effector, a SUMO protease translocated into plant cells. *Mol Plant-*
Microbe Interact **17**:633–643. doi:10.1094/MPMI.2004.17.6.633

Saleh A, Withers J, Mohan R, Marqués J, Gu Y, Yan S, Zavaliev R, Nomoto M, Tada Y, Dong
X. 2015. Posttranslational modifications of the master transcriptional regulator NPR1
enable dynamic but tight control of plant immune responses. *Cell Host Microbe* **18**:169–
182. doi:10.1016/j.chom.2015.07.005

Sanchez-Duran MA, Dallas MB, Ascencio-Ibanez JT, Reyes MI, Arroyo-Mateos M, Ruiz-
Albert J, Hanley-Bowdoin L, Bejarano ER. 2011. Interaction between Geminivirus
Replication Protein and the SUMO-Conjugating Enzyme Is Required for Viral Infection.
J Virol **85**:9789–9800. doi:10.1128/jvi.02566-10

Saracco SA, Miller MJ, Kurepa J, Vierstra RD. 2007. Genetic analysis of SUMOylation in
Arabidopsis: Conjugation of SUMO1 and SUMO2 to nuclear proteins is essential. *Plant*
Physiol **145**:119–134. doi:10.1104/pp.107.102285

Schenk ST, Hernández-Reyes C, Samans B, Stein E, Neumann C, Schikora M, Reichelt M,
Mithöfer A, Becker A, Kogel KH, Schikora A. 2014. N-acyl-homoserine lactone primes
plants for cell wall reinforcement and induces resistance to bacterial pathogens via the
salicylic acid/oxylipin pathway. *Plant Cell* **26**:2708–2723. doi:10.1105/tpc.114.126763

Sharma P, Yamada S, Lualdi M, Dasso M, Kuehn MR. 2013. Senp1 Is Essential for
Desumoylating Sumo1-Modified Proteins but Dispensable for Sumo2 and Sumo3
Deconjugation   in   the   Mouse   Embryo.   *Cell   Rep*   **3**:1640–1650.
doi:10.1016/j.celrep.2013.04.016

Srivastava AK, Orosa B, Singh P, Cummins I, Walsh C, Zhang C, Grant M, Roberts MR,
Anand GS, Fitches E, Sadanandom A. 2018. Sumo suppresses the activity of the jasmonic
acid   receptor   CORONATINE   INSENSITIVE1.   *Plant   Cell*   **30**:2099–2115.
doi:10.1105/tpc.18.00036

Sydorskyy Y, Srikumar T, Jeram SM, Wheaton S, Vizeacoumar FJ, Makhnevych T, Chong
YT, Gingras A-C, Raught B. 2010. A Novel Mechanism for SUMO System Control:
Regulated Ulp1 Nucleolar Sequestration. *Mol Cell Biol* **30**:4452–4462.
doi:10.1128/mcb.00335-10

Tan CM, Li MY, Yang PY, Chang SH, Ho YP, Lin H, Deng WL, Yang JY. 2015. Arabidopsis
HFR1 is a potential nuclear substrate regulated by the xanthomonas type III effector
XopDXcc8004. *PLoS One* **10**:1–18. doi:10.1371/journal.pone.0117067

Tomanov K, Nehlin L, Ziba I, Bachmair A. 2018. SUMO chain formation relies on the amino-
terminal region of SUMO-conjugating enzyme and has dedicated substrates in plants.
Biochem J **475**:61–74. doi:10.1042/BCJ20170472

Tomanov K, Zeschmann A, Hermkes R, Eifler K, Ziba I, Grieco M, Novatchkova M, Hofmann
K, Hesse H, Bachmair A. 2014. Arabidopsis PIAL1 and 2 promote SUMO chain
formation as E4-type SUMO ligases and are involved in stress responses and sulfur
metabolism. *Plant Cell* **26**:4547–60. doi:10.1105/tpc.114.131300

van den Burg HA, Kini RK, Schuurink RC, Takken FLW. 2010. Arabidopsis small ubiquitin-
like modifier paralogs have distinct functions in development and defense. *Plant Cell*
22:1998–2016. doi:10.1105/tpc.109.070961

van den Burg HA, Takken FLW. 2010. SUMO-, MAPK- and resistance protein-signaling
converge at transcription complexes that regulate plant innate immunity. *Plant Signal*
Behav **5**:1597–1601. doi:10.4161/psb.5.12.13913

Verma V, Croley F, Sadanandom A. 2018. Fifty shades of SUMO: its role in immunity and at
the fulcrum of the growth–defence balance. *Mol Plant Pathol* **19**:1537–1544.
doi:10.1111/mpp.12625

Vertegaal ACO, Andersen JS, Ogg SC, Hay RT, Mann M, Lamond AI. 2006. Distinct and
overlapping sets of SUMO-1 and SUMO-2 target proteins revealed by quantitative
proteomics. *Mol Cell Proteomics* **5**:2298–2310. doi:10.1074/mcp.M600212-MCP200

Villajuana-Bonequi M, Elrouby N, Nordström K, Griebel T, Bachmair A, Coupland G. 2014.
Elevated salicylic acid levels conferred by increased expression of ISOCHORISMATE
SYNTHASE 1 contribute to hyperaccumulation of SUMO1 conjugates in the Arabidopsis
mutant early in short days 4. *Plant J* **79**:206–219. doi:10.1111/tpj.12549

Voigt CA. 2014. Callose-mediated resistance to pathogenic intruders in plant defense-related
papillae. *Front Plant Sci* **5**:1–6. doi:10.3389/fpls.2014.00168

Wang J, Taherbhoy AM, Hunt HW, Seyedin SN, Miller DW, Miller DJ, Huang DT, Schulman
BA. 2010. Crystal Structure of UBA2ufd-Ubc9: Insights into E1-E2 Interactions in Sumo

Pathways. *PLoS One* **5**. doi:10.1371/journal.pone.0015805

Watts FZ, Baldock R, Jongjitwimol J, Morley SJ. 2014. Weighing up the possibilities: Controlling translation by ubiquitylation and sumoylation 1–10.

Wildermuth MC, Dewdney J, Wu G, Ausubel FM. 2001. Isochorismate synthase is required to synthesize salicylic acid for plant defence. *Nature* **414**:562–565. doi:10.1038/417571a

Wimmer P, Schreiner S. 2015. Viral mimicry to usurp ubiquitin and sumo host pathways. *Viruses* **7**:4854–4877. doi:10.3390/v7092849

Xiong R, Wang A. 2013. SCE1, the SUMO-conjugating enzyme in plants that interacts with NIb, the RNA-dependent RNA polymerase of Turnip mosaic virus, is required for viral infection. *J Virol* **87**:4704–15. doi:10.1128/JVI.02828-12

Zhao X. 2018. SUMO-Mediated Regulation of Nuclear Functions and Signaling Processes. *Mol Cell* **71**:409–418. doi:10.1016/j.molcel.2018.07.027

Zhu J, Zhu S, Guzzo CM, Ellis NA, Ki SS, Cheol YC, Matunis MJ. 2008. Small ubiquitin-related modifier (SUMO) binding determines substrate recognition and paralog-selective SUMO modification. *J Biol Chem* **283**:29405–29415. doi:10.1074/jbc.M803632200

Zhu S, Goeres J, Sixt KM, Békés M, Zhang XD, Salvesen GS, Matunis MJ. 2009. Protection from Isopeptidase-Mediated Deconjugation Regulates Paralog-Selective Sumoylation of RanGAP1. *Mol Cell* **33**:570–580. doi:10.1016/j.molcel.2009.02.008

Figure Supplements

Additional supporting information may be found in the online version of this article.

Figure supplement 1: *sum3-1* partially alleviates, whereas *eds1-2* and *sid2-1* abolishes growth deficiencies of *sum1-1*.

Figure supplement 2: *sum1-1* and *sum2-1* plants are enhanced resistant whereas *sum3-1* is hypersusceptible to TNL-specific *PstDC3000(hopA1)* but not to CNL-specific *PstDC3000(avrRpm1)* avirulent strains.

Figure supplement 3: *sum1-1* and *sum2-1* leaves display increased whereas *sum3-1* is deficient in callose deposition in response to virulent *PstDC3000(EV)* or avirulent *PstDC3000(avrRps4)* infections.

Figure supplement 4: Developmental phenotype of 4-week-old SD grown plants of indicated genotypes.

Figure supplement 5: *sum1-1* and *sum2-1* display rapid whereas *sum3-1* is delayed than Col-0 in induction of *PR1* and *PR2* transcripts in response to virulent *PstDC3000(EV)* or avirulent *PstDC3000(avrRps4)* infections.

Figure supplement 6: Increased trichome density, elevated expression of PTI markers *FRK1* or *WRKY29*, and enhanced resistance to avirulent *PstDC3000(hopA1)* in *sum1-1* is SA-modulated.

Figure supplement 7: Enhanced basal immunity and ETI defences in *sum2-1* to TNL-specific *PstDC3000* strains is EDS1-dependent.

Figure supplement 8: Developmental defects and enhanced TNL-specific immunity in *sum1-1* is partially *SUM3* regulated.

Figure supplement 9: *SUM3* buffers increased induction of *PR1* in *sum1-1* in response to *PstDC3000* challenges.

Figure supplement 10: Schematic representation of key amino acid conservation and divergences among three *Arabidopsis* SUMO paralogs suggest their functional overlaps/distinctions.

Figure supplement 11: Complemented *sum1-1* or *sum3-1* lines have wild-type levels of SUMO1/2 conjugates upon *PstDC3000* (EV) infection.

Figure supplement 12: SUMO1 and SUMO3 cause reciprocal enhancements of SUMO-conjugates in *E. coli* SUMOylation-reconstitution system.

Figure supplement 13: *SUM3* mutation reduces SUMO1/2-conjugate enhancements during heat-shock.

Table supplement 1: List of primers used in this study.

 Figures with legends

Figure 1: *sum1* and *sum2* **mutants display enhanced whereas** *sum3* **is deficient in defenses against** *Pseudomonas syringae* **pv.** *tomato* **(***PstDC3000***) strains.** (A) Developmental phenotypes of 4-week-old Col-0, *sum1-1, sum2-1,* and *sum3-1* mutants propagated under short day (SD) and long day (LD) growth conditions. (B) Whisker Box plot of trichome densities (#Trichomes/4.1 mm^2) in Col-0 and *sum1-1, sum2-1,* and *sum3-1* mutants grown under SD conditions. Trichomes

in 4-week-old plants were counted after imaging leaves under bright field in a fluorescence microscope. Error bars indicate standard deviation of trichome densities in 10 random images of different leaf sections (n=10). *** indicates $p < 0.001$, ns= non-significant. (C-D) Growth of (C) virulent *PstDC3000(EV)* or (D) avirulent *PstDC3000*(*avrRps4*) strains, respectively in Col-0 and *sum1-1, sum2-1,* and *sum3-1* mutants. Three to four expanded leaves from 4-week-old plants of each genotype were infiltrated with the indicated bacterial suspension at a density of 5 X 10^4 cfu ml^{-1}. Leaf discs (of predefined diameter) were punched from the infiltrated area, macerated in 10mM $MgCl_2$, serially diluted and plated on appropriate antibiotic plates. Bacterial titer was calculated at 0- and 3-dpi (days post-infiltration) for the indicated bacterial infection. Whisker Box plot with Tukey test; n= 12; ANOVA was performed to measure statistical significant differences (at *p-value* <0.001) in growth of bacteria in Log_{10} scale. (E) Whisker Box plot of callose deposits/mm^2 area of infected leaves. Infiltrated leaves at 24-hpi (hours post-infiltration) were bleached in acetic acid: ethanol solution, stained with Aniline blue and were observed under DAPI filter in a fluorescence microscope. The images were analyzed in Imaris 8.0 software. The data is representative of callose deposits median from independent random sections area (n=10). Statistical significance was determined by t-test (*** indicates at $p < 0.001$).

1198

1199

Figure 2: Basal SA levels and expression of several defense-associated markers are elevated

in *sum1-1* or *sum2-1* whereas downregulated in *sum3-1* plants. (A) Total SA (SA + SAG) and

(B) free SA levels in *sum* and combinatorial mutants in 4-week-old plants measured by biosensor

Acinetobacter method. Data is representative of three biological replicates. Statistical significance

calculated by Student t-test is denoted by alphabet on top of each bar. Different alphabets indicate

significance at $p<0.001$. Relative transcript abundance of (C) *PR1*, *PR2*, *ICS1* (E) *FRK1*, *WRKY29*,

SNC1 (F) *SUM1*, *SUM2*, and *SUM3* in 4-week-old SD grown Col-0, *sum* and indicated

combinatorial mutants was determined by qRT-PCR and normalized to *SAND* expression. The

values are represented as fold change relative to Col-0 (WT). Data is representative of mean of

three biological replicates (n=3). Error bars indicate SD. Student *t*-test was performed to calculate

statistical significance; *=$p<0.05$; **=$p<0.01$; ***=$p<0.001$, ns= not significant. (D) Total protein

extracts from 4-week-old SD grown Col-0, *sum1-1*, *sum2-1*, and *sum3-1* plants were

immunoblotted with anti-PR2 antibodies. Ponceau S stain of the Rubisco subunit indicative of

equal protein loading between samples is shown. Migration positions of molecular weight

standards (in kDa) are indicated.

1215

1216

1217

Figure 3. Altered defense responses to *PstDC3000* strains in *sum1-1* or *sum3-1* plants are restored to wild-type levels in the respective complemented lines. (A) Relative transcript abundance of *SUM1, SUM2* and *SUM3* in 4-week-old SD grown Col-0, *sum* mutants and the corresponding complemented line(s) was determined by qRT-PCR and normalized to *SAND* expression. The values are represented as fold change relative to Col-0 (WT). Data is representative of mean of three biological replicates (n=3). Error bars indicate SD. Statistical significance was determined by Student *t*-test; **=*p*<0.01; ***=*p*<0.001, ns= not significant. (B) Total (*left panel*) and free SA (*right panel*) levels in *sum* mutants and the complemented lines. Data is representative of three biological replicates. Statistical significance was determined by Student *t*-test and represented by alphabets with significance at *p*<0.001. (C) Endogenous PR1 protein levels in *sum* and complemented lines. Immunoblot with anti-Actin antibodies and Ponceau S staining of membrane show equal protein loading among the extracts. (D) Growth of (*left panel*) virulent *PstDC3000(EV)* or (*right panel*) avirulent *PstDC3000(hopA1)* strains, respectively in the indicated plants. Leaves from 4-week-old plants of each line were infiltrated with the indicated bacterial suspension at a density of 5×10^4 cfu ml^{-1}. Leaf discs from the infiltrated area was macerated in 10mM MgCl$_2$, serially diluted and plated on appropriate antibiotic plates. Bacterial titers calculated at 0- and 3-dpi (days post-infiltration) for the indicated infection are shown. Different alphabets denote statistical significance at *p-value* <0.001.

1236

1237

Figure 4: *sum1-1* and *sum2-1* display enhanced basal and rapid induction of PR1 and PR2 proteins upon pathogen challenge whereas *sum3-1* plants are deficient in these responses. Total protein from 4-week-old SD grown Col-0, *sum1-1*, *sum2-1*, *sum3-1*, *sum1-1 sum3-1*, *sum1-1 eds1-2*, *sum2-1 eds1-2*, *eds1-2*, *sum1-1 sid2-1*, and *sid2-1* was extracted from (A) un-infiltrated or at 6-, 12-, and 24-hpi (hours post-infiltration) with either (B) virulent *PstDC3000(EV)* or (C) avirulent *PstDC3000(avrRps4)* strains and immunoblotted with anti-PR1 or anti-PR2 antibodies. The membranes were also probed with anti-Actin antibodies or stained with Ponceau S for Rubisco subunit to indicate comparable loading between extracts. Migration position of protein molecular weight standards (in kDa) are indicated.

1247

1248

Figure 5: Growth retardation and enhanced defenses in *sum1-1* are SA-dependent. (A)

Developmental phenotypes of 4-week-old Col-0, *sum1-1*, *sum1-1 eds1-2*, *eds1-2*, *sum1-1 sid2-1*,

and *sid2-1* plants grown under SD and LD growth conditions. (B) Relative transcript abundance

of *PR1* and *ICS1* in 4-week-old SD grown indicated plants was determined by qRT-PCR and

normalized to *SAND* expression. The values are represented as fold change relative to Col-0 (WT).

Data is representative of independent experiments (n=3). Error bars indicate SD. Student *t*-test

significances are mentioned (*=$p<0.05$; **=$p<0.01$; ***=$p<0.001$). Growth of (C) virulent

PstDC3000(EV) and (D) avirulent *PstDC3000* (*avrRps4*) strains, respectively in indicated plants

are shown. Three to four expanded leaves from 4-week-old SD grown plants of each genotype

were infiltrated with the indicated bacterial suspension at a density of 5 X 10^4 cfu ml^{-1}. Leaf discs

were punched from the infiltrated area, macerated in 10mM MgCl$_2$, serially diluted and plated on

appropriate antibiotic plates. Bacterial titer was calculated at 0- and 3-dpi for the indicated

infiltration. Whisker Box plot with Tukey test; n= 12-18; ANOVA was performed to measure

statistical significant differences (at *p-value* <0.001) in growth of bacteria in Log$_{10}$ scale.

1263

1264

Figure 6: *sum3-1* alleviates developmental defects and enhanced defense responses in *sum1-1*. (A) Developmental phenotypes of 4-week-old SD grown Col-0, *sum1-1, sum3-1* and *sum1-1 sum3-1* plants. (B) Total protein extracts from indicated plants were immunoblotted with anti-SNC1 antibodies. Ponceau S stain of the Rubisco subunit and anti-Actin immunoblot indicative of equal protein loading between samples are shown. Migration positions of molecular weight standards (in kDa) are indicated. Growth of (C) virulent *PstDC3000(EV)* or (D) avirulent *PstDC3000(avrRps4)* strains, respectively in indicated plants. Three expanded leaves from 4-week-old SD grown plants of each genotype were infiltrated with the respective bacterial suspension at a density of 5×10^4 cfu ml^{-1}. Leaf discs of predefined diameter were punched from the infiltrated area, macerated in 10mM MgCl$_2$, serially diluted and plated on appropriate antibiotic plates. Bacterial titer was calculated at 0- and 3-dpi with for indicated infiltration. Whisker Box plot with Tukey test; n= 12; ANOVA was performed to measure statistical significance (at *p-value* <0.001) in growth of bacteria in Log$_{10}$ scale.

1278

Figure 7: SUMO3, but not SUMO1, can form oligomers. (A) Green fluorescence indicative of YFP reconstitution between BiFC (Bi-molecular Fluorescence complementation) combinations of SUMOylation proficient (GG) or deficient (AA) forms of SUMO1 and SUMO3 paralogs are shown. *Agrobacterium* GV3101 strains expressing the indicated BiFC constructs were co-infiltrated in expanded *N. benthamiana* leaves. Infiltrated leaf sections were visualized under a confocal microscope at 2-days post-infiltration (dpi). (Scale bar = 50 μm). Images are

representative of pattern observed in two independent experiments. (B) *In vitro* binding assays
between SUMO1 and SUMO3 paralogs. His-or Strep-II-tagged SUMO1 or SUMO3 paralog as
indicated were expressed in *E. coli.* Bacterial lysates from indicated combinations were mixed and
enriched through the Ni^{2+}-NTA matrix. Enrichment of the mentioned His-SUMO and the co-
eluting paralog was investigated via immunoblotting with anti-His or anti-Strep antibodies,
respectively (pull-down panel). The input panel shows the protein levels in the extracts used for
the enrichments. Position of protein molecular weight standards (in kDa) are indicated.

1302

Figure 8: Loss of *SUM3* reduces SUMO1/2-conjugate enhancements in response to *PstDC3000(EV)* infection. (A) Fully expanded 3-4 leaves from 4-week-old SD grown Col-0, *sum1-1, sum3-1* and *sum1-1 sum3-1* plants were infiltrated either with buffer alone (- lanes) or with the virulent *PstDC3000(EV)* strain (+ lanes) at a density of 5 X 10⁶ cfu ml⁻¹. Total protein isolated from the infected leaves at 24 hrs post-infiltration (hpi) was used for immunoblotting with anti-SUMO1/2 antibodies. Approximate positions of SUMO1/2-conjugates are shown. Anti-Actin immunoblot or Ponceau S straining demonstrate comparable protein loadings from the extracts. Relative positions of protein molecular weight standards (in kDa) are indicated. Relative transcript abundance at 24-hpi of (B) *SAE1, SAE2, SCE1, SIZ1, HPY2* and (C) *ESD4, ELS1, OTS1, OTS2* in mock versus *PstDC3000(EV)* infiltrated samples of Col-0, *sum1-1* and *sum3-1* was determined by qRT-PCR and normalized to *SAND* expression. The values are represented as fold change relative to Col-0 (WT). Data is representative of mean of three biological replicates (n=3). Error bars indicate SD. Student *t*-test was performed to calculate statistical significance; *=*p*<0.05; **=*p*<0.01; ***=*p*<0.001, ns= not significant. (D) Genetic model for *SUM1/2* crosstalks with

SUM3 in regulation of SA-dependent defences and heat-shock responses in *Arabidopsis*. *SUM1/2*

function additively but not equally as negative regulators of defences via modulation of SA-

signaling routes and expression of defense-associated markers. Pathogen exposure induce SA-

dependent *SUM3* role as a positive regulator leading to increase in SUMO1/2-conjugation

proficiencies, and expression of response-appropriate markers such as PR proteins, PTI markers,

and callose deposits. Induced SUMO1/2-conjugates modulate SUMO3 functions and prevents

overshooting of responses. *SUM1* also regulates via *SUM3*-independent but SA-dependent

developmental aspects such as trichome production and leaf textures.